Cell type-specific network analysis in Diversity Outbred mice identifies genes potentially responsible for human bone mineral density GWAS associations

Luke J Dillard[1], Gina M Calabrese[1], Larry D Mesner[1,2], Charles R Farber[1,2,3]

[1]Center for Public Health Genomics, School of Medicine, University of Virginia, Charlottesville, VA 22908
[2]Department of Public Health Sciences, School of Medicine, University of Virginia, Charlottesville, VA 22908
[3]Department of Biochemistry and Molecular Genetics, School of Medicine, University of Virginia, Charlottesville, VA 22908

Correspondence to:

Charles R. Farber
E-mail: crf2s@virginia.edu
Center for Public Health Genomics
University of Virginia
P.O. Box 800717
Charlottesville, VA 22908, USA

Tel. 434-243-8584

Abstract

Genome-wide association studies (GWASs) have identified many sources of genetic variation associated with bone mineral density (BMD), a clinical predictor of fracture risk and osteoporosis. Aside from the identification of causal genes, other difficult challenges to informing GWAS include characterizing the roles of predicted causal genes in disease and providing additional functional context, such as the cell type predictions or biological pathways in which causal genes operate. Leveraging single-cell transcriptomics (scRNA-seq) can assist in informing BMD GWAS by linking disease-associated variants to genes and providing a cell type context for which these causal genes drive disease. Here, we use large-scale scRNA-seq data from bone marrow–derived stromal cells cultured under osteogenic conditions (BMSC-OBs) from Diversity Outbred (DO) mice to generate cell type-specific networks and contextualize BMD GWAS-implicated genes. Using trajectories inferred from the scRNA-seq data, we identify networks enriched with genes that exhibit the most dynamic changes in expression across trajectories. We discover 21 network driver genes, which are likely to be causal for human BMD GWAS associations that colocalize with expression/splicing quantitative trait loci (eQTL/sQTL). These driver genes, including *Fgfrl1* and *Tpx2,* along with their associated networks, are predicted to be novel regulators of BMD via their roles in the differentiation of mesenchymal lineage cells. In this work, we showcase the use of single-cell transcriptomics from mouse bone-relevant cells to inform human BMD GWAS and prioritize genetic targets with potential causal roles in the development of osteoporosis.

Keywords: Osteoblasts, Osteocytes, Osteoporosis, Mesenchymal Stem Cells (MSC), Stromal cells, Systems Biology

Introduction

Osteoporosis is a complex disease characterized by low bone mineral density (BMD), bone fragility, and an increased risk of fracture[1]. BMD, a highly heritable trait, is one of the most important clinical predictors of osteoporotic fracture[2,3]. Increasing our understanding of the genetic basis of BMD is critical for the development of approaches for the treatment and prevention of osteoporosis. In recent years, genome-wide association studies (GWAS) have made great progress in unraveling BMD genetics by identifying over 1,100 independent associations[4]. Now the challenge lies in pinpointing causal genes, which is necessary for the translation of genetic findings into novel therapies.

A number of approaches exist to identify genes responsible for GWAS associations[5–8]. Most rely on population-based "-omics" data[9], which are scarce for human bone, to connect associations to causal genes. Recently, our group has used co-expression networks generated from mouse bone transcriptomic datasets to assist in the identification of genes likely responsible for BMD associations. The idea is simple, genes that play a central role in the regulation of a complex trait are often functionally-related and functionally-related genes are often co-expressed[10]. By identifying networks enriched for genes implicated in GWAS we predicted putatively causal genes for hundreds of BMD associations based on their membership in enriched modules. One significant advantage of this approach is its ability to utilize the network connections between candidate genes and genes already established to influence bone function. This enables us to predict how these candidate genes may affect BMD. For example, we generated co-expression networks from bone tissue and primary osteoblasts in mouse genetic reference populations and identified multiple co-expression modules enriched with genes located in BMD associations[11,12]. We then cross-referenced genes in these modules with those regulated by co-localizing expression

quantitative trait loci (eQTLs) from the Gene-Tissue Expression project (GTEx)[13,14] to identify

"high priority" genes. We showed that one of these genes, *MARK3*, a serine-threonine kinase[15],

influenced BMD in an osteoblast-autonomous manner through a role in osteoblast

differentiation[11,15]. Recently, we expanded our approach to include directed networks generated

via a Bayesian approach using cortical bone RNA-seq data from 192 Diversity Outbred (DO) mice.

By combining key driver analysis and GTEx eQTL colocalization data, we identified 19 novel

genes, such as *SERTAD4* and *GLT8D2*, which are likely causal for human BMD GWAS

associations[16].

To date, our analyses have been reliant on networks generated from heterogeneous bulk

transcriptomics (RNA-seq) datasets from mouse bone and primary bone cells. However,

leveraging single-cell transcriptomics (scRNA-seq) data would offer the added benefit of resolving

the transcriptomic profiles of discrete cell types. Additionally, using scRNA-seq data has the

potential to provide context by predicting the specific cell types in which causal genes and their

associated networks operate. In recent work, we demonstrated the utility of bone marrow-derived

stromal cells cultured under osteogenic condition (BMSC-OB) for the generation of population-

scale scRNA-seq data on bone relevant cell types[17]. The BMSC-OB model effectively enriches for

mesenchymal lineage cells (e.g., mesenchymal progenitors, osteoblasts, osteocyte-like cells) that

are highly relevant to the regulation of BMD.

In this work, our goal was to prioritize and contextualize genes implicated by BMD GWAS

using an expanded large-scale (N=80) BMSC-OB scRNA-seq dataset on bone cell types. We

accomplished this by first generating co-expression and Bayesian networks[16] for each BMSC-OB

mesenchymal cell type. We subsequently prioritized networks based on their enrichment for genes

exhibiting the most dynamic changes in expression across trajectories inferred from the scRNA-

seq data, thus highlighting networks likely associated with the differentiation of BMSC-OBs. We

then used these networks linked to osteogenic differentiation to prioritize genes with eQTL and/or

splicing quantitative trait loci (sQTL) which colocalize with BMD GWAS associations[7,18]. In

doing so, this analysis provides additional support for a role of these genes in the regulation of

BMD and highlights their potential roles in differentiation of cell types essential to skeletal health.

97
98   **Results**
99
100   **BMSC-OBs derived from DO mice yield diverse cell types that are enriched for**
101   **mesenchymal lineage cells:**
102
We cultured BMSCs from a total of 80 Diversity Outbred (DO) mice, a genetically diverse

outbred mouse population[19,20] (N=75 from the current study and N=5 from[17]; N = 49 male and

N = 31 females). We cultured BMSCs under osteogenic conditions and subsequently performed

scRNA-seq, as described in[17]. After stringent processing and quality control (Materials and

Methods), the dataset consisted of the expression of 21,831 genes across 139,392 cells. We

manually annotated 15 clusters ranging in size from 270 to 27,291 cells and identified cell types

of the mesenchymal lineage as well as various other cell types (**Fig. 1A**, **Supplementary Table**

S1).

Based on our prior BMSC-OB scRNA-seq study[17], we expected to identify a large

proportion of mesenchymal cells and a smaller fraction of non-mesenchymal cell types. Consistent

with this hypothesis, clusters associated with mesenchymal lineages accounted for 74.1% of all

cells (**Fig. 1A**). These included mesenchymal progenitor cells (MPCs), late mesenchymal

progenitors (LMPs), osteoblast progenitors (OBPs), two mature osteoblast populations (OB1 and

OB2), osteocyte-like cells (Ocy), and marrow adipogenic lineage progenitors (MALPs). The non-

mesenchymal cell types observed included macrophages, monocytes, granulocytes, T-cells, B-

cells, endothelial cells, and osteoclast-like cells (**Fig. 1A**). With regards to the mesenchymal cell

types, the only differences in cell clusters relative to our previous report[17] were the presence of

MPCs and two mature osteoblast clusters. Upon comparing the two distinct osteoblast clusters,

OB1 and OB2 (**Fig. 1A**), both clusters had ubiquitous expression of genes associated with mature

osteoblasts (e.g., *Col1a1*, *Bglap, Sparc*, and *Ibsp*) (**Supplementary Table S1**) while many of the

"canonical" osteoblast markers were more highly expressed in OB1 compared to OB2

(**Supplementary Table S2**). Interestingly, MPCs did not have transcriptomic profiles similar to

other mesenchymal progenitor cells previously identified by our group or others[17,21]. All other

mesenchymal cell types demonstrated specific expression of canonical marker genes (**Fig. 1A, B**).

127   We next assessed the variability in cell type frequencies across DO mice by quantifying

the proportions of each annotated mesenchymal cell type. All other clusters, which mainly

consisted of immune cells of hematopoietic origin, were aggregated into one group (Hem) for each

mouse. We observed high variability in the raw proportional abundances of cell types derived from

each mouse (**Fig. 1C, Supplementary Table S3**). For example, the proportions of osteoblasts

(OB1 and OB2) varied significantly among individual DO mice (**Fig. 1D**). All mice used in the

current experiment had been extensively phenotyped for a wide range of bone traits (microCT,

histomorphometry, biomechanical bone properties, etc.) as part of a previous genetic analysis of

bone strength[16]. We correlated cell type frequencies with bone traits, however, none of the cell

type proportions were strongly correlated with any bone trait (**Supplementary Table S4, S5**).

Figure 1. Analysis of single cell RNA-seq (scRNA-seq) data for BMSC-OBs derived from 80
Diversity Outbred (DO)

Mesenchymal lineage cells are enriched in BMD heritability:

The primary goal of this work was to prioritize and contextualize genes implicated by BMD GWAS. As a first step towards this goal, we sought to determine which cell types were the most relevant to the genetics of BMD. Using the BMD GWAS and the BMSC-OB scRNA-seq data, we performed a CELLECT[22] analysis to identify cell clusters enriched for BMD heritability. We observed that all mesenchymal cell types were nominally significantly ($P < 0.05$) enriched for BMD heritability (**Fig. 1E**, **Supplementary Tables S6**). The OB1, Ocy, and MALP cell types were significantly enriched for BMD heritability after adjusting for multiple comparisons ($P_{adj} < 0.05$). None of the non-mesenchymal cells identified were significant ($P > 0.05$) (**Fig. 1E**). As a result, all downstream efforts focused solely on using data on mesenchymal cell types to inform GWAS.

Generating mesenchymal cell type specific Bayesian networks to inform BMD GWAS:

We have previously shown that network-based approaches using bulk RNA-seq are powerful tools for the identification of putative causal genes from BMD GWAS data[11,12,16]. Here, our goal was to apply these same approaches using the BMSC-OB scRNA-seq data to prioritize and contextualize genes we previously identified as having a colocalizing expression quantitative trait locus (eQTL; N=512) or splicing QTL (sQTL; N=732) in a tissue from the Genotype-Tissue Expression (GTEx) project[7,14,18]. Genes identified in each study (or both) yielded a list of high priority target genes (N = 1,037). While GTEx does not currently contain data for bone tissue, eQTL can be shared across many tissues and may exert their effects in cell types resident to bone[23]. Therefore, utilizing our previous work, we hypothesized that generating cell type-specific networks would yield more biologically relevant representations of processes occurring within

particular mesenchymal cell types. Additionally, by integrating GWAS, cell type specific

networks, and dynamic gene expression as a function of differentiation we anticipated we would

identify points of intervention in which genetic variation impacts genes involved in the

differentiation process.

Our network analysis begins by partitioning genes into groups based on co-expression by

applying iterative weighted gene co-expression network analysis (iterativeWGCNA)[24] to each

mesenchymal cell type (Step 1, **Fig. 2**). In total, 535 modules were identified from the BMSC-OB

scRNA-seq data, and the number of modules identified for each mesenchymal cell cluster ranged

from 26 to 153 with an average of 76 modules per cluster (**Supplementary Table S7, S8**). We

sought to infer causal relationships between genes in each cell type-specific co-expression module

and subsequently identify networks involved in processes relevant to BMSC-OB differentiation.

To this end, we generated Bayesian networks for each co-expression module, thus enabling us to

model directed interactions between co-expressed genes based on conditional independence[16]

(Step 2, **Fig. 2**).

180
181 **Figure 2. Overview of the network analysis pipeline**
182

183 **Identifying putative drivers of mesenchymal cell differentiation:**
184
We hypothesized that many genes impacting BMD do so by influencing osteogenic

differentiation or possibly bone marrow adipogenic differentiation, as suggested by the CELLECT

analysis above. Therefore, the next step of our network analysis was to identify cell type-specific

Bayesian networks enriched for genes potentially driving mesenchymal differentiation (Step 3,

Fig. 2). To accomplish this, we first performed a pseudotime trajectory analysis to infer paths of

differentiation in the mesenchymal lineage cells. We resolved three pseudotime trajectories (two

osteogenic, one adipogenic) originating from the MPC cell cluster and ending in either Ocy, OB2,

or MALP cell fates (**Fig. 3A**). It is important to note that given the identification of multiple

skeletal stem cells[25–28], we do not view these trajectories as lineages, but rather "differentiation

paths" (progenitor to mature and/or terminally differentiated cells) that are likely traversed by

different subsets of skeletal stem cells.

To identify genes likely impacting BMSC-OB differentiation, we used tradeSeq to identify

genes that exhibit dynamic changes in expression along pseudotime[29]. Prior to performing the

tradeSeq analysis, we parsed the pseudotime trajectories into regions that encompass cells

associated with each cell type along their respective trajectories (**Fig. 3B**). We defined multiple

cell type boundaries (nine in total) using pseudotime values, which represent points along a

trajectory. The tradeSeq analysis was performed for each boundary (**Supplementary Table S9**).

For example, trajectories bifurcate in the LMP cell cluster (**Fig. 3A**); therefore, cells belonging to

the LMP cluster can map to adipogenic and/or osteogenic trajectories depending on their

placement along pseudotime. Between a cell type boundary, cells spanning a specific cluster (e.g.,

LMP) and mapping to a specific trajectory (e.g., osteogenic trajectory) are used as input to analyze

gene expression between the start and end points of the cell type boundary (e.g., LMP_to_OBP).

We analyzed gene expression within the established cell type boundaries for all trajectories and

identified genes that exhibit the most significant differences in expression between the start and

end points of the cell type boundaries. The total number of significant trajectory-specific tradeSeq

genes ($P_{adj} < 0.05$) ranged from 87 to 1,697 across the 9 cell type boundaries (**Supplementary**

Table S9, S10-12). The expression of representative marker genes for all cell types as a function

of pseudotime were consistent with boundaries defined for each cell type (**Fig. 3C**).

To provide further support that tradeSeq-identified genes are involved in differentiation,

we performed a cell type-specific expression quantitative trait locus (eQTL) analysis for each

mesenchymal cell type. We identified 563 genes (eGenes) regulated by a significant *cis*-eQTL in

specific cell types of the BMSC-OB scRNA-seq data (**Supplementary Table S14**). In total, 73

eGenes were also tradeSeq-identified genes in one or more cell type boundaries along their

respective trajectories (**Supplementary Table S9**).

Figure 3. Pseudotime Trajectory Inference analysis and establishment of cell type boundaries for tradeSeq analysis

We hypothesized that if tradeSeq genes were responsible for driving mesenchymal

differentiation, then the eQTLs that perturb their expression would also impact the proportion of

cells at different stages along the cell trajectories. Despite being significantly underpowered for

this analysis due to our relatively smaller sample size (N = 80), we identified two cell type-specific

eGenes where the genotype responsible for the *cis*-eQTL effect was also associated with cell type

proportions. The first of these genes was Pyruvate Kinase, muscle (*Pkm*), which was identified as

a significant global tradeSeq gene ($P_{adj} = 8.35 \times 10^{-8}$; **Supplementary Table S13**) associated with

the transition from LMPs to OBPs along an osteogenic trajectory (**Fig. 4A**). Moreover, *Pkm* served

as an eGene in the LMP cell cluster (LOD = 9.72; **Fig. 4B, Supplementary Table S14**). Mice

inheriting at least one PWK allele at this locus (N = 15) demonstrated lower *Pkm* expression (**Fig.**

4C) and a notable reduction in mature osteoblasts (OB1) and osteocyte-like cells (Ocy) proportions

(P = 0.030 and P = 0.026, respectively), while LMP proportions were unaffected (**Fig. 4D,**

Supplementary Table 15).

Similarly, S100 calcium binding protein A1 (*S100a1*) was an OBP to OB1 transition

tradeSeq gene ($P_{adj} = 0.023$; **Fig. 4A, Supplementary Table S13**) and an eGene in the OBP cell

cluster (LOD = 10.12; **Fig. 4B, Supplementary Table S14**). Mice inheriting at least one 129 allele

at this locus (N = 30) had higher *S100a1* expression, while the opposite was observed for mice

inheriting NZO alleles (N = 14) (**Fig. 4C**). Additionally, mice inheriting at least one 129 allele

showed a significant decrease in LMP proportion and increase in OB1 proportion (P = 0.008 and

P = 0.016, respectively) (**Fig. 4D, Supplementary Table S15**), while no significant differences

were observed in cell type proportions among mice inheriting NZO alleles at this locus

(**Supplementary Fig. 2, Supplementary Table S15**). These data support the role of tradeSeq-

identified genes in the differentiation of mesenchymal cell types.

Figure 4. TradeSeq-identified genes associated with BMSC-OB differentiation exhibit eQTL effects.

Identification of differentiation driver genes (DDG):

In order to discover BMSC-OB differentiation genes potentially responsible for BMD GWAS associations, the next step of our network analysis leveraged the trajectory-specific tradeSeq genes identified for each cell type boundary (**Supplementary Table S10-12**) to identify differentiation driver genes (DDGs) (Step 3, **Fig. 2**). We identified DDGs by extracting subnetworks (i.e., large 3-step neighborhoods; see Methods) for each gene in each cell type-specific Bayesian network and identifying those subnetworks enriched ($P_{adj} < 0.05$) for trajectory-specific tradeSeq genes for the cell type boundary. The analysis identified 408 significant DDGs (**Supplementary Table S16, S17-19**). We performed a PANTHER[30] Gene Ontology (GO) analysis for the cell type boundaries yielding a sufficient number of DDGs and found that DDGs for the osteogenic cell type boundaries (LMP_to_OBP, OBP_to_OB1, OBP_to_OB2) were enriched for genes associated with the cell cycle (GO:0007049; N = 23, 18, 23; P = 1.12 x 10^{-6}, 1.29 x 10^{-13}, 1.0 x 10^{-14}, respectively) (**Supplementary Table S20-22**). The DDGs for the adipogenic cell type boundary (LMP_to_MALP, MALP_to_end) were enriched for genes associated with extracellular matrix organization (GO:0030198; N = 10; P = 1.62 x 10^{-7}) and lipid metabolic processes (GO:0006629; N = 25; P = 1.83 x 10^{-11}), respectively (**Supplementary Table S23-24**). Across all 408 DDGs, 49 (12%) were identified in one or more cell type boundaries as genes with a significant alteration (P < 0.05) of whole-body BMD when knocked-out/down in mice, as reported by the International Mouse Knockout Consortium (IMPC)[31] (**Supplementary Table S17-19**).

We used our previously generated list of potentially causal BMD GWAS genes (N=1,037) to subsequently prioritize the DDGs (Step 4, **Fig. 2**). Of the 408 DDGs, 21 DDGs in one or more cell type boundaries were genes that have BMD GWAS associations that colocalize with

sQTL/eQTL (**Table 1**). The majority of these DDGs were identified in LMPs along both the

osteogenic (LMP_to_OBP) and adipogenic (LMP_to_MALP) trajectories (N = 10 and 6,

respectively; **Supplementary Table S16, S25**). The remaining DDGs were identified in OBPs

along both osteoblast trajectories (OBP_to_OB1, OBP_to_OB2; N = 1 and 3, respectively) and

MALPs (MALP_to_end; N = 6). Additionally, 3 of the 21 DDGs (*Tet1*, *Tpx2*, *Timp2*) are IMPC

genes that exhibit a significant alteration of BMD (**Supplementary Table S16, S25**).

Table 1: Prioritized Differentiation Driver Genes (DDGs) that have BMD GWAS
associations that colocalize with splicing/expression QTL (eQTL/sQTL) identified in a
Genotype-Tissue Expression project (GTEx) tissue. The tissue with the most significant
colocalization (RCP and/or H4PP) is listed for each DGG (26 total, 21 distinct), as determined
from Al-Barghouthi *et al.* (2022) and Abood *et al.* (2023) for eQTL and sQTL, respectively[7,18].
RCP = Regional Colocalization Probability (GWAS and eQTL colocalization). H4PP = H4
Posterior Probability (GWAS and sQTL colocalization).

Trajectory	Cell type boundary	DDG	GTEx Tissue with strongest eQTL colocalization (RCP)	GTEx Tissue with strongest sQTL colocalization (H4PP)
1	LMP to OBP	*Tet1*	Adipose (Visceral); 0.3191	-
1	LMP to OBP	*Tpx2*	Testis; 0.2031	-
1	LMP to OBP	*Cdk1*	-	Pituitary; 0.7795
1	LMP to OBP	*Ttyh3*	-	Liver; 0.9350
1	LMP to OBP	*Olfml3*	Artery (Aorta); 0.8048	-
1	LMP to OBP	*Izumo4*	-	Brain (Hypothalamus); 0.9182
1	LMP to OBP	*Sec24d*	Nerve (Tibial); 0.2677	
1	LMP to OBP	*Tmem263*	Adipose (Subcutaneous); 0.5704	Cultured cells (fibroblasts); 0.9716
1	LMP to OBP	*Lmf2*	-	Adrenal Gland; 0.8181
1	LMP to OBP	*Tln2*	Esophagus (Muscularis); 0.9697	-
1	OBP to OB1	*Kremen1*	Heart (Left Ventricle); 0.8686	-
2	OBP to OB2	*Kremen1*	Heart (Left Ventricle); 0.8686	-
2	OBP to OB2	*Ebf1*	-	Testis; 0.8760
2	OBP to OB2	*Lrp4*	Pancreas; 0.7943	-

3	LMP to MALP	*Ttyh3*	-	Liver; 0.9350
3	LMP to MALP	*Fgfrl1*	Cultured cells (fibroblasts); 0.1611	-
3	LMP to MALP	*Ebf1*	-	Testis; 0.8760
3	LMP to MALP	*Ppp1r12b*	-	Nerve (Tibial); 0.8807
3	LMP to MALP	*Rhoj*	Cultured cells (fibroblasts); 0.352	Breast; 0.7844
3	LMP to MALP	*Tln2*	Esophagus (Muscularis); 0.9697	-
3	MALP to end	*Adh1*	-	Esophagus (Gastroesophageal Junction); 0.9999
3	MALP to end	*Fgfrl1*	Cultured cells (fibroblasts); 0.1611	-
3	MALP to end	*Adcy5*	-	Esophagus (Gastroesophageal Junction); 0.8456
3	MALP to end	*Cnn2*	-	Spleen; 0.7743
3	MALP to end	*Mxra8*	-	Pituitary; 0.7545
3	MALP to end	*Timp2*	-	Testis; 0.9429

Network analysis predict Fgfrl1 and Tpx2 as novel regulators of BMD:

Here we highlight two DDGs that putatively impact human BMD via their roles in LMP

differentiation along either an adipogenic (*Fgfrl1*) or osteogenic (*Tpx2*) trajectory. Based on our

previous work[7], *Fgfrl1* (fibroblast growth factor receptor-like 1) was identified as a DDG with

significant human BMD GWAS associations that also colocalized with eQTL identified in the

cultured fibroblast GTEx tissue (RCP = 0.1611, **Table 1**). The *Fgfrl1* network was enriched for

tradeSeq-identified genes (N = 6 genes, $P_{adj} = 7.5 \times 10^{-3}$) for LMPs along an adipogenic trajectory

(**Fig. 5A**). An increase in the expression of all tradeSeq-identified genes for the *Fgfrl1* network

was observed (**Fig. 5B, Supplementary Table S12**). Importantly, the expression pattern for the

tradeSeq-identified genes were consistent with the cell type boundaries established for LMPs

differentiating along the adipogenic trajectory toward the MALP cell state (**Fig. 5B**). Furthermore,

in the surrounding *Fgfrl1* network, two genes (*Plpp3* and *Cfap100*) have significant human BMD

GWAS associations that also colocalized with sQTL in GTEx tissues, as reported in our previous

study[18]. In the *Fgfrl1* network, many other genes can be associated with adipocyte function (e.g.,

Lpl, *Plpp3*, *Igfbp4*)[32–34] and the maintenance of cilia (e.g., *Cfap100*, *St5* (*Denn2b*), *Mark1*)[35–37].

The other network we identified, the *Tpx2* network, was identified for LMPs along an

osteogenic trajectory (**Fig. 5C**). *Tpx2* (TPX2 microtubule nucleation factor) is a DDG with

significant human BMD GWAS associations that also colocalized with eQTL identified in the

Testis GTEx tissue (RCP = 0.2031, **Table 1**). The network was enriched for tradeSeq-identified

genes (N = 9 genes, P_{adj} = 5.7 x 10^{-7}) for LMPs differentiating along the osteogenic trajectory (**Fig.**

5C). Furthermore, the expression of the tradeSeq-identified genes for the *Tpx2* network were

consistent with the cell type boundaries established for LMPs differentiating along the osteogenic

trajectory toward the OBP (osteoblast progenitor) cell state (**Fig. 5D; Supplementary Table S10**).

The expression of these genes increase as LMPs differentiate into OBPs and subsequently decrease

upon reaching an OBP cell state. Additionally, *Tpx2* exhibited a significant alteration of BMD in

both male and female mutant mice (Genotype P-value = 1.03 x 10^{-3}) from IMPC (**Fig. 5E**). In

regards to the constituents of the surrounding *Tpx2* network, additional genes have been tested by

the IMPC and result in a significant impact on BMD, such as *Ube2c*, *Top2a*, and *Papss1*. Many

other genes in the *Tpx2* network can be associated with cellular division and proliferation,

including four of the genes of the kinesin family (*Kif*) motor protein genes[38]: *Kif4, Kif11, Kif15,*

Kif23.

 Figure 5. *Fgfrl1* **and** *Tpx2* **are prioritized DDGs and putative drivers of mesenchymal**
 differentiation.

Discussion

324
325
BMD GWAS has been successful at identifying thousands of SNPs associated with

disease; however, the identification of the causal genes and defining their functional role in disease

remains challenging. The integration of "-omics" data, particularly transcriptomics, can assist in

overcoming this challenge. Leveraging transcriptomics data has proven invaluable to informing

GWAS, as demonstrated in studies that use these data to perform eQTL mapping, transcriptome-

wide association studies (TWASs), and co-expression/gene-regulatory network reconstruction.

GWAS associations can colocalize with predicted sources of genetic variation that perturb causal

gene function or expression, thus providing a potential mechanism through which associations

impact disease. While bulk RNA-seq data has been the foundation of such analyses, scRNA-seq

data can provide valuable biological context by predicting the cell type in which causal genes are

affected. To inform BMD GWAS, the generation of population-scale transcriptomics data at

single-cell resolution in bone-relevant cell types can assist in the discovery of novel gene targets.

Here, we perform scRNA-seq on 80 DO mice to generate single-cell transcriptomics data of

mesenchymal cell types relevant to bone. Using these data, our goal was to prioritize putative

causal genes and provide biological context in which these genes potentially influence disease, at

cell type-specific resolution. Through our temporal gene expression and network analyses, we

identified 21 networks governed by predicted differentiation driver genes (DDGs) that have

corresponding human BMD GWAS associations colocalizing with eQTL/sQTL in a GTEx tissue.

We demonstrate that the BMSC-OB model serves as an effective method to enrich for

mesenchymal lineage cells, particularly bone-relevant cells. We characterized cells from 80 mice

and identified both osteogenic and adipogenic cells derived from the mesenchymal lineage, such

as two populations of osteoblasts (OB1 and OB2), osteocyte-like cells (Ocy), and MALPs. Our

trajectory inference analysis identified three distinct trajectories in which mesenchymal progenitor

cells give rise to both osteogenic and adipogenic cell types, thus portraying biologically relevant

and known paths of differentiation of mesenchymal progenitor cells. Temporal gene expression

was analyzed along each trajectory, in a cell type-specific fashion, to identify genes that were

changing the most as a function of pseudotime (tradeSeq-identified genes). Subsequent *cis*-eQTL

analysis indicated that the expression of some tradeSeq-identified genes were associated with the

relative proportion of cell types. While instances such as these were rare, they illustrate that the

potential consequence of genetic variation impacting the expression of tradeSeq-identified genes

may impact differentiation and the abundances of certain cell types along a trajectory.

To inform BMD GWAS, we utilized the scRNA-seq data in a network analysis to

contextualize causal genes (and their associated networks) by predicting the cell types through

which these genes are likely acting. Towards this goal, we generated cell type-specific Bayesian

networks from our BMSC-OB scRNA-seq data. Our approach was similar to our previous network

analyses where bulk RNA-seq data was leveraged to identify genes with strong evidence of playing

central roles in networks[11,12,16]. In contrast, here we utilized scRNA-seq data to identify DDGs and

prioritize networks based on the likelihood that they are involved in the differentiation of

mesenchymal lineage cells (based on network connections enriched for tradeSeq-identified genes

determined from inferred trajectories). Leveraging our previous work[7,18], we prioritized DDGs if

they were genes with BMD GWAS associations colocalizing with human eQTL/sQTL in a GTEx

tissue. Together, a gene being both a DDG and having BMD GWAS associations that colocalize

with eQTL/sQTL is strong support of causality.

We identified 21 DDGs and associated networks, some of which have little to no known

prior connection to bone. We contextualize these causal genes and their networks by not only

providing cell type predictions in which they likely operate, but also providing information

regarding the biological processes they likely affect. For example, the *Tpx2* network was identified

in LMPs differentiating along an osteogenic trajectory. *Tpx2* is a microtubule nucleation factor

that interacts with spindle microtubules during cellular division[39]. In our previous study, *Tpx2* was

identified as a gene that has BMD GWAS associations that colocalize with eQTL in the Testis

GTEx tissue[7]. While GTEx does not maintain bone tissue, eQTL are shared across many tissues[23];

therefore, non-bone eQTL may exert their effects in cell types associated with bone, such as LMPs,

and evidence of a human eQTL effect indicates that genetic variation can modulate the expression

of *Tpx2*. Additionally, when knocked out by IMPC, *Tpx2* exhibited a significant increase in whole

body BMD in mice; therefore, providing strong support for *Tpx2* influencing the regulation of

BMD in humans. In the surrounding gene neighborhood of the *Tpx2* network, other genes are

associated with cellular division as well, such as Topoisomerase 2A (*Top2a*) and the kinesin family

(*Kif*) genes[38,40]. Taken together, these results indicate a potential role of *Tpx2* as a mediator of

BMD via its role in microtubule maintenance during the expansion of osteogenic cell populations.

Additionally, the *Fgfrl1* network was identified in LMPs differentiating along an

adipogenic trajectory. Fibroblast growth factor receptor-like 1 (*Fgfrl1*) is presumed to function as

a decoy receptor that interacts with FGF ligands[41]. Our previous study also identified *Fgfrl1*, which

has BMD GWAS associations that colocalize with eQTL in the cultured fibroblasts GTEx tissue[16].

In the surrounding neighborhood of the *Fgfrl1* network, *Lpl*, *Plpp3*, *Igfbp4* have well-established

roles in adipocyte function and metabolism[32–34]; however, other genes can be associated with cilia

function, such as *Cfap100*, *St5* (*Denn2b*), *Mark1*[35–37]. Interestingly, the maintenance of cilia is

essential to the function of both LMPs and pre-adipocytes (MALPs) while mature adipocytes lack

cilia[42]. Therefore, the modulation of ciliogenesis and/or cilia function may coincide with *Fgfrl1*

signaling. Additionally, given the prioritization of MALPs in the CELLECT analysis and the well-

established inverse relationship between marrow adiposity and BMD[43,44], skewed balance of LMP

differentiation toward marrow adipogenic cell fates may affect BMD. In summary, the *Fgfrl1*

network harbors genes involved in adipogenic function, including cilia, which may contribute to

LMP differentiation along an adipogenic trajectory. Together, these results indicate a potential role

of *Fgfrl1* as a mediator of BMD via its role in adipogenic differentiation and maintenance of cilia.

Analyses performed here are not without limitations to consider. A pitfall of scRNA-seq is

the sparsity of the resulting data, which yields an increased frequency of zero values for the

expression of some genes in a proportion of cells, also known as "drop-outs"[45]. While statistical

approaches can be employed to impute missing data, the accuracy of such methods and whether

or not the resulting improvement in transcriptomic signal recovery is enough to warrant such

intervention is contentious[46,47]. However, this issue may be partially offset given the larger scale

of the scRNA-seq performed in this study and the average expression approach performed for

network and eQTL analysis. An additional limitation is that the BMSC-OB model does not capture

osteoclasts, another cell type associated with bone tissue. Importantly, results from our CELLECT

analysis indicate that BMD heritability was not enriched for genes whose expression was more

specific to osteoclast-like cells; however, these cell likely represent immature osteoclasts, as

mature multinucleated cells would be too large to be captured for sequencing. Lastly, while our

study employed 80 DO mice, the issue of statistical power is still a limitation; however, we

demonstrate that the BMSC-OB model is amenable to high throughput and the inclusion of

hundreds of mice, thus statistical power will be improved in future studies.

In summary, we showcase the use of large-scale scRNA-seq data to inform GWAS by

performing a network analysis to contextualize BMD GWAS associations. Through the use of

multiple single-cell analyses, we have expanded upon our understanding of the genetics of BMD.

Our work exemplifies the power of single-cell transcriptomics coupled with systems genetics to

discover the genetic determinants of disease.

Methods
422
423   **Sample preparation and scRNA-seq**
424
We prepared our samples in the same fashion as performed previously in Al-Barghouthi

and colleagues[17]. In brief, bone marrow was extracted from the femurs of initially 80 DO mice.

BMSCs were grown to confluence after 3 days of incubation in 48-well plates and then underwent

in vitro osteoblast differentiation for 10 days with osteogenic differentiation media (alpha MEM,

10% FBS, 1% pen/strep, 1% Glutamax, 50 μg/μL ascorbic acid [Sigma, St. Louis, MO, USA],

10 nM B-glycerophosphate [Sigma], 10 nM dexamethasome [Sigma]). After differentiation, single

cells were liberated from mineralizing cultures via incubations with 60 mM

ethylenediaminetetraacetic acid pH 7.4 (EDTA [Thermo Fisher Scientific], made in DPBS),

8 mg/mL collagenase (Gibco) in HBSS/4 mM $CaCl_2$ (Fisher), and 0.25% trypsin–EDTA (Gibco).

After single-cell isolation, cells from mice were pooled into groups containing cells from five mice

total and concentrated to 800 cells/μL in PBS supplemented with 0.1% BSA (bovine serum

albumin). Pooled single cells were prepared for sequencing using the 10× Chromium Controller

(10× Genomics, Pleasanton, CA, USA) with the Single Cell 3' v2 reagent kit, according to the

manufacturer's protocol. Libraries were sequenced on the NextSeq500 (Illumina, San Diego, CA,

USA).

scRNA-seq analysis pipeline
442

The data was subsequently processed using the 10× Genomics Cell Ranger toolkit (version

5.0.0) using the GRCm38 reference genome[48]. Using Seurat[49] (version 4.1.0), a combined Seurat

object containing all cells was generated with the inclusion of features detected in at least three

cells and cells with at least 200 features detected. We used Souporcell[50] (version 2.0.0) to

deconvolve the genotypes of all mice and to remove doublet cells. Cells were assigned to their

associated DO mouse by making a pairwise comparison between allele calls made by the shared

variants captured between Souporcell and GigaMUGA genotype arrays generated for all mice in

the cohort, as previous performed in Dillard and colleagues[17]. We filtered out cells with more than

6200 reads and less than 400 reads, as well as those cells with more than 10% mitochondrial reads.

Further, cells were removed if they expressed greater than 20% *Rpl* and 15% *Rps* reads, which

equates to cells approximately exceeding the 98 percentile. After filtering, 139,392 cells remained

and the resulting object underwent standard normalization, scaling, and the top 3000 features were

modeled from a variance stabilizing transformation (VST) using the Seurat. Cell-cycle markers

based on Tirosh and colleagues[51] were regressed out using the "CellCycleScoring" and scaling

functions. For subsequent dimensionality reduction, 15 principal components (PCs) were

summarized. Then, a kNN ($k = 20$) graph was created and the Louvain algorithm was used to

cluster cells at a resolution of 0.5. Annotation of cell type clusters was performed manually based

on differential gene expression analysis using the Seurat "FindAllMarkers" function

(**Supplementary Table S1**).

For subsequent WGCNA and eQTL mapping, transcriptomic profiles for each cell type

cluster were generated for each sample using a mean expression approach, as performed similarly

by others[52,53]. For each sample contributing at least 5 cells to a given cluster, unnormalized unique

molecular identifier (UMI) counts of gene expression for all cells in the cluster for the sample were

averaged and then rounded to the nearest hundredth decimal place. A total of 80, 80, 77, 67, 50,

76, 80 mice contributed enough cells to the MPC, LMP, OBP, OB1, OB2, Ocy, and MALP cell

type clusters, respectively. Genes with non-zero expression values in fewer than 15 samples were

removed. A total of 11971, 15162, 14857, 13674, 13825, 14136, and 14534 genes remained for

the MPC, LMP, OBP, OB1, OB2, Ocy, and MALP clusters, respectively. Samples were

normalized by computing CPMs (counts per million) without log transformation for each gene

using edgeR[54] (version 4.0.7), then transformed via VST using DESeq2[55] (version 1.42.0), and

quantile normalized using preprocessCore (version 1.60.2).

Trajectory and tradeSeq Analysis

477   Trajectory inference analysis was performed using Slingshot[56] (version 1.8.0) on the

mesenchymal lineage cell clusters (seven total) of the BMSC-OB scRNA-seq data. The starting

cluster was set as the MPC cluster and trajectories were inferred using 15 PCs. TradeSeq[29] (version

1.4.0) was used to analyze gene expression along the trajectories by fitting a negative binomial

generalized additive model (NB-GAM) to each gene using the "fitGAM" function with nknots =

10, which was determined by using the "evaluateK" function. Prior to performing the tradeSeq

analysis, the scRNA-seq data was downsampled to reduce the size of the dataset to approximately

10,000 cells (sampled at random across all seven clusters).

485   All cell type boundaries were established to encompass on average 78% of cells of a cell

cluster (**Supplementary Table S9**). To identify genes significantly changing between boundaries

in a trajectory-specific fashion, we first performed tradeSeq to compare gene expression within

each trajectory (two osteogenic, one adipogenic) to highlight genes with a significant difference

in expression between boundaries using the "startVsEndTest" function (**Supplementary Table**

S9, S10-12). Next, we performed a global test with tradeSeq to compare gene expression between

trajectories in order to highlight genes exhibiting a significant difference in expression using the

"startVsEndTest" function (**Supplementary Table S9, S13**). All tests were performed with the

$\log_2$ fold change threshold (l2fc) = 0.5. For all global and trajectory-specific tests, the P-values

associated with each gene were adjusted to control the false discovery rate using the "p.adjust"

function from the stats (version 4.2.1) R package and genes were filtered to include those with a

$P_{adj} < 0.05$.

CELLECT Analysis
499
CELLECT[22] (CELL-type Expression-specific integration for Complex Traits) (version

1.1.0) was used to identify likely etiologic cell types underlying complex traits of both the BMSC-

OBs scRNA-seq data (**Fig. 1E, Supplementary Table S6**). CELLECT quantifies the association

between the GWAS signal and cell type expression specificity using the S-LDSC genetic

prioritization model[57]. Summary statistics from the UK Biobank eBMD and Fracture GWAS (Data

Release 2018) and cell type annotations from each scRNA-seq data set were used as input. Cell

type expression specificities were estimated using CELLEX[22] (CELL-type EXpression-

specificity) (version 1.2.1) (**Supplementary Table S26**).

WGCNA
509
Cell type-specific mean expression matrices (as obtained above) were used as input to

generate signed co-expression network modules (**Supplementary Table S7, S8**).

IterativeWGCNA[24] (version 1.1.6) was used from a Singularity container built from a Docker hub

image[58]. A soft threshold (power) of 14, which exceeded a R^2 threshold of 0.85 for all cell type

clusters, was selected for module construction (**Supplementary Fig. 1**). Modules were generated

using iterativeWGCNA with default parameters for the "blockwiseModules" function, a minimum

module size of 20 genes, minCoreKME = 0.7, and minKMEtoStay = 0.5.

Bayesian network learning

Bayesian networks were learned from each of the cell type-specific modules of co-

expressed genes with the bnlearn (version 4.8.3). Gene expression matrices containing the genes

for each module were used as input to the "mmhc" function which employs the Max-Min Hill

Climbing algorithm (MMHC) algorithm[59] to learn the underlying structure of the Bayesian

network. From the generated networks, igraph (version 1.6.0) was used to resolve 3-step

neighborhoods[60]. Nodes (genes) that were unconnected to a neighborhood or connected to only

one neighbor were removed. Neighborhoods were filtered to include those with a size greater than

1 standard deviation from the mean across all neighborhoods generated for the network.

DDGs (differentiation driver genes) are genes that yield large 3-step neighborhoods that

are enriched (P_{adj} < 0.05) with tradeSeq-identified genes for a given cell type boundary. We

calculated whether each neighborhood contained more tradeSeq-identified genes (for the

neighborhoods' associated cell type boundary) than would be expected by chance using the

hypergeometric distribution ("phyper" function) from the stats (version 4.2.1) R package. The

arguments were as follows: q: (number of neighbors in a neighborhood that are also tradeSeq-

identified genes for a given cell type boundary) – 1; m: total number of tradeSeq-identified genes

for a given cell type boundary; n: (total number of identified neighborhoods) – m; k: neighborhood

size (total number of neighbors); lower.tail = false. P-values were adjusted to control the false

discovery rate using the "p.adjust" function from the stats (version 4.2.1) R package. These

pruning steps resulted in a total of 408 DDGs and associated networks for all cell types

(**Supplementary Table S16, S17-19**).

DO eQTL mapping
542

Prior to performing the eQTL analysis, DNA was extracted from the tails of the 80 DO

mice, using the PureLink Genomic DNA mini kit (Invitrogen) and genotyped using the

GigaMUGA array by Neogen Genomics (GeneSeek; Lincoln, NE). Processing and quality control

of genotype data, including calculation of genotype/allele probabilities, was performed as

previously described in Al-Barghouthi and colleagues[16]. Cell type-specific mean expression

matrices (as obtained above) for mesenchymal lineage clusters were used as input for the eQTL

mapping, which was performed using a linear mixed model (LMM) via the "scan1" function from

the qtl2[61] (version 0.30) R package with allowances for the following covariates: sex, age at

sacrifice (in days), weight, length, and DO mouse generation. To identify significant eQTL, we

calculated a LOD (logarithm of the odds) threshold; for each cell type cluster, we chose 50 genes

at random and then permuted them 1000 times using the "scan1perm" function from qtl2. We

established the LOD threshold of 9.68 and 9.49 for the autosomal chromosomes and X

chromosome, respectively, by taking the average of the median LOD across each cell type. A total

of 563 eQTL that exceeded the LOD thresholds and were no more than 1 Mbp from the

transcription start site of the associated eGene (**Supplementary Table S14**).

Cell type proportion analysis
560

To account for technical sources of variation often retained in scRNA-seq, cell type

proportions were transformed using the arcsin (asin) square root transformation from the speckle[62]

R package (version 0.0.3). Tests of statistical significance were performed using the propeller t-

test and ANOVA functions with default parameters. Sex of the mice and the batch each mouse

was associated with for sequencing were modeled as covariates. Transformed values were used as

input for computing tests of statistical differences of cell type proportions between mice, as well

as correlation to phenotypic traits (**Supplementary Table S3-S5**).

569
570 **Acknowledgements**

576 **Author Contributions**
577

578   Luke J Dillard: Writing - review & editing; Visualization; Conceptualization;

579 Methodology; Formal analysis; Data curation; Writing - original draft. Gina M Calabrese:

580 Methodology. Larry D Mesner: Methodology. Charles R Farber: Writing - review & editing;

581 Funding acquisition; Supervision; Conceptualization.

583 **Disclosures**
584
585   The authors declare that they have no conflicts of interest with the contents of this article.

Figure Legends

Figure 1. Analysis of single cell RNA-seq (scRNA-seq) data for BMSC-OBs derived from 80 Diversity Outbred (DO)
(A) Uniform Manifold Approximation and Projection (UMAP) of 139,392 single cells (BMSC-OBs). Cell numbers and corresponding percentages for the fifteen (15) annotated cell clusters are listed in parenthesis to the right of the annotated cluster name. **(B)** Dot plot[63] portraying representative and highly expressed genes for all annotated cell clusters. Dot color indicates the scaled average gene expression while the size of the dot corresponds to the percentage of cells of a given cluster that express a given gene. **(C)** The raw proportional abundances of seven (7) mesenchymal cell clusters and one (1) cluster (Hem) representing the remain cells (i.e., hematopoietic immune cells) across all 80 DO mice. **(D)** UMAP plots for mesenchymal lineage cell clusters for DO mouse 50 and DO mouse 233. **(E)** CELLECT (CELL-type Expression-specific integration for Complex Traits) cell type prioritization results. The seven (7) mesenchymal cell clusters were significantly enriched ($P < 0.05$) for BMD heritability and the MALP, Ocy, and OB1 cell clusters were significantly enriched after correcting P-value for multiple testing. All remaining cell clusters (i.e., hematopoietic immune cells) were not significant ($P > 0.05$)

Figure 2. Overview of the network analysis pipeline
Step 1: For all seven (7) of the mesenchymal lineage cell clusters (MPC, LMP, OBP, OB1, OB2, Ocy, MALP), cell type-specific co-expression modules were generated using iterative Weighted Gene Co-expression Network Analysis (iterativeWGCNA). *Step 2:* Bayesian networks were learned to generate directed networks and model causal interactions between co-expressed genes. *Step 3:* Differentiation Driver Genes (DDGs) were identified by extracting subnetworks (i.e., large 3-step neighborhood) for each gene in each cell type-specific Bayesian network and highlighting those subnetworks that were enriched ($P_{adj} < 0.05$) for trajectory-specific tradeSeq genes for the cell type boundary. *Step 4:* DDGs (and associated networks) were prioritized if the DDG was identified previously as an expression/splicing quantitative trait loci (eQTL/sQTL) that colocalized with BMD GWAS associations. Created with Biorender.com.

Figure 3. Pseudotime Trajectory Inference analysis and establishment of cell type boundaries for tradeSeq analysis
(A) Three (3) trajectories (two adipogenic, one adipogenic) were inferred from the mesenchymal cell clusters of the BMSC-OB scRNA-seq data using Slingshot. All trajectories originate from the MPC and end in either osteogenic (Ocy, OB2) or adipogenic (MALP) cell fates. **(B)** For each of the trajectories, cell type boundaries were generated using pseudotime values along the trajectories, which encompass the majority of cells of a cell type mapping to their respective trajectory. **(C)** Normalized gene expression of select genes associated with each cluster are represented in feature plots (*top*) and each gene plotted as a function of pseudotime (*bottom*) for all pseudotime trajectories (color corresponds to cell type annotation observed throughout). Vertical lines (red) represent the cell type (pseudotime) boundaries established for each cell type (label). The cell type boundary for OB1 and OB2 are represented as one red line/label for visualization purposes.

Figure 4. TradeSeq-identified genes associated with BMSC-OB differentiation exhibit eQTL effects.

(A) *Pkm* was identified as a significant global tradeSeq-identified gene ($P_{adj} = 8.35 \times 10^{-8}$) for LMP cells along an osteogenic trajectory (LMP_to_OBP) (*left*). *S100a1* was identified as a significant global tradeSeq-identified gene ($P_{adj} = 0.023$) for OBP cells along an osteogenic trajectory 1 (OBP_to_OB1) (*right*). **(B)** Plots indicating the cell type-specific expression quantitative trait loci (eQTL) signal for both *Pkm* and *S100a1*. A negative eQTL effect on *Pkm* expression was observed in LMPs for Diversity Outbred (DO) mice with a PWK haplotype background at the *Pkm* locus (*left*). A positive eQTL effect on the expression of *S100a1* was observed in OBPs for DO mice with a 129 haplotype background at the *S100a1* locus, while a negative effect was observed for NZO mice (*right*). **(C)** The expression of *Pkm* and *S100a1* based on DO mouse (expression values transformed via variance stabilizing transformation (VST), as described in Methods). Genotype abbreviations correspond to DO haplotype background (legend) at the respective gene locus. Mice with at least one PWK allele (genotype abbreviation G) tend to have decreased expression of *Pkm* (*left*). Mice with at least one 129 allele (genotype abbreviation C) tend to have increased expression of *S100a1*, while NZO mice (genotype abbreviation E) have decreased expression (*right*). **(D)** PWK mice had a significant reduction in mature osteoblasts (OB1) and osteocyte-like cells (Ocy) proportions relative to other mice (P = 0.030 and P = 0.026, respectively; t-test), while LMP proportions were unaffected. Asterisks represent any of the other haplotype backgrounds. 129 mice showed a significant decrease in LMP proportion and increase in OB1 proportion (P = 0.008 and P = 0.016, respectively; t-test), but OBP proportions were unaffected. No significant effects on cell type proportions were observed in NZO mice (**Supplementary Fig. 2**).

Figure 5. Fgfrl1 and Tpx2 are prioritized DDGs and putative drivers of mesenchymal differentiation.

(A) *Fgfrl1* was identified as a Differentiation Driver Gene (DDG) of a network for LMPs differentiating along an adipogenic trajectory. The network is enriched ($P_{adj} = 7.5 \times 10^{-3}$) for trajectory-specific tradeSeq-identified genes for the LMP_to_MALP cell type boundary (*Hnmt, St5, Igfbp4, Cyp1b1, Pdzrn4, Mark1*). *Fgfrl1* was previous identified as a gene that has BMD GWAS associations that colocalize with an eQTL in the cultured fibroblast GTEx tissue. **(B)** *Tpx2* was identified as a DDG of a network for LMPs differentiating along an osteogenic trajectory. The network is enriched ($P_{adj} = 5.7 \times 10^{-7}$) for tradeSeq-identified genes for the LMP_to_OBP cell type boundary (*Tpx2, Top2a, Kif4, Iqgap3, Prc1, Kif11, Ect2, Sgo2a, Ube2c*). *Tpx2* is both a tradeSeq gene and previous identified as a gene that has BMD GWAS associations that colocalize with an eQTL in the Testis GTEx tissue. **(C-D)** An increase in the expression of all tradeSeq-identified genes coincides with the cell type boundary in which they were identified as significant along their respective trajectories. **(E)** Box plot displaying whole-body bone mineral density (BMD) measurements (excluding skull) from the International Mouse Knockout Consortium (IMPC) for *Tpx2* mutant mice, which exhibited a significant increase in BMD (Genotype P-value = 1.03×10^{-3}) in both male and female mice (N = 8 (M) and 8 (F) mutants; N = 2574 (M) and 2633 (F) controls)

References

672
673
1.   Lin, J. T. & Lane, J. M. Osteoporosis: a review. *Clin. Orthop. Relat. Res*. 126–134 (2004).
2.   Peacock, M., Turner, C. H., Econs, M. J. & Foroud, T. Genetics of osteoporosis. *Endocr.*
Rev. **23**, 303–326 (2002).
3.   Johnell, O. *et al*. Predictive value of BMD for hip and other fractures. *J. Bone Miner. Res*.
20, 1185–1194 (2005).
4.   Morris, J. A. *et al*. An atlas of genetic influences on osteoporosis in humans and mice. *Nat.*
Genet. **51**, 258–266 (2019).
5.   Cookson, W., Liang, L., Abecasis, G., Moffatt, M. & Lathrop, M. Mapping complex disease
traits with global gene expression. *Nat. Rev. Genet*. **10**, 184–194 (2009).
6.   Wen, X., Pique-Regi, R. & Luca, F. Integrating molecular QTL data into genome-wide
genetic association analysis: Probabilistic assessment of enrichment and colocalization.
PLoS Genet. **13**, e1006646 (2017).
7.   Al-Barghouthi, B. M. *et al*. Transcriptome-wide association study and eQTL colocalization
identify potentially causal genes responsible for human bone mineral density GWAS
associations. *Elife* **11**, (2022).
8.   Li, B. & Ritchie, M. D. From GWAS to Gene: Transcriptome-Wide Association Studies
and Other Methods to Functionally Understand GWAS Discoveries. *Front. Genet*. **12**,
713230 (2021).
9.   Akiyama, M. Multi-omics study for interpretation of genome-wide association study. *J.*
Hum. Genet. **66**, 3–10 (2020).
10. Goh, K.-I. *et al*. The human disease network. *Proc. Natl. Acad. Sci. U. S. A*. **104**, 8685–
8690 (2007).
11. Calabrese, G. M. *et al*. Integrating GWAS and Co-expression Network Data Identifies Bone
Mineral Density Genes SPTBN1 and MARK3 and an Osteoblast Functional Module. *Cell*
Syst **4**, 46-59.e4 (2017).
12. Sabik, O. L., Calabrese, G. M., Taleghani, E., Ackert-Bicknell, C. L. & Farber, C. R.
Identification of a Core Module for Bone Mineral Density through the Integration of a Co-
expression Network and GWAS Data. *Cell Rep*. **32**, 108145 (2020).
13. GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat. Genet*. **45**, 580–
585 (2013).
14. GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human
tissues. *Science* **369**, 1318–1330 (2020).
15. Zhang, Q. *et al*. Genomic variants within chromosome 14q32.32 regulate bone mass
through MARK3 signaling in osteoblasts. *J. Clin. Invest*. **131**, (2021).
16. Al-Barghouthi, B. M. *et al*. Systems genetics in diversity outbred mice inform BMD GWAS
and identify determinants of bone strength. *Nat. Commun*. **12**, 3408 (2021).
17. Dillard, L. J. *et al*. Single-Cell Transcriptomics of Bone Marrow Stromal Cells in Diversity
Outbred Mice: A Model for Population-Level scRNA-Seq Studies. *J. Bone Miner. Res*. **38**,
1350–1363 (2023).
18. Abood, A. *et al*. Long-read proteogenomics to connect disease-associated sQTLs to the
protein isoform effectors of disease. *bioRxiv* 2023.03.17.531557 (2023)
doi:10.1101/2023.03.17.531557.
19. Bogue, M. A., Churchill, G. A. & Chesler, E. J. Collaborative Cross and Diversity Outbred
data resources in the Mouse Phenome Database. *Mamm. Genome* **26**, 511–520 (2015).

20.   Churchill, G. A., Gatti, D. M., Munger, S. C. & Svenson, K. L. The Diversity Outbred
mouse population. *Mamm. Genome* **23**, 713–718 (2012).

21.   Zhong, L. *et al.* Single cell transcriptomics identifies a unique adipose lineage cell
population that regulates bone marrow environment. *Elife* **9**, (2020).

22.   Timshel, P. N., Thompson, J. J. & Pers, T. H. Genetic mapping of etiologic brain cell types
for obesity. *Elife* **9**, (2020).

23.   GTEx Consortium *et al.* Genetic effects on gene expression across human tissues. *Nature*
550, 204–213 (2017).

24.   Greenfest-Allen, E., Cartailler, J.-P., Magnuson, M. A. & Stoeckert, C. J. iterativeWGCNA:
iterative refinement to improve module detection from WGCNA co-expression networks.
bioRxiv 234062 (2017) doi:10.1101/234062.

25.   Chan, C. K. F. *et al.* Identification of the Human Skeletal Stem Cell. *Cell* **175**, 43-56.e21
(2018).

26.   Mizuhashi, K. *et al.* Resting zone of the growth plate houses a unique class of skeletal stem
cells. *Nature* **563**, 254–258 (2018).

27.   Debnath, S. *et al.* Discovery of a periosteal stem cell mediating intramembranous bone
formation. *Nature* **562**, 133–139 (2018).

28.   Matsushita, Y. *et al.* A Wnt-mediated transformation of the bone marrow stromal cell
identity orchestrates skeletal regeneration. *Nat. Commun.* **11**, 332 (2020).

29.   Van den Berge, K. *et al.* Trajectory-based differential expression analysis for single-cell
sequencing data. *Nat. Commun.* **11**, 1201 (2020).

30.   Thomas, P. D. *et al.* PANTHER: Making genome-scale phylogenetics accessible to all.
Protein Sci. **31**, 8–22 (2022).

31.   Groza, T. *et al.* The International Mouse Phenotyping Consortium: comprehensive
knockout phenotyping underpinning the study of human disease. *Nucleic Acids Res.* **51**,
D1038–D1045 (2023).

32.   Enerbäck, S., Ohlsson, B. G., Samuelsson, L. & Bjursell, G. Characterization of the human
lipoprotein lipase (LPL) promoter: evidence of two cis-regulatory regions, LP-alpha and
LP-beta, of importance for the differentiation-linked induction of the LPL gene during
adipogenesis. *Mol. Cell. Biol.* **12**, 4622–4633 (1992).

33.   Federico, L. *et al.* Lipid phosphate phosphatase 3 regulates adipocyte sphingolipid
synthesis, but not developmental adipogenesis or diet-induced obesity in mice. *PLoS One*
13, e0198063 (2018).

34.   Maridas, D. E., DeMambro, V. E., Le, P. T., Mohan, S. & Rosen, C. J. IGFBP4 Is Required
for Adipogenesis and Influences the Distribution of Adipose Depots. *Endocrinology* **158**,
3488–3500 (2017).

35.   Sigg, M. A. *et al.* Evolutionary Proteomics Uncovers Ancient Associations of Cilia with
Signaling Pathways. *Dev. Cell* **43**, 744-762.e11 (2017).

36.   Kumar, R. *et al.* A cell-based GEF assay reveals new substrates for DENN domains and a
role for DENND2B in primary ciliogenesis. *Sci Adv* **8**, eabk3088 (2022).

37.   Fumoto, K. *et al.* Mark1 regulates distal airspace expansion through type I pneumocyte
flattening in lung development. *J. Cell Sci.* **132**, (2019).

38.   Miki, H., Setou, M., Kaneshiro, K. & Hirokawa, N. All kinesin superfamily protein, KIF,
genes in mouse and human. *Proc. Natl. Acad. Sci. U. S. A.* **98**, 7004–7011 (2001).

39.   Zhang, R., Roostalu, J., Surrey, T. & Nogales, E. Structural insight into TPX2-stimulated
microtubule assembly. *Elife* **6**, (2017).

40. Uusküla-Reimand, L. & Wilson, M. D. Untangling the roles of TOP2A and TOP2B in
transcription and cancer. *Sci Adv* **8**, eadd4920 (2022).
41. Trueb, B. Biology of FGFRL1, the fifth fibroblast growth factor receptor. *Cell. Mol. Life
Sci.* **68**, 951–964 (2011).
42. Hilgendorf, K. I. Primary Cilia Are Critical Regulators of White Adipose Tissue Expansion.
Front. Physiol. **12**, 769367 (2021).
43. Fazeli, P. K. *et al.* Marrow fat and bone--new perspectives. *J. Clin. Endocrinol. Metab.* **98**,
935–945 (2013).
44. Veldhuis-Vlug, A. G. & Rosen, C. J. Clinical implications of bone marrow adiposity. *J.
Intern. Med.* **283**, 121–139 (2018).
45. Haque, A., Engel, J., Teichmann, S. A. & Lönnberg, T. A practical guide to single-cell
RNA-sequencing for biomedical research and clinical applications. *Genome Med.* **9**, 75
(2017).
46. Cheng, Y., Ma, X., Yuan, L., Sun, Z. & Wang, P. Evaluating imputation methods for single-
cell RNA-seq data. *BMC Bioinformatics* **24**, 302 (2023).
47. Yu, X., Abbas-Aghababazadeh, F., Chen, Y. A. & Fridley, B. L. Statistical and
Bioinformatics Analysis of Data from Bulk and Single-Cell RNA Sequencing Experiments.
Methods Mol. Biol. **2194**, 143–175 (2021).
48. Church, D. M. *et al.* Lineage-specific biology revealed by a finished genome assembly of
the mouse. *PLoS Biol.* **7**, e1000112 (2009).
49. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29
(2021).
50. Heaton, H. *et al.* Souporcell: robust clustering of single-cell RNA-seq data by genotype
without reference genotypes. *Nat. Methods* **17**, 615–620 (2020).
51. Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-
cell RNA-seq. *Science* **352**, 189–196 (2016).
52. Neavin, D. *et al.* Single cell eQTL analysis identifies cell type-specific genetic control of
gene expression in fibroblasts and reprogrammed induced pluripotent stem cells. *Genome
Biol.* **22**, 76 (2021).
53. van der Wijst, M. G. P. *et al.* Single-cell RNA sequencing identifies celltype-specific cis-
eQTLs and co-expression QTLs. *Nat. Genet.* **50**, 493–497 (2018).
54. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140
(2010).
55. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).
56. Street, K. *et al.* Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. *BMC Genomics* **19**, 477 (2018).
57. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide
association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).
58. Cartailler, J. P. *Iterativewgcna.* (2022).
59. Tsamardinos, I., Brown, L. E. & Aliferis, C. F. The max-min hill-climbing Bayesian
network structure learning algorithm. *Mach. Learn.* **65**, 31–78 (2006).
60. Porter, M. D. & Smith, R. Network neighborhood analysis. in *2010 IEEE International
Conference on Intelligence and Security Informatics* (IEEE, 2010).
doi:10.1109/isi.2010.5484781.

61. Broman, K. W. *et al*. R/qtl2: Software for Mapping Quantitative Trait Loci with High-
Dimensional Data and Multiparent Populations. *Genetics* **211**, 495–502 (2019).
62. Phipson, B. *et al*. propeller: testing for differences in cell type proportions in single cell
data. *Bioinformatics* **38**, 4720–4726 (2022).
63. Marsh, S., Salmon, M. & Hoffman, P. *Custom Visualizations & Functions for Streamlined*
Analyses of Single Cell Sequencing. (2023). doi:10.5281/zenodo.7534950.