IMPACT OF LIVER-SPECIFIC SURVIVAL MOTOR NEURON (SMN) DEPLETION ON PERIPHERAL AND CENTRAL NERVOUS SYSTEM TISSUE PATHOLOGY

Monique Marylin Alves de Almeida[1,2], Yves De Repentigny[1], Sabrina Gagnon[1], Emma R. Sutton[1,2], Rashmi Kothary[1,2,3,4*]

[1]Regenerative Medicine Program, Ottawa Hospital Research Institute, Ottawa, K1H 8L6, Canada
[2]Centre for Neuromuscular Disease, University of Ottawa, K1H 8M5, Canada
[3]Department of Cellular and Molecular Medicine, University of Ottawa, Ottawa, K1H 8M5, Canada
[4]Department of Medicine, University of Ottawa, Ottawa, K1H 8M5, Canada
*Corresponding author: rkothary@ohri.ca

Abstract

Spinal muscular atrophy (SMA) is an inherited neuromuscular disorder stemming from deletions or mutations in the Survival Motor Neuron 1 (*SMN1*) gene, leading to decreased levels of SMN protein, and subsequent motor neuron death and muscle atrophy. While traditionally viewed as a disorder predominantly affecting motor neurons, recent research suggests the involvement of various peripheral organs in SMA pathology. Notably, the liver has emerged as a significant focus due to the observed fatty liver phenotype and dysfunction in both SMA mouse models and SMA patients. Despite these findings, it remains unclear whether intrinsic depletion of SMN protein in the liver contributes to pathology in the peripheral or central nervous systems. To address this knowledge gap, we developed a mouse model with a liver-specific depletion of SMN by utilizing an *Alb-Cre* transgene together with one *Smn²ᴮ* allele and one *Smn* exon 7 allele flanked by loxP sites. We evaluated phenotypic changes in these mice at postnatal day 19 (P19), a time when the severe model of SMA, the *Smn²ᴮ/⁻* mice, typically exhibit many symptoms of the disease. Our findings indicate that liver-specific SMN depletion does not induce motor neuron death, neuromuscular pathology or muscle atrophy, characteristics typically observed in the *Smn²ᴮ/⁻* mouse at P19. However, mild liver steatosis was observed at this time point, although no changes in liver function were detected. Notably, pancreatic alterations resembled that of *Smn²ᴮ/⁻* mice, with a decrease in insulin producing alpha-cells and an increase in glucagon producing beta-cells, accompanied with a reduction in blood glucose levels. While the mosaic pattern of the Cre-mediated excision precludes definitive conclusions regarding the contribution of liver-specific SMN depletion to overall tissue pathology, our findings highlight an intricate connection between liver function and pancreatic abnormalities in SMA, adding a nuanced layer to our understanding of the disease's complexities.

Keywords
Spinal muscular atrophy, neuromuscular disease, metabolism, pancreas

1. Introduction

Insufficient levels of Survival Motor Neuron (SMN) protein, primarily arising from mutations or deletions in the *SMN1* gene, are the root cause of spinal muscular atrophy (SMA), a hereditary neuromuscular disorder widely acknowledged as one of the leading genetic causes of infant mortality[1-3]. Characterized by the loss of motor neurons, SMA presents as a progressive weakening

of muscle strength, primarily impacting proximal muscles. If left untreated, this deterioration
culminates in respiratory failure and premature mortality[4].

The diverse phenotypes observed in SMA patients can be attributed to the genetic variations within
the *SMN* gene[1]. Humans harbor two copies of the *SMN* gene, *SMN1* and *SMN2*, which share a
nearly identical sequence except for a critical C to T substitution at position 6 of exon 7 in *SMN2*[5].
This single nucleotide change disrupts splicing of the gene, resulting in the predominant
production of an unstable protein lacking exon 7, known as *SMNΔ7*[5]. Despite this, approximately
10% of functional full-length (FL) SMN protein is still produced by the *SMN2* gene[2].
Consequently, the *SMN2* gene serves as a modifier of disease phenotype, with a higher copy
number correlating with milder disease severity[6]. In contrast, mice possess a single copy of the
Smn gene[2]. Notably, while homozygous deletion of both *SMN* genes has not been observed in
humans, the homozygous deletion of the *Smn* gene in mice induces morphological alterations and
degenerative changes in embryos post-morula stage, ultimately resulting in death at
preimplantation stage[7]. This underscores the pivotal role of the *SMN* gene in early embryonic
development[8].

Recent research has unveiled the extensive impact of SMN depletion beyond motor neurons,
implicating additional tissues such as the liver and pancreas in both SMA patients and mouse
models of SMA[9-12]. Despite major therapeutic advancements focusing on restoring SMN levels,
which have notably enhanced both lifespan and quality of life in SMA patients, complete
phenotypic rescue remains elusive[4,13,14]. Additionally, existing therapies primarily target postnatal
treatment, potentially overlooking crucial developmental changes stemming from SMN depletion.
This underscores the likelihood of a broader role for the *SMN* gene in multiple non-neuronal cell
types, necessitating further exploration.
70
Attention has been increasingly drawn to the liver as a critical area of investigation in SMA
research. Both clinical studies and preclinical mouse models have unveiled disruptions in fatty
acid metabolism, leading to an increased susceptibility among SMA patients to dyslipidemia and
liver steatosis[12,15-18]. Given its multifaceted role in regulating numerous biological functions,
including the storage and regulation of lipids, carbohydrates, amino acids, and iron, as well as the
synthesis of essential growth factors and the clearance of toxic metabolites, any disruptions in liver
function may have deleterious effects on other tissues[19].

To explore the role of SMN in the liver, prior studies have utilized a transgenic mouse model
featuring the Cre recombinase transgene driven by the α-fetoprotein promoter (*Alfp-Cre*). This
approach led to near-total depletion of SMN protein specifically in the liver, resulting in severe
impairment of liver development and late embryonic lethality[8]. In our study, to better represent the
level of SMN depletion in the liver observed in SMA and elucidate the intricate interplay between
SMN deficiency in the liver and its systemic impact in SMA-like pathology, we utilized a mouse
model employing *Alb-Cre*, where the albumin promoter drives expression of Cre recombinase in
a liver-specific manner. This model also harbors one *Smn[2B]* allele and one *Smn* exon 7 allele
flanked by loxP sites. The *Smn[2B]* allele contains a 3-base pair substitution in the mouse *Smn* gene,
resulting in aberrant splicing of exon 7[20,21]. Consequently, this genetic approach results in the
production of approximately 30% full-length functional SMN protein production in the liver.

Our findings reveal that our novel liver specific SMN depleted (herein referred to as
$Alb^{Cre/+};Smn^{2B/F7}$) mouse model induces mild liver steatosis and disrupts pancreatic function,
characterized by decreased insulin-positive and increased glucagon-positive cells, alongside
reduced blood glucose levels. Remarkably, this mouse model does not exhibit an overt SMA
phenotype, demonstrating normal lifespan and motor function. Nevertheless, our data suggests an
intricate relationship between hepatic function and pancreatic abnormalities in SMA, thereby
enhancing our comprehension of the disease's pathophysiology.
96
97   ## 2. Results
98
99   *2.1 Generation of liver-specific SMN-depleted mice*
100
To elucidate the impact of liver specific SMN depletion on both CNS and peripheral tissue
pathology, we utilized mice expressing Cre recombinase under the control of the mouse albumin
promoter, referred to as Alb^{Cre}. This strain is commonly employed for liver-specific gene knockout
using the Cre/loxP system[22]. We generated the $Smn^{2B/F7}$ genotype by crossing $Smn^{F7/F7}$ with
$Smn^{2B/2B}$ mice (Fig. 1A). The $Smn^{2B/F7}$ genotype carries one allele harboring a three nucleotide
switch in the exon splicing enhancer within exon 7 of the mouse *Smn* gene (known as Smn^{2B} allele),
mimicking the *SMN2* gene in humans[21], while the other allele features exon 7 flanked by *loxP*
sites[23]. Additionally, crosses between Alb^{Cre} and $Smn^{F7/F7}$ yielded $Alb^{Cre/+};Smn^{F7/+}$ offspring.

The liver-specific mouse line was subsequentially established through further genetic crosses
between $Alb^{Cre/+};Smn^{F7/+}$ and $Smn^{2B/F7}$, resulting in various genotypes, including $+/+;Smn^{2B/F7}$
(Fig. 1B), and $Alb^{Cre/+};Smn^{2B/F7}$ (Fig. 1C). As $+/+;Smn^{2B/F7}$ mice lack Cre recombinase, they were
used as heterozygous controls. In contrast, recombination occurs in $Alb^{Cre/+};Smn^{2B/F7}$ animals
resulting in the specific deletion of one allele of exon 7 in the liver, while the *Smn*2B allele remains
unaffected.

$Smn^{2B/-}$ mice express approximately 15% residual full-length SMN[24] (FLSMN) compared to
$Smn^{2B/+}$ heterozygous mice. Therefore, both genotypes were included in this study as additional
controls. Tissues were collected at postnatal day 19 (P19), corresponding to the symptomatic phase
of the $Smn^{2B/-}$ mouse model. At this stage, $Smn^{2B/-}$ mice show increased liver steatosis and
metabolic defects, ultimately culminating in multiple system defects and mortality, typically
occurring around P25[25].

We first assessed SMN protein levels in the liver and across CNS and peripheral tissues. Our
findings confirmed a reduction in SMN protein levels across various tissues in $Smn^{2B/-}$ mice (Fig.
2A-E). However, in $Alb^{Cre/+};Smn^{2B/F7}$, there was a ~70% decrease in FLSMN protein expression
in the liver (Fig. 2A, A') compared to littermate control counterparts ($+/+;Smn^{2B/F7}$). Conversely,
SMN protein levels in the brain (Fig. 2B, B'), muscle (Fig. 2C, C'), SC (Fig. 2D, D') and pancreas
(Fig. 2E, E') remained unaltered, indicating that deletion of *Smn* exon 7 was restricted to the liver
in $Alb^{Cre/+};Smn^{2B/F7}$ mice.

128
129   *2.2 $Alb^{Cre/+};Smn^{2B/F7}$ mice show mild yet important liver steatosis*
130

We previously showed that $Smn^{2B/-}$ mice exhibit defects in fatty acid metabolism and lipid
accumulation in the liver[12,16-18]. Additionally, SMA patients face an elevated risk of dyslipidemia
and steatosis[12]. Hence, we explored whether the intrinsic decrease in SMN levels restricted to the
liver would lead to increased lipid accumulation. Histology from $Alb^{Cre/+};Smn^{2B/F7}$ mice (Fig. 3A)
qualitatively indicates some level of hepatic microvesicular steatosis and lipid accumulation.
However, there was a noticeable variation in cell morphology and Oil Red-O retention among
hepatocytes, suggesting incomplete recombination.

Consistent with the histological findings, $Alb^{Cre/+};Smn^{2B/F7}$ mice showed variable triglycerides
levels (n1=45.95, n2=10.10; n3=75.72; n4=21.91 µg/mg, mean±SEM 38.44±14.48) (Fig. 3B), and
yet higher than those observed in $+/+;Smn^{2B/F7}$ controls (n1=2.60, n2=0.15; n3=0.66; n4=1.44
µg/mg, mean±SEM 1.213±0.5332), although this difference did not reach statistical significance
(p=0.0824). As expected, $Smn^{2B/-}$ controls exhibited increased lipid accumulation and triglyceride
levels. Taken together, the data suggest an intrinsic role of SMN in regulating fat liver metabolism.

2.3 Liver-specific SMN depletion does not perturb liver homeostatic function

We next sought to identify whether $Alb^{Cre/+};Smn^{2B/F7}$ mice display disturbed liver homeostatic
function. Specifically, we evaluated parameters associated with iron metabolism (Fig. 4A,B),
endoplasmic reticulum (ER) stress (Fig. 4C), and insulin-like growth factor (IGF-1) (Fig. 4D)
levels, all of which have previously been identified as altered in $Smn^{2B/-}$ mice[18] and were further
confirmed in our study. Our analysis revealed no significant differences in the protein levels of
heme oxygenase, transferrin, P62, and IGF-1 between the $Alb^{Cre/+};Smn^{2B/F7}$ and $+/+;Smn^{2B/F7}$
control groups. These findings suggest that liver-specific SMN depletion may not play a significant
role in these pathways or that the mosaic nature of Cre-mediated excision resulted in a partial effect
that was difficult to pick up at this time point.

2.4 Liver-specific SMN depleted mice display pancreatic defects

We have demonstrated pancreatic and glucose abnormalities in the $Smn^{2B/-}$ mouse model of SMA
and type I SMA patients[11,26]. Therefore, we conducted immunohistochemistry to investigate the
cellular profile of the pancreas, aiming to ascertain whether the targeted depletion of SMN in the
liver contributes to pancreatic dysfunction. Surprisingly, double-labeling of insulin-producing β
cells and glucagon-producing α cells revealed a significant decrease in the proportion of β cells
and a notable increase in α cells in the pancreas of $Alb^{Cre/+};Smn^{2B/F7}$ mice compared to
$+/+;Smn^{2B/F7}$ and $Smn^{2B/+}$ heterozygous controls (Fig. 5A-C). This loss of β cells and rise in α cells
could suggest a functional impairment in glucose metabolism. Consequently, we evaluated non-
fasting glucose levels in these mice (Fig. 5D), revealing reduced blood glucose levels in
$Alb^{Cre/+};Smn^{2B/F7}$ mice relative to $+/+;Smn^{2B/F7}$ and $Smn^{2B/+}$. However, blood glucose levels
remained higher than those observed in $Smn^{2B/-}$ mice. Overall, these findings indicate imbalances
in α and β cell fate within pancreatic islets in $Alb^{Cre/+};Smn^{2B/F7}$ mice, resembling those observed in
$Smn^{2B/-}$ mice. Furthermore, the maintenance of SMN protein expression at levels comparable to
those of heterozygous controls in the pancreas (Fig. 2E,E') suggests that selective SMN depletion
in the liver may be implicated in the dysfunction of the pancreas-liver axis.

2.5 Motor neurons, neuromuscular junction (NMJ) or motor fibers are unaffected in liver specific
SMN depleted mice

The hallmark features of SMA include the degeneration of lower spinal motor neurons,
neuromuscular junction (NMJ) pathology, and muscle atrophy[10,25,27-33]. While the exact
mechanisms linking these changes in the motor unit to the clinical manifestations of SMA remain
unclear, diminished levels of SMN protein are recognized as a pivotal factor[1,2,25,33-35]. Thus, we
investigated the impact of liver-specific SMN depletion on these SMA-like features. Our findings
show no significant differences in motor neuron cell body counts (Fig. 6A,B), neurofilament
accumulation at NMJs (Fig. 7A,B), NMJ endplate occupancy (Fig. 7A,C), muscle fiber size (Fig.
8A,B), or muscle fiber frequency distribution (Fig. 8C) among $Alb^{Cre/+};Smn^{2B/F7}$ and $+/+;Smn^{2B/F7}$
and $Smn^{2B/+}$ heterozygous controls. However, as anticipated, $Smn^{2B/-}$ mice exhibited motor neuron
loss, NMJ defects, and muscle atrophy.

2.6 Liver-specific SMN depleted mice have normal survival rate and motor function

We also conducted a comprehensive assessment of survival, weight, and motor function in
$Alb^{Cre/+};Smn^{2B/F7}$ and $+/+;Smn^{2B/F7}$ mice over a 60-day period (Fig. 9A). While the $Smn^{2B/-}$ mice
typically have a mean survival of 25 days[25], both $Alb^{Cre/+};Smn^{2B/F7}$ and $+/+;Smn^{2B/F7}$ mice in our
study survived up to 60 days (Fig. 9B), after which they were euthanized for tissue collection.
Moreover, throughout the observation period, mice from both groups exhibited similar weight gain
trends (Fig. 9C).

To further evaluate motor function at different stages, we implemented several age-specific
behavioral tests commonly used to assess motor function[36] (Fig. 9D-F). We previously showed
$Smn^{2B/-}$ mice have reduced motor function scores, as seen through increased time to right,
prolonged balancing time and diminished muscle strength[26]. Here, we found no alterations in
$Alb^{Cre/+};Smn^{2B/F7}$ compared to $+/+;Smn^{2B/F7}$ mice in time to right, as assessed by the righting reflex
(Fig. 9D), nor in distal muscle strength measured through the inverted mesh grip test (Fig. 9E), or
motor balance and coordination evaluated by the pen test (Fig. 9F). Overall, these findings
collectively demonstrate that liver-specific SMN depletion does not impact survival, weight gain,
or motor function.

202
203

3. Discussion

The liver is pivotal in regulating numerous biological functions in the body. These encompass,
among others, the storage and balance of lipids, carbohydrates, and iron, as well as the synthesis
of vital growth factors and the clearance of non-essential or harmful metabolites[19]. Given its
multifaceted role, any alterations in liver function can potentially have cascading effects on other
organs within the body. Hence, aside from elucidating the impact of liver specific SMN depletion
on hepatic function, it is imperative to explore its effects on other tissues, as alterations in these
tissues have been documented in various mouse models of SMA[11,18,34,37].

Previous work on conditional knockout of *Smn* in the livers of mice resulted in severe impairment
of liver development characterized by iron overload and subsequent liver atrophy, ultimately
leading to late embryonic lethality[8]. In our study, we generated a novel mouse model where instead

of a complete knockout of *Smn* in the liver, we created a scenario where SMN protein was depleted
to 30% of full-length SMN production specifically within the liver, while SMN protein levels in
other tissues remain comparable to those of heterozygous controls. Notably, this mouse line did
not exhibit any signs of developmental delays, liver atrophy, or reduced lifespan. Nonetheless,
within 19 days after birth, *Alb*[Cre/+]*;Smn*[2B/F7] mice showed a mild yet rapid onset of fatty liver
disease and trends towards an increase in lipid levels when maintained on a normal chow diet,
underscoring the importance of SMN in lipid metabolism.

Mild liver steatosis may be accompanied by disruptions in crucial liver function markers, notably
those implicated in iron metabolism, autophagy, and growth factor regulation, such as IGF-1[18].
Accordingly, we initially investigated the expression of HO-1, an enzyme responsible for heme
degradation into iron, carbon monoxide, and biliverdin; and transferrin, a liver-synthesized
glycoprotein involved in iron transport across tissues (reviewed in[38]). Prior studies have
documented alterations in these proteins in *Smn*[2B/-] mouse livers[18]. While our current analysis did
not reveal significant differences in *Alb*[Cre/+]*;Smn*[2B/F7] mice, the observed upward trend in
transferrin and HO levels suggests ongoing changes in iron metabolism, which may not be fully
manifested at P19. In addition, incomplete Cre-mediated excision of *Smn*[F7] allele may be hindering
a comprehensive understanding of SMN's role in liver function.

Furthermore, no alterations were observed for P62 levels, a multifunctional scaffold protein
implicated in autophagy, formation of hepatic inclusion bodies, and hepatocyte cell death
(reviewed in[39]) despite its increased presence in mouse models of SMA[18]. Similarly, there were no
changes detected in IGF-1 protein levels, whose deficiency has been linked to liver pathology and
non-alcoholic fatty liver disease (NAFLD)[40,41], and its decrease is evident in *Smn*[2B/-] mice[18].

Altogether, our findings underscore the necessity of conducting further investigations at later time
points to unveil potential modifications in other pathways and their repercussions on liver
physiology. Additionally, the variability in the data may be due to differences in Cre recombination
efficiency from mouse to mouse due to inconsistent Cre recombination among littermates, or the
absence of Cre expression in certain cells leading to recombination failure[42].
243
Previously, our research unveiled metabolic abnormalities in SMA-like mice, including glucose
intolerance, insulin hypersensitivity, and hyperglucagonemia, accompanied by changes in
pancreatic islet composition characterized by an increase in glucagon-producing α-cells at the
expense of insulin-producing β-cells[11]. Histopathological examination of pancreatic tissue from
infants and children with SMA type I corroborated these findings, revealing a predominant
presence of glucagon-producing α-cells within pancreatic islets[11]. Furthermore, one-year-old
Smn[+/-]mice producing approximately 50% of SMN protein, when exposed to a high-fat diet,
exhibited dysregulation in the proportion of glucagon-producing α-cells within pancreatic islets
and heightened hepatic insulin and glucagon sensitivity[9]. Our study uncovers a novel intrinsic role
of liver-specific SMN in pancreatic function, distinct from IGF-1 signaling, pancreatic SMN levels
(which remained stable), or canonical SMA pathology.
255
In addition, while the hallmark pathological features of SMA include motor neuron loss, NMJ
pathology, muscle atrophy, and consequent motor function alterations[2,30,33-35], these were not
evident in the *Alb*[Cre/+]*;Smn*[2B/F7] mouse model at P19. However, given that glucose metabolism
defects are observed in various neurodegenerative and neuromuscular disorders, subsequent time

points are crucial to discern whether additional alterations arise as the mice age. Although hepatic
and pancreatic abnormalities likely occur independently of disease onset, given that SMN
depletion in motor neurons and muscle alone[23,43,44] is sufficient to induce a SMA-like phenotype,
the potential influence of these anomalies on SMA progression warrants careful consideration.
264
265 Overall, our findings suggest that alterations in SMN production solely within the liver may suffice
266 to induce pathological hepatic and pancreatic function, shedding light on a novel role of SMN in
267 liver physiology.
268
4. **Material and Methods**

4.1 Mouse models

The *Smn^{2B}* mouse, which harbors the *Smn^{2B}* mutation resulting from the substitution of three
nucleotides within the exon splicing enhancer of exon 7, was developed and maintained on a
C57BL/6 background in our laboratory[21]. The FVB.129(B6)-*Smn1^{tm1Jme}* (*Smn$^{F7/F7}$*) and B6.Cg-
Speer6-ps1$^{Tg(Alb-Cre)21Mgn}$/J (*Alb^{Cre}*) mice were acquired from The Jackson Laboratory. Liver-
specific SMN-depleted mice were obtained through successive crosses between *Alb$^{Cre/+}$;Smn$^{F7/+}$*
and *Smn$^{2B/F7}$* mice (see Fig. 1). Line validation was conducted via genotyping of DNA extracted
from mouse ear biopsies using PCR (see Table 1 for list of primers used). Both male and female
mice were included in the studies. All mice were bred and housed in the University of Ottawa
Animal Facility under protocol OHRI-3343, adhering to guidelines established by the Canadian
Council on Animal Care. Additionally, we employed *Smn$^{2B/-}$* mice (a severe SMA mouse model)
and asymptomatic heterozygous *Smn$^{2B/+}$* mice as controls in our experiments.

Table 1. List of primers used for genotyping
283

NAME	SEQUENCE (5'-3')
Alb-Cre	TGCAAACATCACATGCACAC
	TTGGCCCCTTACCATAACTG
	GAAGCAGAAGCTTAGGAAGATGG
SmnF7	AGAAGGAAAGTGCTCACATACAAATT
	TGTCTATAATCCTCATGCTATGGAG
Smn2B	GGTTTCAGACAAAATAAAAAAGAATTTAAG
	TTTGGCAGACTTTAGCAGGGC

284
285
286 *4.2 Tissue harvesting and processing*
287
Following euthanasia, liver, and *tibialis anterior* (TA) muscles underwent fixation in a 1:10
dilution of buffered formalin for 48 h at 4°C, followed by transfer to 70% ethanol at the same
temperature until processing. The pancreas was fixed in 4% paraformaldehyde (PFA) for 48 h at
4°C, then similarly moved to 70% ethanol for processing. For the lumbar spinal cord (SC) and
liver (for Oil Red-O staining), fixation was performed overnight at 4°C in 4% PFA, followed by
immersion in 30% sucrose in PBS. Subsequently, the liver and SC were flash-frozen in optimal
cutting temperature (OCT) embedding medium. Cryosections of the lumbar spinal cord were cut
at a thickness of 16 μm, while liver sections were cut at 10 μm and stored at -80°C until
immunohistochemical analysis.

The abdominal musculature was dissected and fixed in 4% PFA for 15 min at room temperature,
followed by 3 washes in 1X PBS, after which the transversus abdominis (TVA) muscle was
dissected from the abdominal musculature. TA, liver, and pancreas samples were handled at the
University of Ottawa Department of Pathology and Laboratory Medicine. Paraffin blocks
containing tissues were sectioned at 4 μm thickness using a microtome. Sections of TA and liver
were stained with hematoxylin and eosin (H&E) using an XL CV5030 autostainer from Leica.
Liver cryosections were stained with Oil Red-O and counterstained with hematoxylin using
established protocols. Images of H&E and Oil Red-O-stained samples were scanned with a
MIRAX MIDI digital slide scanner manufactured by Zeiss. Image acquisition was performed using
Panoramic Viewer 1.15.4 software.

4.3 Immunoblotting

Total protein was extracted by homogenizing flash-frozen samples of spinal cord, brain, liver,
pancreas and hindlimb muscles using RIPA lysis buffer (Cell Signaling), supplemented with
phenylmethylsulfonyl fluoride (PMSF, Cell Signaling). The concentrations of the extracted
proteins were determined using the Pierce BCA Protein Assay kit (ThermoFisher).

For liver, brain, and spinal cord tissues, 20 μg of protein per sample were separated by 10% sodium
dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE), while 30 μg were used for
pancreas and muscle samples. Following electrophoresis, the proteins were transferred onto a
PVDF membrane (Immobilon-P, Millipore) and blocked for 30 min at room temperature using
intercept blocking buffer (LI-COR). Subsequently, the membranes were incubated overnight at
4°C with primary antibodies.

After incubation, the membranes were rinsed with tris-buffered saline containing 0.1% Tween-20
(TBS-T) and then incubated for 1 h at room temperature with IRDye 680R or 800CW (LI-COR)
secondary antibody. Fluorescent signals were detected using the Odyssey Infrared Imaging System
(LI-COR). Total raw values were normalized to α-tubulin (housekeeping protein) obtained from
the same blot.

4.4 Immunohistochemistry

4.4.1 Pancreas

Pancreatic sections underwent deparaffinization using three washes of Histo-Clear (National
Diagnostics, Atlanta, GA) for 5 min each, followed by two washes of a 50/50 mixture of absolute
ethanol and Histo-Clear for 3 min each. Subsequently, the slides were gradually rehydrated in two
washes of 100% (v/v) ethanol (3 min each), followed by one wash of 95% (v/v), 70% (v/v), 50%
(v/v), and 0% (v/v) ethanol for 3 min each. The slides were then incubated in 0.5% Triton-X-100
(Millipore Sigma, Burlington, MA) in PBS for 5 min, followed by three washes with PBS.
Afterward, the slides were blocked in a solution containing 20% goat serum and 0.3% Triton-X-
100 in TBS for 2 h.

Primary antibodies targeting insulin and glucagon (Table 2) were applied to the slides in a solution
containing 2% goat serum and 0.3% Triton-X-100 in TBS overnight at 4°C. The slides were then

washed three times with PBS and subsequently incubated with secondary antibodies in a solution
containing 2% goat serum and 0.3% Triton-X-100 for 1 h. Following this, the solution was
removed, and DAPI (1:1,000) in PBS was added for 5 min at room temperature. The slides were
washed three times with PBS and mounted in Fluoromount-G™ Mounting Medium (Invitrogen).

Images were taken using an Axio Imager M2 microscope (Zeiss) and captured with a 40X
objective. The number of glucagon and insulin positive cells per islet was counted for each mouse.

347
4.4.2 Spinal cord (SC)

Lumbar spinal cord sections were prepared for choline acetyltransferase (ChAT) staining of motor
neurons. The slides were air-dried at room temperature for 30 min and then rinsed in TBS-T for 5
min. Samples were permeabilized in 0.3% Triton X-100 in PBS for 30 min, followed by blocking
in 1x Power Block (BioGenex, Fremont, CA) for 20 min at room temperature. Subsequently, the
samples were incubated with a goat anti-ChAT (details in Table 2) antibody at a dilution of 1:100
in 1% BSA and 0.3% Triton X-100 in PBS for 3 nights at 4°C.

355
After the initial antibody incubation, slides were washed twice for 10 min with TBS-T at room
temperature. Following the washes, samples were then incubated with Alexa Fluor 555 donkey
anti-goat IgG at a dilution of 1:200 in 10% donkey serum and 0.3% Triton X-100 in PBS for 2 h
at room temperature in a humid chamber. Nuclei were counterstained with 4',6-diamidino-2-
phenylindole (DAPI) at a dilution of 1:1000 in PBS for 5 min.

Finally, the slides were rinsed 3X for 10 min with TBS-T and mounted with Fluoromount-G™
Mounting Medium (Invitrogen). SC were examined under fluorescence using an Axio Imager M2
microscope (Zeiss) and imaged at 20X magnification for quantification, and at 63x for
representative pictures. The number of ChAT-positive motor neurons with a diameter of ≥20 μm
per ventral horn was recorded for 7-10 different sections per animal, each separated by at least 100
μm to prevent re-counting of the same motor neuron. An average number of motor neuron cell
bodies per section was determined.

4.4.3 Neuromuscular Junction (NMJ)
369
To evaluate neuromuscular innervation, we performed immunohistochemistry using established
protocols[26]. After euthanasia, the TVA muscles were promptly dissected and fixed in 4%
paraformaldehyde (Electron Microscopy Science) in PBS for 15 min. Post-synaptic acetylcholine
receptors (AChRs) were labeled for 10 min with alpha-bungarotoxin (aBTX) conjugated to Alexa
Fluor 488. Subsequently, the muscles were blocked in 4% bovine serum albumin (BSA) and 1%
Triton-X-100 in PBS for 30 min.

376
377 Following blocking, the muscles underwent overnight incubation with primary antibodies for
378 neurofilament and synaptic vesicle protein 2 (refer to Table 2). Visualization of the labeled
379 structures was achieved using DyLight-conjugated secondary antibodies. The muscles were then
380 whole-mounted in Dako Fluorescent mounting media and imaged with a 20x objective on a Zeiss
381 Axio Imager M1 microscope.
382

383 A minimum of three fields of view (FOV) were quantified per muscle. For each FOV, the
384 percentage of fully occupied endplates was noted. Fully occupied endplates were defined as motor
385 endplates completely covered by the presynaptic terminal labeled with synaptic vesicle protein 2
386 (SV2) and neurofilament (NF).
387
388 **Table 2. List of antibodies used.**
389

Application	Antibody	Species	Dilution	Company (Catalog #)
Western Blot	SMN	Mouse	1:2 500	BD Transduction (610647)
Western Blot	Alpha-tubulin	Rabbit	1:10 000	Abcam (ab4074)
Western Blot	Alpha-tubulin	Mouse	1:10 000	Sigma-Aldrich (CP06)
Western Blot	Heme oxygenase-1 (HO)	Rabbit	1:10 000	Abcam (ab68477)
Western Blot	P62	Mouse	1:1 000	Abcam (ab56416)
Western Blot	Transferrin	Rabbit	1:1 000	Abcam (ab82411)
Western Blot	Goat anti-mouse IR Dye 800 CW		1:5 000	LI-COR (925-32210)
Western Blot	Goat anti-mouse IR Dye 680 R		1:5 000	LI-COR (926-68070)
Western Blot	Goat anti-rabbit IR Dye 800 CW		1:5 000	LI-COR (925-32211)
Western Blot	Goat anti-rabbit IR Dye 680 R		1:5 000	LI-COR (926-68071)
IHC	ChAT	Goat	1:100	Millipore (AB144P)
IHC	Donkey anti-goat Alexa Fluor 555		1:200	Invitrogen (A21432)
IHC	TRITC conjugated bungarotoxin	N/A	1:1 000	Invitrogen (T1175)
IHC	Neurofilament (NF-M)	Mouse	1:100	(Developmental Studies Hybridoma Bank, P12839)
IHC	Synaptic vesicle glycoprotein 2A (SV2A)	Mouse	1:250	(Developmental Studies Hybridoma Bank, Q7L0J3)
IHC	Goat anti-mouse Alexa Fluor 488		1:250	Invitrogen (A11001)
IHC	Glucagon	Mouse	1:200	Abcam (ab10988)
IHC	Insulin	Rabbit	1:50	Abcam (ab181547)

| IHC | Goat anti-mouse Alexa Fluor 555 | 1:500 | Invitrogen (A21422) |
| IHC | Goat anti-rabbit Alexa Fluor 488 | 1:500 | Invitrogen (A11034) |

4.5 Muscle fiber analysis

Muscle fiber analysis was performed using ImageJ software (version 2.9.0/1.53t). A total of one hundred fibers were evaluated per animal, covering various regions of the muscle section to ensure good representation. The area of each fiber was measured in square micrometers (μm^2) to compute both the average and distribution of fiber sizes for each animal.

4.6 Glucose levels

Non-fasting blood glucose concentrations were assessed using a OneTouch Ultra2 Blood Glucose Test Strips (LifeScan Europe GmbH, Zug, Switzerland) immediately following blood collection. Approximately 2 μL of blood was applied to the test strip for glucose concentration measurement.

4.7 ELISA for Insulin-like Growth Factor I (IGF-1)

The concentration of IGF-1 in the liver was quantified using the Mouse/Rat IGF-I/IGF1 Quantikine ELISA Kit (MG100, R&D Systems, Inc., Minneapolis, MN, USA). Liver protein lysates were diluted at a ratio of 1:6 in calibrator diluent and the assay was performed following the manufacturer's instructions.

4.8 Triglycerides quantification

Liver tissues from P19 mice were promptly dissected and flash frozen. Subsequently, triglyceride analysis was conducted at the Vanderbilt Mouse Metabolic Phenotyping Center, employing established protocols previously utilized by our team[18].

4.9 Motor function, weight, and survival assessments

Motor function, weight, and survival were assessed according to established protocols[36]. Spinal reflexes were evaluated using the righting reflex test, while motor balance and coordination were evaluated with the pen test. Additionally, muscle strength was measured using the inverted mesh grip test (see Fig. 9A). The righting reflex test was conducted from P7 to P13, the inverted mesh grip test from P13 to P25, and the pen test from P19 to P25. Assessments were performed every two days. A maximum duration of 30 sec for the pen test and 60 sec for the inverted mesh grip represented the established threshold. Three consecutive measurements were obtained, and their average was recorded for subsequent analysis. Weight was measured every two days starting at day 7 until day 30, and then weekly until day 60. Animal welfare was monitored throughout the 60-day period during weight measurements.

4.10 Statistical analysis

Survival data were visualized using Kaplan-Meier survival curves, and intergroup differences were evaluated using the Mantel-Cox test. Brown-Forsythe and Welch ANOVA were employed to compare three or more sets of unpaired measurements, given the mosaic-like outcome observed in the $Alb^{Cre/+};Smn^{2B/F7}$ model, where assumptions of equal variances were not tenable. Two-way ANOVA, followed by Šídák's method for multiple comparisons, was utilized to assess the influence of two factors on a response. Statistical analyses were conducted using GraphPad Prism V.10.2.1 (GraphPad Prism Software, San Diego, CA), with significance set at $P < 0.05$. Data were presented as mean ± standard error of the mean. Sample sizes (n) are provided in figure legends, indicating the number of biological replicates analyzed, with each data point corresponding to an individual mouse from at least two different litters. Detailed statistical information is outlined in the respective figure legends, where statistical significance is indicated by *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Data sharing

All authors had access to the study data and reviewed and approved the final manuscript. All data associated with this study are available in the main text or supplementary materials. Raw data can be provided upon request.

Acknowledgements

This work was supported by Muscular Dystrophy Association (USA) [grant number 963652 to R.K.]; the Canadian Institutes of Health Research [grant number PJT-186300 to R.K.]; and the University of Ottawa Brain and Mind Institute TRIMS Award to MMAA.

Declaration of interests
The authors disclose no conflicts of interest. The funders mentioned above were not involved in the study design, data collection and analysis, decision to publish, or manuscript preparation.

References

Lefebvre, S. *et al.* Identification and characterization of a spinal muscular atrophy-determining gene. *Cell* **80**, 155-165 (1995).
Lefebvre, S. *et al.* Correlation between severity and SMN protein level in spinal muscular atrophy. *Nat Genet* **16**, 265-269 (1997).
Crawford, T. O. & Pardo, C. A. The Neurobiology of Childhood Spinal Muscular Atrophy. *Neurobiology of Disease* **3**, 97-110 (1996).
Finkel, R. S. *et al.* Treatment of infantile-onset spinal muscular atrophy with nusinersen: a phase 2, open-label, dose-escalation study. *Lancet* **388**, 3017-3026 (2016). https://doi.org:10.1016/S0140-6736(16)31408-8
Lorson, C. L., Hahnen, E., Androphy, E. J. & Wirth, B. A single nucleotide in the SMN gene regulates splicing and is responsible for spinal muscular atrophy. *Proc Natl Acad Sci U S A* **96**, 6307-6311 (1999).
Feldkotter, M., Schwarzer, V., Wirth, R., Wienker, T. F. & Wirth, B. Quantitative analyses of SMN1 and SMN2 based on real-time lightCycler PCR: fast and highly

476		reliable carrier testing and prediction of severity of spinal muscular atrophy. *Am J Hum*
477		*Genet* **70**, 358-368 (2002). https://doi.org:10.1086/338627
478	7	Schrank, B. *et al.* Inactivation of the survival motor neuron gene, a candidate gene for
479		human spinal muscular atrophy, leads to massive cell death in early mouse embryos.
480		*Proc Natl Acad Sci U S A* **94**, 9920-9925 (1997).
481	8	Vitte, J. M. *et al.* Deletion of murine Smn exon 7 directed to liver leads to severe defect
482		of liver development associated with iron overload. *Am J Pathol* **165**, 1731-1741 (2004).
483		https://doi.org:10.1016/S0002-9440(10)63428-1
484	9	Bowerman, M. *et al.* Defects in pancreatic development and glucose metabolism in
485		SMN-depleted mice independent of canonical spinal muscular atrophy neuromuscular
486		pathology. *Hum Mol Genet* **23**, 3432-3444 (2014). https://doi.org:10.1093/hmg/ddu052
487	10	Bowerman, M., Murray, L. M., Beauvais, A., Pinheiro, B. & Kothary, R. A critical smn
488		threshold in mice dictates onset of an intermediate spinal muscular atrophy phenotype
489		associated with a distinct neuromuscular junction pathology. *Neuromuscul Disord* **22**,
490		263-276 (2012). https://doi.org:S0960-8966(11)01345-9 [pii] 10.1016/j.nmd.2011.09.007
491	11	Bowerman, M. *et al.* Glucose metabolism and pancreatic defects in spinal muscular
492		atrophy. *Ann Neurol* **72**, 256-268 (2012).
493	12	Deguise, M. O. *et al.* Abnormal fatty acid metabolism is a core component of spinal
494		muscular atrophy. *Ann Clin Transl Neurol* **6**, 1519-1532 (2019).
495		https://doi.org:10.1002/acn3.50855
496	13	Chiriboga, C. A. Nusinersen for the treatment of spinal muscular atrophy. *Expert Rev*
497		*Neurother* **17**, 955-962 (2017). https://doi.org:10.1080/14737175.2017.1364159
498	14	McMillan, H. J. *et al.* Onasemnogene abeparvovec for the treatment of spinal muscular
499		atrophy. *Expert Opin Biol Ther* **22**, 1075-1090 (2022).
500		https://doi.org:10.1080/14712598.2022.2066471
501	15	Crawford, T. O., Sladky, J. T., Hurko, O., Besner-Johnston, A. & Kelley, R. I. Abnormal
502		fatty acid metabolism in childhood spinal muscular atrophy. *Ann Neurol* **45**, 337-343
503		(1999).
504	16	Deguise, M. O., Chehade, L. & Kothary, R. Metabolic Dysfunction in Spinal Muscular
505		Atrophy. *Int J Mol Sci* **22** (2021). https://doi.org:10.3390/ijms22115913
506	17	Deguise, M. O., Chehade, L., Tierney, A., Beauvais, A. & Kothary, R. Low fat diets
507		increase survival of a mouse model of spinal muscular atrophy. *Ann Clin Transl Neurol*
508		**6**, 2340-2346 (2019). https://doi.org:10.1002/acn3.50920
509	18	Deguise, M. O. *et al.* SMN Depleted Mice Offer a Robust and Rapid Onset Model of
510		Nonalcoholic Fatty Liver Disease. *Cell Mol Gastroenterol Hepatol* **12**, 354-377 e353
511		(2021). https://doi.org:10.1016/j.jcmgh.2021.01.019
512	19	Trefts, E., Gannon, M. & Wasserman, D. H. The liver. *Curr Biol* **27**, R1147-R1151
513		(2017). https://doi.org:10.1016/j.cub.2017.09.019
514	20	DiDonato, C. J. *et al.* Regulation of murine survival motor neuron (Smn) protein levels
515		by modifying Smn exon 7 splicing. *Hum Mol Genet* **10**, 2727-2736. (2001).
516	21	Hammond, S. M. *et al.* Mouse survival motor neuron alleles that mimic SMN2 splicing
517		and are inducible rescue embryonic lethality early in development but not late. *PLoS One*
518		**5**, e15887 (2010). https://doi.org:10.1371/journal.pone.0015887
519	22	Postic, C. *et al.* Dual roles for glucokinase in glucose homeostasis as determined by liver
520		and pancreatic beta cell-specific gene knock-outs using Cre recombinase. *J Biol Chem*
521		**274**, 305-315 (1999). https://doi.org:10.1074/jbc.274.1.305

522	23	Frugier, T. *et al.* Nuclear targeting defect of SMN lacking the C-terminus in a mouse
523		model of spinal muscular atrophy. *Hum Mol Genet* **9**, 849-858 (2000).
524	24	Eshraghi, M., McFall, E., Gibeault, S. & Kothary, R. Effect of genetic background on the
525		phenotype of the Smn2B/- mouse model of spinal muscular atrophy. *Hum Mol Genet* **25**,
526		4494-4506 (2016). https://doi.org:10.1093/hmg/ddw278
527	25	Reilly, A., Yaworski, R., Beauvais, A., Schneider, B. L. & Kothary, R. Long term
528		peripheral AAV9-SMN gene therapy promotes survival in a mouse model of spinal
529		muscular atrophy. *Hum Mol Genet* **33**, 510-519 (2024).
530		https://doi.org:10.1093/hmg/ddad202
531	26	Reilly, A. *et al.* Central and peripheral delivered AAV9-SMN are both efficient but target
532		different pathomechanisms in a mouse model of spinal muscular atrophy. *Gene Ther* **29**,
533		544-554 (2022). https://doi.org:10.1038/s41434-022-00338-1
534	27	Deguise, M. O. *et al.* Differential induction of muscle atrophy pathways in two mouse
535		models of spinal muscular atrophy. *Sci Rep* **6**, 28846 (2016).
536		https://doi.org:10.1038/srep28846
537	28	Deguise, M. O. *et al.* Motor transmission defects with sex differences in a new mouse
538		model of mild spinal muscular atrophy. *EBioMedicine* **55**, 102750 (2020).
539		https://doi.org:10.1016/j.ebiom.2020.102750
540	29	Gavrilina, T. O. *et al.* Neuronal SMN expression corrects spinal muscular atrophy in
541		severe SMA mice while muscle-specific SMN expression has no phenotypic effect. *Hum*
542		*Mol Genet* **17**, 1063-1075 (2008).
543	30	Murray, L. M., Beauvais, A., Bhanot, K. & Kothary, R. Defects in neuromuscular
544		junction remodelling in the *Smn2B/-* mouse model of spinal muscular atrophy.
545		*Neurobiology of Disease* **49C**, 57-67 (2012).
546	31	Murray, L. M., Beauvais, A., Gibeault, S., Courtney, N. L. & Kothary, R. Transcriptional
547		profiling of differentially vulnerable motor neurons at pre-symptomatic stage in the Smn
548		(2b/-) mouse model of spinal muscular atrophy. *Acta Neuropathol Commun* **3**, 55 (2015).
549		https://doi.org:10.1186/s40478-015-0231-1
550	32	Shababi, M., Lorson, C. L. & Rudnik-Schoneborn, S. S. Spinal muscular atrophy: a
551		motor neuron disorder or a multi-organ disease? *J Anat* **224**, 15-28 (2014).
552		https://doi.org:10.1111/joa.12083
553	33	Lorson, C. L., Rindt, H. & Shababi, M. Spinal muscular atrophy: mechanisms and
554		therapeutic strategies. *Hum Mol Genet* **19**, R111-118 (2010).
555		https://doi.org:10.1093/hmg/ddq147
556	34	Hua, Y. *et al.* Peripheral SMN restoration is essential for long-term rescue of a severe
557		spinal muscular atrophy mouse model. *Nature* **478**, 123-126 (2011).
558		https://doi.org:nature10485 [pii] 10.1038/nature10485
559	35	Mercuri, E., Sumner, C. J., Muntoni, F., Darras, B. T. & Finkel, R. S. Spinal muscular
560		atrophy. *Nat Rev Dis Primers* **8**, 52 (2022). https://doi.org:10.1038/s41572-022-00380-8
561	36	Chehade, L. *et al.* Suppression of the necroptotic cell death pathways improves survival
562		in Smn (2B/-) mice. *Front Cell Neurosci* **16**, 972029 (2022).
563		https://doi.org:10.3389/fncel.2022.972029
564	37	Hua, Y. *et al.* Motor neuron cell-nonautonomous rescue of spinal muscular atrophy
565		phenotypes in mild and severe transgenic mouse models. *Genes Dev* **29**, 288-297 (2015).
566		https://doi.org:10.1101/gad.256644.114

38 Muckenthaler, M. U., Rivella, S., Hentze, M. W. & Galy, B. A Red Carpet for Iron
Metabolism. *Cell* **168**, 344-361 (2017). https://doi.org:10.1016/j.cell.2016.12.034
39 Manley, S., Williams, J. A. & Ding, W. X. Role of p62/SQSTM1 in liver physiology and
pathogenesis. *Exp Biol Med (Maywood)* **238**, 525-538 (2013).
https://doi.org:10.1177/1535370213489446
40 Hribal, M. L. *et al.* Insulin-like growth factor-I, inflammatory proteins, and fibrosis in
subjects with nonalcoholic fatty liver disease. *J Clin Endocrinol Metab* **98**, E304-308
(2013). https://doi.org:10.1210/jc.2012-3290
41 Ichikawa, T. *et al.* Role of growth hormone, insulin-like growth factor 1 and insulin-like
growth factor-binding protein 3 in development of non-alcoholic fatty liver disease.
Hepatol Int **1**, 287-294 (2007). https://doi.org:10.1007/s12072-007-9007-4
42 Heffner, C. S. *et al.* Supporting conditional mouse mutagenesis with a comprehensive cre
characterization resource. *Nat Commun* **3**, 1218 (2012).
https://doi.org:10.1038/ncomms2186
43 Cifuentes-Diaz, C. *et al.* Deletion of murine SMN exon 7 directed to skeletal muscle
leads to severe muscular dystrophy. *J Cell Biol* **152**, 1107-1114. (2001).
44 Cifuentes-Diaz, C. *et al.* Neurofilament accumulation at the motor endplate and lack of
axonal sprouting in a spinal muscular atrophy mouse model. *Hum Mol Genet* **11**, 1439-
1447. (2002).

Figure subtitles

1. **Generation of liver specific SMN depleted mice.** A. Schematic representation of the breeding scheme to generate liver specific SMN depletion. In these mice, the *Smn* gene carries one *Smn^{2B}* allele that harbors a three-nucleotide switch in exon 7, and one *Smn* allele whose exon 7 is floxed by loxP sites (*Smn^{F7}*). The *Alb-Cre* transgene has also been crossed into this background to provide liver-specific expression of the Cre recombinase. B. Littermate controls do not harbor Cre recombinase but do carry one *Smn^{2B}* allele and are thus used as heterozygous controls. C. Upon crossing *Alb$^{Cre/+}$;Smn$^{F7/+}$* with *Smn$^{2B/F7}$* animals, Cre recombination will induce specific excision of exon 7 in the liver, leading to reduced SMN protein production only in the liver.

2. **Liver-specific SMN depletion in *Alb$^{Cre/+}$;Smn$^{2B/F7}$* mice.** A–E. Immunoblots were performed to assess SMN protein levels in various tissues from *Smn$^{2B/+}$*, *Smn$^{2B/-}$*, +/+;*Smn$^{2B/F7}$*, and *Alb$^{Cre/+}$;Smn$^{2B/F7}$* mice at P19. Membranes were probed for SMN and then reprobed for α-tubulin (loading control). A'-E'. Bar graphs show quantification of SMN protein levels in the liver (A'), brain (B'), muscle (C'), spinal cord and pancreas (E'), normalized to α-tubulin. n ≥ 3, mean ± SEM. Statistical significance indicated by *$p < 0.05$, **$p < 0.01$ or p values, following Brown-Forsythe and Welch ANOVA.

3. **Impact of liver specific SMN depletion in hepatocyte morphology and lipid accumulation.** A. Representative images of H&E (top row, scale bar 100 μm) and Oil Red-O (bottom row, scale bar 50 μm) stained liver sections from *Smn$^{2B/+}$*, *Smn$^{2B/-}$*, +/+;*Smn$^{2B/F7}$*, and *Alb$^{Cre/+}$;Smn$^{2B/F7}$* mice at P19. B. Bar graph shows quantification of liver triglycerides. n ≥ 3, mean ± SEM. Statistical significance indicated by *$p < 0.05$, **$p < 0.01$, following Brown-Forsythe and Welch ANOVA.

4. **Levels of key proteins involved in liver homeostatic function are unchanged in liver-specific SMN-depleted mice.** A-C. Immunoblots were performed to assess heme oxygenase (HO, A-A'), transferrin (B-B') and P62 (C-C') protein levels in the liver. α-tubulin was used as a loading control. A'-C' Bar graphs show quantification of HO (A'), transferrin (B'), and P62 (C') levels. D. Bar graph depicts quantification of liver IGF-1. n ≥ 3, mean ± SEM. Statistical significance indicated by *$p < 0.05$, ***$p < 0.001$, following Brown-Forsythe and Welch ANOVA.

5. **Contribution of liver specific SMN depletion to pancreatic pathology.** A. Representative immunofluorescent images of pancreatic islets stained for glucagon (red) and insulin (green) from *Smn$^{2B/+}$*, *Smn$^{2B/-}$*, +/+;*Smn$^{2B/F7}$*, and *Alb$^{Cre/+}$;Smn$^{2B/F7}$* mice at P19. Scale bar 50 μm. B-C. Bar graphs show quantification of insulin (B) and glucagon (C) positive cells relative to the total number of DAPI positive cells within the pancreatic islet. D. Bar graph depicts non-fasting blood glucose levels from P19 mice across different genotypes. n ≥ 3, mean ± SEM. Statistical significance indicated by *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, following Brown-Forsythe and Welch ANOVA.

6. **Impact of liver specific SMN depletion on motor neuron cell body numbers.** A. Representative images of lumbar spinal cord anterior horns stained for ChAT (red) and DAPI

(blue) from $Smn^{2B/+}$, $Smn^{2B/-}$, $+/+;Smn^{2B/F7}$, and $Alb^{Cre/+};Smn^{2B/F7}$ mice at P19. B. Bar graph shows quantification of motor neuron cell body numbers. n ≥ 3, mean ± SEM. Statistical significance indicated by **$p < 0.01$, ***$p < 0.001$, following Brown-Forsythe and Welch ANOVA. Scale bar = 50 μm.

7. **Impact of liver specific SMN depletion on neuromuscular junction pathology.** A. Representative images of *transversus abdominis* (TVA) muscle stained with bungarotoxin (red), neurofilament (NF) (green) and synaptic vesicle protein 2 (green) from $Smn^{2B/+}$, $Smn^{2B/-}$, $+/+;Smn^{2B/F7}$, and $Alb^{Cre/+};Smn^{2B/F7}$ mice at P19. *Depicts occupied endplate, whereas **depicts normal neurofilament distribution; arrowheads show NF accumulation and arrows show unoccupied endplate/denervation. B-C. Bar graphs show quantification of neurofilament accumulation (B) and endplate occupancy (C). n = 3, mean ± SEM. Statistical significance indicated by **$p < 0.01$, ***$p < 0.001$, following Brown-Forsythe and Welch ANOVA. Scale bar = 50 μm.

8. **Impact of liver specific SMN depletion on muscle fiber size and distribution.** A. Representative images of H&E-stained *tibialis anterior* muscle sections from $Smn^{2B/+}$, $Smn^{2B/-}$, $+/+;Smn^{2B/F7}$, and $Alb^{Cre/+};Smn^{2B/F7}$ mice at P19. B. Bar graph shows quantification of muscle fiber cross-sectional area. C. Graph demonstrating frequency of muscle fiber size across different genotypes. n ≥ 3, mean ± SEM. Statistical significance indicated by **$p < 0.01$, following Brown-Forsythe and Welch ANOVA. Scale bar = 20 μm.

9. **Impact of liver specific SMN depletion on motor function**. A. Schematic representation of experimental design. The righting reflex test was conducted from P7 to P13, the inverted mesh grip test from P13 to P25, and the pen test from P19 to P25. Assessments were performed every two days. Weight was measured every two days until day 30, and then weekly until day 60. Animal welfare was monitored throughout the 60-day period during weight measurements. B. Kaplan–Meier survival curve comparing $Alb^{Cre/+};Smn^{2B/F7}$ and $+/+;Smn^{2B/F7}$ mice up to 60 days. C-F. Graphs show weight (C), righting reflex (D), inverted mesh grip (E) and pen test (F). n = 5 per genotype, mean ± SEM. B: Kaplan–Meier survival analysis; C-F: two-way ANOVA, followed by Šídák's method.

Figure 1

Figure 2

Figure 3

Figure 4

A

| $Smn^{2B/+}$ | $Smn^{2B/-}$ | $+/+;Smn^{2B/F7}$ | $Alb^{Cre/+};Smn^{2B/F7}$ |

Glucagon/insulin/DAPI

B

Insulin+ Cells/islet (%)

C

Glucagon+ Cells/islet (%)

D

Glucose (mmol/L)

- $Smn^{2B/+}$
- $Smn^{2B/-}$
- $+/+;Smn^{2B/F7}$
- $Alb^{Cre/+};Smn^{2B/F7}$

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9