1

4 **Structural basis for higher-order DNA binding by a bacterial transcriptional regulator**

6 *Frederik Oskar Graversgaard Henriksen[1], Lan Bich Van[1], Ditlev Egeskov Brodersen[1], and*

7 *Ragnhild Bager Skjerning[1]**

10 *[1]Department of Molecular Biology and Genetics, Universitetsbyen 81, Aarhus University, DK-8000*

11 *Aarhus C, Denmark, *To whom correspondence should be addressed (raba@mbg.au.dk)*

ABSTRACT

Transcriptional regulation by binding of transcription factors to palindromic sequences in promoter regions is a fundamental process in bacteria. Some transcription factors have multiple dimeric DNA-binding domains, in principle enabling interaction with higher-order DNA structures; however, mechanistic and structural insights into this phenomenon remain limited. The *Pseudomonas putida* toxin-antitoxin (TA) system Xre-RES has an unusual 4:2 stoichiometry including two potential DNA-binding sites, compatible with a complex mechanism of transcriptional autoregulation. Here, we show that the Xre-RES complex interacts specifically with a palindromic DNA repeat in the promoter in a 1:1 molar ratio, leading to transcriptional repression. We determine the 2.7 Å crystal structure of the protein-DNA complex, revealing an unexpected asymmetry in the interaction and suggesting the presence of a secondary binding site, which is supported by structural prediction of the binding to the intact promoter region. Additionally, we show that the antitoxin can be partially dislodged from the Xre-RES complex, resulting in Xre monomers and a 2:2 Xre-RES complex, neither of which repress transcription. These findings highlight a dynamic, concentration-dependent model of transcriptional autoregulation, in which the Xre-RES complex transitions between a non-binding (2:2) and a DNA-binding (4:2) form.

KEYWORDS

Crystal structure, HTH domain, RES toxin, Toxin-antitoxin, Transcriptional regulation

AUTHOR SUMMARY

Bacteria regulate their gene expression to respond to environmental stress, to evade antibiotics, and to maintain population stability. In this study, we investigate how the *xre-res* toxin-antitoxin system from *Pseudomonas putida* controls its own expression. Using a combination of microbiology,

structural biology, biophysical assays, and computational modelling, we discover how the Xre-RES

protein complex represses its own transcription through direct binding to a specific DNA element in

the promoter region. We show structurally that the Xre-RES complex adopts a unique 4:2

stoichiometry and binds DNA in an unusual asymmetrical manner. Moreover, the complex was

found to shift between two different forms: one that binds DNA and represses transcription, and one

that does not. We further demonstrate that this shift is dynamic and depends on the relative

concentration of Xre antitoxin. Our findings provide new insight into how bacteria fine-tune gene

expression and offers a model of transcriptional control based on protein stoichiometry and

structural asymmetry.

INTRODUCTION

Regulation of transcription is a fundamental mechanism in bacteria that allows adaptation to

constantly changing environments, and RNA polymerase (RNAP) plays a central part in this. The

core RNAP enzyme has a $\beta\beta'\alpha_2$ subunit composition [1] but in order to form a holoenzyme and

initiate transcription at a particular promoter, the core enzyme needs to interact with a σ subunit [2,

3]. The σ subunit ensures promoter recognition and positioning of the RNAP holoenzyme at the

promoter and unwinding of the DNA duplex near the transcriptional start site (TSS). Transcriptional

activity can be modulated by repressor proteins that bind directly to a segment of the promoter (the

operator) and inhibits transcription [4]. Repression can occur by one of three mechanisms: steric

hindrance of RNAP binding to the promoter, interference with post-recruitment steps in

transcriptional initiation such as the assembly of the RNAP holoenzyme, or DNA looping upon

repressor binding [1]. A common feature of many repressor proteins is the presence of a DNA-

binding helix-turn-helix (HTH) domain [5]. The generic HTH domain is a simple fold comprised of

three core helices of which the second and third helix are connected by a sharp turn and makes up

the HTH motif [5, 6]. The third helix is also called the 'recognition helix' and usually recognizes a

symmetrical sequence element in the DNA by inserting directly into the major groove [7]. The

binding efficiency a repressor protein to an operator site depends on its binding affinity, which

again is determined by base-specific readouts and intrinsic DNA shape, however, in many cases, the

mechanism by with repressor proteins recognise DNA *in vivo* remains unclear [8-10].

Due to the symmetrical nature of DNA, most HTH-containing protein repressors are functional as

dimers, which sometimes depend on even higher-order structures and additional proteins. Thus,

HTH domains often bind palindromic DNA sequences separated by approximately a single helical

turn, allowing for the recognition helices to insert into adjacent major grooves [7]. In the

widespread *xre-res* system [11], the encoded Xre (Xenobiotic response element) protein contains a

putative HTH DNA-binding domain that is stabilised in a higher-order structure through interaction

with the associated RES-domain protein [12-14]. The two proteins are expressed from two-gene

operons that have been classified as toxin-antitoxin (TA) systems, which are widely distributed

cellular regulators in bacteria that encode a protein toxin that inhibits cell growth along with a

cognate protein antitoxin that neutralises it [15]. The RES toxin has an ADP-ribosyltransferase

fold and expression of an orthologue from the insect pathogen, *Photorhabdus luminescens*,

has been shown to result in efficient depletion of intracellular NAD^+ in *E. coli*, suggesting that

the RES toxin is a NAD^+ glycohydrolase (NADase). Similar results were obtained for the RES-

domain toxins MbcT from *Mycobacterium tuberculosis* [12], SlvT from *P. putida* S12 [16],

PA14_51010 from *P. aeruginosa* PA14 [17], and VPA0770 from *Vibrio parahaemolyticus* [18],

while PrsT from *Sinorhizobium meliloti* toxin is structurally similar appears to function as an ADP

ribosyltransferase (mART) targeting and inactivating phosphoribosyl pyrophosphate synthetase

(Prs) [13]. The role of RES domain-containing TA systems in bacterial cell biology is still largely

unknown, but a recent study showed that the *P. aeruginosa* orthologue, PA14_51010 (NatT),

mediates NAD^+ depletion leading to drug-tolerant dormant cell populations *in vitro* and suggesting

that the accumulation of NatT-activating mutations observed during *P. aeruginosa* infections in

human cystic fibrosis patients increases antibiotic tolerance [19].

The crystal structure of a Xre-RES complex from *Pseudomonas putida* revealed an unusual 4:2 TA

stoichiometry in which each of the RES domains of a central toxin dimer interacts with a dimer of

Xre antitoxins, thus generating two complete HTH domains [14]. The HTH domain of this Xre

antitoxin (Xre^{Pp}) is structurally similar to the bacteriophage lambda Cro repressor [20] and could

thus potentially play a part in transcriptional regulation and toxin control through binding of

the antitoxin to the TA promoter region, as observed for many TA systems [21]. For some of

these, the presence of the toxin only stabilises the interaction with DNA up to a certain level

after which a higher stoichiometry of toxin molecules leads to de-repression by a mechanism

termed *conditional cooperativity* [22-24]. However, other modes of autoregulation exist,

including via recruitment of external regulators, three-component TA systems in which the

DNA binding domain is present in a third protein separate from the antitoxin, and TA

modules with reversed gene order and multiple promoter sites [25]. The observation that the

P. putida Xre-RES complex contains two complete DNA-binding domains each composed by

two HTH domains suggests the system may be regulated in a novel way. However, the

organisation of the promoter region and the mechanism of DNA binding and transcriptional

regulation are currently unknown.

In this paper, we dissect the mechanism of transcriptional autoregulation of the *P. putida xre-res*

system. Our findings reveal that a Xre–RES^{Pp} complex with 4:2 stoichiometry binds in a 1:1 ratio to

an imperfect inverted repeat region in the promoter region and effectively represses transcription.

We determine a 2.7 Å crystal structure of the Xre-RES^{Pp} complex bound to the imperfect inverted

repeat, showing that the interaction is unusually asymmetric. Moreover, interactions to one end of a

symmetry-related DNA-duplex suggests a second binding site in the operator region, which is

supported by structure prediction of the interaction to the intact promoter DNA sequence. Finally,

we show that the isolated Xre^{Pp} antitoxin is a monomer that only weakly represses the promoter.

Surprisingly, individual Xre^{Pp} monomers can dislodge from the 4:2 complex, resulting in a 2:2

complex incapable of binding DNA. Together, our data provides a model for how the Xre-RES^{Pp}

complex autoregulates transcription through a balance between binding and non-binding forms,

driven by protein stoichiometry and asymmetry as well as general insights into interaction of

transcription factors with higher-order DNA in bacteria.

RESULTS

The *P. putida xre-res* locus is transcriptionally auto-regulated

The presence of a DNA-binding HTH domain in Xre^{Pp} suggested that the antitoxin and/or the Xre-

RES TA complex could be involved in regulating transcription of the *P. putida xre-res* operon via

direct promoter binding. To test this hypothesis, we analysed the *xre-res*^{Pp} promoter region (P$_{XR}$)

and identified three inverted repeats and one directional repeat (named S1-S4, **Fig 1A and Fig**

S1A). To investigate the promoter activity *in vivo*, we constructed a low-copy GFP reporter plasmid

(pGH254Kgfp) containing the intact P$_{XR}$ promoter region, including the 5' end (30 bp) of *xre*^{Pp},

transcriptionally fused to *gfpmut2* encoding a Green Fluorescent Protein (GFP) variant

(S65A/V68L/S72A) with enhanced fluorescence emission and more efficient folding at 37°C as

compared to the wild type GFP [26]. The resulting construct, as well as a plasmid expressing *xre-*

res^{Pp} under arabinose-inducible control, was introduced into *E. coli* MG1655 by transformation,

and the GFP signal monitored throughout growth in the presence or absence of arabinose (**Fig 1B**).

The results show that the P_{XR} promoter is active in *E. coli* during exponential growth (**Fig 1C**).

Moreover, induction of *xre-res*^{Pp} in this background results in complete absence of GFP signal (**Fig.**

1D, orange curve) compared to the strain containing an empty pBAD33 vector (**Fig. 1D**, green

curve). Together, this supports that *xre-res*^{Pp} negatively regulates expression from P_{XR}.

142
Figure 1. Organisation, activity, and regulation of the *xre-res*^{Pp} operon.
(A) Schematic representation the *P. putida* KT2440 *xre-res*^{Pp} locus including the 172-bp promoter region
(P_{XR}) and the upstream gene (*nfsB*). Top, the *xre*^{Pp} gene (orange) encodes a protein antitoxin with an N-
terminal helix-turn-helix (HTH) domain and a C-terminal domain of unknown function (DUF2384), known
to interact with the RES toxin, while *res*^{Pp} (blue) encodes a NADase toxin with a RES domain. Below,
close-up of the P_{XR} promoter region showing the three inverted repeats (S1, S2, and S4) and one directional
repeat (S3), of which S4 is imperfect and contains one mismatch (yellow box, see also **Fig S1A**). (B)
Overview of the GFP-based assay to measure regulation of P_{XR} in *E. coli* MG1655 *in vivo*. The pBAD33
vector allows for arabinose induction of the *xre-res*^{Pp} genes while P_{XR} coupled to GFP allows for read-out
of the transcriptional activity of the promoter. (C) and (D) Promoter activity assays in *E. coli* MG1655.
Activity is measured as GFP signal in relative fluorescence units (RFU) during growth without ((C), -Ara)
or with ((D), +Ara) arabinose using the promoter reporter pGH254Kgfp::P_{XR} (P_{XR}), as well as empty
pBAD33 (-, green curves) or pBAD33::*xre-res*^{Pp} vector (*xre-res*, orange curves). All curves represent mean-
of-mean values (line, n = 3) with standard error of mean (SEM) shown as a shadow. The dotted vertical
line indicates the 350 min time point

Expression of *xre-res*^{Pp} represses promoter activity via an imperfect repeat upstream of *xre*^{Pp}

To determine the minimal promoter requirements for active transcription from the *xre-res*^{*Pp*} locus,

we repeated the assay using variants of P_{XR} where one or more of the four identified repeats (S1-S4)

were systematically removed (**Fig 2A** and **S1 Table**). To simplify the comparison and since the P_{XR}

promoter was found to be most active during late exponential growth, the GFP fluorescence level

measured at t=350 min was used for this analysis (complete curves are included in **Fig S1** and **S2**).

Removal of S1 (ΔS1) did neither affect promoter activity nor repression by *xre-res*^{Pp}, while removal

of both S1 and S2 (ΔS1-S2), resulted in significantly increased promoter activity that can still be

fully repressed by *xre-res*^{Pp} (**Fig 2B**). In contrast, removal of either S1 through S3 (ΔS1-S3), S3 and

S4 (ΔS3-4), or just S4 (Δ4), completely abolished transcription (**Fig 2B**). Interestingly, removal of

only the central palindromic part of the imperfect inverted repeat in S4 (Δrepeat, **Fig 2C** and **S1**

Table) did not affect promoter activity while expression of *xre-res*^{Pp} only weakly (and non-

significantly) repressed transcription of this variant (**Fig 2D**). Together, this suggests that

transcriptional control by *xre-res*^{*Pp*} depends on the imperfect inverted repeat in the S4 region while

transcription does not. To investigate the importance of the imperfect part of the inverted repeat, we

created two variants with perfect repeats by either introducing G->T on the coding strand in the first

half of the repeat (5'-perf) or A->C in the second half of the repeat (3'-perf) (**Fig 2C** and **S1 Table**).

Both substitutions decreased promoter activity as compared to wild type but did not affect the

ability of *xre-res*^{*Pp*} to repress transcription, suggesting that the imperfectness of the inverted repeat

is required for optimal transcription but not repression (**Fig 2D**).

180
Fig 2. Determination of the minimal promoter requirements for transcription and regulation of P_{XR}.
(A) Detailed overview of the P_{XR} promoter (top) and schematic representation of promoter variants (below).
Full lines indicate the included parts while dotted lines represent regions that have been removed. Complete
growth curves and details of the constructs are found in **Fig S1** and **S1 Table**. (B) Promoter activity assays
in *E. coli* MG1655. The reporter plasmid pGH254Kgfp encoding promoter variations was combined with
pBAD33::*xre-res*^{Pp}. For each variant, the GFP signal (RFU) measured at t=350 min from non-induced
cultures (-Ara, green bars) is compared to that from arabinose-induced cultures (+Ara, orange bars). Mean
RFU values extracted at the same time point from three independent experiments were treated as biological
replicates (dots, $n = 3$) and error bars show SEM. Complete growth curves are included in **Fig S1**. Statistical
analysis was performed using two-way analysis of variance (ANOVA) followed by Dunnett's multiple
comparisons test. *ns*, not significant; ***, $p < 0.01$; ****, $p < 0.0001$. (C) Schematic representation of the
repeat variants of the imperfect inverted repeat in S4 of the P_{XR} promoter region. The DNA sequence of the
imperfect inverted repeat is highlighted (light grey), including the mismatch in the first repeat (red), the
perfect part of the repeat (dark grey) and the alterations (green). The mismatch was removed by a
substitution either in the first (5'-perf) or second (3'-perf) half of the repeat. (D) As b, but using promoter
variations from c. *ns*, not significant; *, $p < 0.05$; **, $p < 0.01$.
197

The Xre-RES^{Pp} complex binds directly to S4 *in vitro*

The presence of an HTH domain in Xre and the observed repression of transcription upon

expression of *xre-res^{Pp}* indicated that the Xre-RES^{Pp} protein complex might bind directly to DNA.

To investigate this, we analysed the binding between the Xre-RES^{Pp} protein complex and different

parts of the P_{XR} promoter *in vitro*. First, we expressed and purified the intact Xre-RES^{Pp} complex

via a N-terminal His$_6$-tag on the RES toxin (Xre-RES^{Pp}$_{His6}$) and used size exclusion

chromatography coupled to multi-angle light scattering (SEC-MALS) to show that Xre and RES$_{His6}$

form a protein complex with an estimated MW of 92 ± 4 kDa (**Fig 3A**), which best matches the

theoretical mass of the 4:2 Xre-RES complex of approximately 103 kDa observed in the crystal

structure (**Fig 3B**) [14]. We then analysed the ability of the purified complex to bind a set of 30 bp

dsDNA probes each covering one of the four sequence elements (S1 to S4) in the P_{XR} promoter (**Fig**

1A and **Fig S1A**). For this, purified Xre-RES$^{Pp}_{His6}$ complex and dsDNA probes were incubated in

1:1 molar ratio followed by analytical SEC (a-SEC). A significant change in elution volume of 0.2

mL and a concomitant change in the 260:280 nm absorbance ratio from 0.54 to 0.68 was observed

for complex mixed with S4 dsDNA (**Fig 3C**), but not for S1, S2 or S3 dsDNA (**Fig S3A-C**),

suggesting that the Xre-RES^{Pp} complex binds specifically to the S4 element of the P_{XR} promoter.

Previously, structure determination of the Xre-RES^{Pp} complex has shown that the 4:2

heterohexamer contains two complete Cro-like Xre^{Pp} dimers [14], each potentially capable of

binding dsDNA via interaction with adjacent major grooves [27]. To investigate whether the S4

element is the only binding site in the promoter, we determined the binding stoichiometry by

analysing the binding at different protein:DNA ratios using SEC-MALS. At a 1:1 ratio of Xre-

RES^{Pp} complex to dsDNA duplex, and thus a ratio of putative binding sites to DNA of 2:1, we

found that the protein complex was unable to bind all DNA present in solution (**Fig S3D**), and

increasing this to a 1:2 ratio of protein complex to DNA, no additional shift in the elution volume or

calculated mass was observed as compared to the 1:1 ratio (**Fig S3D**). To further probe the

stoichiometry of binding between the Xre-RES^{Pp} complex and S4 dsDNA, we saturated the protein

complex with increasing amounts of DNA and used high-resolution SEC (2.4 mL) to check for

indication of a second DNA binding site. Even at a molar ratio of protein complex to dsDNA of

1:10, only a marginal shift in the elution volume of 0.01 ml (0.6% CV) was observed (**Fig S3E**).

Together, this suggests that the 4:2 Xre-RES^{Pp} complex only binds a single copy of S4 dsDNA. To

further quantify the binding interaction between the Xre-RES^{Pp} complex and S4, we determined the

binding coefficient (K_D) using isothermal titration calorimetry (ITC). This showed that the Xre-

RES^{Pp} complex binds with a 1:1 ratio and high affinity to the native S4 element ($K_D = 231 \pm 46$ nM,

Fig 3D), while the affinities towards both the 5'-perf ($K_D = 341 \pm 264$ µM) and 3'-perf variants (K_D

$= 80 \pm 79$ µM) of S4 were drastically reduced (**Table 2**). Given that repression by *xre-res^{Pp}* is intact

for these two variants (5'-perf and 3'-perf) *in vivo*, these data are consistent with the existence of a

second DNA binding site that strengthens the protein-DNA interaction in the context of the full-

length promoter.

238
Figure 3. Xre-RES^{Pp} forms a 4:2 hexameric complex in solution and interacts specifically with the S4
element of the promoter.
(A) SEC-MALS analysis of purified Xre-RES^{Pp}$_{His6}$ complex. The chromatogram shows the elution profile
as measured by absorption at 280 nm (blue) and the molecular mass measured by MALS (red). The average
mass corresponding to the middle of the peak is 92 ± 4 kDa (dotted line). (B) The crystal structure of
heterohexameric Xre-RES^{Pp} (PDB: 6GW6) with total mass (103 kDa) and HTH domains indicated [14].
The RES toxins are coloured in shades of blue, while the Xre antitoxins are in shades of green. (C)
Analytical SEC (a-SEC) analysis of Xre-RES^{Pp}$_{His6}$ (Xre-RES, blue), S4 (S4-DNA only, red), and DNA
mixed with protein in a 1:1 ratio (Xre-RES + S4-DNA, purple). Black arrows represent known molecular

The structure of the Xre-RES^{Pp}-DNA complex reveals an unusual, asymmetrical DNA

binding site

To get a molecular understanding of the interaction between Xre-RES^{Pp} and promoter DNA, we

determined the crystal structure of the Xre-RES^{Pp} complex bound to a 30 bp dsDNA oligo covering

the S4 element to 2.7 Å (R / R$_{free}$ = 22.8% / 26.9%, **Fig 4A, Table 1**). The structure contains four

complete heterohexameric (4:2) Xre-RES^{Pp} complexes and four DNA duplexes in the

crystallographic asymmetric unit (**Fig S4A**). In all molecules, one Xre dimer of each Xre-RES^{Pp}

complex interacts strongly with two adjacent major grooves on dsDNA (**Fig 4A** and **Fig S4B**). The

interaction between the Xre dimer and DNA causes the major groove on the opposing side of the

double helix to expand from 11.8 Å in canonical B-DNA to 14 Å (**Fig 4A**). Significantly, we

observe that the Xre dimer is asymmetrical in the way it interacts with DNA and overall approaches

the double helix in a different way than classical HTH domains such as the Cro-repressor [28].

More specifically, the angle between the helical axis of DNA and that of the recognition helix of the

outer Xre molecule is 61.4° while it is 57.5° for the inner Xre molecule. These values are both

significantly smaller and more different from each other than observed for the canonical structure of

the Cro repressor bound to DNA (PDB 1RPE), in which the the two recognition helices of the Cro

dimer approach DNA at nearly identical angles of 68.8° (**Fig S4C**) (Shimon & Harrison, 1993).

Several residues in Xre, in particular Arg60, Arg67, Thr68, Lys70, and Arg72, interact tightly with

C13, C20, and G21 on the (+) strand, as well as T7, T8, T9, and A14 on the (-) strand of S4 DNA

(**Fig 4B** and **4C**). The dominant interactions are centred around the conserved Arg67 of the

innermost Xre, which engages in hydrogen bonds to the N3 atom of T9 of the DNA (-) strand (**Fig**

4C). Interestingly, Arg67 of the outermost Xre displaces A14 of the (+) strand and forms hydrogen

bonds to the N3 of C13 (+) (**Fig 4C**). Arg60, Thr68, Lys70, and Arg72 also make non-sequence

specific interactions to the phosphodiester backbone of DNA. Of the interacting residues, both

Thr68 and Arg72 are functionally conserved across *xre-res* loci from a wide range of bacterial

species, while Arg67 and Lys70 appear conserved in a subset of species (**Fig 4D**). To support the

importance of Arg67 in DNA binding by Xre, we substituted this residue with alanine (R67A) and

found that transcriptional repression by *xre*^{R67A}-*res in vivo* was abolished (**Fig 4E** and **Fig S4D**).

Intriguingly, inspection of the second Xre dimer of the Xre-RES heterohexamer reveals that there

are interactions to one end of a DNA duplex from a symmetry-related complex (**Fig 4F**). The

specific interactions made at this site vary slightly between complexes in the asymmetric unit but in

all cases include base-specific interactions between Arg67 of the outermost Xre molecule to O2 on

T7 on the (+) strand, and non-sequence specific interactions between Thr68 and Ser71 to the

backbone of A4 and C5 of the (+) strand as well as between Arg75 and Gln77 and C19 of the (-)

strand (**Fig 4B**). In summary, the structure of DNA-bound Xre-RES^{Pp} confirms that one of the two

Xre dimers in the complex recognises S4 dsDNA using base-specific interactions dependent on

Arg67 and moreover, that the second Xre dimer is capable of simultaneously interacting with DNA

supporting the idea that there might be a secondary DNA binding site inside the full operator

region.

295
Figure 4. Crystal structure of the DNA-bound Xre-RES^Pp complex.
(A) Overview of the *P. putida* Xre-RES complex (green/blue cartoon) is bound to a 30 bp dsDNA
(red/orange) with both HTH recognition helices of the of a single Xre dimer interacting with DNA. (B)
Schematic overview of the interactions between Xre and DNA including both sequence specific and
unspecific contacts as well as the contacts from the symmetry related interaction. The shaded DNA region
could not build. (C) Close-up on the interaction between Xre and DNA with interactions between selected
residues of both Xre molecules and the DNA highlighted. (D) Sequence alignment of *P. putida* Xre and a
range of homologous proteins from other bacterial species. UniProt IDs are listed in parentheses for all
proteins, together with and PDB IDs for *P. putida* Xre (PDB ID 6GW6) [14], *M. tuberculosis* MbcA (PDB

ID 6FKG) [12], *V. parahaemolyticus* (PDB ID 8GUG) [18], and *P. aeruginosa* NatR (PDB ID 8QNL) [19].
Residues conserved with *P. putida* Xre are shown in red with functionally similar residues in orange.
Starting and ending residue numbers are listed at each end of the sequences. (E) Promoter activity in *E. coli*
MG1655 for promoter reporter plasmid pGH254Kgfp::P_{XR} (P_{XR}) in combination with pBAD33 (-),
pBAD33::*xre*^{R67A}-*res*^{Pp} (*xre*^{R67A}-*res*), or pBAD33::*xre-res*^{Pp} (*xre-res*), as indicated. Complete growth
curves and details of the constructs are found in **Fig S4**. (F) In the crystal, the Xre-RES complex
(blue/green) interacts with a primary dsDNA molecule (DNA 1) as well as a symmetry-related DNA
molecule (DNA 2).
313

The Xre^{Pp} dimer is stabilised by higher-order complex formation and required for

transcriptional repression

Since the structure shows that only one of the two Xre dimers in the complex is bound to S4

dsDNA, we speculated whether Xre alone can repress transcription. To investigate this, we used the

transcriptional coupling to GFP during to compare the activity of the wild type P_{XR} promoter during

induction of either *xre*^{Pp} or *xre-res*^{Pp}. While expression of the complex fully repressed transcription

as seen before, expression of *xre*^{Pp} alone only partially, and not significantly, repressed transcription

(**Fig 5A**). Moreover, we note that the promoter activity between biological replicates was

considerably more variable upon expression of *xre*^{Pp} (**Fig S5A** and **5B**), indicating that the

Xre^{Pp}:DNA interaction is unstable in the absence of RES^{Pp}. To assess if the weak repression upon

induction of *xre*^{Pp} alone is due to lack of formation of a stable Xre dimer, we expressed and purified

the isolated Xre^{Pp} antitoxin with a C-terminal His-tag (Xre^{Pp}$_{CHis6}$) to avoid disturbing the

dimerisation interface. Purified Xre^{Pp}$_{CHis6}$ was very prone to aggregation and could not be

concentrated above 10 µM, supporting that complex formation contributes to stabilising the

antitoxin. Moreover, the mass of purified Xre^{Pp}$_{CHis6}$ antitoxin was determined by SEC-MALS to 22

± 1 kDa, corresponding to a monomer of Xre^{Pp} (**Fig 5B**). In summary, we conclude that the

presence of RES^{Pp} is required for formation of a functional and DNA-binding competent Xre^{Pp}

dimer.

To further investigate the stability the Xre^{Pp} dimer within the complex, mass photometry (MP), which

is a single-particle mass-determination method, was used to determine the mass of the purified Xre-

RES^{Pp} complex. The particle distribution was not uniform and at 10 nM we observed two populations

of which the larger and most abundant was around ~90 kDa and the smaller was around 76 kDa. At

5 nM, the same two populations were observed, but with the smaller MW complex being most

abundant, however an additional peak around 50 kDa was also seen. Measuring at 2 nM revealed a

mostly homogeneous peak at with 50 kDa, with a trailing tail of larger particles. This concentration

dependent change in mass distribution and median masses (87 ± 15 kDa, 76 ± 14 kDa, 54 ± 14 kDa)

corresponds roughly to a transition from the 4:2 (103 kDa) complex via a 3:2 (86 kDa) to a 2:2 (69

kDa) complex (**Fig 5C**). Together, this suggests that even in presence of RES; the Xre^{Pp} dimer is

unstable and that the Xre-RES^{Pp} complex can exist on both 4:2, 3:2, and 2:2 forms. To confirm the

existence of the 2:2 complex *in vitro*, we expressed and purified a variant of the Xre-RES^{Pp} complex

with a N-terminal His_6-tag on Xre (Xre_{NHis6}-RES^{Pp}), which, based on the crystal structure, should

prevent Xre dimerisation. The expressed protein was highly prone to precipitation during purification,

consistent with a destabilised Xre^{Pp} dimer that would result in free and unstable Xre^{Pp} monomers as

well as a 2:2 complex. This was supported by SEC, which showed two peaks corresponding to the

2:2 complex and isolated Xre^{Pp} antitoxin, respectively (**Fig S6A** and **6B**). The successful isolation of

a 2:2 complex was further confirmed by mass photometry, where the main protein population at both

10 nM and 2 nM exhibited masses of 63 ± 18 kDa (**Fig 5D**).

As the isolated Xre^{Pp} antitoxin is a monomer and induction of xre^{Pp} alone only weakly represses

transcription *in vivo*, we predicted that the isolated 2:2 complex, containing two Xre monomers,

would also have reduced DNA affinity. Analysis by high-resolution SEC column of samples

containing only S4 DNA, only the 2:2 complex, or 2:2 complex and S4 DNA in ratio of 1:10 showed

that the elution volume of the 2:2 complex is not affected by the addition of DNA (**Fig 5E**). This

supports that the 2:2 is a non-binding form of the Xre-RES^{Pp} complex and thus a model in which the

complex exists in a concentration-dependent equilibrium between non-binding (2:2) and DNA-

binding (4:2) forms.

362
Figure 5. DNA binding requires the Xre dimer, which is unstable at low concentrations.
(A) Promoter activity assays in *E. coli* MG1655, where the promoter reporter plasmid pGH254Kgfp::P$_{XR}$
(P$_{XR}$) was combined with either empty pBAD33 (-), pBAD33::*xre*^{Pp} (*xre*), or pBAD33::*xre-res*^{Pp} vector
(*xre-res*), as indicated Complete growth curves and details of the constructs are found in **Fig S4**. Statistical
analysis was performed using a two-sample t -test (with equal variances). *ns*, not significant; *, $p < 0.05$.
(B) SEC-MALS analysis showing that isolated Xre$^{Pp}_{CHis6}$ is a monomer with an approximate mass of $22 \pm$
1 kDa in solution. The chromatogram shows elution profile measured by 280 nm absorbance (blue) and
calculated mass from MALS (red). (C) Single-molecule mass determination of Xre^{Pp}-RES$^{Pp}_{His6}$ at 2, 5, and
10 nM. The violin plots represent the density of all measured single-molecule events. The solid black
horizontal line indicates the median mass, while the vertical line shows the standard deviation. The

theoretical masses of the Xre^{Pp}-RES^{Pp}_{His6} complex with 4:2, 3:2, and 2:2 stoichiometries are indicated by
dotted lines. Significance was tested with the Wilcoxon signed rank test. *****, $p < 5.55e-7$. (D) Single-
molecule mass determination for Xre_{His6}-RES^{Pp} on the 2:2 complex form at 2 and 10 nM, as in c. (E) High
resolution analytical SEC (a-SEC) analysis of Xre_{NHis6}-RES (Xre-RES, blue), S4 (DNA, red), amd DNA
mixed with protein in a 1:10 ratio (Xre-RES + DNA, purple). The full line shows 280 nm absorbance, and
the dotted line is 260 nm absorbance. (F) Alphafold3 predicted structure of Xre-RES complex interacting
with the full 172 bp promoter region (not all nucleotides shown). (G) Structural alignment of crystal
structure of the DNA-bound Xre-RES complex, including the symmetry-related dsDNA molecule
(blue/green/orange) with the AlphaFold 3 prediction (grey).

Structural basis for Xre-RES^{Pp} binding at two sites in the P$_{XR}$ promoter region

Since the crystal structure suggests that both DNA-binding Xre dimers on a Xre-RES

heterohexamer can engage in DNA binding simultaneously, we decided to use Alphafold 3 (AF3) to

predict the interaction between Xre-RES and the full 172 bp promoter [29]. The resulting prediction

showed high confidence for the Xre-RES protein complex (average pLDDT 90.1), with a more

varying confidence for the DNA having pLLDTs between 24.2 and 81.5 (average 53.9). The entire

prediction has a PTM score of 0.64 which, while low, is unsurprising given the variance in pLDDT

and the length of free DNA. The interaction has an iPTM score of 0.6, which is in the grey zone for

confidence and thus would require additional support for validation [29]. This support is provided

by comparison to the crystal structure, as the predicted model has both Xre dimers engaged in DNA

binding about 20-bp apart (**Fig S7A-C**), with one site closely matching the primary S4 DNA

binding site observed in the crystal structure and the other located upstream overlapping partly with

S3 as well as involving the region between S3 and S4 (**Fig 5F**). Comparison of the prediction to the

crystal structure reveals a good correspondence in the details of the interaction at the primary site,

including the central of Arg67 (**Fig S7D**). We consider it, in itself, remarkable that AF3 is able to

precisely map the binding site of a bacterial transcription factor within such a long stretch of DNA

with nucleotide precision, especially given that there are no structures of similar protein complexes

bound to DNA available. However, while the prediction gets the location of the interaction site right

and correctly predicts the unusual, asymmetrical way the Xre recognition helices bind the DNA, it

does not capture all the subtle details of the deformation of the DNA that take place upon induced

fit (**Fig 5F**). Remarkably, however, the predicted interaction of Xre-RES^{Pp} with DNA at the

secondary binding site located upstream at the S3-S4 interface is strikingly similar to the interaction

to a symmetry-related S4 DNA molecules in the crystal structure. We consider this as strong

support for the validity of the overall prediction given the complementarity of the two techniques

(**Fig 5G**). The agreement between the crystal structure and the *in silico* prediction suggests that the

interaction between the second Xre dimer and the symmetry-related DNA molecules reflect a true

biological ability of the complex to interact with higher-order DNA at two sites simultaneously.

Collectively, this analysis allows us to put forward a detailed and testable model for how the Xre-

RES^{Pp} complex interacts with higher-order promoter DNA via the S3 and S4 repetitive elements.

DISCUSSION

In this paper, we present a detailed model for transcriptional autoregulation of the *P. putida xre-res*

operon. We show that the heterohexameric Xre-RES^{Pp} protein complex efficiently represses

transcription from the *xre-res* locus through direct binding to an imperfect inverted repeat region

(S4) in the promoter region, and the structure of the Xre-RES^{Pp} complex bound to this DNA motif

provides a detailed molecular view of how the promoter is specifically recognised. We also confirm

that the unusual 4:2 stoichiometry of the TA complex observed in the isolated crystal structure [14],

in which each of the toxins of a central dimer interacts with a dimer of antitoxins, is present in

solution, and further show that isolated Xre^{Pp} antitoxin is a monomer that only weakly represses

transcription *in vivo*. This suggests that formation of the Xre dimer, which is stabilised through

interactions with the RES toxin, is necessary for efficient binding to DNA. Supportive of this, a 2:2

complex present at low concentrations does not bind DNA. Together, these data allow us to propose

a model for transcriptional regulation of the *xre-res*^{Pp} operon (**Fig 6**). The expressed 4:2 Xre-RES^{Pp}

complex represses transcription through binding via one Xre dimer to the S4 imperfect repeat,

followed by binding of the second Xre dimer to the region between S3 and S4. This serves to

control transcription and maintain a low cellular concentration of RES toxin. But since the RES-

bound Xre-dimer is unstable, dissociation and formation a 2:2 complex will occur over time. In this

2:2 complex, Xre still binds to, and neutralises, the RES toxin, but it is unable to bind DNA and

consequently, transcription of the *xre-res*^{Pp} operon is eventually reactivated. This increases the

cellular concentrations of the Xre antitoxin, leading to the formation of a repressing DNA-binding

4:2 complex, possibly via the 3:2 intermediate.

435
Figure 6. Model for transcriptional regulation of the *xre-res*^{Pp} operon.
Expression of *xre-res*^{Pp} leads to the formation of a 4:2 Xre-RES^{Pp} complex (A), in which one Xre dimer
binds an imperfect inverted repeat in the promoter region (S4), followed binding of the second Xre dimer
to the region between S3 and S4 and subsequently transcriptional repression (B). This lowers the
concentration of the Xre antitoxin, leading to the dissociation of the Xre dimer (C) and formation of a toxin-
neutralising, but non-DNA binding 2:2 complex (D), and thereby initiation of transcription by the RNA
polymerase.

This model resembles the recently proposed translation-responsive model [30], which states that the

expression of TA systems is primarily regulated by the rate of antitoxin production. This model

assumes that degradation of antitoxin over time is constant and that the net level of antitoxin thus is

controlled by transcriptional and translational activity of the TA operon. Here, we propose that

transcriptional autoregulation is based on the concentration of the Xre antitoxin, which again

depends on Xre expression from the *xre-res*^{Pp} operon. Interestingly, the structure of a Xre-RES

system from *V. parahaemolyticus* (VPA0770-VPA0769) also exists in a heterohexameric (4:2)

form that exhibits a W-shaped architecture like Xre-RES^{Pp} and moreover, the antitoxin was also

found in an equilibrium between a dimer and a monomer [18]. The VPA0770-VPA0769 complex

was found to form a hetero-dodecameric complex (4:8) at high protein concentrations ($\geq$ 5 mg/ml),

which was proposed to enhance the stability of the TA complex and thereby facilitate tighter

regulation of toxin activity. However, at biologically relevant levels ($\leq$ 1 mg/ml), the complex

dissociates into heterohexamers similar to Xre-RES^{Pp}. Additionally, higher-order structural analysis

using PISA [31] indicates that the most stable assemblies in solution are either a 2:2 or a 2:4 form.

These observations suggest that the *VPA0770-VPA0769* system might operate in a manner similar

with the model proposed here. The existence of a dynamic equilibrium between two oligomeric

forms of the Xre-RES complex has also been described for the Xre-RES system NatRT from *P.*

aeruginosa [19], which forms a 4:2 inert state and a 2:1 toxic state and in line with the present

findings, the 4:2 NatRT complex autoregulates transcription of *natR-natT*, while the 2:1 state is

unable to bind the promoter region and thus repress transcription. However, in this case, the

interaction surface between NatR and NatT is mediated by a helix-loop-helix domain (referred to as

the 'Flap') not found in RES^{Pp}, and moreover, the switch between the two oligomeric states of

NarRT is due to disintegration of the NatT toxin dimer, which we do not observe. Finally, the

activity of the NatT toxin affects how the complex binds the promoter, and thus how it regulates

transcription. We have not been able to detect NADase activity of RES^{Pp} when bound to Xre,

suggesting that the RES domain toxins are regulated and activated by different mechanisms. Taken

together, while the details of the mechanisms of Xre-RES systems differ between organisms, the

existence of a dynamic equilibrium between oligomeric states, each playing a unique role in activity

or regulation of gene expression, appears to apply more generally across Xre-RES homologs.

The crystal structure of the Xre-RES^{Pp} complex bound to promoter DNA reveals an unusual

structural asymmetry in the way the Xre HTH recognition helix interacts with DNA when

compared to canonical HTH-containing protein repressors, such as the bacteriophage lambda

Cro repressor [20]. This, together with the preference for the imperfect inverted repeat in S4,

suggests that asymmetry is an important feature for optimal protein-DNA interaction. We note

that the bending of the DNA as well as asymmetrical interaction between Xre and the S4

element optimises the involvement of Arg67. The importance of this is supported by the

reduced binding affinity observed for Xre-RES^{Pp} towards the 5'-perf and 3'-perf S4 variants

in vitro (**Table 2**). In addition, the asymmetry of the two recognition helices involved in DNA

binding ensures specificity as well as directional binding of the protein complex, positioning

the second 'free' Xre dimer to bind a potential secondary operator site. While *in vitro*

functional data show that a single Xre-RES^{Pp} complex binds only one copy of S4 dsDNA,

deletion of the central palindromic motif of S4 (Δrepeat) within the full P_{XR} promoter,

including bases identified by our structural data as critical for specific interaction with Arg67,

still results in partial repression by *xre-res*^{Pp} *in vivo*. This finding suggests the existence of a

second binding site within the full promoter and supportive of this, the AF3 prediction

indicates that the second Xre dimer binds between S3 and S4 in the promoter, in a manner

similar to the observed secondary interaction in the crystal. We believe that this interaction is

highly significant because crystal contacts often reveal important functional aspects of the

crystallised macromolecules. However, these can be hard to validate and separate from mere

packing contacts and likewise, hypothetical interactions in predicted models need validation

before any conclusions can be drawn. Here, the results of two completely independent

techniques (AF3 and x-ray crystallography) strongly support each other in suggesting both the

location and specific mode of interaction at the secondary site.

In conclusion, transcriptional regulation of the *xre-res*[Pp] operon relies on an asymmetrical

interaction of the symmetrical Xre-RES[Pp] protein complex to the operator site, as well as on a

dynamic, concentration-dependent stoichiometry of the Xre-RES[Pp] complex. Together, this

puts forward a detailed and testable model for how bacterial transcription factors interacts

with higher-order promoter DNA (**Fig 6**). This model demonstrates how a bacterial protein

repressor and its operator sites co-evolve to ensure optimal and specific transcriptional

regulation and the presence of two DNA-binding domains allows for a tighter transcriptional

repression through avidity, making the system more robust towards both perturbations and

mutations.

MATERIALS AND METHODS

Bacterial strains and plasmids

Bacterial strains and plasmids used in this study are listed in **S2 Table**, and DNA oligonucleotides

in **S3 Table**. Detailed descriptions of plasmid construction are found in **S1 Text**. For site-directed

mutagenesis, the Q5 Site-Directed Mutagenesis Kit from NEB was used according to the protocol

and mutagenesis primers were designed using NEBaseChanger.

Measurements of promoter activities and *in vivo* repression

For promoter activity assays, *E. coli* MG1655 was co-transformed with transcriptional fusion vector

pGH254Kgfp and a pBAD33 vector. Overnight cultures were grown in M9 medium supplemented

with 0.2% Casamino acids, 0.4% glycerol, 50 mg/ml chloramphenicol, 25mg/ml kanamycin and

0.2% glucose. Overnight cultures were washed once in M9 medium and diluted to 0.03 in M9

medium supplemented with 0.2% Casamino acids, 0.4% glycerol, 50 µg/ml chloramphenicol and

25µg/ml kanamycin. From this, cultures were split into non-induced or induced (by 0.2% arabinose)

sub-cultures. Samples of 200 µl were grown in triplicates in a black/clear bottom 96-well plates

(Thermo Scientific, cat. no. 165305), using a Synergy H1 microplate reader (BioTek). Optical

density (600 nm) and fluorescence intensity (excitation, 479 nm; emission, 520 nm) was measured

every 5 min for 12 h. A strain with empty pGH254Kgfp and pBAD33 was used as control and

fluorescence intensity at a specific time point was calculated as described by [32], using the formula

$I(t) = I_u(t) - \frac{A(t)}{B(t)} I_b(t)$ where $I(t)$ is the fluorescence intensity at a specific time point, $I_u(t)$ is the

uncorrected intensity, $A(t)/B(t)$ is the absorbance ration between the measured culture and the

culture used as blank, and $I_b(t)$ is the fluorescence intensity of the culture used as blank. The

calculated fluorescent intensity was divided by OD_{600} to obtain corrected fluorescence intensity per

OD_{600} unit relative to the control strain (relative fluorescence units, RFU). The assay was repeated

three times for each strain. Mean values were calculated for technical triplicates and used to

calculate a mean-of-mean values and standard mean of error (SEM) of biological triplicates.

Statistical analysis was performed in R using Welch's t-test with the assumption of equal variance.

Graphs were prepared with Prism.

Protein expression and purification

For protein expression, plasmids pETDuet::*res*$_{NHis6}$-*xre*^{Pp}, pETDuet::*xre*$_{NHis6}$-*res*^{Pp}, or pET-

29b(+)::*xre*$_{CHis6}$^{Pp} were transformed into *E. coli* BL21(DE3). Cells were grown to an OD_{600} of 0.6-

0.8, induced by the addition of 0.5 mM IPTG for 20-22 h at 20°C, and harvested by centrifugation

(6,000 rpm, 10 min, 4°C).

For purification of the Xre-RES$^{Pp}_{NHis6}$ and the Xre$_{NHis6}$-RES^{Pp} complex, cell pellets were

resuspended in lysis buffer (50 mM Tris-HCl, pH 8.5, 300 mM KCl, 10 mM imidazole, 3 mM β-

mercaptoethanol (BME)) supplemented with 1 ng/µl DNase I and 1 mM phenylmethylsulfonyl

fluoride (PMSF). Cells were lysed by high-pressure homogenization (Avestin) at 15,000 psi, and

clarified by centrifugation (14,000 rpm, 45 min, 4°C). Clarified cell lysates were loaded onto a 1 ml

HisTrap FF column (Cytiva) and columns were washed with lysis buffer including 40 mM

imidazole. Protein complexes were eluted with elution buffer (50 mM Tris-HCl, pH 8.5, 150 mM

KCl, 300 mM imidazole) and further purified using a 1 ml Source 15Q column (GE Healthcare),

running in buffer A (50 mM Tris-HCl, pH 8.5, 5 mM (BME) and a gradient into buffer B (50 mM

Tris-HCl, pH 8.5, 5 mM BME, 1 M KCl). Final separation was achieved on a Superdex 200

Increase 10/300 GL column (GE Healthcare) equilibrated in gel filtration buffer (20 mM HEPES-

KOH, pH 7.5, 150 mM KCl, 5 mM MgCl$_2$, 3 mM BME).

For purification of Xre alone (Xre$_{CHis6}^{Pp}$), cell pellets were resuspended in lysis buffer (50 mM

HEPES-KOH, pH 7.5, 400 mM KCl, 10 mM imidazole) supplemented with 1 ng/µl DNase I and 1

mM PMSF, lysed by high-pressure homogenization (Avestin) at 15,000 psi, and clarified by

centrifugation (14,000 rpm, 45 min, 4°C). Clarified cell lysate was loaded onto a 1 ml HisTrap FF

column (Cytiva), washed with lysis buffer including 30 mM imidazole and eluted using 300 mM

imidazole. Eluate was precipitated overnight at room temperature with 45% ammonium sulphate

(AmS) and clarified by centrifugation (20.000 rpm, 30 min, 20° C). Supernatant was collected and

further precipitated with 60% AmS overnight. Precipitated protein was collected by centrifugation

(20.000 rpm, 30 min, 20° C) and resuspended in gel filtration buffer (20 mM HEPES-KOH, pH 7.5,

150 mM KCl, 3 mM BME) until conductivity was 15 mS/cm. Final separation was achieved on a

Superdex 75 10/300 GL column (GE Healthcare) equilibrated in gel filtration buffer.

Analytical Size Exclusion Chromatography

To screen the interaction between the Xre-RES^{Pp} complex and putative binding sites within the

promoter, 30 bp dsDNA oligos corresponding to S1, S2, S3 or S4 in the promoter region (**Fig 1A**

and **Fig S1A**) were made using oligos specified in **S3 Table**. Briefly, equimolar amounts of

complementary DNA oligos were mixed in milliQ water heated to 95°C and cooling to room

temperature. Concentrations were determined by nanodrop, using calculated extinction coefficients,

and 1.5 nmoles of dsDNA and/or purified Xre-RES$^{Pp}_{His6}$ each were incubated for 5 min at 37° C.

Samples containing protein alone, DNA alone, or mixed protein and DNA were analysed by

analytical gel filtration (a-SEC) using a Superdex 200 Increase 10/300 GL column (GE Healthcare)

running in DNA binding buffer (20 mM HEPES pH 7.5, 150 mM KCl, 5 mM MgCl$_2$ and 3 mM

BME). This method probes only high strength (low K$_D$) interactions which would be necessary for

transcriptional control. To assess stoichiometry of protein to DNA in the interaction, samples with

protein to DNA ratios of 2:1, 1:2, and 1:10 were analyzed by a-SEC and SEC-Multi angle light

scattering (MALS) in DNA binding buffer. For 1:2 and 1:10 ratios of protein to DNA, a Superdex

200 Increase 3.2/300 GL column (GE Healthcare) was used for increased resolution. Data was

plotted in R (version 4.3.1) using the Tidyverse package [33].

SEC-MALS

For ensemble mass determination, purified protein complex was applied to a size-exclusion column

(SEC) coupled to multi angle light scattering (MALS) in DNA binding buffer (20 mM HEPES pH

7.5, 150 mM KCl, 5 mM MgCl$_2$, 3 mM BME). For each sample, 20-40 µg was added to a Superose

6 increase 10/300 column (GE Healthcare) and MALS spectra was recoded using inline DAWN-

HELEOS light scattering and optilab T-rEX refractive index detectors (Wyatt). Mass was

determined by analyzing the differential refractive index using the Debye model for proteins

integrated in the ASTRA VI software [34]. Data was plotted in R (version 4.3.1) using the

Tidyverse package [33].

Structure determination

Xre-RES$_{Nhis6}$ and S4 dsDNA mixed in a 2:1 molar ratio with a final protein concentration of 4

mg/mL. Triagonal plate-shaped crystals were grown in drops containing 0.1M NA HEPES pH 7.5,

10% (w/v) PEG 8000, 8% ethylenglycol, and 0.2 M NaCl. All crystals were cryo-protected in

reservoir solution supplemented with 20% (w/v) sucrose and flash frozen in liquid nitrogen prior to

data collection. A high-quality diffraction dataset was collected at 0.937Å wavelength on a single

crystal cooled to 100 K at the BioMAX beamline at MAX-IV (Lund, Sweden). The dataset was

processed with XDS [35] and molecular replacement carried out using PHASER[36]. Model

building was done manually in coot with iterative refinement using phenix.refine[37], DNA was

built into the model gradually as density became apparent. During the final stages of refinment,

phenix.refine was complemented by NAMDINATOR [38] for improved fitting and handling of

rotamer outliers. The final model contains 32 chains (24 protein and 8 DNA). Xre is complete in

chains B, E, F, H, I, K, N, O, Q, T, W from residues 1-149 (Xre), with chains C and U missing 149,

chain R missing 147-149, and chans L and X missing 146-149. All RES proteins were missing the

13 N terminal residues from the tag, but were otherwise complete from 1-145 with the exepction of

chains A and J which lacks residue 145. DNA chains a and c were built from residue 8-28 (of 30

total), b and d from 3-23, e from 7-30, f from 1-24, g from 11-30, and h was from 1-20. The final

structure contains, 65 water molecules.

Mass photometry

Single molecule mass determination was carried out using the TwoMP mass photometer (Refeyn).

Experimental data was acquired through one-minute videos recorded by AcquireMP (ver 2.5)

software on precleaned high-precision coverslips with Culterwell gaskets (Grace Biolabs). Focus

was automatically found by AcquireMP and was deemed correct when the variance was 0.01. Prior

to protein analysis, a mass calibration was performed using Bovine Serum Albumin (BSA) and

either Immunoglobulin M (IgM) or α2 macroglobulin (α2M) spanning a mass range of 66 kDa to

720 kDa. Standard samples were collected at final concentrations of 10 nM by 10-fold dilution into

gel filtration buffer. Collected videos were analyzed in DiscoverMP (ver 2.5) by fitting Gaussian

functions to ratiometric contrast values of all events in the video. Mass determination was done by

fitting a linear curve between ratiometric contrast and molar mass of standard proteins. The Xre^{Pp}-

RES^{Pp} complex was diluted 10-fold, 20-fold, or 40-fold in gel filtration buffer (20 mM HEPES pH

7.5, 150 mM KCl, 5 mM $MgCl_2$, 3 mM BME) to achieve final concentrations of 10, 5, and 2 nM,

respectively. Videos were collected as with the standards and mass was determined from the

calibration curve. Statistical significance was defined using a Bonferroni corrected p value of

0.0055, to avoid the multiple comparisons problem. Normality was tested using the Kolmogorov-

Smirnov test, and populations were compared using the Wilcoxon test.

Isothermal titration calorimetry

Thermodynamic constant determination was carried out using the PEAQ-ITC (Malvern

panalytical). Purified protein and dsDNA samples were dialyzed into the DNA binding buffer (20

636 mM HEPES pH 7.5, 150 mM KCl, 5 mM $MgCl_2$, 3 mM BME), after which 20 µM of Xre-

637 RES^{Pp}_{Nhis6} was added to the sample chamber and 300-700 µM dsDNA was titrated in through 13 to

638 17 injections. Each dataset was baseline-corrected by independent measurements of buffer into

639 buffer, titrant into buffer, and buffer into sample. Data analysis was done using Microcal PEAQ-

640 ITC Analysis Software (Malvern panalytical). Data quality was evaluated through minimizing

641 residuals, as well as approximating active concentrations of protein and titrant for fitting to correct

642 N. Reported values are a mean of three independent replicates.

644 **Accession codes**

645 Structure factors and coordinates for the Xre-RES^{Pp}:DNA complex structure have been deposited in

646 the Protein Data Bank with PDB ID 9R35.

648 **ACKNOWLEDGEMENTS**

649 This work was supported by grants from the Independent Research Fund Denmark (grant 0135-

650 00072B to DEB) and the Novo Nordisk Foundation (grants NNF18OC0030646 and

651 NNF22OC0079855 to DEB). The authors would like to thank Janni Nielsen and Prof. Daniel E.

652 Otzen for help with SEC-MALS, Camilla G. Andersen for help with XRD data analysis, Rasmus

653 Freund for scripts designed to plot the pLDDT and PAE files from the Alphafold3 webserver, and

654 Dr. Kristoffer Winther for help with identification of the four promoter repeat sequences. We also

655 wish to thank Associate professor Magnus Kjærgaard and the MBG Biophysics Core Facility for

656 use of equipment supported by The Carlsberg Foundation. We acknowledge MAX IV Laboratory

657 for time on Beamline BioMAX under Proposal 20220062. Research conducted at MAX IV, a

658 Swedish national user facility, is supported by the Swedish Research council under contract 2018-

REFERENCES

1.   Browning DF, Busby SJ. The regulation of bacterial transcription initiation. Nat Rev

Microbiol. 2004;2(1):57-65. doi: 10.1038/nrmicro787. PubMed PMID: 15035009.

2.   Burgess RR, Travers AA, Dunn JJ, Bautz EK. Factor stimulating transcription by

RNA polymerase. Nature. 1969;221(5175):43-6. doi: 10.1038/221043a0. PubMed PMID: 4882047.

3.   Busby S, Ebright RH. Promoter structure, promoter recognition, and transcription

activation in prokaryotes. Cell. 1994;79(5):743-6. doi: 10.1016/0092-8674(94)90063-9. PubMed

PMID: 8001112.

4.   Jacob F, Monod J. Genetic regulatory mechanisms in the synthesis of proteins. J Mol

Biol. 1961;3:318-56. doi: 10.1016/s0022-2836(61)80072-7. PubMed PMID: 13718526.

5.   Aravind L, Anantharaman V, Balaji S, Babu MM, Iyer LM. The many faces of the

helix-turn-helix domain: transcription regulation and beyond. FEMS Microbiol Rev.

2005;29(2):231-62. doi: 10.1016/j.femsre.2004.12.008. PubMed PMID: 15808743.

6.   Harrison SC. A structural taxonomy of DNA-binding domains. Nature.

1991;353(6346):715-9. Epub 1991/10/24. doi: 10.1038/353715a0. PubMed PMID: 1944532.

7.   Branden CI, Tooze J. Introduction to Protein Structure: Garland Science; 2012

2012/3//.

8.   Fernandez-Lopez R, Ruiz R, Del Campo I, Gonzalez-Montes L, Boer DR, de la Cruz

F, et al. Structural basis of direct and inverted DNA sequence repeat recognition by helix-turn-helix

transcription factors. Nucleic Acids Res. 2022;50(20):11938-47. doi: 10.1093/nar/gkac1024.

PubMed PMID: 36370103; PubMed Central PMCID: PMCPMC9723621.

9.   John J, Jabbar J, Badjatia N, Rossi MJ, Lai WKM, Pugh BF. Genome-wide promoter

assembly in E. coli measured at single-base resolution. Genome Res. 2022;32(5):878-92. Epub

20220428. doi: 10.1101/gr.276544.121. PubMed PMID: 35483960; PubMed Central PMCID:

PMCPMC9104697.

10.   Slattery M, Zhou T, Yang L, Dantas Machado AC, Gordan R, Rohs R. Absence of a

simple code: how transcription factors read the genome. Trends Biochem Sci. 2014;39(9):381-99.

Epub 20140814. doi: 10.1016/j.tibs.2014.07.002. PubMed PMID: 25129887; PubMed Central

PMCID: PMCPMC4149858.

11.   Makarova KS, Wolf YI, Koonin EV. Comprehensive comparative-genomic analysis

of type 2 toxin-antitoxin systems and related mobile stress response systems in prokaryotes.

Biology direct. 2009;4:19. Epub 20090603. doi: 10.1186/1745-6150-4-19. PubMed PMID:

19493340; PubMed Central PMCID: PMCPMC2701414.

12.   Freire DM, Gutierrez C, Garza-Garcia A, Grabowska AD, Sala AJ, Ariyachaokun K,

et al. An NAD(+) Phosphorylase Toxin Triggers Mycobacterium tuberculosis Cell Death. Mol Cell.

2019;73(6):1282-91 e8. Epub 20190218. doi: 10.1016/j.molcel.2019.01.028. PubMed PMID:

30792174; PubMed Central PMCID: PMCPMC6436930.

13.   Piscotta FJ, Jeffrey PD, Link AJ. ParST is a widespread toxin-antitoxin module that

targets nucleotide metabolism. Proc Natl Acad Sci U S A. 2019;116(3):826-34. Epub 20181231.

doi: 10.1073/pnas.1814633116. PubMed PMID: 30598453; PubMed Central PMCID:

PMCPMC6338849.

14.   Skjerning RB, Senissar M, Winther KS, Gerdes K, Brodersen DE. The RES domain

toxins of RES-Xre toxin-antitoxin modules induce cell stasis by degrading NAD+. Mol Microbiol.

2019;111(1):221-36. Epub 20181111. doi: 10.1111/mmi.14150. PubMed PMID: 30315706.

15.   Harms A, Brodersen DE, Mitarai N, Gerdes K. Toxins, Targets, and Triggers: An

Overview of Toxin-Antitoxin Biology. Mol Cell. 2018;70(5):768-84. Epub 20180203. doi:

10.1016/j.molcel.2018.01.003. PubMed PMID: 29398446.

16.   Kusumawardhani H, van Dijk D, Hosseini R, de Winde JH. Novel Toxin-Antitoxin

Module SlvT-SlvA Regulates Megaplasmid Stability and Incites Solvent Tolerance in

Pseudomonas putida S12. Appl Environ Microbiol. 2020;86(13):1-17. Epub 20200617. doi:

10.1128/AEM.00686-20. PubMed PMID: 32358012; PubMed Central PMCID: PMCPMC7301842.

17.   Zhou J, Li S, Li H, Jin Y, Bai F, Cheng Z, et al. Identification of a Toxin-Antitoxin

System That Contributes to Persister Formation by Reducing NAD in Pseudomonas aeruginosa.

Microorganisms. 2021;9(4). Epub 20210402. doi: 10.3390/microorganisms9040753. PubMed

PMID: 33918483; PubMed Central PMCID: PMCPMC8065639.

18.   Zhang Y, Song X, Chen C, Liu L, Xu Y, Zhang N, et al. Structural insights of the

toxin-antitoxin system VPA0770-VPA0769 in Vibrio parahaemolyticus. Int J Biol Macromol.

2023;242(Pt 3):124755. Epub 20230508. doi: 10.1016/j.ijbiomac.2023.124755. PubMed PMID:

37164131.

19.   Santi I, Dias Teixeira R, Manfredi P, Hernandez Gonzalez H, Spiess DC, Mas G, et al.

Toxin-mediated depletion of NAD and NADP drives persister formation in a human pathogen.

EMBO J. 2024;43(21):5211-36. Epub 20240925. doi: 10.1038/s44318-024-00248-5. PubMed

PMID: 39322758; PubMed Central PMCID: PMCPMC11535050.

20.   Brennan RG, Matthews BW. The helix-turn-helix DNA binding motif. J Biol Chem.

1989;264(4):1903-6. Epub 1989/02/05. PubMed PMID: 2644244.

21.   Fraikin N, Goormaghtigh F, Van Melderen L. Type II Toxin-Antitoxin Systems:

Evolution and Revolutions. J Bacteriol. 2020;202(7):1-14. Epub 20200311. doi: 10.1128/JB.00763-

19. PubMed PMID: 31932311; PubMed Central PMCID: PMCPMC7167474.

22.   Boggild A, Sofos N, Andersen KR, Feddersen A, Easter AD, Passmore LA, et al. The

crystal structure of the intact *E. coli* RelBE toxin-antitoxin complex provides the structural basis for

conditional cooperativity. Structure. 2012;20(10):1641-8. Epub 20120913. doi:

10.1016/j.str.2012.08.017. PubMed PMID: 22981948; PubMed Central PMCID:

PMCPMC3507626.

23.   Garcia-Pino A, Balasubramanian S, Wyns L, Gazit E, De Greve H, Magnuson RD, et

al. Allostery and intrinsic disorder mediate transcription regulation by conditional cooperativity.

Cell. 2010;142(1):101-11. Epub 2010/07/07. doi: 10.1016/j.cell.2010.05.039. PubMed PMID:

20603017.

24.   Overgaard M, Borch J, Jorgensen MG, Gerdes K. Messenger RNA interferase RelE

controls relBE transcription by conditional cooperativity. Mol Microbiol. 2008;69(4):841-57. Epub

20080604. doi: 10.1111/j.1365-2958.2008.06313.x. PubMed PMID: 18532983.

25.   De Bruyn P, Girardin Y, Loris R. Prokaryote toxin-antitoxin modules: Complex

regulation of an unclear function. Protein Sci. 2021;30(6):1103-13. Epub 20210407. doi:

10.1002/pro.4071. PubMed PMID: 33786944; PubMed Central PMCID: PMCPMC8138530.

26.   Cormack BP, Valdivia RH, Falkow S. FACS-optimized mutants of the green

fluorescent protein (GFP). Gene. 1996;173(1 Spec No):33-8. doi: 10.1016/0378-1119(95)00685-0.

PubMed PMID: 8707053.

27.   Takeda Y, Folkmanis A, Echols H. Cro regulatory protein specified by bacteriophage

lambda. Structure, DNA-binding, and repression of RNA synthesis. J Biol Chem.

1977;252(17):6177-83. doi: 10.1016/S0021-9258(17)40047-0. PubMed PMID: 330523.

28.   Shimon LJW, Harrison SC. The phage 434 OR2/R1-69 complex at 2.5 Å resolution.

Journal of Molecular Biology1993. p. 826-38.

29.   Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure

prediction of biomolecular interactions with AlphaFold 3. Nature. 2024;630(8016):493-500. Epub

20240508. doi: 10.1038/s41586-024-07487-w. PubMed PMID: 38718835; PubMed Central

PMCID: PMCPMC11168924.

30.   Ramisetty BCM. Regulation of Type II Toxin-Antitoxin Systems: The Translation-

Responsive Model. Front Microbiol. 2020;11(May):895. Epub 20200505. doi:

10.3389/fmicb.2020.00895. PubMed PMID: 32431690; PubMed Central PMCID:

PMCPMC7214741.

31.   Krissinel E, Henrick K. Inference of macromolecular assemblies from crystalline

state. J Mol Biol. 2007;372(3):774-97. Epub 20070513. doi: 10.1016/j.jmb.2007.05.022. PubMed

PMID: 17681537.

32.   Jurenas D, Fraikin N, Goormaghtigh F, De Bruyn P, Vandervelde A, Zedek S, et al.

Bistable Expression of a Toxin-Antitoxin System Located in a Cryptic Prophage of Escherichia coli

O157:H7. mBio. 2021;12(6):e0294721. Epub 20211130. doi: 10.1128/mBio.02947-21. PubMed

PMID: 34844426; PubMed Central PMCID: PMCPMC8630535.

33.   Wickham H, Averick M, Bryan J, Chang W, McGowan L, François R, et al. Welcome

to the Tidyverse. Journal of Open Source Software. 2019;4(43):1686-. doi: 10.21105/joss.01686.

34.   Edelman J. The low-temperature heat capacity of solid proteins. Biopolymers.

1992;32(3):209-18. doi: 10.1002/bip.360320303. PubMed PMID: 1581544.

35.   Kabsch W. Xds. Acta Crystallogr D Biol Crystallogr. 2010;66(Pt 2):125-32. Epub

20100122. doi: 10.1107/S0907444909047337. PubMed PMID: 20124692; PubMed Central

PMCID: PMCPMC2815665.

36.   McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ.

Phaser crystallographic software. J Appl Crystallogr. 2007;40(Pt 4):658-74. Epub 20070713. doi:

10.1107/S0021889807021206. PubMed PMID: 19461840; PubMed Central PMCID:

PMCPMC2483472.

37.   Adams PD, Afonine PV, Bunkoczi G, Chen VB, Davis IW, Echols N, et al. PHENIX:

a comprehensive Python-based system for macromolecular structure solution. Acta Crystallogr D

Biol Crystallogr. 2010;66(Pt 2):213-21. Epub 20100122. doi: 10.1107/S0907444909052925.

PubMed PMID: 20124702; PubMed Central PMCID: PMCPMC2815670.

38.   Kidmose RT, Juhl J, Nissen P, Boesen T, Karlsen JL, Pedersen BP. Namdinator -

automatic molecular dynamics flexible fitting of structural models into cryo-EM and

785 crystallography experimental maps. IUCrJ. 2019;6(Pt 4):526-31. Epub 20190627. doi:

786 10.1107/S2052252519007619. PubMed PMID: 31316797; PubMed Central PMCID:

787 PMCPMC6608625.

789
790

791 **TABLES**

792 **Table 1**. **Data collection and refinement statistics (molecular replacement).**

	DNA-bound Xre-RES^{Pp} complex (PDB: 9R35)
Data collection	
Space group	P 1 2$_1$ 1
Cell dimensions	
$\quad$ a, b, c (Å)	81.23, 185.11, 170.79
$\quad$ α, β, γ (°)	90, 90.01, 90
Resolution (Å)	26 – 2.7 (2.77 – 2.70)*
R_{sym} (%)	9.2 (122.7)
I / σI	8.29 (0.85)
Completeness (%)	90.3 (83.4)
Redundancy	3.4 (3.0)
CC½ (%)	99.7 (35.8)
Refinement	
Resolution (Å)	26 – 2.7 (2.77 – 2.70)
No. reflections	146,206 (7846)
R_{work} / R_{free} (%)	22.8 (35.2) / 26.9 (36.5)
No. atoms	31,814
$\quad$ Protein	31,524
$\quad$ Solvent	65
Average B-factors (Å^2)	
$\quad$ Protein	89.1
$\quad$ Solvent	79.8
RMS deviations from ideal	
$\quad$ Bond lengths (Å)	0.004
$\quad$ Bond angles (°)	0.628
Rotamer (%)	
$\quad$ Favoured	99.6
$\quad$ Outlier	0.4
Ramachandran values (%)	
$\quad$ Favoured	98.1
$\quad$ Allowed	1.5
$\quad$ Outlier	0.4

793 *Values in parentheses represent highest-resolution shell.

794 **Table 2. Isothermal Titration Calorimetry (ITC).**

dsDNA	K_D [M]	SEM	Enthalpy (H) [kJ /mol]	Entropy (-ST) [J/mol]	Free energy (G) [kJ/mol]
S4	$2.31 \cdot 10^{-7}$	$4.61 \cdot 10^{-8}$	-3.35 ± 1.29	-31.8 ± 1.66	-38.4 ± 0.71
5'-perf	$3.41 \cdot 10^{-4}$	$2.64 \cdot 10^{-4}$	-5.36 ± 0.92	-26 ± 2.09	-31.3 ± 1.25
3'-perf	$8.06 \cdot 10^{-5}$	$7.97 \cdot 10^{-5}$	9.14 ± 7.28	-42.9 ± 8.06	-33.8 ± 0.91

795 S4 dsDNA, or variants, was titrated into Xre-RES[Pp] complex. The calculated binding affinities (K_D), the enthalpy (H), the entropy with
796 temperature (-ST), and the Gibbs free energy (G) are shown. All values are means of three (n=3) independent replicates with standard error
797 of mean (SEM) indicated.
798

SUPPORTIVE INFORMATION

Fig S1. Determination of the minimal promoter requirements of P$_{XR}$.

Fig S2. Functional analysis of the S4 repeat sequence.

Fig S3. Xre-RES^{Pp} interacts specifically with S4 in a 1:1 protein complex to DNA duplex ratio.

Fig S4. Crystal structure of the DNA-bound Xre-RES^{Pp} complex.

Fig S5. Expression of *xre*^{Pp} alone only partially represses transcription from P$_{XR}$.

Fig S6. Isolation of the 2:2 Xre-RES^{Pp} complex.

Fig S7. Alphafold 3 prediction of the interaction between Xre-RES and the full promoter.

S1 Table. Promoter DNA sequences.

S2 Table. Strains and plasmids used in this study.

S3 Tabel. DNA oligonucleotides used in this study.

S1 Text. Supportive Information Methods.

S1 Appendix. Source Data for Figures and Tables.