Peculiar transcriptional reprogramming with functional impairment of dendritic cells upon exposure to transformed HTLV-1-infected cells.

Auriane Carcone[1], Franck Mortreux[2], Sandrine Alais[1], Cyrille Mathieu[3], Chloé Journo[1,#], Hélène Dutartre[1,#]*

Affiliations:
[1] Centre International de Recherche en Infectiologie, Retroviral Oncogenesis, Inserm U1111 - Université Claude Bernard Lyon 1, CNRS, UMR5308, Ecole Normale Supérieure de Lyon, Université Lyon, Lyon, France.
[2] Laboratory of Biology and Modelling of the Cell, University of Lyon, ENS de Lyon, University Claude Bernard, CNRS UMR 5239, Inserm U1210, Lyon, France.
[3] Centre International de Recherche en Infectiologie, équipe Neuro-Invasion, TROpism and VIRal Encephalitis, Inserm U1111 - Université Claude Bernard Lyon 1, CNRS, UMR5308, Ecole Normale Supérieure de Lyon, Université Lyon, Lyon, France.
# Equal contribution
* Corresponding author

Abstract

Manipulation of immune cell functions, independently of direct infection of these cells, emerges as a key process in viral pathophysiology. Chronic infection by Human T-cell Leukemia Virus type 1 (HTLV-1) is associated with immune dysfunctions, including misdirected responses of dendritic cells (DCs). Here, we interrogate the ability of transformed HTLV-1-infected T cells to manipulate human DC functions. We show that exposure to transformed HTLV-1-infected T cells induces a biased and peculiar transcriptional signature in monocyte-derived DCs, associated with an inefficient maturation and a poor responsiveness to subsequent stimulation by a TLR4 agonist. This poor responsiveness is also associated with a unique transcriptional landscape characterized by a set of genes whose expression is either conferred, impaired or abolished by HTLV-1 pre-exposure. Induction of this functional impairment requires several hours of coculture with transformed HTLV-1-infected cells, and associated mechanisms driven by viral capture, cell-cell contacts, and soluble mediators. Altogether, this cross-talk between infected T cells and DCs illustrate how HTLV-1 might co-opt communications between cells to induce a unique local tolerogenic immune microenvironment suitable for its own persistence.

Significance (120words)
Chronic viral infection is associated with an escape from immune surveillance. This may rely on the induction of inappropriate DC responses, which can contribute to immunopathology. Immune dysfunctions have been repeatedly reported in people living with Human T-cell Leukemia Virus type

1 (HTLV-1), years before fatal clinical symptom onset, including misdirected responses of dendritic
cells (DCs). Here, we report that HTLV-1-infected T cells actively manipulate neighboring, uninfected
MDDC functions by rewiring their transcriptional response, leading to a biased, pro-tolerogenic
responsiveness in MDDCs, induced by the bidirectional release of soluble mediators, in cooperation
with mechanisms dependent on cell-cell contacts. This cross-talk illustrate how HTLV-1 might co-opt
communications between cells to induce a local tolerogenic immune microenvironment suitable for its
own persistence

Introduction
Chronic infection by the deltaretrovirus Human T-cell Leukemia Virus type 1 (HTLV-1)
generally remains asymptomatic, but leads to severe diseases in 5 to 10% of infected
individuals[1], in the form of Adult T-cell Leukemia/Lymphoma (ATL), or Tropical Spastic
Paraparesis/HTLV-1 Associated Myelopathy (TSP/HAM). Remarkably, HTLV-1 chronic
infection increases the risk of viral, bacterial or parasitic co-infections[2]. In agreement with
these clinical observations, immune dysfunctions have been repeatedly reported in
asymptomatic HTLV-1 carriers, years before severe diseases were diagnosed, indicating that
chronic infection induces systemic effects and impairs the ability of the immune system to
respond appropriately to infectious and non-infectious triggers[3–7]. Dendritic cells (DCs) are
critical mediators of innate and adaptive immune responses during viral infection. Upon
efficient viral recognition, DCs undergo a maturation program that renders them capable of
inducing specific T-cell responses. The DC maturation process is characterized by a
transcriptional program that translates, among others, in the upregulation of several
maturation markers at the cell surface (*e.g.* CD80, CD86), accompanied by the secretion of
pro-inflammatory cytokines such as tumor necrosis factor alpha (TNF-α), and antiviral
cytokines such as type I interferons (IFN-I)[8]. However, inappropriate DC responses can
contribute to immunopathology, a process that is exacerbated in chronic viral infections[9].
The role of DCs in HTLV-1 infection has been extensively investigated in terms of viral
transmission to CD4[+] T cells, both *in vitro*[10–12] and in animal models[13,14]. Indeed, myeloid
cells such as monocytes and DCs can harbor infectious HTLV-1 genomes in infected humans,
including in asymptomatic individuals[6,16]. They exhibit impaired phenotypes and functions[16–
18] that may correlate with their proviral load[19]. These clinical observations indicate that
HTLV-1 could in fact dampen the responsiveness of infected DCs, which may ultimately
contribute to immune dysfunction during chronic infection. Accordingly, DCs derived from
ATL patients exhibit maturation defects[18] associated with an inability to induce proliferation
of CD4[+] T cells[17]. However, because the proportion of DCs infected *in vivo* is low, the impact
of HTLV-1 on DC function might not be solely attributed to HTLV-1-infected DCs,
suggesting that the functions of non-infected DCs could be also modulated indirectly by
HTLV-1 infection. In this work, we aimed at determining *in vitro* whether transformed
HTLV-1-infected T cells could affect neighboring human DC functions in the absence of their

infection. We show that upon coculture with HTLV-1-transformed T cell lines, human
monocyte-derived DCs (MDDCs) do not fully mature. This impaired maturation is associated
with a unique transcriptional signature in MDDCs, which differs from a typical maturation
program, and is correlated with a dampened responsiveness of HTLV-1-exposed MDDCs to
subsequent stimulation by TLR-4 or TLR-7 agonists. Interestingly, this tolerogenic state can
also be induced by exposing freshly derived MDDCs for several hours to the supernatant from
MDDCs cocultured with HTLV-1-infected T cells for several hours, highlighting a cross-talk
between MDDCs and T cells based on a bidirectional release of soluble mediators. This
unique mechanism of viral-induced tolerance of DCs illustrates how HTLV-1 might co-opt
communications between cells to induce a local immune microenvironment suitable for its
own persistence.
99
100
101
102
103   **Results**
104
Human MDDCs do not fully mature when exposed to HTLV-1-infected T cells
To determine whether chronically infected T cells could directly or indirectly manipulate
human DC functions *in vitro*, we first addressed whether infected T cells were sensed by DCs.
We used human monocyte-derived DCs (MDDCs) cultured *in vitro* with HTLV-1-infected
and transformed T cell lines, and assessed MDDCs maturation by flow cytometry after 24h or
48h of coculture, using the MDDC-specific CD11c marker to identify MDDCs in the
coculture (see **Fig. 1a** for the experimental settings, **and Supplementary Fig. 1** for the gating
strategy). At these time points, we have previously shown that less than 1% of MDDCs have
integrated the viral genome but they do not release any virion[11]. In contrast to MDDCs
exposed to measles virus (MeV), used as a pathogen known to induce MDCC maturation[20,21],
MDDCs exposed to HTLV-1 transformed C91-PL cells, failed to fully upregulate CD86, a
surrogate marker of DC maturation (**Fig. 1b, left panel**). Repeated experiments using MDDC
samples from independent donors showed that while around 90% of MDDCs upregulated
CD86 upon exposure to MeV, regardless of being infected or not by MeV (**Fig. 1b, right**
panel, compare total versus GFP[-] and GFP[+] MDDCs), as well as upon stimulation by the
TLR-4 agonist LPS (**Fig. 1b, right panel**), the percentage of activated MDDCs after
coculture with HTLV-1-transformed C91-PL cells remained low (**Fig. 1b, right panel**). To
exclude a cell line-specific effect, and to thoroughly characterize MDDCs maturation status,
MDDCs were exposed to several control uninfected T cell lines (**Supplementary Fig. 2a-f**;
Jurkat, CEM or Molt-4 T cell lines, green bars), as well as to several HTLV-1-transformed T
cell lines (**Supplementary Fig. 2a-f**; C91-PL, MT-2 or Hut102; blue bars); and the regulation
of both maturation markers CD86, CD83, CD80, and CD40, (**Supplementary Fig. 2a-d**) and
inhibition markers ICOSL and PD-L1 (**Supplementary Fig. 2e-f**) at the cell surface was
monitored by flow cytometry. All the HTLV-1-transformed T cell lines induced either no
change, or only minor changes, in the expression (percentage of positive cells and normalized
mean signal intensity, MFI) of maturation or inhibition markers at the surface of cocultured
MDDCs (**Supplementary Fig. 2a-f**) even after sustained coculture for 48 h (**Supplementary**

Fig. 2a-f), excluding a possible delay in maturation. The inefficient upregulation of MDDC
maturation markers was also accompanied by a low ability to secrete TNF-α or IFN-I (**Fig. 1c**
and **Supplementary Fig. 2g**). Thus, HTLV-1-infected T cells are unable to induce a typical
maturation program in MDDCs, as if they were not efficiently sensed by human DCs. Since
similar observations were made with all HTLV-1-infected T cell lines tested, and with the
tested maturation markers and cytokines, the following experiments were done using the C91-
PL T cell line and CD86 as DC maturation marker.
From our previous work, MDDCs efficiently capture HTLV-1 after coculture with HTLV-1-
infected T cell lines[11], suggesting that in our experimental conditions, HTLV-1 capture itself
is inefficient in driving MDDC maturation. To formally demonstrate this, we assessed HTLV-
1 capture in MDDCs following coculture, by staining the HTLV-1 Gag p19 structural protein.
As expected, MDDCs were efficient at capturing HTLV-1 (**Fig. 1d**), irrespective of the
infected T cell line used in the coculture (**Supplementary Fig. 2h**), with around 30% of
capture observed in a representative coculture experiment with C91-PL cells (**Fig. 1d, left**),
and reaching up to 80% of capture observed in repeated experiments using independent
donors   (**Fig.   1d,   **right**).

148
Figure 1. Human MDDCs do not fully mature when exposed to HTLV-1-infected T cells. a. Schematic
representation of the experimental design. From day 0 (d0), monocytes were differentiated into MDDCs by a 5-
day culture in the presence of IL-4 and GM-CSF. At day 5 (d5), MDDCs were cocultured with control

uninfected or HTLV-1-infected T cell lines for 24h (or 48h, see Supplementary Fig. 2a-f). Alternatively,
MDDCs were stimulated with GFP-expressing MeV or LPS (control stimuli) for 24h (or 48h, see Supplementary
Fig. 2a-f). Cells and culture supernatants were collected at d6 (or d7, see Supplementary Fig. 2a-f). **b.** Flow
cytometry analysis at d6 after CD11c and CD86 staining. The percentage of CD86$^+$ MDDCs (indicated gate)
among total CD11c$^+$ MDDCs was quantified. **Left**: representative experiment. **Right**: results from n=3
independent experiments, analyzed with RM one-way ANOVA as described in Supplementary Table 7. **c.**
Supernatant from the indicated cocultures or from LPS-stimulated MDDCs was collected at d6, and TNF-α (**top**)
and IFN-I (**bottom**) were quantified by Luminex and reporter cell assay, respectively. Results from n=3 or 8
independent experiments, respectively. Presented data are a subset of Supplementary Figure 2g analyzed with
RM one-way ANOVA (top) or Kruskal-Wallis test (bottom) as described in Supplementary Table 7&8. **d.** Viral
capture in MDDCs cocultured with Jurkat or C91-PL T cells was assessed at d6 by Gag p19 staining on CD11c$^+$
MDDCs. The percentage of Gag p19$^+$ and Gag p19$^-$ MDDCs among total CD11c$^+$ MDDCs was determined.
Left: representative experiment. **Right**: results from n=14 independent experiments analyzed with Kruskal-
Wallis test as described in Supplementary Table 7&8 as a subset of Supplementary Figure 2h. **e.** Flow cytometry
analysis after CD11c, Gag p19 and CD86 staining. **Left**: representative experiment. Dashed lines: LPS-exposed
MDDCs. Solid black lines: Jurkat-exposed MDDCs, light blue line: Gag p19$^-$ MDDCs. Magenta lines: Gag p19$^+$
MDDCs. **Right**: the same color code is used to represent results from n=15 independent experiments analyzed
with one-way ANOVA as described in Supplementary Table 7. **f.** Schematic drawing summarizing the results
from Fig. 1 and Supplemental Fig. 2. The drawing was created using BioRender.com

Maturation was then compared in MDDCs that had not captured HTLV-1 (Gag p19-negative
MDDCs, cyan, **Fig 1e**), or had captured HTLV-1 (Gag p19-positive MDDCs, magenta, **Fig.**
1e). Although CD86 expression was consistently higher in p19-positive MDDCs (**Fig. 1e**, left,
compare the two histograms and **right**, compare cyan and magenta bars), the percentage of
CD86 positive Gag p19-positive MDDCs remained low compared to fully matured LPS-
exposed MDDCs (**Fig. 1e**, right compare magenta and red bars), highlighting that maturation
was restricted even in MDDCs that had captured HTLV-1. No significant correlation was
detected among repeated experiments between the level of HTLV-1 capture and the efficiency
of MDDC maturation (**Supplementary Fig. 2i**, p = 0.1412), further confirming that HTLV-1
capture itself is inefficient in driving MDDC maturation.
Taken together, these observations, recapitulated in **Fig. 1f**, demonstrate that cell-associated
HTLV-1 fails to fully mature MDDCs, suggesting that it is poorly sensed by these human
innate cells.

Exposure to HTLV-1-infected T cells induces a unique transcriptional signature in
MDDCs
In contrast to MDDCs, we previously showed that plasmacytoid dendritic cells (pDC) do
efficiently sense cell-associated HTLV-1[22]. Inefficient sensing of HTLV-1-infected T cells by
MDDCs could then be the result of either a stealth behavior of cell-associated virus towards
MDDCs, resulting in a *bona fide* lack of response specifically in this cell type, or,
alternatively, to the active manipulation of MDDC functions by HTLV-1-infected T cells,
resulting in a specific response that differs from a typical maturation program. To
discriminate between both, we aimed at determining the complete transcriptomic landscape of
MDDCs after exposure to HTLV-1-infected T cells, or to control uninfected T cells, using
bulk RNA-seq (**Fig. 2**). After 24h of coculture, MDDCs were magnetically separated from
C91-PL or control Jurkat T cells, based on the exclusive expression of CADM1 on T cells

(**Supplementary Fig. 3a**). Note that although the levels of CADM1 were higher at the surface
of C91-PL cells compared to Jurkat cells (**Supplementary Fig. 3a**), a similar MDDC
enrichment yield of around 99 % was observed after separation in both conditions, with no
significant T cell contamination (**Supplementary Fig. 3b**). LPS-stimulated MDDCs were
also included in the RNA-seq analysis, as a reference for fully matured MDDCs. RNA was
extracted from each MDDC sample, and submitted to quality control followed by sequencing.
First, to show global similarities in gene expression between samples, unsupervised
hierarchical clustering and principal component analysis (PCA) were performed on
normalized transcript count tables. Analysis of sample clustering along the two first PCs (**Fig.
2a,** PC1 and PC2), that explain 81 and 8% of the total variance, respectively, showed that the
main source of variance in the dataset was the stimulation by LPS, followed by the infection
status of cocultured T cells, while the identity of the MDDC donor contributed only weakly to
the global variance. In agreement with flow cytometry data shown in Fig. 1, this confirmed, at
the transcriptional level, the inefficient activation of MDDCs after exposure to HTLV-1-
infected T cells, which contrasts with the typical maturation program of fully activated
MDDCs. This also indicates that exposure to HTLV-1-infected T cells does induce a
detectable, yet subtle, transcriptional response, when compared to exposure to uninfected T
cells.
To characterize the changes in gene expression in MDDCs induced by exposure to HTLV-1-
infected T cells, we used DESeq2 to identify differentially expressed genes (DEGs) between
C91-PL- and Jurkat-exposed MDDCs. As a reference for a typical maturation program, we
identified DEGs between LPS- and Jurkat-exposed MDDCs. Using a fold-change cut-off of 2,
a total number of 474 DEGs were obtained between C91-PL- and Jurkat-exposed MDDCs
(padj<0.05), with 435 genes found significantly upregulated upon HTLV-1 exposure, and 39
genes found significantly downregulated (**Fig. 2b** and **Supplementary Table 1**). A heatmap
of the 474 DEGs showed clustering between samples, as expected (**Supplementary Fig. 4a**).
To gain insight into the processes modulated in MDDCs after exposure to HTLV-1-infected T
cells, we performed gene ontology analysis on the set of 435 upregulated genes. Over-
represented KEGG pathways were retrieved and filtered to reduce redundancy (see
Supplementary Table 2), leading to a final list of 7 pathways (**Fig. 2c** and **Supplementary
Tables 2 and 3**). Strikingly, exposure to HTLV-1-infected T cells was characterized by
significant over-representation of upregulated genes involved in lipid biosynthesis and
metabolism (**Fig. 2c**). Interestingly, genes involved in several pathways linked with response
to viral infection were also over-represented, including genes involved in innate immune
sensing (sub-pathways: RIG-I or Toll-like receptors and NOD-like receptors), in viral
infection (sub-pathways: Influenza A, Measles, Hepatitis C, Hepatitis B, Human
Papillomavirus, Herpes simplex virus 1, Coronavirus, HIV, EBV), and in the NF-κB
signaling pathway. Taken together, these results confirm that exposure to HTLV-1-infected T
cells does induce a transcriptional response in MDDCs, indicating that sensing does occur to
some extent, but does not culminate in a typical maturation and anti-viral program.

239
Figure 2. Exposure to HTLV-1-infected T cells induces a unique transcriptional signature in MDDCs.
MDDCs were cocultured for 24h with control Jurkat cells, or with HTLV-1-infected C91-PL cells, or with
control Jurkat cells in the presence of LPS. Cells were then separated and RNA-seq analysis was performed on
MDDCs in 3 independent replicates. **a.** Principal component analysis of global similarities in gene expression
between samples (DC_ JK: MDDCs after coculture with Jurkat cells, green; DC_JK_LPS: MDDCs after
coculture with Jurkat cells in the presence of LPS, red; DC_C91: MDDCs after coculture with C91-PL cells,
blue). **b.** Volcano plot of DEGs between MDDCs exposed to C91-PL and MDDCs exposed to Jurkat cells.
c. KEGG pathways analysis performed on the 435 upregulated DEGs. Bars represent the fold-enrichments, while

 dots represent the p-values expressed as log10 for each pathway, or their respective mean for grouped pathways
249 (see Supplemental Table 2). **d.** Venn diagram of the DEGs in C91-PL-exposed MDDCs compared to Jurkat-
250 exposed MDDCs (blue circle), and of the LPS-modulated DEGs in Jurkat-exposed MDDCs (red circle). **e-f.**
251 Heatmap of the 373 shared (**e**) and 101 unique (**f**) DEGs. Gene expression was normalized per row. **g.** DESeq2
252 normalized counts for *ELOVL3*, *FADS1*, and *SLC27A6* across samples.
253

To further determine the extent to which this transcriptional response is distinct from a typical
maturation signature, we compared the genes affected by LPS stimulation of Jurkat-exposed
MDDC to those affected by MDDC exposure to C91-PL. Among the 474 DEGs, 373 genes
were also differentially expressed after LPS stimulation (**Fig. 2d**), representing only 5% of all
LPS-modulated genes. Among these shared genes, 7 genes annotated as involved in the NF-
κB signaling pathway were retrieved, including *CCL19, TNFSF13B, TRIM25, BCL2L1* and
BCL2A1 (**Supplementary Fig. 4b**). However, the level of up-regulation of these shared NF-
κB-related genes was strikingly lower after HTLV-1 exposure compared to LPS stimulation
(**Supplementary Fig. 4b**). This observation of a lower magnitude of differential expression
(either up- or downregulation) was general to most of the 373 shared DEGs (**Fig. 2e**). Of note,
the *RELA, RELB* and *NFKB1* genes that were upregulated by LPS stimulation were not
upregulated upon C91-PL exposure (**Supplementary Fig. 4c**), possibly contributing to the
lower activation of the NF-κB-dependent genes observed above (**Supplementary Fig. 4b**). In
addition, among the 377 Interferon-Stimulated Genes (ISG) listed in **Supplementary Table
1**, only 121 were retrieved in the set of significantly upregulated genes, and again, their up-
regulation was lower than that induced by LPS stimulation (**Supplementary Fig 4d**). These
observations strengthen the notion that while sensing of HTLV-1-infected T cells does occur
to some extent, the magnitude of the maturation and antiviral transcriptional response remains
very limited, which is consistent with the inefficient maturation of MDDCs observed at the
protein expression level (see **Fig. 1**).
Interestingly, among the 474 DEGs, 101 genes were not present in the list of DEGs between
LPS- and Jurkat-exposed MDDCs (**Fig. 2d and f**), defining a unique transcriptional signature
associated with the exposure to HTLV-1-infected C91PL cells. Due to the limited number of
genes included in this specific signature, gene ontology analysis was not feasible.
Interestingly however, several genes retrieved in Fig. 2c as those involved in lipid
biosynthesis and metabolism, were also included in this specific signature, such as *ELOVL3*
(Elongation of Very Long Fatty Acid Elongase 3), *FADS1* (Fatty Acid Desaturase 1), and
SLC27A6 (Solute Carrier Family 27 Member 6, a member of the fatty acid transport protein
family) (**Fig. 2g**). Altogether, these transcriptomic analyses show that exposure to HTLV-1-
infected T cells is not completely silent in MDDCs, but rather results in a unique
transcriptional response that differs from a typical maturation program. This supports the
notion that HTLV-1-infected T cells might actively manipulate MDDC functions to limit their
functional response, possibly by rewiring lipid biosynthesis and metabolism.
287
288

**Pre-exposure to HTLV-1-infected T cells dampens the responsiveness of MDDCs to
subsequent stimulation**
We next aimed at investigating whether this unique transcriptional response observed upon
exposure to HTLV-1-infected T cells indeed translates into functional defects in MDDCs,

beyond their inefficient maturation. To this end, we addressed whether pre-exposure to
HTLV-1-infected cells influenced MDDC responsiveness when exposed to a subsequent
stimulation. After 24h of coculture with HTLV-1-infected T cell lines, MDDCs were
restimulated with strong inducers of MDDCs maturation, in the form of LPS (TLR-4 ligand)
or R848 (TLR-7/8 ligand, **Fig. 3a**), for another 24 h. When compared to MDDCs pre-exposed
to uninfected control T cell lines (red bars), MDDCs pre-exposed to HTLV-1-transformed T
cells (dark blue bars) upregulated CD86 (as well as other maturation markers) to a
significantly lower extent when stimulated by LPS or R848 (**Fig. 3b, Supplementary Fig.**
5a-d). In addition, TNF-α secretion by MDDCs upon LPS stimulation was also significantly
reduced by pre-exposure to HTLV-1-transformed T cells (**Fig. 3c**, left panel, **Supplementary**
Fig. 5e), while IFN-I secretion was not affected (**Fig. 3c**, right panel, **Supplementary Fig.**
5e). These results indicate that pre-exposure to HTLV-1-transformed T cells indeed
influences the responsiveness of MDDCs to subsequent stimulation, by specifically
dampening their pro-inflammatory response (monitored here through the upregulation of
maturation markers and through TNF-α secretion), without hampering their antiviral response
(monitored here through IFN-I secretion) (recapitulated in **Fig. 3d**). This suggests that HTLV-
1 might specifically manipulate the responsiveness of certain signaling pathways in MDDCs,
such as the NF-κB pathway upstream of the pro-inflammatory program, while leaving the
responsiveness of others unaffected, such as the IRF3 pathway upstream of the antiviral
program.
313

314
Figure 3. Pre-exposure to HTLV-1-infected T cells dampens the responsiveness of MDDCs to subsequent
stimulation. a. Schematic representation of the experimental design. MDDCs were exposed to uninfected
(Jurkat, green) or HTLV-1-infected T cells (C91-PL, blue) for 24h, before restimulation with LPS or R848 for an
additional 24h. **b.** Flow cytometry analysis after CD11c and CD86 staining, in LPS- (**left**) or R848-restimulated
MDDCs (**right**) pre-exposed to Jurkat (red) or C91-PL (dark blue) cells. Data are represented as the normalized
MFI of CD86, with the MFI in restimulated Jurkat-pre-exposed MDDCs set to 100 (n=30 or 10 independent
experiments, respectively). Presented data are a subset of Supplementary Figure 5a or 5d, respectively and were
analysed with Kruskal-Wallis test or ordinary one-way ANOVA, respectively as described in Supplementary
Table 7&8 **c.** Supernatant from the indicated cocultures was collected after LPS restimulation, and TNF-α (**left**)
and IFN-I (**right**) concentrations were quantified for n=3 or 10 independent experiments, respectively. Presented
data are a subset of Supplementary Figure 5e and were analysed with RM one-way ANOVA or ordinary one-
way ANOVA, respectively as described in Supplementary Table 7&8. **d.** Schematic drawing summarizing the
results from Fig. 2 and Supplemental Fig. 5. The drawing was created using BioRender.com.

328
329

Pre-exposure to HTLV-1-transformed T cells alters the transcriptional response of
MDDCs to subsequent stimulation
To obtain a broader overview of how pre-exposure to HTLV-1-transformed T cells affects the
responsiveness of MDDCs, a second bulk RNA-seq analysis was conducted on MDDCs pre-
exposed to an HTLV-1-infected T cell line (C91-PL), or to a control uninfected T cell line
(Jurkat), and then re-stimulated by LPS (**Fig. 4**). To address whether viral capture was
required for this manipulation of MDDC responsiveness, we also included in the RNA-seq
analysis MDDCs cocultured with the HTLV-1-infected C8166 T cell line that does not

produce viral particles[23]. As expected, no viral particle was captured by C8166-exposed
MDDCs, in contrast to C91-PL-exposed MDDCs (**Supplementary Fig. 6**).
PCA analysis of samples showed that both the donor and the infection status of cocultured T
cells were the main sources of variance in the dataset (**Fig. 4a**), confirming that pre-exposure
to HTLV-1-transformed T cells does alter the transcriptional response of MDDCs to
subsequent stimulation. In contrast, the ability to capture HTLV-1 did not contribute to a
visible extent to the global variance, suggesting that viral capture might not be required for
this alteration.
We then retrieved the lists of genes differentially expressed between the different
experimental conditions. In agreement with the PCA analysis, 1324 genes were found
differentially expressed upon LPS stimulation between C91-PL-pre-exposed MDDCs, and
Jurkat-pre-exposed MDDCs (**Supplementary Table 4**), with a total of 403 DEGs being
downregulated and 921 DEGs being upregulated (**Fig. 4b**). Gene ontology analysis was
conducted on these sets of downregulated and upregulated genes, respectively (**Fig. 4c**, left
and right panels, respectively). The set of downregulated genes was characterized by a
significant over-representation of genes annotated as being involved in TNF-α or NF-κB
signaling and innate immune sensing (**Fig. 4c**, left panel**,** arrows, see **Supplemental Table 5**).
Conversely, the set of upregulated genes was characterized by a significant over-
representation of genes annotated as being involved in the TGF-β signaling pathway (**Fig. 4c**,
right panel, arrow, see **Supplemental Table 6**). This confirmed that pre-exposure to HTLV-
1-transformed T cell does influence the transcriptional response to LPS stimulation. To
further identify the expression patterns of these genes differentially expressed between C91-
PL-pre-exposed MDDCs, and Jurkat-pre-exposed MDDCs upon LPS stimulation, we
classified these genes based on their differential expression across conditions (detailed in
Supplemental Fig. 7a, see also **Supplementary Table 4)** as follows: (i) Genes whose
responsiveness to LPS (be it repression or induction) is specifically conferred by pre-exposure
to infected cells (**Fig. 4d**, upper panel). (ii) Genes whose responsiveness to LPS (be it
induction or repression) is exacerbated by pre-exposure to infected cells (**Fig. 4d**, middle
panel). (iii) Genes whose responsiveness to LPS is attenuated or (iv) abolished by pre-
exposure to infected cells (**Fig. 4d**, lower panel). This classification reveals that pre-exposure
to HTLV-1-transformed T cells induces both a change in the identity of genes that
transcriptionally respond to LPS stimulation, defining a unique LPS-induced transcriptional
signature; and a change in the magnitude of the transcriptional response of genes that are
normally responsive to LPS.
372
In line with the gene ontology analysis presented in **Fig. 4c**, the responsiveness of genes
involved in NF-κB signaling, such as *MYD88* (**Fig. 4d**, lower panel, see graph), or the pro-
inflammatory genes *CCL2*, *IL12B*, *TNF*, *CXCL10* and *CXCL11* (**Supplementary Fig. 7b**,
upper panel), was found drastically (around 10x) attenuated by pre-exposure to infected cells
(C91-PL or C8166). This could account for the inefficient maturation and production of pro-
inflammatory cytokines by HTLV-1-pre-exposed MDDCs after LPS stimulation, which was
observed in Fig. 3. In addition, in line with the gene ontology analysis presented in **Fig. 4c**,
genes of the TGF-β signaling pathway, including *BMP6, BMP7,* and *IL13* (**Supplementary
Fig. 7b**, middle panel), which do not respond to LPS in normal conditions, were found to be

responsive to LPS when MDDCs were pre-exposed to HTLV-1-transformed cells, while the
expression of *TGFB2*, which is repressed upon LPS stimulation in normal conditions, was
maintained by pre-exposure to HTLV-1-infected cells (**Fig. 4d**, lower panel, see graph).
Interestingly, these genes encode cytokines known to participate in the induction of a
tolerogenic immune microenvironment, with Treg and T_H2 responses, which are inefficient at
controlling viral infection. Finally, DCs pre-exposed to HTLV-1-transformed cells strongly
expressed *DDIT4* upon LPS stimulation (**Fig. 4d**, upper panel, see graph). This gene has been
reported in other contexts of tolerogenicity[24]. Of note, and in agreement with the efficient
induction of IFN-I after LPS stimulation in both Jurkat- or C91-PL-pre-exposed MDDCs (see
Fig. 3c), the induction level of *ISGs* such as *ISG15, IFI44* and *IRF2* was not affected by pre-
exposure to HTLV-1-infected cells (**Supplementary Fig. 7b**, lower panel).
393
Altogether, this transcriptomic analysis indicated that pre-exposure to HTLV-1-transformed T
cells influences the responsiveness of MDDCs to subsequent stimulation, by specifically
dampening their pro-inflammatory response at the transcriptional level. In addition, it
uncovered the fact that pre-exposure to HTLV-1-transformed T cells allows a unique set of
genes to respond to LPS stimulation, triggering a biased, pro-tolerogenic response of MDDCs
upon subsequent stimulation.
400

Figure 4. Pre-exposure to HTLV-1-infected T cells alters the transcriptional response of MDDCs to subsequent stimulation. MDDCs were cocultured for 24h with Jurkat cells, or with HTLV-1-infected C91-PL

or C8166 cells, before restimulation with LPS for an additional 24h. Cells were then separated and RNA-seq
analysis was performed on MDDCs in 3 independent replicates. **a.** Principal component analysis of global
similarities in gene expression between samples (DC_JK_LPS: Jurkat-pre-exposed MDDCs after LPS
restimulation, red; DC_C91_LPS: C91-PL-pre-exposed MDDCs after LPS restimulation, pink, DC_C81_LPS:
C8166-pre-exposed MDDCs after LPS restimulation, purple). **b.** Volcano plot of DEGs between C91-PL-pre-
exposed MDDCs after LPS restimulation, and Jurkat-pre-exposed MDDCs after LPS restimulation. **c.** KEGG
pathways analysis performed on the 403 downregulated DEGs (**left**) or on the 921 upregulated DEGs (**right**).
Bars represent the fold-enrichments, while dots represent the p-values expressed as log10 for each pathway, or
their respective mean for grouped pathways. **d.** Heatmaps and DESeq2 normalized counts across samples of
illustrative genes, showing expression across samples of genes belonging to the "conferred" (**top**), "exacerbated"
(**middle**) or "attenuated or abolished" (**bottom**) responsiveness pattern.
415
416

Neither HTLV-1 viral capture, nor cell-cell contact with infected T cells, are strictly
required to dampen the responsiveness of MDDCs to subsequent stimulation
As stated above, RNA-seq analysis using the "non virus producer" C8166 infected cell line
suggests that viral capture might not be required to alter the transcriptional response of
MDDCs to a secondary stimulation. To confirm this notion at the functional level, we
repeated the coculture experiment followed by LPS stimulation, and analyzed MDDC
maturation profile by flow cytometry (**Fig. 5a**). Despite the lack of viral particle capture by
C8166-exposed MDDCs (see **Supplementary Fig. 6**), C8166 pre-exposure still dampened the
responsiveness of MDDCs to LPS and R848 stimulation, as monitored through CD86, CD83
or CD80 upregulation, similar to C91-PL pre-exposure (**Fig. 5a and Supplemental Fig. 8a**),
confirming that viral capture is not strictly required. Although C8166 do not produce viral
particles, they might still engage in cell-cell contacts with MDDCs[25], which could be the
trigger of the dampening of MDDC responsiveness. We addressed this hypothesis by
performing the cocultures in transwells, in which MDDCs were physically separated from
infected T cells by a permeable membrane (**Fig. 5b**). Of note, reduced but detectable levels of
viral capture were detected in MDDCs physically separated from C91-PL (**Supplementary**
Fig. 8b), most probably because the virus is preferentially associated to the cell surface of
infected cells and could be released as large adhesive viral aggregates[26] poorly able to cross
the 0.4µm pores of the permeable membrane. The absence of physical contact between
MDDCs and infected T cells, either producing (C91-PL) or not producing viral particles
(C8166), did not restore a fully efficient maturation after LPS stimulation (**Fig. 5b**), indicating
that cell-cell contacts are not strictly required to allow HTLV-1-infected T cells to manipulate
MDDC responsiveness. Of note however, C8166 cells appeared less efficient than C91PL
cells in dampening MDDC responsiveness upon transwell coculture. This could result from
additive effects of mechanisms dependent on viral capture on the one hand, and of cell-cell
contact on the other hand: indeed, in the presence of cell-cell contacts (**Fig. 5a**), the
contribution of viral capture might be negligible; while in the absence of cell-cell contacts
(**Fig. 5b**), such contribution would become detectable. Consistently, comparison of MDDC
responsiveness in paired experiments of MDDC pre-exposed to infected cells with (coculture)
or without (transwell) physical contacts showed that preventing physical contacts between
MDDCs and infected cells resulted in a slightly higher, yet not significantly different, MDDC
responsiveness (**Fig. 5c**). This indicates that viral capture and/or cell-cell contacts may
participate, but are not strictly required to allow HTLV-1-infected T cells to manipulate

MDDC responsiveness, and suggests the contribution of a distantly acting set of soluble
mediators.
452

Figure 5. Neither HTLV-1 viral capture, nor cell-cell contact with infected T cells, are strictly required to dampen the responsiveness of MDDCs to subsequent stimulation. a-b. MDDCs were cocultured for 24h with Jurkat cells, or with HTLV-1-infected C91-PL or C8166 cells, in the absence (**a**) or presence (**b**) of a transwell insert (0.4µm pore diameter), before re-stimulation with LPS for an additional 24h. Flow cytometry analysis after CD11c and CD86 staining. Data are represented as the normalized MFI of CD86, with the MFI in re-stimulated Jurkat-exposed MDDCs set to 100 (**left**), or as the percentage of CD86[+] MDDCs (**right**), and summarize n=21 (**a**) or n=5 (**b**) independent experiments. Red: Jurkat-exposed MDDCs restimulated with LPS; dark blue: C91-PL pre-exposed MDDCs restimulated with LPS; purple: C8166 pre-exposed MDDCs restimulated with LPS. Detailed statistical analysis is presented Briefly, the statistical difference between means was tested using RM one-way ANOVA (a, left and b, right) or Friedman test (a, right and b, left), in accordance with the distribution. **c.** Paired percentage of CD86[+] MDDCs in standard coculture (as measured in **a**), or with a transwell insert (as measured in **b**), with HTLV-1-infected C91-PL (**left**) or C8166 cells (**right**). Data summarized n=4 or 5 independent experiments, respectively analysed with paired t-test or its non-parametric equivalent Wilcoxon matched-pairs signed rank test as described in Supplementary Table 7. **d.** Schematic drawing summarizing the results from Fig. 5 and Supplemental Fig. 8. The drawing was created using BioRender.com.

Impairment of MDDCs is partially recapitulated by their culture in conditioned medium from HTLV-1-infected T cells and MDDCs co-culture

Since neither viral particles nor cell-cell contacts are strictly required to manipulate MDDC responsiveness, we then tested the conditioning ability of the supernatant of HTLV-1-infected T cells. More specifically, we hypothesized that IL-10 produced by HTLV-infected T cells[27] could be a candidate mediator, as it was reported to have tolerogenic properties[28]. However, except for the supernatant of the MT-2 infected T cell line, IL-10 was not detected in the supernatant of any of the other infected T cell lines (**Supplementary Fig. 9a**). Accordingly, none of these supernatants were able to prevent MDDC responsiveness to LPS stimulation (**Supplementary Fig. 9b**). Because of the indication of the involvement of soluble mediators (see **Fig. 5** above) we speculated that such mediators could be specifically released upon the co-culture between HTLV-1-transformed T cells and MDDCs. To test this hypothesis, we collected the conditioned medium of MDDCs cocultured with infected T cells for 24h, and used it to culture fresh, naïve MDDCs derived from the same monocyte donor (autologous cells) for 24h, before LPS stimulation for another 24h (see **Fig. 6a** for the experimental design). In contrast to the supernatant of infected T cells alone (**Supplementary Fig. 9b**), the supernatant from the co-culture was still able to dampen the responsiveness of MDDCs (**Fig. 6b**), although with less potency compared to coculture with infected T cells, as observed by analysis of paired experiments (**Fig. 6c**). This is consistent with the tendency observed upon transwell cultures (see **Fig. 5c**), and confirms the release of soluble tolerogenic mediators implying a cell-cell crosstalk between infected T cells and MDDCs.

We then addressed the kinetics of the release of these mediators. Conditioned medium of MDDCs cocultured with infected T cells were collected over time, and used to culture fresh autologous MDDCs before the addition of LPS (**Fig. 6d**). A lowered responsiveness of MDDCs was only observed with supernatants collected at least 18h after coculture (**Fig. 6e**), suggesting that the mediators are produced and released after a transcriptional response. Alternatively, we tested the kinetics required for the mediators to manipulate MDDC responsiveness. Conditioned medium of MDDCs cocultured with infected T cells was collected after 24h, and used to culture fresh autologous MDDCs for a varying duration before the addition of LPS (**Fig. 6f**). A lowered responsiveness of MDDCs was only observed

502   when MDDCs were cultured for at least 18h with the conditioned medium before adding the
503   LPS (**Fig. 6g**).

Figure 6. A cross-talk between HTLV-1-infected T cells and MDDCs induces the release of soluble tolerogenic mediators. a. Schematic representation of the experimental design. MDDCs were cocultured for 24h with uninfected Jurkat cells, or with HTLV-1-infected C91-PL or C8166 cells. Conditioned medium was collected and added on naive autologous MDDCs for 24h, before restimulation with LPS for an additional 24h. **b.** Flow cytometry analysis after CD11c and CD86 staining. Data are represented as the normalized MFI of CD86, with the MFI in re-stimulated MDDCs cultured in MDDC/Jurkat-conditioned medium set to 100. The color code is the same as in Fig. 5. Data summarize n=5 (C91-PL) or n=3 (C8166) independent experiments analysed with ordinary one-way ANOVA as described in Supplementary Table 7. **c.** Analysis of paired experiments, after standard coculture with C91-PL (**left**) or C8166 cells (**right**, as measured in **Fig. 5a**), or after culture in conditioned medium from MDDC/C91-PL coculture or from MDDC/C8166 coculture (n=5 or 3 independent experiments, respectively). Data were analysed with Paired t-test as described in Supplementary Table 7. **d.** Schematic representation of the experimental setting. MDDCs were cocultured with C91-PL, and conditioned medium was collected at multiple time points of coculture and added on naïve autologous MDDCs for 24h, before re-stimulation with LPS for another 24h. **e.** Flow cytometry analysis after CD11c and CD86 staining. Data are represented as the normalized MFI of CD86, with the MFI in LPS-stimulated MDDCs cultured in normal medium set to 100 (n=6 and n=3 independent experiments, for <12h and for 18h and 24h conditions, respectively). Data were analysed with ordinary one-way ANOVA as described in Supplementary Table 7. **f.** Schematic representation of the experimental setting. MDDCs were cocultured for 24h with C91-PL. Conditioned medium was collected and added on naive autologous MDDCs for the indicated time, before re-stimulation with LPS for another 24h. **g.** Flow cytometry analysis after CD11c and CD86 staining. Data are represented as the normalized MFI of CD86, with the MFI in LPS-stimulated MDDCs cultured in normal medium set to 100 (independent experiments were performed using n=7 for <12h condition, n=3 for 18h condition and n=2 for 24h condition). Statistical analysis of the two first columns was performed with Unpaired Welch's t-test. **h.** Schematic drawing summarizing the results from Fig. 6 and Supplemental Fig. 9. The drawing was created using BioRender.com.

Thus, the manipulation of MDDC responsiveness might also rely on a transcriptional control, which is consistent with the unique transcriptional signature induced by exposure to HTLV-1-infected T cells observed by RNA-seq.

Altogether, our results (recapitulated in **Fig. 6h**) show that upon coculture, a reciprocal cross-talk is initiated between HTLV-1-transformed T cells and MDDCs, which results in the transcriptionally-controlled release of a set of soluble tolerogenic mediators that manipulates MDDC responsiveness, in cooperation with additional mechanisms dependent on viral capture and/or cell-cell contacts.

Discussion

The remarkable ability of viruses to manipulate intracellular pathways in infected host cells is essential for establishing and maintaining infection. In addition to these direct effects, viruses may also establish a specific microenvironment around the infected cells, which may favor viral spread and persistence by influencing the signaling pathways and behavior of neighboring, uninfected cells, including immune cells[29,30]. As a by-stander effect, this may alter immune responsiveness to other stimuli. Mechanisms by which viruses influence uninfected cells in their local environment are still poorly understood. In this study, we aimed to decipher how HTLV-1-infected cells indirectly manipulate neighboring DCs. We demonstrate that exposure to HTLV-1-infected T cells induces a unique transcriptional signature in DCs, preventing their maturation and impairing their ability to be subsequently activated. This induction of tolerogenicity is not directly caused by infection of DCs, but instead results from a cross-talk between infected T cells and DCs, reminiscent of a viral microenvironment[29,30].

Previous work on the interplay between HTLV-1 and DCs has focused on the role of DCs as
intermediate target cells for HTLV-1 transmission to T cells[10–12]. Several studies investigated
whether viral capture by DCs could lead to T cell cis- and/or trans-infection, and whether
specific DC subtypes or maturation statuses could impact this proviral function. In particular,
in a previous paper, we demonstrated that immature MDDCs do efficiently capture and
transmit HTLV-1 to T cells, while mature MDDCs do not efficiently transmit HTLV-1 to T
cells, despite high levels of viral capture[11]. While these data indicated that the maturation
status of DCs could impact their proviral functions, they did not address whether exposure to
HTLV-1 induced an antiviral immune response by DCs. Such a question was investigated in
pDCs, which we showed to be efficient at sensing HTLV-1-infected cells after cell-cell
contact, and at responding by producing IFN-I[22]. Here, we specifically addressed how
MDDCs respond to HTLV-1 exposure. Intriguingly, our results are in stark contrast to those
obtained in pDCs, as we show that MDDCs do not respond to HTLV-1-infected T cells by
activating a typical maturation and antiviral program, even in the presence of cell-cell contact.
However, a unique transcriptional response is observed in exposed MDDCs, which could be
the result of a specific cross-talk occurring between HTLV-1-infected T cells and MDDCs,
that does not occur with pDCs. Alternatively, this cross-talk could also occur with pDCs, but
be outperformed by the highly efficient capacity of pDCs to produce IFN-I. Investigating how
pDC added to the coculture between HTLV-1-infected T cells and MDDCs could modulate
this induction of tolerogenicity, and how this tolerogenic behavior would shape HTLV-1
transmission to uninfected T cells, would give insight into how these distinct behaviors are
integrated into the microenvironment system. Of note, our results are also in contrast with
data obtained on murine DCs, using cell-free Moloney-pseudotyped chimeric HTLV-1[15],
which however may not fully recapitulate *in vivo* virus / host interactions in Human.
Specifically, given that cell-free HTLV-1 is not detected in HTLV-1 carriers or patients,
human DCs are probably mostly exposed *in vivo* to chronically infected cells rather than cell-
free virus. This underlines the importance of working with human cells when investigating
HTLV-1/host interactions.
Our data point towards the release of tolerogenic mediators following the cross-talk between
infected T cells and MDDCs upon coculture, which could cooperate with other mechanisms
dependent on viral capture and/or cell-cell contacts. The identity of the released mediators
initiating cross-talk observed between HTLV-1-infected T cells and MDDCs, as well as
produced as a result of this cross-talk, is currently under investigation in our laboratory. As
we observed that medium from infected T cells alone did not induce the tolerogenic behavior
of MDDCs, but that conditioned medium from infected T cells cocultured with MDDCs did,
it raises the hypothesis that HTLV-1-infected T cells sense a factor released by MDDCs, that
activates the release of a second factor by infected T cells. This second factor could then
launch a specific transcriptomic response in MDDCs that would lead to tolerogenicity.
Among potential mediators released by HTLV-1-infected cells, exosomes emerge as
compelling candidates. Their composition could be modified by viral infection, explaining the
differences observed in MDDC responses between exposure to uninfected or infected T cells.
Also, their production by T cells could be boosted by coculture with MDDCs, explaining the
requirement of a cross-talk between both cell types. Last, the diameter of exosomes, around
100 nm, is compatible with their diffusion through the pores of the transwell (0.4 µm) used in

our experimental settings. Interestingly, several HTLV-1-infected T cell lines, independently of their ability to produce viral particles or not, do produce exosomes containing cytokines, viral proteins and RNA[31–33] able to modulate target cells, resulting in an increased susceptibility to infection[25]. We can thus hypothesize that in our experimental settings, these exosomes could modulate DC functions, inducing the unique transcriptional signature that we observe. This would be reminiscent of data obtained on Epstein-Barr Virus (EBV), where infection changes the ability of exosomes purified from gastric cancer cell lines to mature MDDCs, leading to a defect in CD86 upregulation and reduced tumor immunity[34].

Whether HTLV-1 proteins contained in exosomes could contribute to the induction of tolerogenicity remains to be investigated. Viral proteins such as p30, HBZ, and Tax are found in exosomes produced by infected cells[25,31], independently of the production of viral particles by these infected cells. In addition, these viral proteins have the potential to interfere with immune functions. For instance, p30 expressed in MDDCs reduces IFN-I responses upon TLR3/4 stimulation, but not upon TLR7/8 stimulation[35], suggesting a specific targeting of the TLR3/4 signaling pathway. However, as we found no alteration in the IFN response of LPS restimulated, HTLV-1-pre-exposed MDDCs, a role of p30 can be excluded. In contrast, HBZ is known to inhibit the NF-κB pathway[36] by degrading the transcription factor p65[37]. As we observed the repression of the NF-κB pathway in HTLV-1-pre-exposed MDDCs upon LPS restimulation, we could suggest that exosome-transferred HBZ could contribute to modulating MDDC functions.

Interestingly, following the cross-talk between infected T cells and MDDCs, we observed the upregulation of genes involved in fatty acid metabolism (such as *ELOVL3*, *FADS1*, and *SLC27A6*) in MDDCs. As these genes are not found in the typical maturation program of MDDCs, it raises the possibility that upon coculture with HTLV-1-infected T cells, MDDCs produce fatty acids, that may act in an autocrine and paracrine manner to modulate MDDC functions. Of note, fatty acid precursors such as squalene and vitamin D are recognized inducers of tolerogenic DCs[38,39]. Polyinsaturated fatty acids have also been shown to block DC activation and functions[40], and inhibition of fatty acid synthesis has been shown to increase their production of pro-inflammatory cytokines[41]. Whether HTLV-1-exposed MDDCs produce higher levels of fatty acids, and whether these could contribute to their observed tolerogenic behavior, is currently under investigation. Thus, our transcriptomic analysis hint at how viruses could indirectly rewire lipid metabolism in immune cells to modulate the immune microenvironment.

Upon restimulation with LPS, HTLV-1-pre-exposed MDDCs undergo a specific transcriptomic response, including an abolished downregulation of *TGFB* expression, which remains highly expressed, and upregulation of *IL13* expression, which could be the basis for their tolerogenic properties. Of note, from the literature, the exact transcriptomic program underlying the tolerogenic behavior of DCs is unclear, as different transcriptomic signatures have been reported[42–44]. Nonetheless, a set of common genes has been defined as a gene signature of tolerogenic DCs[24], including *DDIT4* (DNA-damage-inducible transcript 4), which encodes an inhibitor of mTOR signaling[24]. Interestingly, responsiveness of *DDIT4* was also specifically conferred by pre-exposure to infected cells. Thus, the transcriptomic program induced in HTLV-1-pre-exposed MDDCs by LPS restimulation partially mirrors the tolerogenic DC signature defined in other contexts.

In conclusion, our study demonstrates that exposure to HTLV-1-transformed cells can
indirectly distort the functionality of DCs and impair their response to subsequent stimulation.
Our *in vitro* results are consistent with *in vivo* observations, which report that MDDCs from
HTLV-1-infected patients exhibit deficiencies in both basal maturation and responsiveness to
TNF-α treatment, as well as defects in the induction of T-cell response[16,18]. In addition,
PBMCs from HTLV-1-infected patients stimulated with tuberculin produce fewer TNF-α than
PBMC isolated from healthy individuals[45]. Finally, our results showing that viral production
is not necessary to dampen MDDC responsiveness, might reflect the *in vivo* situation in which
HTLV-1 expression is repressed in chronically infected cells[46]. Therefore, it is plausible that
defects in the myeloid cell response, particularly in DCs, may represent a mechanism of
HTLV-1 pathogenesis *in vivo*. Thus, this work illustrates how HTLV-1 might induce a local
immune microenvironment suitable for its own persistence. Such a microenvironment may
additionally contribute to by-stander immune dysfunctions, in asymptomatic HTLV-1 carriers
as well as symptomatic HTLV-1-infected patients.
658
659
660
Methods
Cells lines
HTLV-1 infected T-cell lines (C91-PL from Cellosaurus ref CVCL_0197, MT-2 ref
CVCL_2631[47], Hut102 ref CVCL_3526[48] and C8166 ref ECACC 88051601[23]) and control
uninfected T-cell lines (Jurkat from ATCC ref ACC 282, CEM/C1 ref CRL-2265, Molt-4
Cellosaurus ref CVCL_0013) were maintained at a cell density of $0.5.10^6$ cells/mL in
complete RPMI medium: RPMI1640 GlutaMAX (Gibco; 61870010) supplemented with 10%
fetal calf serum (FCS) and penicillin-streptomycin (100 U/mL and 100 µg/mL respectively).
The human fibrosarcoma cell line HL116 stably expressing the firefly luciferase reporter gene
under the control of the immediate early IFN-I inducible 6-16 promoter (kindly provided by
Dr. S. Pelligrini, Institut Pasteur, France[49]) was maintained under HAT selection (Gibco;
21060017, used at 1X final concentration) in DMEM GlutaMAX pyruvate medium (Gibco;
10569010) supplemented with 10% FCS and penicillin-streptomycin (100 U/mL and 100
µg/mL respectively).
All cells were grown at 37°C in 5% CO_2 and tested negative for mycoplasma contamination
on a regular basis. None of the cell lines were authenticated.
677
Human primary monocyte-derived dendritic cells (MDDCs)
MDDCs were derived from purified monocytes from healthy blood donors as described
previously[11]. Briefly, blood samples were collected at Etablissement Français du Sang (EFS)
from anonymous healthy blood donors according to the institutional Standard Operating
Procedures for blood donation, including a signed informed consent. Blood was diluted in
PBS 1X (Gibco) and peripheral blood mononuclear PBMCs were isolated using a density
gradient separation using Ficoll-Paque (Fisher Scientific, 11778538). Monocytes were then
isolated from PBMCs using a density gradient separation using Percoll Centrifugation Medium
(Fisher Scientific, 10607095). Freshly or frozen monocytes were cultured in six-well plates at
3.10^6 cells/mL in complete MDDC medium: RPMI1640 GlutaMAX (Gibco) supplemented

with 10% FCS, penicillin-streptomycin (100 U/mL and 100 µg/mL respectively), Hepes
buffer (Gibco, 15630080; 10 mM), MEM non-essential amino acids (2,5mM, Gibco,
11140050), sodium pyruvate (Gibco, 11360070; 1 mM), and beta-mercaptoethanol (Gibco,
31350-010; 0.05 mM). MDDC medium was supplemented with IL-4 and GM-CSF (Miltenyi
Biotec, 130-093-922 and 130-093-866; 100 ng/mL each) for differentiation. On day 3, the
culture medium was refreshed by discarding half of the medium and adding the same volume
of new MDDC medium and twice concentrated IL-4 and GM-CSF to all cell cultures.
Immature MDDCs were harvested on day 5 or 6. Every experiment was repeated on MDDCs
derived from independent blood donors.
697
Reagents and antibodies
Toll-like receptor (TLR)-4 agonist (LPS, tlrl-3pelps; 1µg/mL) and TLR-7/8 agonist (R848,
tlrl-r848; 3µg/mL) were purchased from Invivogen. MeV IC323-eGFP [50] is a recombinant
MeV expressing the gene-encoding eGFP (using the plasmid-encoding MeV IC323-eGFP
kindly provided by Yanagi, Kyushu University, Fukuoka, Japan). MeV IC323 recombinant
virus was rescued in 293-3-46 cells, as previously described[51]. Production of the viruses was
performed at 32°C. All viruses were propagated and titrated in Vero-SLAM/CD150 cells.
MDDCs were infected using a multiplicity of infection (MOI) of 1.
The following antibodies were used: V450-coupled mouse anti-Human CD11c (BD
Biosciences, 560369; 1/100), PE-coupled mouse anti-Human CD86 (Invitrogen, 12-0869-42;
1/100), APC-coupled mouse anti-Human CD83 (Miltenyi Biotec, 130-094-186; 1/50),
BV510-coupled mouse anti-Human CD83 (BD Biosciences, 563223; 1/50), APC-H7-coupled
mouse anti-Human CD80 (BD Biosciences, 561134; 1/50), BB515-coupled mouse anti-
Human PD-L1 (BD Biosciences, 564554; 1/25), PE-Cy7-coupled mouse anti-Human ICOSL
(BioLegend, 309410; 1/100), Mouse anti-Gagp19 clone TP7 (Zeptometrix, 081107; 1/500),
AlexaFluor647-coupled Alpaca anti-Mouse IgG1 (Chromotek, sms1AF647-1-5; 1/500),
Biotin-coupled mouse anti-Human CADM1 (MBL Life Sciences, CM004-6; 1/1000),
AlexaFluor647-coupled mouse streptavidin (Invitrogen, S31374; 1/500).
716
MDDCs coculture experiment
Immature MDDCs (3.10^5 cells) were plated in 48-well plates and cocultured with HTLV-1-
infected or control uninfected T-cells (6.10^4 cells) in 300µL complete MDDC medium
supplemented with IL-4 and GM-CSF for 24h or 48h. When indicated, transwell 24-well
plates with permeable polycarbonate membrane inserts (0.4µm diameter) were used (Fisher
Scientific, 10147291). Alternatively, MDDCs were cultured in supernatant collected from
either HTLV-1-infected or control uninfected T-cell culture, or from coculture of MDDCs
with HTLV-1-infected cells or control uninfected T-cell during 24h (or the indicated
duration). After 24h, cells were harvested, or stimulated with LPS or R848 or medium for an
additional 24h (or the indicated duration). At the indicated time points, cells and supernatants
were collected for phenotyping using flow cytometry and for cytokine quantification,
respectively.
729
Flow cytometry analysis

To assess surface marker expression, cells were fixed using 4% paraformaldehyde (PFA)
diluted from 20% PFA (Electron Microscopy Science, 50-980-493) in PBS 1X, and stained
with the indicated surface markers antibodies diluted in PBS 1X-1% Bovine Serum Albumin
(BSA) for 20 min at 4°C. To assess viral capture, cells were fixed and permeabilized using
the Fix/Perm FoxP3 and Transcription factors kit (Invitrogen, 00-5523-00) according to the
manufacturer's instructions, and stained with mouse anti-Gagp19 antibody followed by the
AlexaFluor647-coupled anti-mouse antibody diluted in the PERM buffer supplemented with
7% Normal Goat Serum (NGS) for 25 min at room temperature. Cells were then stained for the
indicated surface markers diluted in PBS 1X 1% BSA as described above. Compensation beads
(BD Biosciences, 552843) were used to correct signal overlap between the emission spectra of the
different fluorophores.
Data were acquired using a flow cytometer FACS CantoII (BD Biosciences) and analyzed
with FlowJo v10.7 software (BD Life Sciences). The full gating strategy is exemplified in
Supplementary Fig. 1.
745

746 **Cytokine quantification**
747 TNF-α and IL-10 were quantified in the collected supernatants using Bio-Plex Pro Human
748 Luminex kit (Bio-Rad) according to the manufacturer's instructions.
749

Type I interferon quantification
HL116 cells were seeded at 2.10^4 cells/well in 96-U-bottom-well plates and incubated for
24h. Supernatant collected from MDDCs culture (100 µL) or serial dilutions of recombinant
IFN-α (Tebu-Bio, RPA033Gu02) used for standard curve determination were added for an
additional 17h. Cells were then lysed (Promega Passive Lysis Buffer, E1941) and luciferase
activity assayed according to the manufacturer's instruction (Promega Luciferase Assay
System, E1501).
757

MDDCs isolation after coculture
Cells from MDDC / T-cell lines coculture (1.10^6 MDDCs and 2.10^5 T-cells) were harvested
and stained using biotin-coupled anti-CADM1 followed by AlexaFluor647-coupled
streptavidin diluted in PBS 1X-1% BSA for 20 min at 4°C. After several washes in PBS, cells
were then incubated with anti-AlexaFluor647 MicroBeads (Miltenyi Biotec, 130-091-395)
and separated on MACS Separation LD Columns (Miltenyi Biotec, 130-042-901) according
to the manufacturer's instructions. To assess the purity and yield of enrichment of MDDCs after
magnetic separation, 6.10^4 cells were collected before and after separation on LD columns, fixed
in 4% PFA and stained with CD11c-V450 antibody, before analysis using FACS Canto II.
767

RNA preparation
After magnetic separation, MDDCs were lysed and total RNA was extracted using the
NucleoSpin RNA Mini kit for RNA purification (Macherey-Nagel, 740955) according to the
manufacturer's instructions. RNA concentration was determined with a NanoDropND1000
spectrophotometer (Thermo Fisher Scientific) and samples were stored at -80°C until
shipment for external sequencing. Stranded RNA libraries were prepared after removal of
rRNA. High throughput sequencing of 150 bp paired-end reads was performed with an Illumina

HiSeq 2500 platform by Novogene Europe (Cambridge, United Kingdom). Each sample had on
average 50 million matched pairs of reads.
777
RNA-seq data analysis
The quality of sequences was checked using the FastQC tool. The reads were trimmed with
PrinSeq[52] to remove low-quality bases and then mapped to the human reference transcriptome
(hg19) using Kallisto pseudoalignment[53]. Differential gene analysis was carried out with DESeq2
package[54]. Genes with a basemean >10 showing an absolute fold change (FC) $\geq$ 2 with an
adjusted p-value <0.05 (Wald test using Benjamini and Hochberg method) were considered
differentially expressed. All DEGs identified in this study are listed in Supplementary Table 1 and
4. DAVID functional annotation tool using the KEGG pathways database was used for gene
ontology analysis[55,56]. For clarity, KEGG pathways were filtered and grouped as listed in
Supplementary Table 2 to reduce redundancy.
To identify the expression pattern responsiveness in LPS-stimulated MDDC pre-exposed to
C91-PL, we retrieved the genes by Venn analysis using the following list of DEGs:
MDDC/Jurkat restim.w/LPS vs MDDC/Jurkat; MDDC/C91-PL restim.w/LPS vs
MDDC/Jurkat restim.w/LPS; MDDC/C91-PL restim.w/LPS vs MDDC/Jurkat. Then, genes
were classified as follows (summarized in **Supplementary Fig 7a**): (i) responsiveness
confered by pre-exposure to infected cells: genes that are not differentially expressed between
mock- and LPS-stimulated Jurkat-pre-exposed MDDCs, but are differentially expressed
(either up or down) between mock-stimulated Jurkat-pre-exposed MDDCs and LPS-
stimulated C91-PL-pre-exposed MDDCs; (ii) responsiveness exacerbated by pre-exposure to
infected cells: genes that are either up-regulated in LPS-stimulated Jurkat-pre-exposed
MDDCs, compared to mock-stimulated Jurkat-pre-exposed MDDCs, and further up-regulated
in LPS-stimulated C91-PL-pre-exposed MDDCs, compared to LPS-stimulated Jurkat-pre-
exposed MDDCs; or down-regulated in LPS-stimulated Jurkat-pre-exposed MDDCs
compared, to mock-stimulated Jurkat-pre-exposed MDDCs, and further down-regulated in
LPS-stimulated C91-PL-pre-exposed MDDCs, compared to LPS-stimulated Jurkat-pre-
exposed MDDCs; (iii) responsiveness attenuated by pre-exposure to infected cells: genes that
are either up-regulated in LPS-stimulated Jurkat-pre-exposed MDDCs, compared to mock-
stimulated Jurkat-pre-exposed MDDCs, as well as up-regulated in LPS-stimulated C91-PL-
pre-exposed MDDCs, compared to mock-stimulated Jurkat-pre-exposed MDDCs, but down-
regulated in LPS-stimulated C91-PL-pre-exposed MDDCs, compared to LPS-stimulated
Jurkat-pre-exposed MDDCs; or down-regulated in LPS-stimulated Jurkat-pre-exposed
MDDCs, compared to mock-stimulated Jurkat-pre-exposed MDDCs, in LPS-stimulated C91-
PL-pre-exposed MDDCs, compared to mock-stimulated Jurkat-pre-exposed MDDCs, but up-
regulated in LPS-stimulated C91-PL-pre-exposed MDDCs, compared to LPS-stimulated
Jurkat-pre-exposed MDDCs; (iv) responsiveness abolished by pre-exposure to infected cells:
genes that are either up-regulated in LPS-stimulated Jurkat-pre-exposed MDDCs, compared
to mock-stimulated Jurkat-pre-exposed MDDCs, as well as down-regulated in LPS-stimulated
C91-PL-pre-exposed MDDCs, compared to LPS-stimulated Jurkat-pre-exposed MDDCs, but
not differentially expressed in LPS-stimulated C91-PL-pre-exposed MDDCs, compared to
mock-stimulated Jurkat-pre-exposed MDDCs; or down-regulated in LPS-stimulated Jurkat-
pre-exposed MDDCs compared to mock-stimulated Jurkat-pre-exposed MDDCs, as well as

up-regulated in LPS-stimulated C91-PL-pre-exposed MDDCs, compared to LPS-stimulated
Jurkat-pre-exposed MDDCs, but not differentially expressed in LPS-stimulated C91-PL-pre-
exposed MDDCs, compared to mock-stimulated Jurkat-pre-exposed MDDCs.
822

Statistical analysis

Median fluorescence intensity and percentages of gated cells were determined from flow
cytometry analysis with FlowJo software. Data were analyzed using Prism 8 (GraphPad
Software). The central tendency represents the mean, and the error bar represents the standard
error of the mean (SEM).Statistical analysis was performed as follows: normality of the
dataset was tested using Shapiro Wilk test. If the dataset followed a Gaussian distribution,
differences among means were tested using Repeated measures one-way ANOVA, or
Ordinary one-way ANOVA when the dataset contained missing values. Post-hoc comparisons
with the mean of the control condition (exposure to uninfected Jurkat T cells) were assessed
with Dunnet's test. Note that for Figure 1e, Sidak's test was used to compare the means of the
C91-PL-exposed MDDCs depending on their capture status.

When normality of the dataset was not met, analysis was performed using Friedman test, or
Kruskal Wallis test when the dataset had missing values. Post-hoc comparisons with the mean
of the control condition (exposure to uninfected Jurkat T cells) were performed using Dunn's
test.

For main Figures that are subsets of larger supplementary figures, the indicated p-values are
reported from the analysis on the total dataset.
When only two groups were compared, statistical analysis was performed using parametric
paired t-test (Figures 5c and 6c, Sup Figure 8b) or its non-parametric equivalent Wilcoxon
matched-pairs signed rank test (Figure 5c). Note that a parametric unpaired Welch's t-test was
used to analyze Figure 6g due to the uneven number of replicates.
The potential correlation investigated in Supplementary Figure 2i was fitted with a simple
linear regression model. p-value < 0.05 was considered significant, and statistics are denoted
as * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.

Data Availability

849   The RNA-seq raw data generated in the present study were deposited in GEO under the
850   accession number GSE266976.

References

1.   Bangham, C. R. M. Human T Cell Leukemia Virus Type 1: Persistence and Pathogenesis. *Annu. Rev.*
Immunol. **36**, 43–71 (2018).
2.   Gordon, C. A. *et al.* HTLV-I and Strongyloides in Australia: The worm lurking beneath. *Adv. Parasitol.*
111, 119–201 (2021).
3.   Haziot, M. E. *et al.* Detection of clinical and neurological signs in apparently asymptomatic HTLV-1
infected carriers: Association with high proviral load. *PLoS Negl. Trop. Dis.* **13**, e0006967 (2019).
4.   Kagdi, H., Demontis, M. A., Ramos, J. C. & Taylor, G. P. Switching and loss of cellular cytokine
producing capacity characterize in vivo viral infection and malignant transformation in human T- lymphotropic
virus type 1 infection. *PLoS Pathog.* **14**, e1006861 (2018).

5.   Paquin-Proulx, D. *et al.* MAIT cells are reduced in frequency and functionally impaired in human T
lymphotropic virus type 1 infection: Potential clinical implications. *PloS One* **12**, e0175345 (2017).
6.   de Castro-Amarante, M. F. *et al.* Human T Cell Leukemia Virus Type 1 Infection of the Three
Monocyte Subsets Contributes to Viral Burden in Humans. *J. Virol.* **90**, 2195–2207 (2015).
7.   Kannagi, M., Hasegawa, A., Takamori, A., Kinpara, S. & Utsunomiya, A. The roles of acquired and
innate immunity in human T-cell leukemia virus type 1-mediated diseases. *Front. Microbiol.* **3**, 323 (2012).
8.   Dalod, M., Chelbi, R., Malissen, B. & Lawrence, T. Dendritic cell maturation: functional specialization
through signaling specificity and transcriptional programming. *EMBO J.* **33**, 1104–1116 (2014).
9.   Rescigno, M. Dendritic cell functions: Learning from microbial evasion strategies. *Semin. Immunol.* **27**,
119–124 (2015).
10.   Jones, K. S., Petrow-Sadowski, C., Huang, Y. K., Bertolette, D. C. & Ruscetti, F. W. Cell-free HTLV-1
infects dendritic cells leading to transmission and transformation of CD4(+) T cells. *Nat. Med.* **14**, 429–436
(2008).
11.   Rizkallah, G. *et al.* Dendritic cell maturation, but not type I interferon exposure, restricts infection by
HTLV-1, and viral transmission to T-cells. *PLoS Pathog.* **13**, e1006353 (2017).
12.   Alais, S., Mahieux, R. & Dutartre, H. Viral Source-Independent High Susceptibility of Dendritic Cells
to Human T-Cell Leukemia Virus Type 1 Infection Compared to That of T Lymphocytes. *J. Virol.* **89**, 10580–
10590 (2015).
13.   Valeri, V. W. *et al.* Requirement of the human T-cell leukemia virus p12 and p30 products for
infectivity of human dendritic cells and macaques but not rabbits. *Blood* **116**, 3809–3817 (2010).
14.   Galli, V. *et al.* Essential Role of Human T Cell Leukemia Virus Type 1 orf-I in Lethal Proliferation of
CD4+ Cells in Humanized Mice. *J. Virol.* **93**, e00565-19 (2019).
15.   Rahman, S. *et al.* Murine FLT3 ligand-derived dendritic cell-mediated early immune responses are
critical to controlling cell-free human T cell leukemia virus type 1 infection. *J. Immunol. Baltim. Md 1950* **186**,
390–402 (2011).
16.   Nascimento, C. R. *et al.* Monocytes from HTLV-1-infected patients are unable to fully mature into
dendritic cells. *Blood* **117**, 489–499 (2011).
17.   Makino, M., Shimokubo, S., Wakamatsu, S. I., Izumo, S. & Baba, M. The role of human T-
lymphotropic virus type 1 (HTLV-1)-infected dendritic cells in the development of HTLV-1-associated
myelopathy/tropical spastic paraparesis. *J. Virol.* **73**, 4575–4581 (1999).
18.   Makino, M., Wakamatsu, S., Shimokubo, S., Arima, N. & Baba, M. Production of functionally deficient
dendritic cells from HTLV-I-infected monocytes: implications for the dendritic cell defect in adult T cell
leukemia. *Virology* **274**, 140–148 (2000).
19.   Azakami, K. *et al.* Severe loss of invariant NKT cells exhibiting anti-HTLV-1 activity in patients with
HTLV-1-associated disorders. *Blood* **114**, 3208–3215 (2009).
20.   de Witte, L., Abt, M., Schneider-Schaulies, S., van Kooyk, Y. & Geijtenbeek, T. B. H. Measles Virus
Targets DC-SIGN To Enhance Dendritic Cell Infection. *J. Virol.* **80**, 3477–3486 (2006).
21.   Klagge, I. M., Meulen, V. ter & Schneider-Schaulies, S. Measles virus-induced promotion of dendritic
cell maturation by soluble mediators does not overcome the immunosuppressive activity of viral glycoproteins
on the cell surface. *Eur. J. Immunol.* **30**, 2741–2750 (2000).
22.   Assil, S. *et al.* Sensing of cell-associated HTLV by plasmacytoid dendritic cells is regulated by dense β-
galactoside glycosylation. *PLoS Pathog.* **15**, e1007589 (2019).
23.   Bhat, N. K., Adachi, Y., Samuel, K. P. & Derse, D. HTLV-1 gene expression by defective proviruses in
an infected T-cell line. *Virology* **196**, 15–24 (1993).
24.   Avancini, D. *et al.* Aryl hydrocarbon receptor activity downstream of IL-10 signaling is required to
promote regulatory functions in human dendritic cells. *Cell Rep.* **42**, 112193 (2023).
25.   Pinto, D. O. *et al.* Extracellular vesicles from HTLV-1 infected cells modulate target cells and viral
spread. *Retrovirology* **18**, 6 (2021).
26.   Pais-Correia, A.-M. *et al.* Biofilm-like extracellular viral assemblies mediate HTLV-1 cell-to-cell
transmission at virological synapses. *Nat. Med.* **16**, 83–89 (2010).
27.   Mori, N. *et al.* Interleukin-10 gene expression in adult T-cell leukemia. *Blood* **88**, 1035–1045 (1996).
28.   Scheib, N. *et al.* The Dendritic Cell Dilemma in the Skin: Between Tolerance and Immunity. *Front.*
Immunol. **13**, 929000 (2022).
29.   Song, B. *et al.* Intercellular communication within the virus microenvironment affects the susceptibility
of cells to secondary viral infections. *Sci. Adv.* **9**, eadg3433 (2023).
30.   Yogev, O. *et al.* Herpesviruses shape tumour microenvironment through exosomal transfer of viral
microRNAs. *PLOS Pathog.* **13**, e1006524 (2017).
31.   Pinto, D. O. *et al.* HTLV-1 Extracellular Vesicles Promote Cell-to-Cell Contact. *Front. Microbiol.* **10**,
2147 (2019).
32.   Jaworski, E. *et al.* Human T-lymphotropic Virus Type 1-infected Cells Secrete Exosomes That Contain

Tax Protein. *J. Biol. Chem.* **289**, 22284–22305 (2014).

33. Kim, Y. *et al.* Extracellular Vesicles from Infected Cells Are Released Prior to Virion Release. *Cells* **10**,
781 (2021).

34. Hinata, M. *et al.* Exosomes of Epstein-Barr Virus-Associated Gastric Carcinoma Suppress Dendritic
Cell Maturation. *Microorganisms* **8**, 1776 (2020).

35. Fenizia, C. *et al.* Human T-cell leukemia/lymphoma virus type 1 p30, but not p12/p8, counteracts toll-
like receptor 3 (TLR3) and TLR4 signaling in human monocytes and dendritic cells. *J. Virol.* **88**, 393–402
(2014).

36. Huang, J., Ren, T., Guan, H., Jiang, Y. & Cheng, H. HTLV-1 Tax Is a Critical Lipid Raft Modulator
That Hijacks IκB Kinases to the Microdomains for Persistent Activation of NF-κB *. *J. Biol. Chem.* **284**, 6208–
6217 (2009).

37. Zhao, T. *et al.* Human T-cell leukemia virus type 1 bZIP factor selectively suppresses the classical
pathway of NF-κB. *Blood* **113**, 2755–2764 (2009).

38. van Megen, K. M. *et al.* 1,25-dihydroxyvitamin D3 induces stable and reproducible therapeutic
tolerogenic dendritic cells with specific epigenetic modifications. *Cytotherapy* **23**, 242–255 (2021).

39. Navarro-Barriuso, J. *et al.* Vitamin D3-Induced Tolerogenic Dendritic Cells Modulate the
Transcriptomic Profile of T CD4+ Cells Towards a Functional Hyporesponsiveness. *Front. Immunol.* **11**, 599623
(2020).

40. Zeyda, M. *et al.* Polyunsaturated Fatty Acids Block Dendritic Cell Activation and Function
Independently of NF-κB Activation *. *J. Biol. Chem.* **280**, 14293–14301 (2005).

41. Rehman, A. *et al.* Role of Fatty-acid Synthesis in Dendritic Cell Generation and Function. *J. Immunol.*
Baltim. Md 1950 **190**, 4640–4649 (2013).

42. Robertson, H. *et al.* Transcriptomic Analysis Identifies A Tolerogenic Dendritic Cell Signature. *Front.*
Immunol. **12**, 733231 (2021).

43. Navarro-Barriuso, J. *et al.* Comparative transcriptomic profile of tolerogenic dendritic cells
differentiated with vitamin D3, dexamethasone and rapamycin. *Sci. Rep.* **8**, 14985 (2018).

44. Schinnerling, K., García-González, P. & Aguillón, J. C. Gene Expression Profiling of Human
Monocyte-derived Dendritic Cells – Searching for Molecular Regulators of Tolerogenicity. *Front. Immunol.* **6**,
528 (2015).

45. Carvalho, N. B. *et al.* Impaired TNF, IL-1β, and IL-17 production and increased susceptibility to
Mycobacterium tuberculosis infection in HTLV-1 infected individuals. *Tuberculosis* **108**, 35–40 (2018).

46. Kinpara, S. *et al.* Stromal cell-mediated suppression of human T-cell leukemia virus type 1 expression
in vitro and in vivo by type I interferon. *J Virol* **83**, 5101–8 (2009).

47. Morozov, V. A. & Weiss, R. A. Two types of HTLV-1 particles are released from MT-2 cells. *Virology*
255, 279–284 (1999).

48. Bunn, P. A. & Foss, F. M. T-cell lymphoma cell lines (HUT102 and HUT78) established at the
National Cancer Institute: history and importance to understanding the biology, clinical features, and therapy of
cutaneous T-cell lymphomas (CTCL) and adult T-cell leukemia-lymphomas (ATLL). *J. Cell. Biochem. Suppl.*
24, 12–23 (1996).

49. Uzé, G. *et al.* Domains of interaction between alpha interferon and its receptor components. *J. Mol.*
Biol. **243**, 245–257 (1994).

50. Ferren, M. *et al.* Early Permissiveness of Central Nervous System Cells to Measles Virus Infection Is
Determined by Hyperfusogenicity and Interferon Pressure. *Viruses* **15**, 229 (2023).

51. Radecke, F. *et al.* Rescue of measles viruses from cloned DNA. *EMBO J.* **14**, 5773–5784 (1995).

52. Schmieder, R. & Edwards, R. Quality control and preprocessing of metagenomic datasets. *Bioinforma.*
Oxf. Engl. **27**, 863–864 (2011).

53. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification.
Nat. Biotechnol. **34**, 525–527 (2016).

54. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol.* **15**, 550 (2014).

55. Huang, D. W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene lists
using DAVID bioinformatics resources. *Nat. Protoc.* **4**, 44–57 (2009).

56. Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional
annotation of gene lists (2021 update). *Nucleic Acids Res.* **50**, W216-221 (2022).

976
977
978

Acknowledgements

We dedicate this work to the memory of Pr. Renaud Mahieux. We thank the Retroviral
Oncogenesis team, as well as Dr. Arnaud Moris, Dr. Sylvain Baize, Dr. Anne-Sophie
Beignon, Dr. Franck Halary, Dr. Claudine Pique, Pr. Mathias Faure, Dr. Delphine Muriaux
and Dr. Denis Gerlier for helpful discussion and/or critical reading of the manuscript. This
work has benefited from the facilities and expertise of the SFR Biosciences Lyon
(UMS3444/CNRS, US8/Inserm, ENS de Lyon, UCBL). We acknowledge the contribution of
the Etablissement Français du Sang Auvergne - Rhône-Alpes, and we thank Dr. Marlène
Dreux (CIRI) and her team for the help with PBMC isolation. We thank Dr. Carine Rey
(BIBS, CIRI, Lyon) for her help with RNA-seq data analysis. This work was supported by the
"Fondation pour la Recherche Médicale, équipe labellisée" program (grant number DEQ.
20180339200), and by the "Agence Nationale de la Recherche" (grant number ANR-22-
CE15-0044). AC and CJ are funded by the ENS de Lyon. FM, SA, and HD are funded by
Inserm. CM is funded by CNRS.

Author Contributions
Conception and design of the work: HD. Acquisition, analysis, or interpretation of data: AC,
FM, SA, CM, CJ, HD. Initial draft of the manuscript: AC, HD. Revision of the manuscript:
HD, CM, FM and CJ. Approval of manuscript: AC, SA, CM, FM, CJ and HD.
998
999

1000 **Competing Interests statement**
1001 The authors declare no competing interests.

1007 **Supplementary Table 1. Lists of DEGs relative to Fig. 2 and Supplementary Fig. 4.**
1008

1009 **Supplementary Table 2. Lists of filtered and grouped KEGG pathways relative to Fig.**
1010 **2c and Fig. 4c.** Enrichment values and p-values are given for each individual pathway
1011 ("Individual KEGG pathway" sheet), and the calculated means for grouped pathways are
1012 given in the "Grouped KEGG pathway" sheet.
1013

1014 **Supplementary Table 3. Lists of DEGs in each KEGG pathway, relative to Fig. 2c.** All
1015 genes annotated in each KEGG pathway are listed, and those retrieved among the DEGs in
1016 our dataset are highlighted in grey. For grouped pathways, the "Total" column summarizes all
1017 genes from the individual grouped pathways.
1018

1019 **Supplementary Table 4. Lists of DEGs relative to Fig. 4.**
1020

1021 **Supplementary Table 5. List of downregulated DEGs in each KEGG pathway, relative**
1022 **to Fig. 4c.** All genes annotated in each KEGG pathway are listed, and those retrieved among

1023 the downregulated DEGs in our dataset are highlighted in grey. For grouped pathways, the
1024 "Total" column summarizes all genes from the individual grouped pathways.
1025
1026 **Supplementary Table 6. List of upregulated DEGs in each KEGG pathway, relative to**
1027 **Fig. 4c.** All genes annotated in each KEGG pathway are listed, and those retrieved among the
1028 upregulated DEGs in our dataset are highlighted in grey. For grouped pathways, the "Total"
1029 column summarizes all genes from the individual grouped pathways.
1030
1031 **Supplementary Table 7.** List of Source data relative to figures. All raw data corresponding
1032 to the MFI or % for each gating defined by FlowJo, as well as statistical analyses, tests used
1033 and degree of freedom are listed for each individual panel in each figure, with the exception
1034 of RNAseq data from figures 2 and 4.
1035
1036 **Supplementary Table 8.** List of Source data relative to supplementary figures. All raw data
1037 corresponding to the MFI or % for each gating defined by FlowJo, as well as statistical
1038 analyses, tests used and degree of freedom are listed for each individual panel in each figure,
1039 with the exception of RNAseq data from supplementary figures 4 and 7.
1040