Genetically Encoded and Modular SubCellular Organelle Probes (GEM-SCOPe) reveal widespread lysosomal and mitochondrial dysfunction in human *PRKN* knockout astrocytes and neurons

Short title: Fluorescent, live imaging of organelles in human cell models

Camille Goldman[1-5], Tatyana Kareva[1-5], Lily Sarrafha[1-5], Braxton R. Schuldt[1-5], Abhishek

Saharsrabudhe[1-3], Tim Ahfeldt[1-5], Joel W. Blanchard[1-5] *

1. Icahn School of Medicine, Mount Sinai, New York, NY, USA

2. Nash Family Department of Neuroscience, Mount Sinai, New York, NY, USA

3. Friedman Brain Institute, Mount Sinai, New York, NY, USA

4. Ronald M. Loeb Center for Alzheimer's Disease, Mount Sinai, New York, NY USA

5. Black Family Stem Cell Institute, Mount Sinai, New York, NY, USA

* Corresponding author

Email: joel.blanchard@mssm.edu

Abstract

Cellular processes including lysosomal and mitochondrial dysfunction are implicated in the development and progression of many diseases and disorders, including neurodegenerative diseases and cancer. However, there is limited ability to continuously monitor these sub-cellular processes in real-time, and standardized assays to quantify these differences are lacking. Quantitative visualization of mitochondrial and lysosomal networks is crucial to better understand how these organelles are dysregulated during disease pathogenesis. To address this gap, we developed GEM-SCOPe (Genetically Encoded and Modular SubCellular Organelle Probes), an expandable toolbox of genetically encoded fluorescent markers, introduced into cell cultures by lentiviral transduction, which can be used to interrogate changes in proliferation and lysosomal and mitochondrial function in cell models of disease pathogenesis and progression. Genetically encoded fluorescent markers are a powerful tool to consistently visualize subcellular components in live cells without disrupting cellular processes. The fluorophores in GEM-SCOPe are designed to specifically localize to the desired organelle and illuminate information on localization, distribution, organelle turnover, and intracellular oxidative stress. GEM-SCOPe is modular; it can be expanded to include existing or new fluorophores and targeted to other subcellular structures or cell types, increasing the potential for genetically encoded fluorescent markers to probe any cell-model system or subcellular network in live cell cultures. To validate and illustrate the translational functionality of GEM-SCOPe, we applied it to track and quantify sub-cellular pathophysiology changes associated with Parkinson's Disease mediated by loss of function knockout of the *PRKN* gene. We expressed GEM-SCOPe in a *PRKN* knockout induced pluripotent stem cell line, differentiated into astrocytes and neurons, enabling us to track and quantify in real-time disease-associated changes in cellular proliferation, lysosomal distribution, mitochondrial transport and turnover, and reactive oxygen species. Collectively, we demonstrate that GEM-SCOPe is a powerful panel of live-cell fluorescent probes that provide critical insight

39   into the subcellular mechanisms underlying Parkinson's disease in human astrocytes and

40   neurons. GEM-SCOPe can be applied to a diverse range of cellular models of disease to glean

41   an understanding of the underlying mechanisms that promote disease onset and progression.

Abbreviations

BafA1: Bafilomycin A1

COX8A: Cytochrome c Oxidase Subunit 8A

DMSO: Dimethyl Sulfoxide

GEM-SCOPe: Genetically Engineered, Modular, SubCellular Organelle, Probes

H2B: Histone H2B

hiPSC: Human Induced Pluripotent Stem Cell

LAMP1: Lysosomal Associated Membrane Protein 1

PD: Parkinson's Disease

ROS: Reactive Oxygen Species

Introduction

Parkinson's Disease (PD) is a common neurodegenerative disease that robs individuals of their motor and cognitive functions, affecting over 10 million people worldwide and with an estimated 90,000 new diagnoses in the United States each year [1]. PD is clinically characterized by bradykinesia, resting tremor, rigidity, and postural instability [2-5] and pathologically characterized by the loss of dopaminergic neurons in the substantia nigra pars compacta and buildup of intracellular deposits of α-synuclein into insoluble protein aggregates called Lewy Bodies [2, 6-8]. From the first pathological descriptions of PD in the mid-20th century, advancements in many fields, including histology, genetics, cell biology, and neurology enabled insight into many aspects of the disease, from determining genetic and environmental risk factors and mapping disease progression to identifying pathways and proteins to target therapeutically. Despite great progress,

there is still incomplete understanding of the cellular dysfunction that precedes and contributes to

chronic neuronal death.

This limited advancement arises in part due to the absence of tools used to track molecular and

cellular changes in real-time throughout disease progression. Studies on human post-mortem

brain tissue are largely representative of end-stage disease states, making it difficult to dissect

which phenotypes are a primary, causative effect and which phenotypes are a secondary

response [9]. Mouse models offer more opportunities to study multiple time points throughout

disease progression as motor and cognitive symptoms can be monitored over time. However,

targeting specific cell types in mice remains technically challenging and time-consuming.

Emerging technologies such as the miniscope allow continuous monitoring of fluorescence-based

readouts on a cellular level in live mouse and rat brains [10-12]. However, the type of readout

from the miniscope has been restricted to neuronal activity from fluorescent calcium sensors and

resolution along the z-axis remains a limiting factor in analyzing deeper cortical tissue.

Regardless of such technologies, current genetic rodent models of PD fail to completely

recapitulate human disease and lack key pathological hallmarks. Drug-induced models can

recapitulate late-stage disease phenotypes and behaviors, but due to their acute onset, are

inadequate to study the gradual accumulation of cellular dysfunction that precedes clinical

presentation [13, 14]. Advancements in induced pluripotent stem cell (iPSC) differentiation and

CRISPR/Cas9 genome editing have led to the emergence of genetic cellular models of PD [15-

18]. These models enable researchers to study the effects of genetic and environmental factors

on specific human cell populations relevant to PD.

Cellular models of PD provide an exciting opportunity to study subcellular changes and responses

to stress or treatment in live cells. Cells can be genetically engineered to express fluorescent

proteins that can then be visualized by microscopy in live cells. The applicability of GFP as a

reporter gene was immediately appreciated after its identification and isolation in 1962 [19]. Since

then, hundreds of fluorophores have been developed by introducing mutations to naturally

occurring fluorophores isolated from sea anemones and jellyfish to modulate fluorescence

intensity, half-life, and excitation/emission spectra [20]. Additional fluorophores have been

developed as biosensors, by fluorescing or changing fluorescence emission in the presence of a

certain ligand or under specific cellular conditions [20-25]. Cells can express these fluorophores

with virtually limitless possibilities with regards to subcellular localization, cell-type specific

expression, and temporally controlled expression. Genetically encoded fluorescent proteins can

be used to follow a population of cells as they change due to aging, genetics, or environmental

stressors and stimuli.

PD has a complex genetic architecture underlying the disease. Over 90% of PD cases are

idiopathic; over 90 genetic risk loci have been identified by GWAS, but any of those individual

variants confer little risk on their own [26]. The remaining 10% of PD cases can be linked to highly

penetrant mutations in a handful of genes. While extremely rare, studying the function of these

genes and their impact on cellular processes sheds light on the fundamental pathways that drive

PD pathogenesis. Here, we focus on the *PRKN* gene. Autosomal recessive loss of function

mutations in *PRKN* are the most frequent known genetic cause of early onset cases of PD,

accounting for about 15% of PD cases with onset before the age of 50 [27-30]. *PRKN* encodes

for the protein PRKN, an E3 ubiquitin ligase associated with mitochondria. PRKN directs the

ubiquitination of outer mitochondrial membrane proteins on damaged mitochondria, targeting

them for degradation via the autophagy-lysosomal pathway (mitophagy) [31-33]. While the

function of PRKN is well understood, it is still unclear how loss of PRKN function leads to the

gradual disease progression of early onset PD. It is believed that neurodegeneration can begin

over 20 years before clinical onset, with a prolonged prodromal phase associated with several

non-motor symptoms. Thus, it is crucial to study models of early disease to understand what

cellular disturbances in the prodromal phase culminate in neurodegeneration.

Here, we present GEM-SCOPe (**G**enetically **E**ncoded and **M**odular **S**ub**C**ellular **O**rganelle

Probes), an expandable panel of genetically encoded reporters to track and quantify the

subcellular phenotypes in live cells. We developed a library of constructs that localize specifically

to the nucleus, mitochondria, and lysosomes and included fluorophores with a variety of emission

spectra or biosensor capabilities. All the lentiviral constructs were developed in the same

backbone vector to be modular, enabling any component, be it promotor, localization signal,

fluorophore, or antibiotic resistance, to be easily removed or replaced. We validated the

fluorophores with live-imaging of human induced pluripotent stem cell (hiPSC)-derived midbrain

astrocytes employing existing chemical dyes and chemical perturbations. We then applied GEM-

SCOPe to hiPSC-derived astrocytes and neurons from a *PRKN*-knockout model of PD [15]. This

revealed previously unseen changes in cellular proliferation, lysosomal distribution, mitochondrial

motility and turnover, and reactive oxygen species production. Our results demonstrate the

widespread utility of GEM-SCOPe to study a variety of translationally important subcellular

changes affected during PD. The modular, combinatorial flexibility of GEM-SCOPe can be

adapted to investigate any disease, and, therefore, it is a critical new resource that can be broadly

applied in neurodegenerative research and beyond.

Results

131 **Results**

Building a modular toolbox for the subcellular localization of genetically encoded fluorophores

Building a modular toolbox for the subcellular localization of

genetically encoded fluorophores

We developed GEM-SCOPe for live-cell imaging of subcellular organelle dynamics. We aimed to

produce a system that is highly modular to easily customize the expression, localization,

fluorescence, and selection for any given experimental need. We started with the FUW backbone

[41], a 3rd generation lentivirus backbone with a ubiquitin C promoter for high, ubiquitous

expression of the transgene across cell types [41, 42] and a woodchuck hepatitis virus post-

transcriptional regulatory element (WPRE) for improved lentiviral expression [41, 43, 44]. The

designed lentiviral constructs contain four components: promoter, localization sequence,

fluorophore, and antibiotic resistance (**Fig. 1A**). The content of these four components can be

mixed and matched to generate a library of lentiviral constructs to fit a diverse field of experimental

needs. To generate constructs *de novo*, components were combined and inserted into the plasmid

via Gibson assembly (**Fig. 1B, top**). Restriction sites were kept between each domain, so each

component or group of components could be excised via restriction digest and replaced with a

different component, either by restriction site cloning or Gibson assembly (**Fig. 1B, bottom**).

Thus, each component of the lentiviral plasmid can be removed or replaced to accommodate

features such as cell-type specificity with cell-type specific promoters, alternative subcellular

localization, and unique fluorophores.

The FUW backbone has a ubiquitin C promoter, but this can be replaced with lineage or cell-type-

specific promoters for selective expression in specific cell types or with drug-inducible promoters

to temporally control when the construct is transcribed (**Fig. S1A**). Subcellular localization is

determined by including the whole or partial coding sequence of a peptide with a known and

specific subcellular localization. This domain can be omitted to produce a fluorophore expressed

in the cytoplasm. The fluorophore component of the construct offers the most flexibility, from basic

fluorescent proteins to more complex fluorophores that change fluorescent excitation and/or

emission wavelength with time or environment (**Table S1**).

We developed GEM-SCOPe constructs to localize fluorophores to the cytoplasm (no tag), nucleus

(H2B-tag), mitochondria (COX8A-tag), and lysosome (LAMP1-tag) (**Fig. 1C**). Fluorophore

localization was validated using commercially available live cellular dyes. Nuclear localization was

achieved by fusing fluorophores to the C-terminus of histone 2B (H2B). H2B is an integral

chromatin protein that is found in all cells and is localized specifically to the nucleus [45, 46]. H2B

fused to mTagBFP2, a blue fluorophore, (H2B-mTagBFP2) co-localizes with DRAQ5, a far-red

live-cell nuclear dye [47] (**Fig. 1C, top**; Mander's coefficient > 0.99). In addition to the mTagBFP2

construct, we also developed lentiviral plasmids with H2B fused to Emerald (green; H2B-Emerald)

and mCherry (red; H2B-mCherry) (**Fig. 1A, S1B**).

To localize fluorophores to the lysosome, we utilized the N-terminal peptide sequence of

lysosomal-associated membrane protein 1 (LAMP1), a major component of lysosomal

membranes that plays a key role in lysosomal biogenesis and homeostasis. The N-terminus of

LAMP1 fused to mCherry (LAMP1-mCherry) co-localizes with LysoTracker (Thermo Scientific)

(**Fig. 1C, middle**; Mander's coefficient = 0.91). LysoTracker is a fluorophore that is partially

protonated at neutral pH and can readily cross membranes; once protonated it can no longer

diffuse so the fluorescent signal gets trapped in acidic compartments, including, but not limited to,

the lysosome [48, 49]. Overexpression of LAMP1 is associated with numerous cancers and

cancer metastasis [50-53]. By only using a partial sequence of LAMP1 and not the full peptide

sequence, we are able to target the fluorophore specifically to the lysosome without increasing

the expression of endogenous LAMP1 (**Fig. S1C**). We also developed lentiviral plasmids with the

LAMP1 targeting sequence fused to Emerald (green; LAMP1-Emerald; **Fig. 1A, 1D**).

Finally, mitochondrial localization was achieved using the mitochondrial targeting sequence of

COX8A. COX8A is a nuclear-encoded subunit of cytochrome-c oxidase (Complex IV) in the

electron transport chain [54, 55]. The first 25 amino acid residues of the COX8A peptide sequence

target the peptide to the mitochondrial inner membrane and are widely used to specifically localize

proteins and fluorophores to the mitochondria [56-58]. To confirm mitochondrial specificity of

COX8A-Emerald we compared localization with MitoTracker (Thermo Scientific), a dye that

accumulates in the mitochondria due to the negative membrane potential of the inner

mitochondrial membrane [48, 59]. The mitochondrial targeting sequence fused to an Emerald

fluorophore (COX8A-Emerald) exhibited significant co-localization with MitoTracker (**Fig. 1C,**

bottom; Mander's coefficient = 0.97). We developed lentiviral plasmids with the COX8A targeting

sequence fused to mCherry (red; COX8A-mCherry; **Fig. 1A, 1D**), Timer (green/red; COX8A-

Timer; **Fig. 1A, Fig. 5**), and Lemon (cyan/yellow; COX8A-Lemon; **Fig. 1A, Fig. S2**).

Once the localization strategies were finalized, we swapped out the fluorophores with different

fluorescent proteins that are excited and emitted at alternative wavelengths. Mixing localization

sequences and fluorophores with unique emission spectra allows simultaneous imaging of

multiple sub-cellular processes in the same cells such as mitochondrial network and lysosomal

distribution (**Fig. 1D**). Utilizing a combination of subcellular localization and diverse fluorophores,

we developed GEM-SCOPe, a fluorescent, live-cell imaging toolbox of lentiviral constructs that

were systematically validated to quantify cellular proliferation, lysosomal distribution, and

mitochondrial dynamics in human Parkinson's disease astrocytes.

200 **H2B-fused fluorophores offer improvements for efficiently**

201 **labelling nuclei for multi-day live-cell assays**

Quantification of cellular phenotypes by fluorescence microscopy is dependent on normalization

to total cell number. This is often accomplished by counting the number of nuclei. A few fluorescent

dyes that can cross the cellular and nuclear membranes of live cells and bind specifically to DNA

are commercially available. While efficient for short-term and end-point assays, such dyes

intercalate in DNA and therefore have cytotoxic and mutagenic effects [60, 61]. Furthermore,

nuclear dyes do not remain contained within the nuclear compartment, and their signal leaches

into the cytoplasm in just a few hours after exposing the cells. The cytotoxicity and loss of precise

labeling over time render these dyes less useful for long-term microscopy or cell-tracking

experiments. To address these concerns, we developed lentiviral constructs expressing nuclear-

localized fluorophores by fusing the nucleus-specific histone protein, H2B, with either mTagBFP2, Emerald, or mCherry (**Fig. 1C, 2A**). Human induced pluripotent stem-cell (hiPSC)-derived astrocytes transfected with H2B-mTagBFP2 were co-stained with DRAQ5, a far-red DNA-stain for live cells. The localization of the two nuclear signals was compared 3 hours and 24 hours after the addition of DRAQ5. Immediately after adding DRAQ5, there was no difference between the DRAQ5 stain and the H2B-mTagBFP2 (2-way ANOVA with Tukey's HSD; adj-p = 0.83) (**Fig. 2B, C**). Over the experimental period, the DRAQ5 signal spread to where 26% of the DRAQ5 signal was extranuclear (adj-p < 0.0001), while the extranuclear H2B-mTagBFP2 signal did not significantly change (adj-p = 0.99) (**Fig. 2B, C**). While the human eye can still readily distinguish the nucleus from the extranuclear stain in the DRAQ5 images, image analysis programs that rely on grey values for segmentation often fail to correctly differentiate the true nuclear signal from the extranuclear signal. Thus, lentiviral constructs expressing nuclear-localized fluorophores, such as the demonstrated H2B-mTagBFP2, can provide better alternatives for long-term live-imaging experiments with multiple time points.

H2B-fused fluorophores reveal increased cellular proliferation in PRKN knockout astrocytes

While dopaminergic neuron death is the predominant cellular outcome in Parkinson's Disease (PD), other cell types in the brain may change cell cycle regulation under neurotoxic conditions. Tracking cell numbers to monitor proliferation, senescence, or death in vitro can help researchers understand how different cell populations are responding to neurotoxic stress or potential therapeutic strategies.

First, we validated that our nuclear-localized construct, H2B-Emerald, could be used to track cellular proliferation. The proliferation of H2B-Emerald transduced hiPSC-derived astrocytes was

induced using fetal bovine serum (FBS) or blocked by inhibiting the proteasome with MG-132.

Astrocyte proliferation was measured by changes in the number of H2B-Emerald nuclei. As

expected, the astrocytes treated with FBS proliferated twice as rapidly as the untreated astrocytes

over 48 hours (2-way ANOVA with Tukey's HSD, p-adj = 0.005, **Fig. S3**), while the astrocytes

treated with MG-132 did not significantly proliferate over the same period (adj-p = 0.08, **Fig. S3**).

Thus, the GEM-SCOPe H2B-fluorophore fusion proteins can be used to track cellular proliferation

over multiple days in live cell cultures.

To examine the effect of loss of *PRKN* on astrocyte proliferation, we transduced wild-type

(*PRKN*$^{+/+}$) astrocytes with H2B-Emerald and isogenic *PRKN* knockout (*PRKN*$^{-/-}$) astrocytes with

H2B-mCherry (**Fig. 2D**). Leveraging the modularity of GEM-SCOPe, we were able to mix the two

astrocyte populations together and track their proliferation simultaneously. Over 48 hours, the

PRKN$^{-/-}$ population increased significantly by 25% (2-way ANOVA with Tukey's HSD, p-adj = 0.03;

Fig. 2E, F) while the *PRKN*$^{+/+}$ population did not proliferate in the same period (2-way ANOVA

with Tukey's HSD, p-adj = 0.4; **Fig. 2E, F**). These results contradict mouse studies that found that

PRKN-null mice had decreased astroglia proliferation [62]. However, PRKN has also been

described as a tumor suppressor, with loss of function mutations arising in a variety of cancers,

including glioblastomas [63-65]. Studies on PRKN in the context of cancer find that cancer

regulates the degradation of cyclins and cyclin-dependent kinases involved in the G1/S transition,

thus promoting continued proliferation [64, 66, 67]. It has been hypothesized that the outcome of

loss of function PRKN mutations are highly cell-context dependent; the increased half-life of

cyclins leads to cancer in mitotically competent cells while the same cyclins promote apoptosis

upon cell cycle reentry in post-mitotic neurons [64, 68-71]. Our findings highlight the importance

of studying disease-relevant human cells to capture the intricacies and nuances specific to the

human brain that could contribute to PD.

LAMP1-localized fluorophores reveal changes in lysosomal distribution in response to chemical and genetic stressors

Lysosomal dysfunction is widely implicated in PD and other forms of neurodegeneration. Several PD-risk and familial genes are involved in the autophagy-lysosomal-endosomal pathway [72-74], and many hallmarks of lysosomal dysfunction associated with familial PD are present in idiopathic PD [73, 75, 76]. Disruption to the autophagy-lysosomal-endosomal pathway is of broad importance in PD and there is a crucial need to study lysosomal function and dysfunction in the context of PD pathogenesis. As described above, we fused the N-terminus of LAMP1 to a fluorophore (Emerald or mCherry) to specifically target the fluorescent signal to the lysosome (**Fig. 1C, 3A**). To validate the GEM-SCOPe LAMP1-mCherry signal responds to lysosomal stress, hiPSC-derived astrocytes transduced with LAMP1-mCherry lentivirus were exposed to bafilomycinA1 (BafA1) for 15 hours. BafA1 inhibits lysosomal proton pump V-ATPases, which blocks autophagosome-lysosome fusion and lysosomal acidification, impairing lysosomal function [77, 78]. hiPSC-derived astrocytes were simultaneously transduced with a non-localized Emerald fluorophore lentivirus, to provide a cell body marker. We used CellProfiler [39] to evaluate the lysosomal number, size, and distribution within each cell. Compared to vehicle (DMSO) treatment, treatment with BafA1 increased the number of LAMP1-mCherry vesicles (unpaired t-test, p = 0.04; **Fig. 3B, C**), but did not affect lysosome size (unpaired t-test, p = 0.16, **Fig. 3B, C**). Treatment with BafA1 altered the distribution of lysosomes within the astrocytes. Astrocytes treated with BafA1 had a more dispersed distribution of LAMP1-mCherry signal, with only 41% concentrated in the perinuclear area, in contrast to 65% in vehicle-treated astrocytes (unpaired t-test, p = 0.002; **Fig. 3B, C**). Perinuclear and juxtanuclear lysosomes have been posited to be more acidic than peripheral lysosomes [79]. Lysosomal distribution can also be affected by cytosolic pH, with acidification promoting the spreading of lysosomes away from the nucleus [80-82]. Since BafA1 inhibits V-ATPases responsible for establishing and maintaining lysosomal pH, the redistribution

of lysosomes away from the nucleus upon BafA1 treatment might reflect a change in lysosomal

or cytosolic pH, and therefore lysosomal function, in the treated astrocytes.

In addition to its role in ubiquitinating damaged mitochondria for degradation via the autophagy-

lysosomal pathway [31-33], PRKN has been shown to regulate endosome organization [83] and

mitochondria-lysosome contact sites. Although it has been reported that *PRKN* knockout in

hiPSC-derived dopaminergic neurons leads to an increase in lysosome number and size

compared to isogenic, wild-type dopaminergic neurons [84], the effect of *PRKN* knockout on

lysosomes on human astrocytes is largely unknown. Due to PRKN's role in regulating the

autophagy-lysosomal pathway, we hypothesized *PRKN*$^{-/-}$ in human astrocytes would disrupt

lysosomal dynamics. To test this hypothesis, we applied the same CellProfiler analysis on

PRKN$^{+/+}$ and *PRKN*$^{-/-}$ hiPSC-derived astrocytes transduced with GEM-SCOPe Emerald and

LAMP1-mCherry lentivirus. *PRKN*$^{-/-}$ astrocytes had half as many LAMP1-mCherry vesicles per

cell compared to *PRKN*$^{+/+}$ astrocytes (unpaired t-test, p = 0.003, **Fig. 3D, E**). Additionally, the

PRKN$^{-/-}$ lysosomes were smaller than the *PRKN*$^{+/+}$ lysosomes (unpaired t-test, p = 0.055, **Fig. 3D,**

E). Finally, the *PRKN*$^{-/-}$ astrocytes had a decrease in perinuclear LAMP1-mCherry, with only 38%

of LAMP1-mCherry signal located in the perinuclear area, compared to 45% in *PRKN*$^{+/+}$ (unpaired

t-test, p = 0.005, **Fig. 3D, E**). Our results utilizing our GEM-SCOPe LAMP1-mCherry fluorophore

suggest that lysosomal distribution in astrocytes is altered upon loss of *PRKN* and further studies

to examine lysosomal content and activity are needed to understand the implications and how

this could differ from observations in other cell types.

Mitochondria-localized fluorophores capture mitochondrial fragmentation and alterations in mitochondrial trafficking along PRKN knockout dopaminergic neuron axons

Much like lysosomal dysfunction, mitochondrial dysfunction is implicated in PD and neurodegeneration. Therefore, monitoring mitochondrial network dynamics in response to stimuli and stress is important in understanding neurodegenerative diseases. As described above, we fused the N-terminus of the mitochondrial protein COX8A to a fluorophore (Emerald or mCherry) to localize the fluorophore specifically to the mitochondria (**Fig. 1C, 4A**). To validate that GEM-SCOPe COX8A localized fluorophores reflect changes to mitochondrial morphology, we induced mitochondrial stress by treating hiPSC-derived astrocytes transduced with the COX8A-mEmerald lentivirus with 2μM oligomycin for 4 hours. Mitochondrial stress induces fragmentation of the mitochondrial network, which facilitates mitophagy, but also decreases mitochondrial network respiratory capacity [85, 86]. Oligomycin, an inhibitor of ATP synthase, is a known inducer of mitochondrial fragmentation. Treatment with oligomycin induced mitochondrial fragmentation when compared to vehicle-treated astrocytes. The average area of mitochondria was reduced from 1.04 μm^2 in vehicle-treated astrocytes to 0.8 μm^2 in oligomycin-treated astrocytes (1-way ANOVA with Dunett's Test, p-adj = 0.0002; **Fig. 4B, C left)**, and average branch length was reduced from 1.28 μm in vehicle-treated astrocytes to 0.9 μm in oligomycin treated astrocytes (1-way ANOVA with Dunett's Test, p-adj < 0.0001; **Fig. 4B, C middle)**, as expected from a more fragmented mitochondrial network. The aspect ratio is the ratio of the mitochondrial length to width. An aspect ratio of 1 indicates a perfect circle, while larger aspect ratios indicate longer and narrower shapes. The mean aspect ratio decreased from 2.52 in vehicle-treated astrocytes to 2.14 in oligomycin-treated astrocytes (1-way ANOVA with Dunett's Test, p-adj < 0.0001; **Fig. 4B, C right**), another indicator that mitochondria are fragmenting upon oligomycin treatment. Metrics

of mitochondrial network fragmentation (mitochondrial area, branch length, and aspect ratio) are

all affected in a dose-dependent manner, with increasing concentrations of oligomycin having

increasingly more severe effects on mitochondrial network fragmentation (**Fig. S4**).

The transport of mitochondria along neuronal axons is critical for proper synaptic function and

neurotransmitter release. We hypothesized $PRKN^{-/-}$ may impair axonal mitochondrial transport.

To investigate this hypothesis, we transduced wild-type ($PRKN^{+/+}$) and $PRKN$ knock-out ($PRKN^{-/-}$

) hiPSC-derived midbrain organoids with COX8A-mEmerald. To facilitate imaging of axonal

mitochondrial transport, we generated radial axonal arbors by splatting organoids onto the surface

of a plate and allowing the axons to grow out [35]. On day 164 of organoid differentiation,

fluorescent mitochondria were imaged at high magnification over several minutes to monitor the

movement of mitochondria along these axons (**Fig. 4D; Video S1**). In the anterograde direction

(moving away from the soma), $PRKN^{-/-}$ mitochondria moved only 20% of the distance covered by

$PRKN^{+/+}$ mitochondria (Welch's t-test, $p < 0.0001$, **Fig. 4E left**) and took 50% longer (unpaired t-

test, $p = 0.005$; **Fig. 4E right**). Retrograde-moving mitochondria moved the same distance

(unpaired t-test, $p = 0.59$, **Fig. 4F left)** but the $PRKN^{-/-}$ mitochondria moved for 150% longer

compared to the $PRKN^{+/+}$ mitochondria (Welch's t-test, $p = 0.02$; **Fig. 4F right**). PRKN has

previously been implicated in mitochondrial transport due to its ubiquitination of mitochondrial

motor adaptor complexes [87]. Our results demonstrate that PRKN loss of function impairs

mitochondrial trafficking in human neurons, likely reducing the delivery of functional mitochondria

synaptic extremities and ultimately compromising neuronal function.

Dynamic fluorophores measure deficiencies in mitochondrial turnover and glutathione reduction potential in PRKN-knockout astrocytes

We demonstrated the utility of single excitation and emission wavelengths to discern phenotypes regarding cellular and subcellular dynamics. Numerous groups have developed fluorophores that change emission wavelengths under certain conditions or circumstances. We introduced several multi-emission fluorophores into GEM-SCOPe to enable quantification of organelle turnover, acidification, and oxidation, in addition to the localization information described with the previous fluorophores.

Fluorescent Timer is a mutated dsRed protein that irreversibly shifts its fluorescence emission from green to red due to tyrosine oxidation and can be used as a molecular clock to track protein turnover (**Fig. 5A**).[22] We, therefore, localized GEM-SCOPe-Timer to the mitochondria (COX8A-Timer), to further investigate mitochondrial turnover [88, 89] (**Fig. 5B**). To first validate COX8A-Timer, mitophagy was inhibited by treating hiPSC-derived astrocytes transduced with COX8A-Timer with the V-ATPase inhibitor bafilomycinA1 (BafA1). BafA1 treatment led to a significant 11% increase in the ratio of red-fluorescent mitochondria over green-fluorescent mitochondria compared to vehicle control astrocytes (unpaired t-test, p < 0.0001; **Fig. 5C, D**). Since mitochondrial degradation and turnover occur primarily through mitophagy, inhibition of mitophagy and mitochondrial recycling via BafA1 treatment results in a predicted increase in longer-lived mitochondria.

Because PRKN is directly involved in mitophagy, and we observed impaired retrograde mitochondrial transport in *PRKN*$^{-/-}$ neurons, we wanted to measure changes in mitochondrial turnover in human *PRKN*$^{-/-}$ cells. We transduced wild-type (*PRKN*$^{+/+}$) and *PRKN* knockout *(PRKN*$^{-}$

$^{/-}$) hiPSC-derived astrocytes with GEM-SCOPe COX8A-Timer. The ratio of red-fluorescent signal

over green-fluorescent signal increased by 14.5% in *PRKN$^{-/-}$* astrocytes compared to *PRKN$^{+/+}$*

astrocytes (unpaired t-test, p = 0.026, **Fig. 5E, F**). When broken down by individual channel, there

was no change in the green fluorescence (2-way ANOVA with Sidak correction, p-adj = 0.1288,

Fig. 5G), indicating no change in mitochondrial biogenesis. The increase in red to green ratio was

driven by a 27% increase in red-fluorescent signal (2-way ANOVA with Sidak correction, p-adj =

0.0005, **Fig. 5G**), which is in concordance with an accumulation of older mitochondria due to a

failure to properly degrade mitochondria in the absence of PRKN.

An expected consequence of impaired mitochondrial recycling is an accumulation of damaged

mitochondria and an increase in mitochondria-associated reactive oxygen species (ROS) from

inefficient cellular respiration. To measure reactive oxygen levels in live astrocytes, we introduced

GRX1-roGFP2, an established fluorophore for measuring ROS, into GEM-SCOPe [25]. GRX1-

roGFP2 is a fusion protein between GRX1 and roGFP2. When GRX1 is oxidized by glutathione,

it can subsequently oxidize roGFP2. Oxidized roGFP2 shifts its emission spectrum from green

fluorescence to blue fluorescence (**Fig. 6A**) [25, 90]. Thus the ratio of green to blue fluorescence

is an indicator of glutathione oxidation state. When GRX1-roGFP2 is further coupled with a

mitochondrial localization sequence, we can measure the oxidative stress of the mitochondria.

This construct was available as a retroviral vector using the signal sequence from *Neurospora*

crassa ATP synthase protein 9 [25]. When we cloned it into the FUW lentivirus backbone, we

decided to keep that localization sequence rather than replace it with the signal sequence for

COX8A used in the other GEM-SCOPe mitochondria-localized fluorophores (**Fig. 6B**). To validate

GEM-SCOPe GRX1-roGFP2 was working as expected, hiPSC-derived astrocytes transduced

with mito-GRX1-roGFP2 were treated with BafA1 to induce oxidative stress [91]. Astrocytes

treated with BafA1 had a 30% decrease in the ratio of green fluorescence to blue fluorescence

when compared to vehicle (DMSO) treated astrocytes (unpaired t-test, p < 0.0001, **Fig. 6C, D**).

The effect is driven by a 40% increase specifically in blue fluorescence (2-way ANOVA with Sidak

correction, p-adj < 0.0001, **Fig. 6E**) while the green fluorescence remained unchanged (2-way

ANOVA with Sidak correction, p-adj = 0.99, **Fig. 6E**). The increase in blue fluorescence is due to

an increase in roGFP2 oxidation, which is indicative of an increase in glutathione oxidation and

the presence of reactive oxygen species. Thus, GEM-SCOPe-mito-GRX1-roGFP can be used to

monitor glutathione reduction potential and how cells respond to reactive oxygen stress.

Because astrocytes play important metabolically supportive roles in the CNS and we observed

PRKN-related impairments to mitochondrial recycling with COX8A-Timer (**Fig. 5E-G**), we wanted

to understand if PRKN deficiency has effects on ROS production in astrocytes. We transduced

$PRKN^{+/+}$ and $PRKN^{-/-}$ astrocytes with mito-GRX1-roGFP2 lentivirus. Without introducing any

additional chemical mitochondrial stressors, we observed an 18% reduction in the ratio of green-

fluorescence signal to blue-fluorescence signal in $PRKN^{-/-}$ compared to $PRKN^{+/+}$ astrocytes

(unpaired t-test, p = 0.026; **Fig. 6F, G**). This is indicative of an increase in ROS in $PRKN^{-/-}$

astrocytes which correlates with our other observation of decreased mitochondrial turnover in the

same cells. It has been previously demonstrated that $PRKN^{-/-}$ hiPSC-derived dopaminergic

neurons show increased cellular ROS [15] and $PRKN^{-/-}$ mice have decreased antioxidant

capabilities and increased ROS-related protein and lipid damage [92]. By introducing fluorophore-

biosensors into our lentiviral catalog, we were able to expand the phenotypes we can observe in

live cells to measure changes in the intracellular environment in response to chemical and genetic

perturbations.

Discussion

Here, we developed and validated GEM-SCOPe (Genetically Engineered, Modular, SubCellular Organelle Probes), a collection of subcellularly targeted fluorophores encoded in a lentiviral backbone. By exploiting the modular design of the constructs, we easily cloned fluorophores with different excitation/emission spectra specified to localize to the nucleus, lysosomes, or mitochondria. We demonstrated that these fluorophores specifically localize to the desired organelle and can quantitatively assess cellular responses to chemical perturbations. Finally, we applied GEM-SCOPe to human *PRKN*-knockout astrocytes and neurons, which revealed widespread perturbations to cellular proliferation, lysosomal distribution, and mitochondrial dynamics, providing new insights into PD pathogenesis.

Gliosis has been widely reported in human post-mortem brain studies of patients with *PRKN* loss-of-function mutations [94-97]. Similar pro-inflammatory phenotypes have also been observed in mouse models and stem-cell-derived glial models of PRKN deficiency [98, 99]. However, there is still a limited understanding of how genetic mutations related to PD modify astrocytes and their role in neurodegeneration. We used GEM-SCOPe to investigate cellular, mitochondrial, and lysosomal dynamics in astrocytes lacking functional PRKN protein. Using nucleus-localized fluorophores, we show that *PRKN* knockout human astrocytes have increased cellular proliferation compared to isogenic, wild-type astrocytes. LAMP1-localized fluorophores revealed that lysosomes in *PRKN* knockout astrocytes are distributed further from the perinuclear nuclear area. Finally, a variety of mitochondria-localized fluorophores revealed that *PRKN* knockout astrocytes have impaired mitochondrial transport, decreased mitochondrial turnover, increased reactive oxygen species, and oxidized glutathione. While additional experiments should be done to support these findings, GEM-SCOPe highlights that there is extensive dysregulation and disruption to normal organelle dynamics in human *PRKN* astrocytes that warrants further investigation.

GEM-SCOPe was developed in a 3rd generation lentiviral system. We selected this delivery

system because 1) lentiviruses are safe to produce; 2) they infect both dividing and non-dividing

cells; and 3) they can be used to develop stable cell lines. 3rd generation lentiviral systems are

replication-incompetent, so cells that are transduced with lentivirus cannot produce more viral

particles. Only the transfer plasmid, which contains the insert of interest, can be integrated into

the host genome; the other plasmids that encode lentiviral structural and envelope genes do not

integrate into the host genome, preventing viral replication. Most lentiviruses are produced with

VSV-G (Vesicular stomatitis virus G glycoprotein) as the envelope protein, as it provides broad

tropism and enables high transduction efficiency in a wide variety of cell types [100]. Finally, GEM-

SCOPe lentiviruses were developed with the option to include an antibiotic-resistance gene (**Fig.**

1A), which can be used to establish stable cell lines that have incorporated the lentiviral load.

Although lentiviruses provide a lot of flexibility and safety when delivering genetic information,

they also possess some limitations. Lentiviruses have an ideal plasmid size of 9-10kb, and viral

titer can decrease if plasmids exceed this size [101]. The H2B-mTagBFP2, H2B-mCherry, and

H2B-Emerald lentiviruses (**Fig. 1, 2**) are about 11kb each and transduced hiPSC-derived

astrocytes with about 90% efficiency. Coupled with the puromycin resistance that was also

included in this construct, we were still able to generate a population with 100% viral transduction

despite the size limitation. Another drawback of lentiviral gene delivery is that inserts often

integrate into actively transcribed loci [102]. Differentiation of hiPSCs into specific cell types

requires extensive chromatin remodeling, and genomic sequences integrated via lentivirus can

often be silenced in this process. In the case where hiPSCs must be transduced before

differentiation, antibiotic selection during differentiation can select for cells that have integrated

the viral load into a genomic region that is not silenced. The final drawback of lentiviral gene

delivery is that fluorescence intensity is dependent on the viral copy number in each cell and

cannot be used as a readout for organelle abundance. Rather, we need to use measurements

like distribution, size, and ratios of different emissions within the same cell. This is not just a

limitation of lentiviral gene delivery but of other viral and non-viral delivery systems.

In this study, we focused on PD and specifically the alterations to astrocytes with *PRKN* mutations.

However, alterations in proliferation, mitochondria, and lysosomes are by no means unique to PD.

Many reviews cover the extensive roles of mitochondria and lysosomes in health and disease

[103-106]. GEM-SCOPe can be applied beyond neuronal cells and neurodegeneration to

countless other cell types to investigate subcellular dynamics that affect disease development

and progression. Due to its modular design, GEM-SCOPe can be expanded and molded by

research groups to meet specific needs. Different targeting sequences can be integrated to

localize fluorophores to relevant subcellular compartments. New fluorophores and biosensors are

always being developed to measure additional intracellular conditions. By applying GEM-SCOPe

to disease models, we will be able to better understand how cells are disrupted over the course

of disease progression, providing insight for future therapeutic targets.

Acknowledgements

484   The authors thank Deanna Benson for discussions on mitochondrial transport and kymograph

485   analysis. We thank the Microscopy CoRE at Icahn School of Medicine for providing access to the

486   Leica DMi8 and Thermo Scientific CX7 High Content Screening Platform, as well as training and

487   technical expertise for using these microscopes.

Materials and Methods

Molecular cloning

All vectors were derived from pFUW (Addgene plasmid #14882). pFUW was linearized using NheI-HF (NEB R3131) and BamHI-HF (NEB R3136) per manufacturer's protocol, and the desired fragment was gel-purified (Macherey-Nagel 740609). Components of the insert were PCR amplified using Q5-Hot Start DNA Polymerase (NEB M0493) per manufacturer's recommendations. Primers were designed using the NEBuilder Assembly Tool (nebuilder.neb.com) to add 5' and 3' sequences that make the fragments compatible for Gibson assembly and ordered from IDT (Integrated DNA Technologies). Gibson reactions were set up with NEBuilder HiFi DNA Assembly Master Mix (NEB E2621), 50ng of linearized backbone, and 3-fold molar excess of each PCR fragment. For fragments less than 200bp, 5-fold molar excess was used. Reactions were incubated at 50°C for 60 minutes. The Gibson product was transformed in 10-beta competent *E. coli* (NEB C3019) grown on LB Agar plates with 100ug/mL ampicillin. Plasmids from single clones were isolated via miniprep (Macherey-Nagel 740499) and sequenced to confirm successful cloning.

Lentivirus Production

Lentivirus was produced by transfection of HEK293T cells as adapted from Dull et al. 1998 [34]. HEK293T cells were grown in Dulbecco's Modified Eagle Medium (DMEM; Gibco 11965092) with 10% bovine calf serum (Cytiva SH30073) at 37 °C in a 5% CO_2 incubator. The day before transfection, ~10^7 cells were plated per 15cm dish coated in 0.1% gelatin. Media was changed just prior to transfection. For one 15cm dish, 22.5ug of transfer vector was combined with 14.7ug of pMDLg/pRRE (Addgene plasmid #12251), 5.7ug of pRSV.Rev (Addgene plasmid #12253), and

7.9ug of pMD2.G (Addgene plasmid #12259) in a 15mL conical tube. The plasmid mixture was

added to 1mL of 278 uM $CaCl_2$ (Sigma C1016) and mixed thoroughly before adding 1mL of 2x

BBS solution [280 mM NaCl (Fisher Scientific S271-1), 50 mM BES (Millipore 391334), 1.5mM

Na_2HPO_4 (Sigma S5136); pH = 6.95] dropwise. Tubes were mixed by inversion, incubated at room

temperature for 1 min, and then added dropwise to HEK293T dish. Cell culture media was

changed after 16-24 h. Conditioned media was then collected every 24 h for 2 days; conditioned

media from the same plate was combined over the two collections. Conditioned media was

centrifuged at 2000 x g for 10 minutes to pellet debris and filtered through 0.22-um-pore-size PES

filter (Millipore SCGP00525). To concentrate virus, filtered media was incubated with PEG 8000

(Sigma P2139) and NaCl, at a final concentration of 5% and 0.15M respectively, overnight at 4°C,

then centrifuged at 3000 x g for 20 min. The pellet was resuspended in DMEM/F12 with GlutaMAX

(Gibco 10565018) to 1% of the original supernatant volume. The concentrated virus was stored

in aliquots at -80°C.

Midbrain Organoid and Astrocyte Differentiation

hiPSC-derived astrocytes were extracted from 100-day-old midbrain organoids as described in

Sarrafha et al 2021 and Parfitt et al 2024 [35, 36]. In short, hiPSCs were cultured in StemFlex

medium (Gibco A3349401) at 37 °C in a 5% CO_2 incubator. hiPSCs were seeded into 125-mL

spinner flasks (Corning 3152) and allowed to self-aggregate in StemFlex (Gibco A3349401)

supplemented with 10 µM Y-27632 (Tocris 1254) and 1% Penicillin-Streptomycin (Gibco

15140122). Once aggregates were between 300 µm and 500 µm, differentiation was initiated by

dual-SMAD inhibition with 10 µM SB431542 (Stemgent 04-0010), 100nM LDN193189 (Tocris

6053), 1x B27 without Vitamin A (Gibco 12587010) and 1x N2 (Gibco 17502048) in DMEM-F12

with GlutaMAX (Gibco 10565018). Midbrain patterning was achieved with the addition of 3 µM

CHIR99021 (Tocris 4423), 2 µM purmorphamine (Stemcell 72204), and 1 µM SAG (Cayman

11914) starting 4 days after neural induction. After patterning, media was changed to DMEM/F12

with GlutaMAX, supplemented with 1x N2, 1x B27 without Vitamin A, 20 ng/mL GDNF (Peprotech

450-10), 20 ng/mL BDNF (Peprotech 450-02), 200 µM L-ascorbic acid (Fisher Scientific BP351),

100 µM dibutyryl cAMP (Biogems 1698950), and 10 µM DAPT (Cayman 13197). After 35 days,

organoids were transferred to ultra-low attachment plates (Corning 3516), with about 5 organoids

per mL of media and cultured in DMEM/F12 with GlutaMAX supplemented with 1x N2, 1x B27

without Vitamin A, 10 ng/mL GDNF, 10 ng/mL BDNF, and 200 µM ascorbic acid.

Astrocytes were extracted from midbrain organoids starting at day 100. Organoids were gently

dissociated in a trypsin enzyme solution (TrypLE Select, Gibco 12563011) using a glass pipette

to break up organoids into large chunks. The organoid chunks from 5-10 organoids were placed

on 15cm dishes coated with 0.1% gelatin and maintained in Astrocyte Medium (AM; ScienCell

#1801). Astrocytes were allowed to grow out of the organoid chunks until the plate was confluent

(about 1 week). At this point, astrocytes can be cryopreserved. Astrocytes were maintained in AM

and switched to experimental media [1:1 DMEM/F-12:Neurobasal (Gibco 21103049), 1x B27

without Vitamin A, 1x N2, 1x MEM-Non-Essential Amino Acids (Gibco 11140050), 1x GlutaMAX

(Gibco 35050061), 10 ng/mL CNTF (Peprotech 450-13)] 2-3 days before imaging.

554   Astrocyte Viral Transduction

iPSC-derived astrocytes were transduced during regular passaging. In short, astrocytes were

lifted, spun down, resuspended in AM media, and divided into 1.5mL microcentrifuge tubes,

depending on the desired split ratio and number of transductions. The virus was added to the cell

suspension and incubated for 5-10 minutes before plating the mixture. Media was topped off such

that the final dilution of the virus in media was 1:50. Depending on the viral titer, this dilution

sometimes increased or decreased. Astrocytes can also be transduced without passage by

adding the virus directly to the media. Media is changed after 24 hours. Antibiotics were added to

the media, if relevant, starting at least 3 days after transduction.

MitoTracker and LysoTracker

565 For MitoTracker staining, astrocytes were incubated with 100 nM of MitoTracker DeepRed

566 (Thermo Scientific M22425) for 30 min at 37 °C. For LysoTracker staining, astrocytes were

567 incubated 100 nM of LysoTracker DeepRed (Thermo Scientific L12492) for 5 min at 37 °C. Cells

568 were then washed twice with PBS before incubation with Hoechst 33342 (Thermo Scientific

569 62249).

qPCR

RNA was extracted from samples using TRIzol (Invitrogen 15596018) and an RNA extraction kit

following the manufacturer's recommended protocol (Zymo R2062). RNA concentration and purity

were measured on a spectrophotometer. cDNA was produced from 1 ug of RNA per sample, using

Maxima H Minus Reverse Transcriptase (Thermo Scientific EP0752) and following the

manufacturer's recommendations. qPCR reactions were prepared in a 384-well plate (Applied

Biosystems 4309849) with 1:10 dilution of cDNA, 2x PowerUp SYBR Green Master Mix (Applied

Biosystems A25741) and 1 µM each of forward and reverse primers. qPCRs were run on

QuantStudio 7 (Applied Biosystems 4485701) and analyzed using the ΔΔCt method, normalized

to *ACTB*. The following primers were used:

ACTB Forward: CCTGGCACCCAGCACAAT

ACTB Reverse: GCCGATCCACACGGAGTA

LAMP1 Forward: CAGATGTGTTAGTGGCACCCA

LAMP1 Reverse: TTGGAAAGGTACGCCTGGATG

Live Cell Imaging

In general, astrocytes were seeded at 10 x 10^3 cells per well of a 96-well plate with a cover glass

thickness polystyrene bottom (Greiner 655090) coated in 0.1% gelatin 3-5 days before imaging.

Proliferation assays were seeded at 8 x 10^3 cells per well and started 1 day after seeding. Before

imaging, cells were incubated with Hoechst 33342 (Thermo Scientific 62249) or DRAQ5 (Thermo

Scientific 52251) for 10 minutes.

Wide-field live cell imaging was performed using the Thermo Scientific CX7 High Content

Screening Platform with 20x and 40x objective lenses (LUCPLFLN20x and LUCPLFLN40x). The

microscope was equipped with an incubation unit that maintained temperature, CO_2, and humidity

during image acquisition. Confocal images were acquired on a Nikon Ti2E AX R confocal

microscope with a 20x objective lens (PLAN APO λD 20x) and 6x optical zoom. Each well was

imaged in 5 (confocal) or 25-36 (widefield) different areas, with the average of those images

represented as a single data point.

ImageJ Image Analysis

Nuclear Proliferation: Nuclear proliferation was quantified by converting images to a binary based

on a threshold and then using 'Analyze Particles' to obtain a count for the number of nuclei in a

field. The same threshold and parameters were kept for all time points.

Mitochondrial Fragmentation: Mitochondrial fragmentation was measured using the ImageJ

plugin, Mitochondrial Analyzer [37]. For thresholding, the "Block Size" was set to 1.45 microns

and the "C-Value" was set to 5.

Ratiometric Quantification: Ratiometric quantifications for roGFP2, Timer, and Lemon were

achieved by measuring the mean grey value in both channels of interest. The values for each

channel were then divided to generate the ratio of relative intensity between the two channels.

Kymograph Analysis: Kymograph analysis of mitochondrial movement was achieved using the

Image J plugins KymographClear 2.0 and KymographDirect [38]. KymographClear 2.0 was used

to trace mitochondrial trajectories and generate kymographs from which stationary and motile

mitochondria were identified. KymographDirect was then used to extract quantitative information

from the trajectories of the motile mitochondria.

616   Cell Profiler Image Analysis [39, 40]

Subcellular Co-localization: For nuclear co-localization, nuclei in the genetically encoded

fluorophore channel and the DRAQ5 channel were identified, and nuclei from cells that were not

transduced were filtered out. For lysosomal and mitochondrial co-localization, propagation from

the nuclear stain was used to identify cell bodies. Cells were filtered to only analyze those with

lentivirus signal within the cell body. Co-localization was reported as the Mander's coefficient

indicating the overlap of the genetically encoded fluorophore with the commercially available

stain.

Lysosomal Distribution: Cells were defined by nuclei and full cell fluorophore signal. Lamp1-

mCherry signal was assigned to the cell body with which it overlapped. Only cells that were

successfully transduced were included in the analysis. The distribution of the Lamp1-mCherry

signal was calculated using the "MeasureObjectIntensityDistribution" with two bins generated for

each cell using the nuclei as the center of the cell. Bins were scaled such that the inner bin

(perinuclear) was always the same percent of total cell area regardless of cell size.

Figure Legends

Fig. 1: A toolbox of lentiviral plasmids developed and validated for subcellular localization

of fluorophores (A) Schematic of lentiviral plasmids highlighting the combinatorial power of the

toolbox components. Generated using BioRender **(B)** Schematic demonstrating the cloning

strategies used to generate the constructs used in this study as well as any future constructs.

Generated using BioRender **(C)** Representative images of subcellular localization validation with

available dyes in hiPSC-derived astrocytes and Mander's co-efficient for overlap of the genetically

encoded fluorophore with the dye. Top: H2B-mTagBFP2 and DRAQ5 (n = 3; each replicate is an

average over 16 images); Middle: LAMP1-mCherry and LysoTracker Deep Red; nuclei stained

with Hoechst33342 (n = 4; each replicate is an average over 4 images); Bottom: COX8A-Emerald

and MitoTracker Deep Red; nuclei stained with Hoechst33342 (n = 3, each replicate is an average

over 36 images). Scale bars = 50 μm **(D)** Representative images of combinatorial applications of

GEM-SCOPe. Top: Cell expressing LAMP1-Emerald and COX8A-mCherry. Bottom: cell

expression LAMP1-mCherry and COX8A-Emerald. Nuclei were stained with DRAQ5. Scale bars

= 50 μm.

Fig. 2: Nucleus-localized fluorophores offer improvements over available stains for

multiday imaging (A) Schematic of the lentiviral construct localizing mTagBFP2 to the nucleus

with an H2B fusion protein and antibiotic resistance to puromycin used in the following panels **(B)**

Representative images of H2B-mTagBFP2 and DRAQ5 nuclear stain 1 hour (top) and 24 hours

(bottom) after adding DRAQ5 to the cell cultures. **(C)** Quantification of the percent of stain

localized to the nuclear area versus the cytoplasm 1 hour and 24 hours after adding DRAQ5 to

cell cultures. (n = 3 per time point; each replicate is an average over 36 images) **(D)** Schematic

of the lentiviral construct localizing Emerald or mCherry to the nucleus with an H2B fusion protein

and antibiotic resistance to puromycin used in the following panels **(E)** Representative images of

nuclei from $PRKN^{+/+}$ astrocytes, labeled with H2B-Emerald, and $PRKN^{-/-}$ astrocytes, labeled with

H2B-mCherry, cultured together during a proliferation assay. Images were taken 24 hours (top)

after seeding and 72 hours (bottom) after seeding **(F)** Quantification of number of $PRKN^{+/+}$ and

$PRKN^{-/-}$ nuclei per image field 24 hours and 72 hours after seeding (n = 3 per time point; each

replicate is an average over 25 images). Scale bars = 50 μm. All images were acquired on the

CX7 HCS platform with a 20x objective lens.

Fig. 3: Lysosome-localized fluorophores reveal changes in lysosomal distribution upon

chemical and genetic perturbation (A) Schematic of the lentiviral constructs used in the

following panels localizing (top) mCherry to the lysosome with the 5' sequence of *LAMP1* and

(bottom) Emerald to the whole cell by not including a localization sequence. **(B)** Representative

images of astrocytes transduced with Emerald and LAMP1-mCherry lentiviruses and treated with

100 nM bafilomycin A1 (bottom) or a vehicle (DMSO; top). Nuclei were stained with

Hoechst33342. Far right: Output from CellProfiler showing the distribution Lamp1-mCherry signal

as a fraction of signal in each bin. Darker blue indicates a higher fraction of signal in that bin.

Scale bars = 50 μm. **(C)** Quantification of lysosomal number per cell (left), average vesicle area

(middle) and fraction of perinuclear lysosomes (right) using Lamp1-mCherry signal in vehicle

(DMSO) and bafilomycin A1 (100 nM) treated astrocytes. Emerald was used to determine cell

boundaries (n = 4; each replicate is an average over 10 images) **(D)** Representative images of

$PRKN^{+/+}$ (top) and $PRKN^{-/-}$ (bottom) astrocytes transduced with cytoplasmic Emerald and LAMP1-

mCherry lentiviruses. Nuclei were stained with Hoechst33342. Far right: Output from CellProfiler

showing the distribution Lamp1-mCherry signal as a fraction of signal in each bin. Darker blue

indicates a higher fraction of signal in that bin. Scale bars = 25 µm. (**E**) Quantification of Lamp1-

mCherry signal in *PRKN*[+/+] and *PRKN*[-/-] astrocytes to measure lysosomal number per cell (left),

average vesicle area (middle) and fraction of perinuclear lysosomes (right). Emerald fluorescence

was used to determine cell boundaries (n = 6; each replicate is an average over 7 images). All

images were acquired on a Nikon Ti2E AX R confocal microscope with a 20x objective lens and

6x optical zoom.

Fig. 4: Mitochondria-localized fluorophores resolve changes in mitochondrial network

dynamics (A) Schematic of the lentiviral construct used in the following panels localizing Emerald

to the mitochondria with the targeting sequence of COX8A. **(B)** Representative images of

astrocytes transduced with COX8A-Emerald and treated with a vehicle (DMSO; top) or 2 uM

oligomycin (bottom). Nuclei were stained with Hoechst33342. High magnification images (far

right) highlight the changes to the mitochondrial network upon oligomycin treatment. Scale bars

= 50 µm. **(C)** Quantification of mean mitochondrial area, mean mitochondrial branch length, and

mean aspect ratio upon oligomycin treatment (n = 3; each replicate is an average over 36 images)

(D) Representative images of mitochondria labeled with COX8A-Emerald moving along *PRKN*[-/-]

neuron axons in the anterograde (left) and retrograde (right) directions. Scale bars = 5 µm. Images

were taken on a Leica DMi8 microscope for 1 hour per field **(E-F)**. Quantification of mitochondrial

run length (left) and run time (right) of COX8A-Emerald labeled mitochondria in *PRKN*[+/+] and

PRKN[-/-] dopaminergic neuron axons moving anterograde (E) and retrograde (F) as determined

by kymograph analysis (average of n = 37.5 organoids for anterograde and 23.5 organoids for

retrograde; each replicate is an average over 1-3 fields).

Fig. 5: Timer fluorophore measures changes in subcellular mitochondrial turnover in

***PRKN*[-/-](A)** Schematic demonstrating how the fluorescent protein Timer works. Timer undergoes

a fluorescent shift from a green emission to a red emission overtime; analysis of Timer relies on

examining the ratio between the red and green emission. **(B)** Schematic of the lentiviral construct

used in the following panels localizing Timer to the mitochondria by using the targeting sequence

of COX8A. **(C)** Representative images of astrocytes transduced with COX8A-Timer and treated

with a vehicle (DMSO; top) or 100 nM bafilomycin A1 (bottom) for 24 hours. Images were acquired

with a 488-excitation line and a 594-excitation line. Far right: ratiometric representation of the red

channel divided by the green channel and pseudo colored so that orange, yellow and white

indicate more relative red emission and black and purple indicate more relative green emission.

Nuclei were stained with DRAQ5. Scale bars = 50 µm. **(D)** Quantification of the ratio of mean

intensity of emission from excitation with a 594nm laser to the mean intensity of emission from

excitation with a 488 nm laser in astrocytes expressing COX8A-Timer and treated with a vehicle

(DMSO) or 100 nM bafilomycin A1 for 24 hours (n = 6; each replicate is an average over 25

images). (**E**) Representative images of *PRKN*$^{+/+}$ and *PRKN*$^{-/-}$ astrocytes transduced with COX8A-

Timer. Images were acquired with a 488-excitation line and a 594-excitation line. Far right:

ratiometric representation of the red channel divided by the green channel and pseudo colored

so that orange, yellow and white indicate more relative red emission and black and purple indicate

more relative green emission. Nuclei were stained with DRAQ5. Scale bars = 50 µm. **(F)**

Quantification of the ratio of mean intensity of emission from excitation with a 594nm laser to the

mean intensity of emission from excitation with a 488 nm laser in *PRKN*$^{+/+}$ and *PRKN*$^{-/-}$ astrocytes

expressing COX8A-Timer (n = 3; each replicate is an average over 25 images). **(G)** Quantification

of mean intensity normalized by nuclear area of green and red fluorescence in *PRKN*$^{+/+}$ and

PRKN$^{-/-}$ astrocytes expressing COX8A-Timer (n = 3; each replicate is an average over 25 images).

All images were acquired on a CX7 HCS platform with a 20x objective lens.

Fig. 6: GRX1-roGFP2 fluorophore is used to quantify glutathione oxidation in chemical and

genetic models of oxidative stress (A) Schematic demonstrating how the fluorescent protein

fusion Grx1-roGFP2 works. Oxidation of Grx1 upon the oxidation of glutathione results in the

eventual oxidation of roGFP2, shifting its emission spectrum from green to blue. **(B)** Schematic

of the lentiviral construct used in the following panels localizing Grx1-roGFP2 to the mitochondria.

This fluorophore was localized using atp-9 instead of COX8A. **(C)** Representative images of

astrocytes transduced with mito-GRX1-roGFP2 and treated with a vehicle (DMSO; top) or 100

nM bafilomycin A1 (bottom) for 24 hours. Images were acquired with a 405-excitation laser and a

488-excitation laser. Far right: ratiometric representation of the green channel divided by the blue

channel and then pseudo colored so that orange, yellow and white indicate more relative green

emission while black and purple represent more blue emission. Nuclei were stained with DRAQ5.

Scale bars = 50 μm. **(D)** Quantification of the ratio of mean intensity from 488-excitation to the

mean intensity from 405 excitation. In astrocytes expression mito-Grx1-roGFP2 and treated with

a vehicle or 100 nM bafilomycin A1 (n = 6; each replicate is an average over 25 images). **(E)**

Quantification of mean intensity normalized by nuclear area of blue and green fluorescence in

vehicle and bafilomycin A1 treated astrocytes expressing mito-Grx1-roGFP2 (n = 6; each replicate

is an average of 25 images). **(F)** Representative images of $PRKN^{+/+}$ and $PRKN^{-/-}$ astrocytes

transduced with mito-Grx1-roGFP2. Images were acquired with a 405-excitation laser and a 588-

excitation laser. Far right: ratiometric representation of the green channel divided by the blue

channel and pseudo colored so that orange, yellow and white indicate more relative green

emission and black and purple indicate more relative blue emission. Nuclei were stained with

DRAQ5. Scale bars = 50 μm. **(G)** Quantification of the ratio of mean intensity of emission from

excitation with a 488nm laser to the mean intensity of emission from excitation with a 405 nm

laser in $PRKN^{+/+}$ and $PRKN^{-/-}$ astrocytes expressing mito-Grx1-roGFP2 (n = 3; each replicate is

an average over 16 images).

Supporting Information

Fig. S1: GEM-SCOPe supports a diversity of modifications and applications (A) Representative images of astrocytes expressing COX8A-Emerald under the control of a doxycycline inducible promoter. Astrocytes were treated with a vehicle (DMSO; top) or doxycycline (bottom) for 72 hours. Nuclei were stained with Hoechst33342. **(B)** qPCR for endogenous *LAMP1* on RNA from astrocytes transduced with a virus localized to the lysosome (LAMP1-mCherry) or a virus not localized to the lysosome (COX8A-Emerald). qPCR primers were designed to amplify a sequence of *LAMP1* that is not included in the lentivirus to only quantify endogenous *LAMP1* levels. n = 3 independent transductions **(C)** Representative images of a mixed population of astrocytes with nuclei expressing different H2B-fusion fluorophores. Astrocytes were transduced with a single lentivirus and then plated together. Scale bar = 50 μm.

Fig. S2: Mitochondrial pH sensor measures disrupted mitochondrial membrane potential Schematic of the lentiviral construct used in the following panels localizing Lemon, a pH responsive fluorophore, to the mitochondria with the targeting sequence of COX8A. Lemon undergoes Forster Resonance Energy Transfer (FRET) under alkaline conditions. **(B)** Representative images of astrocytes transduced with COX8A-Lemon and treated with a vehicle (DMSO; top) or 2 μM oligomycin (bottom) for 24 hours. Images were acquired with a 455nm-excitation laser and a 515-excitation laser. Far right: ratiometric representation of the emission in the yellow channel divided by the emission in the cyan channel and then pseudo colored so that orange, yellow and white indicate more relative yellow emission (more alkaline) while black and purple represent more cyan emission (more acidic). Nuclei were stained with DRAQ5. Scale bars = 50 μm. **(C)** Quantification of the ratio of yellow emission to cyan emission in astrocytes expressing COX8A-Lemon and treated with DMSO or oligomycin (n = 3; each replicate is an average 36 images).

Fig. S3: Astrocyte proliferation is activated by FBS and inhibited by MG-132 (A)
Representative images of astrocytes transduced with H2B-Emerald lentivirus after 0 hrs (top) or
48 hrs (bottom) of treatment with a vehicle (DMSO; left), 10% fetal bovine serum (FBS; middle),
or MG132 (right). Scale bars = 50 µm. **(B)** Quantification of number of H2B-Emerald nuclei per
image field 0 hours and 48 hours after treatment (n = 6 (n = 3 for MG-132 treatment) per time
point; each replicate is an average over 25 images). Scale bars = 50 µm. All images were acquired
on the CX7 HCS platform with a 20x objective lens.

**Fig. S4: Mitochondrial fragmentation exhibits a dose dependent response to oligomycin-
induced stress (A)** Representative images of astrocytes transduced with COX8A-Emerald and
treated with a vehicle (DMSO), 1µM, 2 µM, or 5 µM oligomycin for 4 hours. Nuclei were stained
with Hoechst33342. Scale bars = 50 µm **(B)** Quantification of mitochondrial features indicative of
mitochondrial network fragmentation: mean area, perimeter, aspect ratio, number of branches,
branch length, and number of branch endpoints (n = 3; each replicate is an average over 36
images).

Table S1: GEM-SCOPe lentiviral construct information A table listing all of the constructs
engineered for GEM-SCOPe, including information on AddGene catalog number, localization,
fluorophore, and antibiotic selection resistance genes.

**Video S1: Mitochondrial axonal transport in *PRKN*+/+ and *PRKN*-/- human iPSC derived
neurons** Representative videos of COX8A-Emerald labeled mitochondria moving within neuronal
axons over the course of 1 hour. Images were taken every 10 seconds and sped up 70x. Scale
bar = 25 µm.

References

1. Willis AW, Roberts E, Beck JC, Fiske B, Ross W, Savica R, et al. Incidence of Parkinson disease in North America. NPJ Parkinsons Dis. 2022;8(1):170.

2. Vazquez-Velez GE, Zoghbi HY. Parkinson's Disease Genetics and Pathophysiology. Annu Rev Neurosci. 2021;44:87-108.

3. Schneider SA, Obeso JA. Clinical and pathological features of Parkinson's disease. Curr Top Behav Neurosci. 2015;22:205-20.

4. Varadi C. Clinical Features of Parkinson's Disease: The Evolution of Critical Symptoms. Biology (Basel). 2020;9(5).

5. Jankovic J. Parkinson's disease: clinical features and diagnosis. J Neurol Neurosurg Psychiatry. 2008;79(4):368-76.

6. Naoi M, Maruyama W. Cell death of dopamine neurons in aging and Parkinson's disease. Mech Ageing Dev. 1999;111(2-3):175-88.

7. Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, Goedert M. Alpha-synuclein in Lewy bodies. Nature. 1997;388(6645):839-40.

8. Ohama E, Ikuta F. Parkinson's disease: distribution of Lewy bodies and monoamine neuron system. Acta Neuropathol. 1976;34(4):311-9.

9. Hartmann A. Postmortem studies in Parkinson's disease. Dialogues Clin Neurosci. 2004;6(3):281-93.

10. Helmchen F, Fee MS, Tank DW, Denk W. A miniature head-mounted two-photon microscope. high-resolution brain imaging in freely moving animals. Neuron. 2001;31(6):903-12.

11. Zong W, Wu R, Li M, Hu Y, Li Y, Li J, et al. Fast high-resolution miniature two-photon microscopy for brain imaging in freely behaving mice. Nat Methods. 2017;14(7):713-9.

12. Chen K, Tian Z, Kong L. Advances of optical miniscopes for in vivo imaging of neural activity in freely moving animals. Front Neurosci. 2022;16:994079.

13. Potashkin JA, Blume SR, Runkle NK. Limitations of animal models of Parkinson's disease. Parkinsons Dis. 2010;2011:658083.

14. Chia SJ, Tan EK, Chao YX. Historical Perspective: Models of Parkinson's Disease. Int J Mol Sci. 2020;21(7).

15. Ahfeldt T, Ordureau A, Bell C, Sarrafha L, Sun C, Piccinotti S, et al. Pathogenic Pathways in Early-Onset Autosomal Recessive Parkinson's Disease Discovered Using Isogenic Human Dopaminergic Neurons. Stem Cell Reports. 2020;14(1):75-90.

16. Raja WK, Neves E, Burke C, Jiang X, Xu P, Rhodes KJ, et al. Patient-derived three-dimensional cortical neurospheres to model Parkinson's disease. PLoS One. 2022;17(12):e0277532.

17. Chlebanowska P, Tejchman A, Sulkowski M, Skrzypek K, Majka M. Use of 3D Organoids as a Model to Study Idiopathic Form of Parkinson's Disease. Int J Mol Sci. 2020;21(3).

18. Kim H, Park HJ, Choi H, Chang Y, Park H, Shin J, et al. Modeling G2019S-LRRK2 Sporadic Parkinson's Disease in 3D Midbrain Organoids. Stem Cell Reports. 2019;12(3):518-31.

19. Shimomura O, Johnson FH, Saiga Y. Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusan, Aequorea. J Cell Comp Physiol. 1962;59:223-39.

20. Wang M, Da Y, Tian Y. Fluorescent proteins and genetically encoded biosensors. Chem Soc Rev. 2023;52(4):1189-214.

21. Ovechkina VS, Zakian SM, Medvedev SP, Valetdinova KR. Genetically Encoded Fluorescent Biosensors for Biomedical Applications. Biomedicines. 2021;9(11).

22. Terskikh A, Fradkov A, Ermakova G, Zaraisky A, Tan P, Kajava AV, et al. "Fluorescent timer": protein that changes color with time. Science. 2000;290(5496):1585-8.

23.   Nakai J, Ohkura M, Imoto K. A high signal-to-noise Ca(2+) probe composed of a single
green fluorescent protein. Nat Biotechnol. 2001;19(2):137-41.
24.   Burgstaller S, Bischof H, Gensch T, Stryeck S, Gottschalk B, Ramadani-Muja J, et al.
pH-Lemon, a Fluorescent Protein-Based pH Reporter for Acidic Compartments. ACS Sens.
2019;4(4):883-91.
25.   Gutscher M, Pauleau AL, Marty L, Brach T, Wabnitz GH, Samstag Y, et al. Real-time
imaging of the intracellular glutathione redox potential. Nat Methods. 2008;5(6):553-9.
26.   Nalls MA, Blauwendraat C, Vallerga CL, Heilbron K, Bandres-Ciga S, Chang D, et al.
Identification of novel risk loci, causal insights, and heritable risk for Parkinson's disease: a
meta-analysis of genome-wide association studies. Lancet Neurol. 2019;18(12):1091-102.
27.   Karimi-Moghadam A, Charsouei S, Bell B, Jabalameli MR. Parkinson Disease from
Mendelian Forms to Genetic Susceptibility: New Molecular Insights into the Neurodegeneration
Process. Cell Mol Neurobiol. 2018;38(6):1153-78.
28.   Jia F, Fellner A, Kumar KR. Monogenic Parkinson's Disease: Genotype, Phenotype,
Pathophysiology, and Genetic Testing. Genes (Basel). 2022;13(3).
29.   Guadagnolo D, Piane M, Torrisi MR, Pizzuti A, Petrucci S. Genotype-Phenotype
Correlations in Monogenic Parkinson Disease: A Review on Clinical and Molecular Findings.
Front Neurol. 2021;12:648588.
30.   Lesage S, Lunati A, Houot M, Romdhan SB, Clot F, Tesson C, et al. Characterization of
Recessive Parkinson Disease in a Large Multicenter Study. Ann Neurol. 2020;88(4):843-50.
31.   Pickrell AM, Youle RJ. The roles of PINK1, parkin, and mitochondrial fidelity in
Parkinson's disease. Neuron. 2015;85(2):257-73.
32.   Tanaka K. The PINK1-Parkin axis: An Overview. Neurosci Res. 2020;159:9-15.
33.   Kamienieva I, Duszynski J, Szczepanowska J. Multitasking guardian of mitochondrial
quality: Parkin function and Parkinson's disease. Transl Neurodegener. 2021;10(1):5.
34.   Dull T, Zufferey R, Kelly M, Mandel RJ, Nguyen M, Trono D, Naldini L. A third-generation
lentivirus vector with a conditional packaging system. J Virol. 1998;72(11):8463-71.
35.   Sarrafha L, Parfitt GM, Reyes R, Goldman C, Coccia E, Kareva T, Ahfeldt T. High-
throughput generation of midbrain dopaminergic neuron organoids from reporter human
pluripotent stem cells. STAR Protoc. 2021;2(2):100463.
36.   Morrone Parfitt G, Coccia E, Goldman C, Whitney K, Reyes R, Sarrafha L, et al.
Disruption of lysosomal proteolysis in astrocytes facilitates midbrain organoid proteostasis
failure in an early-onset Parkinson's disease model. Nat Commun. 2024;15(1):447.
37.   Chaudhry A, Shi R, Luciani DS. A pipeline for multidimensional confocal analysis of
mitochondrial morphology, function, and dynamics in pancreatic beta-cells. Am J Physiol
Endocrinol Metab. 2020;318(2):E87-E101.
38.   Mangeol P, Prevo B, Peterman EJ. KymographClear and KymographDirect: two tools for
the automated quantitative analysis of molecular and cellular dynamics using kymographs. Mol
Biol Cell. 2016;27(12):1948-57.
39.   Carpenter AE, Jones TR, Lamprecht MR, Clarke C, Kang IH, Friman O, et al.
CellProfiler: image analysis software for identifying and quantifying cell phenotypes. Genome
Biol. 2006;7(10):R100.
40.   Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A.
CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics. 2021;22(1):433.
41.   Lois C, Hong EJ, Pease S, Brown EJ, Baltimore D. Germline transmission and tissue-
specific expression of transgenes delivered by lentiviral vectors. Science. 2002;295(5556):868-
72.
42.   Schorpp M, Jäger R, Schellander K, Schenkel J, Wagner EF, Weiher H, Angel P. The
Human Ubiquitin C Promoter Directs High Ubiquitous Expression of Transgenes in Mice.
Nucleic Acids Research. 1996/05/01;24(9).

43. Zufferey R, Donello JE, Trono D, Hope TJ. Woodchuck Hepatitis Virus
Posttranscriptional Regulatory Element Enhances Expression of Transgenes Delivered by
Retroviral Vectors. Journal of Virology. 1999/04;73(4).
44. Higashimoto T, Urbinati F, Perumbeti A, Jiang G, Zarzuela A, Chang LJ, et al. The
woodchuck hepatitis virus post-transcriptional regulatory element reduces readthrough
transcription from retroviral vectors. Gene Therapy. 2007;14(17):1298-304.
45. Mosammaparast N, Jackson KR, Guo Y, Brame CJ, Shabanowitz J, Hunt DF, Pemberton
LF. Nuclear import of histone H2A and H2B is mediated by a network of karyopherins. J Cell
Biol. 2001;153(2):251-62.
46. Marino-Ramirez L, Kann MG, Shoemaker BA, Landsman D. Histone structure and
nucleosome stability. Expert Rev Proteomics. 2005;2(5):719-29.
47. Smith PJ, Wiltshire M, Errington RJ. DRAQ5 labeling of nuclear DNA in live and fixed
cells. Curr Protoc Cytom. 2004;Chapter 7:Unit 7 25.
48. Zhitomirsky B, Farber H, Assaraf YG. LysoTracker and MitoTracker Red are transport
substrates of P-glycoprotein: implications for anticancer drug design evading multidrug
resistance. J Cell Mol Med. 2018;22(4):2131-41.
49. Chazotte B. Labeling lysosomes in live cells with LysoTracker. Cold Spring Harb Protoc.
2011;2011(2):pdb prot5571.
50. Dang Q, Zhou H, Qian J, Yang L, Huang J, Zhang Y, Shi W. LAMP1 Overexpression
Predicts for Poor Prognosis in Diffuse Large B-cell Lymphoma. Clin Lymphoma Myeloma Leuk.
2018;18(11):749-54.
51. Wang QQ, Yao J, Jin Q, Wang XD, Zhu HJ, Huang F, et al. LAMP1 expression is
associated with poor prognosis in breast cancer. Oncol Lett. 2017;14(4):4729-35.
52. Xu Y, Cao X, Zhang S, Zhang Y, Shen Z. High expression of LAMP1 as a prognostic
marker in patients with epithelial ovarian cancer. Int J Clin Exp Pathol. 2017;10(8):9104-11.
53. Ranjbar MA, Jamshidi M. Overexpression of Lysosome-Associated Membrane Protein 1
in Oral Squamous Cell Carcinoma and its Correlation with Tumor Differentiation and Metastasis.
Iran J Otorhinolaryngol. 2022;34(120):3-8.
54. Zong S, Wu M, Gu J, Liu T, Guo R, Yang M. Structure of the intact 14-subunit human
cytochrome c oxidase. Cell Res. 2018;28(10):1026-34.
55. Cunatova K, Reguera DP, Houstek J, Mracek T, Pecina P. Role of cytochrome c oxidase
nuclear-encoded subunits in health and disease. Physiol Res. 2020;69(6):947-65.
56. Partikian A, Olveczky B, Swaminathan R, Li Y, Verkman AS. Rapid diffusion of green
fluorescent protein in the mitochondrial matrix. J Cell Biol. 1998;140(4):821-9.
57. Hoogewijs K, James AM, Smith RA, Gait MJ, Murphy MP, Lightowlers RN. Assessing the
Delivery of Molecules to the Mitochondrial Matrix Using Click Chemistry. Chembiochem.
2016;17(14):1312-6.
58. Candas D, Qin L, Fan M, Li JJ. Experimental Approaches to Study Mitochondrial
Localization and Function of a Nuclear Cell Cycle Kinase, Cdk1. J Vis Exp. 2016(108):53417.
59. Chazotte B. Labeling mitochondria with MitoTracker dyes. Cold Spring Harb Protoc.
2011;2011(8):990-2.
60. Durand RE, Olive PL. Cytotoxicity, Mutagenicity and DNA damage by Hoechst 33342. J
Histochem Cytochem. 1982;30(2):111-6.
61. Wiezorek C. Cell cycle dependence of Hoechst 33342 dye cytotoxicity on sorted living
cells. Histochemistry. 1984;81(5):493-5.
62. Solano RM, Casarejos MJ, Menendez-Cuervo J, Rodriguez-Navarro JA, Garcia de
Yebenes J, Mena MA. Glial dysfunction in parkin null mice: effects of aging. J Neurosci.
2008;28(3):598-611.
63. Rouland L, Duplan E, Ramos Dos Santos L, Bernardin A, Katula KS, Manfioletti G, et al.
Therapeutic potential of parkin as a tumor suppressor via transcriptional control of cyclins in
glioblastoma cell and animal models. Theranostics. 2021;11(20):10047-63.

64.   Wahabi K, Perwez A, Rizvi MA. Parkin in Parkinson's Disease and Cancer: a Double-
Edged Sword. Mol Neurobiol. 2018;55(8):6788-800.
65.   Yeo CW, Ng FS, Chai C, Tan JM, Koh GR, Chong YK, et al. Parkin pathway activation
mitigates glioma cell proliferation and predicts patient survival. Cancer Res. 2012;72(10):2543-
53.
66.   Tay SP, Yeo CW, Chai C, Chua PJ, Tan HM, Ang AX, et al. Parkin enhances the
expression of cyclin-dependent kinase 6 and negatively regulates the proliferation of breast
cancer cells. J Biol Chem. 2010;285(38):29231-8.
67.   Gong Y, Zack TI, Morris LG, Lin K, Hukkelhoven E, Raheja R, et al. Pan-cancer genetic
analysis identifies PARK2 as a master regulator of G1/S cyclins. Nat Genet. 2014;46(6):588-94.
68.   Veeriah S, Taylor BS, Meng S, Fang F, Yilmaz E, Vivanco I, et al. Somatic mutations of
the Parkinson's disease-associated gene PARK2 in glioblastoma and other human
malignancies. Nat Genet. 2010;42(1):77-82.
69.   Frade JM, Ovejero-Benito MC. Neuronal cell cycle: the neuron itself and its
circumstances. Cell Cycle. 2015;14(5):712-20.
70.   Kruman, II, Wersto RP, Cardozo-Pelaez F, Smilenov L, Chan SL, Chrest FJ, et al. Cell
cycle activation linked to neuronal cell death initiated by DNA damage. Neuron. 2004;41(4):549-
61.
71.   Gupta R, Jha A, Ambasta RK, Kumar P. Regulatory mechanism of cyclins and cyclin-
dependent kinases in post-mitotic neuronal cell division. Life Sci. 2021;285:120006.
72.   Gan-Or Z, Dion PA, Rouleau GA. Genetic perspective on the role of the autophagy-
lysosome pathway in Parkinson disease. Autophagy. 2015;11(9):1443-57.
73.   Navarro-Romero A, Montpeyo M, Martinez-Vicente M. The Emerging Role of the
Lysosome in Parkinson's Disease. Cells. 2020;9(11).
74.   Hopfner F, Mueller SH, Szymczak S, Junge O, Tittmann L, May S, et al. Rare Variants in
Specific Lysosomal Genes Are Associated With Parkinson's Disease. Mov Disord.
2020;35(7):1245-8.
75.   Dehay B, Martinez-Vicente M, Caldwell GA, Caldwell KA, Yue Z, Cookson MR, et al.
Lysosomal impairment in Parkinson's disease. Mov Disord. 2013;28(6):725-32.
76.   Klein AD, Mazzulli JR. Is Parkinson's disease a lysosomal disorder? Brain.
2018;141(8):2255-62.
77.   Yoshimori T, Yamamoto A, Moriyama Y, Futai M, Tashiro Y. Bafilomycin A1, a specific
inhibitor of vacuolar-type H(+)-ATPase, inhibits acidification and protein degradation in
lysosomes of cultured cells. J Biol Chem. 1991;266(26):17707-12.
78.   Mauvezin C, Neufeld TP. Bafilomycin A1 disrupts autophagic flux by inhibiting both V-
ATPase-dependent acidification and Ca-P60A/SERCA-dependent autophagosome-lysosome
fusion. Autophagy. 2015;11(8):1437-8.
79.   Johnson DE, Ostrowski P, Jaumouille V, Grinstein S. The position of lysosomes within
the cell determines their luminal pH. J Cell Biol. 2016;212(6):677-92.
80.   Heuser J. Changes in lysosome shape and distribution correlated with changes in
cytoplasmic pH. J Cell Biol. 1989;108(3):855-64.
81.   Parton RG, Dotti CG, Bacallao R, Kurtz I, Simons K, Prydz K. pH-induced microtubule-
dependent redistribution of late endosomes in neuronal and epithelial cells. J Cell Biol.
1991;113(2):261-74.
82.   Pu J, Guardia CM, Keren-Kaplan T, Bonifacino JS. Mechanisms and functions of
lysosome positioning. J Cell Sci. 2016;129(23):4329-39.
83.   Song P, Trajkovic K, Tsunemi T, Krainc D. Parkin Modulates Endosomal Organization
and Function of the Endo-Lysosomal Pathway. J Neurosci. 2016;36(8):2425-37.
84.   Okarmus J, Bogetofte H, Schmidt SI, Ryding M, Garcia-Lopez S, Ryan BJ, et al.
Lysosomal perturbations in human dopaminergic neurons derived from induced pluripotent stem
cells with PARK2 mutation. Sci Rep. 2020;10(1):10278.

85. Twig G, Elorza A, Molina AJ, Mohamed H, Wikstrom JD, Walzer G, et al. Fission and selective fusion govern mitochondrial segregation and elimination by autophagy. EMBO J. 2008;27(2):433-46.

86. Chen H, Chomyn A, Chan DC. Disruption of fusion results in mitochondrial heterogeneity and dysfunction. J Biol Chem. 2005;280(28):26185-92.

87. Wang X, Winter D, Ashrafi G, Schlehe J, Wong YL, Selkoe D, et al. PINK1 and Parkin target Miro for phosphorylation and degradation to arrest mitochondrial motility. Cell. 2011;147(4):893-906.

88. Hernandez G, Thornton C, Stotland A, Lui D, Sin J, Ramil J, et al. MitoTimer: a novel tool for monitoring mitochondrial turnover. Autophagy. 2013;9(11):1852-61.

89. Laker RC, Xu P, Ryall KA, Sujkowski A, Kenwood BM, Chain KH, et al. A novel MitoTimer reporter gene for mitochondrial content, structure, stress, and damage in vivo. J Biol Chem. 2014;289(17):12005-15.

90. Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY, Remington SJ. Investigating mitochondrial redox potential with redox-sensitive green fluorescent protein indicators. J Biol Chem. 2004;279(13):13044-53.

91. Yokomakura A, Hong J, Ohuchi K, Oh SE, Lee JY, Mano N, et al. Increased production of reactive oxygen species by the vacuolar-type (H(+))-ATPase inhibitors bafilomycin A1 and concanamycin A in RAW 264 cells. J Toxicol Sci. 2012;37(5):1045-8.

92. Palacino JJ, Sagi D, Goldberg MS, Krauss S, Motz C, Wacker M, et al. Mitochondrial dysfunction and oxidative damage in parkin-deficient mice. J Biol Chem. 2004;279(18):18614-22.

93. Kim S, Pajarillo E, Nyarko-Danquah I, Aschner M, Lee E. Role of Astrocytes in Parkinson's Disease Associated with Genetic Mutations and Neurotoxicants. Cells. 2023;12(4).

94. Cornejo-Olivas MR, Torres L, Mata IF, Mazzetti P, Rivas D, Cosentino C, et al. A Peruvian family with a novel PARK2 mutation: Clinical and pathological characteristics. Parkinsonism Relat Disord. 2015;21(5):444-8.

95. Orimo S, Amino T, Yokochi M, Kojo T, Uchihara T, Takahashi A, et al. Preserved cardiac sympathetic nerve accounts for normal cardiac uptake of MIBG in PARK2. Mov Disord. 2005;20(10):1350-3.

96. Gouider-Khouja N, Larnaout A, Amouri R, Sfar S, Belal S, Ben Hamida C, et al. Autosomal recessive parkinsonism linked to parkin gene in a Tunisian family. Clinical, genetic and pathological study. Parkinsonism Relat Disord. 2003;9(5):247-51.

97. Hayashi S, Wakabayashi K, Ishikawa A, Nagai H, Saito M, Maruyama M, et al. An autopsy case of autosomal-recessive juvenile parkinsonism with a homozygous exon 4 deletion in the parkin gene. Mov Disord. 2000;15(5):884-8.

98. Schmidt S, Linnartz B, Mendritzki S, Sczepan T, Lubbert M, Stichel CC, Lubbert H. Genetic mouse models for Parkinson's disease display severe pathology in glial cell mitochondria. Hum Mol Genet. 2011;20(6):1197-211.

99. Gerasimova T, Stepanenko E, Novosadova L, Arsenyeva E, Shimchenko D, Tarantul V, et al. Glial Cultures Differentiated from iPSCs of Patients with PARK2-Associated Parkinson's Disease Demonstrate a Pro-Inflammatory Shift and Reduced Response to TNFalpha Stimulation. Int J Mol Sci. 2023;24(3).

100. Hastie E, Cataldi M, Marriott I, Grdzelishvili VZ. Understanding and altering cell tropism of vesicular stomatitis virus. Virus Res. 2013;176(1-2):16-32.

101. Sweeney NP, Vink CA. The impact of lentiviral vector genome size and producer cell genomic to gag-pol mRNA ratios on packaging efficiency and titre. Mol Ther Methods Clin Dev. 2021;21:574-84.

102. Desfarges S, Ciuffi A. Retroviral integration site selection. Viruses. 2010;2(1):111-30.

103. Bonam SR, Wang F, Muller S. Lysosomes as a therapeutic target. Nat Rev Drug Discov. 2019;18(12):923-48.

104.   Zhang Z, Yue P, Lu T, Wang Y, Wei Y, Wei X. Role of lysosomes in physiological
activities, diseases, and therapy. J Hematol Oncol. 2021;14(1):79.
105.   Javadov S, Kozlov AV, Camara AKS. Mitochondria in Health and Diseases. Cells.
2020;9(5).
106.   Chen W, Zhao H, Li Y. Mitochondrial dynamics in health and disease: mechanisms and
potential targets. Signal Transduct Target Ther. 2023;8(1):333.

1052

Figure 1

Figure 2

A.

| UbC | H2B (Nuclear) | mTagBFP2 | P2A - PuromycinR |

B.

| DRAQ5 | H2B-mTagBFP2 | Composite |

1 hour

24 hours

C.

■ DRAQ5 ■ H2B-BFP

%Stain Localized to Cytoplasm

ns

****

ns

****

1 hr   24 hrs

Hours Since DRAQ5 Addition

D.

| UbC | H2B (Nuclear) | mEmerald | P2A - PuromycinR |
| UbC | H2B (Nuclear) | mCherry | P2A - PuromycinR |

E.

| *PRKN*⁺/⁺ H2B-Emerald | *PRKN*⁻/⁻ H2B-mCherry | Composite |

24 hours

72 hours

F.

Number of Nuclei

*

***

ns

24hrs   72hrs

Hours Since Seeding

● $PRKN^{+/+}$ ● $PRKN^{-/-}$

Figure 3

Figure 4

A.

| UbC | Cox8A (Mitochondria) | mEmerald | Empty (No Selection) |

B.

COX8A-Emerald | Composite | High Mag. Composite

0 µM Oligomycin

2 µM Oligomycin

C.

Mean Mitochondrial Area (µm²)
Vehicle   Olig.
***

Mean Branch Length (µm)
Vehicle   Olig.
****

Mean Aspect Ratio
Vehicle   Olig.
****

D.

Anterograde Transport
0s
6s
13s
5um
← Soma   Synapse →

Retrograde Transport
0s
14s
34s
← Soma   Synapse →

E.

Anterograde Transport

Run Length (µm)
$PRKN^{+/+}$ $PRKN^{-/-}$
***

Run Time (s)
$PRKN^{+/+}$ $PRKN^{-/-}$
**

F.

Retrograde Transport

Run Length (µm)
$PRKN^{+/+}$ $PRKN^{-/-}$
ns

Run Time (s)
$PRKN^{+/+}$ $PRKN^{-/-}$
*

Figure 5

Figure 6

Figure S1

A.

| tetCMV COX8A-Emerald | Composite |

Vehicle

Doxycycline

B.

| H2B-mTagBFP2 | H2B-Emerald | H2B-mCherry | Composite |

C.

Endogenous LAMP1

ns

ΔΔCt (Normalized to Actin)

COX8A-Emerald   LAMP1-mCherry

Figure S2

Figure S3

Figure S4

Plasmid Name	Nickname	AddGene Cat#	Localization	Fluorophore	Antibiotic Resistance	Relevant Figure(s)
pFUW H2B-mTagBFP2-p2A-PuroR	H2B-mTagBFP2		Nucleus	mTagBFP2	Puromycin	1, 2, S1, S3
pFUW H2B-Emerald-p2A-PuroR	H2B-Emerald		Nucleus	Emerald	Puromycin	2, S1
pFUW H2B-mCherry-p2A-PuroR	H2B-mCherry		Nucleus	mCherry	Puromycin	2, S1
pFUW LAMP1Sig-Emerald	LAMP1-Emerald		Lysosome	Emerald	NA	1, 3
pFUW LAMP1Sig-mCherry	LAMP1-mCherry		Lysosome	mCherry	NA	1
pFUW COX8ASig-Emerald	COX8A-Emerald		Mitochondria	Emerald	NA	1, 4, S5
pFUW COX8ASig-mCherry	COX8A-mCherry		Mitochondria	mCherry	NA	1
pFUW PuroR-P2A-COX8ASig-Timer	COX8A-Timer		Mitochondria	Timer	Puromycin	5
pFUW COX8ASig-Lemon	COX8A-Lemon		Mitochondria	Lemon	NA	S2
pFUW mito-GRX1-roGFP2	mito-GRX1-roGFP2		Mitochondria	roGFP2	NA	6
pFUW mTagBFP2	mTagBFP2		NA	mTagBFP2	NA	NA
pFUW Emerald	Emerald		NA	Emerald	NA	NA
pFUW mCherry	mCherry		NA	mCherry	NA	NA
pFUW mTagBFP2-P2A-PuroR	mTagBFP2-PuroR		NA	mTagBFP2	Puromycin	NA
pFUW Emerald-P2A-PuroR	Emerald-PuroR		NA	Emerald	Puromycin	3
pFUW mCherry-P2A-PuroR	mCherry-PuroR		NA	mCherry	Puromycin	NA
pFUW mTagBFP2-P2A-NeoR	mTagBFP2-NeoR		NA	mTagBFP2	Neomycin	NA
pFUW Emerald-P2A-NeoR	Emerald-NeoR		NA	Emerald	Neomycin	NA
pFUW mCherry-P2A-NeoR	mCherry-NeoR		NA	mCherry	Neomycin	NA