Genetically Encoded and Modular SubCellular Organelle Probes (GEM-

SCOPe) reveal lysosomal and mitochondrial dysfunction driven by *PRKN*

knockout

Camille Goldman[1-5], Tatyana Kareva[1-5], Lily Sarrafha[1-5], Braxton R. Schuldt[1-5], Abhishek

Sahasrabudhe[1-3], Tim Ahfeldt[1-5] **, Joel W. Blanchard[1-6] *

1.　Icahn School of Medicine, Mount Sinai, New York, NY, USA

2.　Nash Family Department of Neuroscience, Mount Sinai, New York, NY, USA

3.　Friedman Brain Institute, Mount Sinai, New York, NY, USA

4.　Ronald M. Loeb Center for Alzheimer's Disease, Mount Sinai, New York, NY USA

5.　Black Family Stem Cell Institute, Mount Sinai, New York, NY, USA

6.　Lead Contact

* Corresponding author

** 2nd corresponding author

Correspondence: joel.blanchard@mssm.edu

Correspondence: ahfeldt@gmail.com

Summary

Cellular processes including lysosomal and mitochondrial dysfunction are implicated in the development of many diseases. Quantitative visualization of mitochondria and lysosoesl is crucial to understand how these organelles are dysregulated during disease. To address a gap in live-imaging tools, we developed GEM-SCOPe (Genetically Encoded and Modular SubCellular Organelle Probes), a modular toolbox of fluorescent markers designed to inform on localization, distribution, turnover, and oxidative stress of specific organelles. We expressed GEM-SCOPe in differentiated astrocytes and neurons from a human pluripotent stem cell *PRKN*-knockout model of Parkinson's disease and identified disease-associated changes in proliferation, lysosomal distribution, mitochondrial transport and turnover, and reactive oxygen species. We demonstrate GEM-SCOPe is a powerful panel that provide critical insight into the subcellular mechanisms underlying Parkinson's disease in human cells. GEM-SCOPe can be expanded upon and applied to a diversity of cellular models to glean an understanding of the mechanisms that promote disease onset and progression.

Keywords

Live-cell imaging, induced pluripotent stem cells, Parkinson's Disease, astrocytes, PRKN

Introduction

Parkinson's Disease (PD) is a common neurodegenerative disease that robs individuals of their motor and cognitive functions, affecting over 10 million people worldwide and with an estimated 90,000 new diagnoses in the United States each year[1]. PD is clinically characterized by bradykinesia, resting tremor, rigidity, and postural instability[2-5] and pathologically characterized by the loss of dopaminergic neurons in the substantia nigra pars compacta and buildup of intracellular deposits of α-synuclein into insoluble protein aggregates called Lewy Bodies[2,6-8]. From the first pathological descriptions of PD in the mid-20th century, advancements in many fields, including histology, genetics, cell biology, and neurology enabled insight into many aspects of the disease, from determining genetic and environmental risk factors and mapping disease progression to identifying pathways and proteins to target therapeutically. Despite great progress, there is still incomplete understanding of the cellular dysfunction that precedes and contributes to chronic neuronal death.

This limited advancement arises in part due to the absence of tools used to track molecular and cellular changes in real-time throughout disease progression. Studies on human post-mortem brain tissue are largely representative of end-stage disease states, making it difficult to dissect which phenotypes are a primary, causative effect and which phenotypes are a secondary response[9]. Mouse models offer more opportunities to study multiple time points throughout disease progression as motor and cognitive symptoms can be monitored over time. However, targeting specific cell types in mice remains technically challenging and time-consuming.

Emerging technologies such as the miniscope allow continuous monitoring of fluorescence-based

readouts on a cellular level in live mouse and rat brains[10-12]. However, the type of readout from

the miniscope has been restricted to neuronal activity from fluorescent calcium sensors and

resolution along the z-axis remains a limiting factor in analyzing deeper cortical tissue.

Regardless of such technologies, current genetic rodent models of PD fail to completely

recapitulate human disease and lack key pathological hallmarks. Drug-induced models can

recapitulate late-stage disease phenotypes and behaviors, but due to their acute onset, are

inadequate to study the gradual accumulation of cellular dysfunction that precedes clinical

presentation[13,14]. Advancements in induced pluripotent stem cell (iPSC) differentiation and

CRISPR/Cas9 genome editing have led to the emergence of genetic cellular models of PD[15-18].

These models enable researchers to study the effects of genetic and environmental factors on

specific human cell populations relevant to PD.

Cellular models of PD provide an exciting opportunity to study subcellular changes and responses

to stress or treatment in live cells. Cells can be genetically engineered to express fluorescent

proteins that can then be visualized by microscopy in live cells. The applicability of GFP as a

reporter gene was immediately appreciated after its identification and isolation in 1962[19]. Since

then, hundreds of fluorophores have been developed by introducing mutations to naturally

occurring fluorophores isolated from sea anemones and jellyfish to modulate fluorescence

intensity, half-life, and excitation/emission spectra[20]. Additional fluorophores have been

developed as biosensors, by fluorescing or changing fluorescence emission in the presence of a

certain ligand or under specific cellular conditions[20-25]. Cells can express these fluorophores with

virtually limitless possibilities with regards to subcellular localization, cell-type specific expression,

and temporally controlled expression. Genetically encoded fluorescent proteins can be used to

follow a population of cells as they change due to aging, genetics, or environmental stressors and

stimuli.

PD has a complex genetic architecture underlying the disease. Over 90% of PD cases are

idiopathic; over 90 genetic risk loci have been identified by GWAS, but any of those individual

variants confer little risk on their own[26]. The remaining 10% of PD cases can be linked to highly

penetrant mutations in a handful of genes. While extremely rare, studying the function of these

genes and their impact on cellular processes sheds light on the fundamental pathways that drive

PD pathogenesis. Here, we focus on the *PRKN* gene. Autosomal recessive loss of function

mutations in *PRKN* are the most frequent known genetic cause of early onset cases of PD,

accounting for about 15% of PD cases with onset before the age of 50[27-30]. *PRKN* encodes for

the protein PRKN, an E3 ubiquitin ligase associated with mitochondria. PRKN directs the

ubiquitination of outer mitochondrial membrane proteins on damaged mitochondria, targeting

them for degradation via the autophagy-lysosomal pathway (mitophagy)[31-33]. While the function

of PRKN is well understood, it is still unclear how loss of PRKN function leads to the gradual

disease progression of early onset PD. It is believed that neurodegeneration can begin over 20

years before clinical onset, with a prolonged prodromal phase associated with several non-motor

symptoms. Thus, it is crucial to study models of early disease to understand what cellular

disturbances in the prodromal phase culminate in neurodegeneration.

Here, we present GEM-SCOPe (**G**enetically **E**ncoded and **M**odular **S**ub**C**ellular **O**rganelle

Probes), an expandable panel of genetically encoded reporters to track and quantify the

subcellular phenotypes in live cells. We developed a library of constructs that localize specifically

to the nucleus, mitochondria, and lysosomes and included fluorophores with a variety of emission

spectra or biosensor capabilities. All the lentiviral constructs were developed in the same

backbone vector to be modular, enabling any component, be it promotor, localization signal,

fluorophore, or antibiotic resistance, to be easily removed or replaced. We validated the

fluorophores with live-imaging of human induced pluripotent stem cell (hiPSC)-derived midbrain

astrocytes employing existing chemical dyes and chemical perturbations. We then applied GEM-

SCOPe to hiPSC-derived astrocytes and neurons from a *PRKN*-knockout model of PD[15]. This

revealed previously unseen changes in cellular proliferation, lysosomal distribution, mitochondrial

motility and turnover, and reactive oxygen species production. Our results demonstrate the

widespread utility of GEM-SCOPe to study a variety of translationally important subcellular

changes affected during PD. The modular, combinatorial flexibility of GEM-SCOPe can be

adapted to investigate any disease, and, therefore, it is a critical new resource that can be broadly

applied in neurodegenerative research and beyond.

Results

Building a modular toolbox for the subcellular localization of genetically encoded fluorophores

We developed GEM-SCOPe for live-cell imaging of subcellular organelle dynamics. We aimed to

produce a system that is highly modular to easily customize the expression, localization,

fluorescence, and selection for any given experimental need. We started with the FUW

backbone[34], a 3rd generation lentivirus backbone with a ubiquitin C promoter for high, ubiquitous

expression of the transgene across cell types[34,35] and a woodchuck hepatitis virus post-

transcriptional regulatory element (WPRE) for improved lentiviral expression[34,36,37]. The designed

lentiviral constructs contain four components: promoter, localization sequence, fluorophore, and

antibiotic resistance (**Fig. 1A**). The content of these four components can be mixed and matched

to generate a library of lentiviral constructs to fit a diverse field of experimental needs. To generate

constructs *de novo*, components were combined and inserted into the plasmid via Gibson

assembly (**Fig. 1B, top**). Restriction sites were kept between each domain, so each component

or group of components could be excised via restriction digest and replaced with a different

component, either by restriction site cloning or Gibson assembly (**Fig. 1B, bottom**). Thus, each

component of the lentiviral plasmid can be removed or replaced to accommodate features such

as cell-type specificity with cell-type specific promoters, alternative subcellular localization, and

unique fluorophores.

The FUW backbone has a ubiquitin C promoter, but this can be replaced with lineage or cell-type-

specific promoters for selective expression in specific cell types or with drug-inducible promoters

to temporally control when the construct is transcribed (**Fig. S1A**). Subcellular localization is

determined by including the whole or partial coding sequence of a peptide with a known and

specific subcellular localization. This domain can be omitted to produce a fluorophore expressed

in the cytoplasm. The fluorophore component of the construct offers the most flexibility, from basic

fluorescent proteins to more complex fluorophores that change fluorescent excitation and/or

emission wavelength with time or environment. All plasmids generated for GEM-SCOPe have

been deposited to Addgene(**Table S1**).

We developed GEM-SCOPe constructs to localize fluorophores to the cytoplasm (no tag), nucleus

(H2B-tag), mitochondria (COX8A-tag), and lysosome (LAMP1-tag) (**Fig. 1C**). Fluorophore

localization was validated using commercially available live cellular dyes. Nuclear localization was

achieved by fusing fluorophores to the C-terminus of histone 2B (H2B). H2B is an integral

chromatin protein that is found in all cells and is localized specifically to the nucleus[38,39]. H2B

fused to mTagBFP2, a blue fluorophore, (H2B-mTagBFP2) co-localizes with DRAQ5, a far-red

live-cell nuclear dye[40] (**Fig. 1C, top**; Mander's coefficient > 0.99±0.0001). In addition to the

mTagBFP2 construct, we also developed lentiviral plasmids with H2B fused to Emerald (green;

H2B-Emerald) and mCherry (red; H2B-mCherry) (**Fig. 1A, S1B**).

To localize fluorophores to the lysosome, we utilized the N-terminal peptide sequence of

lysosomal-associated membrane protein 1 (LAMP1), a major component of lysosomal

membranes that plays a key role in lysosomal biogenesis and homeostasis. The N-terminus of

LAMP1 fused to mCherry (LAMP1-mCherry) co-localizes with LysoTracker (Thermo Scientific)

(**Fig. 1C, middle**; Mander's coefficient = 0.92±0.03). LysoTracker is a fluorophore that is partially

protonated at neutral pH and can readily cross membranes; once protonated it can no longer

diffuse so the fluorescent signal gets trapped in acidic compartments, including, but not limited to,

the lysosome[41,42]. Overexpression of LAMP1 is associated with numerous cancers and cancer

metastasis[43-46]. By only using a partial sequence of LAMP1 and not the full peptide sequence, we

are able to target the fluorophore specifically to the lysosome without increasing the expression

of endogenous LAMP1 (**Fig. S1C**). We also developed lentiviral plasmids with the LAMP1

targeting sequence fused to Emerald (green; LAMP1-Emerald; **Fig. 1A, 1D**).

Finally, mitochondrial localization was achieved using the mitochondrial targeting sequence of

COX8A. COX8A is a nuclear-encoded subunit of cytochrome-c oxidase (Complex IV) in the

electron transport chain[47,48]. The first 25 amino acid residues of the COX8A peptide sequence

target the peptide to the mitochondrial inner membrane and are widely used to specifically localize

proteins and fluorophores to the mitochondria[49-51]. To confirm mitochondrial specificity of COX8A-

Emerald we compared localization with MitoTracker (Thermo Scientific), a dye that accumulates

in the mitochondria due to the negative membrane potential of the inner mitochondrial

membrane[41,52]. The mitochondrial targeting sequence fused to an Emerald fluorophore (COX8A-

Emerald) exhibited significant co-localization with MitoTracker (**Fig. 1C, bottom**; Mander's

coefficient = 0.97±0.003). We developed lentiviral plasmids with the COX8A targeting sequence

fused to mCherry (red; COX8A-mCherry; **Fig. 1A, 1D**), Timer (green/red; COX8A-Timer; **Fig. 1A,**

Fig. 5), and Lemon (cyan/yellow; COX8A-Lemon; **Fig. 1A, Fig. S2**).

Once the localization strategies were finalized, we swapped out the fluorophores with different

fluorescent proteins that are excited and emitted at alternative wavelengths. Mixing localization

sequences and fluorophores with unique emission spectra allows simultaneous imaging of

multiple sub-cellular processes in the same cells such as mitochondrial network and lysosomal

distribution (**Fig. 1D**). Utilizing a combination of subcellular localization and diverse fluorophores,

we developed GEM-SCOPe, a fluorescent, live-cell imaging toolbox of lentiviral constructs that

were systematically validated to quantify cellular proliferation, lysosomal distribution, and

mitochondrial dynamics in human Parkinson's disease astrocytes.

H2B-fused fluorophores offer improvements for efficiently labelling nuclei for multi-day live-cell assays

H2B-fused fluorophores offer improvements for efficiently

labelling nuclei for multi-day live-cell assays

Quantification of cellular phenotypes by fluorescence microscopy is dependent on normalization

to total cell number. This is often accomplished by counting the number of nuclei. A few fluorescent

dyes that can cross the cellular and nuclear membranes of live cells and bind specifically to DNA

are commercially available. While efficient for short-term and end-point assays, such dyes

intercalate in DNA and therefore have cytotoxic and mutagenic effects[53,54]. Furthermore, nuclear

dyes do not remain contained within the nuclear compartment, and their signal leaches into the

cytoplasm in just a few hours after exposing the cells. The cytotoxicity and loss of precise labeling

over time render these dyes less useful for long-term microscopy or cell-tracking experiments. To

address these concerns, we developed lentiviral constructs expressing nuclear-localized

fluorophores by fusing the nucleus-specific histone protein, H2B, with either mTagBFP2, Emerald,

or mCherry (**Fig. 1C, 2A**). Human induced pluripotent stem-cell (hiPSC)-derived astrocytes

transfected with H2B-mTagBFP2 were co-stained with DRAQ5, a far-red DNA-stain for live cells.

The localization of the two nuclear signals was compared 1 hours and 24 hours after the addition

of DRAQ5. Immediately after adding DRAQ5, there was no difference between the DRAQ5 stain

and the H2B-mTagBFP2 (2-way ANOVA with Tukey's HSD; adj-p = 0.83) (**Fig. 2B, C**). Over the

experimental period, the DRAQ5 signal spread significantly (26±5% extranuclear; adj-p < 0.0001),

while the extranuclear H2B-mTagBFP2 signal did not significantly change (adj-p = 0.99) (**Fig. 2B,**

C). While the human eye can still readily distinguish the nucleus from the extranuclear stain in the

DRAQ5 images, image analysis programs that rely on grey values for segmentation often fail to

correctly differentiate the true nuclear signal from the extranuclear signal. Thus, lentiviral

constructs expressing nuclear-localized fluorophores, such as the demonstrated H2B-

mTagBFP2, can provide better alternatives for long-term live-imaging experiments with multiple

time points.

207 H2B-fused fluorophores reveal increased cellular

208 proliferation in PRKN knockout astrocytes

While dopaminergic neuron death is the predominant cellular outcome in Parkinson's Disease

(PD), other cell types in the brain may change cell cycle regulation under neurotoxic conditions.

Tracking cell numbers to monitor proliferation, senescence, or death in vitro can help researchers

understand how different cell populations are responding to neurotoxic stress or potential

therapeutic strategies.

First, we validated that our nuclear-localized construct, H2B-Emerald, could be used to track

cellular proliferation. The proliferation of H2B-Emerald transduced hiPSC-derived astrocytes was

induced using fetal bovine serum (FBS) or blocked by inhibiting the proteasome with MG-132.

Astrocyte proliferation was measured by changes in the number of H2B-Emerald nuclei. As

expected, the astrocytes treated with FBS proliferated twice as rapidly as the untreated astrocytes

over 48 hours (2-way ANOVA with Tukey's HSD, p-adj = 0.005, **Fig. S3**), while the astrocytes

treated with MG-132 did not significantly proliferate over the same period (adj-p = 0.08, **Fig. S3**).

Thus, the GEM-SCOPe H2B-fluorophore fusion proteins can be used to track cellular proliferation

over multiple days in live cell cultures.

To examine the effect of loss of *PRKN* on astrocyte proliferation, we transduced wild-type

(*PRKN*[+/+]) astrocytes with H2B-Emerald and isogenic *PRKN* knockout (*PRKN*[-/-]) astrocytes with

H2B-mCherry (**Fig. 2D**). Leveraging the modularity of GEM-SCOPe, we were able to mix the two

astrocyte populations together and track their proliferation simultaneously. Over 48 hours, the

PRKN[-/-] population increased significantly (2-way ANOVA with Tukey's HSD, p-adj = 0.03; **Fig.**

2E, F) while the *PRKN*[+/+] population did not proliferate in the same period (2-way ANOVA with

Tukey's HSD, p-adj = 0.4; **Fig. 2E, F**). These results contradict mouse studies that found that

PRKN-null mice had decreased astroglia proliferation[55]. However, PRKN has also been described

as a tumor suppressor, with loss of function mutations arising in a variety of cancers, including

glioblastomas[56-58]. Studies on PRKN in the context of cancer find that cancer regulates the

degradation of cyclins and cyclin-dependent kinases involved in the G1/S transition, thus

promoting continued proliferation[57,59,60]. It has been hypothesized that the outcome of loss of

function PRKN mutations are highly cell-context dependent; the increased half-life of cyclins leads

to cancer in mitotically competent cells while the same cyclins promote apoptosis upon cell cycle

reentry in post-mitotic neurons[57,61-64]. Our findings highlight the importance of studying disease-

relevant human cells to capture the intricacies and nuances specific to the human brain that could

contribute to PD.

LAMP1-localized fluorophores reveal changes in lysosomal

distribution in response to chemical and genetic stressors

Lysosomal dysfunction is widely implicated in PD and other forms of neurodegeneration. Several

PD-risk and familial genes are involved in the autophagy-lysosomal-endosomal pathway[65-67], and

many hallmarks of lysosomal dysfunction associated with familial PD are present in idiopathic PD

[66,68,69]. Disruption to the autophagy-lysosomal-endosomal pathway is of broad importance in PD

and there is a crucial need to study lysosomal function and dysfunction in the context of PD

pathogenesis. As described above, we fused the N-terminus of LAMP1 to a fluorophore (Emerald

or mCherry) to specifically target the fluorescent signal to the lysosome (**Fig. 1C, 3A**). To validate

the GEM-SCOPe LAMP1-mCherry signal responds to lysosomal stress, hiPSC-derived

astrocytes transduced with LAMP1-mCherry lentivirus were exposed to bafilomycinA1 (BafA1) for

15 hours. BafA1 inhibits lysosomal proton pump V-ATPases, which blocks autophagosome-

lysosome fusion and lysosomal acidification, impairing lysosomal function [70,71]. hiPSC-derived

astrocytes were simultaneously transduced with a non-localized Emerald fluorophore lentivirus,

to provide a cell body marker. We used CellProfiler [72] to evaluate the lysosomal number, size, and

distribution within each cell. Compared to vehicle (DMSO) treatment, treatment with BafA1

increased the number of LAMP1-mCherry vesicles (unpaired t-test, p = 0.04; **Fig. 3B, C**), but did

not affect lysosome size (unpaired t-test, p = 0.16, **Fig. 3B, C**). Treatment with BafA1 altered the

distribution of lysosomes within the astrocytes. Astrocytes treated with BafA1 had a more

dispersed distribution of LAMP1-mCherry signal, with only 41±4% concentrated in the perinuclear

area, in contrast to 65±8% in vehicle-treated astrocytes (unpaired t-test, p = 0.002; **Fig. 3B, C**).

Perinuclear and juxtanuclear lysosomes have been posited to be more acidic than peripheral

lysosomes[73]. Lysosomal distribution can also be affected by cytosolic pH, with acidification

promoting the spreading of lysosomes away from the nucleus[74-76]. Since BafA1 inhibits V-

ATPases responsible for establishing and maintaining lysosomal pH, the redistribution of

lysosomes away from the nucleus upon BafA1 treatment might reflect a change in lysosomal or

cytosolic pH, and therefore lysosomal function, in the treated astrocytes.

In addition to its role in ubiquitinating damaged mitochondria for degradation via the autophagy-

lysosomal pathway[31-33], PRKN has been shown to regulate endosome organization[77] and

mitochondria-lysosome contact sites. Although it has been reported that *PRKN* knockout in

hiPSC-derived dopaminergic neurons leads to an increase in lysosome number and size

compared to isogenic, wild-type dopaminergic neurons[78], the effect of *PRKN* knockout on

lysosomes on human astrocytes is largely unknown. Due to PRKN's role in regulating the

autophagy-lysosomal pathway, we hypothesized *PRKN*$^{-/-}$ in human astrocytes would disrupt

lysosomal dynamics. To test this hypothesis, we applied the same CellProfiler analysis on

PRKN$^{+/+}$ and *PRKN*$^{-/-}$ hiPSC-derived astrocytes transduced with GEM-SCOPe Emerald and

LAMP1-mCherry lentivirus. *PRKN*$^{-/-}$ astrocytes had half as many LAMP1-mCherry vesicles per

cell compared to *PRKN*$^{+/+}$ astrocytes (unpaired t-test, p = 0.003, **Fig. 3D, E**). Additionally, the

PRKN$^{-/-}$ lysosomes were smaller than the *PRKN*$^{+/+}$ lysosomes (unpaired t-test, p = 0.055, **Fig. 3D,**

E). Finally, the *PRKN*$^{-/-}$ astrocytes had a decrease in perinuclear LAMP1-mCherry, with only

38±4% of LAMP1-mCherry signal located in the perinuclear area, compared to 45±3% in

PRKN$^{+/+}$ (unpaired t-test, p = 0.005, **Fig. 3D, E**). Our results utilizing our GEM-SCOPe LAMP1-

mCherry fluorophore suggest that lysosomal distribution in astrocytes is altered upon loss of

PRKN and further studies to examine lysosomal content and activity are needed to understand

the implications and how this could differ from observations in other cell types.

287   Mitochondria-localized fluorophores capture mitochondrial

288   fragmentation and alterations in mitochondrial trafficking

289   along PRKN knockout dopaminergic neuron axons

Much like lysosomal dysfunction, mitochondrial dysfunction is implicated in PD and

neurodegeneration. Therefore, monitoring mitochondrial network dynamics in response to stimuli

and stress is important in understanding neurodegenerative diseases. As described above, we

fused the N-terminus of the mitochondrial protein COX8A to a fluorophore (Emerald or mCherry)

to localize the fluorophore specifically to the mitochondria (**Fig. 1C, 4A**). To validate that GEM-

SCOPe COX8A localized fluorophores reflect changes to mitochondrial morphology, we induced

mitochondrial stress by treating hiPSC-derived astrocytes transduced with the COX8A-mEmerald

lentivirus with 2µM oligomycin for 4 hours. Mitochondrial stress induces fragmentation of the

mitochondrial network, which facilitates mitophagy, but also decreases mitochondrial network

respiratory capacity[79,80]. Oligomycin, an inhibitor of ATP synthase, is a known inducer of

mitochondrial fragmentation. Treatment with oligomycin induced mitochondrial fragmentation

when compared to vehicle-treated astrocytes. The average area of mitochondria was reduced

from 1.04 μm^2 in vehicle-treated astrocytes to 0.8 μm^2 in oligomycin-treated astrocytes (1-way

ANOVA with Dunett's Test, p-adj = 0.0002; **Fig. 4B, C left)**, and average branch length was

reduced from 1.28 µm in vehicle-treated astrocytes to 0.9 µm in oligomycin treated astrocytes (1-

way ANOVA with Dunett's Test, p-adj < 0.0001; **Fig. 4B, C middle)**, as expected from a more

fragmented mitochondrial network. The aspect ratio is the ratio of the mitochondrial length to

width. An aspect ratio of 1 indicates a perfect circle, while larger aspect ratios indicate longer and

narrower shapes. The mean aspect ratio decreased from 2.52 in vehicle-treated astrocytes to

2.14 in oligomycin-treated astrocytes (1-way ANOVA with Dunett's Test, p-adj < 0.0001; **Fig. 4B,**

C right), another indicator that mitochondria are fragmenting upon oligomycin treatment. Metrics

of mitochondrial network fragmentation (mitochondrial area, branch length, and aspect ratio) are

all affected in a dose-dependent manner, with increasing concentrations of oligomycin having

increasingly more severe effects on mitochondrial network fragmentation (**Fig. S4**).

The transport of mitochondria along neuronal axons is critical for proper synaptic function and

neurotransmitter release. We hypothesized $PRKN^{-/-}$ may impair axonal mitochondrial transport.

To investigate this hypothesis, we transduced wild-type ($PRKN^{+/+}$) and $PRKN$ knock-out ($PRKN^{-/-}$

) hiPSC-derived midbrain organoids with COX8A-mEmerald. To facilitate imaging of axonal

mitochondrial transport, we generated radial axonal arbors by splatting organoids onto the surface

of a plate and allowing the axons to grow out[81]. On day 164 of organoid differentiation, fluorescent

mitochondria were imaged at high magnification over several minutes to monitor the movement

of mitochondria along these axons (**Fig. 4D; Video S1**). In the anterograde direction (moving

away from the soma), $PRKN^{-/-}$ mitochondria moved only 20% of the distance covered by $PRKN^{+/+}$

mitochondria (Welch's t-test, p < 0.0001, **Fig. 4E left**) and took 50% longer (unpaired t-test, p =

0.005; **Fig. 4E right**). Retrograde-moving mitochondria moved the same distance (unpaired t-

test, p = 0.59, **Fig. 4F left)** but the $PRKN^{-/-}$ mitochondria moved for 150% longer compared to the

$PRKN^{+/+}$ mitochondria (Welch's t-test, p = 0.02; **Fig. 4F right**). PRKN has previously been

implicated in mitochondrial transport due to its ubiquitination of mitochondrial motor adaptor

complexes[82]. Our results demonstrate that PRKN loss of function impairs mitochondrial trafficking

in human neurons, likely reducing the delivery of functional mitochondria synaptic extremities and

ultimately compromising neuronal function.

Dynamic fluorophores measure deficiencies in mitochondrial

turnover and glutathione reduction potential in PRKN-

knockout astrocytes

We demonstrated the utility of single excitation and emission wavelengths to discern phenotypes

regarding cellular and subcellular dynamics. Numerous groups have developed fluorophores that

change emission wavelengths under certain conditions or circumstances. We introduced several

multi-emission fluorophores into GEM-SCOPe to enable quantification of organelle turnover,

acidification, and oxidation, in addition to the localization information described with the previous

fluorophores.

Fluorescent Timer is a mutated dsRed protein that irreversibly shifts its fluorescence emission

from green to red due to tyrosine oxidation and can be used as a molecular clock to track protein

turnover (**Fig. 5A**).[22] We, therefore, localized GEM-SCOPe-Timer to the mitochondria (COX8A-

Timer), to further investigate mitochondrial turnover[83,84] (**Fig. 5B**). To first validate COX8A-Timer,

mitophagy was inhibited by treating hiPSC-derived astrocytes transduced with COX8A-Timer with

the V-ATPase inhibitor bafilomycinA1 (BafA1). BafA1 treatment led to a significant increase in the

ratio of red-fluorescent mitochondria over green-fluorescent mitochondria compared to vehicle

control astrocytes (unpaired t-test, $p < 0.0001$; **Fig. 5C, D**). Since mitochondrial degradation and

turnover occur primarily through mitophagy, inhibition of mitophagy and mitochondrial recycling

via BafA1 treatment results in a predicted increase in longer-lived mitochondria.

Because PRKN is directly involved in mitophagy, and we observed impaired retrograde

mitochondrial transport in $PRKN^{-/-}$ neurons, we wanted to measure changes in mitochondrial

turnover in human $PRKN^{-/-}$ cells. We transduced wild-type ($PRKN^{+/+}$) and $PRKN$ knockout ($PRKN^{-/-}$

) hiPSC-derived astrocytes with GEM-SCOPe COX8A-Timer. The ratio of red-fluorescent signal

over green-fluorescent signal significantly increased in $PRKN^{-/-}$ astrocytes compared to $PRKN^{+/+}$

astrocytes (unpaired t-test, $p = 0.026$, **Fig. 5E, F**). When broken down by individual channel, there

was no change in the green fluorescence (2-way ANOVA with Sidak correction, p-adj = 0.1288,

Fig. 5G), indicating no change in mitochondrial biogenesis. The increase in red to green ratio was

driven by a significant increase in red-fluorescent signal (2-way ANOVA with Sidak correction, p-

adj = 0.0005, **Fig. 5G**), which is in concordance with an accumulation of older mitochondria due

to a failure to properly degrade mitochondria in the absence of PRKN.

An expected consequence of impaired mitochondrial recycling is an accumulation of damaged

mitochondria and an increase in mitochondria-associated reactive oxygen species (ROS) from

inefficient cellular respiration. To measure reactive oxygen levels in live astrocytes, we introduced

GRX1-roGFP2, an established fluorophore for measuring ROS, into GEM-SCOPe[25]. GRX1-

roGFP2 is a fusion protein between GRX1 and roGFP2. When GRX1 is oxidized by glutathione,

it can subsequently oxidize roGFP2. Oxidized roGFP2 shifts its emission spectrum from green

fluorescence to blue fluorescence (**Fig. 6A**)[25,85]. Thus, the ratio of green to blue fluorescence is

an indicator of glutathione oxidation state. When GRX1-roGFP2 is further coupled with a

mitochondrial localization sequence, we can measure the oxidative stress of the mitochondria.

This construct was available as a retroviral vector using the signal sequence from *Neurospora*

crassa ATP synthase protein 9[25]. When we cloned it into the FUW lentivirus backbone, we decided

to keep that localization sequence rather than replace it with the signal sequence for COX8A used

in the other GEM-SCOPe mitochondria-localized fluorophores (**Fig. 6B**). To validate GEM-SCOPe

GRX1-roGFP2 was working as expected, hiPSC-derived astrocytes transduced with mito-GRX1-

roGFP2 were treated with BafA1 to induce oxidative stress[86]. Astrocytes treated with BafA1 had

a significant decrease in the ratio of green fluorescence to blue fluorescence when compared to

vehicle (DMSO) treated astrocytes (unpaired t-test, $p < 0.0001$, **Fig. 6C, D**). The effect is driven

by a significant increase specifically in blue fluorescence (2-way ANOVA with Sidak correction, p-

adj < 0.0001, **Fig. 6E**) while the green fluorescence remained unchanged (2-way ANOVA with

Sidak correction, p-adj $= 0.99$, **Fig. 6E**). The increase in blue fluorescence is due to an increase

in roGFP2 oxidation, which is indicative of an increase in glutathione oxidation and the presence

of reactive oxygen species. Thus, GEM-SCOPe-mito-GRX1-roGFP can be used to monitor

glutathione reduction potential and how cells respond to reactive oxygen stress.

Because astrocytes play important metabolically supportive roles in the CNS and we observed

PRKN-related impairments to mitochondrial recycling with COX8A-Timer (**Fig. 5E-G**), we wanted

to understand if PRKN deficiency has effects on ROS production in astrocytes. We transduced

PRKN[+/+] and *PRKN*[-/-] astrocytes with mito-GRX1-roGFP2 lentivirus. Without introducing any

additional chemical mitochondrial stressors, we observed an siginificant reduction in the ratio of

green-fluorescence signal to blue-fluorescence signal in *PRKN*[-/-] compared to *PRKN*[+/+] astrocytes

(unpaired t-test, $p = 0.026$; **Fig. 6F, G**). This is indicative of an increase in ROS in *PRKN*[-/-]

astrocytes which correlates with our other observation of decreased mitochondrial turnover in the

same cells. It has been previously demonstrated that *PRKN*[-/-] hiPSC-derived dopaminergic

neurons show increased cellular ROS[15] and *PRKN*$^{-/-}$ mice have decreased antioxidant capabilities

and increased ROS-related protein and lipid damage[87]. By introducing fluorophore-biosensors

into our lentiviral catalog, we were able to expand the phenotypes we can observe in live cells to

measure changes in the intracellular environment in response to chemical and genetic

perturbations.

Discussion

Here, we developed and validated GEM-SCOPe (Genetically Engineered, Modular, SubCellular

Organelle Probes), a collection of subcellularly targeted fluorophores encoded in a lentiviral

backbone. By exploiting the modular design of the constructs, we easily cloned fluorophores with

different excitation/emission spectra specified to localize to the nucleus, lysosomes, or

mitochondria. We demonstrated that these fluorophores specifically localize to the desired

organelle and can quantitatively assess cellular responses to chemical perturbations. Finally, we

applied GEM-SCOPe to human *PRKN*-knockout astrocytes and neurons, which revealed

widespread perturbations to cellular proliferation, lysosomal distribution, and mitochondrial

dynamics, providing new insights into PD pathogenesis.

Gliosis has been widely reported in human post-mortem brain studies of patients with *PRKN* loss-

of-function mutations[88-91]. Similar pro-inflammatory phenotypes have also been observed in

mouse models and stem-cell-derived glial models of PRKN deficiency[92,93]. However, there is still

a limited understanding of how genetic mutations related to PD modify astrocytes and their role

in neurodegeneration. We used GEM-SCOPe to investigate cellular, mitochondrial, and lysosomal

dynamics in astrocytes lacking functional PRKN protein. Using nucleus-localized fluorophores,

we show that *PRKN* knockout human astrocytes have increased cellular proliferation compared

to isogenic, wild-type astrocytes. LAMP1-localized fluorophores revealed that lysosomes in *PRKN*

knockout astrocytes are distributed further from the perinuclear nuclear area. Finally, a variety of

mitochondria-localized fluorophores revealed that *PRKN* knockout astrocytes have impaired

mitochondrial transport, decreased mitochondrial turnover, increased reactive oxygen species,

and oxidized glutathione. While additional experiments should be done to support these findings,

GEM-SCOPe highlights that there is extensive dysregulation and disruption to normal organelle

dynamics in human *PRKN* astrocytes that warrants further investigation.

GEM-SCOPe was developed in a 3rd generation lentiviral system. We selected this delivery

system because 1) lentiviruses are safe to produce; 2) they infect both dividing and non-dividing

cells; and 3) they can be used to develop stable cell lines. 3rd generation lentiviral systems are

replication-incompetent, so cells that are transduced with lentivirus cannot produce more viral

particles. Only the transfer plasmid, which contains the insert of interest, can be integrated into

the host genome; the other plasmids that encode lentiviral structural and envelope genes do not

integrate into the host genome, preventing viral replication. Most lentiviruses are produced with

VSV-G (Vesicular stomatitis virus G glycoprotein) as the envelope protein, as it provides broad

tropism and enables high transduction efficiency in a wide variety of cell types[94]. Finally, GEM-

SCOPe lentiviruses were developed with the option to include an antibiotic-resistance gene (**Fig.**

1A), which can be used to establish stable cell lines that have incorporated the lentiviral load.

Although lentiviruses provide a lot of flexibility and safety when delivering genetic information,

they also possess some limitations. Lentiviruses have an ideal plasmid size of 9-10kb, and viral

titer can decrease if plasmids exceed this size[95]. The H2B-mTagBFP2, H2B-mCherry, and H2B-

Emerald lentiviruses (**Fig. 1, 2**) are about 11kb each and transduced hiPSC-derived astrocytes

with about 90% efficiency. Coupled with the puromycin resistance that was also included in this

construct, we were still able to generate a population with 100% viral transduction despite the size

limitation. Another drawback of lentiviral gene delivery is that inserts often integrate into actively

transcribed loci[96]. Differentiation of hiPSCs into specific cell types requires extensive chromatin

remodeling, and genomic sequences integrated via lentivirus can often be silenced in this

process. In the case where hiPSCs must be transduced before differentiation, antibiotic selection

during differentiation can select for cells that have integrated the viral load into a genomic region

that is not silenced. The final drawback of lentiviral gene delivery is that fluorescence intensity is

dependent on the viral copy number in each cell and cannot be used as a readout for organelle

abundance. Rather, we need to use measurements like distribution, size, and ratios of different

emissions within the same cell. This is not just a limitation of lentiviral gene delivery but of other

viral and non-viral delivery systems.

In this study, we focused on PD and specifically the alterations to astrocytes with *PRKN* mutations.

However, alterations in proliferation, mitochondria, and lysosomes are by no means unique to PD.

Many reviews cover the extensive roles of mitochondria and lysosomes in health and disease[97-

100]. GEM-SCOPe can be applied beyond neuronal cells and neurodegeneration to countless other

cell types to investigate subcellular dynamics that affect disease development and progression.

Due to its modular design, GEM-SCOPe can be expanded and molded by research groups to

meet specific needs. Different targeting sequences can be integrated to localize fluorophores to

relevant subcellular compartments. New fluorophores and biosensors are always being

developed to measure additional intracellular conditions. By applying GEM-SCOPe to disease

models, we will be able to better understand how cells are disrupted over the course of disease

progression, providing insight for future therapeutic targets.

Acknowledgements

The authors thank Deanna Benson for discussions on mitochondrial transport and kymograph analysis. We thank the Microscopy CoRE at Icahn School of Medicine for providing access to the Leica DMi8 and Thermo Scientific CX7 High Content Screening Platform, as well as training and technical expertise for using these microscopes.

This work has been funded in part with federal funds from NASA under contract #80ARC022CA004 titled "Identification of Biomarkers and Pathological Mechanisms via Longitudinal Analysis of Neurological and Cerebrovascular Responses to Neurotoxic Stress Using a Multi-cellular Integrated Model of the Human Brain." This work was also supported by funding from Aligning Science Across Parkinsons (02663614), the NIH/NINDS (R010255E131 and UG3NS115064), and the CureAlz Fund. C.G. was supported by funding from the NIH/NIA (T32AG04968) and NIH/NINDS (F31NS13090), and L.S. was supported by the Training Program in Stem Cell Biology fellowship from the New York State Department of Health (NYSTEM-C32561GG).

Author Contributions

Conceptualization, C.G., T.A., and J.W.B.; Methodology, C.G., L.S., and T.A.; Validation, C.G.; Formal Analysis, C.G.; Investigation, C.G., T.K., L.S., and A.S.; Resources, C.G., T.K., L.S., and B.R.S.; Writing – Original Draft, C.G. and J.W.B.; Writing – Review & Editing, C.G., L.S., B.R.S., and J.W.B.; Visualization, C.G.; Funding Acquisition, T.A. and J.W.B.

Declaration of Interests

The authors declare no competing interests

Materials and Methods

Experimental model and study participant details

The human cell lines used in this study are detailed in the key resources table and below.

HEK293FT (RRID: CVCL_6911, fetal kidney origin, female sex) were maintained in DMEM (Gibco 10566) supplemented with 10% bovine calf serum (Cytiva SH30073) at 37 °C in a 5% CO_2 incubator. Cells were passaged ever 4-5 days, when they reached 90% confluency.

H1(WA01) hESCs (male), BJ-SiPS-D iPSCs (male), and AG09173 iPSCs (female) were maintained in StemFlex medium (Gibco A3349401) at 37 °C in a 5% CO_2 incubator. Cells were passaged every 3-4 days, when they reached 90% confluency. WA01 astrocytes were used for all PRKN-genotype dependent experiments. BJ-SiPS-D and AG09173 were used for organelle localization validation and chemical perturbation validation.

Molecular cloning

All vectors were derived from pFUW (Addgene plasmid #14882). pFUW was linearized using NheI-HF (NEB R3131) and BamHI-HF (NEB R3136) per manufacturer's protocol, and the desired fragment was gel-purified (Macherey-Nagel 740609). Components of the insert were PCR amplified using Q5-Hot Start DNA Polymerase (NEB M0493) per manufacturer's recommendations. Fluorophores were amplified from commercially available plasmids (see

Resource Table for complete list) and localization sequences were amplified from cDNA. Primers

were designed using the NEBuilder Assembly Tool (nebuilder.neb.com) to add 5' and 3'

sequences that make the fragments compatible for Gibson assembly and ordered from IDT

(Integrated DNA Technologies). Gibson reactions were set up with NEBuilder HiFi DNA Assembly

Master Mix (NEB E2621), 50ng of linearized backbone, and 3-fold molar excess of each PCR

fragment. For fragments less than 200bp, 5-fold molar excess was used. Reactions were

incubated at 50°C for 60 minutes. The Gibson product was transformed in 10-beta competent *E.*

coli (NEB C3019) grown on LB Agar plates with 100ug/mL ampicillin. Plasmids from single clones

were isolated via miniprep (Macherey-Nagel 740499) and sequenced to confirm successful

cloning.

Lentivirus Production

Lentivirus was produced by transfection of HEK293T cells as adapted from Dull et al. 1998 [101].

HEK293T cells were grown in Dulbecco's Modified Eagle Medium (DMEM; Gibco 11965092) with

10% bovine calf serum (Cytiva SH30073) at 37 °C in a 5% CO_2 incubator. The day before

transfection, ~10^7 cells were plated per 15cm dish coated in 0.1% gelatin. Media was changed

just prior to transfection. For one 15cm dish, 22.5ug of transfer vector was combined with 14.7ug

of pMDLg/pRRE (Addgene plasmid #12251), 5.7ug of pRSV.Rev (Addgene plasmid #12253), and

7.9ug of pMD2.G (Addgene plasmid #12259) in a 15mL conical tube. The plasmid mixture was

added to 1mL of 278 uM $CaCl_2$ (Sigma C1016) and mixed thoroughly before adding 1mL of 2x

BBS solution [280 mM NaCl (Fisher Scientific S271-1), 50 mM BES (Millipore 391334), 1.5mM

Na_2HPO_4 (Sigma S5136); pH = 6.95] dropwise. Tubes were mixed by inversion, incubated at room

temperature for 1 min, and then added dropwise to HEK293T dish. Cell culture media was

changed after 16-24 h. Conditioned media was then collected every 24 h for 2 days; conditioned

media from the same plate was combined over the two collections. Conditioned media was

centrifuged at 2000 x g for 10 minutes to pellet debris and filtered through 0.22-um-pore-size PES

filter (Millipore SCGP00525). To concentrate virus, filtered media was incubated with PEG 8000

(Sigma P2139) and NaCl, at a final concentration of 5% and 0.15M respectively, overnight at 4°C,

then centrifuged at 3000 x g for 20 min. The pellet was resuspended in DMEM/F12 with GlutaMAX

(Gibco 10565018) to 1% of the original supernatant volume. The concentrated virus was stored

in aliquots at -80°C.

535 Midbrain Organoid and Astrocyte Differentiation

hiPSC-derived astrocytes were extracted from 100-day-old midbrain organoids as described in

Sarrafha et al 2021 and Parfitt et al 2024 [81,102]. In short, hiPSCs were cultured in StemFlex medium

(Gibco A3349401) at 37 °C in a 5% CO_2 incubator. hiPSCs were seeded into 125-mL spinner

flasks (Corning 3152) and allowed to self-aggregate in StemFlex (Gibco A3349401)

supplemented with 10 µM Y-27632 (Tocris 1254) and 1% Penicillin-Streptomycin (Gibco

15140122). Once aggregates were between 300 µm and 500 µm, differentiation was initiated by

dual-SMAD inhibition with 10 µM SB431542 (Stemgent 04-0010), 100nM LDN193189 (Tocris

6053), 1x B27 without Vitamin A (Gibco 12587010) and 1x N2 (Gibco 17502048) in DMEM-F12

with GlutaMAX (Gibco 10565018). Midbrain patterning was achieved with the addition of 3 µM

CHIR99021 (Tocris 4423), 2 µM purmorphamine (Stemcell 72204), and 1 µM SAG (Cayman

11914) starting 4 days after neural induction. After patterning, media was changed to DMEM/F12

with GlutaMAX, supplemented with 1x N2, 1x B27 without Vitamin A, 20 ng/mL GDNF (Peprotech

450-10), 20 ng/mL BDNF (Peprotech 450-02), 200 µM L-ascorbic acid (Fisher Scientific BP351),

100 µM dibutyryl cAMP (Biogems 1698950), and 10 µM DAPT (Cayman 13197). After 35 days,

organoids were transferred to ultra-low attachment plates (Corning 3516), with about 5 organoids

per mL of media and cultured in DMEM/F12 with GlutaMAX supplemented with 1x N2, 1x B27

without Vitamin A, 10 ng/mL GDNF, 10 ng/mL BDNF, and 200 µM ascorbic acid.

Astrocytes were extracted from midbrain organoids starting at day 100. Organoids were gently

dissociated in a trypsin enzyme solution (TrypLE Select, Gibco 12563011) using a glass pipette

to break up organoids into large chunks. The organoid chunks from 5-10 organoids were placed

on 15cm dishes coated with 0.1% gelatin and maintained in Astrocyte Medium (AM; ScienCell

#1801). Astrocytes were allowed to grow out of the organoid chunks until the plate was confluent

(about 1 week). At this point, astrocytes can be cryopreserved. Astrocytes were maintained in AM

and switched to experimental media [1:1 DMEM/F-12:Neurobasal (Gibco 21103049), 1x B27

without Vitamin A, 1x N2, 1x MEM-Non-Essential Amino Acids (Gibco 11140050), 1x GlutaMAX

(Gibco 35050061), 10 ng/mL CNTF (Peprotech 450-13)] 2-3 days before imaging.

Astrocyte Viral Transduction

iPSC-derived astrocytes were transduced during regular passaging. In short, astrocytes were

lifted, spun down, resuspended in AM media, and divided into 1.5mL microcentrifuge tubes,

depending on the desired split ratio and number of transductions. The virus was added to the cell

suspension and incubated for 5-10 minutes before plating the mixture. Media was topped off such

that the final dilution of the virus in media was 1:50. Depending on the viral titer, this dilution

sometimes increased or decreased. Astrocytes can also be transduced without passage by

adding the virus directly to the media. Media is changed after 24 hours. Antibiotics were added to

the media, if relevant, starting at least 3 days after transduction.

MitoTracker and LysoTracker

For MitoTracker staining, astrocytes were incubated with 100 nM of MitoTracker DeepRed (Thermo Scientific M22425) for 30 min at 37 °C. For LysoTracker staining, astrocytes were incubated 100 nM of LysoTracker DeepRed (Thermo Scientific L12492) for 5 min at 37 °C. Cells were then washed twice with PBS before incubation with Hoechst 33342 (Thermo Scientific 62249).

qPCR

RNA was extracted from samples using TRIzol (Invitrogen 15596018) and an RNA extraction kit following the manufacturer's recommended protocol (Zymo R2062). RNA concentration and purity were measured on a spectrophotometer. cDNA was produced from 1 ug of RNA per sample, using Maxima H Minus Reverse Transcriptase (Thermo Scientific EP0752) and following the manufacturer's recommendations. qPCR reactions were prepared in a 384-well plate (Applied Biosystems 4309849) with 1:10 dilution of cDNA, 2x PowerUp SYBR Green Master Mix (Applied Biosystems A25741) and 1 µM each of forward and reverse primers. qPCRs were run on QuantStudio 7 (Applied Biosystems 4485701) and analyzed using the ΔΔCt method, normalized to *ACTB*. The following primers were used:

ACTB Forward: CCTGGCACCCAGCACAAT

ACTB Reverse: GCCGATCCACACGGAGTA

LAMP1 Forward: CAGATGTGTTAGTGGCACCCA

LAMP1 Reverse: TTGGAAAGGTACGCCTGGATG

Live Cell Imaging

In general, astrocytes were seeded at 10 x 10^3 cells per well of a 96-well plate with a cover glass thickness polystyrene bottom (Greiner 655090) coated in 0.1% gelatin 3-5 days before imaging. Proliferation assays were seeded at 8 x 10^3 cells per well and started 1 day after seeding. Before imaging, cells were incubated with Hoechst 33342 (Thermo Scientific 62249) or DRAQ5 (Thermo Scientific 52251) for 10 minutes.

Wide-field live cell imaging was performed using the Thermo Scientific CX7 High Content Screening Platform with 20x and 40x objective lenses (LUCPLFLN20x and LUCPLFLN40x). The microscope was equipped with an incubation unit that maintained temperature, CO_2, and humidity during image acquisition. Confocal images were acquired on a Nikon Ti2E AX R confocal microscope with a 20x objective lens (PLAN APO λD 20x) and 6x optical zoom. Each well was imaged in 5 (confocal) or 25-36 (widefield) different areas, with the average of those images represented as a single data point.

ImageJ Image Analysis

Nuclear Proliferation: Nuclear proliferation was quantified by converting images to a binary based on a threshold and then using 'Analyze Particles' to obtain a count for the number of nuclei in a field. The same threshold and parameters were kept for all time points.

Mitochondrial Fragmentation: Mitochondrial fragmentation was measured using the ImageJ plugin, Mitochondrial Analyzer [103]. For thresholding, the "Block Size" was set to 1.45 microns and the "C-Value" was set to 5.

Ratiometric Quantification: Ratiometric quantifications for roGFP2, Timer, and Lemon were

achieved by measuring the mean grey value in both channels of interest. The values for each

channel were then divided to generate the ratio of relative intensity between the two channels.

Kymograph Analysis: Kymograph analysis of mitochondrial movement was achieved using the

Image J plugins KymographClear 2.0 and KymographDirect [104]. KymographClear 2.0 was used

to trace mitochondrial trajectories and generate kymographs from which stationary and motile

mitochondria were identified. KymographDirect was then used to extract quantitative information

from the trajectories of the motile mitochondria.

Cell Profiler Image Analysis [72,105]

Subcellular Co-localization: For nuclear co-localization, nuclei in the genetically encoded

fluorophore channel and the DRAQ5 channel were identified, and nuclei from cells that were not

transduced were filtered out. For lysosomal and mitochondrial co-localization, propagation from

the nuclear stain was used to identify cell bodies. Cells were filtered to only analyze those with

lentivirus signal within the cell body. Co-localization was reported as the Mander's coefficient

indicating the overlap of the genetically encoded fluorophore with the commercially available

stain.

Lysosomal Distribution: Cells were defined by nuclei and full cell fluorophore signal. Lamp1-

mCherry signal was assigned to the cell body with which it overlapped. Only cells that were

successfully transduced were included in the analysis. The distribution of the Lamp1-mCherry

signal was calculated using the "MeasureObjectIntensityDistribution" with two bins generated for

each cell using the nuclei as the center of the cell. Bins were scaled such that the inner bin

(perinuclear) was always the same percent of total cell area regardless of cell size.

Quantification and statistical analysis

All experiments were conducted with 3-6 replicates. All quantifications were represented as mean ± standard deviation, except for the violin plots of the kymograph analysis. Details on data visualization, sample size, and images per replicate are included in figure captions. Details on statistical tests and significance are included in the text. All statistical tests were conducted in *GraphPad Prism* and graphs were generated in *GraphPad Prism*.

Figure Legends

Fig. 1: A toolbox of lentiviral plasmids developed and validated for subcellular localization of fluorophores (A) Schematic of lentiviral plasmids highlighting the combinatorial power of the toolbox components. Generated using BioRender **(B)** Schematic demonstrating the cloning strategies used to generate the constructs used in this study as well as any future constructs. Generated using BioRender **(C)** Representative images of subcellular localization validation with available dyes in hiPSC-derived astrocytes and Mander's co-efficient for overlap of the genetically encoded fluorophore with the dye. Bars represent mean values and error bars represent standard deviation. Top: H2B-mTagBFP2 and DRAQ5 (n = 3; each replicate is an average over 16 images); Middle: LAMP1-mCherry and LysoTracker Deep Red; nuclei stained with Hoechst33342 (n = 4; each replicate is an average over 4 images); Bottom: COX8A-Emerald and MitoTracker Deep Red; nuclei stained with Hoechst33342 (n = 3, each replicate is an average over 36 images). Scale bars = 50 μm **(D)** Representative images of combinatorial applications of GEM-SCOPe.

Top: Cell expressing LAMP1-Emerald and COX8A-mCherry. Bottom: cell expression LAMP1-

mCherry and COX8A-Emerald. Nuclei were stained with DRAQ5. Scale bars = 50 μm.

Fig. 2: Nucleus-localized fluorophores offer improvements over available stains for

multiday imaging (A) Schematic of the lentiviral construct localizing mTagBFP2 to the nucleus

with an H2B fusion protein and antibiotic resistance to puromycin used in the following panels **(B)**

Representative images of H2B-mTagBFP2 and DRAQ5 nuclear stain 1 hour (top) and 24 hours

(bottom) after adding DRAQ5 to the cell cultures. **(C)** Quantification of the percent of stain

localized to the nuclear area versus the cytoplasm 1 hour and 24 hours after adding DRAQ5 to

cell cultures. The central bar represents the mean and the error bars represent the standard

deviation (n = 3 per time point; each replicate is an average over 36 images) **(D)** Schematic of the

lentiviral construct localizing Emerald or mCherry to the nucleus with an H2B fusion protein and

antibiotic resistance to puromycin used in the following panels **(E)** Representative images of

nuclei from $PRKN^{+/+}$ astrocytes, labeled with H2B-Emerald, and $PRKN^{-/-}$ astrocytes, labeled with

H2B-mCherry, cultured together during a proliferation assay. Images were taken 24 hours (top)

after seeding and 72 hours (bottom) after seeding **(F)** Quantification of number of $PRKN^{+/+}$ and

$PRKN^{-/-}$ nuclei per image field 24 hours and 72 hours after seeding. Dots represent mean values

and error bars represent standard deviation (n = 3 per time point; each replicate is an average

over 25 images). Scale bars = 50 μm. All images were acquired on the CX7 HCS platform with a

20x objective lens.

Fig. 3: Lysosome-localized fluorophores reveal changes in lysosomal distribution upon

chemical and genetic perturbation (A) Schematic of the lentiviral constructs used in the

following panels localizing (top) mCherry to the lysosome with the 5' sequence of *LAMP1* and

(bottom) Emerald to the whole cell by not including a localization sequence. **(B)** Representative

images of astrocytes transduced with Emerald and LAMP1-mCherry lentiviruses and treated with

100 nM bafilomycin A1 (bottom) or a vehicle (DMSO; top). Nuclei were stained with

Hoechst33342. Far right: Output from CellProfiler showing the distribution Lamp1-mCherry signal

as a fraction of signal in each bin. Darker blue indicates a higher fraction of signal in that bin.

Scale bars = 50 µm. **(C)** Quantification of lysosomal number per cell (left), average vesicle area

(middle) and fraction of perinuclear lysosomes (right) using Lamp1-mCherry signal in vehicle

(DMSO) and bafilomycin A1 (100 nM) treated astrocytes. Emerald was used to determine cell

boundaries. Central bars represent mean and error bars represent standard deviation (n = 4; each

replicate is an average over 10 images). **(D)** Representative images of $PRKN^{+/+}$ (top) and $PRKN^{-}$

$^{/-}$ (bottom) astrocytes transduced with cytoplasmic Emerald and LAMP1-mCherry lentiviruses.

Nuclei were stained with Hoechst33342. Far right: Output from CellProfiler showing the

distribution Lamp1-mCherry signal as a fraction of signal in each bin. Darker blue indicates a

higher fraction of signal in that bin. Scale bars = 25 µm. (**E**) Quantification of Lamp1-mCherry

signal in $PRKN^{+/+}$ and $PRKN^{-/-}$ astrocytes to measure lysosomal number per cell (left), average

vesicle area (middle) and fraction of perinuclear lysosomes (right). Emerald fluorescence was

used to determine cell boundaries. Central bars represent mean and error bars represent

standard deviation (n = 6; each replicate is an average over 7 images). All images were acquired

on a Nikon Ti2E AX R confocal microscope with a 20x objective lens and 6x optical zoom.

Fig. 4: Mitochondria-localized fluorophores resolve changes in mitochondrial network

dynamics (A) Schematic of the lentiviral construct used in the following panels localizing Emerald

to the mitochondria with the targeting sequence of COX8A. **(B)** Representative images of

astrocytes transduced with COX8A-Emerald and treated with a vehicle (DMSO; top) or 2 uM

oligomycin (bottom). Nuclei were stained with Hoechst33342. High magnification images (far

right) highlight the changes to the mitochondrial network upon oligomycin treatment. Scale bars

= 50 µm. **(C)** Quantification of mean mitochondrial area, mean mitochondrial branch length, and

mean aspect ratio upon oligomycin treatment. Central bars represent mean and error bars

represent standard deviation (n = 3; each replicate is an average over 36 images) **(D)**

Representative images of mitochondria labeled with COX8A-Emerald moving along *PRKN⁻ᐟ⁻*

neuron axons in the anterograde (left) and retrograde (right) directions. Scale bars = 5 µm. Images

were taken on a Leica DMi8 microscope for 1 hour per field **(E-F)**. Quantification of mitochondrial

run length (left) and run time (right) of COX8A-Emerald labeled mitochondria in *PRKN⁺ᐟ⁺* and

PRKN⁻ᐟ⁻ dopaminergic neuron axons moving anterograde (E) and retrograde (F) as determined

by kymograph analysis (average of n = 37.5 organoids for anterograde and 23.5 organoids for

retrograde; each replicate is an average over 1-3 fields).

Fig. 5: Timer fluorophore measures changes in subcellular mitochondrial turnover in

***PRKN⁻ᐟ⁻*(A)** Schematic demonstrating how the fluorescent protein Timer works. Timer undergoes

a fluorescent shift from a green emission to a red emission overtime; analysis of Timer relies on

examining the ratio between the red and green emission. **(B)** Schematic of the lentiviral construct

used in the following panels localizing Timer to the mitochondria by using the targeting sequence

of COX8A. **(C)** Representative images of astrocytes transduced with COX8A-Timer and treated

with a vehicle (DMSO; top) or 100 nM bafilomycin A1 (bottom) for 24 hours. Images were acquired

with a 488-excitation line and a 594-excitation line. Far right: ratiometric representation of the red

channel divided by the green channel and pseudo colored so that orange, yellow and white

indicate more relative red emission and black and purple indicate more relative green emission.

Nuclei were stained with DRAQ5. Scale bars = 50 µm. **(D)** Quantification of the ratio of mean

intensity of emission from excitation with a 594nm laser to the mean intensity of emission from

excitation with a 488 nm laser in astrocytes expressing COX8A-Timer and treated with a vehicle

(DMSO) or 100 nM bafilomycin A1 for 24 hours. Central bars represent mean and error bars

represent standard deviation (n = 6; each replicate is an average over 25 images). (**E**)

Representative images of *PRKN⁺ᐟ⁺* and *PRKN⁻ᐟ⁻* astrocytes transduced with COX8A-Timer.

Images were acquired with a 488-excitation line and a 594-excitation line. Far right: ratiometric

representation of the red channel divided by the green channel and pseudo colored so that

orange, yellow and white indicate more relative red emission and black and purple indicate more

relative green emission. Nuclei were stained with DRAQ5. Scale bars = 50 μm. **(F)** Quantification

of the ratio of mean intensity of emission from excitation with a 594nm laser to the mean intensity

of emission from excitation with a 488 nm laser in *PRKN*$^{+/+}$ and *PRKN*$^{-/-}$ astrocytes expressing

COX8A-Timer. Central bars represent mean and error bars represent standard deviation (n = 3;

each replicate is an average over 25 images). **(G)** Quantification of mean intensity normalized by

nuclear area of green and red fluorescence in *PRKN*$^{+/+}$ and *PRKN*$^{-/-}$ astrocytes expressing

COX8A-Timer. Central bars represent mean and error bars represent standard deviation (n = 3;

each replicate is an average over 25 images). All images were acquired on a CX7 HCS platform

with a 20x objective lens.

Fig. 6: GRX1-roGFP2 fluorophore is used to quantify glutathione oxidation in chemical and

genetic models of oxidative stress (A) Schematic demonstrating how the fluorescent protein

fusion Grx1-roGFP2 works. Oxidation of Grx1 upon the oxidation of glutathione results in the

eventual oxidation of roGFP2, shifting its emission spectrum from green to blue. **(B)** Schematic

of the lentiviral construct used in the following panels localizing Grx1-roGFP2 to the mitochondria.

This fluorophore was localized using atp-9 instead of COX8A. **(C)** Representative images of

astrocytes transduced with mito-GRX1-roGFP2 and treated with a vehicle (DMSO; top) or 100

nM bafilomycin A1 (bottom) for 24 hours. Images were acquired with a 405-excitation laser and a

488-excitation laser. Far right: ratiometric representation of the green channel divided by the blue

channel and then pseudo colored so that orange, yellow and white indicate more relative green

emission while black and purple represent more blue emission. Nuclei were stained with DRAQ5.

Scale bars = 50 μm. **(D)** Quantification of the ratio of mean intensity from 488-excitation to the

mean intensity from 405 excitation. In astrocytes expression mito-Grx1-roGFP2 and treated with

a vehicle or 100 nM bafilomycin A1. Central bars represent mean and error bars represent

standard deviation (n = 6; each replicate is an average over 25 images). **(E)** Quantification of

mean intensity normalized by nuclear area of blue and green fluorescence in vehicle and bafilomycin A1 treated astrocytes expressing mito-Grx1-roGFP2. Central bars represent mean and error bars represent standard deviation (n = 6; each replicate is an average of 25 images). **(F)** Representative images of *PRKN*$^{+/+}$ and *PRKN*$^{-/-}$ astrocytes transduced with mito-Grx1-roGFP2. Images were acquired with a 405-excitation laser and a 588-excitation laser. Far right: ratiometric representation of the green channel divided by the blue channel and pseudo colored so that orange, yellow and white indicate more relative green emission and black and purple indicate more relative blue emission. Nuclei were stained with DRAQ5. Scale bars = 50 μm. **(G)** Quantification of the ratio of mean intensity of emission from excitation with a 488nm laser to the mean intensity of emission from excitation with a 405 nm laser in *PRKN*$^{+/+}$ and *PRKN*$^{-/-}$ astrocytes expressing mito-Grx1-roGFP2. Central bars represent mean and error bars represent standard deviation (n = 3; each replicate is an average over 16 images).

References

1.	Willis, A.W., Roberts, E., Beck, J.C., Fiske, B., Ross, W., Savica, R., Van Den Eeden, S.K., Tanner, C.M., Marras, C., and Parkinson's Foundation, P.G. (2022). Incidence of Parkinson disease in North America. NPJ Parkinsons Dis *8*, 170. 10.1038/s41531-022-00410-y.

2.	Vazquez-Velez, G.E., and Zoghbi, H.Y. (2021). Parkinson's Disease Genetics and Pathophysiology. Annu Rev Neurosci *44*, 87-108. 10.1146/annurev-neuro-100720-034518.

3.   Schneider, S.A., and Obeso, J.A. (2015). Clinical and pathological features of

Parkinson's disease. Curr Top Behav Neurosci *22*, 205-220. 10.1007/7854_2014_317.

4.   Varadi, C. (2020). Clinical Features of Parkinson's Disease: The Evolution of Critical

Symptoms. Biology (Basel) *9*. 10.3390/biology9050103.

5.   Jankovic, J. (2008). Parkinson's disease: clinical features and diagnosis. J Neurol

Neurosurg Psychiatry *79*, 368-376. 10.1136/jnnp.2007.131045.

6.   Naoi, M., and Maruyama, W. (1999). Cell death of dopamine neurons in aging and

Parkinson's disease. Mech Ageing Dev *111*, 175-188. 10.1016/s0047-6374(99)00064-0.

7.   Spillantini, M.G., Schmidt, M.L., Lee, V.M., Trojanowski, J.Q., Jakes, R., and Goedert, M.

(1997). Alpha-synuclein in Lewy bodies. Nature *388*, 839-840. 10.1038/42166.

8.   Ohama, E., and Ikuta, F. (1976). Parkinson's disease: distribution of Lewy bodies and

monoamine neuron system. Acta Neuropathol *34*, 311-319. 10.1007/BF00696560.

9.   Hartmann, A. (2004). Postmortem studies in Parkinson's disease. Dialogues Clin

Neurosci *6*, 281-293. 10.31887/DCNS.2004.6.3/ahartmann.

10.   Helmchen, F., Fee, M.S., Tank, D.W., and Denk, W. (2001). A miniature head-mounted

two-photon microscope. high-resolution brain imaging in freely moving animals. Neuron

31, 903-912. 10.1016/s0896-6273(01)00421-4.

11.   Zong, W., Wu, R., Li, M., Hu, Y., Li, Y., Li, J., Rong, H., Wu, H., Xu, Y., Lu, Y., et al.

(2017). Fast high-resolution miniature two-photon microscopy for brain imaging in freely

behaving mice. Nat Methods *14*, 713-719. 10.1038/nmeth.4305.

12.   Chen, K., Tian, Z., and Kong, L. (2022). Advances of optical miniscopes for in vivo

imaging of neural activity in freely moving animals. Front Neurosci *16*, 994079.

10.3389/fnins.2022.994079.

13.   Potashkin, J.A., Blume, S.R., and Runkle, N.K. (2010). Limitations of animal models of

Parkinson's disease. Parkinsons Dis *2011*, 658083. 10.4061/2011/658083.

14.   Chia, S.J., Tan, E.K., and Chao, Y.X. (2020). Historical Perspective: Models of

808   Parkinson's Disease. Int J Mol Sci *21*. 10.3390/ijms21072464.

809   15.   Ahfeldt, T., Ordureau, A., Bell, C., Sarrafha, L., Sun, C., Piccinotti, S., Grass, T., Parfitt,

810   G.M., Paulo, J.A., Yanagawa, F., et al. (2020). Pathogenic Pathways in Early-Onset

811   Autosomal Recessive Parkinson's Disease Discovered Using Isogenic Human

812   Dopaminergic Neurons. Stem Cell Reports *14*, 75-90. 10.1016/j.stemcr.2019.12.005.

813   16.   Raja, W.K., Neves, E., Burke, C., Jiang, X., Xu, P., Rhodes, K.J., Khurana, V.,

814   Scannevin, R.H., and Chung, C.Y. (2022). Patient-derived three-dimensional cortical

815   neurospheres to model Parkinson's disease. PLoS One *17*, e0277532.

816   10.1371/journal.pone.0277532.

817   17.   Chlebanowska, P., Tejchman, A., Sulkowski, M., Skrzypek, K., and Majka, M. (2020).

818   Use of 3D Organoids as a Model to Study Idiopathic Form of Parkinson's Disease. Int J

819   Mol Sci *21*. 10.3390/ijms21030694.

820   18.   Kim, H., Park, H.J., Choi, H., Chang, Y., Park, H., Shin, J., Kim, J., Lengner, C.J., Lee,

821   Y.K., and Kim, J. (2019). Modeling G2019S-LRRK2 Sporadic Parkinson's Disease in 3D

822   Midbrain Organoids. Stem Cell Reports *12*, 518-531. 10.1016/j.stemcr.2019.01.020.

823   19.   Shimomura, O., Johnson, F.H., and Saiga, Y. (1962). Extraction, purification and

824   properties of aequorin, a bioluminescent protein from the luminous hydromedusan,

825   Aequorea. J Cell Comp Physiol *59*, 223-239. 10.1002/jcp.1030590302.

826   20.   Wang, M., Da, Y., and Tian, Y. (2023). Fluorescent proteins and genetically encoded

827   biosensors. Chem Soc Rev *52*, 1189-1214. 10.1039/d2cs00419d.

828   21.   Ovechkina, V.S., Zakian, S.M., Medvedev, S.P., and Valetdinova, K.R. (2021).

829   Genetically Encoded Fluorescent Biosensors for Biomedical Applications. Biomedicines

830   *9*. 10.3390/biomedicines9111528.

22. Terskikh, A., Fradkov, A., Ermakova, G., Zaraisky, A., Tan, P., Kajava, A.V., Zhao, X.,

Lukyanov, S., Matz, M., Kim, S., et al. (2000). "Fluorescent timer": protein that changes

color with time. Science *290*, 1585-1588. 10.1126/science.290.5496.1585.

23. Nakai, J., Ohkura, M., and Imoto, K. (2001). A high signal-to-noise Ca(2+) probe

composed of a single green fluorescent protein. Nat Biotechnol *19*, 137-141.

10.1038/84397.

24. Burgstaller, S., Bischof, H., Gensch, T., Stryeck, S., Gottschalk, B., Ramadani-Muja, J.,

Eroglu, E., Rost, R., Balfanz, S., Baumann, A., et al. (2019). pH-Lemon, a Fluorescent

Protein-Based pH Reporter for Acidic Compartments. ACS Sens *4*, 883-891.

10.1021/acssensors.8b01599.

25. Gutscher, M., Pauleau, A.L., Marty, L., Brach, T., Wabnitz, G.H., Samstag, Y., Meyer,

A.J., and Dick, T.P. (2008). Real-time imaging of the intracellular glutathione redox

potential. Nat Methods *5*, 553-559. 10.1038/nmeth.1212.

26. Nalls, M.A., Blauwendraat, C., Vallerga, C.L., Heilbron, K., Bandres-Ciga, S., Chang, D.,

Tan, M., Kia, D.A., Noyce, A.J., Xue, A., et al. (2019). Identification of novel risk loci,

causal insights, and heritable risk for Parkinson's disease: a meta-analysis of genome-

wide association studies. Lancet Neurol *18*, 1091-1102. 10.1016/S1474-4422(19)30320-

5.

27. Karimi-Moghadam, A., Charsouei, S., Bell, B., and Jabalameli, M.R. (2018). Parkinson

Disease from Mendelian Forms to Genetic Susceptibility: New Molecular Insights into

the Neurodegeneration Process. Cell Mol Neurobiol *38*, 1153-1178. 10.1007/s10571-

018-0587-4.

28. Jia, F., Fellner, A., and Kumar, K.R. (2022). Monogenic Parkinson's Disease: Genotype,

Phenotype, Pathophysiology, and Genetic Testing. Genes (Basel) *13*.

10.3390/genes13030471.

29.   Guadagnolo, D., Piane, M., Torrisi, M.R., Pizzuti, A., and Petrucci, S. (2021). Genotype-
Phenotype Correlations in Monogenic Parkinson Disease: A Review on Clinical and
Molecular Findings. Front Neurol *12*, 648588. 10.3389/fneur.2021.648588.
30.   Lesage, S., Lunati, A., Houot, M., Romdhan, S.B., Clot, F., Tesson, C., Mangone, G.,
Toullec, B.L., Courtin, T., Larcher, K., et al. (2020). Characterization of Recessive
Parkinson Disease in a Large Multicenter Study. Ann Neurol *88*, 843-850.
10.1002/ana.25787.
31.   Pickrell, A.M., and Youle, R.J. (2015). The roles of PINK1, parkin, and mitochondrial
fidelity in Parkinson's disease. Neuron *85*, 257-273. 10.1016/j.neuron.2014.12.007.
32.   Tanaka, K. (2020). The PINK1-Parkin axis: An Overview. Neurosci Res *159*, 9-15.
10.1016/j.neures.2020.01.006.
33.   Kamienieva, I., Duszynski, J., and Szczepanowska, J. (2021). Multitasking guardian of
mitochondrial quality: Parkin function and Parkinson's disease. Transl Neurodegener *10*,
5. 10.1186/s40035-020-00229-8.
34.   Lois, C., Hong, E.J., Pease, S., Brown, E.J., and Baltimore, D. (2002). Germline
transmission and tissue-specific expression of transgenes delivered by lentiviral vectors.
Science *295*, 868-872. 10.1126/science.1067081.
35.   Schorpp, M., Jäger, R., Schellander, K., Schenkel, J., Wagner, E.F., Weiher, H., and
Angel, P. (1996/05/01). The Human Ubiquitin C Promoter Directs High Ubiquitous
Expression of Transgenes in Mice. Nucleic Acids Research *24*. 10.1093/nar/24.9.1787.
36.   Zufferey, R., Donello, J.E., Trono, D., and Hope, T.J. (1999/04). Woodchuck Hepatitis
Virus Posttranscriptional Regulatory Element Enhances Expression of Transgenes
Delivered by Retroviral Vectors. Journal of Virology *73*. 10.1128/jvi.73.4.2886-
2892.1999.
37.   Higashimoto, T., Urbinati, F., Perumbeti, A., Jiang, G., Zarzuela, A., Chang, L.J., Kohn,
D.B., and Malik, P. (2007). The woodchuck hepatitis virus post-transcriptional regulatory

element reduces readthrough transcription from retroviral vectors. Gene Therapy *14*,

1298-1304. 10.1038/sj.gt.3302979.

38.   Mosammaparast, N., Jackson, K.R., Guo, Y., Brame, C.J., Shabanowitz, J., Hunt, D.F.,

and Pemberton, L.F. (2001). Nuclear import of histone H2A and H2B is mediated by a

network of karyopherins. J Cell Biol *153*, 251-262. 10.1083/jcb.153.2.251.

39.   Marino-Ramirez, L., Kann, M.G., Shoemaker, B.A., and Landsman, D. (2005). Histone

structure and nucleosome stability. Expert Rev Proteomics *2*, 719-729.

10.1586/14789450.2.5.719.

40.   Smith, P.J., Wiltshire, M., and Errington, R.J. (2004). DRAQ5 labeling of nuclear DNA in

live and fixed cells. Curr Protoc Cytom *Chapter 7*, Unit 7 25.

10.1002/0471142956.cy0725s28.

41.   Zhitomirsky, B., Farber, H., and Assaraf, Y.G. (2018). LysoTracker and MitoTracker Red

are transport substrates of P-glycoprotein: implications for anticancer drug design

evading multidrug resistance. J Cell Mol Med *22*, 2131-2141. 10.1111/jcmm.13485.

42.   Chazotte, B. (2011). Labeling lysosomes in live cells with LysoTracker. Cold Spring Harb

Protoc *2011*, pdb prot5571. 10.1101/pdb.prot5571.

43.   Dang, Q., Zhou, H., Qian, J., Yang, L., Huang, J., Zhang, Y., and Shi, W. (2018). LAMP1

Overexpression Predicts for Poor Prognosis in Diffuse Large B-cell Lymphoma. Clin

Lymphoma Myeloma Leuk *18*, 749-754. 10.1016/j.clml.2018.07.288.

44.   Wang, Q.Q., Yao, J., Jin, Q., Wang, X.D., Zhu, H.J., Huang, F., Wang, W., Qiang, J.F.,

and Ni, Q.C. (2017). LAMP1 expression is associated with poor prognosis in breast

cancer. Oncol Lett *14*, 4729-4735. 10.3892/ol.2017.6757.

45.   Xu, Y., Cao, X., Zhang, S., Zhang, Y., and Shen, Z. (2017). High expression of LAMP1

as a prognostic marker in patients with epithelial ovarian cancer. Int J Clin Exp Pathol

10, 9104-9111.

46.   Ranjbar, M.A., and Jamshidi, M. (2022). Overexpression of Lysosome-Associated

908   Membrane Protein 1 in Oral Squamous Cell Carcinoma and its Correlation with Tumor

909   Differentiation and Metastasis. Iran J Otorhinolaryngol *34*, 3-8.

910   10.22038/IJORL.2021.51683.2772.

911   47.   Zong, S., Wu, M., Gu, J., Liu, T., Guo, R., and Yang, M. (2018). Structure of the intact

912   14-subunit human cytochrome c oxidase. Cell Res *28*, 1026-1034. 10.1038/s41422-018-

913   0071-1.

914   48.   Cunatova, K., Reguera, D.P., Houstek, J., Mracek, T., and Pecina, P. (2020). Role of

915   cytochrome c oxidase nuclear-encoded subunits in health and disease. Physiol Res *69*,

916   947-965. 10.33549/physiolres.934446.

917   49.   Partikian, A., Olveczky, B., Swaminathan, R., Li, Y., and Verkman, A.S. (1998). Rapid

918   diffusion of green fluorescent protein in the mitochondrial matrix. J Cell Biol *140*, 821-

919   829. 10.1083/jcb.140.4.821.

920   50.   Hoogewijs, K., James, A.M., Smith, R.A., Gait, M.J., Murphy, M.P., and Lightowlers, R.N.

921   (2016). Assessing the Delivery of Molecules to the Mitochondrial Matrix Using Click

922   Chemistry. Chembiochem *17*, 1312-1316. 10.1002/cbic.201600188.

923   51.   Candas, D., Qin, L., Fan, M., and Li, J.J. (2016). Experimental Approaches to Study

924   Mitochondrial Localization and Function of a Nuclear Cell Cycle Kinase, Cdk1. J Vis Exp,

925   53417. 10.3791/53417.

926   52.   Chazotte, B. (2011). Labeling mitochondria with MitoTracker dyes. Cold Spring Harb

927   Protoc *2011*, 990-992. 10.1101/pdb.prot5648.

928   53.   Durand, R.E., and Olive, P.L. (1982). Cytotoxicity, Mutagenicity and DNA damage by

929   Hoechst 33342. J Histochem Cytochem *30*, 111-116. 10.1177/30.2.7061816.

930   54.   Wiezorek, C. (1984). Cell cycle dependence of Hoechst 33342 dye cytotoxicity on sorted

931   living cells. Histochemistry *81*, 493-495. 10.1007/BF00489756.

55. Solano, R.M., Casarejos, M.J., Menendez-Cuervo, J., Rodriguez-Navarro, J.A., Garcia

de Yebenes, J., and Mena, M.A. (2008). Glial dysfunction in parkin null mice: effects of

aging. J Neurosci *28*, 598-611. 10.1523/JNEUROSCI.4609-07.2008.

56. Rouland, L., Duplan, E., Ramos Dos Santos, L., Bernardin, A., Katula, K.S., Manfioletti,

G., Idbaih, A., Checler, F., and Alves da Costa, C. (2021). Therapeutic potential of parkin

as a tumor suppressor via transcriptional control of cyclins in glioblastoma cell and

animal models. Theranostics *11*, 10047-10063. 10.7150/thno.57549.

57. Wahabi, K., Perwez, A., and Rizvi, M.A. (2018). Parkin in Parkinson's Disease and

Cancer: a Double-Edged Sword. Mol Neurobiol *55*, 6788-6800. 10.1007/s12035-018-

0879-1.

58. Yeo, C.W., Ng, F.S., Chai, C., Tan, J.M., Koh, G.R., Chong, Y.K., Koh, L.W., Foong, C.S.,

Sandanaraj, E., Holbrook, J.D., et al. (2012). Parkin pathway activation mitigates glioma

cell proliferation and predicts patient survival. Cancer Res *72*, 2543-2553. 10.1158/0008-

5472.CAN-11-3060.

59. Tay, S.P., Yeo, C.W., Chai, C., Chua, P.J., Tan, H.M., Ang, A.X., Yip, D.L., Sung, J.X.,

Tan, P.H., Bay, B.H., et al. (2010). Parkin enhances the expression of cyclin-dependent

kinase 6 and negatively regulates the proliferation of breast cancer cells. J Biol Chem

285, 29231-29238. 10.1074/jbc.M110.108241.

60. Gong, Y., Zack, T.I., Morris, L.G., Lin, K., Hukkelhoven, E., Raheja, R., Tan, I.L., Turcan,

S., Veeriah, S., Meng, S., et al. (2014). Pan-cancer genetic analysis identifies PARK2 as

a master regulator of G1/S cyclins. Nat Genet *46*, 588-594. 10.1038/ng.2981.

61. Veeriah, S., Taylor, B.S., Meng, S., Fang, F., Yilmaz, E., Vivanco, I., Janakiraman, M.,

Schultz, N., Hanrahan, A.J., Pao, W., et al. (2010). Somatic mutations of the Parkinson's

disease-associated gene PARK2 in glioblastoma and other human malignancies. Nat

Genet *42*, 77-82. 10.1038/ng.491.

62.   Frade, J.M., and Ovejero-Benito, M.C. (2015). Neuronal cell cycle: the neuron itself and
its circumstances. Cell Cycle *14*, 712-720. 10.1080/15384101.2015.1004937.

63.   Kruman, II, Wersto, R.P., Cardozo-Pelaez, F., Smilenov, L., Chan, S.L., Chrest, F.J.,
Emokpae, R., Jr., Gorospe, M., and Mattson, M.P. (2004). Cell cycle activation linked to
neuronal cell death initiated by DNA damage. Neuron *41*, 549-561. 10.1016/s0896-
6273(04)00017-0.

64.   Gupta, R., Jha, A., Ambasta, R.K., and Kumar, P. (2021). Regulatory mechanism of
cyclins and cyclin-dependent kinases in post-mitotic neuronal cell division. Life Sci *285*,
120006. 10.1016/j.lfs.2021.120006.

65.   Gan-Or, Z., Dion, P.A., and Rouleau, G.A. (2015). Genetic perspective on the role of the
autophagy-lysosome pathway in Parkinson disease. Autophagy *11*, 1443-1457.
10.1080/15548627.2015.1067364.

66.   Navarro-Romero, A., Montpeyo, M., and Martinez-Vicente, M. (2020). The Emerging
Role of the Lysosome in Parkinson's Disease. Cells *9*. 10.3390/cells9112399.

67.   Hopfner, F., Mueller, S.H., Szymczak, S., Junge, O., Tittmann, L., May, S., Lohmann, K.,
Grallert, H., Lieb, W., Strauch, K., et al. (2020). Rare Variants in Specific Lysosomal
Genes Are Associated With Parkinson's Disease. Mov Disord *35*, 1245-1248.
10.1002/mds.28037.

68.   Dehay, B., Martinez-Vicente, M., Caldwell, G.A., Caldwell, K.A., Yue, Z., Cookson, M.R.,
Klein, C., Vila, M., and Bezard, E. (2013). Lysosomal impairment in Parkinson's disease.
Mov Disord *28*, 725-732. 10.1002/mds.25462.

69.   Klein, A.D., and Mazzulli, J.R. (2018). Is Parkinson's disease a lysosomal disorder?
Brain *141*, 2255-2262. 10.1093/brain/awy147.

70.   Yoshimori, T., Yamamoto, A., Moriyama, Y., Futai, M., and Tashiro, Y. (1991). Bafilomycin
A1, a specific inhibitor of vacuolar-type H(+)-ATPase, inhibits acidification and protein
degradation in lysosomes of cultured cells. J Biol Chem *266*, 17707-17712.

71.   Mauvezin, C., and Neufeld, T.P. (2015). Bafilomycin A1 disrupts autophagic flux by

inhibiting both V-ATPase-dependent acidification and Ca-P60A/SERCA-dependent

autophagosome-lysosome fusion. Autophagy *11*, 1437-1438.

10.1080/15548627.2015.1066957.

72.   Carpenter, A.E., Jones, T.R., Lamprecht, M.R., Clarke, C., Kang, I.H., Friman, O.,

Guertin, D.A., Chang, J.H., Lindquist, R.A., Moffat, J., et al. (2006). CellProfiler: image

analysis software for identifying and quantifying cell phenotypes. Genome Biol *7*, R100.

10.1186/gb-2006-7-10-r100.

73.   Johnson, D.E., Ostrowski, P., Jaumouille, V., and Grinstein, S. (2016). The position of

lysosomes within the cell determines their luminal pH. J Cell Biol *212*, 677-692.

10.1083/jcb.201507112.

74.   Heuser, J. (1989). Changes in lysosome shape and distribution correlated with changes

in cytoplasmic pH. J Cell Biol *108*, 855-864. 10.1083/jcb.108.3.855.

75.   Parton, R.G., Dotti, C.G., Bacallao, R., Kurtz, I., Simons, K., and Prydz, K. (1991). pH-

induced microtubule-dependent redistribution of late endosomes in neuronal and

epithelial cells. J Cell Biol *113*, 261-274. 10.1083/jcb.113.2.261.

76.   Pu, J., Guardia, C.M., Keren-Kaplan, T., and Bonifacino, J.S. (2016). Mechanisms and

functions of lysosome positioning. J Cell Sci *129*, 4329-4339. 10.1242/jcs.196287.

77.   Song, P., Trajkovic, K., Tsunemi, T., and Krainc, D. (2016). Parkin Modulates Endosomal

Organization and Function of the Endo-Lysosomal Pathway. J Neurosci *36*, 2425-2437.

10.1523/JNEUROSCI.2569-15.2016.

78.   Okarmus, J., Bogetofte, H., Schmidt, S.I., Ryding, M., Garcia-Lopez, S., Ryan, B.J.,

Martinez-Serrano, A., Hyttel, P., and Meyer, M. (2020). Lysosomal perturbations in

human dopaminergic neurons derived from induced pluripotent stem cells with PARK2

mutation. Sci Rep *10*, 10278. 10.1038/s41598-020-67091-6.

79.   Twig, G., Elorza, A., Molina, A.J., Mohamed, H., Wikstrom, J.D., Walzer, G., Stiles, L.,

1009   Haigh, S.E., Katz, S., Las, G., et al. (2008). Fission and selective fusion govern

1010   mitochondrial segregation and elimination by autophagy. EMBO J *27*, 433-446.

1011   10.1038/sj.emboj.7601963.

1012   80.   Chen, H., Chomyn, A., and Chan, D.C. (2005). Disruption of fusion results in

1013   mitochondrial heterogeneity and dysfunction. J Biol Chem *280*, 26185-26192.

1014   10.1074/jbc.M503062200.

1015   81.   Sarrafha, L., Parfitt, G.M., Reyes, R., Goldman, C., Coccia, E., Kareva, T., and Ahfeldt, T.

1016   (2021). High-throughput generation of midbrain dopaminergic neuron organoids from

1017   reporter human pluripotent stem cells. STAR Protoc *2*, 100463.

1018   10.1016/j.xpro.2021.100463.

1019   82.   Wang, X., Winter, D., Ashrafi, G., Schlehe, J., Wong, Y.L., Selkoe, D., Rice, S., Steen, J.,

1020   LaVoie, M.J., and Schwarz, T.L. (2011). PINK1 and Parkin target Miro for

1021   phosphorylation and degradation to arrest mitochondrial motility. Cell *147*, 893-906.

1022   10.1016/j.cell.2011.10.018.

1023   83.   Hernandez, G., Thornton, C., Stotland, A., Lui, D., Sin, J., Ramil, J., Magee, N., Andres,

1024   A., Quarato, G., Carreira, R.S., et al. (2013). MitoTimer: a novel tool for monitoring

1025   mitochondrial turnover. Autophagy *9*, 1852-1861. 10.4161/auto.26501.

1026   84.   Laker, R.C., Xu, P., Ryall, K.A., Sujkowski, A., Kenwood, B.M., Chain, K.H., Zhang, M.,

1027   Royal, M.A., Hoehn, K.L., Driscoll, M., et al. (2014). A novel MitoTimer reporter gene for

1028   mitochondrial content, structure, stress, and damage in vivo. J Biol Chem *289*, 12005-

1029   12015. 10.1074/jbc.M113.530527.

1030   85.   Hanson, G.T., Aggeler, R., Oglesbee, D., Cannon, M., Capaldi, R.A., Tsien, R.Y., and

1031   Remington, S.J. (2004). Investigating mitochondrial redox potential with redox-sensitive

1032   green fluorescent protein indicators. J Biol Chem *279*, 13044-13053.

1033   10.1074/jbc.M312846200.

1034   86.   Yokomakura, A., Hong, J., Ohuchi, K., Oh, S.E., Lee, J.Y., Mano, N., Takahashi, T.,

1035   Hwang, G.W., and Naganuma, A. (2012). Increased production of reactive oxygen

1036   species by the vacuolar-type (H(+))-ATPase inhibitors bafilomycin A1 and concanamycin

1037   A in RAW 264 cells. J Toxicol Sci *37*, 1045-1048. 10.2131/jts.37.1045.

1038   87.   Palacino, J.J., Sagi, D., Goldberg, M.S., Krauss, S., Motz, C., Wacker, M., Klose, J., and

1039   Shen, J. (2004). Mitochondrial dysfunction and oxidative damage in parkin-deficient

1040   mice. J Biol Chem *279*, 18614-18622. 10.1074/jbc.M401135200.

1041   88.   Cornejo-Olivas, M.R., Torres, L., Mata, I.F., Mazzetti, P., Rivas, D., Cosentino, C., Inca-

1042   Martinez, M., Cuba, J.M., Zabetian, C.P., and Leverenz, J.B. (2015). A Peruvian family

1043   with a novel PARK2 mutation: Clinical and pathological characteristics. Parkinsonism

1044   Relat Disord *21*, 444-448. 10.1016/j.parkreldis.2015.01.005.

1045   89.   Orimo, S., Amino, T., Yokochi, M., Kojo, T., Uchihara, T., Takahashi, A., Wakabayashi, K.,

1046   Takahashi, H., Hattori, N., and Mizuno, Y. (2005). Preserved cardiac sympathetic nerve

1047   accounts for normal cardiac uptake of MIBG in PARK2. Mov Disord *20*, 1350-1353.

1048   10.1002/mds.20594.

1049   90.   Gouider-Khouja, N., Larnaout, A., Amouri, R., Sfar, S., Belal, S., Ben Hamida, C., Ben

1050   Hamida, M., Hattori, N., Mizuno, Y., and Hentati, F. (2003). Autosomal recessive

1051   parkinsonism linked to parkin gene in a Tunisian family. Clinical, genetic and pathological

1052   study. Parkinsonism Relat Disord *9*, 247-251. 10.1016/s1353-8020(03)00016-6.

1053   91.   Hayashi, S., Wakabayashi, K., Ishikawa, A., Nagai, H., Saito, M., Maruyama, M.,

1054   Takahashi, T., Ozawa, T., Tsuji, S., and Takahashi, H. (2000). An autopsy case of

1055   autosomal-recessive juvenile parkinsonism with a homozygous exon 4 deletion in the

1056   parkin gene. Mov Disord *15*, 884-888. 10.1002/1531-8257(200009)15:5<884::aid-

1057   mds1019>3.0.co;2-8.

1058   92.   Schmidt, S., Linnartz, B., Mendritzki, S., Sczepan, T., Lubbert, M., Stichel, C.C., and

1059   Lubbert, H. (2011). Genetic mouse models for Parkinson's disease display severe

pathology in glial cell mitochondria. Hum Mol Genet *20*, 1197-1211.

10.1093/hmg/ddq564.

93.   Gerasimova, T., Stepanenko, E., Novosadova, L., Arsenyeva, E., Shimchenko, D.,

Tarantul, V., Grivennikov, I., Nenasheva, V., and Novosadova, E. (2023). Glial Cultures

Differentiated from iPSCs of Patients with PARK2-Associated Parkinson's Disease

Demonstrate a Pro-Inflammatory Shift and Reduced Response to TNFalpha Stimulation.

Int J Mol Sci *24*. 10.3390/ijms24032000.

94.   Hastie, E., Cataldi, M., Marriott, I., and Grdzelishvili, V.Z. (2013). Understanding and

altering cell tropism of vesicular stomatitis virus. Virus Res *176*, 16-32.

10.1016/j.virusres.2013.06.003.

95.   Sweeney, N.P., and Vink, C.A. (2021). The impact of lentiviral vector genome size and

producer cell genomic to gag-pol mRNA ratios on packaging efficiency and titre. Mol

Ther Methods Clin Dev *21*, 574-584. 10.1016/j.omtm.2021.04.007.

96.   Desfarges, S., and Ciuffi, A. (2010). Retroviral integration site selection. Viruses *2*, 111-

130. 10.3390/v2010111.

97.   Bonam, S.R., Wang, F., and Muller, S. (2019). Lysosomes as a therapeutic target. Nat

Rev Drug Discov *18*, 923-948. 10.1038/s41573-019-0036-1.

98.   Zhang, Z., Yue, P., Lu, T., Wang, Y., Wei, Y., and Wei, X. (2021). Role of lysosomes in

physiological activities, diseases, and therapy. J Hematol Oncol *14*, 79. 10.1186/s13045-

021-01087-1.

99.   Javadov, S., Kozlov, A.V., and Camara, A.K.S. (2020). Mitochondria in Health and

Diseases. Cells *9*. 10.3390/cells9051177.

100.   Chen, W., Zhao, H., and Li, Y. (2023). Mitochondrial dynamics in health and disease:

mechanisms and potential targets. Signal Transduct Target Ther *8*, 333.

10.1038/s41392-023-01547-9.

101.   Dull, T., Zufferey, R., Kelly, M., Mandel, R.J., Nguyen, M., Trono, D., and Naldini, L.

1086   (1998). A third-generation lentivirus vector with a conditional packaging system. J Virol

1087   *72*, 8463-8471. 10.1128/JVI.72.11.8463-8471.1998.

1088   102.   Morrone Parfitt, G., Coccia, E., Goldman, C., Whitney, K., Reyes, R., Sarrafha, L., Nam,

1089   K.H., Sohail, S., Jones, D.R., Crary, J.F., et al. (2024). Disruption of lysosomal

1090   proteolysis in astrocytes facilitates midbrain organoid proteostasis failure in an early-

1091   onset Parkinson's disease model. Nat Commun *15*, 447. 10.1038/s41467-024-44732-2.

1092   103.   Chaudhry, A., Shi, R., and Luciani, D.S. (2020). A pipeline for multidimensional confocal

1093   analysis of mitochondrial morphology, function, and dynamics in pancreatic beta-cells.

1094   Am J Physiol Endocrinol Metab *318*, E87-E101. 10.1152/ajpendo.00457.2019.

1095   104.   Mangeol, P., Prevo, B., and Peterman, E.J. (2016). KymographClear and

1096   KymographDirect: two tools for the automated quantitative analysis of molecular and

1097   cellular dynamics using kymographs. Mol Biol Cell *27*, 1948-1957. 10.1091/mbc.E15-06-

1098   0404.

1099   105.   Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A., and

1100   Goodman, A. (2021). CellProfiler 4: improvements in speed, utility and usability. BMC

1101   Bioinformatics *22*, 433. 10.1186/s12859-021-04344-9.

1102

Figure 1

Figure 2

Figure 3

Figure 4

A.

UbC — Cox8A (Mitochondria) — mEmerald — Empty (No Selection)

B.

| COX8A-Emerald | Composite | High Mag. Composite |

0 µM Oligomycin

2 µM Oligomycin

C.

Mean Mitochondrial Area (µm²) — Vehicle vs Olig. ***

Mean Branch Length (µm) — Vehicle vs Olig. ****

Mean Aspect Ratio — Vehicle vs Olig. ****

D.

Anterograde Transport
0s
6s
13s
⟵ Soma   Synapse ⟶

Retrograde Transport
0s
14s
34s
⟵ Soma   Synapse ⟶

5um

E. **Anterograde Transport**

Run Length (µm) — $PRKN^{+/+}$ vs $PRKN^{-/-}$ ***

Run Time (s) — $PRKN^{+/+}$ vs $PRKN^{-/-}$ **

F. **Retrograde Transport**

Run Length (µm) — $PRKN^{+/+}$ vs $PRKN^{-/-}$ ns

Run Time (s) — $PRKN^{+/+}$ vs $PRKN^{-/-}$ *

Figure 5

Figure 6

Figure S1

Fig. S1: GEM-SCOPe supports a diversity of modifications and applications (A) Representative images of astrocytes expressing COX8A-Emerald under the control of a doxycycline inducible promoter. Astrocytes were treated with a vehicle (DMSO; top) or doxycycline (bottom) for 72 hours. Nuclei were stained with Hoechst33342. **(B)** qPCR for endogenous *LAMP1* on RNA from astrocytes transduced with a virus localized to the lysosome (LAMP1-mCherry) or a virus not localized to the lysosome (COX8A-Emerald). qPCR primers were designed to amplify a sequence of *LAMP1* that is not included in the lentivirus to only quantify endogenous *LAMP1* levels. Bars represent mean and error bars represent standard deviation (n = 3 independent transductions). **(C)** Representative images of a mixed population of astrocytes with nuclei expressing different H2B-fusion fluorophores. Astrocytes were transduced with a single lentivirus and then plated together. Scale bar = 50 µm.

Figure S2

Fig. S2: Mitochondrial pH sensor measures disrupted mitochondrial membrane potential Schematic of the lentiviral construct used in the following panels localizing Lemon, a pH responsive fluorophore, to the mitochondria with the targeting sequence of COX8A. Lemon undergoes Forster Resonance Energy Transfer (FRET) under alkaline conditions. **(B)** Representative images of astrocytes transduced with COX8A-Lemon and treated with a vehicle (DMSO; top) or 2 µM oligomycin (bottom) for 24 hours. Images were acquired with a 455nm-excitation laser and a 515-excitation laser. Far right: ratiometric representation of the emission in the yellow channel divided by the emission in the cyan channel and then pseudo colored so that orange, yellow and white indicate more relative yellow emission (more alkaline) while black and purple represent more cyan emission (more acidic). Nuclei were stained with DRAQ5. Scale bars = 50 µm. **(C)** Quantification of the ratio of yellow emission to cyan emission in astrocytes expressing COX8A-Lemon and treated with DMSO or oligomycin. Central bars represent mean and error bars represent standard deviation (n = 3; each replicate is an average 36 images).

Figure S3

Fig. S3: Astrocyte proliferation is activated by FBS and inhibited by MG-132 (A) Representative images of astrocytes transduced with H2B-Emerald lentivirus after 0 hrs (top) or 48 hrs (bottom) of treatment with a vehicle (DMSO; left), 10% fetal bovine serum (FBS; middle), or MG132 (right). Scale bars = 50 μm. **(B)** Quantification of number of H2B-Emerald nuclei per image field 0 hours and 48 hours after treatment. Dots represent mean values and error bars represent standard deviation (n = 6 (n = 3 for MG-132 treatment) per time point; each replicate is an average over 25 images). Scale bars = 50 μm. All images were acquired on the CX7 HCS platform with a 20x objective lens.

Figure S4

Fig. S4: Mitochondrial fragmentation exhibits a dose dependent response to oligomycin-induced stress

(A) Representative images of astrocytes transduced with COX8A-Emerald and treated with a vehicle (DMSO), 1µM, 2 µM, or 5 µM oligomycin for 4 hours. Nuclei were stained with Hoechst33342. Scale bars = 50 µm **(B)** Quantification of mitochondrial features indicative of mitochondrial network fragmentation: mean area, perimeter, aspect ratio, number of branches, branch length, and number of branch endpoints. Central bars represent mean and error bars represent standard deviation (n = 3; each replicate is an average over 36 images).

Plasmid Name	AddGene Cat#	Localization	Fluorophore	Antibiotic Resistance	Relevant Figures
pFUW H2B-mTagBFP2-p2A-PuroR		Nucleus	mTagBFP2	Puromycin	1, 2, S1, S3
pFUW H2B-Emerald-p2A-PuroR		Nucleus	Emerald	Puromycin	2, S1
pFUW H2B-mCherry-p2A-PuroR		Nucleus	mCherry	Puromycin	2, S1
pFUW LAMP1Sig-Emerald		Lysosome	Emerald	NA	1, 3
pFUW LAMP1Sig-mCherry		Lysosome	mCherry	NA	1
pFUW COX8ASig-Emerald		Mitochondria	Emerald	NA	1, 4, S5
pFUW COX8ASig-mCherry		Mitochondria	mCherry	NA	1
pFUW PuroR-P2A-COX8ASig-Timer		Mitochondria	Timer	Puromycin	5
pFUW COX8ASig-Lemon		Mitochondria	Lemon	NA	S2
pFUW mito-GRX1-roGFP2		Mitochondria	roGFP2	NA	6
pFUW mTagBFP2		NA	mTagBFP2	NA	NA
pFUW Emerald		NA	Emerald	NA	NA
pFUW mCherry		NA	mCherry	NA	NA
pFUW mTagBFP2-P2A-PuroR		NA	mTagBFP2	Puromycin	NA
pFUW Emerald-P2A-PuroR		NA	Emerald	Puromycin	3
pFUW mCherry-P2A-PuroR		NA	mCherry	Puromycin	NA
pFUW mTagBFP2-P2A-NeoR		NA	mTagBFP2	Neomycin	NA
pFUW Emerald-P2A-NeoR		NA	Emerald	Neomycin	NA
pFUW mCherry-P2A-NeoR		NA	mCherry	Neomycin	NA

Table S1: A list of all plasmids generated in this study. Details for all plasmids, including subcellular localization, fluorophore, antibiotic resistance, and in which figure each plasmid was used. More plasmids were generated as part of GEM-SCOPe than used in the final publication, and these are also included here and available on Addgene.

Video S1: Mitochondrial axonal transport in *PRKN⁺/⁺* and *PRKN⁻/⁻* human iPSC derived neurons

Representative videos of COX8A-Emerald labeled mitochondria moving within neuronal axons over the course of 1 hour. Images were taken every 10 seconds and sped up 70x. Scale bar = 25 μm.