Single cell RNA-sequencing of feline peripheral immune cells with V(D)J repertoire and cross species analysis of T lymphocytes

Raneesh Ramarapu[1,2], Judit M. Wulcan[3], Haiyang Chang[4], Peter F. Moore[3], William
Vernau[3], Stefan M. Keller[3]

[1]Department of Surgical and Radiological Sciences, School of Veterinary Medicine, University
of California Davis, Davis, CA, 95616, USA.

[2]Department of Anatomy, Physiology and Cell Biology, School of Veterinary Medicine,
University of California Davis, Davis, CA, 95616, USA.

[3]Department of Pathology, Microbiology and Immunology, School of Veterinary Medicine,
University of California, Davis, CA, United States.

[4]Department of Mathematics and Statistics, University of Guelph, Guelph, ON, Canada.

Abstract

Introduction: The domestic cat (*Felis catus*) is a valued companion animal and the second most
popular pet (over 46.5 million US households). They are also an important model system for
virally-induced cancers (feline leukemia virus) and virally-mediated immunodeficiency (feline
immunodeficiency virus). However, species specific research limitations such as a lack of
reagents and immune cell markers limit our ability to utilize these models to their full capacity.
The goal of this study is to characterize the transcriptomic landscape of circulating feline T cells
and other captured leukocytes utilizing CD5 (only available selective feline T cell marker) flow
cytometry enriched single cell RNA-sequencing and V(D)J repertoire analysis in clinically
healthy domestic shorthair cats between the ages of 6 months and 9 years as well as
contextualize them with the leukocytes of other species.

Methods: 5 mL of peripheral whole blood was collected from 4 healthy cats (6 months, 1 year, 4
years, 9 years) and were housed at the UC Davis Nutrition and Pet Care Center. Samples were
enriched for T cells by flow cytometry using a mouse anti-feline CD5 monoclonal antibody. The
GEX library was constructed using the Chromium Next GEM Single Cell 5' Kit v2 (10x
Genomics). The V(D)J library for all 4 T cell receptor loci was prepared using the Chromium
Single Cell V(D)J Enrichment Kit with custom reverse primers for the feline orthologs.
Sequencing was done via Illumina NovaSeq S4 and pre-processed using the Cell Ranger for
alignment to the feline genome (felis_catus_9.0). Processing and downstream analyses were
conducted via Seurat v5. Additional annotated datasets for cross species T cell integration were
acquired from peer-reviewed literature.

Results: Unsupervised clustering of GEX data revealed 7 major populations – T cells, neutrophils, monocytic cells, B cells, plasmacytoid dendritic cells, mast cells and platelets. Sub cluster analysis of T cells resolved different populations of naive ($CD4^+$ and $CD8^+$), $CD4^+$ effector T cells, $CD8^+$ cytotoxic T cells and γδ T cells for the first time. Cross species analysis revealed relatively high conservation of T cell subtypes along an effector gradient with equitable representation of veterinary species (horse, dog, pig) and humans with the cat. Our V(D)J repertoire analysis demonstrated marked differences in $CD8^+$ cytotoxic T cells from other alpha/beta T cell subsets including productive TRG transcript expression. Among the myeloid cells, we resolved 3 clusters of classical monocytes with polarization into pro- and anti-inflammatory phenotypes in addition to a cluster of conventional dendritic cells. Lastly, our neutrophil sub clustering revealed a larger mature neutrophil cluster and a smaller exhausted/activated cluster.

Discussion: Our study is the first to characterize the subtypes of circulating T cells utilizing an integrative approach of single cell RNA-sequencing, V(D)J repertoire analysis and cross species analysis. Additionally, we also characterize subtypes in the myeloid cells expanding our understanding of the feline immune system. We have also demonstrated species immune cell relatedness which can provide an important foundation for further translational immunology, pathogenesis, translational treatments, and species-tropism of pathogens in veterinary medicine and research.

2   Introduction

The domestic cat, Felis catus, is a valued companion animal and the second most popular pet in the United States. As of the American Veterinary Medicine Association (AVMA) pet ownership statistics, there are upwards of 46.5 million households with at least one cat in 2023 with ownership trends rising in the last decade from 25% in 2016 to 29% in 2022 (1,2). In addition to the growing need for feline veterinary care, the cat is an important model for infectious diseases such as virally-induced cancers (Feline leukemia virus, FeLV) and virally mediated immunodeficiency (Feline immunodeficiency virus, FIV) (3–6). Despite the value of the cat as a companion animal and infectious disease model, there are species specific research limitations including few cat-reactive immunophenotyping reagents and a poor knowledge base of immune cell markers and behavior.

In the advent of deciphering the feline immune system, many studies characterizing feline immune cells have relied strongly on antibody-based assays including flow cytometry and immunohistochemistry. This has allowed the study of immune cells in health and disease conditions such as FeLV or FIV (7–9). This approach depends on the assumption that feline immune cells are similar to other species. However, feline immune mediated diseases behave very differently from those in other small animals such as dogs, exhibiting unique etiologies and pathogenesis (10). In a cross-species context, innate immunity shows relatively high evolutionary conservation; however, this conservation dwindles with progression to adaptive

immunity and complex immune phenotypes (11). Additionally, there is growing evidence demonstrating that evolutionary relationships do not translate to immune transcriptional or cell type relationships. Such an example is the mouse being a better model for human immune cells than the macaque (12). Thus, understanding the species-specific system in the context of our growing evolutionary database is desirable and there is a strong need to characterize the heterogeneous peripheral immune cell population in the cat, especially diverse populations such as T cells.

In efforts to investigate feline specific T cell diseases, our laboratory has contributed to the growing feline specific reagent and immune cell knowledge base including T cell receptor (TCR) repertoire analysis (13). In this trajectory, single cell RNA-sequencing is a logical next step. It enables the efficient large-scale capture and analysis of heterogeneous tissues, overcoming the challenges of species-specific reagent assays (14–17). Consequently, there has been a bloom in single cell RNA-sequencing studies of the peripheral leukocytes in a variety of mammalian species including the dog, cow, horse, and pig (18–22). Additionally, single cell data facilitates the study of cross species variations in immune cell type and evolution of these cells (12). ScRNA-seq of feline circulating leukocytes have been previously performed demonstrating the presence of 5 major cell types (T cells, B cells, NK cells, monocytes, dendritic cells) in the context of other non-model species (23). However, in-depth characterization of key subtypes and their marker conservation is yet to be determined. Thus, the goal of this study is to characterize the heterogeneity of circulating feline T cells and other captured leukocytes utilizing CD5 flow cytometry enriched scRNA-seq and V(D)J repertoire analysis in clinically healthy domestic shorthair cats between the ages of 6 months and 9 years. CD5 was chosen as a selective T-cell marker given the lack of availability of anti-feline CD3 antibodies. Additionally, we performed a cross species transcriptomic integration to assess feline T cells in the context of 4 other mammalian species - dog, horse, human, and pig. Our results provide a foundation for feline immune transcriptomics.

3   Results
3.1   Single cell atlas of the feline circulating immune cells
ScRNA-seq data obtained from 4 representative healthy cats of different ages consisted of 30,073 quality cells. These age groups were selected as they capture the major feline life stages: 6-month-old juvenile (6MO), 1-year-old mature (1YO), 4-year-old adult (4YO), and 9-year-old aged (9YO). The average number of cells captured per cat was 7,518 with an average of 3,655 transcripts per cell. **<Fig1A, Supplementary File1 S1A,B>**. Due to the poor annotation of the cat reference genome, we utilized a custom homologous mapping script based on ENSEMBL homologous mapping between the cat and human to improve gene readability. Our script increased the number of gene symbols mapped from 13,886 to 16,936, improving the quality of cell type assignment and biological interpretation. Following batch correction via reciprocal principal component analysis, unsupervised clustering revealed the presence of 21 clusters **<Fig**

1B>. A majority of cell types are equally represented from each sample **<Fig1C, 1D>**. A quality control violin plot for clusters can be found in the supplementary materials **<Supplementary File1 S2A>**.

These clusters were categorized into 7 major immune cell types based on canonical marker genes from multi-species PBMC single cell literature, top marker genes and over-enrichment gene ontology (GO) **<Supplementary File 2,3>**. The most numerous cell type identified was T cells called by high expression of T cell receptor complex genes (*CD3D, CD3E*) and species conserved T cell markers *LCK*, *ITK* and *TCF7* (12,23) **<Fig 1E-G, Supplementary File1 S2B>**. Across most T cell clusters, top identified GO terms of biological processes (BP) included T leukocyte processes of differentiation, migration, activation, and effector functions **<Supplementary File 3>**.

The second most abundant population was the neutrophils marked by neutrophil-derived proteases including elastase (*ELANE*), granulocyte colony stimulating factor receptor (*CSF3R*) and calprotectin (*S100A8/S100A9*) (24–26) **<Fig 1E,H>**. In most species, neutrophils are not captured in the PBMC layer during density centrifugation (27). Interestingly, our single cell data set captured a significant number of neutrophils from all but the 1YO sample **<Fig 1D>**. This is the first description of feline neutrophils at the single cell transcriptomic level and can provide insights into biological processes of these abundant cells.

A substantial number of monocytic cells were captured with marker genes including scavenger receptor for haptoglobin-hemoglobin complexes *CD163*, antigen presenting cell co-stimulatory molecule *CD83*, lipopolysaccharide detector protein *CD14*, phospholipase B domain containing 1 (*PLBD1*) and colony stimulating factor receptor (*CSF1R*) (28,29) **<Fig 1E,I,J>**. A relatively small but heterogeneous population of B cells were captured which were identified by conserved markers including B cell surface molecule *MS4A1* and Ig-alpha protein of the B-cell antigen component *CD79A* (23) **<Fig 1E,K,L>**. This population also expresses immunoglobulin genes (*IGHG3, IGHM*) indicating a sub-population of plasma cells (12) **<Fig 1E>**.

Unexpectedly, we captured plasmacytoid dendritic cells (pDC) and mast cells, which represented rare but discrete cell populations. pDCs were identified by expression of Fc fragment of gamma immunoglobulin (*FCRLA)*, dendritic marker *PLAC8B*, and immunoglobulin lambda constant 7 (*IGLC7*) and lack of expression of B cell canonical markers *MS4A1* and *CD79A* (30–32) **<Fig 1E,L,M>**. pDcs clustered more closely with B cells than other myeloid cells. This is due to close lineage associations and the shared expression of integral markers *FCRLA* and *IGLC7* among others (33) **<Fig 1E,L>**. Currently, there is debate regarding the reclassification of pDCs as innate lymphocytes and this analysis further supports such an argument (34) **<Fig1B>**. Mast cells were marked by IgE receptor *MS4A2* and a key component of mast cell protease, *CPA3* (35,36) **<Fig 1E,N>**.

Lastly, we captured a population of non-leukocytic immune cells which were annotated as
platelets due to high expression of platelet specific proteins *PPBP*, *VWF* and *GP9* (37)
<Fig1E,O>. This cluster also exhibited an extremely low average count of unique RNA and total
RNA, which is consistent with a highly specialized cell type lacking a nucleus **<Supplementary**
File1 S2A>.

3.2   Feline circulating T cells
To further characterize T cell subsets, we independently analyzed T cell clusters previously
described at the global level (clusters 0,2,9,1,6,4,5,14,8,12, and 15). Unsupervised clustering of
T cells revealed the presence of seven T cell clusters, which were unrecognizable at the global
level **<Fig 2A>**. Largely, these clusters represent greater T cell phenotypes along increasing
pseudotime values (proxy of differentiation or lineage) **<Fig 2B,C>**. The subtle transcriptional
differences, as noted through overlapping differentially expressed genes, make effective
annotation difficult **<Supplementary File 4>**. Nonetheless, we were able to identify several
canonical subsets.

T cell subsets included $CD4^+$, $CD8A^+/CD8B^+$ and γδ T cells and were equally represented in all
age groups **<Fig2D-H>**. A great majority were identified as naive cells, distinguished by their
high expression of *CCR7* and L-selectin (*SELL*) (38–40) **<Fig2I,J>**. These naive populations
segregated into two larger groups which could be distinguished by differential expression of
CD4 and *CD8A*/*CD8B* **<Fig2F-H>**. The $CD8A^+/CD8B^+$ naive population represents a
divergence from the traditionally observed distribution of naive T cells. Most naive T cells in
mammalian species are $CD4^+$ (12). However, a large population of $CD8A^+/CD8B^+$ naive cells
has been described in non-model species such as the horse (22). This highlights a potentially
influential role of $CD8A^+/CD8B^+$ T cells in cats.

The clusters 3 and 4, adjacent to the large naive populations, were identified as effector cells
(T_{EM}) **<Fig2A>**. These clusters show a gradual upregulation of T effector molecules including
S100A11, galectins (*LGALS1, LGALS3),* annexins (*ANXA1, ANXA2*), and granzymes (*GZMA,*
GZMK) (41,42) **<Fig2C,K-M>**. Szabo et al identified *CCL5* as a critical marker for highly
differentiated effector T cells across tissues as well as its antagonistic expression to the naive
marker *SELL* (41). In our data, Cluster 6 expressed the highest levels of cytotoxic markers
including *CCL5* and exhibited the highest pseudotime score with no naive marker expression
<Fig2C,M-O>. This indicates that this cluster is most consistent with the annotation of a
terminally differentiated T cell cluster ($CD8^+$ cytotoxic). Additionally, some effector cells
expressed genes associated with T cell exhaustion such as *PDCD1* and *TIGIT* indicating the
presence of a few of these cells in circulation **<Fig2C,P,Q>**.

Cluster 5 represented γδ T cells as inferred by the overlapping expression of the TCR genes *TRG*
and *TRDC* (13) **<Fig2E>**. Within this population, there was an expression gradient of naive as
well as effector markers, indicating subtypes within the γδ population **<Fig 2I,K,L>**. Although
these T cell clusters represent biologically relevant groups statistically, the literature across
various species identifies significant challenges associated with clustering of transcriptionally
similar but heterogenous T cell populations (43). The γδ cluster exemplifies this challenge in that
the cluster's defining features overlap between two different but not mutually exclusive
phenotypes - γδ recombination and effectorness/naiveness. These cells exhibit overlapping
expressions of *TRG* and *TRDC*, indicating their TCR recombination-based categorization. An
overwhelming majority of these γδ T cells are naive given their expression of naive marker *SELL*
<Fig 2I,J>. However, some cells appear to express features of effectorness **<Fig 2K-N>**. This
suggests some limitations of transcriptionally driven phenotyping of T cells. Nonetheless, this
investigation represents a significant contribution towards deciphering the biology of feline T
cells.
203
3.3   Feline circulating T effector cells
As described previously, subtyping T cells with overlapping phenotypes represents a significant
challenge, which was further exacerbated by the low number of effector T cells in our dataset.
Hence, we clustered effector T cells (clusters 3,4) independently, revealing 13 effector clusters
<Fig3A>. Many of these clusters represent minimally divergent biological states as reflected by
the marked overlap of top differentially expressed genes **<Fig3B, Supplementary File 5>**.
However, direct visualization of known phenotypes demonstrate that the effector populations
captured exhibit phenotypes of human T_H1, T_H2, T_H17 and T_{reg} cells **<Fig3C-J>** (41,42,44).
Gene sets for each module calculation are provided in the supplementary materials
<Supplementary file 1, S3>. These phenotypes do overlap and are not consistent with identified
clusters. Nevertheless, the higher scores among closely positioned cells likely indicates the
presence of these subtypes. However, a greater sample size of effector cells would be necessary
to better resolve subtypes and their trajectory.
217
3.4   Cross species analysis of circulating T cells
To understand how the captured feline T cell populations compared to those of other domestic
animal species and humans, we performed an integrative cross-species analysis. We selected 4
species for which peer-reviewed annotated scRNA-seq data with over 15,000 cells was available.
ScRNA-seq data from three veterinary species (dog, horse and pig) and humans were chosen as
the reference for homologous genome mapping (18,21,22,45). In total, we integrated 95,366 cells
and identified 13 clusters across the 5 species **<Fig4A-F>**. These clusters were functionally
annotated based on markers for T cell identity, naive phenotype, effector phenotype and
cytotoxic phenotype in addition to their differential gene expression analysis **<Fig4G,H,**
Supplementary File 6>.
228

Although feline cells were overrepresented in some naive clusters such as 3 and 8, there was
limited representation in clusters 5, 6, and 9 particularly **<Fig4I>**. To investigate whether this
finding was associated with our enrichment protocol, we assessed *CD5* transcript expression
across all clusters **<Fig4J>**. Our results suggest that the number of $CD5^+$ cells and average *CD5*
expression were lowest in clusters 5 and 9 **<Fig4K,L>**. Additionally, we saw a direct
relationship between the number of feline T cells and the $CD5^+$ cell count in each cluster.
Particularly, clusters 5, 6, 9, 11, and 12 were most underrepresented **<Fig4M>**. These clusters
were identified as cytotoxic effector cells **<Fig4H>**. This highlights a limitation of our CD5
enrichment protocol as it may select against specific effector clusters such as $CD8^+$ cytotoxic
cells. Much of our current understanding of $CD8^+$ feline T cells is from flow cytometric studies
with no further subtyping due to lack of reagents. Thus, further investigation is necessary to
validate whether this low frequency is biological or technical.
241
3.5   V(D)J recombination
To identify patterns in T cell receptor (TR) expression, we sequenced TR alpha (TRA), TR beta
(TRB), TR gamma (TRG) and TR delta (TRD) transcripts simultaneously. The majority
(>87.0%) of $CD4^+$ naïve, $CD4^+$ T_{EM} and $CD8^+$ naïve T cells expressed TRA/TRB transcripts
only **<Fig5A>**. In contrast, only 35.0% of $CD8^+$ cytotoxic T cells expressed TRA/TRB
transcripts while 60.6% expressed TRA/TRB/TRG transcripts. γδ T cells primarily expressed
TRG/TRD transcripts (49.7%) followed by TRG/TRD/TRB (16.4%) transcripts. Of note, 11.5%
of cells had TRA/TRB/TRG transcripts, which might represent αβ T cells with an effector
phenotype similar to gd T cells. $CD8^+$ Cytotoxic T cells also exhibited differential TRAV gene
usage, and to a lesser degree, TRBV gene usage **<Fig5B,C>**. The junctional length of TRG
transcript in $CD8^+$ Cytotoxic T cells was highly skewed towards 16 amino acids and converged
on the conserved motif 'CAAWDPRGYGWAHKVF' **<Fig5D>**. Most TRG rearrangements
involved cassette 2 (TRGV2-2/TRGJ2-2) **<Fig5E>.** Interestingly, a subset of TRGV2-2 genes
rearranged to the TRGJ2-1 gene, which is located upstream of TRGV2-2.
256
257 Next, we characterized the repertoire overlap between cats and the relatedness of junctional
258 regions across cats and T cell subsets **<Fig6>**. The number of shared clonotypes was highest for
259 TRA and TRG but overall low **<Fig6A>**. When expanding the definition of shared repertoires to
260 include clonotypes with one amino acid difference, the TRA locus had the highest number, size
261 and publicity of clusters **<Fig6B,C>**.
262
263 ### 3.6   Feline circulating monocytic cells
264 Although $CD5^+$ enrichment was performed for T cell selectivity, we serendipitously captured
265 neutrophils and monocytic cells, which provides insight into the transcriptome of circulating
266 myeloid cells in cats. Clustering analysis of both cell types was performed independently. Due to
267 the low myeloid cell count in the 1YO, it was excluded from this analysis.
268

Monocytic cells clustered into 5 groups and were largely represented by the 9YO sample **<Fig 7A, Supplementary File 1, S4A>**. Cluster 3 represented neutrophils which is likely due to the close clustering of myeloid cells on the global object **<Fig 1B, Supplementary File 1, S4B>**. Cluster 0, 1 and 2 were identified as classical monocytes (CM) with all clusters having high expression of markers *PLBD1*, *CD83* and classical monocyte markers *CD14* and *VCAN* (28,29,46) **<Fig 7B-D>**. We did not identify any non-classical monocytes, based on absence of *CD16* expression, as seen in other species such as humans, horses and cattle (22,29,46,47). Cluster 4 showed an upregulation of receptor-type tyrosine-protein kinase *FLT3*, a conventional pan-dendritic cell marker (48) **<Fig 7F>**. In addition, these cells exhibited upregulation of MHC class-II molecules (*FECA-DRB*, *HLA-DRA*, *HLA-DOA*, and *HLA-DMD*) and were hence identified as conventional dendritic cells (cDC) (49) **<Fig 7G>**.

In-depth assessment of differentially expressed genes across the four monocytic clusters identified signatures that further characterized these cell types. Cluster 2 demonstrated a proinflammatory phenotype characterized by upregulation of classic proinflammatory cytokines (*IL1A*, *IL1B*), *TNF/NF-KB* pathway (*TNF*, *TNFAIP3, NFKBIA*, *NFKBIZ*, *NFKB1*) and M1-macrophage-associated proinflammatory cytokines (*CXCL10, CXCL16, CCL3, CCL4 and CCL5*) **<Fig 7G,H, Supplementary File 7,8>** (20,50–54). Cluster 1 gene ontology revealed an upregulation of leukocyte migration and chemotaxis terms in addition to upregulation of proinflammatory cytokines *IL1A* and *IL1B* **<Fig 7G,I, Supplementary File 7,8>**. However, these cells also lack *TNF* expression and upregulate anti-inflammatory cytokines *IL10* and *IL18* and proteins which promote anti-inflammatory phenotypes in macrophages such as *LCN2* and *MSRB1* (55,56) **<Fig 7G>**. They have the lowest expression of *CD86*, a T cell co-stimulatory molecule of antigen presenting cells, which is associated with an anti-inflammatory phenotype (57). Most strikingly, these cells were found to upregulate genes similar to a novel monocyte cluster found in SARS-CoV-2 infected and recovered patients. These genes include *AREG*, *EREG*, *BCL6*, *IL10,* and *IL18* which have been described as anti-inflammatory tissue repair genes **<Fig 7G>** (58,59). Thus, these cells were designated as anti-inflammatory, although they do not follow the traditional M2 polarization phenotype described in human monocytes (60). Our data suggests a novel classical monocyte subtype with an anti-inflammatory phenotype.

Cluster 0 (CM) appears the least differentiated with low expression of inflammatory cytokines and the top GO terms include more cellular rather than immune related functions **<Supplementary File 8>**. Cluster 4 (cDC) showed an upregulation of and the GO terms associated with leukocyte activation and T cell activation via TCR contact with antigen bound MHC **<Fig 7G,J, Supplementary File 7,8>**. Given the strong antigen presentation phenotype, the annotation of cDC was further corroborated.

3.7   Feline circulating neutrophils

We identified 2 major neutrophil clusters **<Fig 8A>** based on the expression of canonical
neutrophil markers including *CSF3R* and *ELANE* as well as mature neutrophil functional
markers vimentin (*VIM)*, thioredoxin (*TXN)*, *S100A9* and *S100A12*, suggesting both clusters are
comprised of mature neutrophils (25,61,62) **<Fig 8B-G>**. Cluster 1 exhibited several notable
differences compared to cluster 0. Differential gene expression analysis accompanied by gene
ontology showed an upregulation in cluster 1 of apoptotic signaling and activated neutrophil
functions such as response to lipopolysaccharide and *IL-2* production **<Fig 8H, Supplementary**
File 9,10>. Thus, we annotated this cluster as activated neutrophils. When investigating top
differentially expressed genes, we noted a higher expression of *IL1A* and *TNFAIP3*, suggesting a
proinflammatory phenotype (63) **<Fig 8I,J>**. Similarly, cluster 1 had an increased expression of
VCAN, which is upregulated in the skin in response to damage from ultraviolet light B and
reactive oxygen species (64). Lastly, cluster 1 exhibited lower RNA counts per cell, which is
consistent with findings in murine neutrophils where the number of genes and counts of RNA
decrease with neutrophil differentiation, maturation and eventual activation and function (65).
Since a pathway driving this clustering could not be resolved, we examined an interferon (IFN)
module score, which is a composite score based on expression levels of interferon stimulated
genes **<Fig 8N>**. Cluster 1 did not show an increased IFN score but the adjacent cells in cluster 0
did **<Fig 8N>**. Interferon stimulation can delay neutrophil apoptosis, which could represent one
of the potential driving factors for cells to move from cluster 0 to cluster 1 (66). This analysis is
the first to resolve circulating neutrophil transcriptomic signatures in cats at the single cell level.
328

4   Discussion

The importance of the cat as a companion animal as well as a model for infectious diseases
warrants an in-depth characterization and understanding of the feline immune system. Our study
is one of the first to characterize feline immune cell subpopulations at the single cell level. We
utilized 5' single cell RNA-sequencing with V(D)J analysis to resolve the heterogeneity of CD5$^+$
enriched peripheral blood immune cells. We resolved populations of T cells, neutrophils, B cells
and monocytes. Additionally, this atlas is the first to annotate the single cell transcriptome of
rarer cell types, plasmacytoid dendritic cells and mast cells, which are seldomly captured in
single cell atlases (67,68).
338

Among T cells, we were able to resolve different populations of naive cells including a
significant number of *CD8A*/*CD8B* naive cells. We further identified T effector cells and
terminally differentiated cytotoxic effectors. The transcriptomic gradient of T cell differentiation
was found to fit the paradigm seen with other species (23). Additionally, we captured feline γδ T
cells for the first time. Upon investigation of effector cells, we were able to demonstrate known
T helper effector phenotypes among the most differentiated effectors. Of note, the T$_{reg}$ phenotype
was found to be the most prominent. The importance of T$_{reg}$ populations has been studied in cats,

particularly in the context of feline mammary carcinoma (69–71). Our study corroborates the
large presence of this phenotype and the potential importance of T_{reg} cells in feline immunology.
348
We performed a cross species integrative analysis of T cells to take advantage of the growing
database of single cell RNA data. We found relatively high conservation of T cell subtypes along
an effector gradient with relatively equitable representation of veterinary species (horse, dog,
pig) and humans with the cat. This analysis also revealed the potential negative impacts of $CD5^+$
enrichment for T cells. Effector clusters with high *CD5* expression were found to be
underrepresented in the cat. Thus, a broader T cell selection protocol would be necessary in
future studies to further investigate effector cells in cats.
In addition to characterizing T cells based on their transcriptome, we analyzed TCR expression
on a single cell level for the first time in the cat. Cytotoxic $CD8^+$ T cells differed from other αβ T
cell subsets in multiple aspects. In addition to the expected TRA and TRB transcripts, a high
percentage of cytotoxic $CD8^+$ T cells expressed a productive TRG transcript. Given that this was
not observed in naïve CD8+ T cells, we attribute this phenomenon to differential transcriptional
regulation of naive and effector T cells. Compared to other αβ T cell subsets, cytotoxic T cells
also exhibited divergent TRAV and TRBV gene usage but not TRGV gene usage. Instead, the
junctional length of TRG transcripts was skewed towards a highly conserved16 aa motif. This
finding is of potential significance for clonality testing, an adjunct method for differentiating
reactive from neoplastic lymphoid proliferations. If the majority of cytotoxic $CD8^+$ T cells harbor
a length restricted TRG rearrangement, then reactive $CD8^+$ dominant T cell proliferations might
exhibit reduced TRG diversity, which could result in a higher false positive rate. Of note, since
other αβ T cell subsets did not express a significant number of productive TRG transcripts, it is
unknown if similar length restricted rearrangements occur in these subsets. A previous study
examining the feline TCR transcriptome across thymus, spleen and mesenteric lymph nodes did
not identify any TRG length restrictions (13). However, the previous study aimed to identify
global rearrangement patterns and hence utilized a bulk sequencing strategy, which is not suited
to resolve repertoire characteristics of individual T cell subsets.
374
A cluster analysis of TCR sequences did not identify T cell subsets with invariant TRA or TRB
chains. A previous study utilizing scRNA-seq with VDJ sequencing in dogs identified MAIT-
like cells based on restricted TRAV gene usage and restricted junctional length (19). Although
cytotoxic T cells in the present study exhibited differential TRAV gene and, to a lesser extent,
differential TRBV gene usage, there was no obvious junctional length restriction or dominant
amino acid motif. The fact that we were unable to identify MAIT-like cells in cats could reflect
their true absence or represent an artifact of CD5-based enrichment, which might have resulted
in the exclusion of $CD5^-$ or $CD5^{low}$ subsets. Additional studies with more animals, inclusion of
different tissue types and without selection bias are needed to elucidate the existence of innate-
like alpha/beta T cells in cats.
385

Among the myeloid cells, we resolved 3 clusters of classical monocytes and 1 cluster of conventional dendritic cells. Of the three clusters of monocytes, we found two to be differentiated in a somewhat polarized manner- pro-inflammatory versus anti-inflammatory. The anti-inflammatory population was unconventional in that it had an upregulation of classic pro-inflammatory cytokines *IL-1A* and IL-*1B* but also an upregulation of genes associated with a unique and novel monocytic cluster found in human COVID-19 patients such as *AREG, EREG, IL10* , *IL18* and *BCL6*. (59,72). Sub clustering of neutrophils identified a larger mature cluster and a smaller exhausted/activated cluster which showed upregulation of pro-inflammatory and apoptotic genes. Thus, analysis of these myeloid cells revealed the presence of conventional and non-conventional subtypes, providing a footing for feline circulating myeloid immunology.

Currently, T cell subtypes have only been described at the flow cytometric level, and with a focus on *CD4/CD8* populations due to paucity of species-specific reagents. Our data indicates the presence of unique subtypes of peripheral blood immune cells including $\alpha\beta\gamma$ CD8$^+$ cytotoxic cells, *CD8A/CD8B* naive T cells, plasmacytoid DC, mast cells, conventional DCs, anti-inflammatory monocytes and mature and end stage neutrophils. Our study is the first of many necessary for the further characterization of the feline immune system. Taken together, it has increased our understanding of feline circulating immune cells, including T cell subtyping and capture of rare populations previously undescribed in the cat.

5 Materials and Methods
5.1 Samples and Sorting

Whole blood was obtained from four healthy, female, domestic shorthair cats (13-101/9 years, 18-003/6 years, 21-005/1 year, 22-007/6 months) housed at the UC Davis Nutrition and Pet Care Center in accordance with the UC Davis Policy and Procedure Manual section 290-30 and Animal Use and Care protocol #22150. Blood was collected in purple top EDTA tubes and immediately processed in one batch without freezing until completion of the reverse transcription step during library preparation. Peripheral blood mononuclear cells (PBMC) were isolated from whole blood by density gravity centrifugation. Briefly, 5 mL of whole blood was layered on top of 10 mL Histopaque 1077 (density 1.077g/mL, Sigma-Aldrich, Burlington, MA, USA) in a 50 ml tube. The samples were centrifuged at 850 revolutions per minute (RCF) for 30 minutes at room temperature (RT) without break. After centrifugation, the PBMC layer at the plasma-histopaque interface was aspirated and transferred to a clean 15 ml tube. The PBMCs were then washed twice with 14 ml PBMC isolation media (500 mL Hanks' Balanced Salt Solution (HBSS), 15 mL Fetal Bovine Serum (FBS), 2 mL EDTA) by centrifuging at 850 RCF for 10 minutes at RT. Following the second washing step, the PBMC pellet was resuspended in 200 μl modified flow buffer solution (MFB, 500 ml phosphate buffer saline (PBS), 0.467g EDTA tetrasodium salt and 1% Horse serum).

The isolated PBMCs were first incubated with a primary anti-feline CD5 monoclonal antibody (clone FE 1.1.B11-IgG1-Azide, Peter F. Moore Leukocyte Antigen Biology Laboratory, UC Davis, CA, USA) or a non-cross reacting canine antibody (negative control) for 30 min at RT.

This was followed by a wash step (addition off 3 mL of MFB, followed by centrifugation at 400
RCF for 3 min at RT and resuspension in 100 - 200 μl of supernatant), and incubation with a
secondary horse-anti-mouse FITC antibody (1:99 in MFB) for 15 minutes at RT in darkness.
Finally, the stained PBMC was washed again (as above), resuspended in 500 μl of MFB and
stained with DAPI (1 μg/mL). Cells were sorted at the UC Davis Comprehensive Cancer Center
Flow Cytometry Core. Cells were initially gated based on FSC vs. SSC to exclude debris and
aggregates. Singlet gating was then applied based on SSC to exclude doublets, followed by
discrimination of live and dead cells based on DAPI staining and subsequent gating for CD5
positive (FITC positive) cells **<Supplementary File 11>**.

5.2   Single-cell 5' RNA and V(D)J library preparation, sequencing and mapping
Libraries were constructed using the Chromium Next GEM Single Cell 5' Kit v2 (10x
Genomics). All four feline T cell receptor loci were amplified using the Chromium Single Cell
V(D)J Enrichment Kit according to the manufacturer's instructions but with custom reverse
primers specific for the feline orthologs **<Supplementary File 12>**. Separate libraries were
prepared for the transcriptome, alpha/beta TCRs, and gamma/delta TCRs **<Supplementary File**
12>. Libraries were sequenced using the Illumina NovaSeq S4 platform using 150 paired-end
reads to a target read depth of 30,000 reads per cell. Raw FASTQ sequencing data were
processed by the UC Davis Bioinformatics Core using 10X CellRanger version 7.1 for alignment
to the cat reference genome (*Felis_catus_9.0*). To improve the quality of annotation of the *Felis*
catus genome, we converted unmapped ENSEMBL IDs to human homologs from BioMart for
greater interpretative power. Gene mapping was performed in the following order: (1) Mapping
of one-to-one orthologs from cat to human (2) Mapping of one-to-many and many-to-many was
performed by choosing a representative cat gene based on the following parameters, in respective
order   [orthology_confidence,   gene_order_confidence,   Whole_Genome_alignment,
%_query_identical_target, %target_identical_query]. Additionally, specific unmapped genes
were manually annotated or filtered **<Supplementary File 13>**. Further analysis was then
continued using Seurat v5 (73).

5.3   Count matrix pre-processing and quality control
Filtered feature matrices for the 4 samples were independently normalized and scaled using the
function *SCTransform* v2 (glmGamPoi) (74). Dimensionality reduction (determined using knee
identification in elbow plot), k-means clustering (resolution determined using clustree package)
and neighbor identification (74–76) was performed. Low quality clusters were identified and
filtered based on low gene counts, high mitochondrial read percentage and lack of biologically
relevant markers. To further ensure quality, doublet identification was performed using package
DoubletFinder according to package vignette and ambient RNA decontamination was performed
using the *DecontX* package as per package tutorials adapted to the data (77,78).

5.4   Integration and Downstream analysis

Objects were integrated by sample using the Seurat native reciprocal principal component
analysis (RPCA) (79). The embeddings were utilized for Uniform Manifold Approximation and
Projection (UMAP) and subsequent clustering and neighbor identification as described prior.
Differential gene expression analysis was performed using each cluster identified by the Seurat
function *FindAllMarkers* (Non-parametric Wilcoxon Rank Sum with adjusted P value <0.05).
Gene set enrichment analyses including over-enrichment analysis (ORA) of gene ontology
(biological process) was performed using *ClusterProfiler* on human homologized genes
(adjusted P value <0.05) (80). Additional visualizations and associated statistics were performed
using ggplot2 and gg supplementary packages (81). Figures were illustrated on Bio Render.

5.5 Cell type sub clustering

477 Cell type sub-clustering (example: T cell sub-clustering) analysis was performed in the same
478 manner as described prior. Pseudotime values were calculated using the UMAP embeddings
479 produced via Seurat with RPCA integration using package *monocle3* (82). Gene module scores
480 were calculated via the Seurat function *AddModuleScore*.

5.6 Cross species T cell integrative analysis

Published single cell RNA-sequencing datasets were obtained from NCBI GEO Accession for 4
species including dog (*Canis familiaris*: GSE225599 (18)), human (*Homo sapiens*:
https://support.10xgenomics.com/single-cell-gene-expression/datasets/3.0.0/pbmc_10k_v3),
horse (*Equus caballus*: GSE148416 (22)) and pig (*Sus scrofa*:
https://data.nal.usda.gov/dataset/data-reference-transcriptomics-porcine-peripheral-immune-
cells-created-through-bulk-and-single-cell-rna-sequencing. (21)). All species genes were mapped
to human homologs as described prior for the cat. Only homologized shared genes between all 5
species were utilized for analysis. T cell clusters annotated by authors were validated using
canonical T cell markers (*CD3E*, *CD3D*, *TCF7*, *LCK*, *ITK*). T cells were integrated across
species using Seurat canonical correlation analysis (CCA) and downstream analysis proceeded in
the same manner as described prior for global object analysis.

5.7 VDJ analysis

496 Contigs were mapped to available IMGT reference genes (as of February 2023) using Cell
497 Ranger version 7.1. All analyses were based on the 'all_contig_annotations.csv' file using the R
498 packages tidyverse, circlize and ggseqlogo (83-85). Cluster analysis was performed as previously
499 described (86).

5.8 Data availability

502 All raw and processed data have been submitted to NCBI GEO and are available under
503 GSE267704.

6 Acknowledgements

This project was funded by grant #2022-24-F from the Center for Companion Animal Health,
University of California, Davis. The project was supported by the University of California Davis
Flow Cytometry Shared Resource Laboratory with funding from the NCI P30 CA093373
(Comprehensive Cancer Center) and S10 OD018223, with technical assistance from Bridget
McLaughlin, Jonathan Van Dyke and Ashley Karajeh. The sequencing was carried out by the
DNA Technologies and Expression Analysis Core at the UC Davis Genome Center, supported
by NIH Shared Instrumentation Grant 1S10OD010786-01. We thank Jie (Jessie) Li from the UC
Davis Bioinformatics Core for data analysis and Kristy Harmon from the Leukocyte Biology
Antigen Laboratory for her flow cytometry support.

515
516

7   References

1.   Rowan AN. Companion Animal Statistics in the USA. 2018 [cited 2024 Apr 30];
Available from: https://www.wellbeingintlstudiesrepository.org/demscapop/7/
2.   Larkin M, Radich R. AVMA. Am Vet Med Assoc; 2021 [cited 2024 Apr 5]. Pet
population still on the rise, with fewer pets per household. Available from:
https://avmajournals.avma.org/display/post/news/pet-population-still-on-the-rise--with-
fewer-pets-per-household.xml?print&print&print&print&print
3.   Fromont E, Artois M, Langlais M, Courchamp F, Pontier D. Modelling the feline
leukemia virus (FeLV) in natural populations of cats (Felis catus). Theor Popul Biol
[Internet]. 1997 Aug;52(1):60–70. Available from:
http://dx.doi.org/10.1006/tpbi.1997.1320
4.   Powers JA, Chiu ES, Kraberger SJ, Roelke-Parker M, Lowery I, Erbeck K, et al. Feline
Leukemia Virus (FeLV) Disease Outcomes in a Domestic Cat Breeding Colony:
Relationship to Endogenous FeLV and Other Chronic Viral Infections. J Virol [Internet].
2018 Sep 15;92(18). Available from: http://dx.doi.org/10.1128/JVI.00649-18
5.   Helfer-Hungerbuehler AK, Shah J, Meili T, Boenzli E, Li P, Hofmann-Lehmann R.
Adeno-Associated Vector-Delivered CRISPR/SaCas9 System Reduces Feline Leukemia
Virus Production In Vitro. Viruses [Internet]. 2021 Aug 18;13(8). Available from:
http://dx.doi.org/10.3390/v13081636
6.   Bienzle D. FIV in cats--a useful model of HIV in people? Vet Immunol Immunopathol
[Internet]. 2014 Jun 15;159(3-4):171–9. Available from:
http://dx.doi.org/10.1016/j.vetimm.2014.02.014
7.   Paillot R, Richard S, Bloas F, Piras F, Poulet H, Brunet S, et al. Toward a detailed
characterization of feline immunodeficiency virus-specific T cell immune responses and
mediated immune disorders. Vet Immunol Immunopathol [Internet]. 2005 Jun 15;106(1-
2):1–14. Available from: http://dx.doi.org/10.1016/j.vetimm.2004.12.023
8.   Endo Y, Matsumura S, Washizu T, Ishida T. Alteration of T-cell subsets in the lymph
nodes from cats infected with feline immunodeficiency virus. J Vet Med Sci [Internet].
1997 Sep;59(9):739–46. Available from: http://dx.doi.org/10.1292/jvms.59.739

9.   Hoffmann-Fezer G, Mortelbauer W, Hartmann K, Mysliwietz J, Thefeld S, Beer B, et al.
Comparison of T-cell subpopulations in cats naturally infected with feline leukaemia
virus or feline immunodeficiency virus. Res Vet Sci [Internet]. 1996 Nov;61(3):222–6.
Available from: http://dx.doi.org/10.1016/s0034-5288(96)90067-3
10. Day MJ. Cats are not small dogs: is there an immunological explanation for why cats are
less affected by arthropod-borne disease than dogs? Parasit Vectors [Internet]. 2016 Sep
20;9(1):507. Available from: http://dx.doi.org/10.1186/s13071-016-1798-5
11. Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. Cell
[Internet]. 2006 Feb 24;124(4):783–801. Available from:
http://dx.doi.org/10.1016/j.cell.2006.02.015
12. Jiao A, Zhang C, Wang X, Sun L, Liu H, Su Y, et al. Single-cell sequencing reveals the
evolution of immune molecules across multiple vertebrate species. J Advert Res
[Internet]. 2023 Mar 4; Available from: http://dx.doi.org/10.1016/j.jare.2023.02.017
13. Radtanakatikanon A, Keller SM, Darzentas N, Moore PF, Folch G, Nguefack Ngoune V,
et al. Topology and expressed repertoire of the Felis catus T cell receptor loci. BMC
Genomics [Internet]. 2020 Jan 6;21(1):20. Available from:
http://dx.doi.org/10.1186/s12864-019-6431-5
14. Ginhoux F, Yalin A, Dutertre CA, Amit I. Single-cell immunology: Past, present, and
future. Immunity [Internet]. 2022 Mar 8;55(3):393–404. Available from:
http://dx.doi.org/10.1016/j.immuni.2022.02.006
15. Jovic D, Liang X, Zeng H, Lin L, Xu F, Luo Y. Single-cell RNA sequencing
technologies and applications: A brief overview. Clin Transl Med [Internet]. 2022
Mar;12(3):e694. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1002/ctm2.694
16. Sun X, Lin X, Li Z, Wu H. A comprehensive comparison of supervised and unsupervised
methods for cell type identification in single-cell RNA-seq. Brief Bioinform [Internet].
2022 Mar 10;23(2). Available from: http://dx.doi.org/10.1093/bib/bbab567
17. Wang M, Song WM, Ming C, Wang Q, Zhou X, Xu P, et al. Guidelines for
bioinformatics of single-cell sequencing data analysis in Alzheimer's disease: review,
recommendation, implementation and application. Mol Neurodegener [Internet]. 2022
Mar 2;17(1):17. Available from: http://dx.doi.org/10.1186/s13024-022-00517-z
18. Ammons DT, Harris RA, Hopkins LS, Kurihara J, Weishaar K, Dow S. A single-cell
RNA sequencing atlas of circulating leukocytes from healthy and osteosarcoma affected
dogs. Front Immunol [Internet]. 2023 May 19;14:1162700. Available from:
http://dx.doi.org/10.3389/fimmu.2023.1162700
19. Eschke M, Moore PF, Chang H, Alber G, Keller SM. Canine peripheral blood TCRαβ T
cell atlas: Identification of diverse subsets including CD8A+ MAIT-like cells by
combined single-cell transcriptome and V(D)J repertoire analysis. Front Immunol
[Internet]. 2023 Feb 23;14:1123366. Available from:
http://dx.doi.org/10.3389/fimmu.2023.1123366

20.   Gao Y, Li J, Cai G, Wang Y, Yang W, Li Y, et al. Single-cell transcriptomic and
chromatin accessibility analyses of dairy cattle peripheral blood mononuclear cells and
their responses to lipopolysaccharide. BMC Genomics [Internet]. 2022 Apr 30;23(1):338.
Available from: http://dx.doi.org/10.1186/s12864-022-08562-0
21.   Herrera-Uribe J, Wiarda JE, Sivasankaran SK, Daharsh L, Liu H, Byrne KA, et al.
Reference Transcriptomes of Porcine Peripheral Immune Cells Created Through Bulk
and Single-Cell RNA Sequencing. Front Genet [Internet]. 2021 Jun 23;12:689406.
Available from: http://dx.doi.org/10.3389/fgene.2021.689406
22.   Patel RS, Tomlinson JE, Divers TJ, Van de Walle GR, Rosenberg BR. Single-cell
resolution landscape of equine peripheral blood mononuclear cells reveals diverse cell
types including T-bet+ B cells. BMC Biol [Internet]. 2021 Jan 22;19(1):13. Available
from: http://dx.doi.org/10.1186/s12915-020-00947-5
23.   Li Z, Sun C, Wang F, Wang X, Zhu J, Luo L, et al. Molecular mechanisms governing
circulating immune cell heterogeneity across different species revealed by single-cell
sequencing. Clin Transl Med [Internet]. 2022 Jan;12(1):e689. Available from:
http://dx.doi.org/10.1002/ctm2.689
24.   Touw IP, Palande K, Beekman R. Granulocyte colony-stimulating factor receptor
signaling: implications for G-CSF responses and leukemic progression in severe
congenital neutropenia. Hematol Oncol Clin North Am [Internet]. 2013 Feb;27(1):61–73,
viii. Available from: http://dx.doi.org/10.1016/j.hoc.2012.10.002
25.   Sprenkeler EGG, Zandstra J, van Kleef ND, Goetschalckx I, Verstegen B, Aarts CEM, et
al. S100A8/A9 Is a Marker for the Release of Neutrophil Extracellular Traps and Induces
Neutrophil Activation. Cells [Internet]. 2022 Jan 11;11(2). Available from:
http://dx.doi.org/10.3390/cells11020236
26.   Akgun E, Tuzuner MB, Sahin B, Kilercik M, Kulah C, Cakiroglu HN, et al. Proteins
associated with neutrophil degranulation are upregulated in nasopharyngeal swabs from
SARS-CoV-2 patients. PLoS One [Internet]. 2020 Oct 20;15(10):e0240012. Available
from: http://dx.doi.org/10.1371/journal.pone.0240012
27.   Blanter M, Cambier S, De Bondt M, Vanbrabant L, Pörtner N, Abouelasrar Salama S, et
al. Method Matters: Effect of Purification Technology on Neutrophil Phenotype and
Function. Front Immunol [Internet]. 2022 Feb 10;13:820058. Available from:
http://dx.doi.org/10.3389/fimmu.2022.820058
28.   Buechler C, Ritter M, Orsó E, Langmann T, Klucken J, Schmitz G. Regulation of
scavenger receptor CD163 expression in human monocytes and macrophages by pro- and
antiinflammatory stimuli. J Leukoc Biol [Internet]. 2000 Jan;67(1):97–103. Available
from: https://www.ncbi.nlm.nih.gov/pubmed/10648003
29.   Sampath P, Moideen K, Ranganathan UD, Bethunaickan R. Monocyte Subsets:
Phenotypes and Function in Tuberculosis Infection. Front Immunol [Internet]. 2018 Jul
30;9:1726. Available from: http://dx.doi.org/10.3389/fimmu.2018.01726

30.   Reshetnikova E, Guselnikov S, Volkova O, Baranov K, Taranin A, Mechetina L. B cell-
specific protein FCRLA is expressed by plasmacytoid dendritic cells in humans.
Cytometry B Clin Cytom [Internet]. 2018 Jul;94(4):683–7. Available from:
http://dx.doi.org/10.1002/cyto.b.21611
31.   Rissoan MC, Duhen T, Bridon JM, Bendriss-Vermare N, Péronne C, de Saint Vis B, et
al. Subtractive hybridization reveals the expression of immunoglobulin-like transcript 7,
Eph-B1, granzyme B, and 3 novel transcripts in human plasmacytoid dendritic cells.
Blood [Internet]. 2002 Nov 1;100(9):3295–303. Available from:
http://dx.doi.org/10.1182/blood-2002-02-0638
32.   Edwards AD, Diebold SS, Slack EMC, Tomizawa H, Hemmi H, Kaisho T, et al.
Toll□like receptor expression in murine DC subsets: lack of TLR7 expression by CD8α+
DC correlates with unresponsiveness to imidazoquinolines. Eur J Immunol [Internet].
2003 Apr;33(4):827–33. Available from:
https://onlinelibrary.wiley.com/doi/10.1002/eji.200323797
33.   Medina KL, Tangen SN, Seaburg LM, Thapa P, Gwin KA, Shapiro VS. Separation of
plasmacytoid dendritic cells from B-cell-biased lymphoid progenitor (BLP) and Pre-pro
B cells using PDCA-1. PLoS One [Internet]. 2013 Oct 30;8(10):e78408. Available from:
http://dx.doi.org/10.1371/journal.pone.0078408
34.   Arroyo Hornero R, Idoyaga J. Plasmacytoid dendritic cells: A dendritic cell in disguise.
Mol Immunol [Internet]. 2023 Jul;159:38–45. Available from:
http://dx.doi.org/10.1016/j.molimm.2023.05.007
35.   Atiakshin D, Kostin A, Trotsenko I, Samoilova V, Buchwalow I, Tiemann M.
Carboxypeptidase A3-A Key Component of the Protease Phenotype of Mast Cells. Cells
[Internet]. 2022 Feb 6;11(3). Available from: http://dx.doi.org/10.3390/cells11030570
36.   Cruse G, Kaur D, Leyland M, Bradding P. A novel FcεRIβ-chain truncation regulates
human mast cell proliferation and survival. FASEB J [Internet]. 2010 Oct;24(10):4047–
57. Available from: http://dx.doi.org/10.1096/fj.10-158378
37.   Lee H, Joo JY, Kang J, Yu Y, Kim YH, Park HR. Single-cell analysis of platelets from
patients with periodontitis and diabetes. Res Pract Thromb Haemost [Internet]. 2023
Feb;7(2):100099. Available from: http://dx.doi.org/10.1016/j.rpth.2023.100099
38.   Yang S, Liu F, Wang QJ, Rosenberg SA, Morgan RA. The shedding of CD62L (L-
selectin) regulates the acquisition of lytic activity in human tumor reactive T
lymphocytes. PLoS One [Internet]. 2011 Jul 28;6(7):e22560. Available from:
http://dx.doi.org/10.1371/journal.pone.0022560
39.   Ivetic A, Hoskins Green HL, Hart SJ. L-selectin: A Major Regulator of Leukocyte
Adhesion, Migration and Signaling. Front Immunol [Internet]. 2019 May 14;10:1068.
Available from: http://dx.doi.org/10.3389/fimmu.2019.01068
40.   Bjorkdahl O, Barber KA, Brett SJ, Daly MG, Plumpton C, Elshourbagy NA, et al.
Characterization of CC-chemokine receptor 7 expression on murine T cells in lymphoid

tissues. Immunology [Internet]. 2003 Oct;110(2):170–9. Available from:
http://dx.doi.org/10.1046/j.1365-2567.2003.01727.x

41.   Cano-Gamez E, Soskic B, Roumeliotis TI, So E, Smyth DJ, Baldrighi M, et al. Single-
cell transcriptomics identifies an effectorness gradient shaping the response of CD4+ T
cells to cytokines. Nat Commun [Internet]. 2020 Apr 14;11(1):1801. Available from:
http://dx.doi.org/10.1038/s41467-020-15543-y

42.   Zhu J, Yamane H, Paul WE. Differentiation of Effector CD4 T Cell Populations*. Annu
Rev Immunol [Internet]. 2009 Apr 23 [cited 2024 Mar 23];28(Volume 28, 2009):445–89.
Available from: https://www.annualreviews.org/content/journals/10.1146/annurev-
immunol-030409-101212

43.   Andreatta M, Corria-Osorio J, Müller S, Cubas R, Coukos G, Carmona SJ. Interpretation
of T cell states from single-cell transcriptomics data using reference atlases. Nat
Commun [Internet]. 2021 May 20;12(1):2965. Available from:
http://dx.doi.org/10.1038/s41467-021-23324-4

44.   Wang X, Shen X, Chen S, Liu H, Hong N, Zhong H, et al. Reinvestigation of Classic T
Cell Subsets and Identification of Novel Cell Subpopulations by Single-Cell RNA
Sequencing. J Immunol [Internet]. 2022 Jan 15;208(2):396–406. Available from:
http://dx.doi.org/10.4049/jimmunol.2100581

45.   Vallejo J, Saigusa R, Gulati R, Armstrong Suthahar SS, Suryawanshi V, Alimadadi A, et
al. Combined protein and transcript single-cell RNA sequencing in human peripheral
blood mononuclear cells. BMC Biol [Internet]. 2022 Sep 1;20(1):193. Available from:
http://dx.doi.org/10.1186/s12915-022-01382-4

46.   Schneider L, Marcondes NA, Hax V, da Silva Moreira IF, Ueda CY, Piovesan RR, et al.
Flow cytometry evaluation of CD14/CD16 monocyte subpopulations in systemic
sclerosis patients: a cross sectional controlled study. Adv Rheumatol [Internet]. 2021
May 22;61(1):27. Available from: http://dx.doi.org/10.1186/s42358-021-00182-8

47.   Barut GT, Kreuzer M, Bruggmann R, Summerfield A, Talker SC. Single-cell
transcriptomics reveals striking heterogeneity and functional organization of dendritic
and monocytic cells in the bovine mesenteric lymph node. Front Immunol [Internet].
2022;13:1099357. Available from: http://dx.doi.org/10.3389/fimmu.2022.1099357

48.   Waskow C, Liu K, Darrasse-Jèze G, Guermonprez P, Ginhoux F, Merad M, et al. The
receptor tyrosine kinase Flt3 is required for dendritic cell development in peripheral
lymphoid tissues. Nat Immunol [Internet]. 2008 Jun;9(6):676–83. Available from:
http://dx.doi.org/10.1038/ni.1615

49.   ten Broeke T, Wubbolts R, Stoorvogel W. MHC class II antigen presentation by dendritic
cells regulated through endosomal sorting. Cold Spring Harb Perspect Biol [Internet].
2013 Dec 1;5(12):a016873. Available from:
http://dx.doi.org/10.1101/cshperspect.a016873

50.   Eislmayr K, Bestehorn A, Morelli L, Borroni M, Vande Walle L, Lamkanfi M, et al.
Nonredundancy of IL-1α and IL-1β is defined by distinct regulation of tissues

orchestrating resistance versus tolerance to infection. Sci Adv [Internet]. 2022 Mar
4;8(9):eabj7293. Available from: http://dx.doi.org/10.1126/sciadv.abj7293

51. Lopez-Castejon G, Brough D. Understanding the mechanism of IL-1β secretion.
Cytokine Growth Factor Rev [Internet]. 2011 Aug;22(4):189–95. Available from:
http://dx.doi.org/10.1016/j.cytogfr.2011.10.001

52. Hadadi E, Zhang B, Baidžajevas K, Yusof N, Puan KJ, Ong SM, et al. Differential IL-1β
secretion by monocyte subsets is regulated by Hsp27 through modulating mRNA
stability. Sci Rep [Internet]. 2016 Dec 15;6:39035. Available from:
http://dx.doi.org/10.1038/srep39035

53. Wan J, Shan Y, Fan Y, Fan C, Chen S, Sun J, et al. NF-κB inhibition attenuates LPS-
induced TLR4 activation in monocyte cells. Mol Med Rep [Internet]. 2016
Nov;14(5):4505–10. Available from: http://dx.doi.org/10.3892/mmr.2016.5825

54. Mussbacher M, Derler M, Basílio J, Schmid JA. NF-κB in monocytes and macrophages -
an inflammatory master regulator in multitalented immune cells. Front Immunol
[Internet]. 2023 Feb 23;14:1134661. Available from:
http://dx.doi.org/10.3389/fimmu.2023.1134661

55. Guo H, Jin D, Chen X. Lipocalin 2 is a regulator of macrophage polarization and NF-
κB/STAT3 pathway activation. Mol Endocrinol [Internet]. 2014 Oct;28(10):1616–28.
Available from: http://dx.doi.org/10.1210/me.2014-1092

56. Lee BC, Lee SG, Choo MK, Kim JH, Lee HM, Kim S, et al. Selenoprotein MsrB1
promotes anti-inflammatory cytokine gene expression in macrophages and controls
immune response in vivo. Sci Rep [Internet]. 2017 Jul 11;7(1):5119. Available from:
http://dx.doi.org/10.1038/s41598-017-05230-2

57. Vishnyakova P, Poltavets A, Karpulevich E, Maznina A, Vtorushina V, Mikhaleva L, et
al. The response of two polar monocyte subsets to inflammation. Biomed Pharmacother
[Internet]. 2021 Jul;139:111614. Available from:
http://dx.doi.org/10.1016/j.biopha.2021.111614

58. Zhang Y, Wang S, Xia H, Guo J, He K, Huang C, et al. Identification of Monocytes
Associated with Severe COVID-19 in the PBMCs of Severely Infected Patients Through
Single-Cell Transcriptome Sequencing. Engineering [Internet]. 2022 Oct;17:161–9.
Available from: http://dx.doi.org/10.1016/j.eng.2021.05.009

59. Utrero-Rico A, González-Cuadrado C, Chivite-Lacaba M, Cabrera-Marante O, Laguna-
Goya R, Almendro-Vazquez P, et al. Alterations in Circulating Monocytes Predict
COVID-19 Severity and Include Chromatin Modifications Still Detectable Six Months
after Recovery. Biomedicines [Internet]. 2021 Sep 17;9(9). Available from:
http://dx.doi.org/10.3390/biomedicines9091253

60. Rynikova M, Adamkova P, Hradicka P, Stofilova J, Harvanova D, Matejova J, et al.
Transcriptomic Analysis of Macrophage Polarization Protocols: Vitamin D3 or IL-4 and
IL-13 Do Not Polarize THP-1 Monocytes into Reliable M2 Macrophages. Biomedicines

[Internet]. 2023 Feb 17;11(2). Available from:
http://dx.doi.org/10.3390/biomedicines11020608

61. Wigerblad G, Cao Q, Brooks S, Naz F, Gadkari M, Jiang K, et al. Single-Cell Analysis
Reveals the Range of Transcriptional States of Circulating Human Neutrophils. J
Immunol [Internet]. 2022 Aug 15;209(4):772–82. Available from:
http://dx.doi.org/10.4049/jimmunol.2200154

62. Meng Y, Lin S, Niu K, Ma Z, Lin H, Fan H. Vimentin affects inflammation and
neutrophil recruitment in airway epithelium during Streptococcus suis serotype 2
infection. Vet Res [Internet]. 2023 Jan 30;54(1):7. Available from:
http://dx.doi.org/10.1186/s13567-023-01135-3

63. McColl SR, Paquin R, Ménard C, Beaulieu AD. Human neutrophils produce high levels
of the interleukin 1 receptor antagonist in response to granulocyte/macrophage colony-
stimulating factor and tumor necrosis factor alpha. J Exp Med [Internet]. 1992 Aug
1;176(2):593–8. Available from: http://dx.doi.org/10.1084/jem.176.2.593

64. Kunisada M, Yogianti F, Sakumi K, Ono R, Nakabeppu Y, Nishigori C. Increased
expression of versican in the inflammatory response to UVB- and reactive oxygen
species-induced skin tumorigenesis. Am J Pathol [Internet]. 2011 Dec;179(6):3056–65.
Available from: http://dx.doi.org/10.1016/j.ajpath.2011.08.042

65. Xie X, Shi Q, Wu P, Zhang X, Kambara H, Su J, et al. Single-cell transcriptome profiling
reveals neutrophil heterogeneity in homeostasis and infection. Nat Immunol [Internet].
2020 Sep;21(9):1119–33. Available from: http://dx.doi.org/10.1038/s41590-020-0736-z

66. Glennon-Alty L, Moots RJ, Edwards SW, Wright HL. Type I interferon regulates
cytokine-delayed neutrophil apoptosis, reactive oxygen species production and
chemokine expression. Clin Exp Immunol [Internet]. 2021 Feb;203(2):151–9. Available
from: http://dx.doi.org/10.1111/cei.13525

67. Leylek R, Idoyaga J. Chapter Four - The versatile plasmacytoid dendritic cell: Function,
heterogeneity, and plasticity. In: Lhuillier C, Galluzzi L, editors. International Review of
Cell and Molecular Biology [Internet]. Academic Press; 2019. p. 177–211. Available
from: https://www.sciencedirect.com/science/article/pii/S1937644819300991

68. Varricchi G, de Paulis A, Marone G, Galli SJ. Future Needs in Mast Cell Biology. Int J
Mol Sci [Internet]. 2019 Sep 6;20(18). Available from:
http://dx.doi.org/10.3390/ijms20184397

69. Nascimento C, Ferreira F. Tumor microenvironment of human breast cancer, and feline
mammary carcinoma as a potential study model. Biochim Biophys Acta Rev Cancer
[Internet]. 2021 Aug;1876(1):188587. Available from:
http://dx.doi.org/10.1016/j.bbcan.2021.188587

70. Andrade LM, Sousa ECM, Melo ACS, Pardavil GSS, Silva RA, Valente KF, et al. Gene
expression of cytokines in the blood of domestics cats infected and not infected by Feline
alphaherpesvirus. Available from: http://dx.doi.org/10.4238/gmr19190

71. Smithberg SR, Fogle JE, Mexas AM, Reckling SK, Lankford SM, Tompkins MB, et al.
In vivo depletion of CD4+CD25+ regulatory T cells in cats. J Immunol Methods
[Internet]. 2008 Jan 1;329(1-2):81–91. Available from:
http://dx.doi.org/10.1016/j.jim.2007.09.015
72. Rigamonti A, Castagna A, Viatore M, Colombo FS, Terzoli S, Peano C, et al. Distinct
responses of newly identified monocyte subsets to advanced gastrointestinal cancer and
COVID-19. Front Immunol [Internet]. 2022 Oct 3;13:967737. Available from:
http://dx.doi.org/10.3389/fimmu.2022.967737
73. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell
gene expression data. Nat Biotechnol [Internet]. 2015 May;33(5):495–502. Available
from: http://dx.doi.org/10.1038/nbt.3192
74. Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. Genome Biol [Internet]. 2019 Dec
23;20(1):296. Available from: http://dx.doi.org/10.1186/s13059-019-1874-1
75. Zappia L, Oshlack A. Clustering trees: a visualization for evaluating clusterings at
multiple resolutions. Gigascience [Internet]. 2018 Jul 1;7(7). Available from:
http://dx.doi.org/10.1093/gigascience/giy083
76. Ahlmann-Eltze C, Huber W. glmGamPoi: fitting Gamma-Poisson generalized linear
models on single cell count data. Bioinformatics [Internet]. 2021 Apr 5;36(24):5701–2.
Available from: http://dx.doi.org/10.1093/bioinformatics/btaa1009
77. Yang S, Corbett SE, Koga Y, Wang Z, Johnson WE, Yajima M, et al. Decontamination
of ambient RNA in single-cell RNA-seq with DecontX. Genome Biol [Internet]. 2020
Mar 5;21(1):57. Available from: http://dx.doi.org/10.1186/s13059-020-1950-6
78. McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell
RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst [Internet]. 2019 Apr
24;8(4):329–37.e4. Available from: http://dx.doi.org/10.1016/j.cels.2019.03.003
79. Song Y, Miao Z, Brazma A, Papatheodorou I. Benchmarking strategies for cross-species
integration of single-cell RNA sequencing data. Nat Commun [Internet]. 2023 Oct
14;14(1):6495. Available from: http://dx.doi.org/10.1038/s41467-023-41855-w
80. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal
enrichment tool for interpreting omics data. Innovation (Camb) [Internet]. 2021 Aug
28;2(3):100141. Available from: http://dx.doi.org/10.1016/j.xinn.2021.100141
81. Wickham H. ggplot2 [Internet]. Springer International Publishing; [cited 2023 Oct 16].
16 p. Available from: https://link.springer.com/book/10.1007/978-3-319-24277-4
82. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics
and regulators of cell fate decisions are revealed by pseudotemporal ordering of single
cells. Nat Biotechnol [Internet]. 2014 Apr;32(4):381–6. Available from:
http://dx.doi.org/10.1038/nbt.2859
83. R Core Team (2023). _R: A Language and Environment for Statistical Computing_. R
Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

84.   Gu Z, Gu L, Eils R, Schlesner M, Brors B. circlize Implements and enhances circular
823   visualization in R. Bioinformatics. 2014 Oct;30(19):2811-2. doi:
824   10.1093/bioinformatics/btu393. Epub 2014 Jun 14. PMID: 24930139.
825   85.   Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G,
826   Hayes A, Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K,
827   Ooms J, Robinson D, Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K,
828   Yutani H (2019). "Welcome to the tidyverse." Journal of Open Source Software, 4(43),
829   1686. doi:10.21105/joss.01686.
830   86.   Chang H, Ashlock DA, Graether SP, Keller SM. Anchor Clustering for million-scale
831   immune repertoire sequencing data. BMC Bioinformatics. 2024 Jan 25;25(1):42. doi:
832   10.1186/s12859-024-05659-z. PMID: 38273275; PMCID: PMC10809746.

Figure 1. scRNA-seq atlas of CD5$^+$ enriched circulating feline immune cells revealed 7 major
types across 21 clusters. (A) Table of cell counts from each age group. (B) UMAP plot
demonstrating the unsupervised clustering results for feline circulating immune cells from 4
healthy cats of different age groups. (C) UMAP plot of global clustering colored by age. (D)
Table of cell type counts across clusters by age. (E) Dot plot demonstrating cell type specific
marker expression of the 7 major cell types. (F-O) Feature UMAP plots demonstrating
expression profile of key markers for each of the 7 different cell types.

A

UMAP_2

γδ T

CD4⁺ T_EM

CD8⁺ Naive

CD8⁺ Cytotoxic

CD4⁺ Naive

UMAP_1

B

Pseudotime

C

LAG3
PDCD1
TIGIT
CCL5
NKG7
GZMK
GZMB
GZMA
PRF1
FGL2
ITGB1
VIM
S100A4
LGALS3
LGALS1
ANXA1
ANXA2
IL7R
CCR7
LEF1
SELL
CD8B
CD8A
CD4
TCF7
ITK
LCK
CD3E
CD3D
CD5

CD4+ Naive | CD8+ Naive | CD4+ TEM | gdT | CD8+ Cytotoxic

% Exp
· 0
· 25
· 50
· 75
· 100

Avg Exp
2
1
0
-1
-2

D

	6MO	1YO	4YO	9YO	TOTAL
0	1515	2400	1226	1573	6714
1	1124	1912	1015	1109	5160
2	836	1613	855	728	4032
3	719	727	621	960	3027
4	425	278	301	358	1362
5	111	305	52	63	531
6	178	14	14	20	226

E

TRG | TRDC | OVERLAY

TRCC

TRG

F CD4   2.0 1.5 1.0 0.5 0.0

G CD8A   3 2 1 0

H CD8B   2 1 0

I SELL   3 2 1 0

J CCR7   2.0 1.5 1.0 0.5 0.0

K ANXA2   2.5 2.0 1.5 1.0 0.5 0.0

L *LGALS3*   2 1 0

M *GZMK*   3 2 1 0

N *PRF1*   1.5 1.0 0.5 0.0

O *CCL5*   5 4 3 2 1 0

P TIGIT   2.0 1.5 1.0 0.5 0.0

Q PDCD1   2.0 1.5 1.0 0.5 0.0

Figure 2. CD5$^+$ enriched T cells segregate into conventional T cell subtypes and effectors. Unsupervised clustering of T cells reveals 7 subtypes. (A) UMAP of the scRNA-seq atlas of T cells. (B) UMAP of T cells colored by Pseudotime. (C) Dot plot of marker genes expressed by T cell type. (D) Table of cell type frequencies across ages by cluster. (E) Feature plots of expression and co-expression of *TRG* and *TRDC*. (F-Q) UMAP of T cells colored by classical T marker genes defining *CD4*/*CD8* status, naive (*SELL*, *CCR7*), effectorness (*ANXA2*, *LGALS3*, *GZMK, PRF1*), terminal differentiation (CCL5) and T exhaustion (*TIGIT*, *PDCD1*).

Figure 3. Feline Effector T cells (T_{EM}) segregate into transcriptionally similar clusters and
reveal the presence of helper T phenotypes. Unsupervised clustering of T_{EM} reveals 13
subtypes. (A) UMAP of scRNA-seq atlas of T_{EM}. (B) Dot plot demonstrating up to 3 top
differentially expressed genes for each cluster determined via Wilcoxon rank sum testing
(*Adj P <0.05*). (C,E,G,I) UMAP of T_{EM} colored by helper T subtype gene modules. (D,F,H,J)
Expression UMAP of a representative gene from each helper T gene module presented in
parallel.

Figure 4. Cross-species integrative analysis of T cells reveals missing cytotoxic effectors in
the cat. Unsupervised clustering revealed 12 clusters across 5 species. (A) UMAP of scRNA-
seq atlas of T cells. (B-F) UMAP split by species- dog, horse, cat, human and pig. (G)
UMAP of T cells colored by T cell phenotype. (H) Dot plot of marker gene sets for T cell
subtypes. (I) Percentage bar chart of clusters stacked by species. (J) UMAP colored by *CD5*
expression. (K) Bar chart of percentage *CD5*$^+$ cells per cluster. (L) Bar chart of average *CD5*
expression across cells in each cluster. (M) Scatter plot of percentage *CD5*$^+$ cells per cluster
versus number of cat cells within the corresponding cluster.

Figure 5. (A) Percentage of T cells expressing a certain combination of productive TR chain transcripts. TRA/TRB transcripts dominate in all ab T cell subsets except for CD8$^+$ cytotoxic T cells, which primarily express TRA/TRB/TRG transcripts. (B) TRA & TRB V gene usage of the 4 major ab T cell subsets. CD8$^+$ cytotoxic T cells show differential usage compared to the other T cell subset but are also substantiated by fewer cells. (C) Junctional length of TRA rearrangements of the 4 major ab T cell subsets. Junctional regions with dominant TRAV genes in CD8$^+$ cytotoxic T cells are 13 and 14 amino acids long. (D) TRG transcripts in CD8$^+$ cytotoxic T cells have a skewed junctional length with 16 amino acid length (top) and a conserved motif (bottom) (E) The majority of TRG rearrangements in CD8$^+$ cytotoxic T cells involved cassette 2 (TRGV2-2/TRGJ2-2). Genes are displayed in order of genomic location and colored by cassette.

866 **Figure 6.** (A) Shared clonotypes across 4 cats. (B) Characterization of clusters based on
867 mean centrality and cluster density. (C) Representative examples of TRA clusters with high
868 centrality and density.

Figure 7. Feline circulating monocytes include classical monocytes with more differentiated clusters with traditional proinflammatory and unconventional anti-inflammatory phenotypes. Unsupervised clustering revealed 5 clusters. (A) UMAP of scRNA-seq atlas of monocytes. (B-F) UMAP of monocytes colored by expression levels of canonical markers. (G) Dot plot of genes of monocytic phenotypes across the 4 monocytic clusters. (H-J) Dot plots of top GO biological process terms called based on sets of positive differentially expressed genes identified via Seurat *FindMarker* function (Wilcoxon rank sum, *Adj P <0.05*). Classical monocyte, CM; Conventional dendritic cell, cDC; Pro-inf, proinflammatory; Anti-inf, anti-inflammatory.

Figure 8. Feline circulating neutrophils separate into 2 clusters based on activation state. (A)
UMAP of scRNA-seq atlas of neutrophils. (B-G) UMAP of neutrophils colored by expression
levels of canonical and functional neutrophil markers. (H) Dot plots of top GO biological process
terms in cluster 1 called based on sets of positive differentially expressed genes identified via
Seurat *FindMarker* function (Wilcoxon rank sum, Adj *P <0.05*). (I-K) UMAP of neutrophils
colored by expression levels of select differentially upregulated genes in cluster 1. (L,M) Violin
plots of RNA counts and feature (unique RNA) counts per cell by cluster; (****) indicates P
value less than 0.001 by T-test. (N) UMAP of neutrophils colored by interferon (IFN) gene
composite score per cell; genes included are named in figure.