Single cell RNA-sequencing of feline peripheral immune cells with V(D)J repertoire and cross species analysis of T lymphocytes

Raneesh Ramarapu[1,2], Judit M. Wulcan[3], Haiyang Chang[4], Peter F. Moore[3], William Vernau[3], Stefan M. Keller[3]

[1]Department of Surgical and Radiological Sciences, School of Veterinary Medicine, University of California Davis, Davis, CA, USA.

[2]Department of Anatomy, Physiology and Cell Biology, School of Veterinary Medicine, University of California Davis, Davis, CA, USA.

[3]Department of Pathology, Microbiology and Immunology, School of Veterinary Medicine, University of California, Davis, CA, United States.

[4]Department of Mathematics and Statistics, University of Guelph, Guelph, ON, Canada.

1   Abstract

Introduction. The domestic cat (Felis catus) is a valued companion animal and a model for virally induced cancers and immunodeficiencies. However, species-specific limitations such as a scarcity of immune cell markers constrain our ability to resolve immune cell subsets at sufficient detail. The goal of this study was to characterize circulating feline T cells and other leukocytes based on their transcriptomic landscape and T-cell receptor repertoire using single cell RNA-sequencing.

Methods. Peripheral blood from 4 healthy cats was enriched for T cells by flow cytometry cell sorting using a mouse anti-feline CD5 monoclonal antibody. Libraries for whole transcriptome, alpha/beta T cell receptor transcripts and gamma/delta T cell receptor transcripts were constructed using the 10x Genomics Chromium Next GEM Single Cell 5' reagent kit and the Chromium Single Cell V(D)J Enrichment Kit with custom reverse primers for the feline orthologs.

Results. Unsupervised clustering of whole transcriptome data revealed 7 major cell populations - T cells, neutrophils, monocytic cells, B cells, plasmacytoid dendritic cells, mast cells and platelets. Sub cluster analysis of T cells resolved naive (CD4+ and CD8+), CD4+ effector T cells, CD8+ cytotoxic T cells and gamma/delta T cells. Cross species analysis revealed a high conservation of T cell subsets along an effector gradient with equitable representation of veterinary species (horse, dog, pig) and humans with the cat. Our V(D)J repertoire analysis demonstrated a skewed T-cell receptor alpha gene usage and a restricted T-cell receptor gamma junctional length in CD8+ cytotoxic T cells compared to other alpha/beta T cell subsets. Among myeloid cells, we resolved three clusters of classical monocytes with polarization into pro- and anti-inflammatory phenotypes

in addition to a cluster of conventional dendritic cells. Lastly, our neutrophil sub clustering
revealed a larger mature neutrophil cluster and a smaller exhausted/activated cluster.

Discussion. Our study is the first to characterize subsets of circulating T cells utilizing an
integrative approach of single cell RNA-sequencing, V(D)J repertoire analysis and cross species
analysis. In addition, we characterize the transcriptome of several myeloid cell subsets and
demonstrate immune cell relatedness across different species.

Keywords: feline, T cells, single cell RNA-sequencing (scRNA-seq), T-cell receptor repertoire,
cross species analysis, myeloid cells, V(D)J

2   Introduction
The domestic cat, Felis catus, is a valued companion animal and the second most popular pet in
the United States. As of the American Veterinary Medicine Association (AVMA) pet ownership
statistics, there are upwards of 46.5 million households with at least one cat in 2023 with ownership
trends rising in the last decade from 25% in 2016 to 29% in 2022 (1,2). In addition to the growing
need for feline veterinary care, the cat is an important model for infectious diseases such as virally-
induced cancers (Feline leukemia virus, FeLV) and virally mediated immunodeficiency (Feline
immunodeficiency virus, FIV) (3–6). Despite the value of the cat as a companion animal and
infectious disease model, there are species specific research limitations including few cat-reactive
immunophenotyping reagents and a poor knowledge base of immune cell markers and behavior.

In the advent of deciphering the feline immune system, many studies characterizing feline immune
cells have relied strongly on antibody-based assays including flow cytometry and
immunohistochemistry. This has allowed the study of immune cells in health and disease
conditions such as FeLV or FIV (7–9). This approach depends on the assumption that feline
immune cells are similar to other species. However, feline immune mediated diseases behave very
differently from those in other small animals such as dogs, exhibiting unique etiologies and
pathogenesis (10). In a cross-species context, innate immunity shows relatively high evolutionary
conservation; however, this conservation dwindles with progression to adaptive immunity and
complex immune phenotypes (11). Additionally, there is growing evidence demonstrating that
evolutionary relationships do not translate to immune transcriptional or cell type relationships.
Such an example is the mouse being a better model for human immune cells than the macaque
(12). Thus, understanding the species-specific system in the context of our growing evolutionary
database is desirable and there is a strong need to characterize the heterogeneous peripheral
immune cell population in the cat, especially diverse populations such as T cells.

In efforts to investigate feline specific T cell diseases, our laboratory has contributed to the growing
feline specific reagent and immune cell knowledge base including T cell receptor (TCR) repertoire
analysis (13). In this trajectory, single cell RNA-sequencing is a logical next step. It enables the
efficient large-scale capture and analysis of heterogeneous tissues, overcoming the challenges of
species-specific reagent assays (14–17). Consequently, there has been a bloom in single cell RNA-
sequencing studies of the peripheral leukocytes in a variety of mammalian species including the
dog, cow, horse, and pig (18–22). Additionally, single cell data facilitates the study of cross species
variations in immune cell type and evolution of these cells (12). ScRNA-seq of feline circulating
leukocytes have been previously performed demonstrating the presence of 5 major cell types (T
cells, B cells, NK cells, monocytes, dendritic cells) in the context of other non-model species (23).
However, in-depth characterization of key subtypes and their marker conservation is yet to be
determined. Thus, the goal of this study is to characterize the heterogeneity of circulating feline T
cells and other captured leukocytes utilizing CD5 flow cytometry enriched scRNA-seq and V(D)J
repertoire analysis in clinically healthy domestic shorthair cats between the ages of 6 months and
9 years. CD5 was chosen as a selective T-cell marker given the lack of availability of anti-feline
CD3 antibodies. Additionally, we performed a cross species transcriptomic integration to assess
feline T cells in the context of 4 other mammalian species - dog, horse, human, and pig. Our results
provide a foundation for feline immune transcriptomics.
90
3   Results
3.1   Single cell atlas of the feline circulating immune cells
ScRNA-seq data obtained from 4 representative healthy cats of different ages consisted of 30,073
quality cells. These age groups were selected as they capture the major feline life stages: 6-month-
old juvenile (6MO), 1-year-old mature (1YO), 4-year-old adult (4YO), and 9-year-old aged
(9YO). The average number of cells captured per cat was 7,518 with an average of 3,655
transcripts per cell. **<Fig1A, Supplementary File1 S1A,B>**. Due to the poor annotation of the cat
reference genome, we utilized a custom homologous mapping script based on ENSEMBL
homologous mapping between the cat and human to improve gene readability. Our script increased
the number of gene symbols mapped from 13,886 to 16,936, improving the quality of cell type
assignment and biological interpretation. Following batch correction via reciprocal principal
component analysis, unsupervised clustering revealed the presence of 21 clusters **<Fig 1B>**. A
majority of cell types are equally represented from each sample **<Fig1C, 1D>**. A quality control
violin plot for clusters can be found in the supplementary materials **<Supplementary File1 S2A>**.
105
These clusters were categorized into 7 major immune cell types based on canonical marker genes
from multi-species PBMC single cell literature, top marker genes and over-enrichment gene
ontology (GO) **<Supplementary File 2,3>**. The most numerous cell type identified was T cells
called by high expression of T cell receptor complex genes (*CD3D, CD3E*) and species conserved
T cell markers *LCK*, *ITK* and *TCF7* (12,23) **<Fig 1E-G, Supplementary File1 S2B>**. Across most
T cell clusters, top identified GO terms of biological processes (BP) included T leukocyte

processes of differentiation, migration, activation, and effector functions **<Supplementary File**
3>.

The second most abundant population was the neutrophils marked by neutrophil-derived proteases
including elastase (*ELANE*), granulocyte colony stimulating factor receptor (*CSF3R*) and
calprotectin (*S100A8/S100A9*) (24–26) **<Fig 1E,H>**. In most species, neutrophils are not captured
in the PBMC layer during density centrifugation (27). Interestingly, our single cell data set
captured a significant number of neutrophils from all but the 1YO sample **<Fig 1D>**. This is the
first description of feline neutrophils at the single cell transcriptomic level and can provide insights
into biological processes of these abundant cells.

A substantial number of monocytic cells were captured with marker genes including scavenger
receptor for haptoglobin-hemoglobin complexes *CD163*, antigen presenting cell co-stimulatory
molecule *CD83*, lipopolysaccharide detector protein *CD14*, phospholipase B domain containing 1
(*PLBD1*) and colony stimulating factor receptor (*CSF1R*) (28,29) **<Fig 1E,I,J>**. A relatively small
but heterogeneous population of B cells were captured which were identified by conserved markers
including B cell surface molecule *MS4A1* and Ig-alpha protein of the B-cell antigen component
CD79A (23) **<Fig 1E,K,L>**. This population also expresses immunoglobulin genes (*IGHG3,*
IGHM) indicating a sub-population of plasma cells (12) **<Fig 1E>**.

Unexpectedly, we captured plasmacytoid dendritic cells (pDC) and mast cells, which represented
rare but discrete cell populations. pDCs were identified by expression of Fc fragment of gamma
immunoglobulin (*FCRLA*), dendritic marker *PLAC8B*, and immunoglobulin lambda constant 7
(*IGLC7*) and lack of expression of B cell canonical markers *MS4A1* and *CD79A* (30–32) **<Fig**
1E,L,M>. pDcs clustered more closely with B cells than other myeloid cells. This is due to close
lineage associations and the shared expression of integral markers *FCRLA* and *IGLC7* among
others (33) **<Fig 1E,L>**. Currently, there is debate regarding the reclassification of pDCs as innate
lymphocytes and this analysis further supports such an argument (34) **<Fig1B>**. Mast cells were
marked by IgE receptor *MS4A2* and a key component of mast cell protease, *CPA3* (35,36) **<Fig**
1E,N>.

Lastly, we captured a population of non-leukocytic immune cells which were annotated as platelets
due to high expression of platelet specific proteins *PPBP*, *VWF* and *GP9* (37) **<Fig1E,O>**. This
cluster also exhibited an extremely low average count of unique RNA and total RNA, which is
consistent with a highly specialized cell type lacking a nucleus **<Supplementary File1 S2A>**.

3.2　Feline circulating T cells
To further characterize T cell subsets, we independently analyzed T cell clusters previously
described at the global level (clusters 0,2,9,1,6,4,5,14,8,12, and 15). Unsupervised clustering of T
cells revealed the presence of seven T cell clusters, which were unrecognizable at the global level

<Fig 2A>. Largely, these clusters represent greater T cell phenotypes along increasing pseudotime
values (proxy of differentiation or lineage) **<Fig 2B,C>**. The subtle transcriptional differences, as
noted through overlapping differentially expressed genes, make effective annotation difficult
<Supplementary File 4>. Nonetheless, we were able to identify several canonical subsets.

T cell subsets included $CD4^+$, $CD8A^+/CD8B^+$ and γδ T cells and were equally represented in all
age groups **<Fig2D-H>**. A great majority were identified as naive cells, distinguished by their high
expression of $CCR7$ and L-selectin ($SELL$) (38–40) **<Fig2I,J>**. These naive populations
segregated into two larger groups which could be distinguished by differential expression of $CD4$
and $CD8A/CD8B$ **<Fig2F-H>**. The $CD8A^+/CD8B^+$ naive population represents a divergence from
the traditionally observed distribution of naive T cells. Most naive T cells in mammalian species
are $CD4^+$ (12). However, a large population of $CD8A^+/CD8B^+$ naive cells has been described in
non-model species such as the horse (22). This highlights a potentially influential role of
$CD8A^+/CD8B^+$ T cells in cats.

The clusters 3 and 4, adjacent to the large naive populations, were identified as effector cells (T_{EM})
<Fig2A>. These clusters show a gradual upregulation of T effector molecules including $S100A11,$
galectins ($LGALS1, LGALS3$), annexins ($ANXA1, ANXA2$), and granzymes ($GZMA, GZMK$)
(41,42) **<Fig2C,K-M>**. Szabo et al identified $CCL5$ as a critical marker for highly differentiated
effector T cells across tissues as well as its antagonistic expression to the naive marker $SELL$ (41).
In our data, Cluster 6 expressed the highest levels of cytotoxic markers including $CCL5$ and
exhibited the highest pseudotime score with no naive marker expression **<Fig2C,M-O>**. This
indicates that this cluster is most consistent with the annotation of a terminally differentiated T cell
cluster ($CD8^+$ cytotoxic). Additionally, some effector cells expressed genes associated with T cell
exhaustion such as $PDCD1$ and $TIGIT$ indicating the presence of a few of these cells in circulation
<Fig2C,P,Q>.

Cluster 5 represented γδ T cells as inferred by the overlapping expression of the TCR genes TRG
and $TRDC$ (13) **<Fig2E>**. Within this population, there was an expression gradient of naive as
well as effector markers, indicating subtypes within the γδ population **<Fig 2I,K,L>**. Although
these T cell clusters represent biologically relevant groups statistically, the literature across various
species identifies significant challenges associated with clustering of transcriptionally similar but
heterogenous T cell populations (43). The γδ cluster exemplifies this challenge in that the cluster's
defining features overlap between two different but not mutually exclusive phenotypes - γδ
recombination and effectorness/naiveness. These cells exhibit overlapping expressions of TRG and
$TRDC$, indicating their TCR recombination-based categorization. An overwhelming majority of
these γδ T cells are naive given their expression of naive marker $SELL$ **<Fig 2I,J>**. However, some
cells appear to express features of effectorness **<Fig 2K-N>**. This suggests some limitations of
transcriptionally driven phenotyping of T cells. Nonetheless, this investigation represents a
significant contribution towards deciphering the biology of feline T cells.

3.3 Feline circulating T effector cells

As described previously, subtyping T cells with overlapping phenotypes represents a significant challenge, which was further exacerbated by the low number of effector T cells in our dataset. Hence, we clustered effector T cells (clusters 3,4) independently, revealing 13 effector clusters **<Fig3A>**. Many of these clusters represent minimally divergent biological states as reflected by the marked overlap of top differentially expressed genes **<Fig3B, Supplementary File 5>**. However, direct visualization of known phenotypes demonstrate that the effector populations captured exhibit phenotypes of human T_H1, T_H2, T_H17 and T_{reg} cells **<Fig3C-J>** (41,42,44). Gene sets for each module calculation are provided in the supplementary materials **<Supplementary file 1, S3>**. These phenotypes do overlap and are not consistent with identified clusters. Nevertheless, the higher scores among closely positioned cells likely indicates the presence of these subtypes. However, a greater sample size of effector cells would be necessary to better resolve subtypes and their trajectory.

3.4 Cross species analysis of circulating T cells

To understand how the captured feline T cell populations compared to those of other domestic animal species and humans, we performed an integrative cross-species analysis. We selected 4 species for which peer-reviewed annotated scRNA-seq data with over 15,000 cells was available. ScRNA-seq data from three veterinary species (dog, horse and pig) and humans were chosen as the reference for homologous genome mapping (18,21,22,45). In total, we integrated 95,366 cells and identified 13 clusters across the 5 species **<Fig4A-F>**. These clusters were functionally annotated based on markers for T cell identity, naive phenotype, effector phenotype and cytotoxic phenotype in addition to their differential gene expression analysis **<Fig4G,H, Supplementary File 6>**.

Although feline cells were overrepresented in some naive clusters such as 3 and 8, there was limited representation in clusters 5, 6, and 9 particularly **<Fig4I>**. To investigate whether this finding was associated with our enrichment protocol, we assessed *CD5* transcript expression across all clusters **<Fig4J>**. Our results suggest that the number of $CD5^+$ cells and average *CD5* expression were lowest in clusters 5 and 9 **<Fig4K,L>**. Additionally, we saw a direct relationship between the number of feline T cells and the $CD5^+$ cell count in each cluster. Particularly, clusters 5, 6, 9, 11, and 12 were most underrepresented **<Fig4M>**. These clusters were identified as cytotoxic effector cells **<Fig4H>**. This highlights a limitation of our CD5 enrichment protocol as it may select against specific effector clusters such as $CD8^+$ cytotoxic cells. Much of our current understanding of $CD8^+$ feline T cells is from flow cytometric studies with no further subtyping due to lack of reagents. Thus, further investigation is necessary to validate whether this low frequency is biological or technical.

3.5 V(D)J recombination

To identify patterns in T cell receptor (TR) expression, we sequenced TR alpha (TRA), TR beta (TRB), TR gamma (TRG) and TR delta (TRD) transcripts simultaneously. The majority (>87.0%) of $CD4^+$ naïve, $CD4^+$ T_{EM} and $CD8^+$ naïve T cells expressed TRA/TRB transcripts only **<Fig5A>**. In contrast, only 35.0% of $CD8^+$ cytotoxic T cells expressed TRA/TRB transcripts while 60.6% expressed TRA/TRB/TRG transcripts. γδ T cells primarily expressed TRG/TRD transcripts (49.7%) followed by TRG/TRD/TRB (16.4%) transcripts. Of note, 11.5% of cells had TRA/TRB/TRG transcripts, which might represent αβ T cells with an effector phenotype similar to gd T cells. $CD8^+$ Cytotoxic T cells also exhibited differential TRAV gene usage, and to a lesser degree, TRBV gene usage **<Fig5B,C>**. The junctional length of TRG transcript in $CD8^+$ Cytotoxic T cells was highly skewed towards 16 amino acids and converged on the conserved motif 'CAAWDPRGYGWAHKVF' **<Fig5D>**. Most TRG rearrangements involved cassette 2 (TRGV2-2/TRGJ2-2) **<Fig5E>.** Interestingly, a subset of TRGV2-2 genes rearranged to the TRGJ2-1 gene, which is located upstream of TRGV2-2.

Next, we characterized the repertoire overlap between cats and the relatedness of junctional regions across cats and T cell subsets **<Fig6>**. The number of shared clonotypes was highest for TRA and TRG but overall low **<Fig6A>**. When expanding the definition of shared repertoires to include clonotypes with one amino acid difference, the TRA locus had the highest number, size and publicity of clusters **<Fig6B,C>**.

3.6 Feline circulating monocytic cells

Although $CD5^+$ enrichment was performed for T cell selectivity, we serendipitously captured neutrophils and monocytic cells, which provides insight into the transcriptome of circulating myeloid cells in cats. Clustering analysis of both cell types was performed independently. Due to the low myeloid cell count in the 1YO, it was excluded from this analysis.

Monocytic cells clustered into 5 groups and were largely represented by the 9YO sample **<Fig 7A, Supplementary File 1, S4A>**. Cluster 3 represented neutrophils which is likely due to the close clustering of myeloid cells on the global object **<Fig 1B, Supplementary File 1, S4B>**. Cluster 0, 1 and 2 were identified as classical monocytes (CM) with all clusters having high expression of markers *PLBD1*, *CD83* and classical monocyte markers *CD14* and *VCAN* (28,29,46) **<Fig 7B-D>**. We did not identify any non-classical monocytes, based on absence of *CD16* expression, as seen in other species such as humans, horses and cattle (22,29,46,47). Cluster 4 showed an upregulation of receptor-type tyrosine-protein kinase *FLT3*, a conventional pan-dendritic cell marker (48) **<Fig 7F>**. In addition, these cells exhibited upregulation of MHC class-II molecules (*FECA-DRB*, *HLA-DRA*, *HLA-DOA*, and *HLA-DMD*) and were hence identified as conventional dendritic cells (cDC) (49) **<Fig 7G>**.

In-depth assessment of differentially expressed genes across the four monocytic clusters identified
signatures that further characterized these cell types. Cluster 2 demonstrated a proinflammatory
phenotype characterized by upregulation of classic proinflammatory cytokines (*IL1A*, *IL1B*),
TNF/NF-KB pathway (*TNF*, *TNFAIP3, NFKBIA, NFKBIZ, NFKB1*) and M1-macrophage-
associated proinflammatory cytokines (*CXCL10, CXCL16, CCL3, CCL4 and CCL5*) **<Fig 7G,H,**
Supplementary File 7,8> (20,50–54). Cluster 1 gene ontology revealed an upregulation of
leukocyte migration and chemotaxis terms in addition to upregulation of proinflammatory
cytokines *IL1A* and *IL1B* **<Fig 7G,I, Supplementary File 7,8>**. However, these cells also lack
TNF expression and upregulate anti-inflammatory cytokines *IL10* and *IL18* and proteins which
promote anti-inflammatory phenotypes in macrophages such as *LCN2* and *MSRB1* (55,56) **<Fig**
7G>. They have the lowest expression of *CD86*, a T cell co-stimulatory molecule of antigen
presenting cells, which is associated with an anti-inflammatory phenotype (57). Most strikingly,
these cells were found to upregulate genes similar to a novel monocyte cluster found in SARS-
CoV-2 infected and recovered patients. These genes include *AREG*, *EREG*, *BCL6*, *IL10,* and *IL18*
which have been described as anti-inflammatory tissue repair genes **<Fig 7G>** (58,59). Thus, these
cells were designated as anti-inflammatory, although they do not follow the traditional M2
polarization phenotype described in human monocytes (60). Our data suggests a novel classical
monocyte subtype with an anti-inflammatory phenotype.
288
Cluster 0 (CM) appears the least differentiated with low expression of inflammatory cytokines and
the top GO terms include more cellular rather than immune related functions **<Supplementary**
File 8>. Cluster 4 (cDC) showed an upregulation of and the GO terms associated with leukocyte
activation and T cell activation via TCR contact with antigen bound MHC **<Fig 7G,J,**
Supplementary File 7,8>. Given the strong antigen presentation phenotype, the annotation of
cDC was further corroborated.
295
3.7   Feline circulating neutrophils
We identified 2 major neutrophil clusters **<Fig 8A>** based on the expression of canonical
neutrophil markers including *CSF3R* and *ELANE* as well as mature neutrophil functional markers
vimentin (*VIM*), thioredoxin (*TXN)*, *S100A9* and *S100A12*, suggesting both clusters are comprised
of mature neutrophils (25,61,62) **<Fig 8B-G>**. Cluster 1 exhibited several notable differences
compared to cluster 0. Differential gene expression analysis accompanied by gene ontology
showed an upregulation in cluster 1 of apoptotic signaling and activated neutrophil functions such
as response to lipopolysaccharide and *IL-2* production **<Fig 8H, Supplementary File 9,10>**. Thus,
we annotated this cluster as activated neutrophils. When investigating top differentially expressed
genes, we noted a higher expression of *IL1A* and *TNFAIP3*, suggesting a proinflammatory
phenotype (63) **<Fig 8I,J>**. Similarly, cluster 1 had an increased expression of *VCAN*, which is
upregulated in the skin in response to damage from ultraviolet light B and reactive oxygen species
(64). Lastly, cluster 1 exhibited lower RNA counts per cell, which is consistent with findings in
murine neutrophils where the number of genes and counts of RNA decrease with neutrophil

310 differentiation, maturation and eventual activation and function (65). Since a pathway driving this
311 clustering could not be resolved, we examined an interferon (IFN) module score, which is a
312 composite score based on expression levels of interferon stimulated genes **\<Fig 8N\>**. Cluster 1 did
313 not show an increased IFN score but the adjacent cells in cluster 0 did **\<Fig 8N\>**. Interferon
314 stimulation can delay neutrophil apoptosis, which could represent one of the potential driving
315 factors for cells to move from cluster 0 to cluster 1 (66). This analysis is the first to resolve
316 circulating neutrophil transcriptomic signatures in cats at the single cell level.
317
4 Discussion

318 **4** **Discussion**
319 The importance of the cat as a companion animal as well as a model for infectious diseases
320 warrants an in-depth characterization and understanding of the feline immune system. Our study
321 is one of the first to characterize feline immune cell subpopulations at the single cell level. We
322 utilized 5' single cell RNA-sequencing with V(D)J analysis to resolve the heterogeneity of CD5$^+$
323 enriched peripheral blood immune cells. We resolved populations of T cells, neutrophils, B cells
324 and monocytes. Additionally, this atlas is the first to annotate the single cell transcriptome of rarer
325 cell types, plasmacytoid dendritic cells and mast cells, which are seldomly captured in single cell
326 atlases (67,68).
327
328 Among T cells, we were able to resolve different populations of naive cells including a significant
329 number of *CD8A/CD8B* naive cells. We further identified T effector cells and terminally
330 differentiated cytotoxic effectors. The transcriptomic gradient of T cell differentiation was found
331 to fit the paradigm seen with other species (23). Additionally, we captured feline γδ T cells for the
332 first time. Upon investigation of effector cells, we were able to demonstrate known T helper
333 effector phenotypes among the most differentiated effectors. Of note, the T$_{reg}$ phenotype was found
334 to be the most prominent. The importance of T$_{reg}$ populations has been studied in cats, particularly
335 in the context of feline mammary carcinoma (69–71). Our study corroborates the large presence
336 of this phenotype and the potential importance of T$_{reg}$ cells in feline immunology.
337
338 We performed a cross species integrative analysis of T cells to take advantage of the growing
339 database of single cell RNA data. We found relatively high conservation of T cell subtypes along
340 an effector gradient with relatively equitable representation of veterinary species (horse, dog, pig)
341 and humans with the cat. This analysis also revealed the potential negative impacts of CD5$^+$
342 enrichment for T cells. Effector clusters with high *CD5* expression were found to be
343 underrepresented in the cat. Thus, a broader T cell selection protocol would be necessary in future
344 studies to further investigate effector cells in cats.
345 In addition to characterizing T cells based on their transcriptome, we analyzed TCR expression on
346 a single cell level for the first time in the cat. Cytotoxic CD8$^+$ T cells differed from other αβ T cell
347 subsets in multiple aspects. In addition to the expected TRA and TRB transcripts, a high percentage
348 of cytotoxic CD8$^+$ T cells expressed a productive TRG transcript. Given that this was not observed
349 in naïve CD8+ T cells, we attribute this phenomenon to differential transcriptional regulation of

naive and effector T cells. Compared to other αβ T cell subsets, cytotoxic T cells also exhibited
divergent TRAV and TRBV gene usage but not TRGV gene usage. Instead, the junctional length
of TRG transcripts was skewed towards a highly conserved16 aa motif. This finding is of potential
significance for clonality testing, an adjunct method for differentiating reactive from neoplastic
lymphoid proliferations. If the majority of cytotoxic CD8[+] T cells harbor a length restricted TRG
rearrangement, then reactive CD8[+] dominant T cell proliferations might exhibit reduced TRG
diversity, which could result in a higher false positive rate. Of note, since other αβ T cell subsets
did not express a significant number of productive TRG transcripts, it is unknown if similar length
restricted rearrangements occur in these subsets. A previous study examining the feline TCR
transcriptome across thymus, spleen and mesenteric lymph nodes did not identify any TRG length
restrictions (13). However, the previous study aimed to identify global rearrangement patterns and
hence utilized a bulk sequencing strategy, which is not suited to resolve repertoire characteristics
of individual T cell subsets.
363
A cluster analysis of TCR sequences did not identify T cell subsets with invariant TRA or TRB
chains. A previous study utilizing scRNA-seq with VDJ sequencing in dogs identified MAIT-like
cells based on restricted TRAV gene usage and restricted junctional length (19). Although
cytotoxic T cells in the present study exhibited differential TRAV gene and, to a lesser extent,
differential TRBV gene usage, there was no obvious junctional length restriction or dominant
amino acid motif. The fact that we were unable to identify MAIT-like cells in cats could reflect
their true absence or represent an artifact of CD5-based enrichment, which might have resulted in
the exclusion of CD5[-] or CD5[low] subsets. Additional studies with more animals, inclusion of
different tissue types and without selection bias are needed to elucidate the existence of innate-like
alpha/beta T cells in cats.
374
Among the myeloid cells, we resolved 3 clusters of classical monocytes and 1 cluster of
conventional dendritic cells. Of the three clusters of monocytes, we found two to be differentiated
in a somewhat polarized manner- pro-inflammatory versus anti-inflammatory. The anti-
inflammatory population was unconventional in that it had an upregulation of classic pro-
inflammatory cytokines *IL-1A* and IL-*1B* but also an upregulation of genes associated with a
unique and novel monocytic cluster found in human COVID-19 patients such as *AREG, EREG,*
IL10 , *IL18* and *BCL6.* (59,72). Sub clustering of neutrophils identified a larger mature cluster and
a smaller exhausted/activated cluster which showed upregulation of pro-inflammatory and
apoptotic genes. Thus, analysis of these myeloid cells revealed the presence of conventional and
non-conventional subtypes, providing a footing for feline circulating myeloid immunology.
Currently, T cell subtypes have only been described at the flow cytometric level, and with a focus
on *CD4/CD8* populations due to paucity of species-specific reagents. Our data indicates the
presence of unique subtypes of peripheral blood immune cells including αβγ CD8[+] cytotoxic cells,
CD8A/CD8B naive T cells, plasmacytoid DC, mast cells, conventional DCs, anti-inflammatory
monocytes and mature and end stage neutrophils. Our study is the first of many necessary for the

further characterization of the feline immune system. Taken together, it has increased our
understanding of feline circulating immune cells, including T cell subtyping and capture of rare
populations previously undescribed in the cat.
393
5   Materials and Methods
5.1   Samples and Sorting
Whole blood was obtained from four healthy, female, domestic shorthair cats (13-101/9 years, 18-
003/6 years, 21-005/1 year, 22-007/6 months) housed at the UC Davis Nutrition and Pet Care
Center in accordance with the UC Davis Policy and Procedure Manual section 290-30 and Animal
Use and Care protocol #22150. Blood was collected in purple top EDTA tubes and immediately
processed in one batch without freezing until completion of the reverse transcription step during
library preparation. Peripheral blood mononuclear cells (PBMC) were isolated from whole blood
by density gravity centrifugation. Briefly, 5 mL of whole blood was layered on top of 10 mL
Histopaque 1077 (density 1.077g/mL, Sigma-Aldrich, Burlington, MA, USA) in a 50 ml tube. The
samples were centrifuged at 850 revolutions per minute (RCF) for 30 minutes at room temperature
(RT) without break. After centrifugation, the PBMC layer at the plasma-histopaque interface was
aspirated and transferred to a clean 15 ml tube. The PBMCs were then washed twice with 14 ml
PBMC isolation media (500 mL Hanks' Balanced Salt Solution (HBSS), 15 mL Fetal Bovine
Serum (FBS), 2 mL EDTA) by centrifuging at 850 RCF for 10 minutes at RT. Following the
second washing step, the PBMC pellet was resuspended in 200 μl modified flow buffer solution
(MFB, 500 ml phosphate buffer saline (PBS), 0.467g EDTA tetrasodium salt and 1% Horse
serum).
The isolated PBMCs were first incubated with a primary anti-feline CD5 monoclonal antibody
(clone FE 1.1.B11-IgG1-Azide, Peter F. Moore Leukocyte Antigen Biology Laboratory, UC
Davis, CA, USA) or a non-cross reacting canine antibody (negative control) for 30 min at RT. This
was followed by a wash step (addition off 3 mL of MFB, followed by centrifugation at 400 RCF
for 3 min at RT and resuspension in 100 - 200 μl of supernatant), and incubation with a secondary
horse-anti-mouse FITC antibody (1:99 in MFB) for 15 minutes at RT in darkness. Finally, the
stained PBMC was washed again (as above), resuspended in 500 μl of MFB and stained with DAPI
(1 μg/mL). Cells were sorted at the UC Davis Comprehensive Cancer Center Flow Cytometry
Core. Cells were initially gated based on FSC vs. SSC to exclude debris and aggregates. Singlet
gating was then applied based on SSC to exclude doublets, followed by discrimination of live and
dead cells based on DAPI staining and subsequent gating for CD5 positive (FITC positive) cells
<Supplementary File 11>.
424
425 **5.2   Single-cell 5' RNA and V(D)J library preparation, sequencing and mapping**
426 Libraries were constructed using the Chromium Next GEM Single Cell 5' Kit v2 (10x Genomics).
427 All four feline T cell receptor loci were amplified using the Chromium Single Cell V(D)J
428 Enrichment Kit according to the manufacturer's instructions but with custom reverse primers
429 specific for the feline orthologs **<Supplementary File 12>**. Separate libraries were prepared for

the transcriptome, alpha/beta TCRs, and gamma/delta TCRs **<Supplementary File 12>**. Libraries were sequenced using the Illumina NovaSeq S4 platform using 150 paired-end reads to a target read depth of 30,000 reads per cell. Raw FASTQ sequencing data were processed by the UC Davis Bioinformatics Core using 10X CellRanger version 7.1 for alignment to the cat reference genome (*Felis_catus_9.0*). To improve the quality of annotation of the *Felis catus* genome, we converted unmapped ENSEMBL IDs to human homologs from BioMart for greater interpretative power. Gene mapping was performed in the following order: (1) Mapping of one-to-one orthologs from cat to human (2) Mapping of one-to-many and many-to-many was performed by choosing a representative cat gene based on the following parameters, in respective order [orthology_confidence, gene_order_confidence, Whole_Genome_alignment, %_query_identical_target, %target_identical_query]. Additionally, specific unmapped genes were manually annotated or filtered **<Supplementary File 13>**. Further analysis was then continued using Seurat v5 (73).

5.3   Count matrix pre-processing and quality control

Filtered feature matrices for the 4 samples were independently normalized and scaled using the function *SCTransform* v2 (glmGamPoi) (74). Dimensionality reduction (determined using knee identification in elbow plot), k-means clustering (resolution determined using clustree package) and neighbor identification (74–76) was performed. Low quality clusters were identified and filtered based on low gene counts, high mitochondrial read percentage and lack of biologically relevant markers. To further ensure quality, doublet identification was performed using package *DoubletFinder* according to package vignette and ambient RNA decontamination was performed using the *DecontX* package as per package tutorials adapted to the data (77,78).

5.4   Integration and Downstream analysis

Objects were integrated by sample using the Seurat native reciprocal principal component analysis (RPCA) (79). The embeddings were utilized for Uniform Manifold Approximation and Projection (UMAP) and subsequent clustering and neighbor identification as described prior. Differential gene expression analysis was performed using each cluster identified by the Seurat function *FindAllMarkers* (Non-parametric Wilcoxon Rank Sum with adjusted P value <0.05). Gene set enrichment analyses including over-enrichment analysis (ORA) of gene ontology (biological process) was performed using *ClusterProfiler* on human homologized genes (adjusted P value <0.05) (80). Additional visualizations and associated statistics were performed using ggplot2 and gg supplementary packages (81). Figures were illustrated on Bio Render.

5.5   Cell type sub clustering

Cell type sub-clustering (example: T cell sub-clustering) analysis was performed in the same manner as described prior. Pseudotime values were calculated using the UMAP embeddings produced via Seurat with RPCA integration using package *monocle3* (82). Gene module scores were calculated via the Seurat function *AddModuleScore*.

5.6 Cross species T cell integrative analysis

Published single cell RNA-sequencing datasets were obtained from NCBI GEO Accession for 4 species including dog (*Canis familiaris*: GSE225599 (18)), human (*Homo sapiens*: https://support.10xgenomics.com/single-cell-gene-expression/datasets/3.0.0/pbmc_10k_v3), horse (*Equus caballus*: GSE148416 (22)) and pig (*Sus scrofa*: https://data.nal.usda.gov/dataset/data-reference-transcriptomics-porcine-peripheral-immune-cells-created-through-bulk-and-single-cell-rna-sequencing. (21)). All species genes were mapped to human homologs as described prior for the cat. Only homologized shared genes between all 5 species were utilized for analysis. T cell clusters annotated by authors were validated using canonical T cell markers (*CD3E*, *CD3D*, *TCF7*, *LCK*, *ITK*). T cells were integrated across species using Seurat canonical correlation analysis (CCA) and downstream analysis proceeded in the same manner as described prior for global object analysis.

5.7 VDJ analysis

Contigs were mapped to available IMGT reference genes (as of February 2023) using Cell Ranger version 7.1. All analyses were based on the 'all_contig_annotations.csv' file using the R packages tidyverse, circlize and ggseqlogo (83-85). Cluster analysis was performed as previously described (86).

5.8 Data availability

All raw and processed data have been submitted to NCBI GEO and are available under GSE267355.

6 Acknowledgements

This project was funded by grant #2022-24-F from the Center for Companion Animal Health, University of California, Davis. The project was supported by the University of California Davis Flow Cytometry Shared Resource Laboratory with funding from the NCI P30 CA093373 (Comprehensive Cancer Center) and S10 OD018223, with technical assistance from Bridget McLaughlin, Jonathan Van Dyke and Ashley Karajeh. The sequencing was carried out by the DNA Technologies and Expression Analysis Core at the UC Davis Genome Center, supported by NIH Shared Instrumentation Grant 1S10OD010786-01. We thank Jie (Jessie) Li from the UC Davis Bioinformatics Core for data analysis and Kristy Harmon from the Leukocyte Biology Antigen Laboratory for her flow cytometry support.

7 References

1.   Rowan AN. Companion Animal Statistics in the USA. 2018 [cited 2024 Apr 30];
      Available from: https://www.wellbeingintlstudiesrepository.org/demscapop/7/

2. Larkin M, Radich R. AVMA. Am Vet Med Assoc; 2021 [cited 2024 Apr 5]. Pet
population still on the rise, with fewer pets per household. Available from:
https://avmajournals.avma.org/display/post/news/pet-population-still-on-the-rise--with-
fewer-pets-per-household.xml?print&print&print&print&print
3. Fromont E, Artois M, Langlais M, Courchamp F, Pontier D. Modelling the feline
leukemia virus (FeLV) in natural populations of cats (Felis catus). Theor Popul Biol
[Internet]. 1997 Aug;52(1):60–70. Available from:
http://dx.doi.org/10.1006/tpbi.1997.1320
4. Powers JA, Chiu ES, Kraberger SJ, Roelke-Parker M, Lowery I, Erbeck K, et al. Feline
Leukemia Virus (FeLV) Disease Outcomes in a Domestic Cat Breeding Colony:
Relationship to Endogenous FeLV and Other Chronic Viral Infections. J Virol [Internet].
2018 Sep 15;92(18). Available from: http://dx.doi.org/10.1128/JVI.00649-18
5. Helfer-Hungerbuehler AK, Shah J, Meili T, Boenzli E, Li P, Hofmann-Lehmann R.
Adeno-Associated Vector-Delivered CRISPR/SaCas9 System Reduces Feline Leukemia
Virus Production In Vitro. Viruses [Internet]. 2021 Aug 18;13(8). Available from:
http://dx.doi.org/10.3390/v13081636
6. Bienzle D. FIV in cats--a useful model of HIV in people? Vet Immunol Immunopathol
[Internet]. 2014 Jun 15;159(3-4):171–9. Available from:
http://dx.doi.org/10.1016/j.vetimm.2014.02.014
7. Paillot R, Richard S, Bloas F, Piras F, Poulet H, Brunet S, et al. Toward a detailed
characterization of feline immunodeficiency virus-specific T cell immune responses and
mediated immune disorders. Vet Immunol Immunopathol [Internet]. 2005 Jun 15;106(1-
2):1–14. Available from: http://dx.doi.org/10.1016/j.vetimm.2004.12.023
8. Endo Y, Matsumura S, Washizu T, Ishida T. Alteration of T-cell subsets in the lymph
nodes from cats infected with feline immunodeficiency virus. J Vet Med Sci [Internet].
1997 Sep;59(9):739–46. Available from: http://dx.doi.org/10.1292/jvms.59.739
9. Hoffmann-Fezer G, Mortelbauer W, Hartmann K, Mysliwietz J, Thefeld S, Beer B, et al.
Comparison of T-cell subpopulations in cats naturally infected with feline leukaemia
virus or feline immunodeficiency virus. Res Vet Sci [Internet]. 1996 Nov;61(3):222–6.
Available from: http://dx.doi.org/10.1016/s0034-5288(96)90067-3
10. Day MJ. Cats are not small dogs: is there an immunological explanation for why cats are
less affected by arthropod-borne disease than dogs? Parasit Vectors [Internet]. 2016 Sep
20;9(1):507. Available from: http://dx.doi.org/10.1186/s13071-016-1798-5
11. Akira S, Uematsu S, Takeuchi O. Pathogen recognition and innate immunity. Cell
[Internet]. 2006 Feb 24;124(4):783–801. Available from:
http://dx.doi.org/10.1016/j.cell.2006.02.015
12. Jiao A, Zhang C, Wang X, Sun L, Liu H, Su Y, et al. Single-cell sequencing reveals the
evolution of immune molecules across multiple vertebrate species. J Advert Res
[Internet]. 2023 Mar 4; Available from: http://dx.doi.org/10.1016/j.jare.2023.02.017

13. Radtanakatikanon A, Keller SM, Darzentas N, Moore PF, Folch G, Nguefack Ngoune V,
et al. Topology and expressed repertoire of the Felis catus T cell receptor loci. BMC
Genomics [Internet]. 2020 Jan 6;21(1):20. Available from:
http://dx.doi.org/10.1186/s12864-019-6431-5
14. Ginhoux F, Yalin A, Dutertre CA, Amit I. Single-cell immunology: Past, present, and
future. Immunity [Internet]. 2022 Mar 8;55(3):393–404. Available from:
http://dx.doi.org/10.1016/j.immuni.2022.02.006
15. Jovic D, Liang X, Zeng H, Lin L, Xu F, Luo Y. Single-cell RNA sequencing
technologies and applications: A brief overview. Clin Transl Med [Internet]. 2022
Mar;12(3):e694. Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1002/ctm2.694
16. Sun X, Lin X, Li Z, Wu H. A comprehensive comparison of supervised and unsupervised
methods for cell type identification in single-cell RNA-seq. Brief Bioinform [Internet].
2022 Mar 10;23(2). Available from: http://dx.doi.org/10.1093/bib/bbab567
17. Wang M, Song WM, Ming C, Wang Q, Zhou X, Xu P, et al. Guidelines for
bioinformatics of single-cell sequencing data analysis in Alzheimer's disease: review,
recommendation, implementation and application. Mol Neurodegener [Internet]. 2022
Mar 2;17(1):17. Available from: http://dx.doi.org/10.1186/s13024-022-00517-z
18. Ammons DT, Harris RA, Hopkins LS, Kurihara J, Weishaar K, Dow S. A single-cell
RNA sequencing atlas of circulating leukocytes from healthy and osteosarcoma affected
dogs. Front Immunol [Internet]. 2023 May 19;14:1162700. Available from:
http://dx.doi.org/10.3389/fimmu.2023.1162700
19. Eschke M, Moore PF, Chang H, Alber G, Keller SM. Canine peripheral blood TCRαβ T
cell atlas: Identification of diverse subsets including CD8A+ MAIT-like cells by
combined single-cell transcriptome and V(D)J repertoire analysis. Front Immunol
[Internet]. 2023 Feb 23;14:1123366. Available from:
http://dx.doi.org/10.3389/fimmu.2023.1123366
20. Gao Y, Li J, Cai G, Wang Y, Yang W, Li Y, et al. Single-cell transcriptomic and
chromatin accessibility analyses of dairy cattle peripheral blood mononuclear cells and
their responses to lipopolysaccharide. BMC Genomics [Internet]. 2022 Apr 30;23(1):338.
Available from: http://dx.doi.org/10.1186/s12864-022-08562-0
21. Herrera-Uribe J, Wiarda JE, Sivasankaran SK, Daharsh L, Liu H, Byrne KA, et al.
Reference Transcriptomes of Porcine Peripheral Immune Cells Created Through Bulk
and Single-Cell RNA Sequencing. Front Genet [Internet]. 2021 Jun 23;12:689406.
Available from: http://dx.doi.org/10.3389/fgene.2021.689406
22. Patel RS, Tomlinson JE, Divers TJ, Van de Walle GR, Rosenberg BR. Single-cell
resolution landscape of equine peripheral blood mononuclear cells reveals diverse cell
types including T-bet+ B cells. BMC Biol [Internet]. 2021 Jan 22;19(1):13. Available
from: http://dx.doi.org/10.1186/s12915-020-00947-5

23. Li Z, Sun C, Wang F, Wang X, Zhu J, Luo L, et al. Molecular mechanisms governing
circulating immune cell heterogeneity across different species revealed by single-cell
sequencing. Clin Transl Med [Internet]. 2022 Jan;12(1):e689. Available from:
http://dx.doi.org/10.1002/ctm2.689
24. Touw IP, Palande K, Beekman R. Granulocyte colony-stimulating factor receptor
signaling: implications for G-CSF responses and leukemic progression in severe
congenital neutropenia. Hematol Oncol Clin North Am [Internet]. 2013 Feb;27(1):61–73,
viii. Available from: http://dx.doi.org/10.1016/j.hoc.2012.10.002
25. Sprenkeler EGG, Zandstra J, van Kleef ND, Goetschalckx I, Verstegen B, Aarts CEM, et
al. S100A8/A9 Is a Marker for the Release of Neutrophil Extracellular Traps and Induces
Neutrophil Activation. Cells [Internet]. 2022 Jan 11;11(2). Available from:
http://dx.doi.org/10.3390/cells11020236
26. Akgun E, Tuzuner MB, Sahin B, Kilercik M, Kulah C, Cakiroglu HN, et al. Proteins
associated with neutrophil degranulation are upregulated in nasopharyngeal swabs from
SARS-CoV-2 patients. PLoS One [Internet]. 2020 Oct 20;15(10):e0240012. Available
from: http://dx.doi.org/10.1371/journal.pone.0240012
27. Blanter M, Cambier S, De Bondt M, Vanbrabant L, Pörtner N, Abouelasrar Salama S, et
al. Method Matters: Effect of Purification Technology on Neutrophil Phenotype and
Function. Front Immunol [Internet]. 2022 Feb 10;13:820058. Available from:
http://dx.doi.org/10.3389/fimmu.2022.820058
28. Buechler C, Ritter M, Orsó E, Langmann T, Klucken J, Schmitz G. Regulation of
scavenger receptor CD163 expression in human monocytes and macrophages by pro- and
antiinflammatory stimuli. J Leukoc Biol [Internet]. 2000 Jan;67(1):97–103. Available
from: https://www.ncbi.nlm.nih.gov/pubmed/10648003
29. Sampath P, Moideen K, Ranganathan UD, Bethunaickan R. Monocyte Subsets:
Phenotypes and Function in Tuberculosis Infection. Front Immunol [Internet]. 2018 Jul
30;9:1726. Available from: http://dx.doi.org/10.3389/fimmu.2018.01726
30. Reshetnikova E, Guselnikov S, Volkova O, Baranov K, Taranin A, Mechetina L. B cell-
specific protein FCRLA is expressed by plasmacytoid dendritic cells in humans.
Cytometry B Clin Cytom [Internet]. 2018 Jul;94(4):683–7. Available from:
http://dx.doi.org/10.1002/cyto.b.21611
31. Rissoan MC, Duhen T, Bridon JM, Bendriss-Vermare N, Péronne C, de Saint Vis B, et
al. Subtractive hybridization reveals the expression of immunoglobulin-like transcript 7,
Eph-B1, granzyme B, and 3 novel transcripts in human plasmacytoid dendritic cells.
Blood [Internet]. 2002 Nov 1;100(9):3295–303. Available from:
http://dx.doi.org/10.1182/blood-2002-02-0638
32. Edwards AD, Diebold SS, Slack EMC, Tomizawa H, Hemmi H, Kaisho T, et al. Toll-like
receptor expression in murine DC subsets: lack of TLR7 expression by CD8α+ DC
correlates with unresponsiveness to imidazoquinolines. Eur J Immunol [Internet]. 2003

Apr;33(4):827–33. Available from:
https://onlinelibrary.wiley.com/doi/10.1002/eji.200323797

33. Medina KL, Tangen SN, Seaburg LM, Thapa P, Gwin KA, Shapiro VS. Separation of plasmacytoid dendritic cells from B-cell-biased lymphoid progenitor (BLP) and Pre-pro B cells using PDCA-1. PLoS One [Internet]. 2013 Oct 30;8(10):e78408. Available from: http://dx.doi.org/10.1371/journal.pone.0078408

34. Arroyo Hornero R, Idoyaga J. Plasmacytoid dendritic cells: A dendritic cell in disguise. Mol Immunol [Internet]. 2023 Jul;159:38–45. Available from: http://dx.doi.org/10.1016/j.molimm.2023.05.007

35. Atiakshin D, Kostin A, Trotsenko I, Samoilova V, Buchwalow I, Tiemann M. Carboxypeptidase A3-A Key Component of the Protease Phenotype of Mast Cells. Cells [Internet]. 2022 Feb 6;11(3). Available from: http://dx.doi.org/10.3390/cells11030570

36. Cruse G, Kaur D, Leyland M, Bradding P. A novel FcεRIβ-chain truncation regulates human mast cell proliferation and survival. FASEB J [Internet]. 2010 Oct;24(10):4047–57. Available from: http://dx.doi.org/10.1096/fj.10-158378

37. Lee H, Joo JY, Kang J, Yu Y, Kim YH, Park HR. Single-cell analysis of platelets from patients with periodontitis and diabetes. Res Pract Thromb Haemost [Internet]. 2023 Feb;7(2):100099. Available from: http://dx.doi.org/10.1016/j.rpth.2023.100099

38. Yang S, Liu F, Wang QJ, Rosenberg SA, Morgan RA. The shedding of CD62L (L-selectin) regulates the acquisition of lytic activity in human tumor reactive T lymphocytes. PLoS One [Internet]. 2011 Jul 28;6(7):e22560. Available from: http://dx.doi.org/10.1371/journal.pone.0022560

39. Ivetic A, Hoskins Green HL, Hart SJ. L-selectin: A Major Regulator of Leukocyte Adhesion, Migration and Signaling. Front Immunol [Internet]. 2019 May 14;10:1068. Available from: http://dx.doi.org/10.3389/fimmu.2019.01068

40. Bjorkdahl O, Barber KA, Brett SJ, Daly MG, Plumpton C, Elshourbagy NA, et al. Characterization of CC-chemokine receptor 7 expression on murine T cells in lymphoid tissues. Immunology [Internet]. 2003 Oct;110(2):170–9. Available from: http://dx.doi.org/10.1046/j.1365-2567.2003.01727.x

41. Cano-Gamez E, Soskic B, Roumeliotis TI, So E, Smyth DJ, Baldrighi M, et al. Single-cell transcriptomics identifies an effectorness gradient shaping the response of CD4+ T cells to cytokines. Nat Commun [Internet]. 2020 Apr 14;11(1):1801. Available from: http://dx.doi.org/10.1038/s41467-020-15543-y

42. Zhu J, Yamane H, Paul WE. Differentiation of Effector CD4 T Cell Populations*. Annu Rev Immunol [Internet]. 2009 Apr 23 [cited 2024 Mar 23];28(Volume 28, 2009):445–89. Available from: https://www.annualreviews.org/content/journals/10.1146/annurev-immunol-030409-101212

43. Andreatta M, Corria-Osorio J, Müller S, Cubas R, Coukos G, Carmona SJ. Interpretation of T cell states from single-cell transcriptomics data using reference atlases. Nat

Commun [Internet]. 2021 May 20;12(1):2965. Available from:
http://dx.doi.org/10.1038/s41467-021-23324-4

44. Wang X, Shen X, Chen S, Liu H, Hong N, Zhong H, et al. Reinvestigation of Classic T
Cell Subsets and Identification of Novel Cell Subpopulations by Single-Cell RNA
Sequencing. J Immunol [Internet]. 2022 Jan 15;208(2):396–406. Available from:
http://dx.doi.org/10.4049/jimmunol.2100581

45. Vallejo J, Saigusa R, Gulati R, Armstrong Suthahar SS, Suryawanshi V, Alimadadi A, et
al. Combined protein and transcript single-cell RNA sequencing in human peripheral
blood mononuclear cells. BMC Biol [Internet]. 2022 Sep 1;20(1):193. Available from:
http://dx.doi.org/10.1186/s12915-022-01382-4

46. Schneider L, Marcondes NA, Hax V, da Silva Moreira IF, Ueda CY, Piovesan RR, et al.
Flow cytometry evaluation of CD14/CD16 monocyte subpopulations in systemic
sclerosis patients: a cross sectional controlled study. Adv Rheumatol [Internet]. 2021
May 22;61(1):27. Available from: http://dx.doi.org/10.1186/s42358-021-00182-8

47. Barut GT, Kreuzer M, Bruggmann R, Summerfield A, Talker SC. Single-cell
transcriptomics reveals striking heterogeneity and functional organization of dendritic
and monocytic cells in the bovine mesenteric lymph node. Front Immunol [Internet].
2022;13:1099357. Available from: http://dx.doi.org/10.3389/fimmu.2022.1099357

48. Waskow C, Liu K, Darrasse-Jèze G, Guermonprez P, Ginhoux F, Merad M, et al. The
receptor tyrosine kinase Flt3 is required for dendritic cell development in peripheral
lymphoid tissues. Nat Immunol [Internet]. 2008 Jun;9(6):676–83. Available from:
http://dx.doi.org/10.1038/ni.1615

49. ten Broeke T, Wubbolts R, Stoorvogel W. MHC class II antigen presentation by dendritic
cells regulated through endosomal sorting. Cold Spring Harb Perspect Biol [Internet].
2013 Dec 1;5(12):a016873. Available from:
http://dx.doi.org/10.1101/cshperspect.a016873

50. Eislmayr K, Bestehorn A, Morelli L, Borroni M, Vande Walle L, Lamkanfi M, et al.
Nonredundancy of IL-1α and IL-1β is defined by distinct regulation of tissues
orchestrating resistance versus tolerance to infection. Sci Adv [Internet]. 2022 Mar
4;8(9):eabj7293. Available from: http://dx.doi.org/10.1126/sciadv.abj7293

51. Lopez-Castejon G, Brough D. Understanding the mechanism of IL-1β secretion.
Cytokine Growth Factor Rev [Internet]. 2011 Aug;22(4):189–95. Available from:
http://dx.doi.org/10.1016/j.cytogfr.2011.10.001

52. Hadadi E, Zhang B, Baidžajevas K, Yusof N, Puan KJ, Ong SM, et al. Differential IL-1β
secretion by monocyte subsets is regulated by Hsp27 through modulating mRNA
stability. Sci Rep [Internet]. 2016 Dec 15;6:39035. Available from:
http://dx.doi.org/10.1038/srep39035

53. Wan J, Shan Y, Fan Y, Fan C, Chen S, Sun J, et al. NF-κB inhibition attenuates LPS-
induced TLR4 activation in monocyte cells. Mol Med Rep [Internet]. 2016
Nov;14(5):4505–10. Available from: http://dx.doi.org/10.3892/mmr.2016.5825

54.   Mussbacher M, Derler M, Basílio J, Schmid JA. NF-κB in monocytes and macrophages -
an inflammatory master regulator in multitalented immune cells. Front Immunol
[Internet]. 2023 Feb 23;14:1134661. Available from:
http://dx.doi.org/10.3389/fimmu.2023.1134661
55.   Guo H, Jin D, Chen X. Lipocalin 2 is a regulator of macrophage polarization and NF-
κB/STAT3 pathway activation. Mol Endocrinol [Internet]. 2014 Oct;28(10):1616–28.
Available from: http://dx.doi.org/10.1210/me.2014-1092
56.   Lee BC, Lee SG, Choo MK, Kim JH, Lee HM, Kim S, et al. Selenoprotein MsrB1
promotes anti-inflammatory cytokine gene expression in macrophages and controls
immune response in vivo. Sci Rep [Internet]. 2017 Jul 11;7(1):5119. Available from:
http://dx.doi.org/10.1038/s41598-017-05230-2
57.   Vishnyakova P, Poltavets A, Karpulevich E, Maznina A, Vtorushina V, Mikhaleva L, et
al. The response of two polar monocyte subsets to inflammation. Biomed Pharmacother
[Internet]. 2021 Jul;139:111614. Available from:
http://dx.doi.org/10.1016/j.biopha.2021.111614
58.   Zhang Y, Wang S, Xia H, Guo J, He K, Huang C, et al. Identification of Monocytes
Associated with Severe COVID-19 in the PBMCs of Severely Infected Patients Through
Single-Cell Transcriptome Sequencing. Engineering [Internet]. 2022 Oct;17:161–9.
Available from: http://dx.doi.org/10.1016/j.eng.2021.05.009
59.   Utrero-Rico A, González-Cuadrado C, Chivite-Lacaba M, Cabrera-Marante O, Laguna-
Goya R, Almendro-Vazquez P, et al. Alterations in Circulating Monocytes Predict
COVID-19 Severity and Include Chromatin Modifications Still Detectable Six Months
after Recovery. Biomedicines [Internet]. 2021 Sep 17;9(9). Available from:
http://dx.doi.org/10.3390/biomedicines9091253
60.   Rynikova M, Adamkova P, Hradicka P, Stofilova J, Harvanova D, Matejova J, et al.
Transcriptomic Analysis of Macrophage Polarization Protocols: Vitamin D3 or IL-4 and
IL-13 Do Not Polarize THP-1 Monocytes into Reliable M2 Macrophages. Biomedicines
[Internet]. 2023 Feb 17;11(2). Available from:
http://dx.doi.org/10.3390/biomedicines11020608
61.   Wigerblad G, Cao Q, Brooks S, Naz F, Gadkari M, Jiang K, et al. Single-Cell Analysis
Reveals the Range of Transcriptional States of Circulating Human Neutrophils. J
Immunol [Internet]. 2022 Aug 15;209(4):772–82. Available from:
http://dx.doi.org/10.4049/jimmunol.2200154
62.   Meng Y, Lin S, Niu K, Ma Z, Lin H, Fan H. Vimentin affects inflammation and
neutrophil recruitment in airway epithelium during Streptococcus suis serotype 2
infection. Vet Res [Internet]. 2023 Jan 30;54(1):7. Available from:
http://dx.doi.org/10.1186/s13567-023-01135-3
63.   McColl SR, Paquin R, Ménard C, Beaulieu AD. Human neutrophils produce high levels
of the interleukin 1 receptor antagonist in response to granulocyte/macrophage colony-

stimulating factor and tumor necrosis factor alpha. J Exp Med [Internet]. 1992 Aug 1;176(2):593–8. Available from: http://dx.doi.org/10.1084/jem.176.2.593

64. Kunisada M, Yogianti F, Sakumi K, Ono R, Nakabeppu Y, Nishigori C. Increased expression of versican in the inflammatory response to UVB- and reactive oxygen species-induced skin tumorigenesis. Am J Pathol [Internet]. 2011 Dec;179(6):3056–65. Available from: http://dx.doi.org/10.1016/j.ajpath.2011.08.042

65. Xie X, Shi Q, Wu P, Zhang X, Kambara H, Su J, et al. Single-cell transcriptome profiling reveals neutrophil heterogeneity in homeostasis and infection. Nat Immunol [Internet]. 2020 Sep;21(9):1119–33. Available from: http://dx.doi.org/10.1038/s41590-020-0736-z

66. Glennon-Alty L, Moots RJ, Edwards SW, Wright HL. Type I interferon regulates cytokine-delayed neutrophil apoptosis, reactive oxygen species production and chemokine expression. Clin Exp Immunol [Internet]. 2021 Feb;203(2):151–9. Available from: http://dx.doi.org/10.1111/cei.13525

67. Leylek R, Idoyaga J. Chapter Four - The versatile plasmacytoid dendritic cell: Function, heterogeneity, and plasticity. In: Lhuillier C, Galluzzi L, editors. International Review of Cell and Molecular Biology [Internet]. Academic Press; 2019. p. 177–211. Available from: https://www.sciencedirect.com/science/article/pii/S1937644819300991

68. Varricchi G, de Paulis A, Marone G, Galli SJ. Future Needs in Mast Cell Biology. Int J Mol Sci [Internet]. 2019 Sep 6;20(18). Available from: http://dx.doi.org/10.3390/ijms20184397

69. Nascimento C, Ferreira F. Tumor microenvironment of human breast cancer, and feline mammary carcinoma as a potential study model. Biochim Biophys Acta Rev Cancer [Internet]. 2021 Aug;1876(1):188587. Available from: http://dx.doi.org/10.1016/j.bbcan.2021.188587

70. Andrade LM, Sousa ECM, Melo ACS, Pardavil GSS, Silva RA, Valente KF, et al. Gene expression of cytokines in the blood of domestics cats infected and not infected by Feline alphaherpesvirus. Available from: http://dx.doi.org/10.4238/gmr19190

71. Smithberg SR, Fogle JE, Mexas AM, Reckling SK, Lankford SM, Tompkins MB, et al. In vivo depletion of CD4+CD25+ regulatory T cells in cats. J Immunol Methods [Internet]. 2008 Jan 1;329(1-2):81–91. Available from: http://dx.doi.org/10.1016/j.jim.2007.09.015

72. Rigamonti A, Castagna A, Viatore M, Colombo FS, Terzoli S, Peano C, et al. Distinct responses of newly identified monocyte subsets to advanced gastrointestinal cancer and COVID-19. Front Immunol [Internet]. 2022 Oct 3;13:967737. Available from: http://dx.doi.org/10.3389/fimmu.2022.967737

73. Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of single-cell gene expression data. Nat Biotechnol [Internet]. 2015 May;33(5):495–502. Available from: http://dx.doi.org/10.1038/nbt.3192

74. Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. Genome Biol [Internet]. 2019 Dec
23;20(1):296. Available from: http://dx.doi.org/10.1186/s13059-019-1874-1
75. Zappia L, Oshlack A. Clustering trees: a visualization for evaluating clusterings at
multiple resolutions. Gigascience [Internet]. 2018 Jul 1;7(7). Available from:
http://dx.doi.org/10.1093/gigascience/giy083
76. Ahlmann-Eltze C, Huber W. glmGamPoi: fitting Gamma-Poisson generalized linear
models on single cell count data. Bioinformatics [Internet]. 2021 Apr 5;36(24):5701–2.
Available from: http://dx.doi.org/10.1093/bioinformatics/btaa1009
77. Yang S, Corbett SE, Koga Y, Wang Z, Johnson WE, Yajima M, et al. Decontamination
of ambient RNA in single-cell RNA-seq with DecontX. Genome Biol [Internet]. 2020
Mar 5;21(1):57. Available from: http://dx.doi.org/10.1186/s13059-020-1950-6
78. McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell
RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst [Internet]. 2019 Apr
24;8(4):329–37.e4. Available from: http://dx.doi.org/10.1016/j.cels.2019.03.003
79. Song Y, Miao Z, Brazma A, Papatheodorou I. Benchmarking strategies for cross-species
integration of single-cell RNA sequencing data. Nat Commun [Internet]. 2023 Oct
14;14(1):6495. Available from: http://dx.doi.org/10.1038/s41467-023-41855-w
80. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal
enrichment tool for interpreting omics data. Innovation (Camb) [Internet]. 2021 Aug
28;2(3):100141. Available from: http://dx.doi.org/10.1016/j.xinn.2021.100141
81. Wickham H. ggplot2 [Internet]. Springer International Publishing; [cited 2023 Oct 16].
16 p. Available from: https://link.springer.com/book/10.1007/978-3-319-24277-4
82. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics
and regulators of cell fate decisions are revealed by pseudotemporal ordering of single
cells. Nat Biotechnol [Internet]. 2014 Apr;32(4):381–6. Available from:
http://dx.doi.org/10.1038/nbt.2859
83. R Core Team (2023). _R: A Language and Environment for Statistical Computing_. R
Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
84. Gu Z, Gu L, Eils R, Schlesner M, Brors B. circlize Implements and enhances circular
visualization in R. Bioinformatics. 2014 Oct;30(19):2811-2. doi:
10.1093/bioinformatics/btu393. Epub 2014 Jun 14. PMID: 24930139.
85. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G,
Hayes A, Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K,
Ooms J, Robinson D, Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K,
Yutani H (2019). "Welcome to the tidyverse." Journal of Open Source Software, 4(43),
1686. doi:10.21105/joss.01686.

86.   Chang H, Ashlock DA, Graether SP, Keller SM. Anchor Clustering for million-scale
820   immune repertoire sequencing data. BMC Bioinformatics. 2024 Jan 25;25(1):42. doi:
821   10.1186/s12859-024-05659-z. PMID: 38273275; PMCID: PMC10809746.
822

Figure 1. scRNA-seq atlas of CD5⁺ enriched circulating feline immune cells revealed 7 major
types across 21 clusters. (A) Table of cell counts from each age group. (B) UMAP plot
demonstrating the unsupervised clustering results for feline circulating immune cells from 4
healthy cats of different age groups. (C) UMAP plot of global clustering colored by age. (D)
Table of cell type counts across clusters by age. (E) Dot plot demonstrating cell type specific
marker expression of the 7 major cell types. (F-O) Feature UMAP plots demonstrating
expression profile of key markers for each of the 7 different cell types.

Figure 2. CD5[+] enriched T cells segregate into naïve T cell subtypes and effectors. Unsupervised
clustering of T cells reveals 7 subtypes. (A) UMAP of the scRNA-seq atlas of T cells. (B)
UMAP of T cells colored by Pseudotime. (C) Dot plot of marker genes expressed by T cell type.
(D) Table of cell type frequencies across ages by cluster. (E) Feature plots of expression and co-
expression of *TRG* and *TRDC*. (F-Q) UMAP of T cells colored by classical T marker genes
defining *CD4*/*CD8* status, naïve (*SELL*, *CCR7*), effectorness (*ANXA2*, *LGALS3*, *GZMK, PRF1*),
terminal differentiation (CCL5) and T exhaustion (*TIGIT*, *PDCD1*).

841
842

Figure 3. Feline Effector T cells (T$_{EM}$) segregate into transcriptionally similar clusters and reveal the presence of helper T phenotypes. Unsupervised clustering of T$_{EM}$ reveals 13 subtypes. (A) UMAP of scRNA-seq atlas of T$_{EM}$. (B) Dot plot demonstrating up to 3 top differentially expressed genes for each cluster determined via Wilcoxon rank sum testing (*Adj P <0.05*). (C,E,G,I) UMAP of T$_{EM}$ colored by helper T subtype gene modules. (D,F,H,J) Expression UMAP of a representative gene from each helper T gene module presented in parallel.

849
850
Figure 4. Cross-species integrative analysis of T cells reveals missing cytotoxic effectors in
the cat. Unsupervised clustering revealed 12 clusters across 5 species. (A) UMAP of scRNA-
seq atlas of T cells. (B-F) UMAP split by species- dog, horse, cat, human and pig. (G) UMAP
of T cells colored by T cell phenotype. (H) Dot plot of marker gene sets for T cell subtypes.
(I) Percentage bar chart of clusters stacked by species. (J) UMAP colored by *CD5* expression.
(K) Bar chart of percentage *CD5*+ cells per cluster. (L) Bar chart of average *CD5* expression
across cells in each cluster. (M) Scatter plot of percentage *CD5*+ cells per cluster versus number
of cat cells within the corresponding cluster.

859
860
Figure 5. (A) Percentage of T cells expressing a certain combination of productive TR chain
transcripts. TRA/TRB transcripts dominate in all ab T cell subsets except for CD8[+] cytotoxic
T cells, which primarily express TRA/TRB/TRG transcripts. (B) TRA & TRB V gene usage
of the 4 major ab T cell subsets. CD8[+] cytotoxic T cells show differential usage compared to
the other T cell subset but are also substantiated by fewer cells. (C) Junctional length of TRA
rearrangements of the 4 major ab T cell subsets. Junctional regions with dominant TRAV genes
in CD8[+] cytotoxic T cells are 13 and 14 amino acids long. (D) TRG transcripts in CD8[+]
cytotoxic T cells have a skewed junctional length with 16 amino acid length (top) and a
conserved motif (bottom) (E) The majority of TRG rearrangements in CD8[+] cytotoxic T cells
involved cassette 2 (TRGV2-2/TRGJ2-2). Genes are displayed in order of genomic location
and colored by cassette.

872
873
874 **Figure 6.** (A) Shared clonotypes across 4 cats. (B) Characterization of clusters based on mean
875 centrality and cluster density. (C) Representative examples of TRA clusters with high
876 centrality and density.

Figure 7. Feline circulating monocytes include classical monocytes with more differentiated clusters with traditional proinflammatory and unconventional anti-inflammatory phenotypes. Unsupervised clustering revealed 5 clusters. (A) UMAP of scRNA-seq atlas of monocytes. (B-F) UMAP of monocytes colored by expression levels of canonical markers. (G) Dot plot of genes of monocytic phenotypes across the 4 monocytic clusters. (H-J) Dot plots of top GO biological process terms called based on sets of positive differentially expressed genes identified via Seurat *FindMarker* function (Wilcoxon rank sum, *Adj P <0.05*). Classical monocyte, CM; Conventional dendritic cell, cDC; Pro-inf, proinflammatory; Anti-inf, anti-inflammatory.

887
888
Figure 8. Feline circulating neutrophils separate into 2 clusters based on activation state. (A)
UMAP of scRNA-seq atlas of neutrophils. (B-G) UMAP of neutrophils colored by expression
levels of canonical and functional neutrophil markers. (H) Dot plots of top GO biological
process terms in cluster 1 called based on sets of positive differentially expressed genes
identified via Seurat *FindMarker* function (Wilcoxon rank sum, Adj *P <0.05*). (I-K) UMAP
of neutrophils colored by expression levels of select differentially upregulated genes in cluster
1. (L,M) Violin plots of RNA counts and feature (unique RNA) counts per cell by cluster;
(****) indicates P value less than 0.001 by T-test. (N) UMAP of neutrophils colored by
interferon (IFN) gene composite score per cell; genes included are named in figure.

Figure 2. CD5[+] enriched T cells segregate into naïve T cell subtypes and effectors. Unsupervised clustering of T cells reveals 7 subtypes. (A) UMAP of the scRNA-seq atlas of T cells. (B) UMAP of T cells colored by Pseudotime. (C) Dot plot of marker genes expressed by T cell type. (D) Table of cell type frequencies across ages by cluster. (E) Feature plots of expression and co-expression of *TRG* and *TRDC*. (F-Q) UMAP of T cells colored by classical T marker genes defining *CD4/CD8* status, naïve (*SELL*, *CCR7*), effectorness (*ANXA2*, *LGALS3*, *GZMK, PRF1*), terminal differentiation (CCL5) and T exhaustion (*TIGIT*, *PDCD1*).

841
842

Figure 3. Feline Effector T cells (T_EM) segregate into transcriptionally similar clusters and reveal
the presence of helper T phenotypes. Unsupervised clustering of T_EM reveals 13 subtypes. (A)
UMAP of scRNA-seq atlas of T_EM. (B) Dot plot demonstrating up to 3 top differentially expressed
genes for each cluster determined via Wilcoxon rank sum testing (*Adj P <0.05*). (C,E,G,I) UMAP
of T_EM colored by helper T subtype gene modules. (D,F,H,J) Expression UMAP of a representative
gene from each helper T gene module presented in parallel.

849
850
Figure 4. Cross-species integrative analysis of T cells reveals missing cytotoxic effectors in
the cat. Unsupervised clustering revealed 12 clusters across 5 species. (A) UMAP of scRNA-
seq atlas of T cells. (B-F) UMAP split by species- dog, horse, cat, human and pig. (G) UMAP
of T cells colored by T cell phenotype. (H) Dot plot of marker gene sets for T cell subtypes.
(I) Percentage bar chart of clusters stacked by species. (J) UMAP colored by *CD5* expression.
(K) Bar chart of percentage *CD5*+ cells per cluster. (L) Bar chart of average *CD5* expression
across cells in each cluster. (M) Scatter plot of percentage *CD5*+ cells per cluster versus number
of cat cells within the corresponding cluster.

859
860

Figure 5. (A) Percentage of T cells expressing a certain combination of productive TR chain
transcripts. TRA/TRB transcripts dominate in all ab T cell subsets except for CD8[+] cytotoxic
T cells, which primarily express TRA/TRB/TRG transcripts. (B) TRA & TRB V gene usage
of the 4 major ab T cell subsets. CD8[+] cytotoxic T cells show differential usage compared to
the other T cell subset but are also substantiated by fewer cells. (C) Junctional length of TRA
rearrangements of the 4 major ab T cell subsets. Junctional regions with dominant TRAV genes
in CD8[+] cytotoxic T cells are 13 and 14 amino acids long. (D) TRG transcripts in CD8[+]
cytotoxic T cells have a skewed junctional length with 16 amino acid length (top) and a
conserved motif (bottom) (E) The majority of TRG rearrangements in CD8[+] cytotoxic T cells
involved cassette 2 (TRGV2-2/TRGJ2-2). Genes are displayed in order of genomic location
and colored by cassette.

872
873
Figure 6. (A) Shared clonotypes across 4 cats. (B) Characterization of clusters based on mean
centrality and cluster density. (C) Representative examples of TRA clusters with high
centrality and density.

Figure 7. Feline circulating monocytes include classical monocytes with more differentiated
clusters with traditional proinflammatory and unconventional anti-inflammatory phenotypes.
Unsupervised clustering revealed 5 clusters. (A) UMAP of scRNA-seq atlas of monocytes. (B-
F) UMAP of monocytes colored by expression levels of canonical markers. (G) Dot plot of
genes of monocytic phenotypes across the 4 monocytic clusters. (H-J) Dot plots of top GO
biological process terms called based on sets of positive differentially expressed genes
identified via Seurat *FindMarker* function (Wilcoxon rank sum, *Adj P <0.05*). Classical
monocyte, CM; Conventional dendritic cell, cDC; Pro-inf, proinflammatory; Anti-inf, anti-
inflammatory.

887
888
Figure 8. Feline circulating neutrophils separate into 2 clusters based on activation state. (A)
UMAP of scRNA-seq atlas of neutrophils. (B-G) UMAP of neutrophils colored by expression
levels of canonical and functional neutrophil markers. (H) Dot plots of top GO biological
process terms in cluster 1 called based on sets of positive differentially expressed genes
identified via Seurat *FindMarker* function (Wilcoxon rank sum, Adj *P <0.05*). (I-K) UMAP
of neutrophils colored by expression levels of select differentially upregulated genes in cluster
1. (L,M) Violin plots of RNA counts and feature (unique RNA) counts per cell by cluster;
(****) indicates P value less than 0.001 by T-test. (N) UMAP of neutrophils colored by
interferon (IFN) gene composite score per cell; genes included are named in figure.