***Clockwork Orangutan: microRNAs,
thermoregulatory tradeoffs,
and models of brain size evolution***

Thomas Sorger, PhD*
Department of Biology
Roger Williams University
Bristol, RI 02809

Bastian Fromm, PhD**
UiT - The Arctic University of Norway,
The Arctic University Museum of Norway
9006 Tromsø, NO

* Lead contact
** Second corresponding author

ABSTRACT

The Expensive Tissue Hypothesis was proposed to account for thermal homeostasis during the evolutionary expansion of brain size in the human lineage and abandoned following publication of a study that found no significant anticorrelations among the body mass-independent residuals of metabolically expensive organs in over 100 mammals, including 23 primates. Re-examination of the same dataset reveals a consistent tradeoff between the liver and brain proportions of a four-organ thermogenic core (kidney, heart, liver and brain), an inherent mechanism of thermoregulation that predates the emergence of permanent homeothermy. The ability of current models of brain size evolution to account for thermal homeostasis is limited by two common assumptions: that organ sizes evolve independently, and that the energy cost of the brain is proportional to the log ratio of brain mass with respect to body mass. Instead, arithmetic ratios provide direct experimental evidence for thermoregulatory constraints on brain size, as do organ cellular metabolic rates (*cMR*s). These are inferred without log transformation from the parallel adaptive increases in *MR*/kg and number of microRNA families (*mirFam*) that have accompanied major shifts in mammalian evolution. The *cMR* of the liver, the primary organ of gluconeogenesis, varies inversely with that of the brain, the primary consumer of glucose, a phylogenetic plasticity that appears to recapitulate the liver's unique developmental plasticity. With *mirFam* as a proxy for energy reliability, a positive feedback model of relative brain size detected adaptation to a low-energy regime among the smallest primates and, among the largest primates, adaptation to the physiological limit on the rate of heat dissipation.

INTRODUCTION

With the emergence of 24-hour homeothermy, Boreoeutheria crossed a threshold of energy reliability relative to earlier-branching placentals and marsupials, among which many orders exhibit some degree of heterothermy[1-2]. During the subsequent ecological diversification of Laurasiatheria and Euarchontoglires[3], carnivores, rodents and primates exhibited consistent increases in relative brain mass[4]. The evolution of resting metabolic rate (*rMR*), in concert with that of thermal conductance (C_{min}), allowed Boreoeutheria to occupy nearly every climatic zone regardless of size[5, 6], albeit with a narrower range of tolerance to heat than to cold[7-9], evidence that variation at the higher end of the ranges of body size and ambient temperature has been constrained by a physiological limit on the rate of heat dissipation (HDL)[10-12].

Simulation of two million years of African and Eurasian climate and habitats supports the view that *H. sapiens* evolved from *H. heidelbergensis* in southern Africa during an unusually warm period[13], consistent with the "Savanna hypothesis"[14-16]. The evolution of modern humans appears to have been conditioned by the HDL: among hunter-gatherers and contemporary indigenous populations, both resting and active *MR*s are actually

elevated compared with those of non-human apes[17-18], and these high rates of energy expenditure are

associated with uniquely human adaptations that promote heat loss, such as hairlessness, bipedalism, and a

10-fold increase in the density of eccrine sweat glands relative to the densities found in chimpanzees and

macaques[19]. The latter adaptation may have enabled early humans to run down game in long distance

pursuits[20], but may have also made them uniquely susceptible to dehydration[21], a marathon-enabling tradeoff.

Another potential mechanism to address the thermoregulatory challenge of a larger brain, first proposed four

decades ago, is a tradeoff in energy expenditure between the brain and another metabolically expensive

tissue, such as kidney, heart, liver or intestine[22-23]. Attempts to detect a tradeoff predicted by this "expensive

tissue hypothesis" were abandoned following publication of an influential report by Navarette et al (2011)

based on 100 mammalian species, including 23 primates, that found no significant anticorrelation among the

body mass-independent residuals of seven metabolically expensive organs[24]. However, re-examination of the

same dataset reveals that, of six possible tradeoffs among the four most highly thermogenic organs (kidney,

heart, liver and brain), the proportion of liver mass is strictly anticorrelated with that of the brain. Since this

tradeoff is conserved across 104 placental and 4 marsupial species, it represents a previously unrecognized

plasticity in the control of mammalian organ size[25], an inherent mechanism of thermoregulation that predates

the emergence of permanent homeothermy.

Current models of brain size variation have accounted for phylogenetic, but not developmental

non-independence. Furthermore, we demonstrate that the relative energy consumption of the brain is

proportional to its arithmetic ratio with respect to body size rather than the log ratio, as commonly assumed,

yielding direct experimental evidence for a thermoregulatory constraint on brain size. We develop a model of

organ rates of thermogenesis free of these assumptions based on the parallel adaptive shifts in resting

metabolic rates (rMRs) and number of microRNA families that have characterized major stages in mammalian

evolution: placentation, permanent homeothermy and the divergence of Catarrhini[12].

MirGeneDB, the manually-curated library of microRNA genes[26-27], has previously served as the reference

microRNA database in a wide range of comparative[28-30], phylogenetic[31-32], developmental[33-34] and evolutionary

studies[35-36]. Carnivores and primates are characterized by higher body mass ($Mbdy$) and body temperatures

(T_b) and faster resting metabolic rates (rMRs) than marsupials or earlier-branching placentals

(Atlantogenata)[1-2], differences that coincide with an increased number of conserved microRNA families

($mirFam$)[26-27] and density of microRNA target sites[12]. The expansion of brain size in the hominid lineage also

depended on a significant increase in rMR[17-18], and their clade rank order corresponds to the rank order of

mirFam; conversely, the divergence of smaller- from larger-bodied clades has been repeatedly associated with

decreased *mirFam*: Glires from primates, Strepsirrhini from Haplorhini and Callitrichinae from Platyrrhini[12, 26-27].

Thus, each increase in the range of *mirFam* has defined a new range of variation of *rMR* and *Mbdy* and,

consequently, the potential range of brain size variation, reflecting two potential undelying mechanisms.

Cassidy et al (2019)[37] investigated the nature of the relationship between *rMR* and *mirFam* during *Drosophila*

development. While differentiation of the eye depended on microRNA activity in wild type *Drosophila*,

development proceeded normally without any microRNAs when flies were engineered to have slower *MR*s,

thereby doubling the time to eclosion. The authors reasoned that the normal unfolding of eye development

must depend on the transient expression of transcription factors and the timely decay of their concentrations,

implying that flies with wild type *MR*s and rates of protein synthesis are dependent on microRNA-mediated

repression in order to maintain developmental synchrony[37]. By extension, the adaptive increases in *mirFam*

that have accompanied mammalian shifts to faster *rMR*s may have also served to maintain synchrony among

gene regulatory networks (GRNs). During mouse cranial neural crest differentiation, for example, specification

of neuronal, as opposed to ectomesenchymal cells, depends on prolonging a transient period of chromatin

accessibility in neuronal stem cells, an outcome of the repression of chromatin condensation by *Mir*-302[38].

Furthermore, since intrinsic protein expression noise increases with the rate of translation[39], metabolic

efficiency as well may have depended on matching faster *rMR*s with increases in global microRNA activity:

protein synthesis drives 20% of cellular metabolic rate[40] and global repression by microRNAs has been

estimated to decrease the overall rate of translation by ~ 25%[41-44].

We postulated that the kidney, heart, liver and brain together serve the role of thermogenic "Core" predicted

by a biophysical model of homeothermy[45] and tested the following predictions:

i) differences in measured total energy expenditure among Hominoidae will reflect differences in the

proportion of the brain mass and brain *rMR* in the Core;

ii) the arithmetic ratio of relative brain size will decrease with body size and evolutionary reductions in body

size will be accompanied by increases in relative brain size, irrespective of phylogeny;

iii) Core organ cellular *MR*s (*cMR*s), optimally scaled with respect to body mass and *mirFam*[12], will exhibit a

joint rate of thermogenesis that scales as predicted by the biophysical model of endothermy ($\sim M^{0.50}$)[45].

iii) phylogenetic rates of evolution of liver and brain *cMR*s will reveal adaptive tradeoffs accompanying the

divergence of clades between larger- and smaller-bodied mammals.

Figure 1. The proportion of brain mass in the thermogenic Core varies reciprocally with that of the liver.

A. Variable rate phenogram of body size variation among Therian mammals. The color scale represents the five-fold range of $\log_2$ mass. Branch lengths were obtained by Bayesian inference with the expectation of a Brownian motion model of trait evolution[46-47]. Differences in root-tip branch length from the ultrametric tree[48] represent either accelerated or decelerated rates of trait evolution. In the case of three outliers, branch lengths were arbitrarily shortened by 50% (*Ondatra zibethicus*, *Mustela nivalis*) or 80% (*Elephas maximas*).

B. Mass proportion of each organ relative to the combined Core mass (*Mcor*). *Mkdn*, *Mhrt*, *Mlvr* represent the mass of the kidney, heart, liver and brain, respectively. The mass of each organ was divided by the combined mass of the four thermogenic organs in each species. Dashed lines indicate species with extreme ratios of brain mass/liver mass.

We demonstrate that the *mirFam*-based cellular *MR*s (*cMR*s) of the kidney, heart and brain vary directly with

body mass, while liver *cMR* varies inversely with mass. Since the brain is the primary consumer of glucose

and the liver is the primary organ of gluconeogenesis, these tradeoffs are consistent with a positive feedback

model of brain size evolution governed by energy reliability, as has been previously suggested[4, 49].

RESULTS

The nature of the 108-mammal dataset of organ mass and metabolic rate

For ethical and technical reasons there is a paucity of direct measurements of their oxygen consumption of

the four primary thermogenic organs. Wang et al (2012)[50] derived allometric models based on measurements

of organ metabolic rates in the human, rat, rabbit, cat, and dog[50-52], and applied them to the organ mass

dataset of Navarette et al (2011)[24], the same dataset that provided the basis for refuting the Expensive Tissue

Hypothesis. Together with a term for residual thermogenesis, Wang et al (2012) fit these four organ *rMR*s to an

additive model of whole-body *rMR*: *rMR* = 70 x $M^{0.75}$. Henceforth, we refer to this dataset of measured organ

sizes and predicted *rMR*s as the "Wang dataset"[50].

Tradeoffs among Core organs verify the expensive tissue hypothesis

Using the Wang dataset[50], we obtained the organ/Core ratios of the measured mass of each thermogenic

organ in 108 therian mammals (Figure 1-B). Of the six possible tradeoffs among four thermogenic organs, the

most highly conserved was that between brain and liver, with a dampened tradeoff between heart and kidney.

Brain/Core mass ratios ranged from 0.04 (pig) to 0.36 (human) to 0.70 (squirrel monkey), while Brain/Liver

ratios ranged from 0.05 (opossum) to 0.76 (human) to 1.49 (squirrel monkey) (Figure 1-B). The variation in the

proportions of the Core organs bore no apparent relationship to either body size or phylogenetic clade with

the exception of the heart, whose relative proportion was low among the smallest rodents and high among

the largest carnivores (Figure 1).

A tradeoff between liver and brain mass was also readily demonstrable in log-transformed data, provided that

these were regressed on Core mass (Figure 2-B), rather than body mass (Figure 2-A): the Core

mass-independent residuals for liver and brain were highly anticorrelated (R^2 = 0.62, *P* = 2.32e-23) (Figure

2-B). The inability of the conventional analysis of brain size evolution, the phylogenetic regression of log *Mbrn*

(brain mass) on log *Mbdy* (body mass), to detect such a highly conserved adaptation prompted us to

investigate further the implications of two key assumptions of conventional phylogenetic comparative models.

Figure 2. Variation of liver and brain mass relative to either whole body mass or the mass of the four-organ Core.
A. Liver mass residuals with respect to body mass were plotted against brain mass residuals.
B. Liver mass residuals with respect to Core mass were plotted against brain mass residuals.
All 108 mammals in the dataset are shown, but the regressions represent only 88 carnivores, Glires and primates.

a) While these methods correct for phylogenetic non-independence, they do not take into account developmental non-independence, *i.e.* the gene regulatory programs that ensure that mammalian organ sizes track body size during development[25] (Supplementary Figure S1). For this reason, models of brain size standardized to body size will be blind to tradeoffs in the residual variation among the four thermogenic organs within the Core. Since whole-body *rMR* is the sum of organ *rMR*s, it is likewise not independent of body size, a shortcoming that is compounded when *rMRbdy* is treated as an independent variable in models of the relationship between log *Mbrn* and log *Mbdy*[4].

b) An adaptation that releases a constraint on the rate of heat dissipation may allow a diverging clade to occupy a higher range of body mass, while the opposite adaptation might occur if the energy supply is no longer sufficient to maintain thermal homeostasis. For example, a recent model of stabilizing selection revealed two adaptive shifts in the evolution of primate brain size, corresponding to the divergence of larger-bodied Catarrhini from Haplorhini, and the subsequent divergence of smaller-bodied Callitrichinae from Platyrrhini (Figure 1)[54]. The resulting ranks of brain/body *arithmetic* ratios, Callitrichinae > other Platyrrhini > Cercopithecidae, were consistent with a disproportionate reduction in body mass (Figure 1, Inset), but contrary to the rank order of log ratios (Cercopithecidae > other Platyrrhini > Callitrichinae) (in both cases, $P \cong 0.0$ for Callitrichinae *versus* Cercopithecidae:

Dunnett's Method with control). This interpretation is generalized by the arithmetic mass ratio of the

4-organ thermogenic Core (*Mcor/Mbdy*), which decreases strictly as a function of body size,

consistent with a geometric constraint on the rate of heat dissipation (Figure 3).

Figure 3. Evidence that clade shifts in the range of body mass represent thermoregulatory adaptations.

The main curve describing brain size in 1500 mammals is derived from the quadratic model of Venditti et al (2024)[4].

The circles depict adaptive shifts in the evolution of primate brain size identified by Grabowski et al (2023)[54].

The pink dotted line and confidence band represent *Mcor/Mbdy*, the arithmetic ratio of the combined mass of the

"Core" thermogenic organs to the body mass of 105 Carnivora and Euarchongotlires[50] (R^2 = 0.65, P = 2.23e-25).

Inset: Brain/body mass ratios in Callitrichinae (yellow) > other Platyrrhini (blue) > Cercopithecidae (orange),

consistent with a disproportionate reduction in body size (P = 0.0266 and 0.0, Dunnett's Method with control).

Data for Callitrichinae (N = 18), other Platyrrhini (N = 37), and Cercopithecoidea (N = 83): Grabowski et al (2023)[54].

c) The curvature of the relationship between log *Mbrn* and log *Mbdy* is *prima facie* evidence of
adaptation to thermoregulatory constraints, but owing to the limitations enumerated above, sample
sizes on the order of hundreds of species are required detect a departure from linearity that is
statistically significant[4, 53]. A practical implication of this insensitivity is that unless the allometric model
can resolve a 3% difference in slope ($M^{0.10}$) in the range $M^{0.33}$ - $M^{0.66}$, a value that is likely to be within
the range of measurement error, it will overlook a variation of 7% in the relative energy consumption of
the brain, assuming that it is proportion to the ratio *Mbrn/Mbdy*.

The combined mass of four thermogenic organs constitutes only 6% of mammalian body mass[50], yet
generates between 30 - 80% of the metabolic heat (55 ± 11.8%)[10]. The most significant limitation of current
allometric models is that the evolution of brain size is treated in isolation from that of the other thermogenic
organs. While developmental rules dictate that organ sizes track body size[25] (Supplementary Figure S1),
geometric constraints dictate that core thermogenesis decrease as body size increases (Figure 3).

Differences among Hominoidae in the measured rates of energy expenditure reflect tradeoffs in the
Core proportions of brain and liver mass and rMR
In order to assess the relevance of tradeoffs in the Core model to the evolution of brain size in the hominoid
lineage, we compared the brain/Core and liver/Core ratios of mass and *rMR* with field measurements of the
total energy expenditure (TEE) of orangutans and chimpanzees[17]. Using the double-labelled water method
together with physical activity budgets, these authors found that the TEE of human males was 67% higher
than that of *Pongo* and 25% higher than that of *Pan*, after correction for mass[17]. These differences mirrored a
comparison of Core proportions between a set of three human samples *versus* two other hominoid species
(*Symphalangus syndactylus* and *Hylobates concolor*) in the Wang dataset[50]: the brain/Core mass ratio was
elevated 40% among humans, while the brain/Core *rMR* ratio was 68% higher (*P* = 0.0014 and 0.0004,
respectively). These changes were offset by decreases of 16.4% in the liver/Core mass ratio and 29.6% in the
liver/Core *rMR* ratio (*P* = 0.0033 and 0.0007, respectively; ANOVA with pooled t-test).

The Inference of thermogenic organ cMRs
Previously, we developed an alternative to the log-based method for the estimation of specific *MR* (*sMR*) by
fitting the distribution of rMR/M^{exp} to the distribution of microRNA families (*mirFam*), a proxy for the rate of
energy turnover of a species[12]. Two advantages of this method are the greater resolution of the original
(non-log transformed) values of *rMR* and the independence of *mirFam* from the developmental control of
organ size. Thus, in contrast to the conventional estimation of a fixed *sMR* from the slope of log *rMRbdy* on

log *Mbdy*, the distribution of organ *cMR*s is matched to the distribution of *mirFam* while controlling for the

overall rate of thermogenesis, which is predicted to scale as $M^{0.50}$ [45].

Figure 4. Inference of organ cellular *MR*s (*cMR*s): classification by *mirFam* range or phylogenetic clade.

A. 50 Euarchontoglires were classified into bins according to natural divisions in the distribution of *mirFam*:

140-162 (Glires), 164-182 (Glires, Lemuriformes, Platyrrhini), and 194-210 (Catarrhini). *P* values represent regressions

on Bin x √Mbdy (ANCOVA). Note that the interval $M^{0.70}$ - $M^{0.80}$ defines a zone of minimal correlation for the Core *cMR*,

centered at $M^{0.75}$, which is also the midpoint of the dynamic ranges of liver and brain *cMR*.

B. The same analysis for species classified instead by phylogenetic clade (Glires, Lemuriformes, Platyrrhini, Catarrhini).

Comparison of potential *cMR*s with allometric exponents across the range of 0.65 - 0.85 revealed opposite

trends for liver, on the one hand (decreasing *P* value/increasing correlation), and for kidney, heart and brain,

on the other (increasing *P* value/decreasing correlation), whether species were classified by range of *mirFam*

or phylogenetic clade (Figure 4-A and B, respectively). Notably, the anticorrelation that emerges from this

mirFam-based partitioning of organ rates of thermogenesis is independent of the anticorrelation revealed by

the analysis of organ/Core mass ratios (Figures 1, 2). Assuming an operating range of $M^{0.70}$ - $M^{0.80}$, we

assigned each organ a *cMR* corresponding to the lower limit of the curve of *P* values, hence:

*cMR.**70**kdn*, *cMR.**70**hrt*, *cMR.**70**brn* and *cMR.**80**lvr*

Whole body *cMR* primarily tracked liver *cMR*, while Core *cMR* exhibited a symmetric profile with a zone of

minimal correlation centered at $M^{0.75}$ (Figure 4-A). Since $M^{0.75}$ was also the optimal allometric scaling in a

model of stabilizing selection of mammalian *cMR* with respect to *mirFam*[12], this zone of minimal constraint on

the rate of Core thermogenesis may be interpreted as an evolutionary adaptation for thermal homeostasis.

The same trends were observed when species were classified instead by phylogenetic clade, but in this case

the curves for Core and whole body *cMR* were both shifted toward higher allometric exponents (Figure 4-B).

In order to rule out the possibility that the opposite patterns of correlation of liver and brain *cMR* are unique to

the sample of variation represented by the Wang dataset, we applied the same allometric models to a

completely independent set of *rMR*s measured in 30 primates[10]. Core *rMR* was inferred from its relationship to

body *rMR* and then liver and brain *rMR* were inferred from Core *rMR* (Supplementary Table S3).

Figure 5. Core and organ *rMR*s can be inferred from measured primate *rMR*s.
Based on the allometric models for 22 primates, excluding the human[13], the following chain of inference was applied
to the measured *rMR*s of 30 primates[10] (Supplementary Table S3): organ mass ← organ *rMR* ← Core *rMR* ← Body *rMR*.
A. Anticorrelation between organ/Core *rMR* ratios from independently derived liver and brain *rMR*s.
B. Liver and brain *cMR*s exhibit opposite patterns of correlation with *mirFam* x $\sqrt{Mbdy}$.
Core *cMR*s exhibit minimal correlation with *mirFam* x $\sqrt{Mbdy}$ in the range $M^{0.70}$ - $M^{0.80}$ (compare with Figure 3).

As in the case of organ/Core mass ratios, the *rMR*/Core ratios of liver and brain were anticorrelated in all but

the largest primates (Figure 5-A). Furthermore, liver and brain *cMR*s exhibited opposite correlation patterns

with respect *mirFam* and $\sqrt{M}$, and Core *cMR* exhibited minimal correlation when scaled to $M^{0.75}$ (Figure 5-B).

Thus, the observed anticorrelations between liver and brain are a consistent outcome of the Wang allometric

models, regardless of species sampled or whether the source data represent organ mass or whole-body *rMR*.

Organ cMRs, body mass, and thermal conductance

Plotting organ *cMR*s as a function of body size again revealed an inverse relationship between *cMR.80lvr* and

the *cMR.70*s of the other three organs: while these increased directly with size, *cMR.80lvr* was strictly

anticorrelated with body mass (Figure 6-A). A finer distinction was also evident: while the slope of *cMR.80lvr* was unchanged between smaller-bodied Glires and larger-bodied primates, the slopes for kidney, heart and brain *cMR.70* were significantly higher in the larger-bodied clade (Figure 6-A).

Figure 6. Core organ *cMR*s as a function of body size or thermal conductance (C_{min}).
A. Slope ratios between primates and Glires were compared by ANCOVA (figure legend) (N = 51).
B. Organ rates of thermogenesis as a function of thermal conductance among Boreoeutheria (*N* = 20).
T_a = ambient temperature. Shaded bands: confidence intervals (CI).

Not surprisingly, the same ranking of slopes (heart > kidney > brain) and anticorrelation with liver *cMR.80* was obtained when organ *cMR*s were regressed on the thermal conductance (log C_{min}) of 20 Boreoeutheria (Figure 6-B). C_{min} is an extensive property that depends primarily on total body surface area and secondarily, on the properties of the insulating layer[45, 55]. When Boreoeutheria were ranked according the whole-body rate of heat generation (*cMR.75*), *mirFam* covaried with body size, core thermogenesis (*cMR.53cor*) and thermal conductance (log C_{min}), independent of phylogeny (Supplementary Figure S2-A). As in the case of body size, ranking Boreoeutheria by C_{min} revealed tradeoffs between liver *cMR.80*, on the one hand, and both brain and heart *cMR.70*, on the other (Supplementary Figure S2-B), consistent with the coordinated evolution of body temperature, metabolic rate and thermal conductance[5-6, 56].

Thermodynamic and phylogenetic relevance of Core organ cMRs

We tested two predictions, the first stemming from the allometric model that was the basis of the *rMR*s in our dataset[50], and the second from the thermoregulatory core model of Porter and Kearney (2009)[45]:

i) The joint **sMR** of the thermogenic organs should scale as $M^{0.75}$, since the individual organ *rMR*s were inferred from an additive model assuming that whole body *sMR* = 70 x $M^{0.75}$, and we ignored the residual *rMR*, which averaged only 4.5% of that of the liver in mammals < 75 kg[50];

ii) The joint **cMR** of the four most highly thermogenic organs, although delocalized, should scale approximately as the geometrically centered, thermogenic core of the biophysical model *i.e.* ~ $M^{0.50}$.

Figure 7. Scaling of the Core *cMR* and body *sMR* by linear combinations of either the *mirFam*-based *cMR*s or the log-based *sMR*s of the four thermogenic organs (*N* = 22 primates).
A. % variation (R^2) of linear models of Core *cMR* or body *sMR* composed of the four organ *cMR*s or *sMR*s. *VIP* values (*variable importance of projection*) indicates the number of organs upon which each model depends[57].
B. Comparison of the joint *cMR* of combinations of *cMR.70 - cMR.85* among the four thermogenic organs. "*c*" is the organ *cMR* allometric exponent. Exp_{opt} is the optimal allometric exponent of the additive model.

Applying partial least squares regression[58], we generated additive models in order to predict the optimal scaling of a joint Core *cMR* composed of either log-based *sMR*s:

sMR.Xcor = (rMRkdn + rMRhrt + rMRlvr + rMRbrn) / (Mkdn + Mhrt + Mlvr + Mbrn)^{exp}

= (a' • sMR.**90kdn**) + (b' • sMR.**87hrt**) + (c' • sMR.**67lvr**) + (d' • sMR.**83brn**)

or mirFam-based *cMR*s:

cMR.Xcor = (rMRkdn + rMRhrt + rMRlvr + rMRbrn) / (Mkdn + Mhrt + Mlvr + Mbrn)^{exp}

= (a • cMR.**70kdn**) + (b • cMR.**70hrt**) + (c • cMR.**80lvr**) + (d • cMR.**70brn**)

The log-based joint *sMR* exhibited a single sharp peak at $M^{0.75}$, corresponding exactly to the whole-body *sMR*

of both the biophysical model[45] and the Wang allometric model from which the organ *rMR*s were derived[50].

The composite *cMR* model exhibited two peaks, one on either side of an inversion of Core *cMR* slopes with

respect to $\sqrt{M}$: a broad peak centered at $M^{0.53}$, similar to that predicted by the biophysical model of core

thermogenesis[45], and a second peak between $M^{0.80}$ - $M^{0.85}$ (Figure 7-A). Importantly, the *cMR* and *sMR* models

exhibited critical differences in *VIP* statistics (*Variable Importance of Projection*), a measure of the model

dependence on each variable[57], analogous to the loadings of principal components. The *cMR* models

depended on all four organ *cMR*s across the entire range, with the exception of a zone of minimal correlation

($M^{0.75}$ - $M^{0.80}$), where they depended only on that of liver. In contrast, the *sMR* model depended only liver, heart

and brain ($M^{055.}$ - $M^{0.65}$) or only liver and kidney ($M^{0.70}$ - $M^{0.90}$) (Figure 7-A, bottom).

We applied an additional test of the relevance of these estimated organ *cMR*s by comparing the joint *cMR*s

obtained by permuting combinations of estimated organ *cMR*s, yielding four models with *VIP* = 4 (Figure 7-B).

Of these, we found two combinations that were able to resolve Platyrrhini from Lemuriformes in an

unsupervised classification of major primate clades[59]: either *cMR.80* or *cMR.85* for liver, together with

cMR.70 for heart, kidney and brain. When liver *cMR* was varied between *cMR.75* - *cMR.85*, the original set of

mirFam-based *cMR*s yielded the lowest value for the Bayes information criterion (BIC) (Supplementary Figure

S3), hence the following optimal model of the Core *cMR*:

cMR.53cor = 0.53 · **cMR.70kdn** + 0.69 · **cMR.70hrt** + 0.37 · **cMR.80lvr** + 0.09 · **cMR.70brn**

In the context of the evolution of thermal homeostasis, the coefficient of each organ *cMR* in this model

indicates the degree to which it has been constrained by natural selection, *i.e.* the smaller the coefficient, the

greater the permitted variation of the organ *cMR* since clade divergence.

The original biophysical model for thermoregulation assumed that metabolic heat emanates from the

geometric center of an ellipsoid endotherm. Depending on the wind condition or the presence of fur, the

authors derived a single allometric exponent between 0.46 - 0.58 in order to scale the rate of core

thermogenesis across the entire range of mammalian body size[45]. Since our Core model is delocalized among

four thermogenic organs and makes no geometric assumptions, we expected lower allometric exponents and

higher rates of thermogenesis (rMR/M^c) in smaller mammals with higher ratios of surface area/volume.

The optimal allometric slopes of *cMR* models developed in Carnivora, Primata and Glires were $M^{0.60}$, $M^{0.53}$ and

$M^{0.40}$, respectively (Figure 8). The corresponding distributions of *Mcor/Mbdy* support the notion of geometric

constraints: Glires occupy the upper end of the range, suggesting an adaptation to maintain body

temperature, while primates and carnivores occupy the lower end, consistent with a limit on the rate of heat
dissipation (Supplementary Figure S4-A)[11].

332
333 **Figure 8**. Organ coefficients of additive models of Core thermogenesis in three major clades of Boreoeutheria.
334 Values in parentheses next to each clade indicate the optimal allometric exponent of each Core *cMR* model.
335

The coefficients in the Glires Core model present a striking contrast with those of carnivores and primates: a
reversal in sign of the coefficient for brain *cMR* in Glires, and a reversal of the relative magnitudes of the
coefficients for liver and brain (Figure 8). The Glires model thus penalizes the *cMR* of the brain, suggesting
adaptation to a disproportionate reduction of body size relative to brain size, leaving the arithmetic mass ratio,
Mbrn/Mbdy, higher in Glires than in primates or carnivora ($P < 0.001$, analysis of means) and contrary to the
rank order of the allometric slopes for log *Mbrn versus* log *Mbdy* (primates > Glires) (Supplementary Table S4).
342

343 ***Tradeoffs and adaptations in the rates of evolution of organ cMRs***
344 The preceding evidence suggests that Boreoeutheria have been able to maintain thermal homeostasis while
345 diversifying body mass by fine-tuning the relative sizes of the four Core organs in relation to that of the liver.
346 We applied Bayesian inference in order to investigate the historical timing of the corresponding tradeoffs.

Figure 9. Variable rate phenograms of Liver *cMR* (*cMR.80*) and the *cMRs of* Kidney, Heart and Brain (*cMR.70*).
Bayes factors were computed from the difference in log likelihood (*Lh*) of the 1% most likely trees and the trees with
the smallest non-zero sum of changes to branch length[12], *i.e.* those trees most similar to the reference time tree[48].

In agreement with the Core *cMR* models, we found opposite directions of evolution between the organ

*cMR.70*s that vary directly body size (kidney, heart and brain) and *cMR.80lvr*, which varies inversely with body

mass, both within and between major clades (Figure 9). These observations are supported by a

computational model that found that following an overnight fast, rates of liver gluconeogenesis, lipolysis and proteolysis were all elevated ~ 10-fold in mice, relative to the corresponding rates in humans[60].

Among primates, increased liver *cMR* accompanied the decrease in size of two of the smallest Platyrrhini, one of which (*Callithrix pygmaea*) may be heterothermic[61]. A clade of five small (< 1.0 kg) Platyrrhini exhibited an accelerated *decrease* in brain *cMR.70* paired with elevated levels of *cMR.80lvr*, consistent with adaptation to a low energy regime and greater reliance on gluconeogenesis. Among Glires as well, those species that underwent the most severe reductions in size exhibited the greatest increases in liver *cMR.80*; conversely, within Glires and Haplorhini, the large-bodied clades (Lagomorpha and Catarrhini) exhibited lower liver *cMR*s (Figure 8). Brain *cMR.70* increased in intensity and its rate of evolution accelerated with the divergence of Haplorhini, and then again in humans, in contrast to its stasis or decrease among Glires, Lemuriformes and smaller Platyrrhini (Figure 9). The relative levels of heart *cMR.70* generally tracked body size, with accelerated rates of heart evolution in the clades of the largest Glires (Lagomorpha) and primates (Catarrhini).

The divergence of humans was marked by the accelerated rate of evolution of the *cMR.70* of the kidney in addition to that of the brain (Figure 9), an observation that is noteworthy for two reasons:

a)   The 30% higher level of kidney *cMR* found in humans relative to other hominoid apes is not apparent when kidney *MR* is scaled according to the log-based slope ($M^{0.90}$): the corresponding value for the kidney *sMR* is 10% *lower* in humans than the average value among apes.

b)   The adaptive increase in *cMR.70kdn* is of particular interest since the ability to thermoregulate is highly dependent on water balance, as illustrated in a study of baboons whose fossils appear in the same arid horizons as the fossils of *Australopithecus*[62] and *Homo*[15]. When deprived of water and exposed to elevated ambient temperatures, free-ranging baboons developed a progressive hyperthermia (increasing core temperature) that was rapidly reversed upon rehydration[63].

DISCUSSION

Direct experimental evidence for a thermoregulatory constraint on brain size

It now appears well established that a single rule[4] governs the relationship between brain mass and body mass in mammals, irrespective of phylogeny, rather than multiple, clade-specific rules[64]. This paradigm comports well with the notion of a single developmental program in mammals to scale organ sizes to body size[25]. However, current phylogenetic comparative models have overlooked a thermoregulatory tradeoff embedded in this developmental program, an oversight the present study aims to correct. In order for the variation of a trait to be shaped by thermoregulatory constraints, it must reflect the relative energy cost *in vivo*,

i.e. the arithmetic proportion of mass or heat production, rather than the log-based allometric slope: while

log-based allometric slopes are positive, consistent with the developmental programming of organ sizes,

Core/body mass ratios decrease strictly as a function of size, consistent with geometric constraints.

Venditti et al (2025) recently demonstrated that, irrespective of phylogeny, a quadratic function best fits the

relationship between log brain mass and log body mass in 1,500 mammals, evidence for a universal

constraint on evolution of brain size at the upper end of any range of body mass[4]. Hypothetically, this

constraint could be either anatomic *i.e.* the diameter of the skull in relation to that of the birth canal, or

physiological, such as a limit on the rate of heat dissipation[11].

	Sample	Range Mbdy (g)	Range Ratio (%)	Likelihood	Model	β_1	β_2	R^2
	Low ECV/Mbdy (N = 15)	140 – 22,300	0.80 - 1.86	-5.31	$Log_2ECV = \beta_1 \cdot Log_2 Mbdy$	0.66 ± 0.077		0.85
				-3.08	$Log_2ECV = \beta_1 \cdot Log_2 Mbdy + \beta_2 \cdot Log_2 MR$	0.58 ± 0.079	-0.07± 0.060	0.84
	Middle ECV/Mbdy (N = 15)	140 – 14,150	1.02 – 2.26	-6.22	$Log_2ECV = \beta_1 \cdot Log_2 Mbdy$	0.69 ± 0.072		0.88
				-5.95	$Log_2ECV = \beta_1 \cdot Log_2 Mbdy + \beta_2 \cdot Log_2 MR$	0.70 ± 0.074	-0.04 ± 0.057	0.88
	High ECV/Mbdy (N = 15)	75 - 989	1.49 – 3.59	-3.19	$Log_2ECV = \beta_1 \cdot Log_2 Mbdy$	0.63 ± 0.067		0.88
				-1.88	$Log_2ECV = \beta_1 \cdot Log_2 Mbdy + \beta_2 \cdot Log_2 MR$	0.67 ± 0.069	0.07 ± 0.047	0.90

ECV/Mbdy Distributions (histograms shown with x-axis values 0.005, 0.01, 0.015, 0.02, 0.025, 0.03, 0.035, 0.04)

Table 1. Sliding window analysis of the interaction between relative brain size and whole-body metabolic rate.

The measured *rMR* data for 25 primate species from Clarke et al (2010)[10] were matched with endocranial volumes (*ECV*)

from Grabowski et al (2023)[54]. An exponential model of *rMR ~ Mbdy* was applied to the body mass values of the *ECV*

dataset. Bayesian phylogenetic regression was applied to three overlapping sets of 15 species ranked by *ECV/Mbdy*

ratio in order to test the dependence of Log_2ECV on Log_2Mbdy and Log_2rMR. The first bin in each distribution represents

five species. All statistics represent the median value of the 100 regressions with the greatest likelihood (top 10%). The

primate species included 6 Catarrhini, 5 Platyrrhini, 4 Lemuriformes and 10 Lorisiformes (Supplementary Table S6).

In order to test whether this constraint is related to the overall rate of thermogenesis, we inferred *rMR*s for the

body mass data in a dataset of endocranial volumes (*ECV*)[54] based on a separate set of *rMR*s measured in the

same primate species[10]. Bayesian regressions were then carried out to compare the dependence of log *ECV*

on log *Mbdy* and log *rMR* for 15 species in each of three, overlapping ranges of relative brain size

(*ECV/Mbdy*). As shown in Table 1, the range of *ECV/Mbdy* was inversely related to the range of body mass. In

each case, the model for the dependence of *ECV* on body mass was improved by the inclusion of the *rMR*

term. For the highest range of body mass and lowest brain/body ratios, the regression coefficients for the *rMR*

term were consistently (100%) negative, corresponding to 12.2% of the value of the *Mbdy* term. For the

smallest-bodied species with the highest brain/body ratios, the regression coefficients for the *rMR* term were

consistently (100%) positive, equivalent to 10.5% of the value of the *Mbdy* term. For the middle range of body

size and relative brain size, the regression coefficient for the *rMR* term fluctuated around 0, as was the case for the entire dataset of 25 species. Thus, after correction for phylogenetic covariance, relative brain size was inversely related to metabolic rate in the 15 largest species, *i.e.* those most likely to have adapted to a heat dissipation limit. On the other hand, the *rMR* term was a positive function of relative brain size in the 15 smallest species, consistent with the greater relative energy demand of the brain, and for whom their surface area and rate of heat loss may have required adaptation in order to maintain body temperature.

The developmental and phylogenetic plasticity of the liver

The brain and liver, each accounting for 20% of *bMR* in humans, exhibited highly conserved tradeoffs in mass and *cMR*, a plasticity that appears to be an inherent property of the mammalian evolutionary toolkit, not exclusive to homeotherms. While the negative slope for the dependence of liver *cMR* on body mass remained constant across clades, the positive slopes for kidney, heart and brain *cMR*s were significantly higher in primates than among Glires, suggesting that their combined arithmetic proportion has been adjusted relative to that of the liver in a clade-specific manner.

An inverse relationship between gluconeogenesis and the rate of body mass growth in humans implicates the liver as the pivotal partner in an ontogenetic tradeoff: brain glucose utilization peaks around age five, early in the extended period between weaning and sexual maturity, which corresponds to the period of slowest growth in body mass[65]. Based on the turnover of isotopically-labeled glucose, liver glucose production was proportional to brain mass over the entire lifespan, and therefore highest per body mass among young children[66]. Among prepubertal children (7 - 9 years of age), the rate of gluconeogenesis was still 50% higher than that found among adolescents[67]. The evolutionary plasticity of the liver appears to have co-opted the liver's inherent developmental plasticity: among mammalian organs, only the liver retains full regenerative potential[25]. If the evolution of liver size has been determined by the demand for glucose production, then situations of high energy reliability/low dependence on gluconeogenesis in incipient clades may have favored evolutionary decreases in liver size, such as those associated with the divergence of Lagomorpha from Glires, and of Catarrhini from Haplorhini. Conversely, smaller bodies and tradeoffs favoring the liver appear to signify adaptation to limited energy resources relative to population size, exemplified by the divergence of Glires from Euarchontoglires, Strepsirrhini from primates, Platyrrhini from Haplorhini, and Callitrichinae from Platyrrhini.

Hippo-YAP, the regulator of organ size in Bilateria[68], coordinates liver growth with the suppression of gluconeogenesis[69-72]. At low cell densities, Hippo signaling is normally suppressed, allowing YAP to localize in the nucleus and stimulate cellular proliferation, while at normal organ size and tissue density, YAP

translocation and activity are normally inhibited by Hippo[68]. The global underexpression of microRNAs that is frequently observed in tumor cells has been attributed to aberrant density-dependent regulation of the same Hippo-YAP signaling: constitutively active YAP suppresses microRNA levels in cells and tumors and post-transcriptionally induces oncogenic MYC protein[73-74]. By extension, the evolutionary plasticity of the liver implies that each species divergence involves a reset of the dynamic ranges for the interactions among Hippo-YAP signaling, organ size, and the rate of microRNA biogenesis.

A feedback model for brain size and energy reliability

We modeled a simple feedback loop based on the inverse association between larger brain sizes and *mirFam* and lower liver sizes and *cMR*s, assuming that the latter reflects decreased dependence on gluconeogenesis: a larger brain size leads to greater energy reliability which, in turn, facilitates the evolution of a larger brain, up to the limits imposed by anatomy and physiology, such as the diameter of the birth canal or the physiological limit on the rate of heat dissipation[10-12]. We applied a logistic feedback function to fit the distribution of *mirFam*, a proxy for energy reliability, to the primate brain size dataset of Grabowski et al (2023)[50]:

$$\frac{d\,(mirFam)}{d\,(\log V_{brn}/\log M_{bdy})} = r \cdot mirFam\,(1 - mirFam/mirFam_{max})\,,$$

where **r** is the growth rate, and the range of *mirFam* has been scaled to a range of 1.0.

As shown in Figure 10-A, log *Mbrn*/log *Mbdy* is a predictor of *mirFam,* rather than the converse, consistent with the notion that evolutionary increases in brain size tended to result in greater energy reliability. Since the distribution of *mirFam* can be fit to the mass log ratios of any thermogenic organ, we asked which relationship might be the principal target of selection. Since the ranges of *mirFam* are highly clade-specific (Supplementary Figure S5) it is not a suitable dependent variable for phylogenetic regression. Instead, we inverted the premise and analyzed the dependence of organ *cMR*s on *mirFam*, as the independent variable. All potential kidney and heart *cMR*s (*cMR.60 - cMR.80*) varied independently of phylogenetic distance, as did all but one liver *cMR* (*cMR.60lvr*) (Supplementary Table S5). In contrast, brain *cMR* variation with respect to *mirFam* was, in every case, secondary to phylogenetic distance (Pagel's *lambda* = 1.0); in other words, the divergence of primate clades coincided with changes in *mirFam* range specifically with respect to brain *cMR*.

A variable rate phenogram corresponding to this model was generated using ratios from the linearization of the logistic function[75], such that the color scale signifies rank order (Figure 10-B). Catarrhini exhibited the highest *mirFam* levels with respect to brain size, a relationship that evolved at the same rate as the underlying

Figure 10. Logistic model of the *mirFam* relationship to relative brain size and the primate heat dissipation limit.

A. Comparison of the logistic model with linear and power law models (*N* = 31, excluding the clade of
heterothermic Cheirogaleidae: *M. murinus* and *C. medius*).
Filled symbols: circles = Hominidae; rectangles and diamond: documented (blue) and suspected (purple) heterotherms.

B. Variable rate phenogram based on the linearized values of the logistic model: when ranked in ascending order,
the ratio of *mirFam* values to the cumulative sum decreases in a linear manner (Hubbert, 1982)[75].

C. Comparison of the logistic models with the maximal rate of thermogenesis predicted from whole-body mass and
T_b (Allen et al, 2006)[76] and scaled to the same range as *mirFam*. Shading represents the CI (cubic spline method).

D. Variable rate phenogram based on the difference between predicted *mirFam* and model HDLs. The resulting
distribution was centered and the outlier clade of dwarf and mouse lemurs was assigned an intermediate value of 50.

rate of genomic evolution. In contrast, the three smallest-bodied clades, *Cheirogaleidae, Galago and Callithrix* (marmosets) exhibited lower *mirFam* levels and an accelerated descent in rank (Figure 10-B). The latter observation is consistent with reversion of the brain size allometry of Callitrichines to one more similar to that of Strepsirrhini than that of Platyrrhini[64]. *Callithrix pygmaea* may also be heterothermic[61], and the *Callithrix* clade also exhibited marked increases in the level and rate of evolution of liver *cMR.80*, suggesting adaptation to a regime of reduced energy reliability that is characteristic of low calorie-density folivory or insectivory[77].

Supported by the previous observation that larger-bodied primates and Carnivora occupy the lower end of the range of Core/body ratios, we postulated that the upper plateau of the logistic model would correspond to the physiological limit on the rate of heat dissipation[10-12]. Furthermore, when primate species are ranked by size, *mirFam* varies reciprocally with inverse body temperature, $1/T_b$ (°K); thus, the seemingly random variation of T_b within each clade appears to be tied to the variation in *mirFam*, except among the smallest heterotherms (Supplementary Figure S6). The conserved nature of this reciprocity lends itself to the following formulation of a model of the maximum heat dissipation limit (HDL). The relationship between whole body *rMR* and T_b can be described by the Arrhenius-Boltzmann function, $e^{-E/kT}$, where E represents the mean activation energy of the respiratory complexes and k is the Boltzmann constant[76]; when corrected for total body mass, this function represents the maximal possible rate of thermogenesis and therefore heat dissipation, regardless of phylogeny or thermoregulatory adaptations (Figure 10-C).

In order to detect adaptations that occurred in relation to this limit, we generated variable rate phenograms that depict the direction and rate of evolution of the *difference* between the values of the HDL model and the logistic model (Figure 10-C). In this case, the color scale signifies proximity to the HDL. Interestingly, two of the clades that exhibit accelerated evolution toward the HDL represent the highest relative brain size among Platyrrhini (*Saimiri*) and lowest relative brain size among Lorisiformes (*Galago*), suggesting thermoregulatory adaptation at the extremes of variation (Figure 10-D) (Supplementary Table S2). The divergence of Catarrhini was associated with rapid evolution toward the HDL that was subsequently reversed in two cases: the divergence of the human and colobus monkey, species that again represent the extremes of relative brain size within their clade (Figure 10-D). The apparent "cooling" in the human branch appears to reconcile the disproportionate increase in brain size with our species' unique adaptations for heat dissipation[18].

Our logistic model assigns the ancestral primate a high rank for energy reliability and a low rank relative to the HDL, a propitious combination for the founder of a new order of mammals. In many respects this model fulfills the recent call for an explanation of the distinct pattern of brain size evolution in primates "in terms of the

release of a constraint or an adaptive shift that initiated escalating feedback between brain and behavior"[4],
assuming that such behavior results in greater energy reliability, hence reproductive fitness. Since this logistic
model is agnostic regarding underlying mechanisms, it can also serve as a null hypothesis for comparison
with the evolution of other candidate mechanisms of brain size evolution. More significantly, discovery of the
tradeoff proposed four decades ago in order to account for thermal homeostasis during the evolutionary
expansion of brain size[22-23] depended on the definition of variables that are more sensitive to thermoregulatory
constraints: arithmetic ratios, rather than log ratios or log-based slopes, and cellular MRs matched to an
index of energy turnover and independent of developmental control, rather than log-based specific MRs.

SYNTHESIS

The adaptive value of repression by individual microRNAs operating in gene regulatory networks has been
attributed to their ability to canalize phenotype expression by buffering the effects of environmental or allelic
variation[78-79] and to set thresholds for cell fate determination, or maintain the identity of differentiated cells, by
defining ranges of protein expression[80-81]. Major stages in the diversification of animal cell types,
morphologies, and complexity of behavior have corresponded to significant expansions of microRNA and
target repertoires[82-86]. For example, the emergence of placental mammals is marked by the appearance of 13
novel implantation-related microRNA families[33], and the cognitive sophistication of the genus *Octopus* has
been linked to 34 novel microRNA families that are preferentially expressed in neural tissues[82]. However, these
post hoc associations do not shed light on the preconditions of repertoire expansion, nor on how such
enhanced traits may be related to a higher range of *mirFam*, as opposed to action of individual novel
microRNAs[82]. The consistent pairing of adaptive shifts in MR/kg with expansions of the microRNA repertoire
implies a prerequisite of a high level of energy reliability in an incipient clade. Given that *mirFam* losses have
occurred much less frequently than gains[31], these observations suggest the operation of a ratchet mechanism
that selects, among novel microRNAs, those that confer greater energy reliability. An analogy may be drawn to
the fate of the duplicated genes of transcription factors (TF): owing to their established specificities, they have
been much more likely to be retained than the paralogues of non-TF genes[87].

METHODS

Physiological Data

The masses of the four thermogenic organs were obtained from the 111-species dataset of Wang et al
(2012)[50], originally derived from the compilation of Navarette et al (2011)[24]. Only in humans have the metabolic
rates of the four thermogenic organs been measured directly[51]. Otherwise, they have been inferred by
conventional allometric scaling based on organ mass of rat (all four organs), dog (liver, brain and kidney), cat

and rabbit (liver)[50, 52]. Relative brain sizes for the logistic models were obtained from Grabowski et al (2023)[54],

Smaers et al (2021) (human)[64], and estimated for Dermoptera[88].

Values for thermal conductance (C_{min}) were obtained from the dataset of Fristoe et al (2015)[6], with substitution

of congeneric species from the dataset of Wang et al (2012)[50]. Where C_{min} was available for two or more

congeneric species, we chose the one with the mass nearest to the species in the Wang dataset. Body

temperatures were obtained from the AnAge database[89] and other sources.

MicroRNA Families, *Binning*

Complements of *conserved* microRNA families were obtained from the most complete assembly for each

species using the covariance model MirMachine[90], trained on the manually-curated database of microRNA

genes, MirGeneDB2.1[26] (Supplementary Tables S1 and S2). For Scandentia, Lagamorpha and Rodentia,

mirFam refers to the number of families common to Euarchontoglires. For Lorisiformes, Lemuriformes and

Platyrrhini, *mirFam* refers to the number of families shared with Catarrhini. For the original regression of organ

*cMR*s on *mirFam* and organ size (Figure 3-A), *mirFam* was binned according to natural divisions in its

distribution: 141 - 161 (Scandentia and Glires), 165 - 181 (Glires, Lemuriformes and Platyrrhini), and 194 - 208

`(Catarrhini) (Supplementary Figure S5).

Phylogenetic Trees and Variable Rate Phenograms

Using Mesquite © (2021) we derived ultrametric trees for 108 Theria, 51 Euarchontoglires or 24 primates from

the mammalian phylogeny of Alvarez-Carretero et al (2021)[48]. Variable rate trees and ancestral node values

were inferred using BayesTraits (version 4)[91-92] (software available at www.evolution.rdg.ac.uk). Phenograms

were generated using Phytools[93].

For variable rate trees, Bayes factors were computed from the difference between the median of the top 1%

likelihoods and the median likelihood of the 1% of trees with the smallest non-zero sum of changes to branch

length, *i.e.* those trees most similar to the reference time tree, as previously described[12]. In the case of those

variable rate trees based on linearization of the logistic function[75], artifactual changes of slope at the extremes

were avoided by the addition of dummy values to either end of the range.

Statistics

Linear, logistic, and partial least squares regressions[57-58] and normal mixture clustering[59] were carried out and

visualized using JMP16®. All R^2 values were adjusted for the number of predictors. Other than linear and

logistic regressions, curves were fit empirically using the cubic spline method, with the confidence intervals indicated by shading[94-95]. Phylogenetic linear regressions were carried out using the R: CAPER package[96].

Limitations

　The three principal limitations of this study are:

　i) the indirect origin of the mammalian organ *rMR*s, as described above (*Physiological Data*)[50, 52].

　ii) the small number and variable quality of the body temperature data in the AnAge database[89];

　iii) underestimation of the number of conserved microRNA families by the MirMachine algorithm[90], relative to the reference, manually-curated database[26-27] (currently MirGeneDB 3.0). Excluding Chiroptera, the correlation of computed values with MirGeneDB 3.0 is between 90 - 100% for genomes with scaffold N50 > 10,000[90].

　None of these factors is expected to affect the qualitative interpretation of our results.

REFERENCES

1. Boyles JG, Thompson AB, McKechnie AE, Malan E, Humphries MM and Careau. 2013.
A global heterothermic continuum in mammals. *Global Ecol Biogeogr* **22**: 1029-1039.

2. Ruf T and Geiser F. 2015. Daily torpor and hibernation in birds and mammals. *Biol Rev* **90**: 891-926.

3. Clavel J and Morlon H. 2017. Accelerated body size evolution during cold climatic periods in the Cenozoic.
Proc Natl Acad Sci USA **114**: 4183-4188.

4. Venditti, C, Baker J and Barton RA. 2024. Co-evolutionary dynamics of mammalian brain and body size.
Nat Ecol Evol **8**: 1534–1542.
5. Naya DE, Spangenberg L, Naya H and Bozinovic F. 2013. Thermal conductance and basal metabolic rate are part of a coordinated system for heat transfer regulation. *Proc Roy Soc B* **280**: 20131629.

6. Fristoe TS, Burger JR, Balk MA, Khaliq I, Hof C and Brown JH. 2015. Metabolic heat production and thermal conductance are mass-independent adaptations to thermal environment in birds and mammals.
Proc Natl Acad Sci USA **112**: 15934-15939.

7. Araújo MB, Ferri-Yáñez F, Bozinovic F, Marquet PA, Valladares F and Chown SL. 2013.
Heat freezes niche evolution. *Ecol Letters* **16**: 1206-1219.

8. Khaliq I, Hof C, Prinzinger R, Böhning-Gaese K and Pfenninger M. 2014. Global variation in thermal tolerances and vulnerability of endotherms to climate change. *Proc Roy Soc B* **281**: 20141097.

9. Bennett JM, Sunday J, Calosi P, Villalobos F, Martínez B, Molina-Venegas R, Araújo MB, Algar AC, Clusella-Trulas S, Hawkins BA, Keith SA, Kühn I, Rahbek C, Rodríguez L, Singer A, Morales-Castilla I and Olalla-Tárraga MA. 2021. The evolution of critical thermal limits of life on Earth. *Nature Comm* **12**: 1198.

10. Clarke A, Rothery P and Isaac NJ. 2010. Scaling of basal metabolic rate with body mass and temperature in mammals. *J Anim Ecol* **79**: 610–619.

11. Speakman JR and Król E. 2010. Maximal heat dissipation capacity and hyperthermia risk: neglected key factors in the ecology of endotherms. *J Anim Ecol* **79**: 726-746.

12. Fromm B and Sorger T. 2023. Rapid adaptation of cellular metabolic rate to the microRNA complements of mammals and its relevance to the evolution of endothermy. *iScience* **27**: 108740.

13. Timmermann A, Yun KS, Raia P . . (many others) . . Ganopolski. 2022. Climate effects on archaic human habitats and species succession. *Nature* **604**: 495-501.

14. Dart, RA. *Australopithecus africanus*: the man–ape of South Africa. 1925. *Nature* **115**: 195-199.

15. Reed KE. 1977. Early hominid evolution and ecological change through the African Plio-Pleistocene. *J Hum Evol* **32**: 289-322.

16. Davies TJ, Daru BH, Bezeng BS . . (many others) . . Bond WJ. 2020. Savanna tree evolutionary ages inform the reconstruction of the paleoenvironment of our hominin ancestors. *Sci Rep* **10**: 12430.

17. Pontzer H, Brown MH, Raichlen DA . . (many others) . . Ross SR. 2016. Metabolic acceleration and the evolution of human brain size and life history. *Nature* **533**: 390-393.

18. Yegian AK, Heymsfield SB, Castillo ER, Müller MJ, Redman LM and Lieberman DE. 2024. Metabolic scaling, energy allocation tradeoffs, and the evolution of humans' unique metabolism. *Proc Natl Acad Sci* **121**: e2409674121

19. Kamberova YG, Guhana SM, DeMarchisa A, Jianga J, Wright SS, Morgan BA, Sabetid PC, Tabina CJ, and Lieberman DE. 2018. Comparative evidence for the independent evolution of hair and sweat gland traits in primates. *J Hum Evol* **125**: 99–105.

20. Morin E and Winterhalder B. 2024. Ethnography and ethnohistory support the efficiency of hunting through endurance running in humans. *Nature Hum Behav* **8**: 1065-1075.

21. Hora M, Pontzer H, Wall-Scheffler CM and Sládek V. 2020. Dehydration and persistence hunting in *Homo erectus*. *J Hum Evol* **138**:102682.

22. Armstrong E. 1983. Relative brain size and metabolism in mammals. *Science* **220**: 1302-1304.

23. Aiello LC and Wheeler P. 1995. The expensive-tissue hypothesis: the brain and the digestive system in human and primate evolution. *Curr Anthro* **36**: 199-221.

24. Navarette A, van Schaik CP and Isler K. 2011. Energetics and the evolution of human brain size. *Nature* **480**: 91-93.

25. Penzo-Méndez AI and Stanger BZ. 2015. Organ-size regulation in mammals. *Cold Spring Harb Perspect Biol* **7**: a019240.

26. Fromm B, Høye E, Domanska D . . (many others) . . Friedländer MR and Peterson KJ. 2022.
MirGeneDB 2.1: toward a complete sampling of all major animal phyla. *Nucl Acids Res* **50**: D204-D210.

27. Clarke AW, Høye E, Hembrom AA . . (many others) . . Fromm B. 2025. MirGeneDB 3.0: improved taxonomic
sampling, uniform nomenclature of novel conserved microRNA families and updated covariance models.
Nucl Acid Res **53**: D116–D128.

28. Fromm B, Tarbier M, Smith O, Mármol-Sánchez E, Dalén L, Gilbert MTP and Friedländer MR. 2021.
Ancient microRNA profiles of 14,300-yr-old canid samples confirm taxonomic origin and provide glimpses into
tissue-specific gene regulation from the Pleistocene. *RNA* **27**: 324-334.

29. Rosani U, Bortoletto E, Bai CM, Novoa B, Figueras A, Venier P and Fromm B. 2021.
Digging into bivalve microRNAomes: between conservation and innovation. *Philos Trans R Soc B* **376**: 20200165.

30. Hu RS, Zhang XX, Ma QN, Elsheika HM, Ehsan M, Zhao Q, Fromm B and Zhu XQ. 2021. Differential expression of
microRNAs and tRNA fragments mediate the adaptation of the liver fluke Fasciola gigantica to its intermediate snail and
definitive mammalian hosts. *Int J Parasitol* **51**: 405-414.

31. Tarver JE, Taylor RS, Puttick MN, Lloyd GT, Pett W, Fromm B, Schirrmeister BE, Pisani D, Peterson KJ and Donoghue
PCJ. 2018. Well-annotated microRNAomes do not evidence pervasive miRNA loss. *Genome Biol Evol* **10**: 1457-1470.

32. Ma X, He K, Shi Z, Li M, Li F and Chen XX. 2021. Large-Scale Annotation and Evolution Analysis of
microRNA in Insects. *Genome Biol Evol* **13**: evab083.

33. Taylor AS, Tinning H, Ovchinnikov V, Edge J, Smith W, Pullinger AL, Sutton RA, Constantinides B, Wang D,
Forbes K, Forde N and O'Connell MJ. 2023. A burst of genomic innovation at the origin of placental mammals
mediated embryo implantation. *Comm Biol* **6**:459 (1-8).

34. Jenike AE, Jenike KM, Peterson KJ, Fromm B and Halushka MK. 2023. Direct observation of the evolution
of cell-type-specific microRNA expression signatures supports the hematopoietic origin model of endothelial cells.
Evol Dev **25**: 226-239.

35. Peterson KJ, Beavan A, Chabot PJ, McPeek MA, Pisani D, Fromm B and Simakov O. 2022. MicroRNAs as Indicators
into the Causes and Consequences of Whole-Genome Duplication Events. *Mol Biol Evol* **39**: msab344.

36. Zolotarov G, Fromm B, Legnini I, Ayoub S, Polese G, Maselli V, Chabot PJ, Vinther J, Styfhals S, Seuntjens E,
Di Cosmo A, Peterson KJ and Rajewsky N. 2022. MicroRNAs are deeply linked to the emergence of the
complex octopus brain. *Sci Adv* **8**: eadd9938 (1-12).

37. Cassidy JJ, Bernasek SM, Bakker R, Giri R, Peláez N, Bobrowska A, Bagheri N, Nunes Amaral LA and Carthew RW.
2019. Repressive gene regulation synchronizes development with cellular metabolism. *Cell* **178**: 980-992.

38. Keuls RA, Oh YS, Patel I and Parchem RJ. 2023. Post-transcriptional regulation in cranial neural crest cells expands
developmental potential. *Proc Natl Acad Sci USA* **120**: e2212578120.

39. Ozbudak EM, Thattai M, Kurtser I, Grossman AD and van Oudenaarden. 2002.
Regulation of noise in the expression of a single gene. *Nature Genetics* **31**: 69-73.

40. Buttgereit F and Brand MD. 1995. A hierarchy of ATP-consuming processes in mammalian cells.
Biochem J **312**: 163-167.
731
732 41. Baek D, Villén J, Shin C, Camargo FD, Gygi SP and Bartel DP. 2008. The impact of microRNAs on protein output.
733 *Nature* **455**: 64-72.
734
735 42. Selbach M, Schwanhäusser B, Thierfelder N, Fang Z, Khanin R, Rajewsky N. 2008.
736 Widespread changes in protein synthesis induced by microRNAs. *Nature* **455**: 58–63.
737
738 43. Spies N, Burge CB and Bartel DP. 2013. 3' UTR-isoform choice has limited influence on the stability
739 and translational efficiency of most mRNAs in mouse fibroblasts. *Genome Res* **23**: 2078-2090.
740
741 44. Schmiedel JM, Klemm SL, Zheng Y, Sahay A, Blüthgen N, Marks DS and van Oudenaarden A. 2015.
742 MicroRNA control of protein expression noise. *Science* **348**: 128-132.
743
744 45. Porter WP and Kearney M. 2009. Size, shape, and thermal niche of endotherms.
745 *Proc Natl Acad Sci USA* **106**: 19666-19672.
746
747 46. Pagel M, Meade A and Barker D. 2004. Bayesian estimation of ancestral character states on phylogenies.
748 *Syst Biol* **53**: 673-684.
749
750 47. Venditti C, Meade A. and Pagel M. 2011. Multiple routes to mammalian diversity. *Nature* **479**: 393-396.
751
752 48. Álvarez-Carretero S, Tamuri A, Battini M, Nascimento F, Carlisle E, Asher R, Yang Z, Donoghue P
753 and Dos Reis M. 2022. A species-level timeline of mammal evolution integrating phylogenomic data.
754 *Nature* **602**: 263-267.
755
756 49. Heldstab SA, Isler K, Graber SM, Schuppli C and van Schaik. 2022.
757 The economics of brain size evolution in vertebrates. *Curr Biol* **32**: R697-R708.
758
759 50. Wang ZM, Zhang J, Ying Z and Heymsfield SB. 2012. Organ-tissue level model of resting energy expenditure
760 across mammals: new insights into Kleiber's Law. *ISRN Zool*: 673050.
761
762 51. Elia M. 1992. Organ and tissue contribution to metabolic rate. In *Energy Metabolism:*
763 *Tissue Determinants and Cellular Corollaries*. Kinney JM and Tucker HN (editors), pp: 19-60. Raven Press, New York.
764
765 52. Gallagher D, Belmonte D, Deurenberg P, Wang Z, Krasnow N, Pi-Sunyer FX and Heymsfield SB. 1998. Organ-tissue
766 mass measurement allows modeling of REE and metabolically active tissue mass. *Am J Physiol* **275**: E249–E258.
767
768 53. Kolkotrones T, Savage V, Deeds EJ and Fontana W. 2010. Curvature in metabolic scaling.
769 *Nature* **464**: 753-756.
770
771 54. Grabowski M, Kopperud BT, Tsuboi M and Hansen TF. 2023.
772 Both diet and sociality affect primate brain-size evolution. *Syst Biol* **72**: 404-418.
773
774 55. Scholander PF, Hock F, Walters V, Johnson F and Irving L. 1950.
775 Heat Regulation in Some Arctic and Tropical Mammals and Birds. *Biol Bull* **99**: 237-258.

776 56. Avaria-Llautureo J, Hernandez CE, Rodriguez-Serrano E and Venditti, C. 2019. The decoupled nature
777 of basal metabolic rate and body temperature in endotherm evolution. *Nature* **572**: 651-654.
778
779 57. Wold S, Johansson E. and Cocchi, M. 1993. PLS: partial least squares projections to latent structures.
780 In *Drug Design: Theory, Methods and Applications* (pp. 523-550). Kluwer ESCOM Science Publisher.
781
782 58. de Jong S. SIMPLS: 1993. An alternative approach to partial least squares regression.
783 *Chemometrics and Intelligent Laboratory Systems* **18**: 251-263.
784
785 59. Celeux G and Govaert G. 1995. Gaussian Parsimonious Clustering Models. *Pattern Recognition* **28**: 781-793.
786
787 60. Kummitha CM, Kalhan SC, Saidel GM and Lai N. 2014. Relating tissue/organ energy expenditure to metabolic fluxes
788 in mouse and human: experimental data integrated with mathematical modeling. *Physiol Rep* **2**: e12159.
789
790 61. Tattersall GJ. 2012. Diurnal changes in metabolic rate in pygmy marmosets: Implications for sleep, torpor, and basal
791 metabolism in primates. In: Ruf, Bieber, Arnold, Millesi (eds), *Living in a Seasonal World*, chapter **41**: 471-480. Springer.
792
793 62. Brain CK. 1981. The Hunters or the Hunted?: An Introduction to African Cave Taphonomy.
794 Chicago University Press, Chicago.
795
796 63. Brain C and Mitchell D. 1999. Body temperature changes in free-ranging baboons (*Papio hamadryas ursinus*)
797 in the Namib Desert, Namibia. *Intl J Primatol* **20**: 585-598.
798
799 64. Smaers JB, Rothman RS, Hudson DR . . (many others) . . Safi K. 2021.
800 The evolution of mammalian brain size. *Sci Adv* **7**: eabe2101.
801
802 65. Kuzawa CW, Chugani HT, Grossman LI, Lipovich L, Muzik O, Hof PR, Wildman DE, Sherwood CC, Leonard WR
803 and Lange N. 2021. Metabolic costs and evolutionary implications of human brain development.
804 *Proc Natl Acad Sci USA* **111**: 13010-13015.
805
806 66. Bier DM, Leake RD, Haymond MW, Arnold KJ, Gruenke LD, Sperling MA and Kipnis DM. 1977. Measurement of
807 "true" glucose production rates in infancy and childhood with 6,6-Dideuteroglucose. *Diabetes* **25**: 1016-23.
808
809 67. Sunehag AL, Treuth MS, Toffolo G, Butte NF, Cobelli C, Bier DM and Haymond MW. 2001.
810 Glucose production, gluconeogenesis, and insulin sensitivity in children and adolescents:
811 an evaluation of their reproducibility. *Ped Res* **50**: 115-123.
812
813 68. Camargo FD, Gokhale S, Johnnidis JB, Fu D,1 Bell GW, Jaenisch R, and Brummelkamp TR. 2007.
814 YAP1 increases organ size and expands undifferentiated progenitor cells. *Curr Biol* **17**: 2054-2060.
815
816 69. Hu Y Shin DJ, Pan H, Lin Z, Dreyfuss JM, Camargo FD, Miao J, and Biddinger SB. 2017.
817 YAP suppresses gluconeogenic gene expression via PGC1α. *Hepatol* **66**: 2029–2041.
818
819 70. Ibar C and Irvine KD. 2020. Integration of Hippo-Yap signaling with metabolism. *Dev Cell* **54**: 256-267.
820
821 71. Cox AG, Tsomides A, Yimlamai D . . (many others) . . Goessling W. 2018.
822 Yap regulates glucose utilization and sustains nucleotide synthesis to enable organ growth. *EMBO J* **37**: e100294.

72. Driskill JH and Pan D. 2021. The Hippo pathway in liver homeostasis and pathophysiology. *Ann Rev Pathol Mech Dis* **16**: 299-322.

73. Zhao B, Wei X, Li W . . (many others) . . Guan KL. 2007. Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth control. *Genes Dev* **21**: 2747-2761.

74. Mori M, Triboulet R, Mohseni M, Schlegelmilch K, Shrestha K, Camargo FD and Gregory RI. 2014. Hippo signaling regulates Microprocessor and links cell-density-dependent miRNA biogenesis to cancer. *Cell* **156**: 893–906.

75. Hubbert MK. 1982. Techniques of prediction as applied to the production of oil and gas. In Gass, Saul I. (ed.) *Oil and Gas Supply Modeling* (proceedings of a symposium held at the Department of Commerce, Washington, DC, June 18–20, 1980). National Bureau of Standards Special Publication **631**: 16-141. Washington (DC).

76. Allen AP, Gillooly JF, Savage VM and Brown JH. 2006. Kinetic effects of temperature on rates of genetic divergence and speciation. *Proc Natl Acad Sci USA* **103**: 9130-9135.

77. Schiel N and Souto A. 2017. The Common Marmoset: an overview of its natural history, ecology and behavior. *Dev Neurobiol* **77**: 244

78. Hornstein H and Shomron N. 2006. Canalization of development by microRNAs. *Nature Genetics* **38**: S20-S24.

79. Li X, Cassidy JJ, Reinke CA, Fischboeck S and Carthew RW. 2009. A microRNA imparts robustness against environmental fluctuation during development. *Cell* **137**: 273-282.

80. Mukherji S, Ebert MS, Zheng GXY, Tsang JS, Sharp PA and van Oudenaarden A. 2011. MicroRNAs can generate thresholds in target gene expression. *Nature Genetics* **43**: 854-859.

81. Ebert MS and Sharp PA. 2012. Roles for MicroRNAs in conferring robustness to biological processes. *Cell* **149**: 515-524.

82. Zolboot N, Du JX, Zampa F and Lippi G. 2021. MicroRNAs instruct and maintain cell type diversity in the nervous system. *Front Mol Neurosci* **14**: 646072.

83. Makeyev EV and Maniatis T. 2008. Multilevel Regulation of Gene Expression by MicroRNAs. *Science* **319**: 1789-1790.

84. Erwin DH, Laflamme M, Tweedt SM, Sperling EA, Pisani D and Peterson KJ. 2011.The Cambrian conundrum: early divergence and later ecological success in the early history of animals. *Science* **334**: 1091-1097.

85. Dexheimer PJ and Cochella L. 2020. MicroRNAs: from mechanism to organism. *Front Cell Dev Biol* **8**: 409.

86. Peterson KJ, Beavan A, Chabot PJ, McPeek MA, Pisani D, Fromm B and Simakov O. 2022. MicroRNAs as Indicators into the Causes and Consequences of Whole-Genome Duplication Events. *Mol Biol Evol* **39**: msab344.

87. Baker EA and Woollard A. 2022. The road less travelled? Exploring the nuanced evolutionary consequences of duplicated genes. *Essays in Biochemistry* **66**: 737-744.

88. Martin RD. 1993. Primate origins: plugging the gaps. *Nature* **363**: 223-234.

89. Magalhães JP, Abidi Z, dos Santos GA . . (many others) . . Ka Po To P. 2024. Human Ageing Genomic Resources: updates on key databases in ageing research. *Nucl Acids Res* **52**: D900-D908.

90. Umu SU, Paynter VM, Trondsøn H, Buschmann T, Rounge TB, Peterson KJ and Fromm B. 2023. Accurate microRNA annotation of animal genomes using trained covariance models of curated microRNA complements in MirMachine. *Cell Genomics* **3**: 100348.

91. Pagel M, Meade A and Barker D. 2004. Bayesian estimation of ancestral character states on phylogenies. *Syst Biol* **53**: 673-684.

92. Venditti C, Meade A. and Pagel M. 2011. Multiple routes to mammalian diversity. *Nature* **479**: 393-396.

93. Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol Evol* **3**: 217-223.

94. Reinsch CH. 1967. Smoothing by Spline Functions. *Numerische Mathematik* **10**: 177–183.

95. Eubank RL. 1999. Nonparametric regression and spline smoothing (2nd ed). CRC. Boca Raton, Florida.

96. Orme D, Freckleton RP, Thomas GH and Petzoldt T. 2013. CAPER: comparative analyses of phylogenetics and evolution in R. *Methods Ecol and Evol* **3**: 141-151.

DECLARATIONS

Acknowledgment

BF is supported by the Tromsø forskningsstiftelse (TFS) [20_SG_BF 'MIRevolution'].

Author Contributions

Conceptualization: TS; Methodology: TS; Investigation: BF and TS; Writing – Original Draft: TS; Writing – Review & Editing: BF and TS; Visualization: TS; Funding Acquisition: BF.

Declaration of Interests

The authors declare no competing interests.

Statement on the use of AI

No part of this work was generated or edited using any Large Language Module.