Disentangling acute motor deficits and adaptive responses evoked by the loss of cerebellar output

Nirvik Sinha[1,2,3*]**, Sharon Israely**[1]**, Ora Ben Harosh**[1]**, Ran Harel**[4]**, Julius PA Dewald**[2,3,5,6]**, Yifat Prut**[1*]

[1]Edmond and Lily Safra Center for Brain Sciences, The Hebrew University of Jerusalem, 9190401, Jerusalem, Jerusalem, Israel; [2]Department of Physical Therapy and Human Movement Sciences, Northwestern University Feinberg School of Medicine, 60610, Chicago, Illinois, USA; [3]Interdepartmental Neuroscience Center, Northwestern University Feinberg School of Medicine, 60611, Chicago, Illinois, USA; [4]Department of Neurosurgery, Sheba Medical Center, 5262000, Tel Aviv, Tel Aviv, Israel; [5]Department of Biomedical Engineering, Northwestern University, 60208, Evanston, Illinois, USA; [6]Department of Physical Medicine & Rehabilitation, Northwestern University Feinberg School of Medicine, 60610, Chicago, Illinois, USA

*For correspondence:

Nirvik.Sinha@mail.huji.ac.il (NS)

yifat.prut@mail.huji.ac.il (YP)

Abstract

Patients with cerebellar damage experience various motor impairments, but the specific sequence of primary and compensatory processes that contribute to these deficits remains uncertain. To clarify this, we reversibly blocked cerebellar outflow in monkeys engaged in planar reaching tasks. This intervention led to a spatially selective reduction in hand velocity, primarily due to decreased muscle torque, especially in movements requiring high inter-joint torque coupling. When examining repeated reaches to the same target, we found that the reduced velocity resulted from both an immediate deficit and a gradually developing compensatory slowing strategy designed to reduce passive inter-joint interactions. However, the slowed hand velocity did not account for the fragmented and variable movement trajectories observed during the cerebellar block. Our findings indicate that cerebellar impairment results in motor deficits due to both inadequate muscle torque and an altered compensatory control strategy for managing impaired limb dynamics. Additionally, impaired feedforward control elevates motor noise, which cannot be entirely mitigated through compensatory strategies.

Introduction

The ability to coordinate well-timed movements greatly relies on the cerebellum as evident from studies of individuals with cerebellar ataxia (Beppu, Suda et al. 1984). The cerebellum regulates the timing of movements by predicting the mechanical interactions between adjacent joints (Ivry and Keele 1989, Manto, Bower et al. 2011), a computation that requires an internal model of feedforward control (Shidara, Kawano et al. 1993, Shadmehr and Mussa-Ivaldi 1994, Wolpert, Ghahramani et al. 1995, Wolpert, Miall et al. 1998, Kawato 1999, Sainburg, Ghez et al. 1999, Popa, Hewitt et al. 2013). However, motor impairments exhibited by individuals with cerebellar deficits extend much beyond poor motor timing. For example, during the acute phase of cerebellar injury, there is a loss of muscle tone and weakness of voluntary movements (Holmes 1917, Goldstein 1927, Holmes 1939). Cerebellar patients also exhibit movement-related sensory deficits such as an impaired visual perception of stimulus movement and perception of the limb position sense (i.e., proprioception or state estimation) during active movements (Therrien and Bastian 2015, Weeks, Therrien et al. 2017).

Clinical studies of cerebellar patients have yielded conflicting insights into the pathophysiology caused by a loss of cerebellar function. Some researchers suggest that motor impairments stem primarily from an inability to compensate for inter-joint interactions (Bastian, Martin et al., 1996; Bastian, Zackowski et al., 2000), while others argue that insufficient muscle torque generation is the central issue (Topka, Konczak et al., 1998; Boose, Dichgans et al., 1999). These discrepancies likely arise from the variability among individuals with chronic cerebellar deficits, where primary and compensatory effects are intertwined and difficult to separate. As a result, the specific sequence of events that leads to motor impairments in cerebellar patients remains unclear.

To investigate this issue, we trained monkeys to perform a planar reaching task with arm support provided by an exoskeleton. We then reversibly blocked cerebellar output to the motor cortex using high-frequency stimulation (HFS) of the superior cerebellar peduncle, assessing the impact of this intervention on subsequent movements. Previously, we have shown that HFS can effectively and reversibly interfere with the normal outflow of cerebellar signals (Nashef, Cohen et al. 2019) since this perturbation accurately replicated the behavioral deficits identified in cerebellar patients (Bastian 1997) and the changes in motor cortical activity that appear after dentate cooling in monkeys (Meyer-Lohmann, Conrad et al. 1975). In the present study, we found that the loss of cerebellar outflow led to reduced hand velocity during reaching movements, reflecting two parallel underlying mechanisms. First, there was a reduction in muscle torques, even for movement directions where inter-joint interactions were low. Second, the reduction in hand velocity exhibited a spatial tuning and was greater in directions that generated large inter-joint interactions, thereby reflecting an inability to compensate for limb dynamics. The time course of these two mechanisms was identified by analyzing the sequence of

movements made to the same target during the cerebellar block. During such
movements, hand velocity was low initially, reflecting a primary muscle torque
deficit, and declined further in successive trials, revealing an adaptive motor strategy
to minimize inter-joint interactions. Finally, we found that the decrease in velocity
during cerebellar block could not fully explain the noisy and decomposed movements
observed under these conditions, which may be due to an impaired state estimate of
the limb that leads to the faulty feedforward control of inter-joint interactions. Taken
together, we show that cerebellar output is essential for generating sufficient muscle
torques to allow for fast and stereotypical movements while overcoming the complex
inter-joint interactions inherent in such actions. The loss of this output triggers a
sequence of early primary effects followed by compensatory responses aimed at
mitigating these impairments. Together these processes result in significantly slower
movements.

Results

Slower reaching movements in cerebellar block are due to weakness and
impaired limb dynamics.

We trained four adult monkeys (*Macaca fascicularis*) to sit in a primate chair with
their upper limb supported by an exoskeleton (KINARM system) and perform planar
center-out movements (**Figure 1a**). The spatial location of the targets enabled us to
dissociate between shoulder-dominant (lateral targets) *vs.* shoulder-elbow
movements (diagonal and straight targets respectively).

Cerebellar block significantly reduced the ability of the monkeys to perform the
task as measured by the success rate across all the recorded sessions (mean success
rate in control = 86.5%, CI [85.8, 87.2] vs. cerebellar block = 75.3%, CI [73.9, 76.7], p
< 0.001; **Supplementary table S1**). Since the monkeys had to meet strict timing
criteria to perform the movements successfully during the task, we began by
examining the hand kinematics**. Figure 1b** shows profiles of hand velocity in sample
trials collected during the control (blue) and cerebellar block (vermillion) conditions.
This example shows that movements exhibited a significant reduction in the peak
hand velocity during cerebellar block. However, when compared across the targets,
we found that the reduction in hand velocity was strongly modulated by the target
direction (1-way ANOVA of the change in peak hand velocity, the effect of cerebellar
block: p = 0.01, the effect of targets: p < 0.001; **Supplementary table S2**). *Post hoc*
comparisons further revealed that the effect of cerebellar block on outward reaching
movements i.e., movements to targets 1-4 was significantly higher than on inward
reaching movements i.e., movements to targets 5-8 (**Figure 1c**; mean decrease of
9.7%, CI [8.2, 11.2], p < 0.001 for targets 1-4 compared to mean decrease of 2.5%, CI
[0.9, 4.0], p = 0.21 for targets 5-8) even though there was no difference in the peak
hand velocity between the movements to the two target groups during control (p =
0.88, **Supplementary tables S3a and b**).

In the task used in our study, hand movements were made by controlling the muscle
torque at the shoulder and elbow joints (Scott 2004). However, simultaneous
movements of both joints also generated coupling torques due to passive interactions
between limb inertial properties and joint velocities (Bernshteĭn 1967, Hollerbach
and Flash 1982). To identify the target-specific muscle and coupling torques, we
employed an inverse dynamics model of the upper limb (Fagg, Ojakangas et al. 2009)
which enabled us to distinguish between the different contributions to the net torque
experienced by each joint. According to this model, the net torque at each joint is the
sum of both the active muscle torque and the passive coupling torque acting at that
joint (see Methods section for details).

Using the inverse dynamics model, we found that the target-specific reduction in
hand velocities was produced by changes in muscle commands (e.g., **Figures 2a, b**).
Specifically, we observed that the muscle torque impulses, calculated by integrating
the torque profiles over the early phase of movements when acceleration was
positive, were reduced in a target-specific manner during the cerebellar block
(**Figure 2c**; **Supplementary table S4**-**5**). This reduction was particularly
pronounced at the shoulder joint. Furthermore, as in the case of hand velocities, *post*
hoc comparisons revealed that the shoulder joint experienced a significantly greater
reduction in muscle torque for targets 1-4 vs. 5-8 (**Figure 2d**; mean decrease of
10.6%, CI [9.0, 11.2], p < 0.001 for targets 1-4 compared to mean decrease of 3.3%, CI
[-1.3, 5.2] p = 0.21 for targets 5-8). In contrast, the slight reduction in elbow torques
due to the cerebellar block did not reach significance for any of the targets (**Figure**
2e). Collectively, these findings indicate that during the cerebellar block, there is a
reduction in shoulder muscle torque specifically for targets involved in outward-
reaching movements, which leads to the target-specific reduction in hand velocity.

Several studies have shown that individuals with cerebellar deficits exhibit an
inability to correctly compensate for inter-joint interactions (Bastian, Martin et al.
1996, Bastian, Zackowski et al. 2000). Therefore, we asked whether the reduction in
muscle torques and hand velocities due to the cerebellar block could be associated
with the coupling torques generated during movements in various directions.
Figures 3a and b show the coupling torque profiles at the shoulder and elbow joints
respectively during sample trials. We observed that the coupling torque impulses
varied in a target-dependent manner during both the control and cerebellar block
conditions (**Figure 3c**). Furthermore, the reduction in peak hand velocities during
cerebellar block was correlated to the net coupling torque impulse (calculated as the
sum of absolute coupling torque impulses at both joints) during control, particularly
for the outward reaching movements (**Figure 3d;** targets 1-4: ρ = 0.33, CI [0.25, 0.38],
p < 0.001). This indicated that the effect of the cerebellar block on hand velocity
during the outward reaching movements was strongest for targets with high coupling
torques. Nevertheless, despite the significant correlation between the reduced hand
velocity and coupling torques, the cerebellar block reduced the hand velocity even in
the single-joint outward reaches (i.e., movements to target 1, mean reduction in hand
velocity = 5.8%, CI [2.8, 8.7], p < 0.001) where the elbow was stationary. This finding

indicated a reduction in the motor command which occurred in addition and beyond
the effect attributable to the impaired control of coupling torques alone.

Overall, the cerebellar block produced a significant impact on the kinematics and
dynamics of outward reaching movements. Specifically, the cerebellar block caused
slower hand velocities due to a systematic, target- and joint-specific reduction in
muscle torques for these movements, which strongly correlated with the coupling
torques. Additionally, a reduction in muscle torques was also observed during the
single-joint outward reaching movements, indicating an additional component of
muscle torque insufficiency that further exacerbated motor impairment following
cerebellar block.

Hand velocity exhibits an adaptive trend during repeated movements to the same target during cerebellar block.

The advantage of using a reversible cerebellar block in an animal model is the
ability to track the temporal profile of motor impairments induced by this condition,
allowing us to distinguish between primary and compensatory effects. To this end,
we assessed changes in hand kinematics produced by the cerebellar block across
successive trials. Specifically, we examined the evolution of peak hand velocities for
single-joint (target 1) vs. dual-joint outward reaches (data pooled for movements to
targets 2-4) during blocks of control and cerebellar block trials. **Figure 4a** illustrates
this procedure. Since the targets were presented in a pseudorandom order during
each block of trials, we collected for each block the tested target, while keeping its
order of presentation to generate a temporally accurate sequence of trials, all directed
towards the same single target.

We found that during the control condition, peak hand velocity remained stable
across successive trials. In contrast, cerebellar block affected hand velocity in a trial-
and target-dependent manner (**Figure 4b, c**). For movements to target 1 (i.e., single-
joint movements), hand velocity was lower than control but unaffected by the
temporal sequence of target presentation (trial type × trial sequence interaction
effect: p = 0.69, **Supplementary Table S6**). However, for movements to targets 2-4,
hand velocity was lower than control in the early trials and declined further with
successive trials (trial type × trial sequence interaction effect: p = 0.007,
Supplementary Table S7).

To further quantify the effect of trial sequence on movements to targets 2-4, we
performed *post hoc* comparisons (**Figure 4d**). These comparisons revealed that
during the cerebellar block, hand velocity was 4.3% (CI [1.0, 7.3], p < 0.001) lower
than control during the first 1-2 trials. With successive trials, the hand velocity
declined further before reaching an asymptote during the late trials 11-20 at 8.2% (CI
[5.7, 10.7], p < 0.001) lower than the control.

Taken together these results reveal that the hand velocity of dual-joint outward-
reaching movements during cerebellar block was already low initially, reflecting a

primary deficit. Over and above this, there was a further progressive decline of the
hand velocity during successive movements, most likely due to an adaptive response
to the inability to compensate for coupling torques. Furthermore, this adaptive
response was not observed in the case of single-joint movements where the coupling
torques were low, and therefore inter-joint interactions did not play a significant role.

Slow movements during the cerebellar block cannot fully account for the impaired motor timing.

Individuals with cerebellar deficits suffer from poorly timed movements, including
asynchronous activation of joints during multi-joint movements leading to movement
decomposition (Bastian, Martin et al. 1996) or spatial variability in their movement
trajectories across trials (Day, Thompson et al. 1998). However, these properties of
movement are also velocity-dependent (Bastian, Martin et al. 1996, Osu, Morishige et
al. 2015). Therefore, it is conceivable that the combination of primary deficit and
adaptive reduction in velocity to reduce the effects of coupling torque observed in our
results is the main driver of abnormal motor timing. To evaluate this, we examined
movement decomposition and trajectory variability during the outward reaching
movements (which exhibited a significant reduction in hand velocity during
cerebellar block) and measured the impact of hand velocity on these measures.

To quantify movement decomposition, we normalized the movement times of
individual trials to range from 0 to 1 and identified the bins where either the shoulder
or the elbow joint paused (angular velocity < 20°/s) while the other joint continued
moving (Bastian, Martin et al. 1996). In control trials, decomposition was primarily
observed during the early phase of movement, where joint velocities are low making
this measure more sensitive (**Figure 5a**). However, during cerebellar block trials,
decomposition occurred throughout the movement (**Figure 5b**) similar to individuals
with cerebellar deficits (Bastian, Martin et al. 1996). To quantify this effect, we
calculated a decomposition index for each trial, representing the proportion of
movement time during which the movement was decomposed, as defined above.
Overall, the cerebellar block increased movement decomposition by 53.5% (CI [42.2,
66.0], p < 0.001). Next, we asked whether movement decomposition was mainly due
to lower hand velocities. We therefore selected a subset of control trials that matched
the cerebellar block trials in their peak velocities. However, even though movement
decomposition in these control trials was higher (11.0%, CI [5.2, 17.0], p = 0.03)
compared to all control trials, it was still significantly lower (p < 0.001) than the
velocity-matched cerebellar block trials (in three out of the four monkeys as shown
in **Figure 5c**). This indicates that even at matched velocities, movements during
cerebellar block were more decomposed relative to the control.

Next, we assessed the effect of cerebellar block and hand velocity on trajectory
variability. To this end, we first aligned the hand trajectories to have the same starting
position, and then rotated them so that their endpoint was located on the positive Y-
axis (**Figure 5d**). Subsequently, we defined the error in each trajectory as the

maximal perpendicular distance from the Y-axis. The standard deviation of the errors
across all trials was used to quantify the trial-to-trial trajectory variability. Normally,
slower movements are also less variable due to the speed-accuracy tradeoff
(Plamondon and Alimi 1997). Therefore, we selected the subset of control trials that
matched the cerebellar block trials in their peak velocities. Indeed, the trajectory
variability in this subset of slower control trials was 5.5% (CI [0.9, 9.9], p = 0.02)
lower than that of all control trials. However, when we compared the subset of
velocity matched control and cerebellar block trials, we found that cerebellar block
exhibited 34.6% (CI [26.2, 43.2], p < 0.001) higher trajectory variability (**Figure 5e)**.
In other words, despite slower movements, cerebellar block increased trajectory
variability.

Discussion

Individuals with cerebellar deficits exhibit altered motor behavior related mostly
to poor timing and coordination of voluntary movements. Understanding the
impairments that follow cerebellar lesions is often complicated by the fact that the
studies of this question mostly rely on a heterogeneous population due to various
levels and types of cerebellar pathologies. Here we addressed this question by using
high-frequency stimulation to reversibly block cerebellar output to the motor cortex
in behaving monkeys wearing an exoskeleton (Nashef, Cohen et al. 2019) and
comparing the post-block motor behavior to control conditions. Our results indicate
that cerebellar block leads to decreased hand velocity, particularly during the
outward reaching movements via two mechanisms: a reduction in muscle torques
suggesting motor weakness, and spatially tuned decreases in velocity in targets
where large inter-joint interactions occur, indicating a failure to compensate for limb
dynamics. Analyzing the time course of these impairments revealed partially
overlapping processes: an initial reduction in hand velocity due to the inability to
generate sufficient muscle torques, followed by a further progressive decline in hand
velocities during subsequent movements during the cerebellar block. This secondary
decline was observed only in the case of dual-joint movements, suggesting an
adaptive strategy to minimize inter-joint interactions. Finally, we observed increased
motor noise in terms of movement decomposition and trial-to-trial trajectory
variability which were independent of the slower movements observed during the
cerebellar block.

Target-dependent reduction in hand velocity and coupling torques during
reaching movements during the cerebellar block.

Cerebellar block led to a target-specific reduction in hand velocity, even though the
arm was fully supported by the exoskeleton and thus the limb weight did not play any
role. This observation replicates findings made in previous studies that have
documented the slowing of movements in individuals with cerebellar lesions (Wild
and Dichgans 1993, Topka, Konczak et al. 1998, Konczak, Schoch et al. 2005) as well

as in monkeys that underwent dentate cooling (Tsujimoto, Gemba et al. 1993). It was
argued that the reduction in hand velocity may be a compensatory strategy to
counteract the impaired ability to manage inter-joint coupling torques in the absence
of cerebellar input to the motor cortex (Wild and Dichgans 1993, Bastian and Thach
1995, Bastian, Martin et al. 1996, Beer, Dewald et al. 2000). Since coupling torques in
multi-joint movements are generated in a velocity-dependent manner (Hollerbach
and Flash 1982, Virji-Babul and Cooke 1995), slowing down the movement not only
simplifies its kinetics (by reducing the impact of coupling torques) but also allows
more time for slower visual feedback to aid in guiding the movement (Jeannerod
1988). Our findings, which demonstrate a correlation between the extent of hand
velocity reduction during cerebellar block and the magnitude of coupling torque
under control conditions, further corroborate this hypothesis and confirm our
approach for reversible inactivation of cerebellar output as a model for ataxia.
Furthermore, our model enabled us to study this effect rigorously and systematically
while dissociating it from the other primary effects of cerebellar block as discussed
below.

Muscle torque deficit is an acute response to the cerebellar block.

Poor compensation for coupling torque is a fundamental deficit in cerebellar ataxia
(Bastian, Zackowski et al. 2000). However, early studies have reported the
occurrence of muscle weakness (asthenia) and hypotonia acutely following
cerebellar injury in humans (Haines, Manto et al. 2007) or experimental lesions in
animals (Luciani 1893, Bremer, Roger et al. 1935, Fulton and Dow 1937, Granit,
Holmgren et al. 1955). Both Luciani (Luciani 1893) and Holmes (Holmes 1917)
described a triad of cerebellar deficits to be atonia (low muscle tone), asthenia
(weakness of voluntary movements), and astasia (oscillation of the head and trunk).
Importantly, muscle weakness is typically characteristic of the acute phase of injury,
especially in the upper extremity, and tends to normalize gradually over weeks to
months, depending on the severity of the injury (Konczak, Pierscianek et al. 2010). It
is therefore conceivable that studies in humans with cerebellar deficits, which often
included subjects at the chronic stage of the lesion, may have underestimated the
contribution of muscle weakness which by that time was already less pronounced.

We explored the acute outcome of the cerebellar block, a phase in which muscle
weakness may play a more prominent role. Importantly, we showed that muscle
torque deficits appear immediately after the cerebellar block, resulting in lower hand
velocity already in the first few trials of both single- and dual-joint movements. This
kind of fast response is less likely to be a strategic response aimed at mitigating the
loss of cerebellar signals. This finding is consistent with the initial weakness followed
by a gradual recovery of function observed in acute lesions of other neural structures
involved in movement control, including the primary (Pressman and Rosen 2015)
and premotor cortices (Freund 1985, Freund and Hummelsheim 1985). It also aligns
with the understanding that cerebellar output provides a strong excitatory drive to
the motor cortex (Shinoda, Yamazaki et al. 1982, Hore and Flament 1988, Nashef,

Cohen et al. 2018) and several subcortical descending motor systems (Ruigrok 2013), beyond its well-known role in motor timing and coordination. Following the initial reduction in hand velocity observed in our study, there was a further gradual decline across subsequent trials specifically for the dual-joint movements, a behavior profile that is more consistent with strategic effort, most likely in response to the inability to compensate for limb dynamics.

Therefore, the slower movements in the absence of cerebellar signals are likely due to two main factors: (i) a primary inability to generate sufficient muscle torques due to the lack of synchronized recruitment of the cortical and sub-cortical descending motor systems via the cerebellar output, and (ii) a secondary adaptive strategy aimed at compensating for the impaired prediction of limb dynamics due to the loss of feedforward control signals from the cerebellum.

Insufficient muscle torques vs. poor inter-joint coordination.

Unlike most previous clinical studies that examined motor deficits in individuals with chronic cerebellar injuries, our study utilized a reversible model of cerebellar block. This approach enabled us to track the temporal evolution of motor deficits from the onset of the block within the same study participant. Although several studies have explored the nature of motor deficits following cerebellar injuries, there remains considerable debate regarding the fundamental mechanisms driving these deficits. Some studies have suggested that the inability to compensate for inter-joint interactions is the primary driver of the observed motor abnormalities in cerebellar injury (Bastian, Martin et al. 1996, Bastian, Zackowski et al. 2000). In contrast, other studies have emphasized the inability to generate sufficient muscle torques as the core deficit in cerebellar ataxia while claiming that inter-joint coordination is unaffected following cerebellar lesions (Topka, Konczak et al. 1998, Boose, Dichgans et al. 1999). Given these conflicting findings, our study aimed to systematically explore the entire spectrum of motor deficits using a model of reversible cerebellar block. First, we found that the deficits in muscle torque are more pronounced for the reaching movements as compared to the retrieval movements. We identified both an acute insufficiency in muscle torque generation and the inability to compensate for coupling torques as potential mechanisms underlying the observed slowness during the reaching movements. Moreover, we delineated the time course of these deficits, providing insights into their evolution from the onset of cerebellar block.

The asymmetric effect of cerebellar block on outward vs. inward reaching movements.

We observed a notable distinction in the impact of cerebellar blockage on outward vs. inward reaching movements. At a collective level, this variance could be attributed to the disparity in magnitude of coupling torque impulse, with movements towards targets 2-4 exhibiting 18.7% (CI [13.1, 24.1], $p = 0.008$) higher coupling torque impulse compared to those towards targets 6-8. However, this explanation does not hold consistently when examined at the individual target level. For instance, while

the coupling torque was higher for movements towards target 7, the effects of
cerebellar blockage were less pronounced compared to those observed for target 1,
where the coupling torques are considerably lower. There are two possible
explanations for this discrepancy. One possibility is that the effect of cerebellar block
exhibits a bias in the direction of shoulder flexion (i.e., for movements to targets 1-4).
Several early studies in cats (Hare, Magoun et al. 1936, Chambers 1947), non-human
primates (Magoun, Hare et al. 1935) , and humans (Nashold and Slaughter 1969) have
all reported a bias towards the activation of ipsilateral flexors of the arm following
stimulation of the deep cerebellar nuclei and the superior cerebellar peduncle.
Furthermore, lesioning of these structures led to the reduction of ipsilateral flexor
tone and power (Schneider and Crosby 1963, Nashold and Slaughter 1969). The
cerebellar output influences the excitability of spinal interneurons either directly
(Bantli and Bloedel 1975, Asanuma, Thach et al. 1980, Sathyamurthy, Barik et al.
2020) or via indirect relays in the subcortical descending motor systems like the
rubrospinal and the reticulospinal pathways (Ruigrok 2013). Since the spinal
interneurons influence the activity of flexor muscles twice as often as extensor
muscles (Perlmutter, Maier et al. 1998), this might explain why the cerebellar block
had a stronger effect on movements involving shoulder flexion (i.e., outward reaching
movements) relative to shoulder extension (i.e., inward reaching movements) in our
study. Alternatively, the initial position of the limb in our experimental setup (i.e., on
the central target) might have been favorable for movements in the direction of
shoulder extension, thereby making them easier to perform.

Asynchronous joint movements and motor variability are unaffected by the movement velocity in cerebellar block.

Abnormally high motor variability (Day, Thompson et al. 1998, Timmann, Watts et
al. 1999, Schlerf, Xu et al. 2012) and decomposition of multi-joint movements
(Bastian, Martin et al. 1996) are core deficits in cerebellar patients. We showed that
the cerebellar block decomposed the dual-joint movements into temporally isolated
single-joint actions and led to an increased trial-to-trial spatial noise. These two
measures can be affected by the reduction in movement velocities. Our results
showed that even though movement decomposition increased with a reduction in
velocity in control trials, they were still significantly lower than that of velocity-
matched cerebellar block trials. Second, the trial-to-trial spatial noise in hand
trajectories increased during cerebellar block despite the slower movement speeds.
The cerebellum is known to participate in the planning of movements by providing
an internal model of the limb which is necessary for predicting the sensory outcomes
of motor commands (Wolpert, Miall et al. 1998). For this purpose, the cerebellum aids
in estimating the state of the limb by integrating the sensory information of its last
known state with predictions of its response to the latest motor command (Miall,
Christensen et al. 2007, Miall and King 2008). Therefore, in the absence of cerebellar
signals, there are programming errors in the feedforward command required to
launch the limb accurately toward a target (Day, Thompson et al. 1998). These
feedforward errors may in turn translate into the noisy joint activation patterns and

variable hand trajectories observed in our results. Moreover, the persistence of these deficits independently of the movement speeds observed during cerebellar block indicates that they cannot be compensated for, unlike the impaired prediction of inter-joint interactions which are strategically dealt with by reducing the movement velocity.

In summary, our study systematically investigated the temporal progression and distinct mechanisms of motor deficits in a controlled, reversible animal model of cerebellar block. These include an acute onset muscle torque deficit, impaired compensation of limb dynamics, and increased motor noise. We aim to further explore the neural correlates of these deficits in future studies.

Materials and Methods

Experimental subjects

This study was performed on four adult female monkeys (*Macaca fascicularis*, weight 4.5-8 kg). The care and surgical procedures of the subjects will be in accordance with the Hebrew University Guidelines for the Use and Care of Laboratory Animals in Research, supervised by the Institutional Committee for Animal Care and Use.

Behavioral task

Each monkey was trained to sit in a primate chair and perform planar, center-out movements. During the task, the left upper arm was abducted 90° and rested on cushioned troughs secured to links of a two-joint exoskeleton (KINARM, BKIN Technologies Ltd.). The sequence of events in each trial of this task was as follows. First, the monkey located a cursor (projected on a horizontal screen immediately above its arm) within a central target. After 500-800 ms, a randomly selected peripheral target (out of 8 evenly distributed targets) was displayed. The monkey had to wait until the central target disappeared (i.e., the 'go' signal) and then reached the cued peripheral target. This delay between the cue and the 'go' signal was 450-600 ms. The monkey was rewarded with a drop of applesauce if it moved the cursor to the correct target within 1 s of the 'go' signal.

Insertion of the stimulating electrode into the superior cerebellar peduncle

After the training was completed, a square recording chamber (24x24 mm) was attached to the monkey's skull above the upper limb-related area of the motor cortex in a surgical procedure under general anesthesia. To insert a chronic stimulating electrode into the ipsilateral superior cerebellar peduncle (SCP), a round chamber (diameter = 19 mm) was also implanted above the estimated insertion point (as per stereotactic measurements) during the surgery for the cortical chamber implantation. Subsequently, a post-surgery MRI was used to plan the trajectory for

the electrode insertion into the SCP. After a recovery and re-training period, a bi-polar
concentric electrode (Microprobes for Life Science Inc.) was inserted through the
round chamber along the planned trajectory. During the process of insertion, the
evoked multi-unit activity in the primary motor cortex was simultaneously
monitored to identify the precise location where the stimulating electrode should be
secured (Nashef, Cohen et al. 2018, Nashef, Rapp et al. 2018).

High frequency stimulation to block the cerebellar outflow to the motor cortex

The superior cerebellar peduncle was blocked using high-frequency stimulation
(HFS). This stimulation protocol consisted of a train of biphasic rectangular-pulse
stimuli (where each phase was 200 µs) applied at 130 Hz. Each train was delivered at
a fixed intensity of 150-250 µA. In our earlier study, we demonstrated that by using
this manipulation we can successfully replicate the impaired motor timing and
coordination characteristic of cerebellar ataxia (Nashef, Cohen et al. 2019).
Furthermore, we also demonstrated that at the motor cortical level, this manipulation
leads to a loss of response transients at movement onset and decoupling of task-
related activity, similar to findings observed in studies where the dentate nucleus was
cooled (Meyer-Lohmann, Conrad et al. 1975). Deep brain stimulation (DBS) is a
perturbation similar in properties to the HFS we used, which is routinely used in
clinical practice for similar purposes of interfering with neural circuitries that
underwent pathological changes (Perlmutter and Mink 2006) although in both
applications there is no accurate measure for the extent of the block (i.e., the fraction
of fibers which become ineffective in these conditions) other than the behavioral
consequences. HFS probably imposes only a partial block of the CTC system, but the
behavioral outcomes we found here during an adaptation task further confirm the
validity of this approach.

Data collection protocol

Each monkey performed the experimental task for several days (sessions). A total
of 46, 29, 36, and 54 sessions of data were collected from Monkeys S, C, M, and P,
respectively. Each session was divided into three or four sub-sessions. Each sub-
session included blocks of control trials (~80 trials) and trials with the superior
cerebellar peduncle blocked using high-frequency stimulation (~50 trials) as
described above. During the task performance, the following behavioral data was
obtained at 1000 Hz using the KINARM's motor encoders: (i) Hand position in the x-
y plane of movement, (ii) Elbow and shoulder joint angular positions, velocities, and
accelerations. The lengths of the upper arm and forearm of each monkey were
measured as the distances from the head of the humerus to the lateral epicondyle and
from the lateral epicondyle to the palm, respectively. The inertial properties of the
upper limb were estimated using the monkey's body weight (Cheng and Scott 2000).

Processing of the behavioral data

All data analysis was performed in MATLAB R2021b (MathWorks Inc.). The raw
kinematic data was low-pass filtered by using a second order Butterworth filter
(cutoff = 10Hz) and then epoched into trials. For this purpose, the start and end events
were taken as the time of presentation of the cue (i.e., the peripheral target) and the
time of entry of the control cursor into the peripheral target, respectively. This cutoff
meant that a major portion of the extracted segment of each trial was the acceleration
phase of the movement. The movement onset was defined as the time when the radial
hand velocity exceeded 5% of its peak. Subsequently, all the trials were zero-centered
around the calculated movement onset. We then applied a two-link inverse dynamics
model of the upper arm to the kinematic data to compute the coupling and muscle
torques acting on the elbow and shoulder joints (Fagg, Ojakangas et al. 2009). The
inverse dynamics equations of motion for the combined system consisting of the
monkey's arm and the associated moving components of the KINARM were used to
compute the joint torques:

$$T_{me} = T_{ne} - T_{ce} \tag{i}$$

$$T_{ms} = T_{ns} - T_{cs} \tag{ii}$$

where T_{ne} and T_{ns} are the net torques (proportional to the joint's acceleration), T_{ce}
and T_{cs} are passive coupling torques, and T_{me} and T_{ms} are the active muscle torques
acting at the elbow and shoulder joints respectively. The expanded formulas for
computing each of the above terms are provided by equations 1a-2e in the methods
section C (Fagg, Ojakangas et al. 2009). It must be noted that the 'muscle' torques
computed here include the effects of the actively generated muscle forces as well as
the viscoelastic effects that result from the musculoskeletal system and the KINARM
(Smith and Zernicke 1987).

Statistical analysis

We used linear mixed effects analysis to evaluate the effects of predictor variables
of interest on the computed response variables. This allowed us to account for the
variability across the monkeys which was modeled as a random effect. Each data
sample for this model was obtained by calculating the median of all the trials for each
unique combination of the response variable(s) per session. We computed the change
in the measured outcome variable due to the cerebellar block per session using the
following formula:

$$\frac{Median\ response\ during\ cerebellar\ block - median\ response\ during\ control}{Median\ response\ during\ control}$$

$\times\ 100$

For post-hoc comparisons, we used the 'coefTest' function from MATLAB's 'fitlme'
to perform hypothesis testing on the fixed effects. This involved specifying the
contrast matrices to evaluate specific hypotheses regarding the differences in the
response variables across the levels of the predictor variables. The 'coefTest' function

allowed us to obtain p-values for these comparisons. The computed p-values were
then corrected using the Benjamini-Hochberg procedure to control the false
discovery rate for multiple post-hoc comparisons, wherever applicable (Haynes
2013). A significance level of 0.05 was used for all comparisons in this study. For
estimation of the group means and their confidence intervals, we pooled the per
session data from all monkeys after ensuring that effects are consistent across the
monkeys. For further clarity, we have also overlayed the means of the measured
variables from individual monkeys on all our plots of summary statistics.

Acknowledgements

This work was funded by the Israel Science Foundation (ISF-1801/18 and ISF
1207/23) to Y.P., the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation Project-ID 431549029–SFB 1451) to Y.P., National Institute of
Neurological Disorders and Stroke of the National Institutes of Health (NIH
R01NS11090) to. Y.P., the National Institutes of Health (NIH R01NS105759) to
J.P.A.D., the Council for Higher Education PhD Sandwich Fellowship, Government of
Israel to N.S. for undertaking his dual-PhD at the Hebrew University and
Northwestern University, and the generous support of the Baruch Foundation to Y.P.
The authors would like to thank Dr. Abdulraheem Nashef, a postdoctoral fellow at the
University of Colorado for collecting part of the data used in this study.

Data availability statement

539 Data and code used in this manuscript is available publicly as a GitHub repository
540 (Sinha et al. 2024).

Competing interests

542 The authors declare no financial or non-financial competing interests.

References

Asanuma, C., W. T. Thach and E. G. Jones (1980). "Nucleus interpositus projection to
spinal interneurons in monkeys." Brain Research **191**(1): 245-248.
Bantli, H. and J. R. Bloedel (1975). "The action of the dentate nucleus on the
excitability of spinal motoneurons via pathways which do not involve the primary
sensorimotor cortex." Brain Research **88**(1): 86-90.
Bastian, A. J. (1997). "Mechanisms of Ataxia." Physical Therapy **77**(6): 672-675.
Bastian, A. J., T. A. Martin, J. G. Keating and W. T. Thach (1996). "Cerebellar ataxia:
abnormal control of interaction torques across multiple joints." Journal of
Neurophysiology **76**(1): 492-509.

Bastian, A. J. and W. T. Thach (1995). "Cerebellar outflow lesions: A comparison of
movement deficits resulting from lesions at the levels of the cerebellum and
thalamus." Annals of Neurology **38**(6): 881-892.
Bastian, A. J., K. M. Zackowski and W. T. Thach (2000). "Cerebellar Ataxia: Torque
Deficiency or Torque Mismatch Between Joints?" Journal of Neurophysiology **83**(5):
3019-3030.
Beer, R. F., J. P. A. Dewald and W. Z. Rymer (2000). "Deficits in the coordination of
multijoint arm movements in patients with hemiparesis: evidence for disturbed
control of limb dynamics." Experimental Brain Research **131**(3): 305-319.
Beppu, H., M. Suda and R. Tanaka (1984). "ANALYSIS OF CEREBELLAR MOTOR
DISORDERS BY VISUALLY GUIDED ELBOW TRACKING MOVEMENT." Brain **107**(3):
787-809.
Bernshteĭn, N. A. (1967). The co-ordination and regulation of movements. Oxford,
Pergamon Press Oxford.
Boose, A., J. Dichgans and H. Topka (1999). "Deficits in phasic muscle force
generation explain insufficient compensation for interaction torque in cerebellar
patients." Neuroscience Letters **261**(1): 53-56.
Bremer, F., G.-H. Roger and L. Binet (1935). "Le cervelet." Traité de physiologie: 39-
134.
Chambers, W. W., Jr. (1947). "Electrical stimulation of the interior of the cerebellum
in the cat." Am. J. Anat. **80**(1): 55-93.
Cheng, E. J. and S. H. Scott (2000). "Morphometry of Macaca mulatta forelimb. I.
Shoulder and elbow muscles and segment inertial parameters." Journal of
Morphology **245**(3): 206-224.
Day, B. L., P. D. Thompson, A. E. Harding and C. D. Marsden (1998). "Influence of
vision on upper limb reaching movements in patients with cerebellar ataxia." Brain
121(2): 357-372.
Fagg, A., G. Ojakangas, L. Miller and N. Hatsopoulos (2009). "Kinetic Trajectory
Decoding Using Motor Cortical Ensembles." IEEE transactions on neural systems
and rehabilitation engineering : a publication of the IEEE Engineering in Medicine
and Biology Society **17**: 487-496.
Freund, H.-J. (1985). "Clinical aspects of premotor function." Behavioural Brain
Research **18**(2): 187-191.
Freund, H. J. and H. Hummelsheim (1985). "Lesions of premotor cortex in man."
Brain : a journal of neurology **108 (Pt 3)**: 697-733.
Fulton, J. F. and R. S. Dow (1937). "The Cerebellum: A Summary of Functional
Localization *." The Yale Journal of Biology and Medicine **10**: 89 - 119.
Goldstein, K. (1927). "Das Kleinhirn." Handbuch der normalen und pathologischen
Physiologie **10**: 222-222.
Granit, R., B. Holmgren and P. A. Merton (1955). "The two routes for excitation of
muscle and their subservience to the cerebellum." The Journal of Physiology **130**(1):
213-224.
Haines, D. E., M.-U. Manto and M. Glickstein (2007). "Clinical symptoms of cerebellar
disease and their interpretation." The Cerebellum **6**(2): 141-156.

Hare, W. K., H. W. Magoun and S. W. Ranson (1936). "ELECTRICAL STIMULATION OF THE INTERIOR OF THE CEREBELLUM IN THE DECEREBRATE CAT." American Journal of Physiology-Legacy Content **117**(2): 261-266.

Haynes, W. (2013). Benjamini–Hochberg Method. Encyclopedia of Systems Biology. W. Dubitzky, O. Wolkenhauer, K.-H. Cho and H. Yokota. New York, NY, Springer New York**:** 78-78.

Hollerbach, J. M. and T. Flash (1982). "Dynamic interactions between limb segments during planar arm movement." Biological Cybernetics **44**(1): 67-77.

Holmes, G. (1917). "The symptoms of acute cerebellar injuries due to gunshot injuries." Brain: A Journal of Neurology **40**: 461-535.

Holmes, G. (1939). "THE CEREBELLUM OF MAN1." Brain **62**(1): 1-30.

Hore, J. and D. Flament (1988). "Changes in motor cortex neural discharge associated with the development of cerebellar limb ataxia." Journal of Neurophysiology **60**(4): 1285-1302.

Ivry, R. B. and S. W. Keele (1989). "Timing Functions of The Cerebellum." Journal of Cognitive Neuroscience **1**(2): 136-152.

Jeannerod, M. (1988). The neural and behavioural organization of goal-directed movements. New York, NY, US, Clarendon Press/Oxford University Press.

Kawato, M. (1999). "Internal models for motor control and trajectory planning." Current Opinion in Neurobiology **9**(6): 718-727.

Konczak, J., D. Pierscianek, S. Hirsiger, U. Bultmann, B. Schoch, E. R. Gizewski, D. Timmann, M. Maschke and M. Frings (2010). "Recovery of Upper Limb Function After Cerebellar Stroke." Stroke **41**(10): 2191-2200.

Konczak, J., B. Schoch, A. Dimitrova, E. Gizewski and D. Timmann (2005). "Functional recovery of children and adolescents after cerebellar tumour resection." Brain **128**(6): 1428-1441.

Luciani, L. (1893). "Il Cervelletto. Nuovi Studi di Fisiologia Normale e Patologica." Philosophical Review **2**(4).

Magoun, H. W., W. K. Hare and S. W. Ranson (1935). "ELECTRICAL STIMULATION OF THE INTERIOR OF THE CEREBELLUM IN THE MONKEY." American Journal of Physiology-Legacy Content **112**(2): 329-339.

Manto, M., J. Bower, A. Conforto, J. Delgado-García, S. Guarda, M. Gerwig, C. Habas, N. Hagura, R. Ivry, P. Mariën, M. Molinari, E. Naito, D. Nowak, N. Taib, D. Pélisson, C. Tesche, C. Tilikete and D. Timmann (2011). "Consensus Paper: Roles of the Cerebellum in Motor Control-The Diversity of Ideas on Cerebellar Involvement in Movement." Cerebellum (London, England) **11**: 457-487.

Meyer-Lohmann, J., B. Conrad, K. Matsunami and V. B. Brooks (1975). "Effects of dentate cooling on precentral unit activity following torque pulse injections into elbow movements." Brain Research **94**(2): 237-251.

Miall, R. C., L. O. D. Christensen, O. Cain and J. Stanley (2007). "Disruption of State Estimation in the Human Lateral Cerebellum." PLOS Biology **5**(11): e316.

Miall, R. C. and D. King (2008). "State Estimation in the Cerebellum." The Cerebellum **7**(4): 572-576.

Nashef, A., O. Cohen, R. Harel, Z. Israel and Y. Prut (2019). "Reversible Block of
Cerebellar Outflow Reveals Cortical Circuitry for Motor Coordination." Cell Reports
27(9): 2608-2619.e2604.
Nashef, A., O. Cohen, Z. Israel, R. Harel and Y. Prut (2018). "Cerebellar Shaping of
Motor Cortical Firing Is Correlated with Timing of Motor Actions." Cell Reports
23(5): 1275-1285.
Nashef, A., H. Rapp, M. P. Nawrot and Y. Prut (2018). "Area-specific processing of
cerebellar-thalamo-cortical information in primates." Biological Cybernetics **112**(1):
141-152.
Nashold, B. S. and D. G. Slaughter (1969). "Effects of Stimulating or Destroying the
Deep Cerebellar Regions in Man." Journal of Neurosurgery **31**(2): 172-186.
Osu, R., K.-i. Morishige, J. Nakanishi, H. Miyamoto and M. Kawato (2015). "Practice
reduces task relevant variance modulation and forms nominal trajectory." Scientific
Reports **5**(1): 17659.
Perlmutter, J. S. and J. W. Mink (2006). "DEEP BRAIN STIMULATION." Annual
Review of Neuroscience **29**(Volume 29, 2006): 229-257.
Perlmutter, S. I., M. A. Maier and E. E. Fetz (1998). "Activity of Spinal Interneurons
and Their Effects on Forearm Muscles During Voluntary Wrist Movements in the
Monkey." Journal of Neurophysiology **80**(5): 2475-2494.
Plamondon, R. and A. M. Alimi (1997). "Speed/accuracy trade-offs in target-directed
movements." Behavioral and Brain Sciences **20**(2): 279-303.
Popa, L., A. Hewitt and T. Ebner (2013). "Purkinje Cell Simple Spike Discharge
Encodes Error Signals Consistent with a Forward Internal Model." Cerebellum
(London, England) **12**.
Pressman, P. and H. J. Rosen (2015). Chapter 33 - Disorders of Frontal Lobe
Function. Neurobiology of Brain Disorders. M. J. Zigmond, L. P. Rowland and J. T.
Coyle. San Diego, Academic Press**:** 542-557.
Ruigrok, T. J. H. (2013). Cerebellar Influences on Descending Spinal Motor Systems.
Handbook of the Cerebellum and Cerebellar Disorders. M. Manto, J. D. Schmahmann,
F. Rossi, D. L. Gruol and N. Koibuchi. Dordrecht, Springer Netherlands**:** 497-528.
Sainburg, R. L., C. Ghez and D. Kalakanis (1999). "Intersegmental Dynamics Are
Controlled by Sequential Anticipatory, Error Correction, and Postural Mechanisms."
Journal of Neurophysiology **81**(3): 1045-1056.
Sathyamurthy, A., A. Barik, C. I. Dobrott, K. J. E. Matson, S. Stoica, R. Pursley, A. T.
Chesler and A. J. Levine (2020). "Cerebellospinal Neurons Regulate Motor
Performance and Motor Learning." Cell Reports **31**(6): 107595.
Schlerf, J. E., J. Xu, N. M. Klemfuss, T. L. Griffiths and R. B. Ivry (2012). "Individuals
with cerebellar degeneration show similar adaptation deficits with large and small
visuomotor errors." Journal of Neurophysiology **109**(4): 1164-1173.
Schneider, R. C. and E. C. Crosby (1963). "The Interplay Between Cerebral
Hemispheres and Cerebellum in Relation to Tonus and Movements." Journal of
Neurosurgery **20**(3): 188-198.
Scott, S. H. (2004). "Optimal feedback control and the neural basis of volitional
motor control." Nature Reviews Neuroscience **5**(7): 532-545.

Shadmehr, R. and F. Mussa-Ivaldi (1994). "Adaptive representation of dynamics
during learning of a motor task." <u>The Journal of Neuroscience</u> **14**(5, Pt 2): 3208-
3224.
Shidara, M., K. Kawano, H. Gomi and M. Kawato (1993). "Inverse-dynamics model
eye movement control by Purkinje cells in the cerebellum." <u>Nature</u> **365**(6441): 50-
52.
Shinoda, Y., M. Yamazaki and T. Futami (1982). "Convergent inputs from the dentate
and the interpositus nuclei to pyramidal tract neurons in the motor cortex."
<u>Neuroscience Letters</u> **34**(2): 111-115.
[dataset] Sinha, N., S. Israely, O. Ben Harosh, R. Harel, J. P. A. Dewald and Y. Prut
(2024). Disentangling-acute-motor-deficits-and-adaptive-responses-evoked-by-the-
loss-of-cerebellar-output - (Version 1.0.0) https://doi.org/10.5281/zenodo.1234
Smith, J. L. and R. F. Zernicke (1987). "Predictions for neural control based on limb
dynamics." <u>Trends in Neurosciences</u> **10**(3): 123-128.
Therrien, A. S. and A. J. Bastian (2015). "Cerebellar damage impairs internal
predictions for sensory and motor function." <u>Current Opinion in Neurobiology</u> **33**:
127-133.
Timmann, D., S. Watts and J. Hore (1999). "Failure of Cerebellar Patients to Time
Finger Opening Precisely Causes Ball High-Low Inaccuracy in Overarm Throws."
<u>Journal of Neurophysiology</u> **82**(1): 103-114.
Topka, H., J. Konczak and J. Dichgans (1998). "Coordination of multi-joint arm
movements in cerebellar ataxia:Analysis of hand and angular kinematics."
<u>Experimental Brain Research</u> **119**(4): 483-492.
Topka, H., J. Konczak, K. Schneider, A. Boose and J. Dichgans (1998). "Multijoint arm
movements in cerebellar ataxia: Abnormal control of movement dynamics."
<u>Experimental Brain Research</u> **119**: 493-503.
Tsujimoto, T., H. Gemba and K. Sasaki (1993). "Effect of cooling the dentate nucleus
of the cerebellum on hand movement of the monkey." <u>Brain Research</u> **629**(1): 1-9.
Virji-Babul, N. and J. D. Cooke (1995). "Influence of joint interactional effects on the
coordination of planar two-joint arm movements." <u>Experimental Brain Research</u>
103(3): 451-459.
Weeks, H. M., A. S. Therrien and A. J. Bastian (2017). "Proprioceptive Localization
Deficits in People With Cerebellar Damage." <u>The Cerebellum</u> **16**(2): 427-437.
Wild, B. and J. Dichgans (1993). "Cerebellar ataxia in ataxic hemiparesis? A
kinematic and EMG analysis." <u>Movement Disorders</u> **8**(3): 363-366.
Wolpert, D. M., Z. Ghahramani and M. I. Jordan (1995). "An Internal Model for
Sensorimotor Integration." <u>Science</u> **269**(5232): 1880-1882.
Wolpert, D. M., R. C. Miall and M. Kawato (1998). "Internal models in the
cerebellum." <u>Trends in Cognitive Sciences</u> **2**(9): 338-347.

Figures and Tables

Figure 1: Effect of cerebellar block on peak hand velocity. (a) Schematic of the monkey performing the delayed reaching task in the KINARM exoskeleton to eight peripheral targets on the horizontal plane. The locations of the eight targets are numbered. (b) Movement-onset aligned hand velocity profiles during a sample control and cerebellar block trial performed by monkey C to target 2. Entry of the control cursor into the peripheral target marked the end of the trial. (c) Target-wise effect of the cerebellar block on peak hand velocity. For each session, the target-wise reduction in the median peak hand velocity during the cerebellar block trials was computed relative to that of control trials. The depicted values are the means ± 95% confidence intervals across all sessions pooled from all the four monkeys. The means of individual monkeys are overlaid. The dashed circle indicates no change. Statistical significance is denoted as follows: $p \geq 0.05$NS, $p < 0.05$*, $p < 0.01$**, $p < 0.001$***. [T1-8: Targets 1-8]

Figure 2: Effect of cerebellar block on muscle torques. Movement-onset aligned shoulder (a) and elbow (b) muscle torque profiles during a sample control and cerebellar block trial performed by monkey C to target 2. Entry of the control cursor into the peripheral target marked the end of the trial. (c) Normalized muscle torque impulse at the shoulder vs. elbow per target during control and cerebellar block. For each session, we computed the target-wise median muscle torque impulse during the acceleration phase of the movement across the control and cerebellar block trials. They were then normalized by the maximum absolute torque impulse across all the targets. The depicted values are the means across all the sessions pooled from the data of all four monkeys. The 95% confidence interval of the means is indicated by the horizontal and vertical bars for the shoulder and elbow joints respectively. Target-wise effect of cerebellar block on shoulder (d) and elbow (e) muscle torque impulse. For each session, the target-wise reduction in the median torque impulse during the cerebellar block trials was computed relative to the maximum absolute value of the target-wise medians observed during the control trials. The depicted values are the means ± 95% confidence intervals across all sessions pooled from all four monkeys. The means of individual monkeys are

overlaid. The dashed circle indicates no change. Statistical significance is denoted as follows: p ≥
0.05NS, p < 0.05*, p < 0.01**, p < 0.001***. [T1-8: Targets 1-8]

Figure 3: Effect of cerebellar block on coupling torques. Movement-onset aligned shoulder (a) and
elbow (b) coupling torque profiles during a sample control and cerebellar block trial performed by
monkey C to target 2. Entry of the control cursor into the peripheral target marked the end of the trial.
(c) Normalized coupling torque impulse at the shoulder vs. elbow per target during control and
cerebellar block. For each session, we computed the target-wise median coupling torque impulse
during the acceleration phase of the movement across the control and cerebellar block trials. They
were then normalized by the maximum absolute torque impulse across all the targets. The depicted
values are the means across all the sessions pooled from the data of all four monkeys. The 95%
confidence interval of the means is indicated by the horizontal and vertical bars for the shoulder and
elbow joints respectively.(d) Change in peak hand velocity due to cerebellar block plotted against the
coupling torque in control for each session for the reaching movements (T1-4). The target-wise
reduction in the median peak hand velocity during the cerebellar block trials was computed relative
to that of control trials. The target-wise median coupling torque during the control trials of each
session was normalized to the median across all trials of all sessions per monkey. The depicted data
contains all the sessions pooled from all four monkeys. The black line overlayed on the scatter indicates
the least-squares linear regression fit to the data points. [T1-8: Targets 1-8]

Figure 4: Effect of the cerebellar block on hand velocity across successive trials. (a) Schematic of the process of extraction of peak hand velocity for movements to a particular target from blocks of control/cerebellar block trials while preserving the order in which the target was presented. For each block, the peak velocity for movements to a particular target (in this example target 1) was extracted while retaining the trial numbers in which it was presented in the block. This enabled us to examine the evolution of the peak velocity across the sequence of presentation of the target during control vs. cerebellar block. (b) Mean ± 95% confidence intervals of the peak hand velocity for movements to target 1 across all the trial blocks pooled from all four monkeys. For each monkey, the peak hand velocities were normalized to the median peak velocity of the early trials 1-2 from all the control blocks. (c) Same as in (b) but for targets 2-4. (d) Post hoc comparisons between early trials 1-2 and late trials 11-20 during control and cerebellar block. Statistical significance is denoted as follows: $p \geq 0.05$NS, $p < 0.05$*, $p < 0.01$**, $p < 0.001$***. [E: Early, L: Late, C: Control, Cbl: Cerebellar block; T1-8: Targets 1-8]

Figure 5a-c: Effect of the cerebellar block on the decomposition of dual-joint movements. (a)
Binary decomposition raster of sample trials for movements made to target 2 by monkey C during
control. The duration of each trial was normalized to 0-1 (bin width = 0.001) and then decomposition
was computed for each bin based on whether either (but not both) of the shoulder or the elbow joint
velocity was less than 20°/s. (b) Same as in (a) but for movements to target 2 by monkey C during
cerebellar block. (c) Change in decomposition index (i.e., the proportion of the movement time during
which the movement was decomposed, as defined above) for movements to targets 2-4 during
velocity-matched control and cerebellar block trials relative to all control trials. The change in median
decomposition was computed for each session. The depicted values are the mean ± 95% confidence
intervals across all sessions pooled from all four monkeys. The individual means of each monkey are
overlaid. Figure 5d-e: Effect of the cerebellar block on inter-trial trajectory variability. (d)
Sample hand trajectories during control vs. cerebellar block trials for movements to target 2 by
monkey C. The starting point of the trajectories was shifted to the origin and then they were rotated
about the origin so that their endpoints lie on the positive Y-axis. (e) Change in Inter-trial trajectory
variability for movements to targets 2-4 during velocity-matched control and cerebellar block trials

relative to all control trials. The trajectory variability was measured as the standard deviation of the
maximum perpendicular distance of the trajectories from the Y-axis after transforming them as in (d).
The change in trajectory variability for the cerebellar block trials was computed relative to the control
trials for each session. The depicted values are the mean ± 95% confidence intervals across all sessions
pooled from all four monkeys. The individual means of each monkey are overlaid. Statistical
significance is denoted as follows: $p \geq 0.05$NS, $p < 0.05$*, $p < 0.01$**, $p < 0.001$***. [C: Control, Cbl:
Cerebellar block; T1-8: Targets 1-8; DI: Decomposition index]