Disentangling acute motor deficits and adaptive responses evoked by the loss of cerebellar output

Nirvik Sinha[1,2,3*], **Sharon Israely**[1], **Ora Ben Harosh**[1], **Ran Harel**[4], **Julius PA Dewald**[2,3,5,6], **Yifat Prut**[1*]

[1]Edmond and Lily Safra Center for Brain Sciences, The Hebrew University of Jerusalem, 9190401, Jerusalem, Jerusalem, Israel; [2]Department of Physical Therapy and Human Movement Sciences, Northwestern University Feinberg School of Medicine, 60610, Chicago, Illinois, USA; [3]Interdepartmental Neuroscience Center, Northwestern University Feinberg School of Medicine, 60611, Chicago, Illinois, USA; [4]Department of Neurosurgery, Sheba Medical Center, 5262000, Tel Aviv, Tel Aviv, Israel; [5]Department of Biomedical Engineering, Northwestern University, 60208, Evanston, Illinois, USA; [6]Department of Physical Medicine & Rehabilitation, Northwestern University Feinberg School of Medicine, 60610, Chicago, Illinois, USA

*For correspondence:

Nirvik.Sinha@mail.huji.ac.il (NS)

yifat.prut@mail.huji.ac.il (YP)

Abstract

Patients with cerebellar damage experience various motor impairments, but the specific sequence of primary and compensatory processes that contribute to these deficits remains unclear. To clarify this, we reversibly blocked cerebellar outflow in monkeys engaged in planar reaching tasks. This intervention led to a spatially selective reduction in hand velocity, primarily due to decreased muscle torque, especially in movements requiring high inter-joint torque coupling. When examining repeated reaches to the same target, we found that the reduced velocity resulted from both an immediate deficit and a gradually developing compensatory slowing to reduce passive inter-joint interactions. However, the slowed hand velocity did not account for the fragmented and variable movement trajectories observed during the cerebellar block. Our findings indicate that cerebellar impairment results in motor deficits due to both inadequate muscle torque and an altered motor control strategy for managing impaired limb dynamics. Additionally, impaired motor control elevates noise, which cannot be entirely mitigated through compensatory strategies.

Introduction

The ability to coordinate well-timed movements greatly relies on the cerebellum as evident from studies of individuals with cerebellar ataxia (Beppu, Suda et al. 1984).

The cerebellum regulates the timing of movements by predicting the mechanical
interactions between adjacent joints (Ivry and Keele 1989, Manto, Bower et al. 2011),
a computation that requires an internal model of feedforward control (Shidara,
Kawano et al. 1993, Shadmehr and Mussa-Ivaldi 1994, Wolpert, Ghahramani et al.
1995, Wolpert, Miall et al. 1998, Kawato 1999, Sainburg, Ghez et al. 1999, Popa,
Hewitt et al. 2013). However, motor impairments exhibited by individuals with
cerebellar deficits extend much beyond poor motor timing. For example, during the
acute phase of cerebellar injury, there is a loss of muscle tone and weakness of
voluntary movements (Holmes 1917, Goldstein 1927, Holmes 1939). Cerebellar
patients also exhibit movement-related sensory deficits such as an impaired visual
perception of stimulus movement and perception of the limb position sense (i.e.,
proprioception or state estimation) during active movements (Therrien and Bastian
2015, Weeks, Therrien et al. 2017).

Clinical studies of cerebellar patients have yielded conflicting insights into the
pathophysiology caused by a loss of cerebellar function. Some researchers suggest
that motor impairments stem primarily from an inability to compensate for inter-
joint interactions (Bastian, Martin et al., 1996; Bastian, Zackowski et al., 2000), while
others argue that insufficient muscle torque generation is the central issue (Topka,
Konczak et al., 1998; Boose, Dichgans et al., 1999). These discrepancies likely arise
from the variability among individuals with chronic cerebellar deficits, where
primary and compensatory effects are intertwined and difficult to separate.
Moreover, individuals with cerebellar lesions often have differences in lesion etiology
and associated damage beyond the cerebellum itself. As a result, the specific sequence
of events that leads to motor impairments following the loss of cerebellar signals
remains unclear.

To investigate this issue, we trained monkeys to perform a planar reaching task with
arm support provided by an exoskeleton. We then reversibly disrupted cerebellar
communication with other neural structures using high-frequency stimulation (HFS)
of the superior cerebellar peduncle, assessing the impact of this perturbation on
subsequent movements. Although our approach primarily affects cerebellar output
to the motor cortex, it also disrupts fibers carrying input signals (e.g., spinocerebellar)
and pathways to various subcortical targets (e.g., cerebello-rubrospinal). Thus, our
manipulation broadly interferes with cerebellar communication. Previously, we have
shown that HFS can effectively and reversibly interfere with the normal outflow of
cerebellar signals (Nashef, Cohen et al. 2019) since this perturbation accurately
replicated the behavioral deficits identified in cerebellar patients (Bastian 1997) and
the changes in motor cortical activity that appear after dentate cooling in monkeys
(Meyer-Lohmann, Conrad et al. 1975). In the present study, we found that the loss of
cerebellar outflow led to reduced hand velocity during reaching movements,
reflecting two parallel underlying mechanisms. First, there was a reduction in muscle
torques, even for movement directions where inter-joint interactions were low.
Second, the reduction in hand velocity exhibited a spatial tuning and was greater in
directions that generated large inter-joint interactions, thereby reflecting an inability

to compensate for limb dynamics. The time course of these two mechanisms was identified by analyzing the sequences of movements made to the same target during the cerebellar block. During such movements, hand velocity was low initially, reflecting a primary muscle torque deficit, and declined further in successive trials, revealing an adaptive motor strategy to minimize inter-joint interactions. Finally, we found that the decrease in velocity during cerebellar block could not fully explain the noisy and decomposed movements observed under these conditions, which may be due to an impaired state estimate of the limb that leads to the faulty control of inter-joint interactions. Taken together, we show that cerebellar output is essential for generating sufficient muscle torques to allow for fast and stereotypical movements while overcoming the complex inter-joint interactions inherent in such actions. The loss of this output triggers a sequence of early primary effects followed by compensatory responses aimed at mitigating these impairments. Together these processes result in significantly slower movements.

Results

Slower reaching movements in cerebellar block are due to weakness and impaired limb dynamics.

We trained four adult female monkeys (Macaca fascicularis) to sit in a primate chair with their upper limb supported by an exoskeleton (KINARM system) and perform planar center-out movements (Figure 1a). The spatial location of the targets enabled us to dissociate between shoulder-dominant (lateral targets) vs. shoulder-elbow movements (diagonal and straight targets respectively). Each trial began with the monkey holding a cursor within a central target. After 500–800 ms, one of eight peripheral targets appeared (cue). Following a 450–850 ms delay, the central target disappeared ('go' signal), prompting the monkey to reach the peripheral target within 500-900 ms and briefly hold the cursor there (100-150 ms) to receive a reward. Between successive trials, monkeys repositioned the cursor back to the central target. The experiment was conducted across several sessions (i.e., days). Each session was divided into 3-4 sub-sessions. Each sub-session comprised of a block of control trials (~80) followed by a block of trials (~50) with high-frequency stimulation (130 Hz) of the superior cerebellar peduncle.

Cerebellar block significantly reduced the ability of the monkeys to perform the task as measured by the success rate across all the recorded sessions (mean success rate in control = 86.5%, CI [85.8, 87.2] vs. cerebellar block = 75.3%, CI [73.9, 76.7], p < 0.001; **Supplementary table S1**). Since the monkeys had to meet strict timing criteria to perform the movements successfully during the task, we began by examining the hand kinematics. **Figure 1b** shows profiles of hand velocity in sample trials collected during the control (blue) and cerebellar block (vermillion) conditions. This example shows that movements exhibited a significant reduction in the peak hand velocity during cerebellar block. However, when compared across the targets,

we found that the reduction in hand velocity was strongly modulated by the target
direction (1-way ANOVA of the change in peak hand velocity, the effect of cerebellar
block: p = 0.01, the effect of targets: p < 0.001; **Supplementary table S2**). *Post hoc*
comparisons further revealed that the effect of cerebellar block on outward reaching
movements (targets 1-4) i.e., movements involving shoulder flexion was significantly
higher than on inward reaching movements (targets 5-8) i.e., movements involving
shoulder extension (**Figure 1c**; mean decrease of 9.7%, CI [8.2, 11.2], p < 0.001 for
targets 1-4 compared to mean decrease of 2.5%, CI [0.9, 4.0], p = 0.21 for targets 5-8)
even though there was no difference in the peak hand velocity between the
movements to the two target groups during control (p = 0.88, **Supplementary tables**
S3a and b).

Previous studies have demonstrated that Purkinje neurons of the cerebellar cortex
exhibit weaker modulation during task-irrelevant movements compared to task-
related movements (Pi, Fakharian et al. 2024, Hage, Fakharian et al. 2025). Therefore,
we examined whether the impact of the cerebellar block was limited specifically to
target-directed, reward-associated movements or if it generalized to all movements
performed during the task. To this end, we compared peak hand velocities between
target-directed (within-trial, reward-relevant) movements and subsequent return-
to-center (out-of-trial) movements. Our analysis revealed that cerebellar block
resulted in a significantly larger reduction in peak velocity for target-directed
movements compared to return-to-center movements (7.5%, CI [6.5, 9.0] vs. 1.7%, CI
[-1.8, 3.0]; p < 0.001; **Supplementary Figure S1**). These findings suggest that the
cerebellum plays a preferential role in optimizing reward-associated, within-trial
movements rather than equally influencing all movements.

In the task used in our study, hand movements were made by controlling the muscle
torque at the shoulder and elbow joints (Scott 2004). However, simultaneous
movements of both joints also generated coupling torques due to passive interactions
between limb inertial properties and joint velocities (Bernshteĭn 1967, Hollerbach
and Flash 1982). To identify the target-specific muscle and coupling torques, we
employed an inverse dynamics model of the upper limb (Fagg, Ojakangas et al. 2009)
which enabled us to distinguish between the different contributions to the net torque
experienced by each joint. According to this model, the net torque at each joint is the
sum of both the active muscle torque and the passive coupling torque acting at that
joint (see Methods section for details).

Using the inverse dynamics model, we found that the target-specific reduction in
hand velocities was produced by changes in muscle commands (e.g., **Figures 2a, b**).
Specifically, we observed that the muscle torque impulses, calculated by integrating
the torque profiles over the early phase of movements when acceleration was
positive, were reduced in a target-specific manner during the cerebellar block
(**Figure 2c**; **Supplementary Table S4-5**). This reduction was particularly
pronounced at the shoulder joint. Furthermore, as in the case of hand velocities, *post*
hoc comparisons revealed that the shoulder joint experienced a significantly greater

reduction in muscle torque for targets 1-4 vs. 5-8 (**Figure 2d**; mean decrease of 10.6%, CI [9.0, 11.2], p < 0.001 for targets 1-4 compared to mean decrease of 3.3%, CI [-1.3, 5.2] p = 0.21 for targets 5-8). In contrast, the slight reduction in elbow torques due to the cerebellar block did not reach significance for any of the targets (**Figure 2e**). Collectively, these findings indicate that during the cerebellar block, there is a reduction in shoulder muscle torque specifically for targets involved in outward-reaching movements, which leads to the target-specific reduction in hand velocity.

Several studies have shown that individuals with cerebellar deficits exhibit an inability to correctly compensate for inter-joint interactions (Bastian, Martin et al. 1996, Bastian, Zackowski et al. 2000). Therefore, we asked whether the reduction in muscle torques and hand velocities due to the cerebellar block could be associated with the coupling torques generated during movements in various directions. **Figures 3a and b** show the coupling torque profiles at the shoulder and elbow joints respectively during sample trials. We observed that the coupling torque impulses varied in a target-dependent manner during both the control and cerebellar block conditions (**Figure 3c**). Furthermore, the reduction in peak hand velocities during cerebellar block was correlated to the net coupling torque impulse (calculated as the sum of absolute coupling torque impulses at both joints) during control, particularly for the outward reaching movements (**Figure 3d;** targets 1-4: partial correlation coefficient to account for the variability across monkeys, $\rho = 0.32$, CI [0.23, 0.41], p < 0.001). This indicated that the effect of the cerebellar block on hand velocity during the outward reaching movements was strongest for targets with high coupling torques. Nevertheless, despite the significant correlation between the reduced hand velocity and coupling torques, cerebellar block reduced the hand velocity even in outward reaches where the overall coupling torque was low i.e., movements to target 1, mean reduction in hand velocity = 5.8%, CI [2.8, 8.7], p < 0.001 (movements to all outward targets involved both shoulder and elbow joints, but the contribution of each joint varied in a target dependent manner. Specifically, movements to targets 1 exhibited negligible net torque at the elbow, meaning the elbow remained largely stationary during these movements as shown in Supplementary Figure S2. Consequently, the shoulder experienced correspondingly low coupling torques as shown in Figure 3c). This finding indicated a reduction in the motor command which occurred in addition and beyond the effect attributable to the impaired control of coupling torques alone.

Overall, the cerebellar block produced a significant impact on the kinematics and dynamics of outward reaching movements. Specifically, the cerebellar block caused slower hand velocities due to a systematic, target- and joint-specific reduction in muscle torques for these movements, which strongly correlated with the coupling torques. Additionally, a reduction in muscle torques was also observed during the outward reaching movements with low coupling torques, indicating an additional component of muscle torque insufficiency that further exacerbated motor impairment following cerebellar block.

Hand velocity exhibits an adaptive trend during repeated movements to the same target during cerebellar block.

The advantage of using a reversible cerebellar block in an animal model is the ability to track the temporal profile of motor impairments induced by this condition, allowing us to distinguish between primary and compensatory effects. To this end, we assessed changes in hand kinematics produced by the cerebellar block across successive trials. Specifically, we examined the evolution of peak hand velocities for outward reaches with low (target 1) vs. high coupling torques (data pooled for movements to targets 2-4) during blocks of control and cerebellar block trials. **Figure 4a** illustrates this procedure. Since the targets were presented in a pseudorandom order during each block of trials, we collected for each block the tested target, while keeping its order of presentation to generate a temporally accurate sequence of trials, all directed towards the same single target.

We found that during the control condition, peak hand velocity remained stable across successive trials. In contrast, cerebellar block affected hand velocity in a trial- and target-dependent manner (**Figure 4b, c**). For movements to target 1, hand velocity was lower than control but unaffected by the temporal sequence of target presentation (trial type × trial sequence interaction effect: p = 0.69, **Supplementary Table S6**). However, for movements to targets 2-4, hand velocity was lower than control in the early trials and declined further with successive trials (trial type × trial sequence interaction effect: p = 0.007, **Supplementary Table S7**).

To further quantify the effect of trial sequence on movements to targets 2-4, we performed *post hoc* comparisons (**Figure 4d**). These comparisons revealed that during the cerebellar block, hand velocity was 4.3% (CI [1.0, 7.3], p < 0.001) lower than control during the first 1-2 trials. With successive trials, the hand velocity declined further before reaching an asymptote during the late trials 11-20 at 8.2% (CI [5.7, 10.7], p < 0.001) lower than the control.

Taken together these results reveal that the hand velocity of outward-reaching movements with high coupling torques was already low initially during cerebellar block, reflecting a primary deficit. Over and above this, there was a further progressive decline of the hand velocity during successive movements, most likely due to an adaptive response to the inability to compensate for coupling torques. Furthermore, this adaptive response was not observed in the case of movements where the coupling torques were low, and therefore inter-joint interactions did not play a significant role.

Slow movements during the cerebellar block cannot fully account for the impaired motor timing.

Individuals with cerebellar deficits often exhibit impaired motor timing, leading to movements that are poorly coordinated. Normally, smooth execution of multi-joint movements relies on precise temporal synchronization across joints. However,

cerebellar lesions disrupt this temporal coordination, resulting in asynchronous,
sequential joint activation resulting in movement decomposition as described earlier
(Bastian, Martin et al. 1996). Moreover, the disrupted temporal control of joint
activation introduces variability from trial to trial, causing increased spatial
variability in the resulting movement trajectories (Day, Thompson et al. 1998). Thus,
movement decomposition and trajectory variability can both be viewed as direct
consequences of impaired temporal coordination due to cerebellar dysfunction.
These properties of movement are also velocity-dependent (Bastian, Martin et al.
1996, Osu, Morishige et al. 2015). Therefore, it is conceivable that the combination of
primary deficit and adaptive reduction in velocity to reduce the effects of coupling
torque observed in our results is the main driver of abnormal motor timing. To
evaluate this, we examined movement decomposition and trajectory variability
during the outward reaching movements (which exhibited a significant reduction in
hand velocity during cerebellar block) and measured the impact of hand velocity on
these measures.

To quantify movement decomposition, we normalized the movement times of
individual trials to range from 0 to 1 and identified the bins where either the shoulder
or the elbow joint paused (speed < 20°/s) while the other joint continued moving
(Bastian, Martin et al. 1996). In control trials, decomposition was primarily observed
during the early phase of movement, where joint velocities are low making this
measure more sensitive (**Figure 5a** and **Supplementary Figure S3a**). However,
during cerebellar block trials, decomposition occurred throughout the movement
(**Figure 5b** and **Supplementary Figure S3b**) similar to individuals with cerebellar
deficits (Bastian, Martin et al. 1996). To quantify this effect, we calculated a
decomposition index for each trial, representing the proportion of movement time
during which the movement was decomposed, as defined above. Overall, the
cerebellar block increased movement decomposition by 53.5% (CI [42.2, 66.0], p <
0.001). Next, we asked whether movement decomposition was mainly due to lower
hand velocities. We therefore selected a subset of control trials that matched the
cerebellar block trials in their peak velocities. Movement decomposition in these
control trials was significantly lower than velocity-matched cerebellar block trials (p
< 0.001; **Figure 5c;** see **Supplementary Figure S3c** for target-wise effects), even
though these control trials themselves had 11.0% (CI [5.2, 17.0], p = 0.03) higher
decomposition than all control trials together. This indicates that even at matched
velocities, movements during cerebellar block were more decomposed relative to the
control.

Next, we assessed the effect of cerebellar block and hand velocity on trajectory
variability. To this end, we first aligned the hand trajectories to have the same starting
position and then rotated them so that their endpoint was located on the positive Y-
axis (**Figure 5d;** for visualization in joint position space see **Supplementary Figures
S4a** and **b**). Subsequently, we defined the error in each trajectory as the maximal
perpendicular distance from the Y-axis. The standard deviation of the errors across
all trials was used to quantify the trial-to-trial trajectory variability. When we

compared the subset of velocity matched control and cerebellar block trials, we found that cerebellar block exhibited 34.6% (CI [26.2, 43.2], p < 0.001) higher trajectory variability (**Figure 5e;** see **Supplementary Figure S4c** for target-wise effects**)**. Normally, slower movements are also less variable due to the speed-accuracy tradeoff (Plamondon and Alimi 1997). Indeed, the trajectory variability in this subset of slower control trials was 5.5% (CI [0.9, 9.9], p = 0.02) lower than that of all control trials. In other words, despite slower movements, cerebellar block increased trajectory variability.

Finally, we quantified how cerebellar block affected movement decomposition across successive trials similar to our analysis of hand velocities in Figure 4. The cerebellar block induced increase in movement decomposition was unaffected by the trial-sequence (trial type × trial sequence interaction effect: p = 0.43, **Supplementary Table S8** and **Supplementary Figure S5**). This further demonstrates that movement decomposition remained high during cerebellar block despite slower movement speeds.

Discussion

Individuals with cerebellar deficits exhibit altered motor behavior related mostly to poor timing and coordination of voluntary movements. Understanding the impairments that follow cerebellar lesions is often complicated by the fact that the studies of this question mostly rely on a heterogeneous population due to various levels and types of cerebellar pathologies. Here we addressed this question by using high-frequency stimulation to reversibly block cerebellar output to the motor cortex in behaving monkeys wearing an exoskeleton (Nashef, Cohen et al. 2019) and comparing the post-block motor behavior to control conditions. Our results indicate that cerebellar block leads to decreased hand velocity, particularly during the outward reaching movements via two mechanisms: a reduction in muscle torques suggesting motor weakness, and spatially tuned decreases in velocity in targets where large inter-joint interactions occur, indicating a failure to compensate for limb dynamics. Analyzing the time course of these impairments revealed partially overlapping processes: an initial reduction in hand velocity due to the inability to generate sufficient muscle torques, followed by a further progressive decline in hand velocities during subsequent movements during the cerebellar block. This secondary decline was observed only in the case of movements with high inter-joint interactions, suggesting an adaptive strategy to minimize these interactions. Finally, we observed increased motor noise in terms of movement decomposition and trial-to-trial trajectory variability which were independent of the slower movements observed during the cerebellar block.

Target-dependent reduction in hand velocity and coupling torques during reaching movements during the cerebellar block.

Cerebellar block led to a target-specific reduction in hand velocity, even though the
arm was fully supported by the exoskeleton and thus the limb weight did not play any
role. This observation replicates findings made in previous studies that have
documented the slowing of movements in individuals with cerebellar lesions (Wild
and Dichgans 1993, Topka, Konczak et al. 1998, Konczak, Schoch et al. 2005) as well
as in monkeys that underwent dentate cooling (Tsujimoto, Gemba et al. 1993). It was
argued that the reduction in hand velocity may be a compensatory strategy to
counteract the impaired ability to manage inter-joint coupling torques in the absence
of cerebellar input to the motor cortex (Wild and Dichgans 1993, Bastian and Thach
1995, Bastian, Martin et al. 1996, Beer, Dewald et al. 2000). Since coupling torques in
multi-joint movements are generated in a velocity-dependent manner (Hollerbach
and Flash 1982, Virji-Babul and Cooke 1995), slowing down the movement not only
simplifies its kinetics (by reducing the impact of coupling torques) but also allows
more time for slower visual feedback to aid in guiding the movement (Jeannerod
1988). Our findings, which demonstrate a correlation between the extent of hand
velocity reduction during cerebellar block and the magnitude of coupling torque
under control conditions, further corroborate this hypothesis and confirm our
approach for reversible inactivation of cerebellar output as a model for ataxia.
Furthermore, our model enabled us to study this effect rigorously and systematically
while dissociating it from the other primary effects of cerebellar block as discussed
below.

Muscle torque deficit is an acute response to the cerebellar block.

Poor compensation for coupling torque is a fundamental deficit in cerebellar ataxia
(Bastian, Zackowski et al. 2000). However, early studies have reported the
occurrence of muscle weakness (asthenia) and hypotonia acutely following
cerebellar injury in humans (Haines, Manto et al. 2007) or experimental lesions in
animals (Luciani 1893, Bremer, Roger et al. 1935, Fulton and Dow 1937, Granit,
Holmgren et al. 1955). Both Luciani (Luciani 1893) and Holmes (Holmes 1917)
described a triad of cerebellar deficits to be atonia (low muscle tone), asthenia
(weakness of voluntary movements), and astasia (oscillation of the head and trunk).
Importantly, muscle weakness is typically characteristic of the acute phase of injury,
especially in the upper extremity, and tends to normalize gradually over weeks to
months, depending on the severity of the injury (Konczak, Pierscianek et al. 2010). It
is therefore conceivable that studies in humans with cerebellar deficits, which often
included subjects at the chronic stage of the lesion, may have underestimated the
contribution of muscle weakness which by that time was already less pronounced.

We explored the acute outcome of the cerebellar block, a phase in which agonist
muscle weakness may play a more prominent role. Importantly, we showed that
deficits in the self-generated muscle torque appear immediately after the cerebellar
block, resulting in lower hand velocity already in the first few trials of movements.
This kind of fast response is less likely to be a strategic response aimed at mitigating
the loss of cerebellar signals. This finding is consistent with the initial weakness

followed by a gradual recovery of function observed in acute lesions of other neural
structures involved in movement control, including the primary (Pressman and
Rosen 2015) and premotor cortices (Freund 1985, Freund and Hummelsheim 1985).
It also aligns with the understanding that cerebellar output provides a strong
excitatory drive to the motor cortex (Shinoda, Yamazaki et al. 1982, Hore and Flament
1988, Nashef, Cohen et al. 2018) and several subcortical descending motor systems
(Ruigrok 2013), beyond its well-known role in motor timing and coordination.
Following the initial reduction in hand velocity observed in our study, there was a
further gradual decline across subsequent trials specifically for the movements with
high coupling torques, a behavior profile that is more consistent with strategic effort,
most likely in response to the inability to compensate for limb dynamics.

The muscle torque, representing the sum of all muscle forces acting at a joint, can
also be reduced by co-contraction of agonist and antagonist muscles rather than by
agonist weakness alone. However, in cerebellar ataxia, co-contraction has been
proposed as a strategy to stabilize stationary joints during decomposed multi-joint
movements (Bastian, Martin et al. 1996). In our experiments, this strategy would
likely emerge gradually following cerebellar block similar to the adaptive slowing of
movements aimed at reducing inter-joint interactions. We therefore argue that the
observed acute deficiency in muscle torque was mainly driven by agonist weakness.
Our argument is further supported by previous studies which attributed reduced
agonist muscle activity as a cause for slowing of voluntary movements in individuals
with cerebellar lesions (Hallett, Berardelli et al. 1991, Wild, Klockgether et al. 1996).

In summary, the slower movements in the absence of cerebellar signals are likely
due to two main factors: (i) a primary inability to generate sufficient muscle torques
due to the lack of synchronized recruitment of the cortical and sub-cortical
descending motor systems via the cerebellar output, and (ii) a secondary adaptive
strategy aimed at compensating for the impaired prediction of limb dynamics due to
the loss of signals from the cerebellum.

Insufficient muscle torques vs. poor inter-joint coordination.

Unlike most previous clinical studies that examined motor deficits in individuals
with chronic cerebellar injuries, our study utilized a reversible model of cerebellar
block. This approach enabled us to track the temporal evolution of motor deficits
from the onset of the block within the same study participant. Although several
studies have explored the nature of motor deficits following cerebellar injuries, there
remains considerable debate regarding the fundamental mechanisms driving these
deficits. Some studies have suggested that the inability to compensate for inter-joint
interactions is the primary driver of the observed motor abnormalities in cerebellar
injury (Bastian, Martin et al. 1996, Bastian, Zackowski et al. 2000). In contrast, other
studies have emphasized the inability to generate sufficient muscle torques as the
core deficit in cerebellar ataxia while claiming that inter-joint coordination is
unaffected following cerebellar lesions (Topka, Konczak et al. 1998, Boose, Dichgans

et al. 1999). Given these conflicting findings, our study aimed to systematically explore the entire spectrum of motor deficits using a model of reversible cerebellar block. First, we found that the deficits in muscle torque are more pronounced for the reaching movements as compared to the retrieval movements. We identified both an acute insufficiency in muscle torque generation and the inability to compensate for coupling torques as potential mechanisms underlying the observed slowness during the reaching movements. Moreover, we delineated the time course of these deficits, providing insights into their evolution from the onset of cerebellar block.

The asymmetric effect of cerebellar block on outward vs. inward reaching movements.

We observed a notable distinction in the impact of cerebellar blockage on outward vs. inward reaching movements. At a collective level, this variance could be attributed to the disparity in magnitude of coupling torque impulse, with movements towards targets 2-4 exhibiting 18.7% (CI [13.1, 24.1], p = 0.008) higher coupling torque impulse compared to those towards targets 6-8. However, this explanation does not hold consistently when examined at the individual target level. For instance, while the coupling torque was higher for movements towards target 7, the effects of cerebellar blockage were less pronounced compared to those observed for target 1, where the coupling torques are considerably lower. There are two possible explanations for this discrepancy. One possibility is that the effect of cerebellar block exhibits a bias in the direction of shoulder flexion (i.e., for movements to targets 1-4). Several early studies in cats (Hare, Magoun et al. 1936, Chambers 1947), non-human primates (Magoun, Hare et al. 1935) , and humans (Nashold and Slaughter 1969) have all reported a bias towards the activation of ipsilateral flexors of the arm following stimulation of the deep cerebellar nuclei and the superior cerebellar peduncle. Furthermore, lesioning of these structures led to the reduction of ipsilateral flexor tone and power (Schneider and Crosby 1963, Nashold and Slaughter 1969). The cerebellar output influences the excitability of spinal interneurons either directly (Bantli and Bloedel 1975, Asanuma, Thach et al. 1980, Sathyamurthy, Barik et al. 2020) or via indirect relays in the subcortical descending motor systems like the rubrospinal and the reticulospinal pathways (Ruigrok 2013). Since the spinal interneurons influence the activity of flexor muscles twice as often as extensor muscles (Perlmutter, Maier et al. 1998), this might explain why the cerebellar block had a stronger effect on movements involving shoulder flexion (i.e., outward reaching movements) relative to shoulder extension (i.e., inward reaching movements) in our study. Alternatively, the initial position of the limb in our experimental setup (i.e., on the central target) might have been favorable for movements in the direction of shoulder extension, thereby making them easier to perform.

Asynchronous joint movements and motor variability occur despite slower movements in cerebellar block.

Abnormally high motor variability (Day, Thompson et al. 1998, Timmann, Watts et
al. 1999, Schlerf, Xu et al. 2012) and decomposition of multi-joint movements
(Bastian, Martin et al. 1996) are core deficits in cerebellar patients. We showed that
the cerebellar block decomposed the dual-joint movements into temporally isolated
single-joint actions and led to an increased trial-to-trial spatial noise. These two
measures can be affected by the reduction in movement velocities. Our results
showed that even though movement decomposition increased with a reduction in
velocity in control trials, they were still significantly lower than that of velocity-
matched cerebellar block trials. Second, the trial-to-trial spatial noise in hand
trajectories increased during cerebellar block despite the slower movement speeds.
The cerebellum is known to participate in the planning of movements by providing
an internal model of the limb which is necessary for predicting the sensory outcomes
of motor commands (Wolpert, Miall et al. 1998). For this purpose, the cerebellum aids
in estimating the state of the limb by integrating the sensory information of its last
known state with predictions of its response to the latest motor command (Miall,
Christensen et al. 2007, Miall and King 2008). Therefore, in the absence of cerebellar
signals, there are programming errors in the motor command required to launch the
limb accurately toward a target (Day, Thompson et al. 1998). These errors may in
turn translate into the noisy joint activation patterns and variable hand trajectories
observed in our results. Moreover, the persistence of these deficits independently of
the movement speeds observed during cerebellar block indicates that they cannot be
compensated for, unlike the impaired prediction of inter-joint interactions which are
strategically dealt with by reducing the movement velocity.

In summary, our study systematically investigated the temporal progression and
distinct mechanisms of motor deficits in a controlled, reversible animal model of
cerebellar block. These include an acute onset muscle torque deficit, impaired
compensation of limb dynamics, and increased motor noise. We aim to further
explore the neural correlates of these deficits in future studies.

Materials and Methods

Experimental subjects

477   This study was performed on four adult female monkeys (*Macaca fascicularis*,
478   weight 4.5-8 kg). The care and surgical procedures of the subjects will be in
479   accordance with the Hebrew University Guidelines for the Use and Care of Laboratory
480   Animals in Research, supervised by the Institutional Committee for Animal Care and
481   Use.

Behavioral task

Each monkey was trained to sit in a primate chair and perform planar, center-out
movements. During the task, the left upper arm was abducted 90° and rested on
cushioned troughs secured to links of a two-joint exoskeleton (KINARM, BKIN

Technologies Ltd.). The sequence of events in each trial of this task was as follows.
First, the monkey located a cursor (projected on a horizontal screen immediately
above its arm) within a central target (2-3 cm in diameter). After 500-800 ms, a
randomly selected peripheral target (out of 8 evenly distributed targets) was
displayed. The monkey had to wait 450-850 ms until the central target disappeared
(i.e., the 'go signal') and then reached the cued peripheral target (2-3 cm in diameter)
located 4-5 cm away. For monkeys C and M, we inserted a 200 ms grace period before
the 'go' signal to encourage predictive timing. The onset of movement within this time
before the 'go' signal did not abort the trial. We limited the time to reach the
peripheral target after the 'go' signal to 500 ms for monkeys M and C and 800-900 ms
for monkeys S and P. Following successful reach, the monkeys were rewarded with a
drop of applesauce. It is important to note that the monkeys were required to hold at
the target briefly—100 ms for Monkeys S and P, and 150 ms for Monkeys C and M—
before receiving the reward. However, given the size of the targets and the velocity of
movements, it often happened that the monkeys didn't have to stop their movements
fully to obtain the reward. Importantly, we relaxed the task's requirements (by
increasing the target size and reducing the temporal constraints) to enable the
monkeys to perform a sufficient number of successful trials under both the control
and the cerebellar block conditions. This was necessary as we found that strict
criteria for these parameters yielded a very low success rate in the cerebellar block
condition. Given these practical constraints, we note that this task design is likely
suboptimal for studying endpoint accuracy which is an important aspect of cerebellar
control (Low, Thanawalla et al. 2018, Becker and Person 2019, Calame, Becker et al.
2023).

Insertion of the stimulating electrode into the superior cerebellar peduncle

After the training was completed, a square recording chamber (24x24 mm) was
attached to the monkey's skull above the upper limb-related area of the motor cortex
in a surgical procedure under general anesthesia. To insert a chronic stimulating
electrode into the ipsilateral superior cerebellar peduncle (SCP), a round chamber
(diameter = 19 mm) was also implanted above the estimated insertion point (as per
stereotactic measurements) during the surgery for the cortical chamber
implantation. Subsequently, a post-surgery MRI was used to plan the trajectory for
the electrode insertion into the SCP. After a recovery and re-training period, a bi-polar
concentric electrode (Microprobes for Life Science Inc.) was inserted through the
round chamber along the planned trajectory. During the process of insertion, the
evoked multi-unit activity in the primary motor cortex was simultaneously
monitored to identify the precise location where the stimulating electrode should be
secured (Nashef, Cohen et al. 2018, Nashef, Rapp et al. 2018).

High-frequency stimulation to block the cerebellar outflow to the motor cortex

The superior cerebellar peduncle was blocked using high-frequency stimulation
(HFS). This stimulation protocol consisted of a train of biphasic rectangular-pulse

stimuli (where each phase was 200 μs) applied at 130 Hz. Each train was delivered at
a fixed intensity of 150-250 μA. In our earlier study, we demonstrated that by using
this manipulation we can successfully replicate the impaired motor timing and
coordination characteristic of cerebellar ataxia (Nashef, Cohen et al. 2019).
Furthermore, we also demonstrated that at the motor cortical level, this manipulation
leads to a loss of response transients at movement onset and decoupling of task-
related activity, similar to findings observed in studies where the dentate nucleus was
cooled (Meyer-Lohmann, Conrad et al. 1975). Deep brain stimulation (DBS) is a
perturbation similar in properties to the HFS we used, which is routinely used in
clinical practice for similar purposes of interfering with neural circuitries that
underwent pathological changes (Perlmutter and Mink 2006) although in both
applications there is no accurate measure for the extent of the block (i.e., the fraction
of fibers which become ineffective in these conditions) other than the behavioral
consequences. It is possible that HFS probably imposes only a partial block of the CTC
system.

It is important to note that the superior cerebellar peduncle is known to carry input
and output fibers to and from several neural structures (Naidich, Stein et al. 2013).
Therefore, we cannot rule out the fact that the effect of HFS may be mediated in part
through pathways other than the cerebello-thalamo-cortical pathway (as mentioned
in the Discussion section). However, in primates, the cerebellar-thalamo-cortical
pathway greatly expanded (at the expense of the cerbello-rubro-spinal tract) in
mediating cerebellar control of voluntary movements (Horne and Butler 1995). The
cerebello-subcortical pathways diminished in importance throughout evolution
(Padel, Angaut et al. 1981, Nathan and Smith 1982, ten Donkelaar 1988). Previously
we found that the ascending spinocerebellar axons which enter the cerebellum
through the superior cerebellar peduncle (SCP) are weakly task-related and the
descending system is quite small (Cohen et al, 2017). Nevertheless, we acknowledge
that HFS disrupts cerebellar communication to the rest of the nervous system
broadly, rather than solely the cerebello-thalamo-cortical pathway.

Data collection protocol

Each monkey performed the experimental task for several days (sessions). A total
of 46, 29, 36, and 54 sessions of data were collected from Monkeys S, C, M, and P,
respectively. Each session was divided into three or four sub-sessions. Each sub-
session included blocks of control trials (~80 trials) and trials with the superior
cerebellar peduncle blocked using high-frequency stimulation (~50 trials) as
described above. During the task performance, the following behavioral data was
obtained at 1000 Hz using the KINARM's motor encoders: (i) Hand position in the x-
y plane of movement, (ii) Elbow and shoulder joint angular positions, velocities, and
accelerations. The lengths of the upper arm and forearm of each monkey were
measured as the distances from the head of the humerus to the lateral epicondyle and
from the lateral epicondyle to the palm, respectively. The inertial properties of the
upper limb were estimated using the monkey's body weight (Cheng and Scott 2000).

Processing of the behavioral data

All data analysis was performed in MATLAB R2021b (MathWorks Inc.). The raw kinematic data was low-pass filtered by using a second order Butterworth filter (cutoff = 10Hz) and then epoched into trials. For this purpose, the start and end events were taken as the time of presentation of the cue (i.e., the peripheral target) and the time of entry of the control cursor into the peripheral target, respectively. This cutoff meant that a major portion of the extracted segment of each trial was the acceleration phase of the movement. The movement onset was defined as the time when the radial hand velocity exceeded 5% of its peak. Subsequently, all the trials were zero-centered around the calculated movement onset. We then applied a two-link inverse dynamics model of the upper arm to the kinematic data to compute the coupling and muscle torques acting on the elbow and shoulder joints (Fagg, Ojakangas et al. 2009). The inverse dynamics equations of motion for the combined system consisting of the monkey's arm and the associated moving components of the KINARM were used to compute the joint torques:

$$T_{me} = T_{ne} - T_{ce} \tag{i}$$

$$T_{ms} = T_{ns} - T_{cs} \tag{ii}$$

where T_{ne} and T_{ns} are the net torques (proportional to the joint's acceleration), T_{ce} and T_{cs} are passive coupling torques, and T_{me}, T_{ms} are the active muscle torques acting at the elbow and shoulder joints respectively. The expanded formulas for computing each of the above terms are provided by equations 1a-2e in the methods section C (Fagg, Ojakangas et al. 2009). It must be noted that the 'muscle' torques computed here include the effects of the actively generated muscle forces as well as the viscoelastic effects that result from the musculoskeletal system and the KINARM (Smith and Zernicke 1987). Data from some of the same animals was presented in prior work where we showed that during reaching, cerebellar block resulted in reduced reach velocity, increased trajectory curvature, and shoulder-elbow dyscoordination (Nashef, Cohen et al. 2019). In this study, we extend these findings by using inverse-dynamics modeling to explore the relationship between altered kinematics and joint torques. Furthermore, the analyses exploring progressive velocity changes across trials under the cerebellar block and the relationship of motor noise to movement velocity are newly reported in this study.

To strictly focus on feedforward control, we could have examined the measured variables in the first 50-100 ms of the movement which has been shown to be unaffected by feedback responses (Todorov and Jordan 2002, Pruszynski, Kurtzer et al. 2008, Pruszynski and Scott 2012, Crevecoeur and Scott 2013). However, in our task, the amplitude of movements made by our monkeys was small, and therefore the response measures we used were too small in the first 50-100 ms for a robust estimation. Also, fixing a time window led to an unfair comparison between control and cerebellar block trials, in which velocity was significantly reduced and therefore movement time was longer. Therefore, we used the peak velocity, torque-impulse at

the peak velocity, and maximum deviation of the hand trajectory as response
measures.

Statistical analysis

We used linear mixed effects analysis to evaluate the effects of predictor variables
of interest on the computed response variables. This allowed us to account for the
variability across the monkeys which was modeled as a random effect. Each data
sample for this model was obtained by calculating the median of all the trials for each
unique combination of the response variable(s) per session. We computed the change
in the measured outcome variable due to the cerebellar block per session using the
following formula:

620
$$\frac{Median\ response\ during\ cerebellar\ block - median\ response\ during\ control}{Median\ response\ during\ control}$$

$\times\ 100$

For post-hoc comparisons, we used the 'coefTest' function from MATLAB's 'fitlme'
to perform hypothesis testing on the fixed effects. This involved specifying the
contrast matrices to evaluate specific hypotheses regarding the differences in the
response variables across the levels of the predictor variables. The 'coefTest' function
allowed us to obtain p-values for these comparisons. The computed p-values were
then corrected using the Benjamini-Hochberg procedure to control the false
discovery rate for multiple post-hoc comparisons, wherever applicable (Haynes
2013). A significance level of 0.05 was used for all comparisons in this study. For
estimation of the group means and their confidence intervals, we pooled the per
session data from all monkeys after ensuring that effects are consistent across the
monkeys. For further clarity, we have also overlayed the means of the measured
variables from individual monkeys on all our plots of summary statistics.

Acknowledgements

This work was funded by the Israel Science Foundation (ISF-1801/18 and ISF
1207/23) to Y.P., the Deutsche Forschungsgemeinschaft (DFG, German Research
Foundation Project-ID 431549029–SFB 1451) to Y.P., National Institute of
Neurological Disorders and Stroke of the National Institutes of Health (NIH
R01NS110901) to. Y.P., the National Institutes of Health (NIH R01NS105759) to
J.P.A.D., the Council for Higher Education PhD Sandwich Fellowship, Government of
Israel to N.S. for undertaking his dual-PhD at the Hebrew University and
Northwestern University, and the generous support of the Baruch Foundation to Y.P.
The authors would like to thank Dr. Abdulraheem Nashef, a postdoctoral fellow at the
University of Colorado for collecting part of the data used in this study.

Data availability statement

Data and code used in this manuscript are available publicly as a GitHub repository (Sinha, Israely et al. 2024).

Competing interests

The authors declare no financial or non-financial competing interests.

References

Asanuma, C., W. T. Thach and E. G. Jones (1980). "Nucleus interpositus projection to spinal interneurons in monkeys." Brain Research **191**(1): 245-248.

Bantli, H. and J. R. Bloedel (1975). "The action of the dentate nucleus on the excitability of spinal motoneurons via pathways which do not involve the primary sensorimotor cortex." Brain Research **88**(1): 86-90.

Bastian, A. J. (1997). "Mechanisms of Ataxia." Physical Therapy **77**(6): 672-675.

Bastian, A. J., T. A. Martin, J. G. Keating and W. T. Thach (1996). "Cerebellar ataxia: abnormal control of interaction torques across multiple joints." Journal of Neurophysiology **76**(1): 492-509.

Bastian, A. J. and W. T. Thach (1995). "Cerebellar outflow lesions: A comparison of movement deficits resulting from lesions at the levels of the cerebellum and thalamus." Annals of Neurology **38**(6): 881-892.

Bastian, A. J., K. M. Zackowski and W. T. Thach (2000). "Cerebellar Ataxia: Torque Deficiency or Torque Mismatch Between Joints?" Journal of Neurophysiology **83**(5): 3019-3030.

Becker, M. I. and A. L. Person (2019). "Cerebellar Control of Reach Kinematics for Endpoint Precision." Neuron **103**(2): 335-348 e335.

Beer, R. F., J. P. A. Dewald and W. Z. Rymer (2000). "Deficits in the coordination of multijoint arm movements in patients with hemiparesis: evidence for disturbed control of limb dynamics." Experimental Brain Research **131**(3): 305-319.

Beppu, H., M. Suda and R. Tanaka (1984). "ANALYSIS OF CEREBELLAR MOTOR DISORDERS BY VISUALLY GUIDED ELBOW TRACKING MOVEMENT." Brain **107**(3): 787-809.

Bernshteĭn, N. A. (1967). The co-ordination and regulation of movements. Oxford, Pergamon Press Oxford.

Boose, A., J. Dichgans and H. Topka (1999). "Deficits in phasic muscle force generation explain insufficient compensation for interaction torque in cerebellar patients." Neuroscience Letters **261**(1): 53-56.

Bremer, F., G.-H. Roger and L. Binet (1935). "Le cervelet." Traité de physiologie: 39-134.

Calame, D. J., M. I. Becker and A. L. Person (2023). "Cerebellar associative learning underlies skilled reach adaptation." Nat Neurosci **26**(6): 1068-1079.

Chambers, W. W., Jr. (1947). "Electrical stimulation of the interior of the cerebellum in the cat." Am. J. Anat. **80**(1): 55-93.

Cheng, E. J. and S. H. Scott (2000). "Morphometry of Macaca mulatta forelimb. I. Shoulder and elbow muscles and segment inertial parameters." Journal of Morphology **245**(3): 206-224.

Crevecoeur, F. and S. H. Scott (2013). "Priors Engaged in Long-Latency Responses to Mechanical Perturbations Suggest a Rapid Update in State Estimation." PLOS Computational Biology **9**(8): e1003177.

Day, B. L., P. D. Thompson, A. E. Harding and C. D. Marsden (1998). "Influence of vision on upper limb reaching movements in patients with cerebellar ataxia." Brain **121**(2): 357-372.

Fagg, A., G. Ojakangas, L. Miller and N. Hatsopoulos (2009). "Kinetic Trajectory Decoding Using Motor Cortical Ensembles." IEEE transactions on neural systems and rehabilitation engineering : a publication of the IEEE Engineering in Medicine and Biology Society **17**: 487-496.

Freund, H.-J. (1985). "Clinical aspects of premotor function." Behavioural Brain Research **18**(2): 187-191.

Freund, H. J. and H. Hummelsheim (1985). "Lesions of premotor cortex in man." Brain : a journal of neurology **108 (Pt 3)**: 697-733.

Fulton, J. F. and R. S. Dow (1937). "The Cerebellum: A Summary of Functional Localization *." The Yale Journal of Biology and Medicine **10**: 89 - 119.

Goldstein, K. (1927). "Das Kleinhirn." Handbuch der normalen und pathologischen Physiologie **10**: 222-222.

Granit, R., B. Holmgren and P. A. Merton (1955). "The two routes for excitation of muscle and their subservience to the cerebellum." The Journal of Physiology **130**(1): 213-224.

Hage, P., M. A. Fakharian, A. M. Shoup, J. S. Pi, E. Sedaghat-Nejad, S. P. Orozco, I. K. Jang, V. Looi, H. Y. Elseweifi, N. Mohammadrezaei, A. N. Vasserman, T. Arginteanu and R. Shadmehr (2025). "Control of tongue movements by the Purkinje cells of the cerebellum." bioRxiv: 2024.2007.2025.604757.

Haines, D. E., M.-U. Manto and M. Glickstein (2007). "Clinical symptoms of cerebellar disease and their interpretation." The Cerebellum **6**(2): 141-156.

Hallett, M., A. Berardelli, J. Matheson, J. Rothwell and C. D. Marsden (1991). "Physiological analysis of simple rapid movements in patients with cerebellar deficits." J Neurol Neurosurg Psychiatry **54**(2): 124-133.

Hare, W. K., H. W. Magoun and S. W. Ranson (1936). "ELECTRICAL STIMULATION OF THE INTERIOR OF THE CEREBELLUM IN THE DECEREBRATE CAT." American Journal of Physiology-Legacy Content **117**(2): 261-266.

Haynes, W. (2013). Benjamini–Hochberg Method. Encyclopedia of Systems Biology. W. Dubitzky, O. Wolkenhauer, K.-H. Cho and H. Yokota. New York, NY, Springer New York**:** 78-78.

Hollerbach, J. M. and T. Flash (1982). "Dynamic interactions between limb segments during planar arm movement." Biological Cybernetics **44**(1): 67-77.

Holmes, G. (1917). "The symptoms of acute cerebellar injuries due to gunshot
injuries." <u>Brain: A Journal of Neurology</u> **40**: 461-535.
Holmes, G. (1939). "THE CEREBELLUM OF MAN1." <u>Brain</u> **62**(1): 1-30.
Hore, J. and D. Flament (1988). "Changes in motor cortex neural discharge
associated with the development of cerebellar limb ataxia." <u>Journal of
Neurophysiology</u> **60**(4): 1285-1302.
Horne, M. K. and E. G. Butler (1995). "The role of the cerebello-thalamo-cortical
pathway in skilled movement." <u>Progress in Neurobiology</u> **46**(2): 199-213.
Ivry, R. B. and S. W. Keele (1989). "Timing Functions of The Cerebellum." <u>Journal of
Cognitive Neuroscience</u> **1**(2): 136-152.
Jeannerod, M. (1988). <u>The neural and behavioural organization of goal-directed
movements</u>. New York, NY, US, Clarendon Press/Oxford University Press.
Kawato, M. (1999). "Internal models for motor control and trajectory planning."
<u>Current Opinion in Neurobiology</u> **9**(6): 718-727.
Konczak, J., D. Pierscianek, S. Hirsiger, U. Bultmann, B. Schoch, E. R. Gizewski, D.
Timmann, M. Maschke and M. Frings (2010). "Recovery of Upper Limb Function
After Cerebellar Stroke." <u>Stroke</u> **41**(10): 2191-2200.
Konczak, J., B. Schoch, A. Dimitrova, E. Gizewski and D. Timmann (2005). "Functional
recovery of children and adolescents after cerebellar tumour resection." <u>Brain</u>
128(6): 1428-1441.
Low, A. Y. T., A. R. Thanawalla, A. K. K. Yip, J. Kim, K. L. L. Wong, M. Tantra, G. J.
Augustine and A. I. Chen (2018). "Precision of Discrete and Rhythmic Forelimb
Movements Requires a Distinct Neuronal Subpopulation in the Interposed Anterior
Nucleus." <u>Cell Reports</u> **22**(9): 2322-2333.
Luciani, L. (1893). "Il Cervelletto. Nuovi Studi di Fisiologia Normale e Patologica."
<u>Philosophical Review</u> **2**(4).
Magoun, H. W., W. K. Hare and S. W. Ranson (1935). "ELECTRICAL STIMULATION OF
THE INTERIOR OF THE CEREBELLUM IN THE MONKEY." <u>American Journal of
Physiology-Legacy Content</u> **112**(2): 329-339.
Manto, M., J. Bower, A. Conforto, J. Delgado-García, S. Guarda, M. Gerwig, C. Habas, N.
Hagura, R. Ivry, P. Mariën, M. Molinari, E. Naito, D. Nowak, N. Taib, D. Pélisson, C.
Tesche, C. Tilikete and D. Timmann (2011). "Consensus Paper: Roles of the
Cerebellum in Motor Control-The Diversity of Ideas on Cerebellar Involvement in
Movement." <u>Cerebellum (London, England)</u> **11**: 457-487.
Meyer-Lohmann, J., B. Conrad, K. Matsunami and V. B. Brooks (1975). "Effects of
dentate cooling on precentral unit activity following torque pulse injections into
elbow movements." <u>Brain Research</u> **94**(2): 237-251.
Miall, R. C., L. O. D. Christensen, O. Cain and J. Stanley (2007). "Disruption of State
Estimation in the Human Lateral Cerebellum." <u>PLOS Biology</u> **5**(11): e316.
Miall, R. C. and D. King (2008). "State Estimation in the Cerebellum." <u>The Cerebellum</u>
7(4): 572-576.
Naidich, T. P., E. G. Stein, P. A. Lento, G. M. Kleinman, M. E. Fowkes and D. M.
Carpenter (2013). CHAPTER 16 - Cerebellum. <u>Imaging of the Brain</u>. T. P. Naidich, M.
Castillo, S. Cha and J. G. Smirniotopoulos. Philadelphia, W.B. Saunders**:** 328-347.

Nashef, A., O. Cohen, R. Harel, Z. Israel and Y. Prut (2019). "Reversible Block of
Cerebellar Outflow Reveals Cortical Circuitry for Motor Coordination." <u>Cell Reports</u>
27(9): 2608-2619.e2604.
Nashef, A., O. Cohen, Z. Israel, R. Harel and Y. Prut (2018). "Cerebellar Shaping of
Motor Cortical Firing Is Correlated with Timing of Motor Actions." <u>Cell Reports</u>
23(5): 1275-1285.
Nashef, A., H. Rapp, M. P. Nawrot and Y. Prut (2018). "Area-specific processing of
cerebellar-thalamo-cortical information in primates." <u>Biological Cybernetics</u> **112**(1):
141-152.
Nashold, B. S. and D. G. Slaughter (1969). "Effects of Stimulating or Destroying the
Deep Cerebellar Regions in Man." <u>Journal of Neurosurgery</u> **31**(2): 172-186.
Nathan, P. W. and M. C. Smith (1982). "The rubrospinal and central tegmental tracts
in man." <u>Brain : a journal of neurology</u> **105**(Pt 2): 223-269.
Osu, R., K.-i. Morishige, J. Nakanishi, H. Miyamoto and M. Kawato (2015). "Practice
reduces task relevant variance modulation and forms nominal trajectory." <u>Scientific</u>
<u>Reports</u> **5**(1): 17659.
Padel, Y., P. Angaut, J. Massion and R. Sedan (1981). "Comparative study of the
posterior red nucleus in baboons and gibbons." <u>Journal of Comparative Neurology</u>
202(3): 421-438.
Perlmutter, J. S. and J. W. Mink (2006). "DEEP BRAIN STIMULATION." <u>Annual</u>
<u>Review of Neuroscience</u> **29**(Volume 29, 2006): 229-257.
Perlmutter, S. I., M. A. Maier and E. E. Fetz (1998). "Activity of Spinal Interneurons
and Their Effects on Forearm Muscles During Voluntary Wrist Movements in the
Monkey." <u>Journal of Neurophysiology</u> **80**(5): 2475-2494.
Pi, J. S., M. A. Fakharian, P. Hage, E. Sedaghat-Nejad, S. Z. Muller and R. Shadmehr
(2024). "The olivary input to the cerebellum dissociates sensory events from
movement plans." <u>Proc Natl Acad Sci U S A</u> **121**(17): e2318849121.
Plamondon, R. and A. M. Alimi (1997). "Speed/accuracy trade-offs in target-directed
movements." <u>Behavioral and Brain Sciences</u> **20**(2): 279-303.
Popa, L., A. Hewitt and T. Ebner (2013). "Purkinje Cell Simple Spike Discharge
Encodes Error Signals Consistent with a Forward Internal Model." <u>Cerebellum</u>
<u>(London, England)</u> **12**.
Pressman, P. and H. J. Rosen (2015). Chapter 33 - Disorders of Frontal Lobe
Function. <u>Neurobiology of Brain Disorders</u>. M. J. Zigmond, L. P. Rowland and J. T.
Coyle. San Diego, Academic Press**:** 542-557.
Pruszynski, J. A., I. Kurtzer and S. H. Scott (2008). "Rapid Motor Responses Are
Appropriately Tuned to the Metrics of a Visuospatial Task." <u>Journal of</u>
<u>Neurophysiology</u> **100**(1): 224-238.
Pruszynski, J. A. and S. H. Scott (2012). "Optimal feedback control and the long-
latency stretch response." <u>Exp Brain Res</u> **218**(3): 341-359.
Ruigrok, T. J. H. (2013). Cerebellar Influences on Descending Spinal Motor Systems.
<u>Handbook of the Cerebellum and Cerebellar Disorders</u>. M. Manto, J. D. Schmahmann,
F. Rossi, D. L. Gruol and N. Koibuchi. Dordrecht, Springer Netherlands**:** 497-528.

Sainburg, R. L., C. Ghez and D. Kalakanis (1999). "Intersegmental Dynamics Are
Controlled by Sequential Anticipatory, Error Correction, and Postural Mechanisms."
Journal of Neurophysiology **81**(3): 1045-1056.
Sathyamurthy, A., A. Barik, C. I. Dobrott, K. J. E. Matson, S. Stoica, R. Pursley, A. T.
Chesler and A. J. Levine (2020). "Cerebellospinal Neurons Regulate Motor
Performance and Motor Learning." Cell Reports **31**(6): 107595.
Schlerf, J. E., J. Xu, N. M. Klemfuss, T. L. Griffiths and R. B. Ivry (2012). "Individuals
with cerebellar degeneration show similar adaptation deficits with large and small
visuomotor errors." Journal of Neurophysiology **109**(4): 1164-1173.
Schneider, R. C. and E. C. Crosby (1963). "The Interplay Between Cerebral
Hemispheres and Cerebellum in Relation to Tonus and Movements." Journal of
Neurosurgery **20**(3): 188-198.
Scott, S. H. (2004). "Optimal feedback control and the neural basis of volitional
motor control." Nature Reviews Neuroscience **5**(7): 532-545.
Shadmehr, R. and F. Mussa-Ivaldi (1994). "Adaptive representation of dynamics
during learning of a motor task." The Journal of Neuroscience **14**(5, Pt 2): 3208-
3224.
Shidara, M., K. Kawano, H. Gomi and M. Kawato (1993). "Inverse-dynamics model
eye movement control by Purkinje cells in the cerebellum." Nature **365**(6441): 50-
52.
Shinoda, Y., M. Yamazaki and T. Futami (1982). "Convergent inputs from the dentate
and the interpositus nuclei to pyramidal tract neurons in the motor cortex."
Neuroscience Letters **34**(2): 111-115.
Sinha, N., S. Israely, O. Ben Harosh, R. Harel, J. P. A. Dewald and Y. Prut (2024).
Disentangling-acute-motor-deficits-and-adaptive-responses-evoked-by-the-loss-of-
cerebellar-output.
Smith, J. L. and R. F. Zernicke (1987). "Predictions for neural control based on limb
dynamics." Trends in Neurosciences **10**(3): 123-128.
ten Donkelaar, H. J. (1988). "Evolution of the red nucleus and rubrospinal tract."
Behavioural Brain Research **28**(1): 9-20.
Therrien, A. S. and A. J. Bastian (2015). "Cerebellar damage impairs internal
predictions for sensory and motor function." Current Opinion in Neurobiology **33**:
127-133.
Timmann, D., S. Watts and J. Hore (1999). "Failure of Cerebellar Patients to Time
Finger Opening Precisely Causes Ball High-Low Inaccuracy in Overarm Throws."
Journal of Neurophysiology **82**(1): 103-114.
Todorov, E. and M. I. Jordan (2002). "Optimal feedback control as a theory of motor
coordination." Nature Neuroscience **5**(11): 1226-1235.
Topka, H., J. Konczak and J. Dichgans (1998). "Coordination of multi-joint arm
movements in cerebellar ataxia:Analysis of hand and angular kinematics."
Experimental Brain Research **119**(4): 483-492.
Topka, H., J. Konczak, K. Schneider, A. Boose and J. Dichgans (1998). "Multijoint arm
movements in cerebellar ataxia: Abnormal control of movement dynamics."
Experimental Brain Research **119**: 493-503.

Tsujimoto, T., H. Gemba and K. Sasaki (1993). "Effect of cooling the dentate nucleus
of the cerebellum on hand movement of the monkey." Brain Research **629**(1): 1-9.
Virji-Babul, N. and J. D. Cooke (1995). "Influence of joint interactional effects on the
coordination of planar two-joint arm movements." Experimental Brain Research
103(3): 451-459.
Weeks, H. M., A. S. Therrien and A. J. Bastian (2017). "Proprioceptive Localization
Deficits in People With Cerebellar Damage." The Cerebellum **16**(2): 427-437.
Wild, B. and J. Dichgans (1993). "Cerebellar ataxia in ataxic hemiparesis? A
kinematic and EMG analysis." Movement Disorders **8**(3): 363-366.
Wild, B., T. Klockgether and J. Dichgans (1996). "Acceleration deficit in patients with
cerebellar lesions. A study of kinematic and EMG-parameters in fast wrist
movements." Brain Research **713**(1): 186-191.
Wolpert, D. M., Z. Ghahramani and M. I. Jordan (1995). "An Internal Model for
Sensorimotor Integration." Science **269**(5232): 1880-1882.
Wolpert, D. M., R. C. Miall and M. Kawato (1998). "Internal models in the
cerebellum." Trends in Cognitive Sciences **2**(9): 338-347.

Figures and Tables

Figure 1: Effect of cerebellar block on peak hand velocity. (a) Schematic of the monkey performing
the delayed reaching task in the KINARM exoskeleton to eight peripheral targets on the horizontal
plane. The locations of the eight targets are numbered. (b) Movement-onset aligned hand velocity
profiles during a sample control and cerebellar block trial performed by monkey C to target 2. Entry of
the control cursor into the peripheral target marked the end of the movement for the trial. Since the
monkeys did not have to stop their movements fully for the trial to end (see Methods for a detailed
explanation of this), the traces appear cutoff at the beginning of the deceleration/stopping phase of
the movement. (c) Target-wise effect of the cerebellar block on peak hand velocity. For each session,
the target-wise reduction in the median peak hand velocity during the cerebellar block trials was
computed relative to that of control trials. The depicted values are the means ± 95% confidence
intervals across all sessions pooled from all four monkeys. The means of individual monkeys are
overlaid. The dashed circle indicates no change. Statistical significance is denoted as follows: $p \geq$
0.05NS, $p < 0.05$*, $p < 0.01$**, $p < 0.001$***. [T1-8: Targets 1-8]

Figure 2: Effect of cerebellar block on muscle torques. Movement-onset aligned shoulder (a) and elbow (b) muscle torque profiles during a sample control and cerebellar block trial performed by monkey C to target 2. Entry of the control cursor into the peripheral target marked the end of the movement for the trial. Since the monkeys did not have to stop their movements fully for the trial to end (see Methods for a detailed explanation of this), the traces appear cutoff at the beginning of the deceleration/stopping phase of the movement. (c) Normalized muscle torque impulse at the shoulder vs. elbow per target during control and cerebellar block. For each session, we computed the target-wise median muscle torque impulse during the acceleration phase of the movement across the control and cerebellar block trials. They were then normalized by the maximum absolute torque impulse across all the targets. The depicted values are the means across all the sessions pooled from the data of all four monkeys. The 95% confidence interval of the means is indicated by the horizontal and vertical bars for the shoulder and elbow joints respectively. Target-wise effect of cerebellar block on shoulder (d) and elbow (e) muscle torque impulse. For each session, the target-wise reduction in the median torque impulse during the cerebellar block trials was computed relative to the maximum

904 absolute value of the target-wise medians observed during the control trials. The depicted values are
905 the means ± 95% confidence intervals across all sessions pooled from all four monkeys. The means of
906 individual monkeys are overlaid. The dashed circle indicates no change. Statistical significance is
907 denoted as follows: p ≥ 0.05NS, p < 0.05*, p < 0.01**, p < 0.001***. [T1-8: Targets 1-8]

Figure 3: Effect of cerebellar block on coupling torques. Movement-onset aligned shoulder (a) and
elbow (b) coupling torque profiles during a sample control and cerebellar block trial performed by
monkey C to target 2. Entry of the control cursor into the peripheral target marked the end of the trial.
(c) Normalized coupling torque impulse at the shoulder vs. elbow per target during control and
cerebellar block. For each session, we computed the target-wise median coupling torque impulse
during the acceleration phase of the movement across the control and cerebellar block trials. They
were then normalized by the maximum absolute torque impulse across all the targets. The depicted
values are the means across all the sessions pooled from the data of all four monkeys. The 95%
confidence interval of the means is indicated by the horizontal and vertical bars for the shoulder and
elbow joints respectively.(d) The effect of the cerebellar block on peak hand velocity (y-axis) is plotted
against the total coupling torque experienced across both joints during control (x-axis) for the outward
reaches (T1-4). The effect of the cerebellar block on peak hand velocity was measured by the target-
wise change in the median peak hand velocity during the cerebellar block trials relative to that of
control trials per session (negative percentage values indicate a stronger reduction due to the
cerebellar block). This measure was then correlated to the target-wise median coupling torque during
the control trials of the corresponding sessions. The median coupling torque was taken as the sum of
the absolute coupling torque impulse at both the shoulder and elbow joints per trial. These values were
then normalized by the median across all trials of all sessions per monkey (to yield percentage units
like the y-axis). The depicted data contains all the sessions pooled from all four monkeys with
individual points representing each of the four targets (color-coded) per session. The black line
overlayed on the scatter indicates the least-squares linear regression fit to the data points. [T1-8:
Targets 1-8]

4a: Process of extraction of peak velocities in trial sequence

4b: Hand velocity vs. trial no., single-joint movements (T1)

4c: Hand velocity vs. trial no., dual-joint movements (T2-4)

4d: *Post hoc* comparisons (T2-4)

Figure 4: Effect of the cerebellar block on hand velocity across successive trials. (a) Schematic of
the process of extraction of peak hand velocity for movements to a particular target from blocks of
control/cerebellar block trials while preserving the order in which the target was presented. For each
block, the peak velocity for movements to a particular target (in this example target 1) was extracted
while retaining the trial numbers in which it was presented in the block. This enabled us to examine
the evolution of the peak velocity across the sequence of presentation of the target during control vs.
cerebellar block. (b) Mean ± 95% confidence intervals of the peak hand velocity for movements to
target 1 across all the trial blocks pooled from all four monkeys. For each monkey, the peak hand
velocities were normalized to the median peak velocity of the early trials 1-2 from all the control
blocks. (c) Same as in (b) but for targets 2-4. (d) Post hoc comparisons between early trials 1-2 and
late trials 11-20 during control and cerebellar block. Statistical significance is denoted as follows: p ≥
0.05NS, p < 0.05*, p < 0.01**, p < 0.001***. [E: Early, L: Late, C: Control, Cbl: Cerebellar block; T1-8:
Targets 1-8]

Figure 5a-c: Effect of the cerebellar block on the decomposition of movements. (a) Binary decomposition raster of sample trials for movements made to target 2 by monkey C during control. The duration of each trial was normalized to 0-1 (bin width = 0.001) and then decomposition was computed for each bin based on whether either (but not both) of the shoulder or the elbow joint velocity was less than 20°/s. (b) Same as in (a) but for movements to target 2 by monkey C during cerebellar block. (c) Change in decomposition index (i.e., the proportion of the movement time during which the movement was decomposed, as defined above) for movements to targets 2-4 during velocity-matched control and cerebellar block trials relative to all control trials. The change in median decomposition was computed for each session. The depicted values are the mean ± 95% confidence intervals across all sessions pooled from all four monkeys. The individual means of each monkey are overlaid. **Figure 5d-e: Effect of the cerebellar block on inter-trial trajectory variability.** (d) Sample hand trajectories during control vs. cerebellar block trials for movements to target 2 by monkey C. The starting point of the trajectories was shifted to the origin and then they were rotated about the origin so that their endpoints lie on the positive Y-axis. (e) Change in inter-trial trajectory variability for movements to targets 2-4 during velocity-matched control and cerebellar block trials

relative to all control trials. The trajectory variability was measured as the standard deviation of the
maximum perpendicular distance of the trajectories from the Y-axis after transforming them as in (d).
The change in trajectory variability for the cerebellar block trials was computed relative to the control
trials for each session. The depicted values are the mean ± 95% confidence intervals across all sessions
pooled from all four monkeys. The individual means of each monkey are overlaid. Statistical
significance is denoted as follows: $p \geq 0.05$NS, $p < 0.05$*, $p < 0.01$**, $p < 0.001$***. [C: Control, Cbl:
Cerebellar block; T1-8: Targets 1-8; DI: Decomposition index