TITLE: Occupancy of the HbYX hydrophobic pocket is sufficient to

induce gate opening in the archaeal 20S proteasomes.

SHORT TITLE: Occupying hydrophobic pocket activates proteasome

Janelle J.Y. Chuah[1], Madalena R. Daugherty[1], David M. Smith[1,2*]

1: Department of Biochemistry and Molecular Medicine, West Virginia University School of

Medicine, 64 Medical Center Dr., Morgantown, WV USA

2: Department of Neuroscience, Rockefeller Neuroscience Institute, West Virginia University,

Morgantown, WV USA

Address correspondence to: David M. Smith, E-mail: dmsmith@hsc.wvu.edu.

ABSTRACT

Enhancing proteasome function has been a long-standing but challenging target of interest for the potential treatment of neurodegenerative diseases, emphasizing the importance of understanding proteasome activation mechanisms. Most proteasome activator complexes use the C-terminal HbYX (hydrophobic-tyrosine-almost any residue) motif to bind and trigger gate-opening in the 20S proteasome. This study defines a critical molecular interaction in the HbYX mechanism that triggers gate opening. We focus on the Hb site interaction and find it plays a surprisingly central and crucial role in driving the allosteric conformational changes that induce gate opening in the archaeal 20S. We examined the cryo-EM structure of two mutant archaeal proteasomes, αV24Y T20S and αV24F T20S. These two mutants were engineered to place a bulky aromatic residue in the HbYX hydrophobic pocket; both mutants are highly active, though their mechanisms of activation are undefined. Collectively, our findings indicate that the interaction between the Hb group of the HbYX motif and its corresponding hydrophobic pocket is sufficient to induce gate opening in a mechanistically similar way to the HbYX motif. The activation mechanism studied here involves the expansion of the hydrophobic binding site, allosterically altering the state of the IT switch thus triggering gate-opening. Furthermore, we show that the canonical αK66 residue, previously understood to be critical for proteasome activator binding, also plays a key role in stabilizing the open gate, irrespective of activator binding. This study differentiates between the residues in the HbYX motif that support binding interactions ("YX") versus those that allosterically contribute to gate opening ("Hb"). The insights reported here will guide future drug development efforts, particularly in designing small molecule proteasome activators, by targeting the identified hydrophobic pocket.

INTRODUCTION

The proteasome is a molecular machine responsible for degrading misfolded, damaged, and unneeded proteins that are targeted via multiple mechanisms including the conventional pathway of ubiquitination(*1–6*). Proteasomal degradation regulates major cellular processes and functions, such as synaptic plasticity and memory formation(*7–11*). Impairment of proteasome function has correlated with areas of the brain affected by neurodegenerative diseases, supporting its implication in these diseases(*12–17*). Furthermore, increasing proteasome activity has been shown to induce protective or rescue effects in models of neurodegenerative diseases(*18–23*), strengthening the rationale to develop small molecule proteasome activators. The objective of this study is to provide a clear mechanistic framework of how proteasome activation occurs to inform drug discovery efforts.

The eukaryotic proteasome, also referred to as the 20S or the Core Particle (CP), consists of four stacked heteroheptameric (homoheptameric in archaea) rings arranged as α − β − β − α(*24*). Three of the seven β subunits carry protease sites with distinct protease activities: caspase-like (β1), trypsin-like (β2), and chymotrypsin-like (β5). The α subunits have an N-termini that extends toward the central pore, forming a gate that regulates substrate entry. The gate is primarily formed by the N-termini of α2, 3, and 4 with α3's N-termini (the longest) acting as the lynchpin(*24*). The interactions that stabilize the open conformation have been reported in crystal and cryo-EM structures(*24–31*). A YDR motif, found on the α N-termini, interacts with the neighboring α subunit's N-terminal tail, to stabilize the gate in the open state. Most recently, we discovered that the open and closed states of the α N-termini conformation are stabilized by two alternating conformations of the IT switch(*31*). Two α subunit residues, I12 and T13, in the archaeal *Thermoplasma acidophilum* proteasome (T20S), switch positions to occupy the pocket adjacent to Helix 0 and stabilize the open and closed gate conformation respectively. This IT switch is functionally conserved in the eukaryotic 20S(*31*). Basal kinetics of the archaeal proteasome gate,

quantified using NMR(*32*), indicates that the gate fluctuates between open and close on a time

scale of seconds; however, the gate primarily exists in the closed state of isolated 20S.

The 20S proteasome is known to associate with regulatory complexes, collectively referred to as

Proteasome Activators (PAs)(*33*). PA700 or 19S, which associates with the 20S to form the 26S,

has been identified as a critical component of the Ubiquitin-Proteasome Pathway. PA700

recognizes ubiquitinated substrates and unfolds them to be degraded by the 20S. Like many PAs,

PA700 induces 20S gate-opening via the HbYX-dependent mechanism, as does the archaeal

homolog of PA700 called PAN(*33, 34*). Three of the six C-terminal tails of PA700's ATPases carry

a HbYX motif (hydrophobic-tyrosine-almost any residue), whereby the "X" residue must be the C-

terminal residue. These HbYX motifs interact with the inter-subunit pockets in the 20S α ring and

allosterically induce gate-opening. The precise molecular mechanism of how the motif's binding

in inter-subunit pockets induces gate opening is not known, though high-resolution structures of

the 26S(*26–29, 35*) and the 20S interacting with a di-peptide HbYX-mimetic(*31*) have generated

plausible models. Structures of the 26S reveal that the C-terminal tails bind in a particular order

during activation, though the correlation between the number of C-terminal tails/HbYX-bound to

induce gate opening is controversial. A study indicated that partial gate opening occurs when Rpt2,

3, and 5, which have HbYX motifs, dock in their corresponding α inter-subunit pockets and full

gate opening occurs when Rpt6, which does not have a HbYX motif, additionally binds(*26*).

Another study indicated that the binding of Rpt2, 3, and 5 did not open the gate, but the additional

binding of Rpt6 causes full gate opening(*28*). Interestingly, multiple studies suggest that Rpt1,

which carries the penultimate tyrosine, but not the Hb group (partial-HbYX), must also bind in

combination with Rpt2, 3, 5, and 6 for gate opening(*27, 29, 35*). Collectively, these structures

suggest that each inter-subunit pocket contributes to gate opening to varying extents, which

correlates with the fact that each α subunit's N-terminal tails differentially participate in the closed

state, though the minimum occupancy of inter-subunit pockets required for gate opening is

unresolved.

While some PAs are HbYX-dependent like PA700, the 11S family, such as PA26 or PA28 activate

the proteasome via a HbYX-independent mechanism. Despite the divergence between

mechanisms, they converge on the critical molecular interactions and conformational changes

(e.g. the IT switch) that stabilize the open gate state(*31*). For example, both HbYX-dependent and

HbYX-independent PAs have been shown to interact with αK66 side chain and mutating αK66 to

alanine impedes PAs association and activation, validated by velocity sedimentation assay and

proteasomal peptidase activity(*30*, *34*). Both mechanisms also involve a displacement of

αPro17(*30*, *31*) and switching of the IT switch(*31*). Therefore, though the cause of the

conformational changes mentioned may differ between HbYX-dependent and HbYX-independent

mechanisms, they share in some of the PA-20S interactions that support the open conformational

state of the gate.

Our early studies showed that the HbYX motif could function even on short 6-7 residue

peptides(*34*); recently we showed that HbYX-dependent gate opening could be reduced to a

carboxybenzyl (CBZ)-blocked YA dipeptide called ZYA—a minimal HbYX mimetic. Our ZYA-T20S

structure(*31*) indicated the minimum molecular interactions necessary for optimal HbYX-

dependent gate-opening by a small molecule. In our 1.9Å resolution structure, ZYA was observed

to interact with residues in the α subunit (G19, K66, L81, etc.) in a similar orientation and fashion

as the native HbYX motif of various PA complexes. Our cryo-EM data indicated that αV24 (and

αA154) hydrophobically interacts with the carboxybenzyl of ZYA and the hydrophobic side chain

of the PAs HbYX motif. Interestingly, our previous study also indicated that a single residue

mutation on αV24 (i.e., αV24Y, αV24F) increased proteasome activity, though the mechanism of

activation by this mutation was not determined. To assess the significance of these mutations in

activating the proteasome via the HbYX-dependent pathway, we determined the cryo-EM

structure of αV24Y T20S and αV24F T20S. The results of this study indicate that occupancy of a

hydrophobic pocket by a bulky hydrophobic group such as αV24Y within the α inter-subunit pocket

is sufficient to trigger 20S gate opening via the HbYX-dependent mechanism, demonstrating the

surprising importance of the HbYX motif's single Hb group for allosteric activation of the 20S. The

findings reported here are expected to translate to the human proteasome as ZYA is a robust

activator of the mammalian 20S and the HbYX-dependent mechanism is conserved; thereby,

uncovering a highly specific pharmaceutical target for proteasome activation.

RESULTS

Cryo-EM map of αV24Y T20S shows HbYX-dependent-like conformational changes and an open
gate state.

To elucidate how the mutation, αV24Y, induce gate-opening in the archaeal proteasome (αV24Y

T20S), we resolved its structure using cryo-EM (Fig. 1; Fig. S1 & S2). The *Thermoplasma*

acidophilum 20S (T20S) has D7 symmetry, which was applied during reconstruction, and the final

map resolved to 2.38Å (EMD-44914). Comparing the αV24Y T20S electron density map against

our wild-type T20S (WT T20S) map (EMD-28878) revealed conformational changes similar to

those previously noted in our ZYA-T20S model(*31*) (PDB:8F7K) (Fig. 1). Alignment of the maps

indicate the αV24Y mutation induced minimal conformational changes in the β rings, while the α

subunits exhibited rotational changes that closely mimicked HbYX-dependent conformational

changes, such as those induced by ZYA binding(*31, 36*) (Fig. 1A & B). Comparison of the maps

show the presence of the expected larger density of the mutant tyrosine side chain in the

hydrophobic pocket of the αV24Y T20S map with proximity to the docking location of the

carboxybenzyl group (Z) of ZYA (Fig. 1C, D, & E).

Our prior biochemical analysis showed that the αV24Y mutation caused substantial acceleration

of peptide degradation, suggesting that this mutation might stimulate gate-opening(*31*); therefore,

we sought to determine the conformational changes in the structure of αV24Y T20S. We observed

significant densities corresponding to the YDR motif in the open state, designating an open gate

conformation (Fig. 1F, G, & H). Additionally, the densities corresponding to the "IT switch" were in

similar confirmation to previously known open gate proteasome cryo-EM structures: ZYA-T20S

(Fig. 1I, J, & K) and L81Y-T20S(31). It is also apparent in the map of αV24Y T20S that there is a

lack of densities in the central cavity of the α ring that corresponding to the closed state of the α

N-termini, definitively confirming that V24Y T20S's gate is open (Fig. 1A & B). Collectively, the

comparison of maps demonstrates that V24Y T20S is in an open state with global conformational

changes in the α-subunits that mirror those of a HbYX-bound proteasome (Fig. 1).

αV24Y T20S shows HbYX dependent conformational changes that are greater in magnitude than

those seen in the ZYA-T20S structure.

To gain a more detailed understanding of the conformational changes induced by αV24Y, we

generated an atomic model using the αV24Y T20S map. To evaluate the global conformational

changes in the α subunits, we aligned the αV24Y T20S and the ZYA-T20S to the WT T20S model

via their β subunits and overlayed all three models (Fig. 2). The α subunits in αV24Y T20S rotate

roughly as a rigid body on a pivot around Helix 2 (Fig. 2A, B, C, & 3A), using WT T20S as a

reference. This rotation around Helix 2 is also true for the ZYA-T20S structure and was noted

previously(31), but the rotation in αV24Y T20S is greater than in ZYA-T20S (see arrows in Fig.

2A). The helical bundle composed of Helices 4, 5, and 6 (see asterisks in Fig. 2A) move the most

with some residues in these helices moving 3-4Å in the αV24Y T20S model relative to WT, which

is about twice the distance moved in the ZYA-T20S model(31). In addition, αP17 in the αV24Y

moves about 1.6Å and Helix 0 moves ~1.4Å compared to WT (Fig. 2C), which is also about twice

the distance these residues move in the ZYA-T20S structure. Taken together, the conformational

changes induced by αV24Y are highly similar to those induce by binding of the HbYX mimetic

ZYA, but they are of greater magnitude in these structures. The observed greater change is likely

attributable to the αV24Y mutation affecting the entire proteasome population within that structure.

In contrast, ZYA is a low-affinity ligand, anticipated to occupy only a subset of the binding sites in

its structure's population. Since cryo-EM analysis averages all structures, the reduced population

with conformational changes in the ZYA-bound structure represents an average of closed and

open states and thus, apparent conformational changes are smaller in the ZYA structure relative

to the αV24Y mutant.

αK66 has long been known to be critical for binding to the C-terminal carboxyl of PA's(*30, 31, 34*).

We previously observed that αK66 was repositioned by the binding of ZYA's carboxyl(*31*). Since

the inter-subunit pocket in the αV24Y T20S has no ligand bound, we expected αK66 to have a

similar conformation to WT, but we were surprised to find that K66 was similarly repositioned in

the αV24Y T20S mutant, as it was in ZYA-T20S, except to a slightly greater extent (Fig. 2D). This

suggests αK66, by itself, may play mechanistic roles in gate opening, in addition to its role in PAs

binding. Based on the comparison of the WT, ZYA bound and αV24Y T20S structures, we

conclude that the αV24Y mutation by itself induces HbYX-dependent conformational changes in

the 20S proteasome.

Comparison of V24Y T20S to the WT T20S reveals specific intra-subunit conformational changes
induced by αV24Y mutation.

It is remarkable that a single point mutation in the T20S can mimic a 3-residue motif, let alone the

activation capacity of a large complex like PAN or PA26. While the α ring is an allosteric system

with each neighboring α subunit affecting the next (e.g., a rigid body rotation of one α subunits is

expected to affect its neighbor and so on), it's also likely that HbYX binding or the αV24Y mutation

could cause conformational changes in a single α subunit as well, which could trigger allosteric

transitions to the open state. Thus, we sought to identify the intra-subunit (within a single subunit)

conformational changes that could lead to larger inter-subunit (between subunits) conformational

changes (Fig. 3A & B) that perpetuate around the entire α-ring and induce gate-opening (Fig. 2B).

The substitution of αV24 for tyrosine presents a much larger and aromatic side chain in the

hydrophobic pocket. This tyrosine could impose intra-subunit conformational changes on the

surrounding secondary structures. To measure these changes, we aligned a single α subunit from

the WT T20S to αV24Y T20S. We found that the distance between αL21 and αA154, residues

that surround the hydrophobic pocket, indicate that the pocket is widened by ~1.3A due to the

αV24Y mutation (Fig. 3C vs. D). Another measurement of this pocket width between T124 and

L21 also widens by ~1Å (Fig. 4G). However, the distance between αV24 or αV24Y to αL81 is

consistent (Fig. 3C vs. D), suggesting that the hydrophobic pocket has not enlarged in all

directions.

We hypothesize that the widening of the hydrophobic pocket that αV24Y occupies is the causal

point of origin triggering the inter-subunit conformational changes that we observed in Fig 2. The

HbYX hydrophobic pocket occupied by αV24Y side chain is adjacent to another hydrophobic

pocket found on the opposite side of Helix 0 (on the 20S central channel), where the side chain

of αI12 (an IT Switch residue) occupies in the open state (Fig. 3E & F). The αV24Y T20S model

coincides with the αV24Y T20S map, showing that the IT switch is in the open state (Fig. 1), where

the αI12 side chain is oriented to occupy the IT switch hydrophobic pocket adjacent to Helix 0

(Fig. 3E vs. F). Due to the adjacent proximity of both hydrophobic pockets (on either side of and

below Helix 0), a conformational change in the HbYX hydrophobic pocket would also affect the IT

switch's hydrophobic pocket since these pockets share common space. It is highly plausible that

the change in the HbYX hydrophobic pocket, resulting from the larger side chain of tyrosine

(compared to valine), could facilitate the switch between αT13 and αI12 in the IT switch

hydrophobic pocket (Fig. 3E vs. F). The αV24Y mutation increases the hydrophobicity and size

of the pocket, likely enhancing the binding affinity for the larger hydrophobic side chain of

isoleucine. In conclusion, it appears that occupancy of the HbYX hydrophobic pocket with an

aromatic ring of tyrosine allosterically affects the IT switch hydrophobic pocket, as intra-subunit

conformational changes allow the IT switch to reconfigure to an open state.

αProline17 (P17) displacement is one of the first identified inter-subunit conformational changes

associated with triggering gate opening, specifically by the 11S family members(*30, 33, 37*).

Interestingly, when the IT switch switches to an open configuration, it generates "slack" between

the IT switch and P17(*31*) (Fig. S4A & S4B). In αV24Y T20S, this "slack" appears to allow for an

increase in the intra-subunit distance between Helix 0 and αP17 (Fig. S4B). In addition, we

observe an intra-subunit displacement (not rotation based) of P17 by 1.5Å (Fig. 3G) consistent

with the additional slack from the switching of the IT switch. In addition, intra-subunit analysis

shows αP17 is displaced by 1Å away from its neighbor's Helix 0 (Fig. S4C); however, this distance

between P17 and the neighboring Helix 0 does not change globally (Fig. S4A & S4B) due to the

rigid body rotation of the α subunits which maintain packing between the P17 loop and the

neighboring Helix 0. In short, the conformational change of the IT switch to the open state appears

to provide flexibility for the Pro17 loop to absorb, or perhaps allow, the rigid body rotation

associated with HbYX induced gate-opening. Moreover, our structure suggests the widening of

the hydrophobic pocket between αL21 and αA154 may also contribute to the shift in αP17, as

αL21 is connected on the other side of the αP17 loop (Fig. 3G).

We observed that the side chain of αE25 is involved in an intrahelical hydrogen(H)-bond with

αR28 in the WT structure (Fig. S5A & S5B), which is displaced by the αV24Y mutation and instead

H-bonds with αR20 (Fig. S5C & S5D). The displacement slightly increases the distance between

αE25 and αR20 (Fig. SB vs. S5D) further providing "slack" for the αP17 movement. We previously

showed the mutation of αE25 to alanine, activated gate opening and prevented PAs from being

able to induce gate opening in the T20S(*31*). Therefore, biochemical evidence also supports an

important role for this αE25 rearrangement in destabilizing the closed state and perhaps

stabilizing the open state.

We also considered other mechanisms by which the αV24Y mutation might influence gate

opening such as its side chain's hydroxyl polar interactions. We find that αY24 side chain

interacted with the side chain of αD152 (Fig. 3H); however, no significant conformational changes

were observed along the β-sheet that αD152 is positioned on. Thus, it is not apparent how the

polar interaction between αY24 and αD152 might be associated with gate opening, other than the

fact that the H-bond may help orient this tyrosine rotamer.

Comparison of activated T20S forms: αV24Y T20S and ZYA-T20S

The cryo-EM structure presented here (Fig. 1 & 2) and our previously published biochemical

data(*31*) clearly demonstrate the open state of the αV24Y-T20S mutant. The substitution of αV24

for tyrosine was engineered to mimic how the "Hb" group of the HbYX motif binds to its

hydrophobic pocket, but does it? As previously mentioned, the global conformational changes in

the αV24Y 20S were similar to a HbYX-bound 20S (i.e., minimal conformational changes in the β

subunits, rigid body rotation of α subunits pivoting on Helix 2) (Fig. 1B & 2), suggesting that the

mechanism of gate-opening by αV24Y mutation has mechanistic overlap with the HbYX

mechanism (e.g. ZYA-T20S). Therefore, we sought to identify, at the molecular level, how these

two open T20S models compare and identify any intra-subunit conformational changes that can

be specifically attributed to the occupancy of the hydrophobic pocket by the Hb component of the

HbYX motif, without influence from the Y and X residues of the HbYX motif. The αV24Y T20S

model allows for analysis of the isolated effects of Hb occupancy in the hydrophobic pocket.

Comparison of αV24Y T20S against ZYA-T20S reveals that the intra-subunit conformational

changes induced by αV24Y were highly similar, as there were few differences between the α

subunits when aligned to one another (Fig. 4A, B & C). Comparison of the hydrophobic pocket

between the two open models, ZYA-T20S and αV24Y T20S, shows that the Z group of ZYA and

the αV24Y tyrosine both clearly bind to the HbYX hydrophobic pocket and overlap in space,

though they bind in different orientations (Fig. 4D). In addition, the measurements of αV24Y's side

chain or Z of ZYA to surrounding residues within each individual subunit provide the most relevant

visual approximation and overlap comparison of the "Z" and "αV24Y" aromatic group positions in

the HbYX hydrophobic pocket (Fig. 4E & F). For comparison, the distance between αV24 or

αV24Y to αL81 marginally differs by ~0.3A while the distance between αL21 and αL81 is

comparable in both models (Fig. S6A & S6B). The measurement of the hydrophobic pocket

between αL21 and αT124 is widened in the αV24Y T20S (~0.7Å; Fig 4G). This pocket

measurement also enlarges in the ZYA T20S structure compared to WT, though to a lesser extent

(Fig. 4G), consistent with the observed extent of changes in these models (Fig. 2). However, the

largest difference in this comparison between ZYA bound and αV24Y is in the measurement

between αL21 and αA154, as the V24Y tyrosine is better positioned to influence this specific

measurement. (Fig. 4H). To validate this important conformational change in the maps raw data

in a statistically significant way we used CCP-EM to generate confidence maps of WT and V24Y

structures at an FDR 0.01. These confidence maps clear show conformational differences in A154

in the V24Y structure (Fig. S3B). Lastly, the placement of αI12's side chain in the IT switch

hydrophobic pocket between Helix 0 and Helix 2 protrudes ~0.8Å deeper in the pocket in V24Y

T20S compared to ZYA-T20S (Fig. 4H), consistent with the larger hydrophobic pocket under Helix

0 in the mutant.

Collectively, evidence suggests that ZYA's mechanism as an activator is due to its hydrophobic

group occupying the space between HbYX hydrophobic pocket, similar to αV24Y, but ZYA's

mechanism also relies on others parts of the dipeptide as modifications to tyrosine affected the

small molecule's ability to activate(*38*). In combination with the evidence shown here, we conclude

that the hydrophobic group of ZYA is likely the primarily feature allosterically affecting gate-

opening but the insertion of this Hb group into the hydrophobic pocket also relies on interactions

of the Y and X residues with the inter-subunit pocket to provide sufficient binding affinity. As for

αV24Y T20S, the hydrophobic component (tyrosine's side chain) is covalently held in place in the

hydrophobic pocket and thus does not require the "YX" interactions and therefore, sufficiently

supports stable gate opening through a similar mechanism to ZYA.

αK66 supports gate opening through a previously unknown mechanism.

Many studies in the past have highlighted the importance of αK66, found in the back loop of the

inter-subunit pocket (Fig. 2D & 5A), and our αV24Y T20S structure shows that αK66 is

reorganized similar to when ZYA binds. These observations inspired us to elucidate how intra-

subunit conformational changes in αK66 may be affected by αV24Y. αK66 is well known to form

hydrogen or salt bonds with the C-termini of PAs and is required for PA binding to the 20S(*30*),

but how is αK66 affected when there is no bound PA? The back loop of the inter-subunit pocket

is naturally a more flexible loop but in αV24Y T20S, this loop does have significant intra-subunit

conformational changes (Fig. 5A), with this loop being displaced by ~3Å (inter-subunit

measurements are similar). Additionally, we noticed αK66 moved ~1Å towards αT78 and its side

chain is oriented to be in closer proximity to hydrogen bond with both αT78 and αE211 (Fig. 5A &

B). Such interactions are not observed in the WT T20S (Fig. 5C) but are seen in the ZYA-T20S

(Fig. 5D). αK66A mutants do not exhibit activation by HbYX-dependent or HbYX-independent

activators leading prior studies to conclude that αK66 was important for binding of proteasome

activators(*30, 34*). However, the repositioning of αK66 in the αV24Y T20S structure, which does

not have a proteasome activator bound, indicates that αK66 may also be involved in stabilizing

the open state on its own. To test this hypothesis, we generated a double mutant proteasome,

αV24Y-αK66A T20S. We isolated this mutant and measured fluorogenic peptide hydrolysis activity

(LFP). We found that αV24Y-αK66A T20S has similar activity to WT and is not hyperactive like

αV24Y T20S (Fig. 5E). Thus, αK66 is essential for αV24Y induced gate-opening. These enzymatic

results support the conclusion that αK66 does not only interact with proteasome activators to

support gate opening but is also critically involved in stabilizing the open state of the α subunit

gate by forming new interactions with αT78 and αE211 in the open state.

αV24Y mutation affected the hydrophilicity of the inter-subunit pocket.

Our previous study of ZYA-T20S suggested that the HbYX mechanism of activation involved the

rearrangement of water molecules, so we sought to determine if the same applies in αV24Y T20S.

The mutation essentially introduces a larger hydrophobic side chain to the inter-subunit pocket

and therefore, we expect rearrangement of water molecules. Our map and model suggest that

the αV24Y mutation either excluded or rearranged water molecules found in the WT T20S inter-

subunit pocket (Fig. 7A vs. B). Additionally, none of the water molecules placement in αV24Y

T20S coincided with the exact placement and interactions as in ZYA-T20S (Fig. 6C), suggesting

that waters involved in ZYA's binding to the inter-subunit pocket may not be relevant in αV24Y

T20S. This is unsurprising given that ZYA is a molecule that occupies a good portion of the inter-

subunit pocket, relative to the small change caused by the αV24Y mutation. Interestingly the inter-

subunit pockets of αV24Y appears to be more dehydrated relative to the WT, but structures with

equivalent resolutions would be necessary to confirm. The results collectively suggest that while

water interactions in the inter-subunit pockets change substantially in the open versus closed

states (at these resolutions), we do not observe specific waters shared between these two open

20S models that would suggest waters being critically involved in stabilization of the open gate

by these two different activation mechanisms (Fig. 6).

αV24F T20S is not activated similarly to αV24Y T20S.

Similar to αV24Y T20S, the αV24F T20S mutant showed increased basal activity compared to

WT T20S biochemically(*31*). Therefore, we sought to elucidate whether the mechanism of

activation would be similar if the substitution was to a phenylalanine as opposed to a tyrosine.

Comparison of the αV24F T20S map (Fig. 7A, Fig. S7 & S8) to αV24Y T20S and WT T20S maps

and models reveal a distinct lack of densities in the α subunits when αV24 is mutated to

phenylalanine (Fig. 7B & C). The lack of densities is most apparent from the N-termini and through

Helix 0, though the map was also lacking densities in other selected regions of the α subunit.

Globally analyzing the αV24F T20S map suggests that the αV24F mutation affected the proper

folding of secondary structures surrounding the αV24F mutation (e.g. on Helix 0), but interestingly,

this lack of Helix 0 folding did not affect the assembly of the proteasome itself (Fig. 7B & C). The

difference between maps indicates that the β subunits are minimally affected in the assembly of

the proteasome and that the lack of densities affected the α subunit throughout, though they were

most apparent around the gating regions (Fig. 7D & E). The lack of densities in αV24F T20S map

suggests that substrates could enter the mutant proteasome more easily through the larger

opening and therefore, biochemically exhibited kinetics comparable to or even exceeding the

activated proteasome(*31*).

DISCUSSION

Proteasome impairment and reduced proteostasis are common observations in clinical cases and

animal models of neurodegenerative diseases such as Alzheimer's and Parkinson's, which has

inspired the search for small molecule proteasome activators for years. Yet, there is a lack of

robust small molecule proteasome activators to date, in part due to a lack of understanding in

proteasome activation mechanisms. The α inter-subunit pockets have long been known to be the

binding site of PA complexes but the exact molecular interactions between the inter-subunit

pockets and the PAs that cause gate-opening have been elusive, preventing efficacious design

of proteasome-activating small molecules. While no FDA-approved *in vivo* proteasome-activating

small molecule exists as a tool for proof of concept, our previous work(*18*) to express constitutively

active 20S proteasomes in *C. elegans* indicated that the mutant proteasomes supported an

increased resistance to proteotoxicities and increased lifespan, supporting the rationale for this

study. Here, we present a single specific binding site within the inter-subunit pocket that when

occupied, is sufficient to induce gate opening and therefore, a plausible and well-defined

pharmaceutical target for proteasome activators.

To identify the fundamental molecular interaction(s) required for gate-opening, we focused on the

archaeal proteasome as it is homoheptameric, which is particularly an advantage when desiring

a high resolution cryo-EM structure to map detailed interactions. Moreover, the archaeal

proteasome still carries the conserved structures involved in gating, and both human and archaeal

20S can be opened by the same HbYX peptides (e.g. ZYA), implying that the findings of this study

could be applied to the human system with some caveats discussed below. We recently

demonstrated that the αV24Y mutant archaeal proteasomes exhibited upregulated activity(*31*).

Our αV24Y T20S structure here reveals inter-subunit conformational changes essentially identical

to a HbYX-bound archaeal proteasome (ZYA-T20S)(*31*), demonstrating that it is in an open state,

allowing us to further elucidate how this single residue mutation opens the gate, and by extension,

better understand the mechanism of gate opening by the HbYX motif itself.

Collectively, our findings in αV24Y T20S focus on a mechanism of gate-opening originating at the

occupancy of a hydrophobic pocket between αL21 and αT124. The αV24Y mutation places an

aromatic side chain in between αL21 and αT124 (as well αL21 and αA154), similar to the position

of the benzyl group (Z) of ZYA in the ZYA-T20S structure. The occupancy of this hydrophobic

pocket by αV24Y expands the adjacent pocket where αI12 of the IT Switch is previously found to

bind in open gate proteasome structures. αL21, displaced by αV24Y's larger side chain compared

to αV24 (Fig. 3C), is connected to the αP17 loop, which has been commonly shown to be

displaced in open gate proteasome structures. The results here also suggest that this mechanism

applies to the ZYA-T20S (Fig. 4G), though it may not be the only mechanism supporting ZYA's

gate-opening as ZYA is a dipeptide as opposed to a single residue mutation (αV24Y) that occupies

the hydrophobic pocket. For example, the YA residues bridge the subunit from αK66 to the αP17

loop, which also likely stabilizes the open state(*31*). The interaction of ZYA's carboxy group with

αK66 may also help stabilize the open-state conformation of αK66 (Fig. 5B), which we showed is

essential for gate-opening here (Fig. 5E). The conformational changes between the two models

are consistent, although αV24Y T20S exhibits larger conformational changes likely due cryo-EM

averaging effects in the lower populated ligand-bound states (ZYA) compared to the V24Y mutant.

ZYA as a dipeptide has the ability to and perhaps, even rely on interacting with other parts of the

inter-subunit pocket to supports its binding and orient the Z group in the hydrophobic pocket.

Results from both αV24Y T20S and ZYA-T20S both suggest that occupying the hydrophobic

pocket is a key to gate opening in the archaeal proteasome, and potentially the mammalian

proteasome if placed in the right pocket(s).

The two key residues that "sandwiches" αV24Y's side chain and forms the hydrophobic pocket,

αL21 and αA154, are conserved in the human 20S α inter-subunit pockets (Fig. S9), indicating

that a similar mechanism of occupying this hydrophobic pocket is used for gate-opening in the

human system. Five out of seven H20S α subunits carry leucine in a similar position to αL21 in

the T20S. The remaining two subunits carry an isoleucine (α6: PSMA1) and valine (α7: PSMA3),

which are hydrophobic like leucine. Additionally, while αA154 is only conserved as alanine in α1:

PSMA6, the rest of the subunits carry a methionine in this position, which is also hydrophobic.

Previous analysis(*31*) also indicated that the H20S has analogs of the IT switch in selected α

subunits, which suggests that targeting this hydrophobic pocket at selected α subunits with the

conserved IT switch would have a greater effect on gate-opening than others. Any application of

this mechanism in the human system is still complicated and should also consider the prior

prediction that multiple inter-subunit pockets likely need to be occupied by the HbYX motifs for

gate-opening(*38*). The conservation of activation motif, residues, and conformational changes

between archaeal and human proteasomes for gate opening demonstrates potential for the

findings in this study to translate to activating the human 20S.

Interestingly, our αV24F T20S structure indicates that the position of this residue in the protein

chain plays a role in α subunit folding. Our αV24F T20S cryo-EM map reveals that the α subunits

are partially assembled, with the majority of regions proximal to the gating residues lacking

densities, suggesting disorder. Yet, the β subunits appear functional, both structurally and

biochemically(*31*), which suggests that the proteasome does not require perfectly formed α

subunits for complete proteasome assembly. We notice that the hydroxyl group of αV24Y's side

chain is involved in polar interactions; however, we do not presume this to be involved in α subunit

assembly as valine lacks the side chain to interact similarly. Rather, we speculate that αV24F may

introduce too large of a hydrophobic group that, without a hydroxyl to interact with αD152 (Fig.

3H) prevents proper folding interactions.

Collectively, our analysis indicates that the HbYX hydrophobic pocket expands due to αV24Y's

more bulky side chain, which likely sets of a chain of events starting with the IT switch, which are

modeled and schematized in Figure 8. The wider and now more hydrophobic IT switch pocket

likely promotes IT switch to switch to the open state, providing flexibility for the αP17 loop intra-

subunit motions, or "slack".   In addition, the widening of the hydrophobic pocket and

rearrangement of the αE25 H-bonding pattern also appears to allow more flexibility for αP17 loop

motions. Combined, these intra-subunit conformational changes collectively elongate the αP17

loop, increasing its flexibility. This enhanced flexibility is crucial because the αP17 loop interfaces

with Helix 0 of neighboring α subunits. The loop's increased plasticity accommodates rigid-body

rotations against the adjacent α subunits, which would otherwise be constrained. As a result,

these rotational movements can propagate allosterically around the proteasome α ring. Though

this causal chain of events is supported by these structures, the timing and order of these events

cannot be elucidated with these static structures. The structures here suggest a plausible

molecular mechanism of how a single residue mutation in the HbYX hydrophobic pocket can

induce 20S gate opening (Fig. 8).

Our findings in this study also highlight and revise an understanding of the αK66's role in

proteasomal gate-opening. Previous findings have consistently alluded to its critical role for the

association of HbYX-dependent and HbYX-independent PAs, as αK66 was shown to interact with

and orient the C-terminal tails of PAs for gate-opening(*30*, *31*, *35*, *37*, *39*). To our surprise, the

αV24Y T20S structure demonstrates that αK66's side chain conformation changes, even though

it does not interact with a PA or HbYX mimetic such as ZYA. Our double mutant αV24Y&K66A

T20S's basal activity supports the necessity of the αK66 conformational change observed in the

αV24Y model and indicates the involvement of αK66 in stabilizing the open-gate conformation

through a previously unreported mechanism. The conformational change in αK66 is supported by

changes in H-bonding interactions between αK66 and surrounding residues, which are absent in

the closed structure (WT T20S). These findings therefore demonstrate that αK66's conformational

changes support other conformational changes such as the conformation of the back loop, which

it is connected to, and has been shown to be involved in HbYX-dependent gate opening(*31*).

Essentially, αK66 is deduced to contribute more than interacting with activators, and also appears

to be necessary to stabilize the open state of the 20S.

Our prior study showed the possible roles waters could play in proteasome activation and gating

regulation. Our ZYA-T20S structure indicated that the hydroxyl of ZYA's tyrosine side chain

interacted with a water molecule that further hydrogen bonded with αL21 and αE25. To elucidate

the relevance and importance of this water interaction, we mutated αE25(*31*). αE25A proteasome

mutants exhibited an upregulated basal activity that was not further activated by PAN or PA26,

suggesting it plays an important role in gating. To elucidate whether a similar mechanism occurs

in αV24Y T20S open gate conformation, we sought to identify where waters might be found within

the inter-subunit pocket. Broadly, we observed significantly fewer waters in the inter-subunit

pockets of αV24Y T20S than we did in the WT T20S and ZYA-T20S. We note that αV24Y T20S's

resolution is not as high as ZYA-T20S but is reasonable for predicting water coordinates. Because

no common water was found in the two different open states, the necessity and role of specific

water molecule interactions in gate-opening could not be elucidated from these structures.

Ultimately, the findings described indicate that the occupancy of the hydrophobic pocket within

the α inter-subunit pocket is sufficient to induce gate-opening in archaeal proteasomes. The

application of this finding in the mammalian system relies on further identification of the critical

inter-subunit pockets that need to be occupied by the HbYX motif in that system to stabilize gate-

opening. Our prior studies involving ZYA have indicated a minimum of two HbYX motifs binding

for gate-opening in the mammalian system(*38*). However, the mammalian 20S has 7 different

pockets that can be occupied in different ways by 6 different Rpt1-6 C-termini, so the specific

inter-subunit pockets that need C-termini/HbYX motifs bound to activate gate opening has not

been worked out. All cryo-EM studies to date agree that occupancy of five Rpt C-termini is

sufficient for gate opening, but which of those are required for gate opening is not known. Further

progress on the HbYX mechanism based on the results thus far would be to identify the specific

inter-subunit pockets that need to be occupied by the HbYX motif to induce gate-opening in the

mammalian proteasomes.

The work presented here is conducted with the intentions of understanding proteasome activation

mechanisms and developing the molecular framework for restoring or activating proteasome

activity to combat the impairment consistently observed in neurodegenerative diseases, which is

barred by our limited understanding of proteasome regulation. Here, we advance the field's

understanding of proteasomal gate-opening by extending beyond the vague idea that HbYX motif

binding induces gate opening, to pinpointing a sole molecular interaction that significantly and

sufficiently opens the proteasome's gate. Our findings inform proteasome activating drug

discovery efforts to focus on small molecules that occupies the HbYX hydrophobic pocket.

MATERIALS AND METHODS

Proteasome purifications

T. acidophilum wild-type 20S, αV24Y 20S, or αV24F 20S proteasomes were similarly purified as described(*40*), using the 8XHis tags on the C-terminus of the β subunits. All T20S mutants were generated by overlapping PCR site-directed mutagenesis.

Proteasome activity assays

Fluorogenic substrate peptide, LFP (Mca-AKVYPYPME-Dpa(Dnp)-amide), were synthesized by from EZBiolabs. LFP was dissolved in DMSO and incubated with proteasomes at 3µM final concentration. The final concentration of DMSO in activity assays was 1%. Protein concentrations were determined by Bradford assay (Thermo Scientific). 0.25ug archaeal proteasomes and LFP peptides were added to 25uL Tris-based reaction buffer at 45ºC. Assays were performed for 30mins to an hour and analyzed using BioTek Gen5 Data Analysis software. Activity was measured as relative fluorescence units/minute (rfu/min), generating a curve which was used to calculate the initial velocity, according to the slope of the curves. "Fold Activation" was calculated by dividing the average initial velocity of mutant proteasomes against the average initial velocity of wild type proteasomes.

Cryo-EM Sample Preparation and Data Collection

Copper Quantifoil R 1.2/1.3 300 mesh (EMS) grids were cleaned using a PELCO easiGlow Glow Discharge cleaning system. A volume of 3 uL of 0.75mg/mL αV24Y-T20S or αV24F T20S (suspended in 50mM Tris pH 7.4, 150mM NaCl) sample was placed onto a grid, and then flash frozen in liquid ethane using a manual plunge freeze apparatus. Data collection was done using a Titan Krios transmission electron microscope (Thermo Fisher) operated at 300kW and a magnification of x81,000, which resulted in 0.503Å/px. Images were collected using a Falcon IIIEC

direct electron detector camera equipped with a K3/GIF operating in counting and super resolution

modes. Electron dose per pixel of 50 e-/Å2 was saved as 40 frame movies within a target defocus

range of -2.5 to -1.25. All the data was collected using cryoSPARC software (Structura

Biotechnology Inc)(*41*).

Cryo-EM Single Particle Analysis

Cryo-EM images of the αV24Y T20S and αV24F T20S proteasome were analyzed using

cryoSPARC. Schematic for cryo-EM single-particle data processing available in supplement.

αV24Y T20S: From 3663 movies collected, we picked 396542 particles after two rounds of 2D

Classification to generate an Ab-initio model, which was used for homogeneous refinement (using

D7 symmetry). Particle processing scheme and validations are shown in Fig. S1 & S2.

αV24F T20S: From 4920 movies collected, we picked 542722 after two rounds of 2D

Classification to generate an Ab-initio model, which was used for homogenous refinement (using

D7 symmetry). Particle processing scheme and validations are shown in Fig. S6 & S7.

All representations of the T20S proteasome complex were created using PyMOL 2.5.4 and UCSF

ChimeraX v1.7.1(*42, 43*).

Confidence map generation

Small conformational changes (e.g. ~1Å) with larger allosteric effects altering 20S gating and

enzymatic activity are common features in the 20S proteasome, which has been published

extensively. However, to more rigorously demonstrate that these changes are statistically

significant at the map data level, we used CPP-EM to generate confidence maps of our WT and

V24Y structures (Fig. S3B). Confidence maps statistically interpret cryo-EM densities by testing

each voxel against background noise and controlling the false discovery rate (FDR). We

thresholded our maps of WT and V24Y at an FDR of 0.01 to highlight regions with statistically

significant density. We then overlayed these maps in Chimera to show they clearly show

conformational changes even at an FDR of 0.01, especially at the specific regions the reviewer

was concerned about.

Atomic model building

The atomic models were built using PDB: 8F7K as a template, rigid body fitting into the electron

density map using PHENIX 1.19.2-4158(*44–46*). The docked models were subjected to a cycle

of morphing and simulated annealing, five real-space refinement macrocycles with atomic

displacement parameters, secondary structure restraints, and local grid searched in PHENIX.

Consequently, the models were refined by oscillating between manual real-space refinement in

WinCoot(*47*) 0.9.8.1 EL and real-space refinement in PHENIX (five macrocycles, without

morphing and simulated annealing). Waters were added to models using PHENIX Douse. The

final model of αV24Y T20S was deposited at wwPDB: 9BUZ.

Statistical analysis

Data were analyzed in Graph Pad or excel using an unpaired Student's t-test (Prism). For all

statistical analyses, a value of $p < 0.05$ was considered significant.

REFERENCES

1. V. Cohen-Kaplan, I. Livneh, N. Avni, C. Cohen-Rosenzweig, A. Ciechanover, The ubiquitin-proteasome system and autophagy: Coordinated and independent activities. *Int. J. Biochem. Cell Biol.* **79**, 403–418 (2016).

2. W. Bialek, J. F. Collawn, R. Bartoszewski, Ubiquitin-Dependent and Independent Proteasomal Degradation in Host-Pathogen Interactions. *Molecules* **28**, 6740 (2023).

3. M. Demasi, F. M. da Cunha, The physiological role of the free 20S proteasome in protein degradation: A critical review. *Biochim. Biophys. Acta BBA - Gen. Subj.* **1862**, 2948–2954 (2018).

4. A. Khaminets, C. Behl, I. Dikic, Ubiquitin-Dependent And Independent Signals In Selective Autophagy. *Trends Cell Biol.* **26**, 6–16 (2016).

5. Y. Makaros, A. Raiff, R. T. Timms, A. R. Wagh, M. I. Gueta, A. Bekturova, J. Guez-Haddad, S. Brodsky, Y. Opatowsky, M. H. Glickman, S. J. Elledge, I. Koren, Ubiquitin-independent proteasomal degradation driven by C-degron pathways. *Mol. Cell* **83**, 1921-1935.e7 (2023).

6. H. Tsuchiya, A. Endo, Y. Saeki, Multi-Step Ubiquitin Decoding Mechanism for Proteasomal Degradation. *Pharm. Basel Switz.* **13**, 128 (2020).

7. A. N. Hegde, Proteolysis, synaptic plasticity and memory. *Neurobiol. Learn. Mem.* **138**, 98–110 (2017).

8. H. Park, B.-K. Kaang, Balanced actions of protein synthesis and degradation in memory formation. *Learn. Mem.* **26**, 299–306 (2019).

9. M. B. Patrick, N. Omar, C. T. Werner, S. Mitra, T. J. Jarome, The ubiquitin-proteasome system and learning-dependent synaptic plasticity – A 10 year update. *Neurosci. Biobehav. Rev.* **152**, 105280 (2023).

10. P. Z. Y. Lip, M. Demasi, D. Bonatto, The role of the ubiquitin proteasome system in the memory process. *Neurochem. Int.* **102**, 57–65 (2017).

11. T. J. Jarome, R. K. Devulapalli, The Ubiquitin-Proteasome System and Memory: Moving Beyond Protein Degradation. *The Neuroscientist* **24**, 639–651 (2018).

12. J. N. Keller, K. B. Hanni, W. R. Markesbery, Impaired Proteasome Function in Alzheimer's Disease. *J. Neurochem.* **75**, 436–439 (2000).

13. A. Ciechanover, P. Brundin, The ubiquitin proteasome system in neurodegenerative diseases: sometimes the chicken, sometimes the egg. *Neuron* **40**, 427–446 (2003).

14. K. S. McNaught, C. W. Olanow, B. Halliwell, O. Isacson, P. Jenner, Failure of the ubiquitin-proteasome system in Parkinson's disease. *Nat. Rev. Neurosci.* **2**, 589–594 (2001).

15. Z. Ortega, M. Díaz-Hernández, J. J. Lucas, Is the ubiquitin-proteasome system impaired in
Huntington's disease? *Cell. Mol. Life Sci. CMLS* **64**, 2245–2257 (2007).

16. C. McKinnon, S. J. Tabrizi, The ubiquitin-proteasome system in neurodegeneration.
Antioxid. Redox Signal. **21**, 2302–2321 (2014).

17. K. S. P. McNaught, L. M. Björklund, R. Belizaire, O. Isacson, P. Jenner, C. W. Olanow,
Proteasome inhibition causes nigral degeneration with inclusion bodies in rats. *Neuroreport*
13, 1437–1441 (2002).

18. R. T. Anderson, T. A. Bradley, D. M. Smith, Hyperactivation of the proteasome in
Caenorhabditis elegans protects against proteotoxic stress and extends lifespan. *J. Biol.*
Chem. **298**, 102415 (2022).

19. E. Njomen, J. J. Tepe, Proteasome Activation as a New Therapeutic Approach To Target
Proteotoxic Disorders. *J. Med. Chem.* **62**, 6469–6481 (2019).

20. J. J. S. VerPlank, S. Lokireddy, J. Zhao, A. L. Goldberg, 26S Proteasomes are rapidly
activated by diverse hormones and physiological states that raise cAMP and cause Rpn6
phosphorylation. *Proc. Natl. Acad. Sci.* **116**, 4228–4237 (2019).

21. J. J. S. VerPlank, S. D. Tyrkalska, A. Fleming, D. C. Rubinsztein, A. L. Goldberg, cGMP via
PKG activates 26S proteasomes and enhances degradation of proteins, including ones
that cause neurodegenerative diseases. *Proc. Natl. Acad. Sci.* **117**, 14220–14230 (2020).

22. N. Myeku, H. Wang, M. E. Figueiredo-Pereira, cAMP stimulates the ubiquitin/proteasome
pathway in rat spinal cord neurons. *Neurosci. Lett.* **527**, 126–131 (2012).

23. X. Wang, J. Yen, P. Kaiser, L. Huang, Regulation of the 26S Proteasome Complex During
Oxidative Stress. *Sci. Signal.* **3**, ra88–ra88 (2010).

24. M. Groll, M. Bajorek, A. Köhler, L. Moroder, D. M. Rubin, R. Huber, M. H. Glickman, D.
Finley, A gated channel into the proteasome core particle. *Nat. Struct. Biol.* **7**, 1062–1067
(2000).

25. J. Rabl, D. M. Smith, Y. Yu, S.-C. Chang, A. L. Goldberg, Y. Cheng, Mechanism of gate
opening in the 20S proteasome by the proteasomal ATPases. *Mol. Cell* **30**, 360–368
(2008).

26. Z. Ding, C. Xu, I. Sahu, Y. Wang, Z. Fu, M. Huang, C. C. L. Wong, M. H. Glickman, Y.
Cong, Structural Snapshots of 26S Proteasome Reveal Tetraubiquitin-Induced
Conformations. *Mol. Cell* **73**, 1150-1161.e6 (2019).

27. Y. Dong, S. Zhang, Z. Wu, X. Li, W. L. Wang, Y. Zhu, S. Stoilova-McPhie, Y. Lu, D. Finley,
Y. Mao, Cryo-EM structures and dynamics of substrate-engaged human 26S proteasome.
Nature **565**, 49–55 (2019).

28. M. Wehmer, T. Rudack, F. Beck, A. Aufderheide, G. Pfeifer, J. M. Plitzko, F. Förster, K.
Schulten, W. Baumeister, E. Sakata, Structural insights into the functional cycle of the
ATPase module of the 26S proteasome. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 1305–1310
(2017).

29.   M. R. Eisele, R. G. Reed, T. Rudack, A. Schweitzer, F. Beck, I. Nagy, G. Pfeifer, J. M.
Plitzko, W. Baumeister, R. J. Tomko, E. Sakata, Expanded Coverage of the 26S
Proteasome Conformational Landscape Reveals Mechanisms of Peptidase Gating. *Cell*
Rep. **24**, 1301-1315.e5 (2018).

30.   A. Förster, E. I. Masters, F. G. Whitby, H. Robinson, C. P. Hill, The 1.9 A structure of a
proteasome-11S activator complex and implications for proteasome-PAN/PA700
interactions. *Mol. Cell* **18**, 589–599 (2005).

31.   J. J. Y. Chuah, M. S. Rexroad, D. M. Smith, High resolution structures define divergent and
convergent mechanisms of archaeal proteasome activation. *Commun. Biol.* **6**, 1–16
(2023).

32.   T. L. Religa, R. Sprangers, L. E. Kay, Dynamic regulation of archaeal proteasome gate
opening as studied by TROSY NMR. *Science* **328**, 98–102 (2010).

33.   B. M. Stadtmueller, C. P. Hill, Proteasome activators. *Mol. Cell* **41**, 8–19 (2011).

34.   D. M. Smith, S.-C. Chang, S. Park, D. Finley, Y. Cheng, A. Goldberg, Docking of the
Proteasomal ATPases' C-termini in the 20S Proteasomes alpha Ring Opens the Gate for
Substrate Entry. *Mol. Cell* **27**, 731–744 (2007).

35.   A. H. de la Peña, E. A. Goodall, S. N. Gates, G. C. Lander, A. Martin, Substrate-engaged
26S proteasome structures reveal mechanisms for ATP-hydrolysis–driven translocation.
Science **362** (2018).

36.   Y. Yu, D. M. Smith, H. M. Kim, V. Rodriguez, A. L. Goldberg, Y. Cheng, Interactions of
PAN's C-termini with archaeal 20S proteasome and implications for the eukaryotic
proteasome–ATPase interactions. *EMBO J.* **29**, 692–702 (2010).

37.   C. P. Hill, E. I. Masters, F. G. Whitby, The 11S regulators of 20S proteasome activity. *Curr.*
Top. Microbiol. Immunol. **268**, 73–89 (2002).

38.   J. J. Chuah, T. A. Thibaudeau, D. M. Smith, Minimal mechanistic component of HbYX-
dependent proteasome activation that reverses impairment by neurodegenerative-
associated oligomers. *Commun. Biol.* **in press** (2023).

39.   Y. Dong, S. Zhang, Z. Wu, X. Li, W. L. Wang, Y. Zhu, S. Stoilova-McPhie, Y. Lu, D. Finley,
Y. Mao, Cryo-EM structures and dynamics of substrate-engaged human 26S proteasome.
Nature **565**, 49–55 (2019).

40.   I. T. Dorn, R. Eschrich, E. Seemüller, R. Guckenberger, R. Tampé, High-resolution AFM-
imaging and mechanistic analysis of the 20 S proteasome. *J. Mol. Biol.* **288**, 1027–1036
(1999).

41.   A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

42.   T. D. Goddard, C. C. Huang, E. C. Meng, E. F. Pettersen, G. S. Couch, J. H. Morris, T. E.
Ferrin, UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein*
Sci. Publ. Protein Soc. **27**, 14–25 (2018).

43.   E. F. Pettersen, T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, T. I. Croll, J. H.
Morris, T. E. Ferrin, UCSF ChimeraX: Structure visualization for researchers, educators,
and developers. *Protein Sci. Publ. Protein Soc.* **30**, 70–82 (2021).

44.   D. Liebschner, P. V. Afonine, M. L. Baker, G. Bunkóczi, V. B. Chen, T. I. Croll, B. Hintze, L.-
W. Hung, S. Jain, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, B. K. Poon, M. G. Prisant, R.
J. Read, J. S. Richardson, D. C. Richardson, M. D. Sammito, O. V. Sobolev, D. H.
Stockwell, T. C. Terwilliger, A. G. Urzhumtsev, L. L. Videau, C. J. Williams, P. D. Adams,
Macromolecular structure determination using X-rays, neutrons and electrons: recent
developments in Phenix. *Acta Crystallogr. Sect. Struct. Biol.* **75**, 861–877 (2019).

45.   P. D. Adams, P. V. Afonine, G. Bunkóczi, V. B. Chen, I. W. Davis, N. Echols, J. J. Headd,
L.-W. Hung, G. J. Kapral, R. W. Grosse-Kunstleve, A. J. McCoy, N. W. Moriarty, R. Oeffner,
R. J. Read, D. C. Richardson, J. S. Richardson, T. C. Terwilliger, P. H. Zwart, PHENIX: a
comprehensive Python-based system for macromolecular structure solution. *Acta
Crystallogr. D Biol. Crystallogr.* **66**, 213–221 (2010).

46.   P. V. Afonine, B. K. Poon, R. J. Read, O. V. Sobolev, T. C. Terwilliger, A. Urzhumtsev, P. D.
Adams, Real-space refinement in PHENIX for cryo-EM and crystallography. *Acta
Crystallogr. Sect. Struct. Biol.* **74**, 531–544 (2018).

47.   P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta
Crystallogr. D Biol. Crystallogr.* **66**, 486–501 (2010).

ACKNOWLEDGEMENTS

We thank the members of the Smith Lab for the helpful and valuable discussions, especially Giovanni Howells for his meticulous review of this manuscript. We also thank Thomas (Tom) C. Terwilliger, PhD at the Los Alamos National Laboratory for his gracious guidance and assistance in using Phenix. Transmission electron micrographs were recorded at the University of Virginia Molecular Electron Microscopy Core facility (RRID:SCR_019031), which is supported in part by the School of Medicine and built with NIH grant G20-RR31199. In addition, the Titan Krios (S10-RR025067), Falcon II/3EC direct detector (S10-OD018149), and K3/GIF (U24-GM116790) were purchased in part or in full using the designated NIH grants. Molecular graphics and analyses performed with UCSF ChimeraX, developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy, and Infectious Diseases(*43*).

FUNDING

This work was supported by National Institute of Health grant R01AG064188 (DMS).

AUTHOR CONTRIBUTIONS

JJYC and **MRD** prepared WT T20S, αV24Y T20S, and αV24F T20S for cryo-EM and activity assays. **JJYC** and **DMS** analyzed cryo-EM data, reconstructed the αV24Y T20S and αV24F T20S structures. **JJYC** and **MRD** performed enzymatic assays of the archaeal proteasomes. **JJYC** and **DMS** generated PyMOL figures. Results were analyzed and interpreted by **JJYC** and **DMS.** Manuscript was prepared by **JJYC**, **MRD**, and **DMS.** All authors reviewed the results and approved the final version of this manuscript.

COMPETING INTERESTS

All authors declare that they have no competing interests.

DATA AND MATERIALS AVAILABILITY

Cryo-EM maps are deposited in the Electron Microscopy Data Bank (EMDB) under accession

codes, EMD-44914 (αV24F T20S) and coordinates are available from the RCSB Protein Data

Bank under accession codes, 9BUZ (αV24Y T20S), 8F7K (ZYA-T20S), 8F6A (WT T20S). The

authors declare that data supporting the findings of this study are available within the paper and

its supplementary information files and are available from the corresponding author upon request.

Main Figures Legend

Fig. 1: Overlay of cryo EM maps shows αV24Y T20S has similar conformational changes as ZYA-bound T20S and is in the gate-open state.

Fig. 2: Atomic model of αV24Y T20S compared to WT and ZYA-T20S shows it has undergone HbYX-dependent conformational changes associated with T20S gate-opening.

Fig. 3: αV24Y mutation induces intrasubunit conformational changes in the HbYX hydrophobic pocket, IT switch pocket, and αP17 loop.

Fig. 4: αV24Y mutation and ZYA binding both induce similar intrasubunit conformational changes.

Fig. 5: αLysine66 plays a key role in stabilizing the open state of the T20S, even in the absence of proteasome activator binding.

Fig. 6: The intersubunit pocket of the αV24Y T20S is less populated with waters, relative to ZYA-T20S and WT T20S.

Fig. 7: A large portion of the N-terminal domain of the αV24F T20S mutant is unstructured.

Fig. 8: Model for induction of gate opening by binding of the Hb group binding to the HbYX hydrophobic site in the 20S intersubunit pockets.

WT T20S **αV24Y T20S** **ZYA-T20S**

Figure 1

Fig. 1: Overlay of cryo EM maps shows αV24Y T20S has similar conformational changes as ZYA bound T20S and is in the gate-open state.

A Overlay of wild-type (WT) T20S (yellow) and αV24Y T20S (red) electron density maps, showing the α and β rings. Whole maps were aligned to one another.

B Overlay of αV24Y T20S (red) and ZYA-T20S (blue) electron density maps, showing the α and β rings. V24Y T20S shows similar but larger conformational changes than does ZYA-T20S.

C Close up of the intersubunit pocket in the WT T20S (yellow) electron density map, highlighted in **A** in the larger neon green box outline.

D, E Same as **C**, but for αV24Y T20S (red) and ZYA-T20S (blue). Arrows point to the density corresponding to αV24Y and Z of ZYA, respectively.

F Close up of the WT T20S (yellow) electron density map, corresponding to the YDR motif, the missing density for tyrosine is highlighted by the neon green dotted circle, which is expected in the closed state. The relative position of the close-up is indicated by the smaller neon blue box outline in **A**.

G, H Same as **F**, but for αV24Y T20S (red) and ZYA-T20S (blue). Neon green dotted circle highlight the YDR tyrosine side chain density, which is expected to be in this position in the open gate conformation.

I Close up of the WT T20S (yellow) electron density map, corresponding to the IT Switch, highlighted by the neon green dotted circle.

J, K Same as **I** but for αV24Y T20S (red) and ZYA-T20S (blue). Changes in the conformation of the IT switch indicates switching from the closed to the open state.

WT T20S   **αV24Y T20S**   **ZYA-T20S**

Fig. 2: Atomic model of αV24Y T20S compared to WT and ZYA-T20S shows it has undergone HbYX-dependent conformational changes associated with 20S gate-opening.

A Atomic model overlay of WT T20S (yellow), αV24Y T20S (red), and ZYA-T20S (blue) with bound ZYA (gold), aligned by the β rings to show conformational changes in the α subunits. Top half of sideview of 20S is shown. Conformational changes in the α-ring are clearly visible (black arrows). Helix 4, 5, and 6 are labeled with black asterisks outlined in white.

B Top view of **A**. Conformational changes noted by (black arrows).

C Closeup view of one of the α intersubunit pockets as in **B**. αPro17 is shown with a black arrow. Helix 0 is labeled with a black asterisk outlined in white.

D Closeup view of αK66 which is labeled and shown in sticks.

Models in panel b through d are aligned to show intersubunit conformational changes (i.e., aligned by β rings)

Fig. 3: αV24Y mutation induces intrasubunit conformational changes in the HbYX hydropbobic pocket, IT switch pocket, and P17 loop.

A Overlay of WT T20S (yellow) and αV24Y T20S (red) α subunit models, aligned by the β rings.

B Same as **A** except the models were aligned according to individual α subunit.

C Close up of the intersubunit pocket in αV24Y T20S (red), showing distances between the α-carbon selected residues (labeled and shown in sticks) to demonstrate conformational changes when compared to the WT T20S (yellow) model in **D**. Models are aligned by individual α subunit.

D Same as **C** except it is the WT T20S (yellow) model.

E Close up on the residues of the IT Switch (sticks) and αV24Y (sticks), showing the distance between the α-carbon of αV24Y and αA154 (sticks) to demonstrate conformational changes in the open state, when compared to the WT T20S (yellow) model in **F**. Models are aligned by individual α subunit.

F Same as **E** except it is the WT T20S (yellow) model.

G Close up of Helix 0 and selected surrounding residues (shown in sticks) in an overlay of WT T20S (yellow) and αV24Y T20S (red) α subunits, aligned by individual α subunit. Distances are measured between α-carbons and colored according to the model it corresponds to.

H Close up of αV24Y (shown in sticks) in the αV24Y T20S model, demonstrating its side chain's polar interaction (yellow dotted line) with αD152 (shown in sticks).

Figure 4

Fig. 4: αV24Y mutation and ZYA binding both induce similar intrasubunit conformational changes.

A Same as **3A** except for αV24Y T20S (red) and ZYA-T20S (blue). Models are aligned by the β ring.

B Same as **3B** except for αV24Y T20S (red) and ZYA-T20S (blue). Models are aligned by individual α subunit.

C Same as **B** except showing a closeup of the Helix 4, 5, and 6 bundle.

D αV24Y T20S and ZYA-T20S are aligned by β subunits and αV24Y side chain and carboxybenzyl group (Z) of ZYA are both shown in space-fill model to appreciate the level of spatial overlap between the αV24Y tyrosine ring (red) and the Z ring (gold) of ZYA, demonstrating they each occupy the same HbYX hydrophobic pocket in the T20S.

E Close up of the αV24Y T20S (red) α intersubunit pocket showing measurements between the αV24Y tyrosine and labeled α carbons (shown in sticks). Comparison against **F** indirectly demonstrates the position of αV24Y's side chain relative to the carboxybenzyl group of ZYA in the ZYA-T20S model.

F Measurements of distances between ZYA's carboxybenzyl group (gold) and the α-carbon of surrounding residues (shown in sticks) in the ZYA-T20S model (blue).

G The HbYX hydrophobic pocket measurement: αT124 to αL21 (main chain shown in sticks) distance in WT T20S (yellow), αV24Y T20S (red), and ZYA-T20S (blue).

H Same as **3E & F**, except with alignment of ZYA-T20S (blue) and αV24Y T20S (red) α subunit models. Distances noted are colored according to the models they correspond to representing intrasubunit differences.

Figure 5

Fig. 5: αK66 plays a key role in stabilizing the open state of the T20S, even in the absence of proteasome activator binding.

A Overlay of WT T20S (yellow) and αV24Y T20S (red), aligned by individual α subunits, showing a close-up of the αK66 (shown in sticks) back loop. Distance measured demonstrates the extent of intrasubunit conformational changes in the "back loop" and K66 α carbon.

B Close up of αV24Y T20S (red) model, showing the polar interactions (yellow dotted lines) between αK66 and surrounding residues (shown in sticks). Distances are measured between the exact atoms involved in the polar interactions.

C Same as **B** except it is the WT T20S (yellow) model.

D Same as **B** except it is the ZYA-T20S (blue) model and ZYA (gold).

E WT T20S and mutants were measured for activity based on LFP degradation rate (rfu/min), normalized to the WT T20S as a control. Data (means) are representative of three or more independent experiments, each performed in triplicate. Error bars represent ± standard deviation.

A

Water	Residue
	α_1G166
	α_1T124
	α_1A154
	α_1I157
	α_1N158
	α_1T156
	α_2A29
	α_2A83
	α_2G80
	α_2V82

B

Water	Residue
	α_1D150
	α_1T156
	α_1N158
	α_2K33
	α_2G80

C

Water	Residue
	ZYA-Tyr
	α_1L21
	α_1E25
	α_1E29
	α_1D150
	α_1T156
	ZYA-backbone
	α_2E29
	α_2K33
	α_2V82
	α_2A83
	α_2D84

Figure 6

Fig. 6: The interunit pocket of the αV24Y T20S is less populated with waters, relative to ZYA T20S and WT T20S.

A Intersubunit pocket of WT T20S model (yellow) showing waters (spheres) and their interactions (yellow dotted lines) with the residues (shown in sticks, listed in table) in the intersubunit pocket. Spheres are colored individually and match the colors on the table (right) to identify residues it interacts with.

B, C Same as **A** except αV24Y T20S model (red) and ZYA-T20S model (blue).

A

b

αV24F T20S

B

α ring

β ring

C

D

αV24Y T20S
subtract
αV24F T20S Map

E

WT T20S (EMDB-28878)
subtract
αV24F T20S Map

Figure 7

Fig. 7: A large portion of the N-terminal domain of the αV24F T20S mutant is unstructured.

A Angled view of αV24F T20S 2.2Å electron density map (green), showing the α and β rings.

B Overlay of αV24F T20S map with αV24Y T20S map (red; transparent surface; left) and WT T20S map (yellow; transparent surface; right).

C Same as **B** except αV24Y T20S model (red, cartoon, left) and WT T20S model (yellow; cartoon; right).

E Difference map generated by subtracting αV24F T20S map from αV24Y T20S map, using ChimeraX.

F Same as **E** except WT T20S map instead of αV24Y T20S map.

Fig. 8: Model for induction of gate opening by binding of the Hb group binding to the HbYX hydrophobic site in the 20S intersubunit pockets.

Binding of the Hb group of the HbYX motif, or tyrosine of αV24Y, is expected to initiate an allosteric chain of events that leads to 20S gate opening. Hb group binding causes:

1) Increase the size of the Hb pocket
2) Affects IT switch Hb pocket promoting IT switching to open state
3) IT switching lengthens P17 loop (increases loop slack)
4) E25 H-bond rearrangement further lengthens P17 loop
5) Combine effects on P17 loop slack promotes rigid body rotations against the neighboring Helix 0.
6) Rigid body rotations effect back loop (not shown), which repositions K66 to further stabilize open state.

A

3663 micrographs
965963 particles

↓

2 rounds of 2D
Classification

↓

396542 particles

↓

Homogenous
refinement
D7

B

C

2.38Å

Fig. S1: αV24Y T20S Processing Scheme
A Cryo-EM workflow for alphaV24Y T20S. All steps were performed in Cryosparc.
B Final 2D class averages after 2 rounds of 2D Classification shown with a mixture of top and size views.
C Homogenous refinement of class of particles that showed the open gate conformation.

Fig. S2: αV24Y T20S Validation

A Micrograph representative showing αV24Y T20S. Scale bar represents 5nm.

B. Standard FSC-0.143 graph showing corrected 2.38Å resolution.

C View of particle angle distribution showing a mixture of side and top view particles.

D Particles from selected ab-initio model were refined with homogenous refinement and D7 symmetry applied. D7 symmetry was confirmed as appropriate by homogenous refinement with C1 symmetry; the resulting C1 map was copied then flipped 180⁰ around the 2-fold axis then rotated 180⁰ about the 7-fold axis. The flipped and rotated map was essentially identical to the original, confirming D7 symmetry of the complex. Final 2.38Å D7 map was colored in Chimera to show range of resolution of 3D reconstructed map.

Fig. S3: αV24Y T20S differ from WT T20S

A Close up of αV24Y T20S sharpened map and model, demonstrating quality of map and model fit in the shown region.

B. Overlay of CCP-EM generated confidence maps for αV24Y T20S (yellow) and WT T20S (cyan), indicating statistically significant conformational changes. Maps were set at a threshold of 0.01 FDR. View is similar to Figure 4H showing clear conformational changes in A154 relative to helix 0. Arrows point out the distance differences between the same two residues in the WT or the V24Y maps. Gaussian smoothing was applied to both maps.

Fig. S4: Additional measurements in the intersubunit pocket in V24Y T20S. Complementary to Figure 3.
A Measurement between P17 and neighboring Helix 0 for β ring-aligned WT and αV24Y T20S structures representing absence of intersubunit conformational differences in this distance.
B Measurement between P17 and its own Helix 0 (E25) show intrasubunit changes to this distance.
C Close up of intrasubunit conformational change in the IT switch and αP17 loop providing flexibility to allow for intersubunit rigid body rotation. Distance shown are intrasubunit conformational change in the αV24Y T20S model and do not represent the global conformational changes, as the P17 packing against the neighboring Helix 0 is similar in open and closed states (SFig. 3a)

WT T20S   **αV24Y T20S**   **ZYA-T20S**

Sig. S5: V24Y mutation and ZYA binding both alter E25's H- bonding partner.
E25 hydrogen bonding is affected by hydrophobic group occupancy in the HbYX hydrophobic pocket as it is positioned at the top of this pocket.
A E25 is H-bonded to R28 in the WT T20S.
B Distance between E25 and R20 in the WT T20S.
C E25 H-bonds with R20 in the V24Y T20S.
D Distance between E25 and R20 in the V24Y T20S
E E25 H-bonds with R20 in ZYA-T20S structure.
F Distance between E25 and R20 in the ZYA-T20S.

Fig. S6: Comparison of HbYX hydrophobic pocket in V24Y T20S and ZYA-T20S structures
A Distances between the α-carbon of selected residues (shown in sticks) to be compared against **B** to demonstrate the similarities and minimal level of intrasbunit conformational changes between these structures.
B Same as **A** except the ZYA-T20S model (blue) was used.

A

4920 micrographs
1031789 particles

↓

2 rounds of 2D
Classification

↓

542722 particles

↓

Homogenous
refinement
D7

B

C

2.20Å

Fig. S7: αV24F T20S Processing Scheme
A Cryo-EM workflow for αV24F T20S. All steps were performed in Cryosparc.
B Final 2D class averages after 2 rounds of 2D Classification shown with a mixture of top and size views.
C Homogenous refinement of class of particles that showed the open gate conformation.

Fig. S8: αV24Y T20S Validation

A Micrograph representative showing αV24F T20S. Scale bar represents 5nm.
B Standard FSC-0.143 graph showing corrected 2.20Å resolution.
C View of particle angle distribution showing a mixture of side and top view particles.
D Particles from selected ab-initio model were refined with homogenous refinement and D7 symmetry applied (D7 symmetry was appropriately confirmed as discussed in SFig. 2d). Final 2.20Å D7 map was colored in Chimera to show range of resolution of 3D reconstructed map.

Fig. S9: αV24Y T20S Validation

A Alignment showing sequences in the first 30 positions of the α subunits of T20S and H20S (listed according to gene names). Alignment was performed using ClustalΩ. Conversation and figure was generated using Jalview 2.11.2.5. Alignments are colored using the Clustal color scheme and color intensity indicates conservation (darkest: highly conserved). The bar graph in yellow indicates the conservation score, accounting for size, charge, hydrophobicity, and polarity. Red arrow points to αL21 and the corresponding residues in the alignment.

B Same as **A** except the alignment shows sequences in the positions 145 to 176 of the α subunits of T20S and H20S (listed according to gene names). The red arrow points to αA154 and the corresponding residues in the alignment.