Comparative metagenomics of tropical reef fishes show conserved core gut functions across

hosts and diets with diet-related functional gene enrichments

Derek G. Wu[1], Cassandra R. Harris[1], Katie M. Kalis[1], Malique Bowen[1], Jennifer F. Biddle[1],

Ibrahim F. Farag[1]*

[1]School of Marine Science and Policy, University of Delaware, Lewes DE 19958 USA

*Corresponding author: faragif@udel.edu

Running title: Fish gut metagenome

Abstract

Fish gut microbial communities are important for the breakdown and energy harvesting

of the host diet. Microbes within the fish gut are selected by environmental and evolutionary

factors. To understand how fish gut microbial communities are shaped by diet, three tropical fish

species (hawkfish, *Paracirrhites arcatus*; yellow tang, *Zebrasoma flavescens*; and triggerfish,

Rhinecanthus aculeatus) were fed piscivorous (fish meal pellets), herbivorous (seaweed), and

invertivorous (shrimp) diets, respectively. From fecal samples, a total of 43 metagenome

assembled genomes (MAGs) were recovered from all fish diet treatments. Each host-diet

treatment harbored distinct microbial communities based on taxonomy, with *Proteobacteria*,

Bacteroidota, and *Firmicutes* being the most represented. Based on their metagenomes, MAGs

from all three host-diet treatments demonstrated a baseline ability to degrade proteinaceous, fatty

acid, and simple carbohydrate inputs and carry out central carbon metabolism, lactate and

formate fermentation, acetogenesis, nitrate respiration, and B vitamin synthesis. The herbivorous

yellow tang harbored more functionally diverse MAGs with some complex polysaccharide

degradation specialists, while the piscivorous hawkfish's MAGs were more specialized for the

degradation of proteins. The invertivorous triggerfish's gut MAGs lacked many carbohydrate

degrading capabilities, resulting in them being more specialized and functionally uniform.

Across all treatments, several MAGs were able to participate in only individual steps of the

degradation of complex polysaccharides, suggestive of microbial community networks that

degrade complex inputs.

Importance

The benefits of healthy microbiomes for vertebrate hosts include the breakdown of food

into more readily usable forms and production of essential vitamins from their host's diet.

Compositions of microbial communities in the guts of fish in response to diet have been studied,

but lack a comprehensive understanding of the genome-based metabolic capabilities of how they

support their hosts. Therefore, we assembled genomes of several gut microbes collected from the

feces of three fish species that were being fed different diets to illustrate how individual

microbes can carry out specific steps in the degradation and energy utilization of various food

inputs and support their host. Herbivorous fish harbored a functionally diverse microbes with

plant matter degraders, while the piscivorous and invertivorous fish had microbes that were more

specialized in protein degradation.

Introduction

There are many factors that affect the composition of gut microbes in fish, including the

microbial composition of the surrounding water, diet, and environmental conditions such as

salinity, temperature, and pH (1, 2). Additionally, gut microbiomes are dynamic, where fish

hatch with few microbes in their gut as larvae (3), which increase in quantity and diversity as

they take in water and food from the surrounding environment (1). The composition of the

aquatic microbial environment has been shown to correlate with the gut microbiomes of the

fishes in those aquatic environments (4, 5, 6), while other studies have shown a divergence

between gut and environmental microbial communities (7). Gut microbiomes in individual fishes

in the same environment often possess distinct microbial communities (5), and these

communities can remain relatively stable over time in the absence of disturbances (8).

In marine fish, *Proteobacteria*, *Fusobacteria*, *Firmicutes*, *Bacteroidetes*, *Actinobacteria*

and *Verrucomicrobia* are among the most commonly identified phyla in intestinal samples (9).

At lower taxonomic scales, *Vibrio*, *Pseudomonas*, *Achromobacter*, *Corynebacterium*,

Alteromonas, *Flavobacterium*, and *Micrococcus* are considered to be the predominant gut

colonizers (1). Although the existence of baseline similarities in core gut microbial communities

among congeneric fish hosts has been suggested (10, 11, 12, 13), there are also studies that show

that microbiomes between congeneric hosts can diverge based on differences in environment and

trophic level (14). For example, in a study of migratory rabbitfish, less than half of the gut

microbial community remained consistent across the fish's migratory path (15). There are also

reported convergences of gut microbial communities between surgeonfish and nonsurgeonfish,

suggesting that core microbiomes across fish species may exist (16, 17).

Diet is an important factor that affects the gut microbiome. It has been shown that the

midgut microbial community of convict surgeonfish reflects the microbial community of the

algae that it was eating, suggesting a direct gut microbiome seeding related to diet (18). Yet,

most diet-gut microbiome connections are more indirect. For example, larval diet

supplementation provided to gilthead sea bream, European sea bass, and rainbow trout shifted

their gut microbial communities in different ways, varying by host species (19). Similarly,

changing the diet of sea bream from fishmeal to vegetables resulted in clear changes in microbial

community composition, but not overall community diversity (20).

Gut microbial communities also vary by feeding strategy and trophic level (6, 21). In the

microbial community composition of surgeonfish, herbivorous and algivorous diets are most

associated with distinct communities, while carnivorous and omnivorous diets tend to have

dynamic and less distinct compositions (16). Not only does diet affect the composition of

microbes found in the gut, but it also affects the metabolism that occurs in the gut. Carnivorous

diets are classified by the presence of pathways that relate to catabolism and biosynthesis of

amino acids (22). Herbivorous diets are classified by the pathways linked to pyruvate

metabolism and a high abundance of short chain fatty acids, which are indicators of fermentation

happening in the gut (15, 23). Because herbivores do not intake an abundance of protein in their

diet, they go through the process of fermentation to produce amino acids (22). To date,

invertivorous diets have been less studied for gut microbiome functions.

Several studies have shown how individual gut microbes play putative roles in the

degradation of key diet inputs (24). For example, in guts from carp fed grass diets, hundreds of

bacterial isolates have been cultured that demonstrate cellulolytic activity (25). A study of three

herbivorous fishes found that the activity of amylase enzymes in the guts positively correlated

with diets that were higher in rhodophytes, while laminarases were negatively correlated with

phaephytic diets (26). Using a genomic content estimator, PICRUSt, based on 16S rRNA

amplicon libraries, and enzymatic tests, herbivorous diets were associated with fish gut microbes

that produced more cellulases, while carnivorous diets were correlated with gut microbes that

produced more trypsins (27). In rabbitfish, seaweed diets resulted in diverse gut microbial

communities that were able to produce nonstarch polysaccharide degrading enzymes (28).

However, these patterns likely vary by host species and diet, considering that there are

alternative studies that have found little functional ability of microbial communities to degrade

diet inputs (29).

The majority of studies on fish gut microbiomes have utilized only small subunit

ribosomal gene amplicon sequencing for analysis, and they have clearly shown that microbiomes

vary based on diet (14, 15, 19, 20, 30, 31). Few studies have examined in detail the metabolic

pathways that exist across the gut microbiota of different fish species, which metagenomic

analyses can provide unique insights into (32, 33). As such, we examined the gut inhabitants

from multiple coral reef fish to probe at putative microbial functions by maintaining these wild

caught fish in aquaria while controlling their diets. We examine three fish species in this study:

yellow tang (*Zebrasoma flavescens*), humu humu Triggerfish (*Rhinecanthus aculeatus*), and arc-

eye Hawkfish (*Paracirrhites arcatus*). Yellow tangs are herbivores that naturally feed on algae

and marine plants; in the experiment they were fed seaweed, a natural herbivorous diet.

Triggerfish are invertivores that naturally feed on shrimp, clams and snails; for this experiment

these fish were fed mysis shrimp. Hawkfish are generalist predators feeding on small fishes and

invertebrates; they were fed fish pellets for this study. The overarching goal of this study was to

determine the extent to which fish gut MAGs corresponded to selection pressures from the host

diet using a comparative metagenomic approach. We hypothesized that the fish gut communities

would primarily be determined by the biochemical capabilities of recovered metagenomes to

degrade and harvest energy from the host's specific diet, thus the major energy harvesting

pathways exhibited by the MAGs would largely reflect the substrate inputs (i.e., fatty acid

oxidation for high fat diets and mixed acid fermentation pathways for high polysaccharide diets).

In this work, we show that the core potential metabolism of each fish gut microbiome is

redundant across diets and species of fish, with greater accessory metabolisms for the

herbivorous diet. We also show that the method by which one analyzes gut samples (i.e., 16S

rRNA amplicon sequencing, metagenome construction, binning) has a strong role in determining

which microbes are detected and included in downstream analyses, underscoring the importance

of diverse sequence methodology for the continued study of gut microbial communities.

Results

Diet Composition

The majority of both the fish meal pellet (56.1%) and shrimp (69.3%) diets was

composed of protein (Supplemental Table 1). The fish meal pellets had a total fat content of

16.6%, while the shrimp diet was 6.87% fat (Supplemental Table 1). Of the fats, both the fish

pellets and shrimp had a broad composition of fatty acids. From the fish pellet diet, the largest

contributions of fats were C22:6n3 docosahexaenoic, C20:1n9 cis eicosenoic, and C16:1n7

palmitoleic acids (Supplemental Table 2). From the shrimp diet, the largest contributions of fats

were C16:0 palmitic, C22:6n3 docosahexaenoic, and C20:5n3 eicosapentaenoic acids

(Supplemental Table 2). Both diets had a small component of fiber (2.42% and 3.57% for fish

pellets and shrimp, respectively) (Supplemental Table 1). Both diets could provide gut microbes

with biologically relevant minerals, such as potassium, calcium, phosphorus, and metals like

iron, magnesium, zinc, manganese, and copper (Supplemental Table 1). The seaweed, fed as nori

138 paper, contained almost equivalent protein and carbohydrates (5.7% and 5%, respectively), and

139 minor components of fiber and sugar (both 0.4%). Seaweed had large amounts of iodine,

140 vitamins C, A and B12 and also contained calcium, potassium, iron and magnesium.

141 *Initial 16S rRNA gene amplicon and metagenome communities*

142  We first examined the fish comparatively using 16S rRNA gene amplicon sequencing,

143 which identifies only the taxonomic representation of the sample. With this analysis, the

144 hawkfish gut community was least diverse, and all samples showed an abundance of

145 *Enterobacteriales* within the *Gammaproteobacteria* (Supplemental Figure 1). We then

146 proceeded with metagenomic analysis of a later timed sample. Using coverage of a single copy

147 ribosomal gene, *rps3*, as our taxonomic and abundance indicator, we saw that the microbial

148 composition varies, with a high representation of *Vibrionales* in each sample that was not

149 initially observed by amplicons (Figure 1). The greatest diversity of taxa was within the tang

150 sample, which was also not observed in amplicon data (Figure 1).

151 *Metagenome-Assembled Genome (MAG) Recovery*

152  After assembly and binning (Table 1), we recovered MAGs, quality checked and

153 identified them (Table 2). A total of 38 MAGs were recovered from the piscivorous hawkfish

154 treatment, and 18 passed our quality threshold with an average genome completeness and

155 contamination of 91.44% and 4.47%, respectively (Table 2). These 18 MAGs contained 4

156 taxonomic classes: *Alphaproteobacteria* (8), *Bacteroidia* (3), *Gammaproteobacteria* (6), and

157 *Gracilibacteria* (1) (Table 2). The MAGs recovered were, for the most part, not reflective of the

158 potential diversity of each sample (Figure 1), as the Vibrionales in each sample were likely too

159 diverse to allow the MAG clustering. However, *Bacteroidales* were recovered in the tang sample

160 and were also abundant in the unassembled metagenome (Figure 1) and the amplicon data

(Supplementary Figure 1). *Enterobacterales*, which was abundant in the amplicon data and

MAG data for the triggerfish, were not seen in the unassembled metagenome (Supplementary

Figure 1; Figure 1). These comparisons show how the microbial community can seem to shift

based on the analysis method used and highlights how studies of fish guts are best done using

multiple methods of analysis. Since our MAG analysis is not comprehensive of the total

microbial community, we take each analysis as a first-pass in the comprehensive metabolic

analysis of fish guts.

Our MAG analysis does highlight the ability to see microbial groups that may be

overlooked due to primer bias in 16S rRNA amplicon studies or in a representation that allows

them to be overlooked in an assembled metagenome. Additionally, we are able to determine how

similar these MAG genomes are to known genomes. Of these MAGs, 52% are within the

similarity metrics to be considered the same as previously discovered species (average nucleotide

identity; ANI >95), leaving the remainder to be potentially new species or genera (Table 2). In

particular, the *Gracilibacteria* MAG is within the *Patescibacteria* phylum at only 76 ANI to its

neighbors. While this phylum has been seen in numerous fish guts, this MAG can only be

classified to the family level.

From the herbivorous yellow tang treatment, 18/62 MAGs were also recovered and

passed quality checks, with an average genome completeness and contamination of 94.19% and

3.63%, respectively (Table 2). In contrast to the piscivorous gut community, the herbivorous

community appeared to have more taxonomic diversity at higher taxonomic ranks (e.g., phylum

or order levels). The 18 MAGs represented 7 taxonomic classes: *Alphaproteobacteria* (1), *Bacilli*

(1), *Bacteroidia* (5), *Clostridia* (3), *Gammaproteobacteria* (6), *Gracilibacteria* (1), and

Spirochaetia (1) (Table 2). The phylogenetic novelty of this sample was also higher, with only

23% of MAGs being classified within a known species, and the majority of MAGs were only

able to be classified to the family level. Some of the most novel MAGs are within the

Patescibacteria, Firmicutes and *Bacteroidota,* based on MAG ANI values (Table 2).

Despite a similar sized dataset and what appears to be a similar assembly (Table 1), only

7/33 MAGs passed our quality threshold and were recovered from the invertivorous triggerfish

treatment, with an average genome completeness and contamination of 90.0% and 3.3%,

respectively (Table 2). The 7 MAGs represented 5 taxonomic classes: *Alphaproteobacteria* (2),

Bacilli (1), *Bacteriovoracia* (1), *Campylobacter* (1), and *Gammaproteobacteria* (2) (Table 2).

Only one of these MAGs was within a known species, with the majority of MAGs only being

classified to the family level and one classified only to class (MAG Tr3LTN.016; *Bacillus;* Table

2). The taxonomic novelty of the MAGs within this project highlights the need for genome-level

understanding of fish gut communities, beyond the small subunit ribosomal gene. Further

description of novel MAGs was not done for this study, as we focused on function across diet

types.

Diet-Based Analyses

With the MAGs recovered, we analyzed each sample and its genomes for a response to

diet types by performing principal component analysis on the copy numbers of enzymes within

each MAG, normalizing by genome completeness. These data suggest that fish host diet may be

a major driving selection factor for MAG composition (PERMANOVA; $F = 6.3078$; $p = 0.003$)

(Figure 2). Hemicellulose, cellulose, chitin, and starch were forces loaded towards the

herbivorous diet whereas fatty acids and proteins were loaded towards the piscivorous and

invertivorous diets (Figure 2). The herbivorous gut community displayed a greater variance in

gene copies, while the invertivore gut community had the narrowest variation in substrate

degradation gene compositions (homogeneity of dispersion; $F = 9.3049$; $p = 0.001$) (Figure 2).

Although several of the MAGs collected from the herbivorous yellow tang demonstrated

specialization for the degradation of complex polysaccharides, it also contained MAGs that were

more specialized in protein and fatty acid degradation. In contrast, the invertivorous triggerfish

MAGs had a narrower range of substrate degradation functions, suggestive of a more specialized

community that is adapted to the high protein shrimp diet.

Carbohydrates

Overall genes encoding for carbohydrate degrading enzymes are more enriched in the

MAGs recovered from herbivorous fish compared to fishes following other diets. For example,

the degradation of cellulose could be completed in both the herbivorous and piscivorous diets

(Figure 3a). However, MAGs recovered from the herbivorous yellow tang had significantly more

gene copies for enzymes involved in the cellulose degradation pathway than both piscivorous

and invertivorous hosts (Kruskal Wallis, $p = 0.01$) (Figure 4a). Within the piscivorous hawkfish

gut, 7/18 MAGs [005, 009, 011, 022, 028, 030, and 033] could facilitate the hydrolysis of

cellulose to glucose monomers (Figure 3a). Of these 7 MAGs, 4 are *Alphaproteobacteria,* 2 are

Bacteroidia, and 1 is a *Gammaproteobacteria*. From the herbivorous diet, 9/18 MAGs [001, 002,

005, 006, 007, 009, 014.3, 015, and 016] could catalyze these reactions. Of these 9 MAGs, 3 are

Gammaproteobacteria, 3 are *Bacteroidia*, 2 are *Clostridia*, and 1 is *Gracilibacteria*. In contrast,

no MAGs recovered from the invertivorous triggerfish could degrade cellulose to simple

monomers (Figure 3a). Additionally, in all treatments, there were several MAGs that could

potentially catalyze at least one step of the larger cellulose degradation pathway but not the

entire process, hereby referred to as "partial degraders" (Figure 3a). From the piscivorous

treatment, MAGs 001, 010, and 015 were partial degraders, while from the herbivorous

treatment, MAGs 003, 011, 012, and 017 were partial degraders of cellulose. The existence of

these MAGs suggests that they may be opportunists that require other MAGs to initiate the

breakdown of cellulose into simpler substrates for them to use.

The MAGs recovered from the herbivorous yellow tang had significantly higher gene

copies of enzymes involved in hemicellulosic degradation than both the piscivorous and

invertivorous treatments (Kruskal Wallis, p = 0.0074) (Figure 4b). From the herbivorous

treatment, 6/18 MAGs [001, 005, 006, 007, 011, and 015] could completely degrade at least one

of the three screened hemicelluloses (Figure 3b-d). Of these 6 MAGs, 2 are *Clostridia*, 3 are

Bacteroidia, and 1 is a *Gammaproteobacteria*. From the piscivorous treatment, only 2/18 [022

and 030] were able to fully degrade at least one hemicellulosic compound (Figure 3b-d). No

MAGs recovered from the invertivorous treatment could completely degrade hemicellulose

(Figure 3b-d). However, in all treatments, there were several MAGs that were partial degraders

of hemicellulose backbone or one of the side chains (Figure 3b-d). From the herbivorous

treatment, all MAGs except the complete degraders could be classified as partial degraders of

various hemicelluloses, wherein they were capable of performing one or more steps in a

hemicellulose degradation pathway. Similarly, all the MAGs recovered from the piscivorous diet

could catalyze at least one step of a hemicellulose degradation pathway; however, there was a

lack of capacity to hydrolyze and ferment many of the auxiliary sugars from the hemicellulose

compounds.

The herbivorous MAGs had significantly greater numbers of genes involved in the

degradation of chitin than both the invertivorous and piscivorous samples: 2.2 fold and 13.1 fold,

respectively (Kruskal Wallis, p = 0.0082) (Figure 4c). In fact, the invertivorous MAGs showed

an inability to catalyze the initial steps to either of the chitin degradation pathways (Figure 3e).

Meanwhile, 2/18 MAGs [009 and 030] from the piscivorous treatment and 5/18 MAGs [001,

003, 005, 006, and 015] from the herbivorous treatment could fully catalyze the breakdown of

chitin to simpler glucosamine units. The 2 MAGs from the piscivorous treatment that can

degrade chitin are both in the *Cytophagales* order within the *Bacteroidia* class. Of the 5 chitin-

degrading MAGs from the herbivorous treatment, 2 are *Bacteroidia*, 2 are *Clostridia*, and 1 is a

Gammaproteobacteria.

All three samples had MAGs that encoded amylases for the degradation of starches.

However, the herbivorous gut sample had significantly more amylase gene copies per MAG than

the piscivorous sample (Kruskal Wallis, p = 0.05) (Figure 4d).

Proteins

All three hosts contained MAGs that produced exopeptidases for protein degradation,

allowing for basal levels of protein breakdown in the gut environment. The piscivorous gut

MAGs had 1.69 fold more exopeptidase gene copies per MAG than those of the herbivorous gut

community (Figure 4e), suggesting that protein degradation pathways were enriched in response

to the high protein fish pellet diet. From the piscivorous community, 16/18 MAGs had more than

10 exopeptidase genes, while 11/16 and 3/7 MAGs from the herbivorous and invertivorous guts

had more than 10 exopeptidase genes, respectively (Table 3). Additionally, MAG 005 from the

piscivorous gut (*Phaeomarinobacter* sp.) was the only MAG recovered across the fish treatments

to have a gene for collagenase, which could highlight a specialized niche for collagen

degradation in response to the fish pellet diet.

Fatty Acids

All three hosts had MAGs that could carry out complete beta oxidation of fatty acids. The

screen for fatty acid breakdown via beta oxidation included genes for acyl-CoA dehydrogenase,

enoyl-CoA hydratase, 3-hydroxyacyl-CoA dehydrogenase, and acetyl-CoA acyltransferase. In

the piscivorous gut community, 9/18 MAGs [003, 006, 007, 009, 012, 015, 018, 022, 030] were

able to carry out fatty acid oxidation to completion to produce acetyl CoA (Figure 5a). These

MAGs represent the *Gammaproteobacteria* (2), *Alphaproteobacteria* (3), and *Bacteroidia* (3)

classes. In contrast, only 4/18 MAGs [003, 004, 012, and 016] from the herbivorous community

and 3/7 MAGs [002, 005, and 006] from the invertivorous community could independently

complete all of fatty acid oxidation (Figure 5b,c). All four MAGs from the herbivorous yellow

tang microbiome that could complete beta oxidation were *Gammaproteobacteria*, and 3 of them

were in the *Enterobacterales* order. These data highlight a potential enrichment of fatty acid

oxidation capacities in the piscivorous hawkfish treatment, whose fish pellets contained high

amounts of various fatty acids (Supplemental Table 1).

Energy Harvesting and Fermentation

MAGs recovered from all three samples were analyzed for their metabolic capacities.

13/18, 14/18 and 5/7 MAGs from the piscivorous, herbivorous, and invertivorous treatments,

respectively, could carry all of glycolysis from glucose to pyruvate (Figure 5). Following

glycolysis, 17/18, 17/18, and 7/7 MAGs from the piscivorous, herbivorous, and invertivorous gut

communities could carry out pyruvate oxidation, bridging glycolysis to the citric acid cycle

(Figure 5). 13/18, 6/18, and 4/7 MAGs from the piscivorous, herbivorous, and invertivorous

guts, respectively, possessed more than 80% of the genes for the entire TCA cycle (Figure 5).

Within each treatment, there are some MAGs that can complete individual energy-harvesting

steps within the TCA cycle, but not the complete process, which may be a product of genome

incompleteness (Table 3). Since the piscivorous hawkfish sample contained a higher proportion

of MAGs that could carry out the TCA cycle, particularly the second oxidation reactions, these

data could suggest that these MAGs are more predominantly metabolizing the amino acids from

their proteinaceous diet via central carbon metabolism.

The central carbon pathways observed in these MAGs produce reduced electron carriers

that need to be oxidized in order to sustain the catabolic pathways. From the piscivorous

treatment, 4/18 MAGs [001, 010, 012, and 030] could perform dissimilatory nitrate reduction to

ammonia through a nitrite intermediate as a mechanism for anaerobic respiration (Figure 5a).

5/18 MAGs [003, 004, 007, 012, and 016] from the herbivorous treatment are dissimilatory

nitrate respirers, while 2/7 MAGs [002 and 008] from the invertivorous treatment can carry out

nitrate respiration (Figure 5b-c). Lactate fermentation was rare among all three treatments, with

only 2/18, 1/18, and 0/7 MAGs from the piscivorous, herbivorous, and invertivorous diets,

respectively, being able to produce a L-lactate dehydrogenase (Figure 5).

All three gut communities demonstrated an ability to perform the latter half of the

acetogenesis pathway to reduce acetyl-CoA to acetate. From the piscivorous treatment, 6/18

MAGs had genes for both phosphate acetyltransferase and acetate kinase, synthesizing ATP

Figure 5a; Supplemental Figure 2). Similarly, 11/18 MAGs from the herbivorous diet and 4/7

MAGs from the invertivorous diet could produce these enzymes (Figure 5b-c; Supplemental

Figures 3-4). Using the acetate products, 5/18, 7/18, and 3/7 MAGs from the piscivorous,

herbivorous, and invertivorous communities, respectively, could produce alcohol dehydrogenase

homologs for the production of ethanol and consumption of reduced NADH (Figure 5;

Supplemental Figures 2-4).

Alternatively, all three treatments had MAGs that had genes for formate acetyltransferase

and formate dehydrogenase, which are involved in the fermentation of formate to carbon dioxide

and hydrogen gas. From the piscivorous treatment, 3/18 MAGs could produce formate

acetyltransferase while 8/18 MAGs could produce formate dehydrogenase (Figure 5a;

Supplemental Figure 2). In contrast, 11/18 MAGs from the herbivorous diet could produce

formate acetyltransferase, while only 3/18 MAGs could produce formate dehydrogenase (Figure

5b; Supplemental Figure 3). From the invertivorous diet, 4/7 MAGs possessed formate

acetyltransferase genes, while 3/7 MAGs possessed genes for formate dehydrogenase (Figure 5c;

Supplemental Figure 4). None of the MAGs in this study could perform dissimilatory sulfate

respiration or methanogenesis.

Vitamin and Hydrogenase Production

All communities had MAGs that could produce vitamins B1, B2, B7, B9, and B12.

Thiamine (B1) could be produced by 7/18, 10/18, and 3/7 MAGs from the piscivorous,

herbivorous, and invertivorous communities, respectively (Table 3). Riboflavin (B2) could be

synthesized by 4/18, 6/18, and 2/7 MAGs from the piscivorous, herbivorous, and invertivorous

hosts guts, respectively (Table 3). Biotin (B7) could be synthesized by 12/18, 10/18, and 3/7

MAGs from the piscivorous, herbivorous, and invertivorous diet communities, respectively,

using at least one of the various biotin biosynthetic pathways (Table 3). Hydrotetrafolate, a

precursor to folic acid (B9), could be produced by 14/18, 9/18, and 3/7 MAGs from the

piscivorous, herbivorous, and invertivorous hosts guts, respectively (Table 3). Cobalamin (B12)

could be made by 7/18, 4/18, and 2/7 MAGs from the piscivorous, herbivorous, and

invertivorous hosts guts, respectively (Table 3).

From the piscivorous gut community, the only MAG capable of producing a component

of a nickel-iron hydrogenase is 001 (*Photobacterium aphoticum*), being able to produce a

periplasmic Group 1 hydrogenase, which is generally responsible for catalyzing steps in

anaerobic respiration using metals, nitrates, or sulfates as terminal electron acceptors

(Supplemental Figure 5). Similarly only 1 MAG [016] from the invertivorous treatment produces

a group A1 and A3 iron-iron hydrogenase, which produces fermentative hydrogen gas

(Supplemental Figure 5). Also from the invertivorous diet, 2/7 MAGs [002 and 008] from this

diet produce group 1 nickel-iron hydrogenases., which also act as respiratory electron receptors

(Supplemental Figure 5).

In contrast, MAGs from the herbivorous treatment synthesize various iron and nickel

hydrogenases. 6/18 MAGs [005, 010, 015, 017, 023, and 006] produce group A1 and A3 iron-

iron hydrogenases (Supplemental Figure 5). 7/18 MAGs [001, 005, 010, 015, 017, 023, and 006]

produce group B iron-iron hydrogenases involved in hydrogen gas fermentation (Supplemental

Figure 5). 4/18 MAGs [010, 015, 017, and 006] produce group C1 iron-iron hydrogenases

(Supplemental Figure 5). Only 2/18 MAGs [017 and 006] produce group C2 iron-iron

hydrogenases, which are transcriptional regulators (Supplemental Figure 4). 4/18 MAGs [010,

015, 023, and 006] produce group C3 iron-iron hydrogenases (Supplemental Figure 5). 3/18

MAGs [001, 004, and 005] produce group 1 nickel-iron hydrogenases, also transcriptional

regulators (Supplemental Figure 5). Within the herbivorous yellow tang gut, MAGs 006, 010,

015, and 017 produce the greatest variety of hydrogenases, and 3 of them belong to the

Clostridia class within *Firmicutes*.

Discussion

The present study used metagenome-assembled genomes recovered from the feces of

tropical reef fishes that were collected from the wild and housed in aquaria under controlled

diets. We found evidence of core functions in the examined microbiomes. Our study benefited

from the aquarium-based approach, as we had defined food sources that the fish were fed for

weeks in captivity, compared to the uncontrolled diet that may be found in nature. Core functions

include protein degradation, fatty acid oxidation, central carbon metabolism, nitrate respiration,

acetogenesis, and formate oxidation. Considering our sampling design and dataset limits,

additional functions may be possible in fish guts and more taxa should be closely examined;

however, our results show that previous work based on small subunit ribosomal sequences or the

interpretation of metabolism based solely on taxonomic data is a limited view of a fish gut

microbiome.

Despite the number of shared functions, diet types also showed selective trends. The

microbial MAGs associated with the piscivorous and invertivorous diets were the most

functionally similar, as expected, considering the compositions of the fish pellets and shrimp

diets, consistent with the model of diet-based microbiomes (34). Both diets were high in protein

and fat content, and the MAGs recovered from these diets showed high gene copy numbers for

exopeptidases and the abilities to carry out beta oxidation of fatty acids. The piscivorous gut

community, although specialized for the degradation of proteins and fatty acids, could also

degrade plant matter. It could be that plant material is indirectly ingested during fish

consumption. In contrast, the invertivorous triggerfish is primarily digesting shrimp, which likely

do not contain substantial amounts of complex plant polysaccharides in their intestines.

Therefore, the triggerfish MAGs encode for digesting proteins and fats with some versatility for

simple sugars. As a caveat, examination of additional MAGs after additional sequencing effort

could show a broader digestion ability for the triggerfish.

As expected, the herbivorous yellow tang gut MAGs demonstrated a strong ability to

degrade complex plant polysaccharides, likely due to their nori seaweed diet. Unexpectedly, the

MAGs recovered from the invertivorous triggerfish diet demonstrated an inability to degrade

chitinous compounds. This contrasts with previous results which found that rainbow trout being

fed chitinaceous insect exuviae had gut microbiomes enriched in chitinolytic bacterial genera;

however, these results were based on predictive metagenomes based solely on 16S rRNA gene

amplicons (35). Exochitinase production has been observed in various fish species being fed

insect diets, although the source of these enzymes (host versus gut microbiome) was not

specified (36, 37).

With the higher fatty acid diets of the piscivorous hawkfish and the invertivorous

triggerfish, their MAGs carried a strong signature of beta oxidation. The beta oxidation of fatty

acids is a reductive cycle that rapidly degrades long chain fatty acids to directly produce ATP

and acetyl-CoA. This high energy-yielding process explains its abundance in MAGs recovered

from these treatments, while the herbivorous yellow tang gut microbial community had fewer

MAGs capable of fatty acid degradation. Instead of relying on this high yield process, our data

suggest that the herbivorous gut community relies more on mixed acid fermentation and acetate-

producing pathways for the production of ATP and recycling of reduced electron carriers. This

establishes a major difference in the proteinaceous and fatty acid diet economy expected in the

piscivorous and invertivorous treatments and the polysaccharide diet economy expected in the

herbivorous treatment.

Further supporting the polysaccharide economy expected in the herbivorous yellow tang

gut, the MAGs collected from this treatment possessed a deep redundancy and breadth of

hydrogenases. These hydrogenases varied in function, including being final electron acceptors in

anaerobic respiration and hydrogen gas production at the end of fermentation. Since the

herbivorous treatment is associated with higher concentrations of fibrous substrates, the

constituent sugars would rely on electron mediated processes to degrade the substrates and

recycle cofactors. The higher proportion of MAGs that produce hydrogenases supports this

expectation, with these hydrogenases playing a putative role in balancing electron budgets and

gradients to support respiration and fermentation pathways.

A limitation of this study that is worth noting is the possibility for there to be host-

specific factors such as gut physiology and chemistry that could also be selecting for taxon and

function-specific MAGs. While the present study found substantial evidence for the linkage

between diet inputs and community composition, it has been shown that host species may harbor

distinct microbial communities (7). Future research should study model fish species that have

adaptable diets or whose diets change with development (38). Then, diets could be shifted and

changes in the microbial community composition could be followed as a function of time and

treatment. The methodology used here shows that fish under treatment can have their

microbiome analyzed while remaining viable for longer treatments of study. If these diet changes

are considered in the context of ecological disturbances, these studies would be indicative of the

likelihood of adjusting to rapidly changing environments, which is important to understand given

the cascade of environmental changes being mediated by anthropogenic activity.

We recognize that keeping fish viable for this study meant that internal shifts of microbes

along gut structures were unable to be observed (5, 39, 40). As such, these data serve as a

preliminary basis for further studies whose experimental design could involve more intensive

sampling. However, our data does show strongly that single-gene taxonomic analyses fall short

of determining differences in fish gut microbiomes, and many lineages may be more or less

represented based on the method of molecular analysis. The ability to sample the full

metagenome and build MAG-based models of metabolism shows that fish guts are more similar

in functional potential than expected. In order to get insight into the functions of these MAGs

within these microbiomes, a full study should be done that addresses biomass abundance within

the gut, potentially by transcriptomics or proteomics, which can provide knowledge of how these

microbes are behaving and regulating the expression of their genomes.

Materials and Methods

Fish Rearing, Diets and Sample Collection

The yellow tangs, triggerfish, and hawkfish used for this study were wild caught and

obtained from Live Aquaria ® (Rhinelander, WI) and cared for according to IACUC approval

through the University of Delaware (A1319). Fish were held in 37L aquaria on a large

recirculating system at the UD Lewes Campus (Lewes, DE). Hawkfish (n=4) were kept

individually in aquarium tanks due to aggressive behavior. Yellow tangs (n=5) and triggerfish

(n=5) were kept in communal aquarium tanks with their same species. Hawkfish are piscivorous

fish and were fed commercial fish meal pellets (Sustainable Aquatics Dry Hatchery Diet,

Jefferson City TN). Yellow tangs are herbivorous fish and were fed commercially purchased nori

seaweed (sushi wrapping, various vendors). Triggerfish are invertivores and were fed

commercially purchased mysis shrimp (San Francisco Bay Brand Frozen Mysis Shrimp). We

opted to collect gut microbiomes via fecal collection to preserve the fish for other experiments

(41).

When collecting feces, which occurred multiple times due to the fish not being sacrificed,

fish were placed in individual 2.5L aquaria. To minimize a seawater microbial signal, artificial

seawater was created with Milliq-filtered water and Instant Ocean (Instant Ocean; Blacksburg,

VA) to 33 ppm salinity. Fish were moved to these tanks 10 min after feeding and remained there

overnight with air stones and no shared water flow to ensure that water quality remained high.

After 12-15 hours, the fish were placed back in their 37L tanks. From each small tank, 1500 mL

of water was sampled. The tank waters were filtered through an 8.0 micron cellulose filter

(Merck Millipore; Burlington, MA). This filter size was selected to ensure fecal particles were

preferentially collected instead of free-living bacteria in the water. The filters were labeled and

placed in a 60mm x 15mm petri plate with a stackable lid and stored in a -80C freezer until used

for DNA extractions.

Chemical Analysis of Food Sources

To determine major dietary differences between the fish-meal pellets and mysis shrimp,

20g of crushed, freeze dried mysis shrimp and 20g of fish-meal pellets were shipped to NP

Analytics Laboratory (St. Louis, MO) for testing. Nutrition information for nori seaweed was

retrieved from the commercial packaging.

DNA Extraction, Amplicon and Metagenomic Sequencing

DNA was extracted from the feces samples collected on the filter using the DNeasy

PowerSoil Kit (Qiagen, Valencia CA). A quarter of the filter paper was used for the extractions.

After each extraction, the amount of DNA in each sample was measured using the Qubit dsDNA

HS assay kit with the Qubit fluorometer (Invitrogen; Waltham, MA). The DNA was stored in a -

20C freezer. Samples with the highest DNA yield were selected from each species and sent to

the University of Connecticut MARS facility for amplification and sequencing of the 16S rRNA

gene (primers 515F, 806R) (42) using the Illumina MiSeq platform via their standard protocol.

After 1 month of daily feeding, additional samples were taken in the same manner, processed the

same way and sent to the University of Delaware DNA Sequencing and Genotyping Center at

the Delaware Biotechnology Institute (Newark, DE) for library creation and sequencing on an

Illumina HiSeq 2500.

Data Accessibility Statement

Raw metagenome data associated with these samples have been deposited in the NCBI

database under BioProject 1112800. Amplicon sequences are under BioProject

PRJNA1174778. Individual metagenome assembled genomes (MAGs) and annotation files are

available at https://figshare.com/projects/FishGutMAGs_AEM2024/223746.

Amplicon Sequence Quality Control and Processing

Raw forward and reverse amplicon sequences were paired and processed using 16S

rRNA analysis pipeline in MOTHUR version 1.46.1 (43). Paired sequences were trimmed,

ambiguous nucleotides were removed, and then operational taxonomic units (OTUs) with a 3%

dissimilarity were created. OTUs were then aligned and classified using the Silva138 database

version 138 (44).

Metagenome Quality Control, Assembly and Binning

Data processing, including reads quality checking and trimming, was performed using

Nesoni (45) under default parameters and applying a quality score of Q20. Qualified reads for

each dataset were assembled into contigs using IDBA-UD (46), metaSPAdes (47), and Megahit

v.1.1 (48) with default settings and k-mer lengths varying from 27 to 117. Community

composition was determined via annotation and retrieval of all 30S ribosomal protein (rps3)

sequences from assembled contigs. Reads were mapped to each rps3 gene using gbtools (49) and

BLAST (50) against UniProt (51) to identify the taxonomy of the gene. Community composition

was determined by the coverage of each gene compared to the total read coverage of all rps3

genes in the dataset. Contigs longer than 1000 bp were binned using binning tools MaxBin2.0

(52), Metabat (53), and Metabat2 (54). The resulting metagenome-assembled genomes (MAGs)

were quality assessed using CheckM (55) with the "lineage_wf" option. Only bacterial MAGs,

meeting criteria of $\geq 50\%$ completion and $\leq 20\%$ contamination, were retained. These MAGs

were further refined by removing outlier contigs based on t-SNE signatures, GC content,

taxonomic assignments, and differential coverages. Coverage calculations were performed by

mapping trimmed reads against the contigs using BBMap v.37.61 (56). MAGs with $< 10\%$

contamination were deemed acceptable, while those with 10-20% contamination were manually

cleaned again.

Phylogenetic Placement

The phylogenetic placements of the MAGs were estimated by creating a phylogenomic

tree from concatenated ribosomal proteins, aligned using Muscle v3.8 (57) and analyzed with

FastTree v2.1 (58). MAG identities were confirmed by calculating average nucleotide identity

(ANI) and average amino acid identity (AAI) with custom reference genomes closely related to

the screened lineages based on their single marker gene(s) closest matches. Taxonomic profiles

were verified using GTDB-Tk v2.1.0 (59).

Metabolic Reconstruction

Protein sequences from all MAGs were predicted using Prodigal v2.6.3 (60) using the

default translation table 11. The collective metabolic pathways for all MAGs present in each

depth were deduced via screening the predicted proteins against a custom hidden Markov model

database composed of 126 key metabolic genes covering most of the substrate utilization,

biosynthesis and energy metabolism related pathways (61). These pathways were further

assessed for completion through querying against the Kyoto Encyclopedia of Genes and

Genomes (KEGG) (https://www.genome.jp/kegg/) (62) database using the BlastKoala tool (63).

The pathway was deemed complete if the MAG encodes for at least 80% of the enzymes and

other involved proteins of a given pathway. We validated our results using METABOLIC-C

(64). METABOLIC-C integrates gene calling and annotation based on HMM-based databases,

motif validation, and annotation via specific databases.

Substrate Degradation Analysis

CAZyme sequences were identified via HMM searches against dbCAN V8 (65).

Enzymes involved in the degradation of cellulose, hemicelluloses (xylan, mannan, and

xyloglucan), and starch were identified as presented in Kumla et al. (66). Enzymes for the

degradation of chitin were identified as presented in De Tender et al. (67). Corresponding

CAZyme families for these enzymes were identified on the Carbohydrate-Active enZYmes

Database (http://www.cazy.org/) (68). Exopeptidases were identified using the MEROPS

Peptidase Database (https://www.ebi.ac.uk/merops/) (69). Enzymes involved in lactate, acetate,

and ethanol fermentation pathways were organized as described in the EcoCyc Database

(https://ecocyc.org/) (70). Formate fermentation enzymes were identified as described in Knappe

& Sawers (71). Hydrogenase enzyme sequences were identified through specific HMM searches

against the Kofam database V2024-01-01 (72). Hydrogenase functions were identified on the

HydDB database (https://services.birc.au.dk/hyddb/) (73).

Statistical Analysis

To visualize clustering patterns in substrate degradation enzymes in response to host

diets, a principal component analysis of all screened substrate pathways was performed on all

samples using the prcomp function in R Version 4.3.2 (74). The total number of gene copies for

all enzymes involved in the degradation of proteins, cellulose, hemicellulose, starch, and chitin

normalized to genome completeness and a binary system for the presence/absence of fatty acid

degradation were scaled and used for PCA processing. The loadings of the first two components

are reported, explaining 77.0% of detected variance. PCA data were visualized using the

fviz_pca_biplot function from the factoextra package (75). To test if the centroids of diet

treatments were different, a permutational multivariate analysis of variance (PERMANOVA)

was performed using the adonis2 function from the vegan package (76). To assess the

homogeneity of multivariate dispersions across diets, the betadisper and permutest functions

were used to compute the distances from group centroids to individual samples with 999

permutations.

To determine if MAGs recovered from each treatment differed in the number of substrate

degrading gene copies, the total gene copies for each degradation pathway normalized to MAG

genome completeness were totaled. Then, a Kruskal Wallis test was performed to determine if

there were significant differences among the three diets using the kruskal.test function. If

significant, a post-hoc Wilcoxen test was performed to identify significantly different pairwise

comparisons between diets.

Ethics Statement

The handling and care protocols of the fish used in this study were approved by the

University of Delaware Institutional Animal Care and Use Committee (A1319), AUP #1319-

2019-2.

References:

1. Nayak SK. 2010. Role of gastrointestinal microbiota in fish. *Aquac Res* 41:1553-1573.

2. Yukgehnaish K, Kumar P, Sivachandran P, Marimuthu K, Arshad A, Paray BA, Arockiaraj J.

2020. Gut microbiota metagenomics in aquaculture: Factors influencing gut microbiome

and its physiological role in fish. *Rev Aquacult* 12:1903–1927.

3. Reid HI, Treasurer JW, Adam B, Birkbeck TH. 2009. Analysis of bacterial populations in the

gut of developing cod larvae and identification of *Vibrio logei*, *Vibrio anguillarum* and

Vibrio splendidus as pathogens of cod larvae. *Aquaculture* 288:36–43.

4. Arockiaraj J, Kumaresan V, Chaurasia MK, Bhatt P, Palanisamy R, Pasupuleti M, Gnanam

AJ, Kasi M. 2014. Molecular characterization of a novel cathepsin B from striped murrel

Channa striatus: Bioinformatics analysis, gene expression, synthesis of peptide and

antimicrobial property. *Turk J Fish Aquat Sc* 14:379-389.

5. León-Zayas R, McCargar M, Drew JA, Biddle JF. 2020. Microbiomes of fish, sediment and

seagrass suggest connectivity of coral reef microbial populations. *PeerJ*, 8:e10026.

6. Pan B, Han X, Yu K, Sun H, Mu R, Lian CA. 2022. Geographical distance, host

evolutionary history and diet drive gut microbiome diversity of fish across the Yellow

River. *Molec Ecol* 32:1183-1196.

7. Li X, Yu Y, Feng W, Yan Q, Gong Y. 2012. Host species as a strong determinant of the

intestinal microbiota of fish larvae. *J Microbiol* 50:29–37.

8. Pisaniello A, Handley KM, White WL, Angert ER, Boey JS, Clements KD. 2023. Host

individual and gut location are more important in gut microbiota community composition

than temporal variation in the marine herbivorous fish *Kyphosus sydneyanus*. *BMC*

Microbiol 23:275.

9. Llewellyn MS, Boutin S, Hoseinifar SH, Derome N. 2014. Teleost microbiomes: The state of

the art in their characterization, manipulation and importance in aquaculture and

fisheries. *Front Microbiol* 5:207.

10. Shade A, Handelsman J. 2011. Beyond the Venn diagram: The Hunt for a core microbiome.

Environ Microbiol 14:4-12.

11. Givens C, Ransom B, Bano N, Hollibaugh J. 2015. Comparison of the gut microbiomes of 12

bony fish and 3 shark species. *Mar Ecol Prog Ser* 518:209–223.

12. Le D, Nguyen P, Nguyen D, Dierckens K, Boon N, Lacoere T, Kerckhof FM, De Vrieze J,

Vadstein O, Bossier P. 2019. Gut microbiota of migrating wild rabbit fish (*Siganus*

guttatus) larvae have low spatial and temporal variability. *Microb Ecol* 79:539–551.

13. Roeselers G, Mittge EK, Stephens WZ, Parichy DM, Cavanaugh CM, Guillemin K, Rawls

JF. 2011. Evidence for a core gut microbiota in the zebrafish. *ISME J* 5:1595–1608.

14. Sullam KE, Essinger SD, Lozupone CA, O'Connor MP, Rosen GL, Knight ROB, Kilham

SS, Russell JA. 2012. Environmental and ecological factors that shape the gut bacterial

communities of fish: a meta-analysis. *Molec Ecol* 21:3363-3378.

15. Jones J, DiBattista JD, Stat M, Bunce M, Boyce MC, Fairclough DV, Travers MJ, Huggett

MJ. 2018. The microbiome of the gastrointestinal tract of a range-shifting marine

herbivorous fish. *Front Microbiol* 9:2000.

16. Miyake S, Ngugi DK, Stingl U. 2014. Diet strongly influences the gut microbiota of

surgeonfishes. *Mol Ecol* 24:656–672.

17. Diwan AD, Harke SN, Panche AN. 2023. Host-microbiome interaction in fish and shellfish:

An overview. *Fish Shellfish Immunol Rep* 4:100091.

18. Parata L, Nielsen S, Xing X, Thomas T, Egan S, Vergés A. 2019. Age, gut location and diet

impact the gut microbiome of a tropical herbivorous surgeonfish. *FEMS Microbiol Ecol*

96:fiz179.

19. Antonopoulou E, Nikouli E, Piccolo G, Gasco L, Gai F, Chatzifotis S, Mente E, Kormas KA.

2019. Reshaping gut bacterial communities after dietary *Tenebrio molitor* larvae meal

supplementation in three fish species. *Aquaculture* 503:628–635.

20. Estruch G, Collado MC, Peñaranda DS, Tomás Vidal A, Jover Cerdá M, Pérez Martínez G,

Martinez-Llorens S. 2015. Impact of fishmeal replacement in diets for Gilthead Sea

Bream (*Sparus aurata*) on the gastrointestinal microbiota determined by pyrosequencing

the 16S rRNA gene. *PLoS ONE* 10e0136389.

21. Egerton S, Culloty S, Whooley J, Stanton C, Ross RP. 2018. The gut microbiota of marine

fish. *Front Microbiol* 9:873.

22. Sanders JG, Beichman AC, Roman J, Scott JJ, Emerson D, McCarthy JJ, Girguis PR. 2015.

Baleen whales host a unique gut microbiome with similarities to both carnivores and

herbivores. *Nat Commun* 6:8285.

23. Mountfort DO, Campbell J, Clements KD. 2002. Hindgut fermentation in three species of

marine herbivorous fish. *Appl Environ Microb* 68:1374–1380.

24. Ray AK, Ghosh K, Ringø E. 2012. Enzyme-producing bacteria isolated from fish gut: A

review. *Aquacult Nutr* 18:465–492.

25. Li H, Wu S, Wirth S, Hao Y, Wang W, Zou H, Li W, Wang G. 2014. Diversity and activity

of cellulolytic bacteria, isolated from the gut contents of grass carp (*Ctenopharyngodon*

idellus) (Valenciennes) fed on Sudan grass (*Sorghum sudanense*) or artificial feedstuffs.

Aquac Res 47:153–164.

26. Moran D, Clements KD. 2002. Diet and endogenous carbohydrases in the temperate marine

herbivorous fish *Kyphosus sydneyanus*. *J Fish Biol* 60:1190–1203.

27. Liu H, Guo X, Gooneratne R, Lai R, Zeng C, Zhan F, Wang W. 2016. The gut microbiome

and degradation enzyme activity of wild freshwater fishes influenced by their trophic

levels. *Sci Rep*, 6:24340.

28. Zhang X, Wu H, Li Z, Li Y, Wang S, Zhu D, Wen X, Li S. 2017. Effects of dietary

supplementation of *Ulva pertusa* and non-starch polysaccharide enzymes on gut

microbiota of *Siganus canaliculatus*. *J Oceanol Limnol* 36:438–449.

29. McCauley M, German DP, Lujan NK, Jackson CR. 2020. Gut microbiomes of sympatric

Amazonian wood-eating catfishes (Loricariidae) reflect host identity and little role in

wood digestion. *Ecol Evol* 10:7117–7128.

30. Liu Y, Li X, Li Y, Li J, Zhu S. 2022. Gut microbiomes of cyprinid fish exhibit host-species

symbiosis along gut trait and diet. *Front Microbiol* 13:936601.

31. Kashinskaya EN, Simonov EP, Kabilov MR, Izvekova GI, Andree KB, Solovyev MM. 2018.

Diet and other environmental factors shape the bacterial communities of fish gut in an

eutrophic lake. *J Appl Microbiol* 125:1626–1641.

32. Tarnecki AM, Burgos FA, Ray CL, Arias CR. 2017. Fish intestinal microbiome: Diversity

and symbiosis unravelled by metagenomics. *J Appl Microbiol* 123:2–17.

33. Stevenson SJR, Lee KC, Handley KM, Angert ER, White WL, Clements KD. 2022.

Substrate degradation pathways, conserved functions and community composition of the

hindgut microbiota in the herbivorous marine fish *Kyphosus sydneyanus*. *Comp Biochem*

Phys A 272:111283.

34. Wong S, Rawls JF. 2012. Intestinal microbiota composition in fishes is influenced by host

ecology and environment. *Molec Ecol* 21:3100–3102.

35. Rimoldi S, Ceccotti C, Brambilla F, Faccenda F, Antonini M, Terova G. 2023. Potential of

shrimp waste meal and insect exuviae as sustainable sources of chitin for fish feeds.

Aquaculture 567:739256.

36. German DP, Nagle BC, Villeda JM, Ruiz AM, Thomson AW, Contreras Balderas S, Evans

DH. 2010. Evolution of herbivory in a carnivorous clade of minnows (Teleostei:

Cyprinidae): Effects on gut size and digestive physiology. *Physiol Biochem Zool* 83:1–

18.

37. Eggink KM, Pedersen PB, Lund I, Dalsgaard J. 2022. Chitin digestibility and intestinal

exochitinase activity in Nile tilapia and rainbow trout fed different black soldier fly

larvae meal size fractions. *Aquac Res* 53:5536–5546.

38. Benavides AG, Cancino JM, Ojeda FP. 1994. Ontogenetic change in the diet of *Aplodactylus*

punctatus (Pisces: Aplodactylidae): An ecophysiological explanation. *Mar Biol* 118:1–5.

39. Pardesi B, Roberton AM, Lee KC, Angert ER, Rosendale DI, Boycheva S, White WL,

Clements KD. 2022. Distinct microbiota composition and fermentation products indicate

functional compartmentalization in the hindgut of a marine herbivorous fish. *Molec Ecol*

31:2494–2509.

40. Zhang Z, Li D, Refaey MM, Xu W. 2017. High spatial and temporal variations of microbial

community along the southern catfish gastrointestinal tract: Insights into dynamic food

digestion. *Front Microbiol* 8:1531.

41. Harris CR. 2019. Interactions between predators, diets, and the gut microbiome of tropical

reef fish. Master's thesis. University of Delaware, Newark, DE.

42. Parada AE, Needham DM, Fuhrman JA. 2016. Every base matters: assessing small subunit

rRNA primers for marine microbiomes with mock communities, time series and global

field samples. *Environ Microbiol* 18:1403-1414.

43. Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, Lesniewski RA,

Oakley BB, Parks DH, Robinson CJ, et al. 2009. Introducing mothur: Open-source,

platform-independent, community-supported software for describing and comparing

microbial communities. *Appl Environ Microbiol* 75:7537-7541.

44. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Oliver Glöckner F.

2013. The SILVA ribosomal RNA gene database project: improved data processing and

web-based tools. *Nucleic Acids Res* 41:D590–D596.

45. Victorian Bioinformatics Consortium. 2014. *Nesoni* https://github.com/Victorian-

Bioinformatics-Consortium/nesoni.

46. Peng Y, Leung HC, Yiu SM, Chin FY. 2012. IDBA-Ud: A de novo assembler for single-cell

and metagenomic sequencing data with highly uneven depth. *Bioinformatics* 28:1420–

1428.

47. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. Metaspades: A new versatile

metagenomic assembler. *Genome Res* 27:824–834.

48. Li D, Luo R, Liu CM, Leung CM, Ting HF, Sadakane K, Yamashita H, Lam TW. 2016.

Megahit v1.0: A fast and scalable metagenome assembler driven by advanced

methodologies and community practices. *Methods* 102:3–11.

49. Seah BK, Gruber-Vodicka HR. 2015. gbtools: Interactive visualization of metagenome bins

in R. *Front Microbiol*. 6:1451.

50. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search

tool. *J Molec Biol* 215:403-410.

51. The UniProt Consortium. 2023. UniProt: the Universal Protein Knowledgebase in 2023.

Nucleic Acids Res 51:D523-D531.

52. Wu YW, Simmons BA, Singer SW. 2015. MaxBin 2.0: An automated binning algorithm to

recover genomes from multiple metagenomic datasets. *Bioinformatics* 32:605–607.

53. Kang DD, Froula J, Egan R, Wang Z. 2015. MetaBAT, an efficient tool for accurately

reconstructing single genomes from complex microbial communities. *PeerJ* 3:e1165.

54. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: An

adaptive binning algorithm for robust and efficient genome reconstruction from

metagenome assemblies. *PeerJ*, 7:e7359.

55. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. Checkm: Assessing

the quality of microbial genomes recovered from isolates, single cells, and metagenomes.

Genome Res 25:1043–1055.

56. Bushnell B. 2014. BBMap: A fast, accurate, splice-aware aligner. *Lawrence Berkeley*

National Laboratory. LBNL Report #: LBNL-7065E

https://escholarship.org/uc/item/1h3515gn.

57. Edgar RC. 2004. Muscle: Multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res* 32:1792–1797.

58. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – approximately maximum-likelihood trees

for large alignments. *PLoS ONE*: 5:e9490.

59. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. 2022. GTDB-TK V2: Memory friendly

classification with the genome taxonomy database. *Bioinformatics* 38:5315–5316.

60. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:

Prokaryotic gene recognition and translation initiation site identification. *BMC*

Bioinformatics 11:119.

61. Farag IF, Zhao R, Biddle JF. 2021. "*Sifarchaeota*," a novel Asgard phylum from Costa Rican

sediment capable of polysaccharide degradation and anaerobic methylotrophy. *Appl*

Environ Microb 87:e02584-20.

62. Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. 2023. KEGG for

taxonomy-based analysis of pathways and genomes. *Nucleic Acids Res* 51:D587–D592.

63. Kanehisa M, Sato Y, Morishima K. 2016. Blastkoala and Ghostkoala: KEGG tools for

functional characterization of genome and metagenome sequences. *J Mol Biol* 428:726–

731.

64. Zhou Z, Tran PQ, Breister AM, Liu Y, Kieft K, Cowley ES, Karaoz U, Anantharaman K

2022. METABOLIC: high-throughput profiling of microbial genomes for functional

traits, metabolism, biogeochemistry, and community-scale functional networks.

Microbiome 10:33.

65. Zheng J, Ge Q, Yan Y, Zhang X, Huang L, Yin Y. 2023. dbCAN3: Automated carbohydrate-

active enzyme and substrate annotation. *Nucleic Acids Res* 51:W115–W121.

66. Kumla J, Suwannarach N, Sujarit K, Penkhrue W, Kakumyan P, Jatuwong K, Vadthanarat S,

Lumyong S. 2020. Cultivation of mushrooms and their lignocellulolytic enzyme

production through the utilization of agro-industrial waste. *Molecules* 25:12.

67. De Tender C, Mesuere B, Van der Jeugt F, Haegeman A, Ruttink T, Vandecasteele B,

Dawyndt P, Debode J, Kuramae EE. 2019. Peat substrate amended with chitin modulates

the N-cycle, siderophore and chitinase responses in the lettuce rhizobiome. *Sci Rep*

9:9890.

68. Drula E, Garron ML, Dogan S, Lombard V, Henrissat B, Terrapon N. 2022. The

carbohydrate-active enzyme database: Functions and literature. *Nucleic Acids Res*

50:D571–D577.

69. Rawlings ND, Waller M, Barrett AJ, Bateman A. 2014. MEROPS: The database of

proteolytic enzymes, their substrates and inhibitors. *Nucleic Acids Res* 42:D503–D509.

70. Keseler IM, Collado-Vides J, Santos-Zavaleta A, Peralta-Gil M, Gama-Castro S, Muñiz-

Rascado L, Bonavides-Martinez C, Paley S, Krummenacker M, Altman T, et al. 2011.

EcoCyc: A comprehensive database of *Escherichia coli* biology. *Nucleic Acids Res*

39:D583–D590.

71. Knappe J, Sawers G. 1990. A radical-chemical route to acetyl-CoA: The anaerobically

induced pyruvate formate-lyase system of *Escherichia coli*. *FEMS Microbiol Rev* 6:383–

398.

72. Aramaki T, Blanc-Mathieu R, Endo H, Ohkubo K, Kanehisa M, Goto S, Ogata H. 2020.

KofamKOALA: KEGG ortholog assignment based on profile HMM and adaptive score

threshold. *Bioinformatics* 36:2251–2252.

73. Søndergaard D, Pedersen CNS, Greening C. 2016. HydDB: A web tool for hydrogenase

classification and analysis. *Sci Rep* 6:34212.

74. R Core Team. 2023. R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. https://www.R-project.org/

75. Kassambara A, Mundt F. 2020. factoextra: Extract and visualize the results of multivariate

data analyses. R package version 1.0.7.999.

http://www.sthda.com/english/rpkgs/factoextra

76. Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P,

Stevens M, Szoecs E, et al. 2022. vegan: Community ecology package. R package

version 2.6-4. https://CRAN.R-project.org/package=vegan

Acknowledgements:

We would like to thank our colleagues at UD in Lewes for assistance with fish care and

sampling. This work was supported by The Mindlin Foundation, funding to CR via the UD

Graduate Scholars Fellowship, funding to KK and JFB from the WM Keck Foundation and

funding to IF and JFB by ExxonMobil Research and Engineering Company. DW was supported

as a UD School of Marine Science & Policy scholar. Support from the University of Delaware

CBCB Bioinformatics Data Science Core Facility (RRID:SCR_017696), including use of the

BIOMIX and BioStore computational resources was made possible through funding from

Delaware INBRE (NIGMS P20GM103446), NIH Shared Instrumentation Grant (S10OD028725)

the State of Delaware, and the Delaware Biotechnology Institute.

Figures

Figure 1: Coverage of 30S ribosomal protein annotations from the assembled metagenomes

compared to coverage within each MAG. Samples shown are a) piscivorous hawkfish, b)

herbivorous yellow tang, and c) invertivorous triggerfish. The comparison highlights which taxa

are represented by MAGs. Presumably those not in MAGs have a high degree of species

diversity and could not be accurately binned. The taxonomy of the 30S ribosomal protein gene

was determined by best hit in Uniprot, the taxonomy of the MAG was determined by GTDB.

Figure 2: Principal component analysis of normalized enzyme gene copy numbers across

all MAGs recovered from the three diet treatments. The total numbers of enzymes of each

degradation pathway (proteins, cellulose, hemicellulose, starch, and chitin) normalized to the

MAG's genome completeness were used to generate components. Fatty acids were included as a

binary where 1 designated that the MAG could complete all four steps of beta oxidation, while 0

designated that it could not. Ellipses represent t distributions around the centroids. Ellipses are

labeled for clarity. Individual points represent individual MAGs. The larger shapes in the ellipses

are group centroids.

Figure 3: Polysaccharide degradation pathway to monosaccharide products. Shown are a)

cellulose, b) xylan, c) mannan, d) xyloglucan, and e) chitin. Enzymes and their CAZyme IDs are

noted above arrows connecting substrates and products. MAGs that possess genes for each

enzyme are denoted beneath the arrow, where dark brown (left), green (center), and pink (right)

font represent the piscivorous, herbivorous, and invertivorous treatments, respectively.

Figure 4: Number of substrate degradation genes per MAG normalized to genome

completeness. Shown are a) cellulose, b) hemicellulose, c) chitin, d) starch, and e) proteins in

each diet treatment. Kruskal Wallis values above bars represent group comparisons. Pairwise

comparison values denoted by brackets are the result of post-hoc Wilcoxon tests between diets.

Figure 5: Heatmaps of metabolic functions that can be performed by recovered MAGs.

Samples shown are a) piscivorous hawkfish, b) herbivorous yellow tang, and c) invertivorous

triggerfish. Red indicates the ability to perform a metabolic pathway, while blue indicates

inability.

Table 1: Metagenome statistics

	Dataset	Size (bp)	Contigs in assembly	Max contig (bp)	Ave contig (bp)	N50 (bp)	Bins generated[1]
Hawkfish	H1N1	371954018	791521	31299	470	459	18
Tang	T4LTN	332236806	687584	125302	483	474	18
Triggerfish	Tr3LTN	310457717	640083	67851	485	471	7

[1]Bins that passed quality threshold

Table 2: MAG taxonomic assignments, completeness, contamination, coverage and ANI values

MAG ID[1]	Phylum	Class	Order	Family	Genus	Species	Genome Completeness (%)	Contamination (%)	Coverage	ANI to nearest neighbor (or RED)[2]	Neighbor reference code in GTDB
H1N1.001	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Vibrionaceae*	*Photobacterium*	*aphoticum*	97.58	2.55	94.31	95.41	GCF_001029435.1
H1N1.003	*Proteobacteria*	*Gammaproteobacteria*	*Pseudomonadales*	*Alcanivoracaceae*	*Alcanivorax*	sp.	100	0.74	61.69	97.91	GCA_002706085.1
H1N1.005	*Proteobacteria*	*Alphaproteobacteria*	*Parvibaculales*	*Phaeomarinobacteraceae*	*Phaeomarinobacter*	sp.	86.52	5.77	35.76	79.02	GCF_000689395.1
H1N1.006	*Proteobacteria*	*Alphaproteobacteria*	*Rhodobacterales*	*Rhodobacteraceae*	*Phaeobacter*	*italicus*	99.66	6.79	35.06	98.07	GCF_001258055.1
H1N1.007	*Proteobacteria*	*Alphaproteobacteria*	*Rhodospirillales*	*Thalassospiraceae*	*Thalassospira*	sp.	100	1	33.33	85.04	GCF_003326795.1
H1N1.009	*Bacteroidota*	*Bacteroidia*	*Cytophagales*	*Cyclobacteriaceae*	*Fabibacter*	*pacificus*	98.73	2.89	31.61	96.83	GCF_900111145.1
H1N1.010	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Vibrionaceae*	*Vibrio*	*fortis*	74.03	9.2	41.62	96.01	GCF_000695685.1
H1N1.011	*Patescibacteria*	*Gracilibacteria*	UBA1369	GCA-2718715	-	-	73.06	0.93	18	76.01	GCA_002718715.1
H1N1.012	*Proteobacteria*	*Alphaproteobacteria*	*Rhodobacterales*	*Rhodobacteraceae*	*Ruegeria*	sp.	98.27	3.69	16.12	RED, 0.98	n/a
H1N1.015	*Bacteroidota*	*Bacteroidia*	*Cytophagales*	*Cyclobacteriaceae*	*Fabibacter*	*spongicola*	98.01	0.6	12.06	96.92	GCF_001592965.1
H1N1.018	*Proteobacteria*	*Alphaproteobacteria*	*Minwuiales*	*Minwuiaceae*	*Minwuia*	sp.	98.91	3.77	13.26	79.34	GCF_002924445.1
H1N1.021	*Proteobacteria*	*Gammaproteobacteria*	*Pseudomonadales*	*Porticoccaceae*	*Porticoccus*	sp.	89.33	3.83	7.95	95.8	GCA_006226995.1
H1N1.022	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Alteromonadaceae*	*Alteromonas*	sp.	92.49	9.21	7.86	90.14	GCF_003731635.1
H1N1.025	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Alteromonadaceae*	*Pseudoalteromonas*	*spongiae*	70.69	6.03	9.77	96.51	GCF_000238255.3
H1N1.026	*Proteobacteria*	*Alphaproteobacteria*	*Sphingomonadales*	*Sphingomonadaceae*	*Erythrobacter*	*flavus*	94.33	0.86	8.16	97.62	GCF_002237615.1
H1N1.028	*Proteobacteria*	*Alphaproteobacteria*	*Caulobacterales*	*Maricaulaceae*	*Maricaulis*	sp.	93.44	1.68	6.04	81.73	GCF_005871165.1
H1N1.030	*Bacteroidota*	*Bacteroidia*	*Cytophagales*	*Flammeovirgaceae*	*Sediminitomix*	sp.	93.49	8.17	4.71	80.04	GCF_003149185.1
H1N1.033	*Proteobacteria*	*Alphaproteobacteria*	UBA8366	GCA-2696645	GRI0909	sp.	87.3	12.75	4.31	80.5	GCF_004005845.1
T4LTN.001	*Bacteroidota*	*Bacteroidia*	*Bacteroidales*	*Bacteroidaceae*	*Bacteroides*	sp.	98.51	0.5	240.58	98.79	GCA_002471185.1
T4LTN.002	*Bacteroidota*	*Bacteroidia*	*Bacteroidales*	*Paludibacteraceae*	-	-	85.3	5.67	45.25	RED, 0.86	n/a
T4LTN.003	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Vibrionaceae*	*Vibrio*	*owensii*	92.93	1.51	51.4	96.86	GCF_000817815.1
T4LTN.004	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Shewanellaceae*	*Shewanella*	sp.	94.4	3.15	39.3	84.16	GCF_003353085.1
T4LTN.005	*Bacteroidota*	*Bacteroidia*	*Bacteroidales*	*Tannerellaceae*	-	sp.	96.92	7.31	34.71	79.39	GCF_000969825.1
T4LTN.006	*Firmicutes*	*Clostridia*	*Lachnospirales*	*Lachnospiraceae*	-	sp.	99.11	4.83	21.97	80.86	GCF_000733755.1
T4LTN.007	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Alteromonadaceae*	-	-	91.46	10.06	33.35	77.83	GCF_002954545.1
T4LTN.009	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Alteromonadaceae*	-	-	92.4	3.7	17.73	78.04	GCF_002954545.1
T4LTN.010	*Spirochaetota*	*Spirochaetia*	*Sphaerochaetales*	*Sphaerochaetaceae*	-	-	100	2.3	21.93	77.22	GCA_001604275.1
T4LTN.011	*Bacteroidota*	*Bacteroidia*	*Bacteroidales*	UBA1402	UBA8389	sp.	100	2.02	18.54	76.81	GCA_003506275.1

ID	Phylum	Class	Order	Family	Genus	Species					Accession
T4LTN.012	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Vibrionaceae*	*Enterovibrio*	sp.	97.84	0.9	17.66	82.02	GCA_001310415.1
T4LTN.014.2	*Firmicutes*	*Bacilli*	RF39	CAG-822	UBA5364	sp.	96.63	0.61	19.86	76.06	GCA_002410935.1
T4LTN.014.3	*Patescibacteria*	*Gracilibacteria*	BD1-5	UBA6164	MAAV01	sp.	83.15	1.69	11.87	77.15	GCA_004563595.1
T4LTN.015	*Firmicutes*	*Clostridia*	*Oscillospirales*	*Ruminococcaceae*	UMGS363	sp.	97.32	6.64	22.67	RED, 0.92	n/a
T4LTN.016	*Proteobacteria*	*Gammaproteobacteria*	*Pseudomonadales*	*Saccharospirillaceae*	*Oleispira*	sp.	87.84	3.6	16.05	RED, 0.96	n/a
T4LTN.017	*Firmicutes*	*Clostridia*	*Oscillospirales*	*Ruminococcaceae*	-	-	99.32	0.91	9.67	RED, 0.79	n/a
T4LTN.022	*Proteobacteria*	*Alphaproteobacteria*	*Sphingomonadales*	*Sphingomonadaceae*	*Erythrobacter*	*flavus*	95.9	7.03	5.72	97.4	GCF_002237615.1
T4LTN.023	*Bacteroidota*	*Bacteroidia*	*Bacteroidales*	P3	-	-	86.31	2.89	5.4	RED, 0.90	n/a
Tr3LTN.002	*Proteobacteria*	*Gammaproteobacteria*	*Enterobacterales*	*Shewanellaceae*	*Shewanella*	sp.	98.87	1.08	74.15	84.49	GCF_003353085.1
Tr3LTN.005	*Proteobacteria*	*Alphaproteobacteria*	*Rhodobacterales*	*Rhodobacteraceae*	*Epibacterium*	sp.	98.94	3.55	11.73	96.78	GCF_001681715.1
Tr3LTN.006	*Proteobacteria*	*Gammaproteobacteria*	*Psuedomonadales*	*Cellvibrionaceae*	GCA-2707785	sp.	95.52	1.92	26.73	76.39	GCA_002707785.1
Tr3LTN.008	*Campylobacterota*	*Campylobacteria*	*Campylobacterales*	*Arcobacteraceae*	UBA4036	sp.	99.19	2.37	11.66	77.9	GCA_002382325.1
Tr3LTN.016	*Firmicutes*	*Bacilli*	-	-	-	-	84.8	7.11	5.09	RED, 0.70	n/a
Tr3LTN.017	*Proteobacteria*	*Alphaproteobacteria*	UBA1280	UBA12223	UBA12223	-	81.93	2.99	7.53	81.5	GCA_002937495.1
Tr3LTN.026	*Bdellovibrionota*	*Bacteriovoracia*	*Bacteriovoracales*	*Bacteriovoracaceae*	-	-	70.65	4.1	4.87	RED, 0.78	n/a

[1] Naming: H1N1, Hawkfish; T4LTN, Yellow tang; Tr3LTN, Triggerfish

[2]GTDB-tk determines values based on Average Nucleotide Identity (ANI; calculated by either FastANI or closest placement ANI) or Relative Evolutionary Distance (RED). If an organism cannot be processed with ANI, it is automatically processed with RED.

Table 3: Summary of biochemical functions that could be performed by all recovered MAGs.

MAG ID[1]	Family	Host Diet	Substrate Utilization[2]	Energy Production[3]	Vitamin Production	Benefits to Host
H1N1.001	Vibrionaceae	Piscivorous	Proteins, Starch	Glycolysis, Pyruvate Oxidation, TCA Cycle, Nitrate reduction, Formate fermentation, Acetogenesis	B1, B2, B7, B9, B12	Protein Degradation, Vitamin Production
H1N1.003	Alcanivoraceae	Piscivorous	Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle, Beta Oxidation, Acetogenesis (p)	B2, B7, B9, B12	Protein Degradation, Vitamin Production
H1N1.005	Phaeomarinobacteraceae	Piscivorous	Collagen, Proteins, Cellulose	Glycolysis, TCA Cycle (p), Pyruvate Oxidation	B7	Vitamin Prroduction
H1N1.006	Rhodobacteraceae	Piscivorous	Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Beta Oxidation, Acetogenesis	B7, B9, B12	Protein Degradation, Vitamin Production
H1N1.007	Thalassospiraceae	Piscivorous	Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle, Beta Oxidation, Lactate fermentation, Acetogenesis (p)	B1, B9, B12	Protein Degradation, Vitamin Thraduction
H1N1.009	Cyclobacteriaceae	Piscivorous	Proteins, Cellulose, Starch, Chitin	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Acetogenesis (p)	B7, B9	Protein Degradation, Vitamin Cyroduction
H1N1.010	Vibrionaceae	Piscivorous	Proteins, Starch	Pyruvate Oxidation, TCA Cycle (p), Nitrate reduction, Acetogenesis, Formate fermentation	B1, B7, B9, B12	Protein Degradation, Vitamin Prroduction
H1N1.011	GCA-2718715	Piscivorous	Cellulose	Glycolysis (p), Lactate fermentation, Acetogenesis (p)	-	-
H1N1.012	Rhodobacteraceae	Piscivorous	Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Beta Oxidation, Nitrate reduction, Acetogenesis	B7, B9, B12	Protein Degradation, Vitamin Prroduction
H1N1.015	Cyclobacteriaceae	Piscivorous	Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Beta Oxidation, Acetogenesis (p)	B7, B9	Vitamin Cyroduction
H1N1.018	Minwuiaceae	Piscivorous	Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle, Beta Oxidation, Acetogenesis(p)	B12	Protein Degradation, Vitamin Prroduction
H1N1.021	Porticoccaceae	Piscivorous	Proteins	Glycolysis (p), Pyruvate Oxidation, Acetogenesis (p)	B9	Protein Degradation, Vitamin Porticoduction
H1N1.022	Alteromonadaceae	Piscivorous	Proteins, Fatty Acids, Cellulose, Hemicellulose, Starch	Glycolysis, Pyruvate Oxidation, TCA Cycle, Beta Oxidation, Acetogenesis (p)	B1, B2, B7, B9	Protein Degradation, Vitamin Atroduction
H1N1.025	Alteromonadaceae	Piscivorous	Proteins, Starch	Glycolysis, Pyruvate Oxidation, Acetogenesis	B2, B9	Protein Degradation, Vitamin Production
H1N1.026	Sphingomonadaceae	Piscivorous	Proteins	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Acetogenesis (p)	B1, B7	Protein Degradation, Vitamin Sphroduction
H1N1.028	Maricaulaceae	Piscivorous	Proteins, Cellulose	Glycolysis (p), Pyruvate Oxidation, Acetogenesis (p)	B1, B7, B9	Protein Degradation, Vitamin Prroduction
H1N1.030	Flammeovirgaceae	Piscivorous	Proteins, Fatty Acids, Cellulose, Hemicellulose, Starch, Chitin	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Beta Oxidation, Acetogenesis	B1, B7, B9	Protein Degradation, Vitamin Flroduction
H1N1.033	GCA-2696645	Piscivorous	Proteins, Cellulose	Glycolysis (p), Pyruvate Oxidation, TCA Cycle (p), Acetogenesis (p)	-	Protein Degradation
T4LTN.001	Bacteroidaceae	Herbivorous	Cellulose, Hemicellulose, Starch, Proteins, Chitin	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), H2-Producing Fermentation, Acetogenesis (p)	B1, B2, B7, B9	Plant Matter Degradation, Vitamin Production
T4LTN.002	Paludibacteraceae	Herbivorous	Cellulose, Proteins	Glycolysis, Pyruvate Oxidation, Acetogenesis (p)	-	Plant Matter Degradation
T4LTN.003	Vibrionaceae	Herbivorous	Starch, Proteins, Fatty Acids, Chitin	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Nitrate reduction, Acetogenesis, Formate fermentation, Beta Oxidation	B1, B2, B7, B9	Plant Matter Degradation, Vitamin Production
T4LTN.004	Shewanellaceae	Herbivorous	Starch, Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle, Nitrate reduction, Acetogenesis, Formate fermentation, Beta Oxidation	B1, B2, B7, B9, B12	Plant Matter Degradation, Vitamin Shroduction
T4LTN.005	Tannerellaceae	Herbivorous	Cellulose, Hemicellulose, Starch, Proteins, Chitin	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), H$_2$-Producing Fermentation, Acetogenesis (p)	B1, B2, B7, B9, B12	Plant Matter Degradation, Vitamin Production
T4LTN.006	Lachnospiraceae	Herbivorous	Cellulose, Hemicellulose, Starch, Proteins, Chitin	Glycolysis, Pyruvate Oxidation, Lactate fermentation, H$_2$-Producing Fermentation, Acetogenesis (p)	B7, B12	Plant Matter Degradation, Vitamin Production
T4LTN.007	Alteromonadaceae	Herbivorous	Cellulose, Hemicellulose, Starch	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Nitrate reduction, Acetogenesis	B1, B2, B7, B9	Plant Matter Degradation, Vitamin Atroduction
T4LTN.009	Alteromonadaceae	Herbivorous	Cellulose, Starch	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Acetogenesis	B1, B7	Plant Matter Degradation, Vitamin Production
T4LTN.010	Sphaerochaetaceae	Herbivorous	Proteins	Glycolysis, Pyruvate Oxidation, H$_2$-Producing Fermentation, Acetogenesis (p)	-	-
T4LTN.011	UBA1402	Herbivorous	Hemicellulose	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Acetogenesis (p)	B9	Plant Matter Degradation, Vitamin Production
T4LTN.012	Vibrionaceae	Herbivorous	Starch, Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle, Nitrate reduction, Acetogenesis, Formate fermentation, Beta Oxidation	B1, B2, B7, B9, B12	Plant Matter Degradation, Vitamin Production
T4LTN.014.2	CAG-822	Herbivorous	Starch	Glycolysis (p), Pyruvate Oxidation, Acetogenesis (p)	-	Plant Matter Degradation
T4LTN.014.3	UBA6164	Herbivorous	Simple Saccharides	-	-	-
T4LTN.015	Ruminococcaceae	Herbivorous	Cellulose, Hemicellulose, Chitin	Glycolysis, Pyruvate Oxidation, H$_2$-Producing Fermentation, Acetogenesis (p)	-	Plant Matter Degradation

T4LTN.016	Saccharospirillaceae	Herbivorous	Cellulose, Proteins, Fatty Acids	Pyruvate Oxidation, Nitrate reduction, Acetogenesis, Beta Oxidation	B1, B7, B9	Plant Matter Degradation, Vitamin Production
T4LTN.017	Ruminococcaceae	Herbivorous	Simple Saccharides	Glycolysis, Pyruvate Oxidation, H_2-Producing Fermentation, Acetogenesis (p)	B7	Vitamin Production
T4LTN.022	Sphingomonadaceae	Herbivorous	Proteins	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Acetogenesis (p)	B1	Vitamin Sproduction
T4LTN.023	P3	Herbivorous	Proteins	Glycolysis, Pyruvate Oxidation, H_2-Producing Fermentation, Acetogenesis (p)	B1	Vitamin Production
Tr3LTN.002	Shewanellaceae	Invertivorous	Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle, Nitrate reduction, Acetogenesis, Formate fermentation, Beta Oxidation	B1, B2, B7, B9, B12	Protein Degradation, Vitamin Production
Tr3LTN.005	Rhodobacteraceae	Invertivorous	Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, TCA Cycle (p), Beta Oxidation	B2, B9, B12	Protein Degradation, Vitamin Production
Tr3LTN.006	Cellvibrionaceae	Invertivorous	Proteins, Fatty Acids	Glycolysis, Pyruvate Oxidation, Beta Oxidation	B1, B7, B9	Protein Degradation, Vitamin Production
Tr3LTN.008	Arcobacteraceae	Invertivorous	Simple Saccharides	Glycolysis, Pyruvate Oxidation, TCA Cycle (p)	B1, B7	Vitamin Production
Tr3LTN.016	Class: Bacilli	Invertivorous	Simple Saccharides	Glycolysis, Pyruvate Oxidation, H_2-Producing Fermentation	-	-
Tr3LTN.017	UBA12223	Invertivorous	Simple Saccharides	Glycolysis (p), Pyruvate Oxidation, TCA Cycle (p)	-	-
Tr3LTN.026	Bacteriovoracaceae	Invertivorous	Simple Saccharides	Glycolysis (p), Pyruvate Oxidation	-	-

[1] Naming: H1N1, Hawfish; T4LTN, Yellow tang; Tr3LTN, Triggerfish

[2]Carbohydrate substrates are included if the MAG was capable of completely degrading it from start to finish. Starch is listed if the MAG possesses more than 5 amylase genes. Proteins are listed if the MAG possesses more than 10 exopeptidase genes. If the MAG could only produce β-glucosidase, simple saccharides is listed.

[3]For energy production, the pathway is included if the MAG can complete it entirely. If the MAG can complete an energy yielding/harvesting step of a pathway but not the entire process, the function is listed with a "p" (partial).