Parkinson's disease-associated shifts between DNA methylation and DNA hydroxymethylation in human brain

Juliana I. Choza, BA[3], Mahek Virani, BA[3], Nathan C. Kuhn, BS[1], Marie Adams, MS[2], Joseph Kochmanski, PhD[1], and Alison I. Bernstein, PhD[1,3,4]

[1] Department of Translational Neuroscience, College of Human Medicine, Michigan State University, Grand Rapids, MI
[2] Genomics Core, Van Andel Research Institute, Grand Rapids, MI
[3] Department of Pharmacology and Toxicology, and
[4] Environmental and Occupational Health Sciences Institute, Ernest Mario School of Pharmacy, Rutgers University, Piscataway, NJ

Corresponding Author: Alison Bernstein, bernstein.alison@rutgers.edu
Environmental and Occupational Health Sciences Institute
Ernest Mario School of Pharmacy, Rutgers University
170 Freylinghuysen Rd
Piscataway, NJ 08854

Orcid IDs
Juliana I. Choza: 0000-0001-7038-9750
Mahek Virani: 0009-0006-6094-4478
Marie Adams: 0000-0001-7909-2339
Joseph Kochmanski: 0000-0002-8472-3032
Alison I. Bernstein: 0000-0002-5589-4318

Word count:

Running title: DNA hydroxymethylation in Parkinson's disease

Conflict of interest statement: All authors declare that they have no conflicts of interest

Funding sources: This work was supported by the National Institute of Environmental Health Sciences of the National Institutes of Health (R01 ES031237-03S1 to AIB). The Banner Sun Health Research Institute Brain and Body Donation Program has been supported by the National Institute of Neurological Disorders and Stroke (U24 NS072026 National Brain and Tissue Resource for Parkinson's Disease and Related Disorders), the National Institute on Aging (P30 AG19610 Arizona Alzheimer's Disease Core Center), the Arizona Department of Health Services (contract 211002, Arizona Alzheimer's Research Center), the Arizona Biomedical Research Commission (contracts 4001, 0011, 05-901 and 1001 to the Arizona Parkinson's Disease Consortium) and the Michael J. Fox Foundation for Parkinson's Research.

Abstract

Background: The majority of PD cases are due to a complex interaction between aging, genetics, and environmental factors; epigenetic mechanisms are thought to act as important mediators of these risk factors. While multiple studies to date have explored the role of DNA modifications in PD, few focus on 5-hydroxymethylcytosine (5hmC), which is thought to be particularly important in the brain and the response to environmental exposures.

Objectives: The goal of this study was to identify paired changes in 5hmC and 5-methylcytosine (5mC) in PD in enriched neuronal nuclei isolated from PD post-mortem parietal cortex and age- and sex-matched controls.

Methods: We performed oxidative bisulfite (BS) conversion and paired it with our previously published BS-based EWAS to identify cytosines with significant shifts between these two related epigenetic marks. Interaction differentially modified cytosines (iDMCs) were identified using our recently published mixed effect model for co-analyzing β_{mC} and β_{hmC} data.

Results: We identified 1,030 iDMCs with paired changes in 5mC and 5hmC (FDR < 0.05) that map to 695 genes, including the PD risk gene, DNAJC6 (*PARK19*).

Conclusions: These data potentially link epigenetic regulation of the *PARK19* locus in the pathogenesis of idiopathic PD. In addition, iDMC-containing genes have known functions in synaptic formation and function, cell cycle and senescence, neuroinflammation, and epigenetic regulation. These data suggest that there are significant shifts between 5mC and 5hmC associated with PD in genes relevant to PD pathogenesis that are not captured by analyzing BS-based data alone or by analyzing each mark as a distinct dataset.

Introduction

An estimated 5-10% of Parkinson's disease (PD) cases are familial and caused by monogenically inherited mutations, while the remaining ~90% of sporadic cases (sPD) are likely due to a complex interaction between age, genes, and environmental factors.[1-4] While the relative contribution of genetic and environmental risk factors in the etiology of sPD is debated, it is well documented that they play critical roles in the large majority of PD cases. Epigenetic mechanisms have emerged as critical mediators of the complex interactions between aging, genetics, the environment, and disease risk because they are dynamic with age, sensitive to the environment, and regulate gene expression throughout the lifespan.[5-10] Evidence for the role of epigenetic regulation in PD has been building, particularly for DNA modifications.[10-18]

5-methylcytosine (5mC), the addition of a methyl group to the 5′-carbon of cytosine, is one of the most well-studied epigenetic marks.[19] 5-hydroxymethylcytosine (5hmC) is formed via oxidation of 5mC by ten-eleven translocation (TET) enzymes and is a stable, independent epigenetic mark that has its highest levels in the brain, recruits a distinct set of DNA binding proteins from 5mC, differs in its genomic distribution in the brain compared 5mC, and is enriched in transcriptionally active gene bodies in the nervous system, suggesting a specific regulatory role for 5hmC in the brain.[20-29] As such, 5hmC is now thought to be particularly important in gene regulation in brain, particularly in the response to neurotoxicants.[19, 22, 30, 31] However, most studies of DNA modifications in PD brain have relied on bisulfite (BS) conversion, which cannot distinguish between 5mC and 5hmC.[10, 32-40]

Recently, studies have begun to explore links between 5hmC and PD.[41-43] First, rare variants in *TET1*, which encodes an enzyme responsible for conversion of 5mC to 5hmC, were associated with increased risk of PD in a Chinese PD cohort.[41] Second, a targeted analysis of DNA modifications within known enhancers in human postmortem prefrontal cortex identified epigenetic disruption of an enhancer targeting the *TET2* gene, which encodes another enzyme responsible for conversion of 5mC to 5hmC, in PD patients.[43] This study also performed hydroxymethylated DNA immunoprecipitation-sequencing (hMeDIP-Seq) and found that PD-associated-hydroxymethylated peaks were enriched in gene bodies, promoters, and enhancers. Third, a small study in human postmortem substantia nigra (SN) used hMe-Seal, a selective chemical labeling method, and identified thousands of differentially hydroxymethylated regions in genes related to CNS and neuronal differentiation, neurogenesis, and development and maintenance of neurites and axons, although the widespread neurodegeneration in the SN by the time of PD diagnosis complicates interpretation of these results.[42] Regardless, taken together, these initial studies support a role for 5hmC in regulation of expression of genes important for PD pathogenesis and indicate that additional research is warranted.

In our previous study, we performed a neuron-specific epigenome-wide association study (EWAS) with the Illumina EPIC BeadChip array paired with BS conversion using enriched neuronal nuclei from human postmortem parietal cortex obtained from the Banner Sun Health Research Institute Brain Bank.[44] We identified largely sex-specific PD-associated changes in DNA modification in 434 unique genes, including genes previously implicated in PD, including *PARK7* (DJ-1), *SLC17A6* (VGLUT2), *PTPRN2* (IA-2β), and *NR4A2* (NURR-1), as well as genes involved in developmental pathways, neurotransmitter packaging and release, and axon/neuron projection guidance. However, we could not differentiate between 5mC and 5hmC because we used BS.

Here, we report the results of an epigenome-wide analysis of 5hmC and 5mC in enriched neurons from PD brain using our recently proposed method for reconciling base-pair resolution 5mC and 5hmC data.[45-47] This method considers 5mC and 5hmC as paired data since these marks are biologically and statistically dependent on each other. We utilized additional DNA isolated from the same neuron-enriched samples and performed oxidative BS (oxBS) conversion paired with the Illumina EPIC BeadChip array to specifically measure 5mC.[48, 49] oxBS adds an oxidation step with potassium perruthenate ($KRuO_4$) that specifically oxidizes 5hmC, forming 5-formylcytosine, prior to BS conversion. BS then deaminates 5mC only, but not 5-formylcytosine, such that C and 5hmC are read as thymine, while only the 5mC is read as cytosine, providing a readout of "true" methylation. Comparison of BS and oxBS results allows estimation of 5hmC. To our knowledge, this is the largest epigenome-wide analysis of 5hmC to date in neurons enriched from PD post-mortem brain at single base pair resolution.

Methods

Human brain tissue

De-identified tissue samples from control (n=50) and sPD (n=50) human brain samples were obtained from archival human autopsy specimens provided by the Banner Sun Health Research Institute (BSHRI), using BSHRI's approved institutional review board (IRB) protocols. Further details about the BSHRI's brain samples and sample selection are available in a previous publication.[50] We selected PD patients with mid-stage disease (Braak stage = II–III), as defined by Lewy pathology.[51] Postmortem parietal cortex was selected because this region develops pathology in the late stages of PD and is expected to still have robust populations of neurons from which disease-associated, pre-degenerative gene regulatory marks can be measured.

Magnetic-activated cell sorting

DNA samples previously isolated from an enriched neuronal population from de-identified parietal cortex samples from the BSHRI Brain Bank were used for this study. NeuN-positive (NeuN$^+$) nuclei were enriched from 100 mg of flash frozen parietal cortex tissue using a two-stage magnetic-assisted cell sorting (MACS) method as previously described.[44]

DNA extraction

DNA was isolated from enriched NeuN$^+$ nuclei using the Qiagen QIAamp DNA Micro Kit (Cat. # 56304) as previously described. [44]

Oxidative bisulfite treatment and EPIC arrays

Intact genomic DNA yield was quantified by Qubit fluorometry (Life Technologies). oxBS conversion was performed on 1 µg genomic DNA using the TrueMethyl Array kit (Cambridge Epigenetix). While the recommended amount of 500 ng DNA was sufficient for BS conversions, this was insufficient for oxBS, likely due to the additional harsh oxidation step. Cleanup and preparation of DNA, and all steps for the EPIC bead chip protocol were performed as previously described per the manufacturer's protocol.[44]

EPIC array data processing of oxBS data

IDAT files were imported into R and processed using an in-house bioinformatics pipeline that utilizes *minfi* (version 1.48.0), *ChAMP* (version 2.32.0), *posibatch* (version 1.0), and *ENmix* (version 1.38.01), and *CETS* (version 3.03) packages in R, as previously described (Supplementary File 1). [44, 45, 49, 52-59] After QC, eleven female samples and six male samples were

removed due to a high level (>10%) of failed probes, leaving 57 male (28 PD, 29 control) and 26 female (13 PD, 13 control) samples. Failed probes (53,305) were removed from remaining samples when detection p-value was > 0.01 in more than 5% of samples. Cross-reactive probes and probes containing SNPs (93,527) were masked based on previous identification.[60]

We continued with male samples only due to small sample size of the remaining female samples. Samples with estimated glial cell proportion > 0.50 were also removed from analysis. This removed one sample from the male data set, leaving 56 males (27 PD, 29 control). Data for the included samples are summarized in Table 1 and full metadata is in Supplementary File 3.

5hmC β values ($β_{hmC}$) were estimated by pairing oxBS β values with our previously published BS data using the maximum likelihood estimate function (oxBS.MLE) from the *ENmix* package, which returns true methylation β values ($β_{mC}$) and estimates $β_{hmC}$. Raw BS and oxBS values, as well as MLE-corrected $β_{mC}$ and $β_{hmC}$ values, are shown in Figure 1A,B.

After MLE, probes with mean $β_{mC}$ or $β_{hmC}$ < 0.01 across all samples were removed due to increased variability and decreased interpretability of β values at such low levels, as well as to remove the issue of zero inflation for 5hmC β values. We then produced three data sets: one with all probes where $β_{mC}$ > 0.01 (714,199 probes), one with all probes where $β_{hmC}$ > 0.01 (588,147), and one with only probes common to the two sets, where $β_{mC}$ > 0.01 and $β_{hmC}$ > 0.01 (588,123). Of these, only 12 probes had $β_{hmC}$ data only, and 126,076 had $β_{mC}$ data only.

Differential testing for individual CpG sites: The *gamlss* R package (version 5.4-22) was used to test for interaction differentially methylated cytosines (iDMCs) as previously described (Supplementary File 2). [44, 45, 61, 62] All models included glial cell proportion estimated from BS data and PMI as covariates.[44] An FDR < 0.05 was used as the cutoff for significance and annotation of significant differential probes was performed using the Illumina EPIC array manifests. P-value histograms generated to confirm accuracy of modeling show that the interaction model provided more appropriate histograms (uniform with an overabundance of low p-values), whereas the separate models produced histograms with an overabundance of high p-values (Supplementary Figure 1).[63] For separate analyses of $β_{mC}$ and $β_{hmC}$, neither normal nor beta regression modeling accurately modeled the data. All code for data visualizations is provided in Supplementary Files 7,8.

Annotation of interaction DMCs: Gene IDs corresponding to each iDMC were extracted from the EPIC array manifest provided by Illumina (v1.0 B5). Annotation of universal chromatin states was also performed.[64] Since this study utilized neuronally enriched nuclei, neuronal expression of iDMC-containing genes was verified using the Allen Brain Cell Atlas.[65] The Harmonize and GeneCards databases were used to explore gene function.[66, 67]

Gene ontology pathway enrichment and protein-protein interaction networks: Gene ontology (GO) term enrichment testing and pathway analysis was performed on genes annotated to iDMCs identified in the interaction model using the ClueGO application in Cytoscape (version 3.10.1).[68, 69] "Groups" was selected as the visual style, and the GO biological process (GOBP) term was selected. Network specificity was set to "Medium", with the GO Tree Interval minimum set at 3 and maximum at 8. Only terms with at least 3 genes and a Bonferroni-corrected p-value < 0.05 were included in pathway visualizations. The connectivity score (Kappa) was set at 0.4, and default GO Term Grouping settings were used in all analyses. The genes found in enriched GOBP terms by ClueGO were used for protein-protein interaction network analysis using STRING (version 12.0).[70] STRING network analysis was performed

using default parameters, including a minimum required interaction score = 0.9 (Figure 2) or 0.7 (Supplementary Figure 2).

Results

We identified 1,030 iDMCs with significant shifts in the proportions of 5mC and 5hmC associated with PD in DNA isolated from an enriched neuronal population derived from parietal cortex (FDR < 0.05) (Figure 1C; Supplementary File 4). More iDMCs (652) have a negative than a positive beta coefficient (378). In this output, a positive beta coefficient (interaction term) indicates a relative decrease in 5mC and increase in 5hmC in PD brains, as illustrated by the most significant DMC with a positive beta coefficient and the DMC with the most positive beta coefficient (Figure 1F,H). In contrast, a negative interaction term indicates a relative increase in 5mC and a decrease in 5hmC in PD brains, as illustrated by the most significant DMC with a negative beta coefficient and the most negative beta coefficient (Figure 1G,I). Corresponding raw BS β-values for each iDMCs demonstrate that these probes were not identified by BS data alone (Figure 1F-J). Of these 1,030 iDMCs, 681 are found within genes and annotate to 695 genes, including the PD risk gene *PARK19* (DNAJC6) (Figure 1J; Supplementary File 5). Only 43 of these genes were also identified in our previous BS-only EWAS, but at different cytosines (Supplementary File 6).[44]

Of the 695 genes, 375 annotated to 16 enriched GO terms in 7 GO term groups, which are enriched for genes involved in nervous system development, cell-cell adhesion, regulation of signal transduction and cell communication, morphogenesis, and cell differentiation (Figure 2A; Table 2). Genes within the enriched GO terms were used for STRING network analysis to identify potential functional interactions between proteins encoded by differentially modified genes. For clarity, networks of proteins within enriched GO terms with the highest stringency settings are shown in Figure 2B; less stringent networks for each individual GO term group are shown in Supplementary Figure 2 to highlight the high degree of potential connectivity within each group.

While there are major differences in existing PD EWAS related to sample size, sample selection, brain region assessed, methods used to measure DNA modifications, and statistical modeling, as well as inconsistent reporting between studies, we compared the list of genes in this study to recent brain-specific EWAS studies for PD, including ours, for data was provided.[34, 35, 38, 43] One study performed hMe-DIP in prefrontal cortex to assess genome-wide 5hmC.[43] In addition, three BS-based studies used either the EPIC array or the previous 450K array in SN, frontal cortex, the dorsal motor nucleus of the vagus (DMV), and cingulate gyrus (CG).[34, 35, 38] Not surprisingly, given the differences between these studies overlap between our data and these studies was minimal (Table 3, Supplementary File 7). The gene most consistently identified in these studies is *PTPRN2*, which we previously reported as showing sexually dimorphic alterations in DNA modifications and has previously been implicated in PD.[44, 71-73] The methodological differences between these studies complicate the comparison of these studies and underscore the need for rigorous and reproducible methodology and analysis, as discussed in our recent chapter on rigor and reproducibility in EWAS.[74]

Discussion

Significant shifts between 5mC and 5hmC associated with PD

Here, we performed an integrated genome-wide analysis of 5mC and 5hmC using our novel application of mixed effects modeling in enriched neuronal nuclei from PD post-mortem parietal cortex samples.[45-47] These PD-associated iDMCs were largely unique from DMCs identified in our previous BS-based EWAS: 43 genes were identified in both studies, 652 only in the current analysis, and 480 genes identified only in the previous analysis (Supplementary Files 5,6).[44] Collectively, these data suggest that there are significant PD-associated shifts between 5mC and 5hmC at iDMCs that are not captured by analyzing BS-based data alone or by analyzing each mark as distinct datasets (Figure 1C).[44, 45] Because many of these iDMCs are located within genes previously implicated in PD and in genes within pathways thought to play a role in PD, these data suggests that shifts in the balance between DNA modifications may play an important but unrecognized role in PD etiology in both known and novel PD-related genes.

An important caveat of these findings is that this study does not address the biological significance of these epigenetic shifts. While shifts between 5mC and 5hmC may potentially impact the binding of proteins that regulate gene expression and/or other epigenetic marks, this study does not examine these functional impacts; however, it does provide multiple avenues for further study of the impact of these changes on gene expression, alternate promoter usage, differential isoform expression, and neuronal function and susceptibility.

Epigenetic regulation of the PARK19 (DNAJC6) PD risk allele

As with our previous study, we identified epigenetic changes in the known PD risk genes. Here, we identified an iDMC within the 3′ UTR of *PARK19,* which encodes DNAJC6. This iDMC shows increased 5mC and decreased 5hmC in PD and is located in a region annotated by universal chromatin state annotation as a transcribed enhancer (Figure 1G; Supplementary File 5).[64] Autosomal recessive mutations in *PARK19* predominantly cause juvenile- or early-onset PD, with atypical signs and symptoms characterized by poor L-DOPA responsiveness, pyramidal signs, dystonia, rapid disease course, mental retardation, and seizures and is one of multiple DNAJC genes associated with familial forms of PD. [2, 75-80] DNAJC6 works with multiple other genes linked to familial and sporadic PD, including *PARK20* (SYNJ1), *PARK8* (LRRK2), *PARK2* (PRKN), *SNCA* (α-synuclein), and other DNAJC proteins (*DNAJC26*, RAK; *DNAJC13,* REM8), to regulate key steps in synaptic vesicle trafficking, synaptic vesicle endocytosis, and clathrin-dynamics in endocytosis.[80-84] While mutations in this gene have only been associated with rare familial forms of PD, these data potentially links epigenetic regulation of this locus with the pathogenesis of sPD and are consistent with previous findings from our lab and others demonstrating epigenetic changes in familial and GWAS PD genes in brain tissue from sPD cases. [10, 36-39, 44]

PD-associated interaction DMCs are enriched in pathways related to synaptic formation and function, senescence, neuroinflammation, and epigenetic regulation

Within the 695 iDMC-containing genes, our functional annotation found that these gene products function within specific pathways related to neuronal dysfunction in PD (Figure 2, Table 2). These pathways are described here with STRING subnetworks in Figure 2B indicated by their numbers; expanded connections with additional iDMC-containing genes are shown in Supplementary Figure 2.

Multiple iDMC-containing genes encode proteins that converge on functions related to proper development and function of synapses, including synaptogenesis, synaptic plasticity, neurotransmitter release, and clathrin-mediated endocytosis (2,4,7,8,9,10,13,14). In addition, a

set of genes play roles in the myelination and maintenance of myelin (3,5), as well as in the synthesis of dopamine, serotonin, and norepinephrine (4). Together, this suggests the regulation of genes related to the formation of synapses and the maintenance of synaptic and vesicular integrity are disrupted prior to the onset of pathology within the parietal cortex. We observed epigenetic changes in similar key neurodevelopmental pathways across the lifespan in our mouse model of increased PD susceptibility.[85] Together, these pathways coordinate the formation, structure, function, and plasticity of synapses, consistent with the idea that the integrity of synapses is critical for proper neurotransmission and that multiple PD-related mechanisms converge on the disruption of synaptic function.[86-89] This is also consistent with findings in our mouse model of increased PD susceptibility, showing that exposure to the organochlorine pesticide dieldrin is associated with epigenetic changes in similar key neurodevelopmental pathways across the lifespan.[85]

A large body of evidence now supports a role of neuroinflammatory pathways in PD.[90-92] Here, we identified multiple iDMC-containing genes important in the neuronal response to environmental stressors and neuroinflammatory signals, including multiple genes within the NFκB signaling pathway (3,11). Altered epigenetic regulation of these genes could lead to impropriate activation of these pathways in response to cellular and environmental stressors. We also identified genes involved in cell cycle and senescence pathways (1,12), consistent with evidence that neuronal cell cycle reentry events and cellular senescence in the aging brain can lead to degeneration. [93-95] Finally, we observed epigenetic regulation of multiple epigenetic regulators themselves (6).

While these changes were not assessed in the SN, the region most commonly studied in the context of PD, by using the parietal cortex, which does not have widespread degeneration, we were able to assess neuron-specific DNA modifications associated with PD in a region without widespread neuron loss prior to the onset of pathology within the parietal cortex. Despite this, overall, these data suggest that PD-associated alterations in the epigenetic regulation of these genes may alter gene expression, promoter usage, or isoform expression of these genes and may represent early events that precede the onset of degeneration.

Technical limitations of the oxBS method

In addition, it is also important to note that this study included a high level of failed probes and samples that was unique to the oxBS reactions. In contrast, no samples were excluded due to high levels of failed probes, and far fewer probes failed from the BS data. As noted in the methods, we started with 1 µg of input DNA for oxBS because when we used the recommended starting amount of 500 ng, all oxBS probes failed. Together, this suggests that oxBS is much harsher on the DNA, requiring high input amounts that may limit the utility of this method as the field moves towards cell type-specific methods and lower input amounts.

Data Availability

This study was preregistered with Open Science Framework: https://osf.io/z4vbw.

Raw and processed data will be available in GEO (GSE267937)

Supplementary material availability

All supplementary material, including additional figures, tables of results, and code used for analyses, are available as supplementary files.

Acknowledgements

The authors thank the Van Andel Genomics Core, especially Marie Adams, for providing consultation, library preparation, and next-generation sequencing facilities and services (Illumina EPIC array). We are grateful to the Banner Sun Health Research Institute Brain and Body Donation Program of Sun City, Arizona for the provision of human brain tissue.

CRediT Author Statement

Juliana I. Choza: Formal Analysis, Visualization, Writing – original draft, Writing – review and editing; **Mahek Virani:** Visualization**,** Data Curation; Writing – review and editing; **Nathan C. Kuhn:** Investigation, Software, Formal Analysis; **Marie Adams:** Investigation, Methodology; **Joseph Kochmanski:** Conceptualization, Formal Analysis, Investigation, Methodology, Writing – review and editing; **Alison I. Bernstein:** Conceptualization, Methodology, Visualization, Data Curation, Writing - Review & Editing, Supervision, Project Administration, Funding Acquisition

	Male (n = 56)	
Variables	**Mean ± SD or**	**Range**
Disease Status		
Control	29 (51.8%)	
Parkinson's disease	27 (48.2%)	
Age at Death		
Control	79.3 ± 9.1	53-93
Parkinson's disease	79.4 ± 7.1	64-91
PMI		
Control	3.25 ± 0.81	2.16-5.5
Parkinson's disease	3.27 ±0.83	1.83-4.92
Race		
White	55 (98.2%)	
Unidentified	1 (1.8%)	

Table 1: Cohort characteristics including disease status, age at death in years, postmortem interval (PMI) in hours, and race of samples remaining after QC. Seven male samples and eleven female samples were removed during quality control and pre-processing, leaving 56 male and 26 female samples.

Figure 1: Identification of PD-associated DMCs. Frequency histograms of raw BS and oxBS β-values (A) and MLE-corrected β_{mC} and β_{hmC} (B). (C) Volcano plots of DMCs derived from MLE-corrected β values input into differential modification analysis (FDR < 0.05). Red indicates an increased interaction term; blue indicates a decreased interaction term. (D,E) Frequency histograms showing annotations of significant iDMCs for intragenic annotations (D) and universal chromatin state annotations (E). (F-J) MLE-corrected β vales graphed by modification (top) and raw BS β values with MLE-corrected β values graphed by disease status (bottom) for the most significant iDMC with a positive interaction term (F), the most significant iDMC with a negative interaction term (G), the iDMC with the most positive interaction term (H), the iDMC with the most negative interaction term (I) and the iDMC annotated to the PD gene *PARK19/DNAJC6* (J). The corresponding beta coefficient/interaction term and FDR value are indicated for each iDMC.

Figure 2: Gene ontology and protein interaction networks for DMC-containing genes. (A) Gene ontology network from ClueGO shows enriched groups of GO terms. Bolded text indicates the most significant GO term in each group. Grouped GO terms share at least 50% of their genes. (B) String protein interaction network for 374 unique genes (confidence < 0.9; >2 genes per network). All but 1 gene mapped to the STRING database (RTEL1-TNFRSF6B). PPI enrichment value = 5.91 x 10^{-6}. Individual subnetworks are numbered for reference within the text.

GO Groups	GO Terms	Associated Genes Found
Group0	L-histidine transmembrane transport	SLC15A4, SLC25A29, SLC38A3, SLC7A1
Group1	anatomical structure morphogenesis	ABCA2, ACAN, ACTN2, AGRN, ALPL, ALX4, ANO1, APOA1, ARHGEF15, ARX, ATP10A, BARX2, BCR, BMP1, CACNA1C, CACNA1H, CASQ1, CCBE1, CCDC141, CCNA2, CD34, CDH11, CDH13, CDH23, CDH4, CLRN2, CNMD, COL22A1, COL27A1, COL4A3, CSF1R, CSNK2B, CTSZ, DCHS1, DDR1, DIO3, DISC1, DLL1, DNM3, DOCK2, DSCAML1, EDAR, EFNA5, EGFLAM, EIF2AK4, ELN, EMX1, EPHB4, ETS1, F11R, FAM20C, FAT3, FBXW8, FGFR2, FKBP10, FOXG1, FRMD6, FZD1, GAA, GAS7, GLI2, GNAS, HAND1, HGS, HOXA3, HS3ST3B1, HSBP1, IGF2BP2, INF2, IRX2, ITGB5, KCNQ1, KDM5B, LAMA2, LIMD1, LRIG3, LTBP3, LYVE1, MAP2K5, MAP3K13, MBP, MECP2, MRTFA, MYH3, MYMK, MYOD1, NCAM1, NFKB1, NYAP1, OBSCN, OTX1, PAX6, PCDHGB4, PLEKHA1, PLXNA2, PLXND1, POFUT1, PPFIA2, PRAG1, RAB23, RADIL, RB1, RERE, RFX2, RHOD, RIPOR2, ROR2, SASH1, SDK1, SKI, SMOC2, SOX15, SOX2, SP100, SPARC, SS18L1, ST6GAL1, STAB1, STAT1, STK24, SYT2, TAFAZZIN, TERT, TFAP2A, TGFB1, TIAM2, TLR3, TMEM100, TNFSF11, TNR, TOPORS, TSPEAR, VAV2, VSX1, WNT3, WNT9A, WWOX, ZFP36L1
Group2	cell differentiation	ABCA2, ACAN, ACTN2, AGRN, AKNA, ALPL, ANO1, APC2, APOA1, ARX, ASAP1, ATP11A, BARX2, BCR, BFSP1, BICRA, BMP1, BPGM, BRINP2, BTBD2, C9orf24, CACNA1H, CASQ1, CCDC141, CCNA2, CD34, CD38, CDH1, CDH11, CDH23, CDH4, CDYL, CEBPE, CEP85L, CFAP157, CLRN2, CNMD, COL22A1, COL27A1, CSF1R, CSF3R, CSK, DCHS1, DDR1, DIAPH2, DIO3, DISC1, DLK1, DLL1, DNASE1L3, DNM3, DOCK2, DPF2, DSCAML1, DYSF, EDAR, EFNA5, EID2B, EIF2AK4, EMX1, EPHB4, ERG, ETS1, F11R, F2RL1, FAM20C, FAT3, FBXW8, FCER1G, FGFR2, FOXO6, FRMD6, FZD1, GAK, GAS7, GCNT2, GLI2, GPAT4, GRM5, HAND1, HDAC4, HSBP1, IL18R1, IL4I1, IRF4, IRF7, IRX2, ITGA11, ITGB5, KCNQ1, KDM5B, KRT8, LAMA2, LDLRAD4, LIMD1, LTBP3, MAP2K6, MAP3K13, MAP3K4, MBP, MECP2, MLH1, MMD2, MRTFA, MTURN, MYH3, MYMK, MYOD1, NAV2, NCAM1, NEU4, NFKB1, NFKBID, NHSL2, NKX6-3, NPHP4, NTM, NTRK3, NUMA1, NXN, NYAP1, OBSCN, OSBP2, PAX6, PIEZO1, PIWIL3, PLCB1, PLEKHA1, PLXNA2, PLXND1, POU6F2, PPFIA2, PRAG1, PRDM16, PRKG1, PRLR, PRM2, PRM3, PRRC2A, PRRC2C, RAB32, RADIL, RAPGEF1, RB1, RERE, RFX2, RIPOR2, ROR2, RTN4RL1, SCUBE1, SDK1, SKI, SLC12A5, SLC4A5, SLCO4C1, SMOC1, SOCS2, SOX15, SOX2, SPAG16, SS18L1, ST6GAL1, STAT1, STK24, SYNE2, SYT2, TBX21, TBX22, TCF7L1, TERT, TESMIN, TFAP2A, TGFB1, TIAM2, TLR3, TMEM100, TMEM91, TNFSF11, TNR, TOPORS, TP73, TRAPPC9, TST, TYK2, VCAN, VSX1, WDR38, WNT3, WNT9A, WWOX, ZBTB7B, ZFP36L1
Group3	multicellular organism development, system development, nervous system development	AATK, ABCA2, ACAN, ACTN2, AGRN, AK8, AKNA, ALPL, ALX4, ANKLE2, ANO1, APBA3, APC2, APOA1, ARHGEF15, ARNT2, ARX, ASAP1, ATN1, ATP11A, BARX2, BCR, BFSP1, BMP1, BPGM, BRINP2, BTBD2, CACNA1C, CCBE1, CCDC141, CCNA2, CD34, CD38, CDH1, CDH11, CDH13, CDH23, CDH4, CEBPE, CEP85L, CLRN2, CNMD, COL22A1, COL27A1, COL4A3, CRYGN, CSF3R, CSK, CSNK2B, CTNS, CTSZ, DCHS1, DDC, DDR1, DHX35, DIO3, DISC1, DLL1, DNAJB1, DNM3, DOCK2, DPF2, DSCAML1, EDAR, EFNA5, EIF2AK4, ELN, EMX1, EPHB4, ETS1, F11R, F2RL1, FAM20C, FAT3, FBXW8, FCER1G, FGFR2, FKBP10, FOXG1, FOXO6, FZD1, GAA, GAK, GAS7, GCNT2, GLI2, GNAS, GRM5, HAND1, HDAC4, HGS, HOXA3, HS3ST3B1, HSBP1, HSD17B3, IGF2BP2, IL18R1, IL4I1, INPP5K, IRF4, IRF7, IRX2, ITGB5, KCNQ1, KDM5B, KIF26A, LAMA2, LRIG3, LTBP3, MAD1L1, MALL, MAOB, MAP2K5, MAP2K6, MAP3K13, MAP3K4, MBP, MECP2, MLH1, MMD2, MPST, MRTFA, MTURN, MYH3, MYOD1, NAV2, NCAM1, NEU4, NFKBID, NKX6-3, NPHP4, NRXN2, NTM, NTRK3, NUMA1, NXN, NYAP1, OTX1, PAX6, PCDHGA1, PCDHGA10, PCDHGA2, PCDHGA3, PCDHGA4, PCDHGA5, PCDHGA6, PCDHGA7, PCDHGA8, PCDHGA9, PCDHGB1, PCDHGB2, PCDHGB3, PCDHGB5, PCDHGB6, PDCD1, PDE6B, PDGFD, PKD1L1, PLCB1, PLCE1, PLEKHA1, PLXNA2, PLXND1, POFUT1, POU6F2, PPFIA2, PRAG1, PRDX1, PRKG1, PRLR, PRRC2C, RAB23, RAPGEF1, RB1, RBM19, RD3, RERE, RGS7, RIPOR2, ROR2, RTN4RL1, SASH1, SBF2, SCUBE1, SDK1, SKI, SLC12A5, SLC15A4, SLC34A2, SLC38A3, SLC4A5, SMOC1, SMOC2, SOCS2, SOX15, SOX2, SP100, SPARC, SS18L1, STAB1, STAT1, STK24, SYNE2, SYNJ2, SYT2, TAFAZZIN, TBX21, TCF7L1, TEAD1, TERT, TFAP2A, TGFB1, TIAM2, TLR3, TMEM100, TMEM91, TNFSF11, TNR, TOPORS, TP73, TRAPPC9, TRIM26, UMODL1, VAV2, VCAN, VSX1, WDR38, WNT3, WNT9A, WWOX, XYLT1, ZBTB7B, ZFP36L1
Group4	regulation of GTPase activity, regulation of small GTPase mediated signal	ABR, AGAP1, AGRN, APC2, APOA1, ARHGAP42, ARHGAP45, ARHGAP6, ARHGEF15, ARHGEF4, ASAP1, BCR, BMP1, CGNL1, CHN2, DENND1A, DENND1C, DOCK2, DOCK6, EFNA5, F11R, F2RL1, GNAS, HERC2, INF2, MCF2L, MCF2L2,

	transduction, guanyl-nucleotide exchange factor activity	MMD2, NTRK3, NUP62, OBSCN, PLCE1, PLEKHG4B, PLEKHG7 PLXNA2, PLXND1, PRAG1, PRKG1, RAPGEF1, RGS7, RHOD, RIPOR2, SBF2, SGSM2, SLC6A5, TBC1D10B, TIAM2, VAV2
Group5	cell-cell adhesion, cell-cell adhesion via plasma-membrane adhesion molecules, homophilic cell adhesion via plasma membrane adhesion molecules	AKNA, APOA1, ARVCF, CCNA2, CD34, CD93, CDH1, CDH11, CDH13, CDH23, CDH4, CLDN10, CLDN2, CSK, DCHS1, DSCAML1, EFNA5, ELANE, ETS1, F11R, FAT3, GCNT2, GLI2, GNAS, IGDCC4, IL4I1, ITGA11, ITGB5, MAD1L1, MAP2K5, MBP, NFKBID, NPHP4, NRXN2, OBSCN, PCDHGA1, PCDHGA10, PCDHGA2, PCDHGA3, PCDHGA4, PCDHGA5, PCDHGA6, PCDHGA7, PCDHGA8, PCDHGA9, PCDHGB1, PCDHGB2, PCDHGB3, PCDHGB4, PCDHGB5, PCDHGB6, PIEZO1, PKD1L1, PKP3, PPM1F, PRKG1, RAPGEF1, RIPOR2, SDK1, SLC7A1, SOX2, TBX21, TGFB1, TNFSF11, TNR, TYK2, ZBTB7B, ZFP36L1
Group6	regulation of signaling, regulation of response to stimulus, regulation of cell communication, regulation of signal transduction	ABCB1, ABR, ADGRE2, AGPAT2, AKNA, ANO1, APC2, APOA1, ARHGAP42, ARHGAP45, ARHGAP6, ARHGEF15, ARHGEF4, BCR, C10orf71, CAMTA1, CASQ1, CCBE1, CCNA2, CCND3, CD34, CD38, CD59, CDH1, CDH11, CDH13, CDK14, CFB, CFD, CFHR5, CGNL1, CHN2, CIDEA, CLEC16A, CLPB, CRTC3, CSF1R, CSK, CSNK2B, CTNS, DAPK2, DENND1A, DISC1, DLGAP4, DLK1, DLL1, DNAJB1, DNASE1L3, DOCK2, DPF2, EDAR, EFNA5, EIF2AK4, ELANE, ETS1, F11R, F2RL1, FAM107A, FAM20C, FCER1G, FGFR2, FGFR4, FRMD8, FZD1, GCNT2, GCSAML, GLI2, GNAS, GRAMD4, GRID1, GRM4, GRM5, GTF2IRD1, HDAC4, HEBP2, HGS, IL18R1, IL4I1, INF2, INPP5K, IRAK2, IRF4, IRF7, JPH3, KCTD6, KDM5B, KIF26A, KLHL15, LAMA2, LDLRAD4, LIMD1, LRPAP1, MAD1L1, MAGI3, MAP2K5, MAP2K6, MAP3K1, MAP3K13, MAP3K4, MAP4K2, MCF2L, MECP2, MLH1, MMD2, MRTFA, MTURN, MYMK, MYOD1, NCAM1, NCOA7, NFKB1, NFKBID, NFKBIL1, NLRP7, NPHP4, NTRK3, NTSR1, NUMA1, NUP62, NXN, OBSCN, P2RY6, PDCD1, PDGFD, PFKL, PLCB1, PLCE1, PLEKHA1, PLEKHF1, PLEKHG4B, PMEPA1, POFUT1, PPM1F, PRAG1, PRDM16, PRDX1, PRKG1, PRLR, RAPGEF1, RB1, RGS7, RHOD, RIPOR2, RNF126, RNF185, ROR2, RPTOR, RTEL1-TNFRSF6B, RTEL1, RTN4RL1, SASH1, SCUBE1, SETD4, SHISA7, SIGIRR, SKI, SLC15A4, SLC6A5, SMOC2, SOCS2, SOX15, SOX2, SP100, SREBF2, SSTR5, ST6GAL1, STAT1, STK24, STK39, STX4, STXBP2, TBX21 TCF7L1, TEDC2, TERT, TGFB1, TIAM2, TLR3, TMEM100, TNFSF11, TNR, TP73, TRABD2B, TRAP1, TRIM33, TSPEAR, TYK2, UBR5, USP44, VAV2, VWF, WNT3, WWOX, ZBTB7B, ZFYVE28, ZNF366

Table 2: Enriched GO Terms and GO Term Groups of interaction DMC-containing genes. Results of ClueGO gene ontology enrichment analysis and significant GO terms are shown (p < 0.05). GO terms are grouped when they share >50% of their genes. "Associated Genes" shows all interaction DMC containing genes that map to each GO Term Group.

	5hmC	5mC and 5hmC (BS)							Paired BS/oxBS
	Marshall 2020	**Kia 2021**		**Masliah 2013**	**Young 2019**			**Kochmanski 2022**	**Current study**
Control (M/F)	23*	not specified		6 (2/4)	41 (all male)			49 (29/20)	29 (all male)
PD (M/F)	20*	134*		5 (5/0)	38 (all male)			50 (33/17)	27 (all male)
Region	PFC	SN	CTX	FC	DMV	CG	SN	PC	PC
Method	hMe-DIP	BS-450K		BS-450K	BS-450K/EPIC			BS-EPIC	BS/oxBS-EPIC
# genes	5157	154	125	155	203	119	1466	523	695
Overlap with BS	121	4	8	12	19	14	67	-	43
Overlap with interaction	173	6	11	14	20	8	105	43	-

Table 3: Summary of comparisons between the current study and previous PD EWAS in brain tissue. *Sex not specified, Abbreviations: SN, substantia nigra; PFC, prefrontal cortex; CTX, cortex; FC, frontal cortex; DMV, dorsal motor nucleus of the vagus; CG, cingulate gyrus; PC, parietal corte

References Cited

1. Keller MF, Saad M, Bras J, Bettella F, Nicolaou N, Simón-Sánchez J, Mittag F, Büchel F, Sharma M, Gibbs JR, Schulte C, Moskvina V, Durr A, Holmans P, Kilarski LL, Guerreiro R, Hernandez DG, Brice A, Ylikotila P, Stefánsson H, Majamaa K, Morris HR, Williams N, Gasser T, Heutink P, Wood NW, Hardy J, Martinez M, Singleton AB, Nalls MA, Plagnol V, Sheerin UM, Lesage S, Sveinbjörnsdóttir S, Arepalli S, Barker R, Ben-Shlomo Y, Berendse HW, Berg D, Bhatia K, Bie RMAd, Biffi A, Bloem B, Bochdanovits Z, Bonin M, Brockmann K, Brooks J, Burn DJ, Charlesworth G, Chen H, Chinnery PF, Chong S, Clarke CE, Cookson MR, Cooper JM, Corvol JC, Counsell C, Damier P, Dartigues JF, Deloukas P, Deuschl G, Dexter DT, Dijk KDv, Dillman A, Durif F, Dürr A, Edkins S, Evans JR, Foltynie T, Gao J, Gardner M, Goate A, Gray E, Gústafsson Ó, Harris C, Hilten JJv, Hofman A, Hollenbeck A, Holton J, Hu M, Huang X, Huber H, Hudson G, Hunt SE, Huttenlocher J, Illig T, Jónsson PV, Lambert JC, Langford C, Lees A, Lichtner P, Limousin P, Lopez G, Lorenz D, McNeill A, Moorby C, Moore M, Morrison KE, Mudanohwo E, O'Sullivan SS, Pearson J, Perlmutter JS, Pétursson H, Pollak P, Post B, Potter SC, Ravina B, Revesz T, Riess O, Rivadeneira F, Rizzu P, Ryten M, Sawcer SJ, Schapira A, Scheffer H, Shaw K, Shoulson I, Sidransky E, Smith C, Spencer CCA, Steinberg S, Stockton JD, Strange A, Talbot K, Tanner CM, Tashakkori-Ghanbaria A, Tison F, Trabzuni D, Traynor BJ, Uitterlinden AG, Velseboer D, Vidailhet M, Walker R, Warrenburg Bvd, Wickremaratchi M, Williams-Gray CH, Winder-Rhodes S, Stefánsson K, Donnelly P, Barroso I, Blackwell JM, Bramon E, Brown MA, Casas JP, Corvin A, Duncanson A, Jankowski J, Markus HS, Mathew CG, Palmer CNA, Plomin R, Rautanen A, Trembath RC, Viswanathan AC, Band G, Bellenguez C, Freeman C, Hellenthal G, Giannoulatou E, Pirinen M, Pearson R, Su Z, Vukcevic D, Gwilliam R, Blackburn H, Bumpstead SJ, Dronov S, Gillman M, Hammond N, Jayakumar A, McCann OT, Liddle J, Ravindrarajah R, Ricketts M, Waller M, Weston P, Widaa S, Whittaker P, McCarthy MI. (2012) "Using Genome-Wide Complex Trait Analysis to Quantify 'Missing Heritability' in Parkinson's Disease". *Human molecular genetics.* 21(22):4996-5009. DOI:10.1093/HMG/DDS335.

2. Lill CM. (2016) "Genetics of Parkinson's Disease". *Molecular and cellular probes.* 30(6):386-396. DOI:10.1016/j.mcp.2016.11.001.

3. Trinh J, Farrer M. (2013) "Advances in the Genetics of Parkinson Disease". *Nature Reviews Neurology.* 9(8):445-454. DOI:10.1038/nrneurol.2013.132.

4. Dorsey ER, Sherer T, Okun MS, Bloem BR. (2018) "The Emerging Evidence of the Parkinson Pandemic". *Journal of Parkinson's Disease.* 8(s1):S3-S8. DOI:10.3233/JPD-181474.

5. Faulk C, Dolinoy DC. (2011) "Timing is Everything: The when and how of Environmentally Induced Changes in the Epigenome of Animals.". *Epigenetics.* 6(7):791-797. DOI:10.4161/epi.6.7.16209.

6. Allis CD, Jenuwein T. (2016) "The Molecular Hallmarks of Epigenetic Control". *Nature Reviews Genetics.* 17(8):487-500. DOI:10.1038/nrg.2016.59.

7. Cavalli G, Heard E. (2019) "Advances in Epigenetics Link Genetics to the Environment and Disease". *Nature.* 571(7766):489-499. DOI:10.1038/s41586-019-1411-0.

8. Wu H, Eckhardt CM, Baccarelli AA. (2023) "Molecular Mechanisms of Environmental Exposures and Human Disease". *Nature reviews.Genetics.* 24(5):332-344. DOI:10.1038/S41576-022-00569-3.

9. Boyce WT, Sokolowski MB, Robinson GE. (2020) "Genes and Environments, Development and Time". *Proceedings of the National Academy of Sciences.* 117(38):23235-23241. DOI:10.1073/pnas.2016710117.

10. Schaffner SL, Kobor MS. (2022) "DNA Methylation as a Mediator of Genetic and Environmental Influences on Parkinson's Disease Susceptibility: Impacts of Alpha-Synuclein, Physical Activity, and Pesticide Exposure on the Epigenome". *Frontiers in genetics.* 13:971298. DOI:10.3389/FGENE.2022.971298.

11. Jakovcevski M, Akbarian S. (2012) "Epigenetic Mechanisms in Neurological Disease". *Nature medicine.* 18(8):1194-204. DOI:10.1038/nm.2828.

12. Marques S, Outeiro TF. (2013) "Epigenetics in Parkinson's and Alzheimer's Diseases". *Sub-cellular biochemistry.* 61:507-25. DOI:10.1007/978-94-007-4525-4_22.

13. Miranda-Morales E, Meier K, Sandoval-Carrillo A, Salas-Pacheco J, Vazquez-Cardenas P, Arias-Carrion O. (2017) "Implications of DNA Methylation in Parkinson's Disease". *Front Mol Neurosci.* 10:225. DOI:10.3389/fnmol.2017.00225.

14. Lardenoije R, Pishva E, Lunnon K, van den Hove DL. (2018) "Neuroepigenetics of Aging and Age-Related Neurodegenerative Disorders". *Prog Mol Biol Transl Sci.* 158:49-82. DOI:10.1016/bs.pmbts.2018.04.008.

15. Labbe C, Lorenzo-Betancor O, Ross OA. (2016) "Epigenetic Regulation in Parkinson's Disease". *Acta Neuropathol.* 132(4):515-530. DOI:10.1007/s00401-016-1590-9.

16. Wullner U, Kaut O, DeBoni L, Piston D, Schmitt I. (2016) "DNA Methylation in Parkinson's Disease". *J Neurochem.* 139 Suppl:108-120. DOI:10.1111/jnc.13646.

17. van Heesbeen HJ, Smidt MP. (2019) "Entanglement of Genetics and Epigenetics in Parkinson's Disease". *Frontiers in neuroscience.* 13:277. DOI:10.3389/fnins.2019.00277.

18. Tsalenchuk M, Gentleman SM, Marzi SJ. (2023) "Linking Environmental Risk Factors with Epigenetic Mechanisms in Parkinson's Disease". *npj Parkinson's Disease.* 9(1):123. 37626097. DOI:10.1038/s41531-023-00568-z.

19. Smith ZD, Meissner A. (2013) "DNA Methylation: Roles in Mammalian Development". *Nature Reviews Genetics 2013 14:3.* 14(3):204-220. DOI:10.1038/nrg3354.

20. Bachman M, Uribe-Lewis S, Yang X, Williams M, Murrell A, Balasubramanian S. (2014) "5-Hydroxymethylcytosine is a Predominantly Stable DNA Modification". *Nature chemistry.* 6(12):1049-55. DOI:10.1038/nchem.2064.

21. Nestor CE, Ottaviano R, Reddington J, Sproul D, Reinhardt D, Dunican D, Katz E, Dixon JM, Harrison DJ, Meehan RR. (2012) "Tissue Type is a Major Modifier of the 5-Hydroxymethylcytosine Content of Human Genes". *Genome research.* 22(3):467-477. DOI:10.1101/gr.126417.111.

22. Cheng Y, Bernstein AI, Chen D, Jin P. (2014) "5-Hydroxymethylcytosine: A New Player in Brain Disorders?". *Experimental neurology.* 268:3-9. DOI:10.1016/j.expneurol.2014.05.008.

23. Chen Y, Damayanti NP, Irudayaraj J, Dunn K, Zhou FC. (2014) "Diversity of Two Forms of DNA Methylation in the Brain". *Frontiers in genetics.* 5:46. DOI:10.3389/fgene.2014.00046.

24. Globisch D, Münzel M, Müller M, Michalakis S, Wagner M, Koch S, Brückl T, Biel M, Carell T. (2010) "Tissue Distribution of 5-Hydroxymethylcytosine and Search for Active Demethylation Intermediates". *PLoS ONE.* 5(12) DOI:10.1371/journal.pone.0015367.

25. Szulwach KE, Li X, Li Y, Song C, Wu H, Dai Q, Irier H, Upadhyay AK, Gearing M, Levey AI, Vasanthakumar A, Godley La, Chang Q, Cheng X, He C, Jin P. (2011) "5-hmC–mediated Epigenetic Dynamics during Postnatal Neurodevelopment and Aging". *Nature neuroscience.* 14(12):1607-1616. DOI:10.1038/nn.2959.

26. Khare T, Pai S, Koncevicius K, Pal M, Kriukiene E, Liutkeviciute Z, Irimia M, Jia P, Ptak C, Xia M, Tice R, Tochigi M, Moréra S, Nazarians A, Belsham D, Wong AHC, Blencowe BJ, Wang SC, Kapranov P, Kustra R, Labrie V, Klimasauskas S, Petronis A. (2012) "5-hmC in the Brain is Abundant in Synaptic Genes and shows Differences at the Exon-Intron Boundary". *Nature structural & molecular biology.* 19(10):1037-1044. DOI:10.1038/NSMB.2372.

27. Wen L, Li X, Yan L, Tan Y, Li R, Zhao Y, Wang Y, Xie J, Zhang Y, Song C, Yu M, Liu X, Zhu P, Li X, Hou Y, Guo H, Wu X, He C, Li R, Tang F, Qiao J. (2014) "Whole-Genome Analysis of 5-Hydroxymethylcytosine and 5-Methylcytosine at Base Resolution in the Human Brain". *Genome biology.* 15(3):R49. DOI:10.1186/gb-2014-15-3-r49.

28. Mellén M, Ayata P, Heintz N. (2017) "5-Hydroxymethylcytosine Accumulation in Postmitotic Neurons Results in Functional Demethylation of Expressed Genes". *Proceedings of the National Academy of Sciences.* 114(37):E7812-E7821. 28847947. DOI:10.1073/pnas.1708044114.

29. Spruijt CG, Gnerlich F, Smits AH, Pfaffeneder T, Jansen PWTC, Bauer C, Münzel M, Wagner M, Müller M, Khan F, Eberl HC, Mensinga A, Brinkman AB, Lephikov K, Müller U, Walter Jö, Boelens R, Van Ingen H, Leonhardt H, Carell T, Vermeulen M. (2013)

"Dynamic Readers for 5-(Hydroxy)Methylcytosine and its Oxidized Derivatives". *Cell.* 152(5):1146-1159. DOI:10.1016/j.cell.2013.02.004.

30. Kochmanski J, Bernstein AI. (2020) "The Impact of Environmental Factors on 5-Hydroxymethylcytosine in the Brain". *Current environmental health reports.* 7(2):109-120. 32020534.PMC7809708. DOI:10.1007/s40572-020-00268-3.

31. Al-mahdawi S, Virmouni SA, Pook Ma, Evans-galea M. (2014) "The Emerging Role of 5-Hydroxymethylcytosine in Neurodegenerative Diseases". 8:1-8. DOI:10.3389/fnins.2014.00397.

32. Jin S, Kadam S, Pfeifer GP. (2010) "Examination of the Specificity of DNA Methylation Profiling Techniques Towards 5-Methylcytosine and 5-Hydroxymethylcytosine". *Nucleic acids research.* 38(11):e125. DOI:10.1093/nar/gkq223.

33. Bernstein AI, Jin P. High-throughput sequencing-based mapping of cytosine modifications. In: Epigenetic Technological Applications. ; 2015.

34. Young JI, Sivasankaran SK, Wang L, Ali A, Mehta A, Davis DA, Dykxhoorn DM, Petito CK, Beecham GW, Martin ER, Mash DC, Pericak-Vance M, Scott WK, Montine TJ, Vance JM. (2019) "Genome-Wide Brain DNA Methylation Analysis Suggests Epigenetic Reprogramming in Parkinson Disease". *Neurology. Genetics.* 5(4):e342. 31403079.PMC6659138. DOI:10.1212/NXG.0000000000000342.

35. Masliah E, Dumaop W, Galasko D, Desplats P. (2013) "Distinctive Patterns of DNA Methylation Associated with Parkinson Disease: Identification of Concordant Epigenetic Changes in Brain and Peripheral Blood Leukocytes". *Epigenetics.* 8(10):1030-1038. 23907097.PMC3891683. DOI:10.4161/epi.25865.

36. Jowaed A, Schmitt I, Kaut O, Wüllner U. (2010) "Methylation Regulates Alpha-Synuclein Expression and is Decreased in Parkinson's Disease Patients' Brains". *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience.* 30(18):6355-6359. 20445061.PMC6632710. DOI:10.1523/JNEUROSCI.6119-09.2010.

37. Li P, Ensink E, Lang S, Marshall L, Schilthuis M, Lamp J, Vega I, Labrie V. (2020) "Hemispheric Asymmetry in the Human Brain and in Parkinson's Disease is Linked to Divergent Epigenetic Patterns in Neurons". *Genome Biology.* 21(1):61. 32151270. DOI:10.1186/s13059-020-01960-1.

38. Kia DA, Zhang D, Guelfi S, Manzoni C, Hubbard L, Reynolds RH, Botía J, Ryten M, Ferrari R, Lewis PA, Williams N, Trabzuni D, Hardy J, Wood NW. (2021) "Identification of Candidate Parkinson Disease Genes by Integrating Genome-Wide Association Study, Expression, and Epigenetic Data Sets". *JAMA neurology.* 78(4):464-472. 33523105.PMC7851759. DOI:10.1001/jamaneurol.2020.5257.

39. Go RCP, Corley MJ, Ross GW, Petrovitch H, Masaki KH, Maunakea AK, He Q, Tiirikainen MI. (2020) "Genome-Wide Epigenetic Analyses in Japanese Immigrant Plantation Workers with Parkinson's Disease and Exposure to Organochlorines Reveal Possible Involvement of Glial Genes and Pathways Involved in Neurotoxicity". *BMC neuroscience.* 21(1):31. 32650713. DOI:10.1186/s12868-020-00582-4.

40. Kaut O, Schmitt I, Stahl F, Fröhlich H, Hoffmann P, Gonzalez FJ, Wüllner U. (2022) "Epigenome-Wide Analysis of DNA Methylation in Parkinson's Disease Cortex". *Life.* 12(4):502. DOI:10.3390/life12040502.

41. Shu L, Qin L, Min S, Pan H, Zhong J, Guo J, Sun Q, Yan X, Chen C, Tang B, Xu Q. (2019) "Genetic Analysis of DNA Methylation and Hydroxymethylation Genes in Parkinson's Disease". *Neurobiology of aging.* 84:242.e13-242.e16. DOI:10.1016/J.NEUROBIOLAGING.2019.02.025.

42. Min S, Xu Q, Qin L, Li Y, Li Z, Chen C, Wu H, Han J, Zhu X, Jin P, Tang B. (2022) "Altered Hydroxymethylome in the Substantia Nigra of Parkinson's Disease". *Human molecular genetics.* 31(20):3494-3503. 35661211. DOI:10.1093/hmg/ddac122.

43. Marshall LL, Killinger BA, Ensink E, Li P, Li KX, Cui W, Lubben N, Weiland M, Wang X, Gordevicius J, Coetzee GA, Ma J, Jovinge S, Labrie V. (2020) "Epigenomic Analysis of Parkinson's Disease Neurons Identifies Tet2 Loss as Neuroprotective". *Nature neuroscience.* 23(10):1203-1214. DOI:10.1038/s41593-020-0690-y.

44. Kochmanski J, Kuhn NC, Bernstein AI. (2022) "Parkinson's Disease-Associated, Sex-Specific Changes in DNA Methylation at PARK7 (DJ-1), SLC17A6 (VGLUT2), PTPRN2 (IA-2β), and NR4A2 (NURR1) in Cortical Neurons". *npj Parkinson's Disease.* 8(1):120. DOI:10.1038/s41531-022-00355-2.

45. Kochmanski J, Savonen C, Bernstein AI. (2019) "A Novel Application of Mixed Effects Models for Reconciling Base-Pair Resolution 5-Methylcytosine and 5-Hydroxymethylcytosine Data in Neuroepigenetics". *Frontiers in Genetics.* 10:801. DOI:10.3389/fgene.2019.00801.

46. Petroff RL, Cavalcante RG, Langen ES, Dolinoy DC, Padmanabhan V, Goodrich JM. (2023) "Mediation Effects of DNA Methylation and Hydroxymethylation on Birth Outcomes After Prenatal Per- and Polyfluoroalkyl Substances (PFAS) Exposure in the Michigan Mother-Infant Pairs Cohort". *Clinical epigenetics.* 15(1) DOI:10.1186/S13148-023-01461-5.

47. Rygiel CA, Goodrich JM, Solano-González M, Mercado-García A, Hu H, Téllez-Rojo MM, Peterson KE, Dolinoy DC. (2021) "Prenatal Lead (Pb) Exposure and Peripheral Blood DNA Methylation (5mC) and Hydroxymethylation (5hmC) in Mexican Adolescents from the ELEMENT Birth Cohort". *Environmental health perspectives.* 129(6):67002. DOI:10.1289/EHP8507.

48. Booth MJ, Branco MR, Ficz G, Oxley D, Krueger F, Reik W, Balasubramanian S. (2012) "Quantitative Sequencing of 5-Methylcytosine and 5-Hydroxymethylcytosine at Single-Base Resolution". *Science.* 336(6083):934-937. 22539555. DOI:10.1126/science.1220671.

49. Houseman EA, Johnson KC, Christensen BC. (2016) "OxyBS: Estimation of 5-Methylcytosine and 5-Hydroxymethylcytosine from Tandem-Treated Oxidative Bisulfite and Bisulfite DNA". *Bioinformatics (Oxford, England).* 32(16):2505-2507. 27153596. DOI:10.1093/bioinformatics/btw158.

50. Beach TG, Adler CH, Sue LI, Serrano G, Shill HA, Walker DG, Lue L, Roher AE, Dugger BN, Maarouf C, Birdsill AC, Intorcia A, Saxon-Labelle M, Pullen J, Scroggins A, Filon J, Scott S, Hoffman B, Garcia A, Caviness JN, Hentz JG, Driver-Dunckley E, Jacobson SA, Davis KJ, Belden CM, Long KE, Malek-Ahmadi M, Powell JJ, Gale LD, Nicholson LR, Caselli RJ, Woodruff BK, Rapscak SZ, Ahern GL, Shi J, Burke AD, Reiman EM, Sabbagh MN. (2015) "Arizona Study of Aging and Neurodegenerative Disorders and Brain and Body Donation Program ". *Neuropathology.* 35(4):354-389. 25619230. DOI:10.1111/NEUP.12189.

51. Michel Goedert, Maria Grazia Spillantini, Kelly Del Tredici, Heiko Braak. (2012) "100 Years of Lewy Pathology". *Nature Reviews Neurology.* 9(1):13-24. DOI:10.1038/nrneurol.2012.242.

52. "Bioconductor - ENmix".

53. Xu Z, Taylor JA, Leung YK, Ho SM, Niu L. (2016) "oxBS-MLE: An Efficient Method to Estimate 5-Methylcytosine and 5-Hydroxymethylcytosine in Paired Bisulfite and Oxidative Bisulfite Treated DNA". *Bioinformatics (Oxford, England).* 32(23):3667-3669. DOI:10.1093/BIOINFORMATICS/BTW527.

54. Fortin J, Triche TJ, Hansen KD, Hansen KD. (2017) "Preprocessing, Normalization and Integration of the Illumina HumanMethylationEPIC Array with Minfi". *Bioinformatics (Oxford, England).* 33(4):558-560. DOI:10.1093/bioinformatics/btw691.

55. Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, Irizarry RA. (2014) "Minfi: A Flexible and Comprehensive Bioconductor Package for the Analysis of Infinium DNA Methylation Microarrays". *Bioinformatics.* 30(10): 1363-1369.DOI:10.1093/BIOINFORMATICS/BTU049.

56. "Bioconductor - ChAMP".

57. Tian Y, Morris TJ, Webster AP, Yang Z, Beck S, Feber A, Teschendorff AE. (2017) "ChAMP: Updated Methylation Analysis Pipeline for Illumina BeadChips". *Bioinformatics (Oxford, England).* 33(24):3982-3984. DOI:10.1093/bioinformatics/btx513.

58. Jiao C, Zhang C, Dai R, Xia Y, Wang K, Giase G, Chen C, Liu C. (2018) "Positional Effects Revealed in Illumina Methylation Array and the Impact on Analysis". *Epigenomics.* 10(5):643-659. DOI:10.2217/EPI-2017-0105.

59. Guintivano J, Aryee MJ, Kaminsky Za. (2013) "A Cell Epigenotype Specific Model for the Correction of Brain Cellular Heterogeneity Bias and its Application to Age, Brain Region and Major Depression". *Epigenetics : official journal of the DNA Methylation Society.* 8(3):290-302. DOI:10.4161/epi.23924.

60. Zhou W, Hinoue T, Barnes B, Mitchell O, Iqbal W, Lee SM, Foy KK, Lee KH, Moyer EJ, VanderArk A, Koeman JM, Ding W, Kalkat M, Spix NJ, Eagleson B, Pospisilik JA, Szabó PE, Bartolomei MS, Vander Schaaf NA, Kang L, Wiseman AK, Jones PA, Krawczyk CM, Adams M, Porecha R, Chen BH, Shen H, Laird PW. (2022) "DNA Methylation Dynamics and Dysregulation Delineated by High-Throughput Profiling in the Mouse". *Cell Genomics.* 2(7):100144. DOI:10.1016/J.XGEN.2022.100144.

61. Wahl S, Fenske N, Zeilinger S, Suhre K, Gieger C, Waldenberger M, Grallert H, Schmid M. (2014) "On the Potential of Models for Location and Scale for Genome-Wide DNA Methylation Data". *BMC Bioinformatics.* 15(1):232. DOI:10.1186/1471-2105-15-232.

62. Rigby RA, Stasinopoulos DM, Lane PW. (2005) "Generalized Additive Models for Location, Scale and Shape". *Journal of the Royal Statistical Society: Series C (Applied Statistics).* 54(3):507-554. DOI:10.1111/J.1467-9876.2005.00510.X.

63. Breheny P, Stromberg A, Lambert J. (2018) "P-Value Histograms: Inference and Diagnostics". *High-throughput.* 7(3):23. 30200313. DOI:10.3390/ht7030023.

64. Vu H, Ernst J. (2022) "Universal Annotation of the Human Genome through Integration of Over a Thousand Epigenomic Datasets". *Genome Biology.* 23(1):9. 34991667. DOI:10.1186/s13059-021-02572-z.

65. Yao Z, van Velthoven CTJ, Kunst M, Zhang M, McMillen D, Lee C, Jung W, Goldy J, Abdelhak A, Aitken M, Baker K, Baker P, Barkan E, Bertagnolli D, Bhandiwad A, Bielstein C, Bishwakarma P, Campos J, Carey D, Casper T, Chakka AB, Chakrabarty R, Chavan S, Chen M, Clark M, Close J, Crichton K, Daniel S, DiValentin P, Dolbeare T, Ellingwood L, Fiabane E, Fliss T, Gee J, Gerstenberger J, Glandon A, Gloe J, Gould J, Gray J, Guilford N, Guzman J, Hirschstein D, Ho W, Hooper M, Huang M, Hupp M, Jin K, Kroll M, Lathia K, Leon A, Li S, Long B, Madigan Z, Malloy J, Malone J, Maltzer Z, Martin N, McCue R, McGinty R, Mei N, Melchor J, Meyerdierks E, Mollenkopf T, Moonsman S, Nguyen TN, Otto S, Pham T, Rimorin C, Ruiz A, Sanchez R, Sawyer L, Shapovalova N, Shepard N, Slaughterbeck C, Sulc J, Torkelson A, Tung H, Valera Cuevas N, Vance S, Wadhwani K, Ward K, Levi B, Farrell C, Young R, Staats B, Wang MM, Thompson CL, Mufti S, Pagan CM, Kruse L, Dee N, Sunkin SM, Esposito L, Hawrylycz MJ, Waters J, Ng L, Smith K, Tasic B, Zhuang X, Zeng H. (2023) "A High-Resolution Transcriptomic and Spatial Atlas of Cell Types in the Whole Mouse Brain". *Nature (London).* 624(7991):317-332. 38092916. DOI:10.1038/s41586-023-06812-z.

66. Rouillard AD, Gundersen GW, Fernandez NF, Wang Z, Monteiro CD, McDermott MG, Ma'ayan A. (2016) "The Harmonizome: A Collection of Processed Datasets Gathered to Serve and Mine Knowledge about Genes and Proteins". *Database.* 2016:baw100. DOI:10.1093/database/baw100.

67. Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, Stein TI, Nudel R, Lieder I, Mazor Y, Kaplan S, Dahary D, Warshawsky D, Guan-Golan Y, Kohn A, Rappaport N, Safran M, Lancet D. (2016) "The GeneCards Suite: From Gene Data Mining to Disease Genome Sequence Analyses". *Current Protocols in Bioinformatics.* 54:1.30.1-1.30.33. 27322403. DOI:10.1002/cpbi.5.

68. Bindea G, Mlecnik B, Hackl H, Charoentong P, Tosolini M, Kirilovsky A, Fridman W, Pagès F, Trajanoski Z, Galon J. (2009) "ClueGO: A Cytoscape Plug-in to Decipher Functionally Grouped Gene Ontology and Pathway Annotation Networks". *Bioinformatics.* 25(8):1091-1093. DOI:10.1093/bioinformatics/btp101.

69. Smoot ME, Ono K, Ruscheinski J, Wang P, Ideker T. (2011) "Cytoscape 2.8: New Features for Data Integration and Network Visualization.". *Bioinformatics (Oxford, England).* 27(3):431-432. DOI:10.1093/bioinformatics/btq675.

70. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, Gable AL, Fang T, Doncheva NT, Pyysalo S, Bork P, Jensen LJ, von Mering C. (2023) "The STRING Database in 2023: Protein–protein Association Networks and Functional Enrichment Analyses for any Sequenced Genome of Interest ". *Nucleic acids research.* 51(D1):D638-D646. 36370105. DOI:10.1093/nar/gkac1000.

71. Sandor C, Robertson P, Lang C, Heger A, Booth H, Vowles J, Witty L, Bowden R, Hu M, Cowley SA, Wade-Martins R, Webber C. (2017) "Transcriptomic Profiling of Purified Patient-Derived Dopamine Neurons Identifies Convergent Perturbations and Therapeutics for Parkinson's Disease". *Human Molecular Genetics.* 26(3):552-566. 28096185.PMC5409122. DOI:10.1093/hmg/ddw412.

72. Grünblatt E, Mandel S, Jacob-Hirsch J, Zeligson S, Amariglo N, Rechavi G, Li J, Ravid R, Roggendorf W, Riederer P, Youdim MBH. (2004) "Gene Expression Profiling of Parkinsonian Substantia Nigra Pars Compacta; Alterations in Ubiquitin-Proteasome, Heat Shock Protein, Iron and Oxidative Stress Regulated Proteins, Cell Adhesion/Cellular Matrix and Vesicle Trafficking Genes". *Journal of Neural Transmission (Vienna, Austria: 1996).* 111(12):1543-1573. 15455214. DOI:10.1007/s00702-004-0212-1.

73. Chuang Y, Lu AT, Paul KC, Folle AD, Bronstein JM, Bordelon Y, Horvath S, Ritz B. (2019) "Longitudinal Epigenome-Wide Methylation Study of Cognitive Decline and Motor Progression in Parkinson's Disease". *Journal of Parkinson's Disease.* 9(2):389-400. 30958317.PMC6946688. DOI:10.3233/JPD-181549.

74. Kochmanski J, Bernstein AI. Best practices for epigenome-wide DNA modification data collection and analysis . In: Rigor and Reproducibility in Genetics and Genomics. Academic Pressp. 261-284.

75. Blauwendraat C, Nalls MA, Singleton AB. (2020) "The Genetic Architecture of Parkinson's Disease". *Lancet neurology.* 19(2):170-178. 31521533. DOI:10.1016/S1474-4422(19)30287-X.

76. Çiğdem Köroğlu, Leyla Baysal, Murat Cetinkaya, Hatice Karasoy, Aslıhan Tolun. (2013) "DNAJC6 is Responsible for Juvenile Parkinsonism with Phenotypic Variability". *Parkinsonism & Related Disorders.* 19(3):320-324. DOI:10.1016/j.parkreldis.2012.11.006.

77. Edvardson S, Cinnamon Y, Ta-Shma A, Shaag A, Yim Y, Zenvirt S, Jalas C, Lesage S, Brice A, Taraboulos A, Kaestner KH, Greene LE, Elpeleg O. (2012) "A Deleterious Mutation in DNAJC6 Encoding the Neuronal-Specific Clathrin-Uncoating Co-Chaperone Auxilin, is Associated with Juvenile Parkinsonism". *PLOS ONE.* 7(5):e36458. DOI:10.1371/journal.pone.0036458.

78. Liena Elbaghir Omer Elsayed, Valérie Drouet, Tatiana Usenko, Inaam N. Mohammed, Ahlam AbdAlrahman Ahmed Hamed, Maha Abdelmoneim Elseed, Mustafa A.M. Salih, Mahmoud Eltayeb Koko, Ashraf Yahia Osman Mohamed, Rayan Abubaker Siddig, Mustafa Idris Elbashir, Muntaser Eltayeb Ibrahim, Alexandra Durr, Giovanni Stevanin, Suzanne Lesage, Ammar Eltahir Ahmed, Alexis Brice. (2016) "A Novel Nonsense Mutation in DNAJC6Expands the Phenotype of Autosomal-Recessive Juvenile-Onset Parkinson's Disease". *Annals of Neurology.* 79(2):335-337. DOI:10.1002/ana.24591.

79. Simone Olgiati, Marialuisa Quadri, Mingyan Fang, Janneke P.M.A. Rood, Jonas A. Saute, Hsin Fen Chien, Christian G. Bouwkamp, Josja Graafland, Michelle Minneboo, Guido J. Breedveld, Jianguo Zhang, Frans W. Verheijen, Agnita J.W. Boon, Anneke J.A. Kievit, Laura Bannach Jardim, Wim Mandemakers, Egberto Reis Barbosa, Carlos R.M. Rieder, Klaus L. Leenders, Jun Wang, Vincenzo Bonifati. (2016) "DNAJC6Mutations Associated with Early-Onset Parkinson's Disease". *Annals of Neurology.* 79(2):244-256. DOI:10.1002/ana.24553.

80. Roosen DA, Blauwendraat C, Cookson MR, Lewis PA. (2019) "DNAJC Proteins and Pathways to Parkinsonism". *The FEBS journal.* 286(16):3080-3094. 31120186.PMC9048129. DOI:10.1111/febs.14936.

81. Quadri M, Fang M, Picillo M, Olgiati S, Breedveld GJ, Graafland J, Wu B, Xu F, Erro R, Amboni M, Pappatà S, Quarantelli M, Annesi G, Quattrone A, Chien HF, Barbosa ER, Oostra BA, Barone P, Wang J, Bonifati V. (2013) "Mutation in the SYNJ1 Gene Associated with Autosomal Recessive, Early-Onset Parkinsonism". *Human Mutation.* 34(9):1208-1215. 23804577. DOI:10.1002/humu.22373.

82. Krebs CE, Karkheiran S, Powell JC, Cao M, Makarov V, Darvish H, Di Paolo G, Walker RH, Shahidi GA, Buxbaum JD, De Camilli P, Yue Z, Paisán-Ruiz C. (2013) "The Sac1

Domain of SYNJ1 Identified Mutated in a Family with Early-Onset Progressive Parkinsonism with Generalized Seizures". *Human Mutation.* 34(9):1200-1207. 23804563.PMC3790461. DOI:10.1002/humu.22372.

83. Jacquemyn J, Kuenen S, Swerts J, Pavie B, Vijayan V, Kilic A, Chabot D, Wang Y, Schoovaerts N, Corthout N, Verstreken P. (2023) "Parkinsonism Mutations in DNAJC6 Cause Lipid Defects and Neurodegeneration that are Rescued by Synj1". *NPJ Parkinson's disease.* 9(1):19. 36739293.PMC9899244. DOI:10.1038/s41531-023-00459-3.

84. Ng XY, Wu Y, Lin Y, Yaqoob SM, Greene LE, De Camilli P, Cao M. (2023) "Mutations in Parkinsonism-Linked Endocytic Proteins Synaptojanin1 and Auxilin have Synergistic Effects on Dopaminergic Axonal Pathology". *NPJ Parkinson's Disease.* 9(1):26. 36792618. DOI:10.1038/s41531-023-00465-5.

85. Kochmanski J, Virani M, Kuhn NC, Boyd SL, Becker K, Adams M, Bernstein AI. Developmental Origins of Parkinson's Disease Risk: Perinatal Exposure to the Organochlorine Pesticide Dieldrin Leads to Sex-Specific DNA Modifications in Critical Neurodevelopmental Pathways in the Mouse Midbrain. bioRxiv; 2024. DOI:10.1101/2024.04.26.590998.

86. Alter SP, Lenzi GM, Bernstein AI, Miller GW. (2013) "Vesicular Integrity in Parkinson's Disease". *Current Neurology and Neuroscience Reports.* 13(7):362. 23690026.PMC4019229. DOI:10.1007/s11910-013-0362-3.

87. Brooker SM, Naylor GE, Krainc D. (2024) "Cell Biology of Parkinson's Disease: Mechanisms of Synaptic, Lysosomal, and Mitochondrial Dysfunction". *Current Opinion in Neurobiology.* 85:102841. 38306948. DOI:10.1016/j.conb.2024.102841.

88. Soukup S, Vanhauwaert R, Verstreken P. (2018) "Parkinson's Disease: Convergence on Synaptic Homeostasis". *The EMBO journal.* 37(18):e98960. 30065071. DOI:10.15252/embj.201898960.

89. Gcwensa NZ, Russell DL, Cowell RM, Volpicelli-Daley LA. (2021) "Molecular Mechanisms Underlying Synaptic and Axon Degeneration in Parkinson's Disease". *Frontiers in cellular neuroscience.* 15:626128. 33737866. DOI:10.3389/fncel.2021.626128.

90. Anderson FL, Coffey MM, Berwin BL, Havrda MC. (2018) "Inflammasomes: An Emerging Mechanism Translating Environmental Toxicant Exposure into Neuroinflammation in Parkinson's Disease". *Toxicological Sciences: An Official Journal of the Society of Toxicology.* 166(1):3-15. 30203060.PMC6659017. DOI:10.1093/toxsci/kfy219.

91. Hirsch EC, Standaert DG. (2021) "Ten Unsolved Questions about Neuroinflammation in Parkinson's Disease". *Movement Disorders.* 36(1):16-24. 32357266. DOI:10.1002/mds.28075.

92. Tansey MG, Wallings RL, Houser MC, Herrick MK, Keating CE, Joers V. (2022) "Inflammation and Immune Dysfunction in Parkinson Disease". *Nature reviews. Immunology.* 22(11):657-673. 35246670. DOI:10.1038/s41577-022-00684-6.

93. Wang Y, Kuca K, You L, Nepovimova E, Heger Z, Valko M, Adam V, Wu Q, Jomova K. (2024) "The Role of Cellular Senescence in Neurodegenerative Diseases". *Archives of Toxicology.*38744709. DOI:10.1007/s00204-024-03768-5.

94. Holloway K, Neherin K, Dam KU, Zhang H. (2023) "Cellular Senescence and Neurodegeneration". *Human genetics.* 142(8):1247-1262. 37115318. DOI:10.1007/s00439-023-02565-x.

95. Wu D, Sun JK, Chow KH. (2024) "Neuronal Cell Cycle Reentry Events in the Aging Brain are More Prevalent in Neurodegeneration and Lead to Cellular Senescence". *PLOS Biology.* 22(4):e3002559. DOI:10.1371/journal.pbio.3002559.

Supplementary Figures

Supplementary Figure 1: P-value histograms for hypothesis testing in interaction and separate analyses. P-value histograms can be used as a quality control measure in association studies with a large number of hypothesis tests. Using a normal distribution model, the interaction hypothesis test histogram (A) is appropriate, with a peak of low p-values as would be expected from a well-calibrated hypothesis test. Histograms from separate modeling for 5mC (B) and 5hmC (C) alone show an overabundance of high p-values. Beta regression modeling did not perform well in interaction (D) or separate 5mC (E) or 5hmC (F) modeling.

Supplementary Figure 2: String protein interaction network for each GO Term Group. Protein interaction networks of genes within each GO Term Group (confidence < 0.7; >2 genes per network). The following groups that correspond to GO terms in Table 2 are displayed: (A) group 1, (B) group 2, (C) group 3, (D) group 5, (E) group 6. Groups 0 and 4 are not shown because they had no connected networks with >2 genes.