1 **Comparative and population genomics analyses of eared pheasants**
2 **inhabiting highly varying altitudes**
3

Siwen Wu[1,3†], Kun Wang[2,4†], Xuehai Ge[2†], Sisi Yuan[3], Dong-Dong Wu[4], Changrong Ge[2*], Junjing

Jia[2*], Zhengchang Su[3*], Tengfei Dou[2*]

1. Institute of Medical Biology, Chinese Academy of Medical Sciences, Peking Union Medical

College, Kunming, Yunnan, China.

2. Faculty of Animal Science and Technology, Yunnan Agricultural University, Kunming 650201,

Yunnan, China.

3. Department of Bioinformatics and Genomics, College of Computing and Informatics, the

University of North Carolina at Charlotte, Charlotte, NC 28223, USA.

4. State Key Laboratory of Genetic Resources and Evolution/Key Laboratory of Healthy Aging

Research of Yunnan Province, Kunming Institute of Zoology, Chinese Academy of Sciences,

Kunming, Yunnan, China.

†These authors contributed equally to this work.

[*]Corresponding author(s): Changrong Ge (gcrzal@126.com), Junjing Jia

(junjingli2009@hotmail.com), Zhengchang Su (zcsu@uncc.edu), Tengfei Dou

(tengfeidou@sina.com)

Abstract

Oxygen pressure varies dramatically with altitudes on Earth; however, humans and animals

thrive at almost all altitudes. To better understand genetic basis underlying adaptation of

closely related species to varying altitudes, we annotated and compared the genome of a white

eared pheasant (WT) (*Crossoptilon crossoptilon*) inhabiting high altitudes and the genome of a

brown eared pheasant (BR) (*C. mantchuricum*) inhabiting low altitudes. Moreover, we

compared genetic variations in populations of WT and BR as well as of blue eared pheasants (BL)

(*C. auritum*) inhabiting intermediate altitudes, and identified thousands of selective sweeps in

each species. Intriguingly, the unique genes and pseudogenes in the two genomes converge on

the same set of altitude adaptation-related pathways of four functional categories as genes in

selective sweeps in each species. Thus, these species appear to adapt to highly varying altitudes

by diverging selection on the same traits via loss-of-function mutations and fine-tuning genes in

common pathways.

Introduction

Faunas have gone through an evolutionary process to gain physiological feature for the adaptation of highly varying oxygen abundance. Populations of humans[1-3] and many animal species that are naturally adapted to lowlands thrive in highland environments, such as Tibetan chickens[4-6], Tibetan pigs[7,8], Tibetan goats[9], Tibetan sheep[10,11], Tibetan mastiffs[12], Tibet cattle[13,14], ground tits[15], and yaks[16], to name a few. The major challenge faced by these populations of humans and animals in highland environments is the low oxygen pressure-induced stress on their physiological systems[17-19]. The long-term adaptation to low oxygen pressure environments has resulted in genetic changes in these human[1-3,14,17,20-27] and animal[4-7,10-16,19,28-31] populations. Adaptive mutations of many genes of humans[2,14,17,18,26,32,33] and animals[16,19,30,31,33,34] have been linked to highland environments. Moreover, introgression has been shown as a major mechanism for human[1,3,27] and domesticated animals[9,10,13] to adapt to highland environments. Despite these great progresses, the understanding of human and animal adaptation to highly varying altitudes is still limited[33]. Particularly, some closely related animal species can rapidly adapt to highly varying altitudes, thus, it is interesting to understand the genetic basis of their adaptation from low- to intermediate- and high-altitude environments.

Crossoptilon, belonging to the Phasianidae family in the Galliformes order, is a rare but important genus endemic in China[35]. There are four species in the Crossoptilon genus, including Tibetan eared pheasant (TB) (*C. harmani*), white eared pheasant (WT) (*C. crossoptilon*), blue eared pheasant (BL) (*C. auritum*) and brown eared pheasant (BR) (*C. mantchuricum*)[35,36]. These species are found in coniferous forests, mixed broadleaf-conifer forests and alpine scrubs in various parts of China with very different altitudes ranging from 800 m to 5,000 m[35,37-42]. TB and

WT are believed to be conspecifics, diverging about 0.5 million years ago[43]. They sympatrically

inhabit montane forest at a high altitude of 3,000-5,000 m[35,36,38,40-42]. TB is distributed in

southeastern Tibet and adjacent northern India (3,000-5,000 m)[35,38,41], while WT is found in

Qinghai, Sichuan, Yunnan and Tibet (3,000-4,300 m)[35,40]. BL and BR are closely related,

diverging about 0.3 million years ago[43], but allopatrically inhabit different areas[35,36]. BL is only

encountered in the mountains of Qinghai, Gansu, Sichuan and Ningxia, at an intermediate

altitude of 2,000-3,000 m[38,44], while BR is mainly distributed in mountains in Shanxi and Hebei

and near Beijing, at an altitude of 800-2,600 m[37,39,42]. Thus, WT, BL and BR are excellent models

to study the genetic basis for closely related species to adapt to highly varying altitudes.

Recently, Wang et al.[45] assembled a BR genome and re-sequenced three BR

subpopulations and a BL population. Although they provide new insights into genomic changes

during the course of BR's long-term population decline[45], genetic basis underlying the

adaptation of the BR and BL to low (800-2,600 m) and intermediated altitudes (1500-3000 m),

respectively, were not investigated. Moreover, with a small contig N50 (0.11 Mb), this BR

genome assembly has limitations as a reference genome to study the *Crossoptilon* species. To

fill the gap, we recently sequenced and assembled the genome of a WT individual at

chromosome-level with a contig N50 of 19.6 Mb, and re-sequenced 10 WT individuals and a BL

individual[46]. Combining our data in WT and BL[46] with those of Wang et al. in BL and BR[45], we

were well positioned to investigate the genetic bases of these closely related avian species to

adapt to highly varying altitudes using comparative and population genomics approaches. This

study is based in part upon SW's dissertation[47].

Results

Annotation of protein-coding genes in the WT and BR genomes

To see whether gene contents are related to adaption of closely related avian species to different altitudes, we first annotated the assembled WT[46] and BR[45] genomes using a combination of reference- and RNA-based methods (Materials and Methods). Using the coding DNA sequences (CDSs) of 53 well-represented Aves (Table S1) as the templates and a total of 23 RNA-seq datasets from 20 tissues of WT individuals and three tissues of BR individuals (Materials and Methods), we annotated 16,377 and 15,410 protein-coding genes as well as 1,519 and 1,976 pseudogenes in the WT and BR genomes, respectively (Table 1). The vast majority (15,565 and 14,727) of these genes and all the pseudogenes (1,519 and 1,976) in the WT and BR genomes were predicted based on the CDSs in the reference genomes (reference-based), while a small portion (812 and 683) of the genes were predicted based on the RNA-seq datasets in the two species (RNA-based) (Table 1). To evaluate the effects of the unbalanced RNA-seq datasets on the numbers of annotated genes in the two genomes, we repeated the annotation process by 100 times with each time using the CDSs in the 53 Aves genomes for reference-based annotations and a total of six RNA-seq datasets for RNA-based annotation. Of the six RNA-seq datasets, three were those from the BR tissues and the other three were randomly selected in each annotation from the 20 datasets of the WT tissues. As shown in Table S2, in each of the 100 times of annotations, the total number of annotated genes in WT is larger than that in BR, which is mainly due to the larger number of genes annotated by the reference-based approach in WT (15,565) than in BR (14,727). On the other hand, the two genomes have a similar average number (350 vs 355) of genes annotated by the RNA-based approach (p-value = 0.33, two-tailed t-test). Interestingly, in 73 of the 100 times of annotations,

the BR genome has even more RNA-annotated genes than does the WT genome, suggesting

that the assembly quality of the CDSs in BR is good enough for finding genes. However, when

the union of the 100 annotations is taken, WT still has more RNA-based annotated genes (812

vs 683). Taken together, these results suggest that the difference in the numbers of genes in

the two species is not mainly due to the difference in the quality of genome assemblies and/or

the bias of RNA-seq datasets.

Most of the reference-based genes (14,815 and 14,518) in both species have an intact

open reading frame (ORF) (intact genes), while the remaining small numbers (750 and 209)

contain either premature stop codon or ORF-shift mutations, which are not supported by the

corresponding short DNA reads, we thus called them partially supported genes, as the

mutations might be caused by sequencing errors, particularly, in long reads. Vast majority of

the intact genes (99.01%), partially supported genes (97.73%), and pseudogenes (95.13%) in

WT were transcribed in at least one of the 20 tissues examined (Table S3). These percentages

(96.54%, 93.78% and 90.84%) are smaller in BR (Table S4) as RNA-seq data from only three

tissues are available[45]. Most (751 and 631) of the RNA-based genes in both species can be

mapped to the NT database[48] (NT-supported new genes), suggesting that they are likely true

genes. The remaining small numbers (61 and 52) that cannot be mapped to the NT database

are likely novel genes (Table 1).

We compared the RNA-based genes annotated in the WT and BR assemblies, and found

450 pairs with an identity > 96.8%, comprising 55.4% and 65.9% of their RNA-based genes,

respectively. They are likely true orthologous genes in both species, as the same patterns of

transcriptional noise are unlikely to occur in two different species. Moreover, most of the RNA-

based genes in both species (Tables S3 and S4) were varyingly expressed in different tissues

with RNA-seq data available (Figures 1a-1d), indicating that they might be authentic and

functional. In addition, we randomly selected 17 of the 812 RNA-based genes in WT and

measured their expression levels in 16 tissues using RT-qPCR. We found that 15 (88.24%) of

them were transcribed in at least one of the tissues and five of which were putative novel

genes (Table S5, Figure 1e), further suggesting that most of the RNA-based genes at least in WT

are likely authentic, although the expression patterns of some genes are different from those

seen in the RNA-seq data due probably to the different sensitivity of RT-qPCR and RNA-seq

methods (Figure 1f). Moreover, we annotated 12 and eight in WT and BR, respectively, of the

274 presumed missing genes in avian species[49] (Table S6).

Most pseudogenes found in the WT and BR genomes are unprocessed and unitary and thus

might lose gene functions

Vast majority (for WT, 1,310 or 86.2%; for BR, 1,701 or 86.1%) of pseudogenes in both WT and

BR genomes are unprocessed, i.e., they arose due to direct mutations in the parent genes. We

failed to find any functional copy of these unprocessed pseudogenes; thus, they are also

unitary[50]. To see whether the pseudogenization alleles in the 1,519 and 1,976 pseudogenes in

the WT and BR genomes, respectively, are under selection, we mapped the short reads from 10

WT individuals and 41 BR individuals to the corresponding genomes (Materials and Methods).

We found that most of the first pseudogenization alleles along the parental genes were fixed in

the population of the two species (Figures 2a and 2b), suggesting that these pseudogenization

events might be under positive selection and thus contribute to the phenotypic traits of the two

species. Interestingly, the synonymous mutations in true genes in the two species, which are

generally assumed to be selectively neutral, are largely uniformly distributed along the CDSs,

except at the two ends, where the numbers of synonymous mutations decrease (Figures 2c and

2d), consistent with an earlier report in chickens[51]. The reduced synonymous mutations at the

two ends of CDSs suggest that they might harbor functional elements not related to their

coding functions, such as transcriptional and post-transcriptional regulatory elements[52]. In

contrast, pseudogenization mutations are strongly biased to the 5'-ends and 3'-ends in both

species (Figures 2c and 2d). The same phenomenon was also found in the other species such as

human[51,53] and chickens[51]. Therefore, the strongly biased pseudogenization mutations to the 5'-

end and 3'-end in the two species also suggest that they are under positive selection.

Finally, of the 1,519 pseudogenes in the WT genome, 727 (47.9%) have alternative

transcripts, but only 10 (1.4%) have functional alternative transcripts (Table S7). Of the 1,976

pseudogenes in BR, 1,712 have alternative transcripts, but only 17 (1.0%) have functional

alternative transcripts (Table S8). These results suggest that most pseudogenes in both species

do not have functional alternative transcripts, further suggesting that most of them have lost

the functions of their parent genes.

Unique genes and pseudogenes in WT and BR might be related to their adaptation to distinct

ecological niches

We found that the WT and BR genomes shared 14,507 of their genes (88.6% and 94.1%) and

1,040 of their pseudogenes (68.5% and 52.6%) (Table 1). Meanwhile, the WT genome contains

1,870 unique genes and 479 unique pseudogenes, while the BR genome harbors 903 unique

genes and 936 unique pseudogenes (Tables S9-S12, Figure 3a). Of the 1,870 unique genes in the

WT genome, 658 (35.2%) are unique pseudogenes and 1,212 (64.8%) are missing in the BR

genome. Of the 936 unique pseudogenes in BR, 658 (70.3%) are unique genes and 278 (29.7%)

are missing in the WT genome (Figure 3a). Moreover, of the 903 unique genes in the BR

genome, 173 (19.2%) are unique pseudogenes and 730 (80.8%) are missing in the WT genome.

Of the 479 unique pseudogenes in the WT genome, 173 (36.1%) are unique genes and 306

(63.9%) are missing in the BR genome (Figure 3a). There are a total of 3,357 genes that either

are unique genes or become unique pseudogenes in the WT or BR genomes. To see whether

the unique genes and pseudogenes in each species are related to their unique traits and reflect

the evolutionary pressure received from their niches, we assigned GO terms to the unique

genes and pseudogenes in each species based on their homologs in chickens or humans. The

unique genes and pseudogenes in WT are involved in 69 and 25 biological pathways (Table S13),

respectively, while those in BR are involved in 27 and 59 biological pathways, respectively

(Table S13). Interestingly, the unique genes and pseudogenes with GO term assignments are

mainly involved in and affect, respectively, pathways related to four major physiological

functions: cardiovascular functions, energy metabolism, neuronal functions and immunity

(Table S13, Figure 3b). It has been shown that alterations in these functions are critical for

animals to adapt to different altitudes[33]. Thus, the unique genes and loss of functions of the

missing genes and pseudogenes in BR and WT might play a role in the adaption of the two

species to their strikingly different altitudes. A few examples of such genes and pseudogenes

are given below.

1) Cardiovascular functions: As shown in Figure 3b, unique gene *PIK3C2G* in BR, which is

pseudogenized in WT, is involved in the HIF (hypoxia-inducible factor) activation pathway[54-56].

Unique genes *SH2D2A*, *PRKCD*, *PRKD3* and *NOS3* in WT, which are all missing in BR, are involved

in the VEGF (vascular endothelial growth factor) signaling pathway that plays critical roles in

proliferation, survival and migration of blood endothelial cells required for the angiogenesis

pathway[57-59]. Unique genes *SH2D2A*, *MAP3K1*, *PRKCD*, *WNT1*, *PRKD3*, *NOS3*, *EPHA3*, *NOTCH1*

and *GRB14* in WT, which are missing (*SH2D2A*, *PRKCD*, *PRKD3*, *NOS3* and *EPHA3*) or

pseudogenized (*MAP3K1*, *WNT1*, *NOTCH1* and *GRB14*) in BR, are involved in the angiogenesis

pathway. Unique genes *DOK1* and *PIK3C2G* in BR, which are pseudogenized in WT, are involved

in the angiogenesis pathway. Unique genes *PRKCD*, *NOS3* and *NPR2* in WT are missing (*PRKCD*

and *NOS3*) or pseudogenized (*NPR2*) in BR, are involved in the endothelin signaling pathway.

Unique gene *ARRB1* in BR, which is missing in WT, is involved in the angiotensin II-stimulated

signaling through G proteins and beta-arrestin pathway. Unique gene *VWF* in WT, which is

pseudogenized in BR, is involved in blood coagulation pathway.

2) Energy metabolism: As shown in Figure 3b, unique gene *PFKM* in BR, which is

pseudogenized in WT, is involved in glycolysis. Unique genes *PRKCD*, *CACNB1* and *CACNB3* in

WT, which are missing (*PRKCD* and *CACNB3*) or pseudogenized (*CACNB1*) in BR, are involved in

the thyrotropin-releasing hormone receptor signaling pathway. Unique genes *IRS2* and *TSC2* in

WT, which are missing (*TSC2*) or pseudogenized (*IRS2*) in BR, are involved in the insulin/IGF

pathway-protein kinase B signaling cascade pathway. Unique gene *TSC2* in WT, which is missing

in BR, is involved in the p53 pathway by glucose deprivation.

3) Neuronal functions: As shown in Figure 3b, unique genes *CACNA1D*, *CACNB1* and

CACNB3 in WT, which are missing (*CACNA1D* and *CACNB3*) or pseudogenized (*CACNB1*) in BR,

are members of the beta1/2 adrenergic receptor signaling pathway. Unique gene *ADRB1* in BR,

which is pseudogenized in WT, is involved in the beta1/2 adrenergic receptor signaling pathway.

Unique genes *GRIK1* and *GRIN2C* in WT, which are pseudogenized in BR, are involved in the

metabotropic glutamate receptor group I pathway. Unique gene *CACNB1* (metabotropic

glutamate receptor group III pathway) in WT is pseudogenized in BR. Unique gene *GABBR1*

(GABA-B receptor II signaling pathway) in WT is missing in BR. Unique genes *CAMK2G*, *GRIK1*,

SHANK2, *GRIN2C* and *CAMK2B* in WT, which are missing (*SHANK2* and *CAMK2B*) or

pseudogenized (*CAMK2G*, *GRIK1* and *GRIN2C*) in BR, are involved in the ionotropic glutamate

receptor pathway. Unique genes *CACNA1D*, *MYO15A* and *CACNB1* in WT, which are missing

(*CACNA1D*) or pseudogenized (*MYO15A*, *CACNB1*) in BR, are involved in the nicotinic

acetylcholine receptor signaling pathway.

4) Immunity: As shown in Figure 3b, unique gene *MAP3K1* in WT, which is

pseudogenized in BR, is involved in the T cell activation pathway. Unique gene *PIK3C2G* in BR,

which is pseudogenized in WT, is involved in the T cell activation pathway. Unique genes *CD79A*

and *PRKCD* in WT, which are missing in BR, are involved in the B cell activation pathway. Unique

genes *PREX1*, *CAMK2G*, *ITGB7*, *VWF*, *TYK2*, *IL6* and *CAMK2B* in WT, which are missing (*ITGB7*,

VWF, *TYK2* and *CAMK2B*) or pseudogenized (*PREX1*, *CAMK2G*, *VWF* and *IL6*) in BR, are members

of the inflammation mediated by chemokine and cytokine signaling pathway. Unique gene

ARRB1 in BR, which is missing in WT is involved in inflammation mediated by chemokine and

cytokine signaling pathway.

In addition, some unique genes or unique pseudogenes in the two species are involved

in pathways that might be related to the adaptation of the two species to other aspects of their

distinct ecological niches. For example, unique genes *CACNA1D*, *PRKCD*, *CACNB1* and *CACNB3* in

WT are missing (*CACNA1D*, *PRKCD* and *CACNB3*) or pseudogenized (*CACNB1*) in BR (Table S13).

These genes are involved in oxytocin receptor mediated signaling pathway that has been shown

to play critical roles in social behaviors of animals[60]. Moreover, unique genes *IRS2*, *SMAD4*,

FOSB, *CACNA1D*, *MAP3K1*, *PRKCD*, *MAP2K7*, *EP300*, *NR5A1*, *NPR2*, *MAP3K5* and *CAMK2B* in WT

are missing (*SMAD4*, *FOSB*, *CACNA1D*, *PRKCD*, *MAP2K7*, *NR5A* and *CAMK2B*) or pseudogenized

(*IRS2*, *MAP3K1*, *EP300*, *NPR2* and *MAP3K5*) in BR (Table S13), these genes are in involved in

gonadotropin releasing hormone receptor pathway that regulate reproduction.

The populations of the three species are strongly structured

To further understand the genetic basis of altitude adaptation, we next compared the SNPs of

10 WT individuals, 41 BR individuals and 12 BL individuals (Table 2). Figure 4a shows the

geographical locations where the populations were sampled. Specifically, the three BR

subpopulations were from Shaanxi province, Shanxi province, and Hebei province & Beijing

metropolitan, respectively[45]; the BL population were from Gansu province[45]; and the WT

population were from Yunnan province. Using our assembled WT genome[61] as the reference,

we identified SNPs in the BL, BR and WT populations (Table 2). Based on the bi-allelic SNPs (16

million) in the populations of the three species (Table 2), we performed principal component

analysis (PCA), and found that individuals of the same species form distinct clusters (Figure 4b).

Notably, each of the three BR subpopulations forms a distinct compact subcluster, which is

consistent with the previous result[45]. We also run the admixture algorithm[62] on the SNPs to

infer their ancestral relationships. For K=2, individuals of BL and BR formed a uniformly colored

cluster, while WT individuals formed the other cluster sharing small portions of ancestries with

the former two species (Figure 4c), suggesting that BL and BR were derived from the same

ancestor, while WT branched out earlier. This result is consistent with previous phylogenetic

analysis that BL and BR are more closely related to each other[43]. For K=3, individuals of each

species formed an almost uniformly colored cluster, indicating their distinct recent ancestries.

255   To see possible gene introgression among the three species, we calculated the D-

statistic[63] and f3-statistic[64]. We obtained a D value of 0.57 (z-score = 36.18, and p-value= 2.3e-

16) using a Daweishan chicken population as the outgroup (O) on the rooted tree

(((BR,BL),WT),O), suggesting introgression between BL and WT[63]. The f3 values and associated

z-scores for the three choices of (target; source 1, source 2) are shown in Table 3. The

significant (z-score = -26.45) negative f3 value (-0.0586) for (target=BL; source 1 = WT, source 2

= BR) suggests introgression in BL from both WT and BR[64]. However, the positive f3 values for

the other two scenarios (Table 3) could not indicate there is gene introgression in BR from WT

and BL, and in WT from BR and BL[64]. Thus, although gene introgression might have occurred in

BL from both BR and WT, the data does not support gene introgression in BR from WT and BL,

and in WT from BR and BL. Thus, gene introgression might play a role in the evolution of BL, but

this might not be the case for BR and WT.

Genes in selective sweeps of each species might be related to their distinct ecological niches

To identify genomic regions and genes that might be related to the adaptation of the three

species to their distinct ecological niches, particularly, to different altitudes, we identified

selective sweeps in each species by comparing its SNPs with those of the other two species

using both the genetic differentiation (F_{ST}) and difference in nucleotide diversity ($\Delta\pi$)

parameters[65-68]. To rigorously evaluate the statistical significance of the F_{ST} and $\Delta\pi$ values, we

normalized them (ZF_{ST} and $Z\Delta\pi$) using a permutation test[69] (Materials and Methods). Figures

5a-5c show the distribution of ZF_{ST} and $Z\Delta\pi$ as well as their values for each 40 kb genome

windows with a 20 kb-step size for the three pairwise comparisons. We consider a window with

$ZF_{ST} > 2.33$ (P=0.01) and $Z\Delta\pi > 1.64$ (P=0.05) or $Z\Delta\pi < -1.64$ (P=0.05) as a selective sweep

(Figures 5a-5c). Figures 6a-6f show the Manhattan plots of ZF_{ST} and $Z|\Delta\pi|$ scores for the three

pairwise comparisons. Since adjacent selective sweeps can overlap with one another, we

merged the overlapping ones as a discrete selection sweep (DSS) in each species. For the BL vs

BR comparison, we identified 208 and 638 selective sweeps, and 128 and 463 DSSs containing

143 and 474 genes in BL and BR, respectively (Tables 4 and S14). For the BL vs WT comparison,

we identified 92 and 274 selective sweeps, and 72 and 181 DSSs containing 67 and 167 genes in

BL and WT, respectively (Tables 4 and S15). For the BR vs WT comparison, we identified 306

and 359 selective sweeps, and 184 and 188 DSSs containing 196 and 167 genes in BR and WT,

respectively (Tables 4 and S16).

To further confirm our identified selective sweeps, we computed Tajima's D values for

each 40 kb window. As shown in Figures 6g-6i and Table S17, BR has the largest number (38,488)

of windows with positive Tajima's D values among the three species, suggesting that BR has

experienced a large-scale population contraction, consistent with the previous report[45]. For BL

and WT, most of their selective sweeps identified above in each pair-wise comparison (78.8%,

53.3%, 90% and 87.2% for BL vs BR, BL vs WT, BL vs WT and BR vs WT, respectively) have a

negative Tajima's D value (Table S17), suggesting negative or positive selections and confirming

the selective sweeps identified above.

To find the duration and intensity of the selective sweep in each species, we calculated

the run of homozygosity (ROH) in each selective sweep for each pair of comparison. As shown

in Figures 6j-6l and Table S17, BR has the largest number of selective sweeps (66.7%-70.5%)

with a high ratio of individuals that have a long ROH (>100kb), suggesting that BR has the most

recent inbreeding, their selection sweeps arose recently, and the selection intensity of BR is the

largest compared to the other two species. By contrast, both BL and WT have only small

portions (12%-16.8%) of selective sweeps with a long ROH (Table S17). These results suggest

that BR has undergone the most intensive selection recently compared to BL and WT, which is

consistent with the previous report[45].

To see whether genes in the selective sweeps are under positive or negative selections,

we calculated the ratio of the number of nonsynonymous mutations over the number of

synonymous mutations (dN/dS) for each gene in the DSSs for each pair of comparison. As

shown in Table S17, of the genes in the identified DSSs in each pair of comparison, 76.1-90.4%

are under purifying selection (dN/dS<1), while 6.6-19.4% are under positive selection (dN/dS>1).

Thus, most (90.5-97.6%) of the genes in the identified DSSs are under selection.

Genes in the DSSs of each comparison might be related to adaptation to their different

ecological niches. About 55% of genes in these selective sweeps identified in each comparison

do not harbor fixed nonsynonymous mutations, thus, natural selection might act upon their

regulatory sequences, thereby changing their expression levels. The remaining about 45% of

the genes contain fixed nonsynonymous mutations that might alter the functions of encoded

proteins. For example, gene *MYO3B* is located in a selective sweep in the BR population based

on the BL vs BR comparison (Figure 5a). *MYO3B* in BR contains a fixed G-to-A nonsynonymous

mutation in exon 7, leading to an Arg-to-Lys substitution. Since the AGA codon for Arg at this

position in *MYO3B* is shared by WT, BL, quail (*Coturnix japonica*) and human (*Homo sapiens*)

(Figure 5d), the G-to-A mutation is BR lineage specific. Thus, the Arg-to-Lys substitution in

MYO3B in BR might be related to their unique traits. It is reported that gene *MYO3B* is required

for normal cochlear hair bundle development and hearing[70]. As BR has been hunted by human

for hundreds of years[48], it is highly likely that the mutation might increase their auditory

sensation, thereby helping them to escape earlier when perceiving hunters approaching. It is

interesting to investigate the effects of the Arg-to-Lys mutation on the auditory functions of BR

and its implication for natural selection and adaptation. Moreover, gene *PTPRC* is located in a

selective sweep in the WT populations based on both the BL vs WT and the BR vs WT

comparisons. *PTPRC* in WT contains a fixed A-to-G nonsynonymous mutation in exon 7,

resulting in a Lys-to-Arg substitution. As the AAA codon for Lys at the position in *PTPRC* is

shared by BL, BR, quail and human (Figure 5d), the mutation is WT lineage specific. Interestingly,

we found that the same mutation existed in the *PTPRC* genes of Lance-tailed manakin

(*Chiroxiphia lanceolata*) and White-ruffed manakin (*Corapipo altera*) (Figure 5d), both are

altitudinal migrants living in subtropical Andes highlands[71], due probably to convergent

evolution. It has been reported that *PTPRC* is related to the T cell activation[72]. Therefore, it is

highly likely that the nonsynonymous mutation in the *PTPRC* gene in WT and manakins might

help them adapt to high altitudes where there are fewer pathogens, and thus, the pressure on

the immune system in these species might be relaxed.

Genes under selection tend to be involved in the same altitude adaptation-related pathways

in the three species

There are a total of 1,106 genes that are in a selective sweep predicted based on at least one of

the three comparisons. They share only 105 genes with the 3,357 unique genes and

pseudogenes in the WT and BR genomes (Figure 7a). Intriguingly, the genes with GO term

assignments in the selective sweeps in each species tend to be involved in the same pathways

in the four functional categories (Tables S14-S16) as are the unique genes and unique

pseudogenes with GO term assignments in the WT and BR genomes (Table S13), although the

two sets only share four genes (Figure 7b). For example, selection on cardiovascular function-

related genes *MAPK1* and *PRKAR2B* in BL might be related to its adaptation to intermediate

altitudes, while selection on neuronal function related gene *PRKAR2B* might be beneficial for BL

to survive in its ecological niches (Figure 7b). Moreover, selection on cardiovascular (*PLA2G4A*),

immunity (*ITPR2*), energy metabolism (*PRKCD*) and neuronal function (*SLC17A6*) related genes

might be beneficial for BR to adapt to low altitude (Figure 7b). Furthermore, selection on the

hypoxia response (*PPARG*) related genes might help WT to adapt to the high-altitude niches,

while selection on immunity (*PPP3CB*, *MAPK8* and *PTPRC*) and neuronal function (*GRIN3A*)

related genes might be beneficial for WT to adapt to their unique ecological niches (Figure 7b).

More examples of genes in the selective sweeps, which are involved in altitude-adaptation

related pathways, are given as follows.

For BL, genes in selective sweeps identified in the BL vs BR and BL vs WT comparisons

are involved in the same number of 22 GO pathways (Table 4). Compared to BR, six genes in the

selection sweeps in BL have GO pathway assignments (Table S14). Among these genes, *MAPK1*

is involved in cardiovascular functions such as angiogenesis pathway, VEGF signaling pathway[57-

59], endothelin signaling pathway[73-75], and angiotensin II-stimulated signaling through G proteins

and beta-arrestin pathway (Figure 7b). Compared to WT, 76 genes in the selection sweeps in BL

have GO pathway assignments (Table S15). Among these genes, *PRKAR2B* is involved in

neuronal functions such as beta1/2 adrenergic receptor signaling pathway, metabotropic

glutamate receptor group III pathway, GABA-B receptor II signaling pathway and endothelin

signaling (Figure 7b).

For BR, genes in the selective sweeps identified in the BL vs. BR and BR vs WT

comparisons are involved in 27 and 32 GO pathways, respectively (Table 4). Compared to BL,

175 genes in the selection sweeps in BR have GO pathway assignments (Table S14). Among

these genes, some are involved in cardiovascular functions, such as *PLA2G4A* (VEGF pathway

and angiogenesis pathway), *PLA2G4A* and *ITPR2* (endothelin signaling pathway), *ITPR2*

(angiotensin II-stimulated signaling through G proteins & beta-arrestin pathway), *PLA2G4A*

(oxidative stress response pathway); some are involved in immunity such as *PLA2G4A* and

ITPR2 (B cell activation pathway); some are involved in neuronal functions, such as *RYR2*

(beta1/2 adrenergic receptor signaling pathway), *SLC17A6* (ionotropic glutamate receptor

pathway) (Figure 7b). Compared to WT, 78 genes in the selection sweeps in BR have GO

pathway assignment (Table S16). Some of these genes are involved in cardiovascular functions,

such as *PRKCD* (VEGF pathway, angiogenesis pathway, endothelin signaling pathway) that is

missing in the selection sweep in BR. *PRKCD* is also involved in energy metabolism (thyrotropin-

releasing hormone receptor signaling pathway), immunity (B-cell activation pathway) and

neuronal functions (alpha adrenergic receptor signaling pathway) (Figure 7b). Moreover,

PRKAR2B is also under selection in BR, which is involved in neuronal functions (Figure 7b).

For WT, genes in the selective sweeps identified in the BL vs WT and BR vs WT

comparisons are involved in 62 and 48 GO pathways, respectively (Table 4). Compared to BL,

341 genes in the selection sweeps in WT have GO pathway assignments (Table S15). Of these

genes, some are involved in cardiovascular functions such as *ARNT2* and *PPARG* (hypoxia

induced factor pathway[54-56,76]), *FZD1* and *MAPK8* (angiogenesis pathway) and *MAPK8* (oxidative

stress response pathway); some are involved in energy metabolism pathway, such as *PPP2CB*

(p53 pathway by glucose deprivation); some are involved in neuronal functions, such as *GRIN3A*

(metabotropic glutamate receptor group I pathway, metabotropic glutamate receptor group III

pathway and ionotropic glutamate receptor pathway) and *GABBR2* (GABA-B receptor II

signaling pathway); and some are involved in immunity, such as *PPP3CB*, *MAPK8* and *PTPRC*

(T/B cell activation pathway), *MAPK8* and *UBE2V1* (toll receptor signaling pathway and

interferon-gamma signaling pathway) (Figure 7b).

Moreover, compared to BR, 254 genes in the selection sweeps in WT have GO pathway

assignments (Table S16). Of the these genes, some are involved in cardiovascular functions,

such as *MAPK8* (oxidative stress response pathway[77-81]), *TEK*, *FZD1* and *MAPK8* (angiogenesis

pathway), and *VWF* (blood coagulation pathway) that is missing in BR (Figure 3b); Some are

involved in immunity, such as *MAPK8* (interferon-gamma signaling pathway and toll receptor

signaling pathway), *PPP3CB*, *MAPK8* and *PTPRC* (T/B cell activation pathway and toll receptor

signaling pathway and interferon-gamma signaling pathway[82-84]), *CAMK2G* (pseudogenized in

BR) and *VWF* (inflammation mediated by chemokine and cytokine signaling pathway[85,86]); some

are involved in neuronal functions, such as *GRIN3A* (metabotropic glutamate receptor group I

pathway, metabotropic glutamate receptor group III pathway), *GABBR2* (GABA-

B_receptor_II_signaling pathway), *CAMK2G* and *GRIN3A* (ionotropic glutamate receptor

pathway) (Figure 7b).

Discussion

To better understand the genetic bases of adaptation of closely related eared pheasant species

to highly varying altitudes, we combined our own published data in WT and BL[46] with those in

BR and BL published by Wang et al.[45]. We first annotated the assembled WT[46] and BR[45]

genomes, and compared their gene compositions. We identified similar numbers of protein-

coding genes in the WT (16,447) and BR (15,410) genomes to those annotated in other avian

species[87], but surprisingly large numbers of pseudogenes in them (1,519 in WT and 1,976 in BR),

not previously reported in other avian genomes, to our best knowledge. We provided evidence

that most pseudogenization mutations are under positive selection and the resulting

pseudogenes might have lost gene functions.

We found that many unique genes in one species (WT or BR) are often either missing or

pseudogenized in the other species (BR or WT). Interestingly, these unique genes and genes

with loss-of-function mutations (pseudogenization or missing) are mainly involved in and affect,

respectively, pathways related to four functional categories: cardiovascular, neuronal, energy

metabolic, and immune functions. This finding is quite interesting but not very surprising, since

cardiovascular, neuronal, and metabolic functions are highly sensitive to blood oxygen levels

which is directly affected by oxygen pressures at varying altitudes. Moreover, varying

pathogens and ultraviolet radiations at varying altitudes might require different immune

responses to infections and DNA damages. Physiological systems often contain both positive

and negative regulatory pathways to achieve a homeostasis of the functions. Thus, both the

presence of a gene involved in the positive pathway and the loss of a gene involved in the

negative pathway can enhance the relevant function. Conversely, both the presence of a gene

involved in the negative pathway and the loss of gene involved in positive pathway can

suppress the relevant functions. More specifically, the presence or the loss of genes in WT

might lead to changes in these four functional categories that are beneficial to its adaptation to

high altitude, as have been reported for other high-altitude adapted species and populations of

humans and animals[4,7,11,15,18,19,22,23,31]. And the loss or the presence of the same genes in BR

might lead to opposite changes of these four functional categories that are beneficial to its

adaptation to low altitude. For example, we found that gene *PIK3C2G* in BR is pseudogenized in

WT. This gene is involved in many pathways, including the HIF pathway, endothelin signaling

pathway, angiogenesis pathway, VEGF signaling pathway, T cell activation pathway and

inflammation mediated by chemokine and cytokine signaling pathway (Figure 3b). It is likely

that functional *PIK3C2G* in BR plays a role in BR's possibly low demands for these cardiovascular

and immune functions to adapt to low altitudes, while the loss-of-function of *PIK3C2G* in WT

might better meet WT's possibly high demands for these functions to adapt to high altitudes.

Moreover, gene *CACNB1* in WT is pseudogenized in BR. This gene is involved in the beta1/2

adrenergic receptor signaling pathway, metabotropic glutamate receptor group III pathway,

nicotinic acetylcholine receptor signaling pathway and thyrotropin-releasing hormone receptor

signaling pathway (Figure 3b). It is likely that functional *CACNB1* in WT might play a role in WT's

possibly high demands for these neuronal and energy metabolic functions to adapt to high

altitudes or other aspects of its unique niches, while the loss-of-function of *CACNB1* in BR might

better meet BR's possibly low demands for these functions to adapt to low altitudes or other

aspects of its unique niches.

Next, we compared the SNPs in the WT, BL and BR populations. Interestingly, we found

that genes in the selective sweeps in each species are mainly involved in the same GO pathways

of the four functional categories as are the unique genes and pseudogenes in the WT and BR

genomes, even though the two sets of genes have little overlaps (Figure 7a). Most genes in the

selective sweeps do not have amino acid-altering mutations, although some may have missense

mutations such as *MYO3B* in BR and *PTPRC* in WT (Figure 5d), become pseudogenes such as

CACNB2, or be missing such as *PRKCD* in BR. These results suggest that natural selection might

act on their *cis*-regulatory sequences, thereby altering their expression levels in relevant tissues.

It appears that natural selection on these genes might be related to the adaptation of the

species to different niches, particularly different altitudes. For example, in the BL vs BR

comparison, gene *MAPK1* in BL is included in a selective sweep. This gene is involved in the

VEGF signaling pathway, angiogenesis pathway, endothelin signaling pathway, angiotensin II-

stimulated signaling through G proteins and beta-arrestin pathway, insulin/IGF pathway-

mitogen activated protein kinase kinase/MAP kinase cascade pathway, and many pathways of

the immunity (Figure 7b). It is likely that selection on *MAPK1* might be related to the adaptation

of BL to intermediate altitude. In the WT vs BL comparison, gene *PPARG* is included in a

selective sweep, and the gene is involved in the HIF pathway[76] (Figure 7b), thus, selection on

the gene might be related to the adaptation of WT to low oxygen pressure at high altitudes.

It is not surprising that the unique genes/pseudogenes in WT and BR and genes in the

selective sweeps of BL, BR and WT converge on the same sets of pathways involved in

cardiovascular, energy metabolic, neuronal and immune functions. On the one hand, high

altitude niches with low oxygen pressure, low temperature and less availability of food would

pose a direct pressure on WT's cardiovascular system and energy metabolism functions, while

such pressure would be somewhat relaxed on BL and even more relaxed on BR. On the other

hand, other altitude related ecological factors might exert different pressures on the three

species for their neuronal functions for food foray and escaping for predators, and for immune

functions for different pathogens at varying altitudes. Indeed, it has been shown that high

altitude niches can have effects on the neuronal functions such as sensory perception in

domestic yaks (*Bos grunniens*)[16], and olfaction in Tibetan pig[7], wild boars[7] and ground tit (*Parus*

humilis)[15]. In this study, we found that a nonsynonymous mutation in *MYO3B* in the BR

population might alter their auditory sensation. It has also been shown that high altitude niches

can have effects on immune functions in ground tit[15]. In this study, we found that a

nonsynonymous mutation in *PTPRC* in the WT population and Manakins might alter their T-cell

functions. Thus, it appears that the three eared pheasant species might adapt to highly varying

altitudes or other aspects of their distinct ecological niches by loss-of-function mutations and

fine-tuning genes involved in the same set of pathways to meet their varying demands for the

four major functions. In other words, genes in this set of pathways might play roles in the

adaptation of the three pheasant species to different niches, particularly, different altitudes by

complete deletion (gene missing), pseudogenization, alteration of CDSs, or alteration of *cis*-

regulatory sequences.

Finally, to our best knowledge, we provided thus far the most comprehensive

annotation of genes and pseudogenes in the WT and BR genomes. We also identified numerous

selection sweeps in the WT, BR, and BL genomes. These results can be valuable resources for

researchers to better understand the biology of these uniquely adapted species and for policy

makers to design strategies for the preservation of these endangered species.

Materials and Methods

Datasets

We downloaded our previously published data[46], including the assembled genome of the WT

individual from GenBank with accession number GCA_036346035.1, re-sequencing data of 10

WT individuals from NCBI SRA with accession number SRP433016, re-sequencing data of a BL

individual from NCBI SRA with accession number SRP471294, and RNA-seq data from 20 tissues

(chest muscle, leg muscle, liver, kidney, lung, spleen, heart, cerebellum, brain cortex, ovary,

glandular, pancreas, abdominal fat, skin from chest, crops, rectum, cecum, duodenum, small

intestine and gizzard) of the WT individual from NCBI SRA with accession number SRP432961.

We downloaded the published datasets[45], including the assembled genome of the BR individual

from China National Genomics Data Center (CNGDC) with accession number

GWHAOPW00000000, re-sequencing data of 41 BR individuals from CNGDC with project

number PRJCA003284, RNA-seq data from three BR tissues (blood, developing primaries, and

developing tail feathers) from CNGDC with project number PRJCA003284, and re-sequencing

data of 12 BL individuals from CNGDC with project number PRJCA003284. We downloaded re-

sequencing data of 24 individuals of an indigenous chicken (Daweishan chicken) (*Gallus gallus*)

breed[67] from the NCBI SRA database with project number PRJNA893352.

Ethics approval

All the experimental procedures were approved by the Animal Care and Use Committee of the

Yunnan Agricultural University (approval ID: YAU202103047). The care and use of animals fully

complied with local animal welfare laws, guidelines, and policies.

Real-time quantitative PCR (RT-qPCR) analysis

Three female adult WT individuals were collected from Diqing Tibet Autonomous Prefecture,

Yannan Province, China. The birds were killed by electric shock to unconsciousness followed by

neck artery bleeding. One to two grams of 20 tissues were aseptically collected from each

individual bird in a centrifuge tube within 20 mins after sacrifice and immediately frozen in

liquid nitrogen, then stored at -80°C until use. The collected tissues included chest muscle, leg

muscle, liver, kidney, lung, spleen, heart, cerebellum, brain cortex, ovary, glandular, pancreas,

abdominal fat, skin from chest, crops, rectum, cecum, duodenum, small intestine and gizzard.

Total RNA from each tissue sample was extracted using TaKaRa MiniBEST Universal RNA

Extraction Kit (TaKaRa Biotechnology Co., Ltd., Dalian, P. R. China) according to the vendor's

instructions. cDNA was synthetized from the total RNA by using a PrimeScript RT reagent Kit

(TaKaRa Biotechnology Co., Ltd., Dalian, P. R. China) following the vendor's instructions as

previously described[88]. RT-qPCR was performed using the Bio-Rad CFX96 real-time PCR

platform (Bio-Rad Laboratories, Inc.) and SYBR Green master mix (iQTM SYBRGreen® Supermix,

Dalian TaKaRa Biotechnology Co. Ltd.). We randomly selected 17 putative new genes for

validation. The primers of the 17 putative new genes and their annealing temperatures are

listed in Supplementary Note. The β-actin gene was used as a reference. Primers were

commercially synthesized (Shanghai Shenggong Biochemistry Company, P.R.C.). Each PCR

reaction was performed in 25 μl volumes containing 12.5 μl of iQ™ SYBR Green Supermix, 0.5 μl

(10 mM) of each primer, and 1 μl of cDNA. Detection of products was performed with the

following cycle profile: one cycle of 95 °C for 2 min, and 40 cycles of 95 °C for 15 s, the

annealing temperature (Supplementary Note) for 30 s, and 72 °C for 30 s, followed by a final

cycle of 72 °C for 10 min. The $2^{-\Delta Ct}$ method was used to analyze mRNA abundance. All tissues

were analyzed with three biological replicates and each biological replicate with five technical

replicates.

Protein-coding gene annotation

To annotate the protein-coding genes and pseudogenes in the WT and BR genomes, we used a

combination of reference-based and RNA-based method as previously described[89]. Briefly, for

the reference-based method, we used the CDSs of 53 well-represented avian genomes in NCBI

(Table S1) as the templates. We mapped all the CDSs isoforms of genes in the 53 genomes to

the assembly using Splign (2.0.0)[90] using default settings. For each template gene whose CDSs

could be mapped to the assembly, we concatenated all the mapped parts on the assembly. If

the CDSs of the template gene could be mapped to multiple loci on the assemblies, we chose

the locus with the highest mapping identity and checked whether the resulting concatenated

sequence formed an intact ORF (it had a length of an integer multiple of three and contained no

premature stop codon). If resulting concatenated sequence formed an intact ORF, we called the

locus an intact gene. If resulting concatenated sequence did not form an intact ORF, i.e. its

length was not an integer multiple of three and/or it contained a premature stop codon, we

mapped the Illumina DNA short reads from the same individual to the locus in the assembly

allowing no mismatch and gaps using bowtie (v2.4.1)[91] with default settings. If each nucleotide

position in the locus can be mapped by at least 10 short reads, we considered the locus as a

pseudogene; otherwise, we called it a partially supported gene since the pseudogenization

mutation might be caused by the sequencing errors in the assembly that cannot be corrected

by the polishing process.

560   For the RNA-based method, we first mapped all the RNA-seq reads from the 20 tissues

561  of WT (Table S3) and the three tissues of BR (Table S4) to the rRNA database SILVA_138[92] using

562  bowtie (v2.4.1)[91] with default settings, and filtered out the mapped reads. We then mapped the

563  unaligned reads to each genome using STAR (2.7.0c)[93] with default settings. As the *Crossoptilon*

564  species are closely related[35,36], here, we used a total of 23 RNA-seq datasets from both species

565  in annotating each genome to fully utilize the available data. Particularly, the datasets of BR

566  were collected from tissues (blood, developing primaries, and developing tail feathers) that

567  were not collected from WT, thus, they were complementary to facilitate gene finding. Based

568  on the mapped reads to each genome, we assembled transcripts using genome-guided method

569  in Trinity (2.8.5)[94] with default settings. Then we mapped the assembled transcripts to their

570  corresponding genome using Splign (2.0.0)[90] with default settings, and defined their gene

571  structures, and removed those that partially overlap the genes and pseudogenes predicted by

572  the reference-based method or the non-coding RNA genes (see below). For each unmapped

573  assembled transcripts, we found the longest ORF in it, and called it a protein-coding gene if it

574  was at least 300 bp long.

575   For completeness, we also annotated tRNA, miRNA, rRNA, snoRNA, telomerase RNA and

576  SRP RNA genes in each genome using infernal (1.1.2)[95] with the Rfam (v.14) database[96] as the

577  reference using default settings.

578  **Single nucleotide variants calling**

579  To call the variants in the populations of BL, BR, WT and Daweishan chicken, we first mapped

580  the short reads of each individual to the WT genome using BWA (0.7.17)[97] and SAMtools (1.9)[98]

581  using default settings, and then called the variants for each individual using GATK-

HaplotypeCaller (4.1.6)[99] using default settings and merged the variants of each individual from

the same species using GATK-CombineGVCFs (4.1.6)[99] using default settings. We removed the

SNPs with Quality by depth (QD) < 2, Fisher strand (FS) > 60, Root mean square mapping quality

(MQ) < 40, Strand odd ratio (SOR) > 3, Rank Sum Test for mapping qualities (MQRankSum) < -

12.5 and Rank Sum Test for site position within reads (ReadPosRankSum) < -8 and indels with

QD < 2, FS > 200, SOR > 10, Likelihood-based test for the consanguinity among samples

(InbreedingCoeff) < -0.8 and ReadPosRankSum < -20.

To calculate the fixation rate of the pseudogenes in the population of each species, we

called the SNPs in the pseudogenes of each species using the DNA short reads from the species'

population (n=10 for WT and n=41 for BR) using the same method mentioned above. The

annotated pseudogenes of each species were used as the reference for the SNP calling.

PCA, ancestry, and introgression analyses

The called biallelic SNPs in each species located in strong linkage disequilibrium blocks and

those with a MAF<0.05 were filtered. Principal component analysis (PCA) was performed on the

remaining SNPs using PLINK (1.90)[100] with default settings. The same SNPs were used to infer

the ancestral relationships of the three species using ADMIXTURE (1.3.0)[62] with default settings.

The results for the number of ancestral groups K = 2 and 3 were tested. The same SNPs were

used to infer possible gene introgression among the three species by computing the D-statistic

using Dsuite[63] with default settings and the f3-statistic using ADMIXTOOLS[64] with default

settings.

Calculation of allele frequencies of pseudogenes

We computed allele frequencies of the first pseudogenization mutation of each pseudogene in

each species using GATK (4.1.6)[99] based on called SNPs and indels.

Selective sweeps detection

Selective sweeps in a species relative to another were detected using two methods including

genetic differentiation (F_{ST})[101] and difference in nucleotide diversity $(\Delta\pi)$[68]. These two methods

were widely used to detect selective sweeps by many studies[65-68]. We estimated F_{ST} for each

pair of comparison using VCFtools (0.1.16)[102] with a sliding window of 40 kb and a step size of

20 kb with default settings. We estimated π for each species using VCFtools (0.1.16)[102] with a

sliding window of 40 kb and a step size of 20 kb with default settings, and calculated the

difference in nucleotide diversity $(\Delta\pi)$ in each window for each pair of comparison. To evaluate

the statistical significance of the F_{ST} and π values for a comparison, we generated a Null model

by shuffling the allele frequency data for 100 times while keeping the SNP positions fixed[69]. We

then computed F_{ST} and $\Delta\pi$ for the permuted windows as well as their means $(\mu(F_{ST_{Null}})$ and

$\mu(\Delta\pi_{Null}))$ and standard deviations $(\sigma(F_{ST_{Null}})$ and $\sigma(\Delta\pi_{Null}))$. We computed the Z value for

each F_{ST} and $\Delta\pi$ value for the comparison by using the following formulas:

$ZF_{ST}(i) = (F_{ST}(i) - \mu(F_{ST_{Null}}))/\sigma(F_{ST_{Null}})$ and

$Z\Delta\pi(i) = (\Delta\pi(i) - \mu(\Delta\pi_{Null}))/\sigma(\Delta\pi_{Null})$.

We considered a window with $ZF_{ST} > \alpha$ and $Z\Delta\pi > \beta$ or $Z\Delta\pi < -\beta$ ($\alpha>0$, $\beta>0$) as a

selective sweep. For the selective sweeps in each pair of comparison, we used the value of $Z\Delta\pi$

to help identify the selective sweeps of each species. Windows with $Z\Delta\pi$ scores < -β represent

selective sweeps of the minuend species, and windows with $Z\Delta\pi$ scores > β represent selective

sweeps of the subtrahend species. We set α = 2.33 and β = 1.64, corresponding to a one-tailed

and a two-tailed z-test, respectively.

Runs of homozygosity (ROH) analysis

ROH analysis in each species was done using BCFtools (1.10)[103] with the default settings based

on the SNPs called in each individual bird of the species.

Tajima's D value calculations

Tajima's D in each species was calculated using ANGSD (0.930)[104] with default settings based on

the SNPs called in each individual bird of the species.

RNA-seq data analyses

RNA-seq reads of each tissue were mapped to the WT or BR genome using STAR (2.7.0c)[93] with

default settings. The expression level of each gene or pseudogene g_i in the tissue was

computed as TPM (transcript per million), defined as $TPM_i = \frac{r_i/l_i}{\sum_j r_j/l_j} * 10^6$, where r_i denotes

reads mapped to g_i, l_i is the length of g_i, and $\sum_j r_j/l_j$ corresponds to the sum of mapped reads

to each g_j normalized by its length.

Hierarchical clustering of gene expression levels

The expression level of each gene g_i in a tissue was rescaled as $s_i = \frac{e_i - e_{min}}{e_{max} - e_{min}}$,

where e_i denotes the expression level of g_i, and e_{min} and e_{max} the minimal and maximal

expression levels of all genes in the tissue, respectively. The clustermap program from seaborn

in Python was used to generate the heatmaps in Figure 1.

Data Availability
The WT genome is available at GenBank with accession number GCA_036346035.1; the BR
genome is available at CNGDC with accession number GWHAOPW00000000; re-sequencing
data of 10 WT individuals are available at NCBI SRA with accession number SRP433016; re-

sequencing data of 41 BR individuals are available at CNGDC with project number PRJCA003284;
re-sequencing data of 12 BL individuals are available at CNGDC with project number
PRJCA003284 and at NCBI SRA with accession number SRP471294; RNA-seq data from 20
tissues of the WT individual are available at NCBI SRA with accession number SRP432961; RNA-
seq data from the BR individual's blood, developing primaries and developing tail feathers
tissues are available at CNGDC with project number PRJCA003284. Our gene annotation results
of the WT and BR genomes are available at FigShare
(https://doi.org/10.6084/m9.figshare.24438295.v1).
655

656 **Code Availability**
657 All gene annotation code and the corresponding pipeline description are available at
658 https://github.com/zhengchangsulab/A-genome-assebmly-and-annotation-pipeline.
659

660 **Author Contributions**
661 CG, JJ, ZS and TD supervised and conceived the project; KW, XG and DW collected tissue
662 samples and conducted molecular biology experiments; SW and SY assembled and annotated
663 the genomes; SW and ZS performed data analysis; and SW and ZS wrote the manuscript.
664

665 **Competing interests**
666 The authors declare that they have no competing interests.
667

Funding
This work was supported by the National Natural Science Foundation of China (U2002205 and
U1702232), Yunling Scholar Training Program of Yunnan Province (2014NO48), Yunling Industry
and Technology Leading Talent Training Program of Yunnan Province (YNWR-CYJS-2015-027),
Natural Science Foundation of Yunnan Province (2019IC008 and 2016ZA008), and Department
of Bioinformatics and Genomics of the University of North Carolina at Charlotte.

References
676

1   Zhang, P. *et al.* Denisovans and Homo sapiens on the Tibetan Plateau: dispersals and
adaptations. *Trends Ecol Evol* **37**, 257-267 (2022).
https://doi.org:10.1016/j.tree.2021.11.004
2   Peng, Y. *et al.* Genetic variations in Tibetan populations and high-altitude adaptation at
the Himalayas. *Mol Biol Evol* **28**, 1075-1081 (2011).
https://doi.org:10.1093/molbev/msq290
3   Hu, H. *et al.* Evolutionary history of Tibetans inferred from whole-genome sequencing.
PLoS Genet **13**, e1006675 (2017). https://doi.org:10.1371/journal.pgen.1006675
4   Wang, M. S. *et al.* Genomic Analyses Reveal Potential Independent Adaptation to High
Altitude in Tibetan Chickens. *Mol Biol Evol* **32**, 1880-1889 (2015).
https://doi.org:10.1093/molbev/msv071
5   Li, S. *et al.* A non-synonymous SNP with the allele frequency correlated with the altitude
may contribute to the hypoxia adaptation of Tibetan chicken. *PLoS One* **12**, e0172211
(2017). https://doi.org:10.1371/journal.pone.0172211

6   Yuan, J. *et al.* Genome-wide run of homozygosity analysis reveals candidate genomic
regions associated with environmental adaptations of Tibetan native chickens. *BMC*
Genomics **23**, 91 (2022). https://doi.org:10.1186/s12864-021-08280-z

7   Li, M. *et al.* Genomic analyses identify distinct patterns of selection in domesticated pigs
and Tibetan wild boars. *Nat.Genet* **45**, 1431-1438 (2013).
https://doi.org:10.1038/ng.2811

8   Zhang, B. *et al.* Comparative transcriptomic and proteomic analyses provide insights into
the key genes involved in high-altitude adaptation in the Tibetan pig. *Sci Rep* **7**, 3654
(2017). https://doi.org:10.1038/s41598-017-03976-3

9   Li, C. *et al.* Markhor-derived Introgression of a Genomic Region Encompassing PAPSS2
Confers High-altitude Adaptability in Tibetan Goats. *Mol Biol Evol* **39** (2022).
https://doi.org:10.1093/molbev/msac253

10   Hu, X. J. *et al.* The Genome Landscape of Tibetan Sheep Reveals Adaptive Introgression
from Argali and the History of Early Human Settlements on the Qinghai-Tibetan Plateau.
Mol Biol Evol **36**, 283-303 (2019). https://doi.org:10.1093/molbev/msy208

11   Liu, J. *et al.* Genetic signatures of high-altitude adaptation and geographic distribution in
Tibetan sheep. *Sci Rep* **10**, 18332 (2020). https://doi.org:10.1038/s41598-020-75428-4

12   Li, Y. *et al.* Population variation revealed high-altitude adaptation of Tibetan mastiffs.
Mol Biol Evol **31**, 1200-1205 (2014). https://doi.org:10.1093/molbev/msu070

13   Wu, D. D. *et al.* Pervasive introgression facilitated domestication and adaptation in the
Bos species complex. *Nature ecology & evolution* **2**, 1139-1145 (2018).
https://doi.org:10.1038/s41559-018-0562-y

14   Simonson, T. S., McClain, D. A., Jorde, L. B. & Prchal, J. T. Genetic determinants of Tibetan
high-altitude adaptation. *Hum Genet* **131**, 527-533 (2012).
https://doi.org:10.1007/s00439-011-1109-3

15   Qu, Y. *et al.* Ground tit genome reveals avian adaptation to living at high altitudes in the
Tibetan plateau. *Nat Commun* **4**, 2071 (2013). https://doi.org:10.1038/ncomms3071

16   Qiu, Q. *et al.* The yak genome and adaptation to life at high altitude. *Nat.Genet* **44**, 946-
949 (2012). https://doi.org:10.1038/ng.2343

17   Simonson, T. S. *et al.* Genetic evidence for high-altitude adaptation in Tibet. *Science* **329**,
72-75 (2010). https://doi.org:10.1126/science.1189406

18   Ji, L. D. *et al.* Genetic adaptation of the hypoxia-inducible factor pathway to oxygen
pressure among eurasian human populations. *Mol Biol Evol* **29**, 3359-3370 (2012).
https://doi.org:10.1093/molbev/mss144

19   Xiong, Y. *et al.* Physiological and genetic convergence supports hypoxia resistance in
high-altitude songbirds. *PLoS Genet* **16**, e1009270 (2020).
https://doi.org:10.1371/journal.pgen.1009270

20   Peng, Y. *et al.* Down-Regulation of EPAS1 Transcription and Genetic Adaptation of
Tibetans to High-Altitude Hypoxia. *Mol Biol Evol* **34**, 818-830 (2017).
https://doi.org:10.1093/molbev/msw280

21   Xin, J. *et al.* Chromatin accessibility landscape and regulatory network of high-altitude
hypoxia adaptation. *Nat Commun* **11**, 4928 (2020). https://doi.org:10.1038/s41467-020-
18638-8

22 Xu, D. *et al.* A single mutation underlying phenotypic convergence for hypoxia
adaptation on the Qinghai-Tibetan Plateau. *Cell Res* **31**, 1032-1035 (2021).
https://doi.org:10.1038/s41422-021-00517-6
23 Zhu, X. *et al.* Divergent and parallel routes of biochemical adaptation in high-altitude
passerine birds from the Qinghai-Tibet Plateau. *Proc Natl Acad Sci U S A* **115**, 1865-1870
(2018). https://doi.org:10.1073/pnas.1720487115
24 Guo, Y. B. *et al.* GCH1 plays a role in the high-altitude adaptation of Tibetans. *Zool Res* **38**,
155-162 (2017). https://doi.org:10.24272/j.issn.2095-8137.2017.037
25 Heinrich, E. C. *et al.* Genetic variants at the EGLN1 locus associated with high-altitude
adaptation in Tibetans are absent or found at low frequency in highland Andeans.
Annals of human genetics **83**, 171-176 (2019). https://doi.org:10.1111/ahg.12299
26 Xiang, K. *et al.* Identification of a Tibetan-specific mutation in the hypoxic gene EGLN1
and its contribution to high-altitude adaptation. *Mol Biol Evol* **30**, 1889-1898 (2013).
https://doi.org:10.1093/molbev/mst090
27 Jeong, C. *et al.* Admixture facilitates genetic adaptations to high altitude in Tibet. *Nat*
Commun **5**, 3281 (2014). https://doi.org:10.1038/ncomms4281
28 Li, D. *et al.* Population genomics identifies patterns of genetic diversity and selection in
chicken. *BMC Genomics* **20**, 263 (2019). https://doi.org:10.1186/s12864-019-5622-4
29 Liu, X., Wang, X., Liu, J., Wang, X. & Bao, H. Identifying Candidate Genes for Hypoxia
Adaptation of Tibet Chicken Embryos by Selection Signature Analyses and RNA
Sequencing. *Genes* **11** (2020). https://doi.org:10.3390/genes11070823
30 Tang, Q. *et al.* HIF-1 regulates energy metabolism of the Tibetan chicken brain during
embryo development under hypoxia. *Am J Physiol Regul Integr Comp Physiol* **320**, R704-
r713 (2021). https://doi.org:10.1152/ajpregu.00052.2020
31 Zhang, Y., Zhang, H., Zhang, B., Ling, Y. & Zhang, H. Identification of key HIF-1α target
genes that regulate adaptation to hypoxic conditions in Tibetan chicken embryos. *Gene*
729, 144321 (2020). https://doi.org:10.1016/j.gene.2019.144321
32 Scheinfeldt, L. B. & Tishkoff, S. A. Living the high life: high-altitude adaptation. *Genome*
Biol **11**, 133 (2010). https://doi.org:10.1186/gb-2010-11-9-133
33 Pamenter, M. E., Hall, J. E., Tanabe, Y. & Simonson, T. S. Cross-Species Insights Into
Genomic Adaptations to Hypoxia. *Front Genet* **11**, 743 (2020).
https://doi.org:10.3389/fgene.2020.00743
34 Friedrich, J. & Wiener, P. Selection signatures for high-altitude adaptation in ruminants.
Anim Genet **51**, 157-165 (2020). https://doi.org:10.1111/age.12900
35 Li, X., Huang, Y. & Lei, F. Comparative mitochondrial genomics and phylogenetic
relationships of the Crossoptilon species (Phasianidae, Galliformes). *BMC Genomics* **16**,
42 (2015). https://doi.org:10.1186/s12864-015-1234-9
36 Li, X. *et al.* Assessment of genetic diversity in Chinese eared pheasant using fluorescent-
AFLP markers. *Mol Phylogenet Evol* **57**, 429-433 (2010).
https://doi.org:10.1016/j.ympev.2010.05.024
37 Pang, X., Ma, J. & Liu, W. The research of the main variables of habitat selection in brown
eared-pheasant breeding in XiaoWutai Mountain. *Journal of Hebei Forestry Science and*
Technology **(Supplementary Issue)**, 27-29 (2009).

38   Zhao, Z. *Ornithology of China (Non-Passeriformes)*. Vol. 1 376-381 (Jilin Science and
Technology Press, 2001).
39   Xu, Z., Lei, Y., Jin, X. & Yuan, H. Brown eared-pheasant population was found in
Huanglong Mountain, Shanxi. *Chinese Wildlife* **19**, 13 (1998).
40   TH, C. *et al. Fauna of China*. Vol. IV 106–107 (Beijing: Science Press, 1978).
41   *Crossoptilon harmani IUCN Red List of Threatened Species*, 2016).
42   Li, Y., Li, X., Song, Z. & Ding, C. Determining the distribution loss of brown eared-
pheasant (Crossoptilon mantchuricum) using historical data and potential distribution
estimates. *PeerJ* **4**, e2556 (2016). https://doi.org:10.7717/peerj.2556
43   Wang, P. *et al.* The role of niche divergence and geographic arrangement in the
speciation of Eared Pheasants (Crossoptilon, Hodgson 1938). *Mol Phylogenet Evol* **113**,
1-8 (2017). https://doi.org:10.1016/j.ympev.2017.05.003
44   Zheng, Z. *A complete checklist of species and subspecies of the Chinese birds.* (Science
Press, 1994).
45   Wang, P. *et al.* Genomic Consequences of Long-Term Population Decline in Brown Eared
Pheasant. *Mol Biol Evol* **38**, 263-273 (2021). https://doi.org:10.1093/molbev/msaa213
46   Wu, S. *et al.* High-quality genome assembly of a C. crossoptilon and related functional
and genetics data resources. *Sci Data* **11**, 247 (2024). https://doi.org:10.1038/s41597-
024-03087-5
47   Wu, S. De Novo Assemblies of Genomes of Four Indigenous Chickens and a White Eared
Pheasant Reveal Genetic Basis of Chicken Domestication and Altitude Adaptation.
Dissertation (2023). https://doi.org:https://graduateschool.charlotte.edu/de-novo-
assemblies-genomes-four-indigenous-chickens-and-white-eared-pheasant-reveal-
genetic-basis/
48   Sayers, E. W. *et al.* Database resources of the national center for biotechnology
information. *Nucleic Acids Res* **50**, D20-d26 (2022).
https://doi.org:10.1093/nar/gkab1112
49   Lovell, P. V. *et al.* Conserved syntenic clusters of protein coding genes are missing in birds.
Genome Biol **15**, 565 (2014). https://doi.org:10.1186/s13059-014-0565-1
50   Zhang, Z. D., Frankish, A., Hunt, T., Harrow, J. & Gerstein, M. Identification and analysis of
unitary pseudogenes: historic and contemporary gene losses in humans and other
primates. *Genome Biol* **11**, R26 (2010). https://doi.org:10.1186/gb-2010-11-3-r26
51   Derks, M. F. L. *et al.* A survey of functional genomic variation in domesticated chickens.
Genet Sel Evol **50**, 17 (2018). https://doi.org:10.1186/s12711-018-0390-1
52   Parmley, J. L. & Hurst, L. D. How do synonymous mutations affect fitness? *Bioessays* **29**,
515-519 (2007). https://doi.org:10.1002/bies.20592
53   Ng, P. C. *et al.* Genetic variation in an individual human exome. *PLoS Genet* **4**, e1000160
(2008). https://doi.org:10.1371/journal.pgen.1000160
54   Semenza, G. L. Targeting HIF-1 for cancer therapy. *Nat Rev Cancer* **3**, 721-732 (2003).
https://doi.org:10.1038/nrc1187
55   Giaccia, A., Siim, B. G. & Johnson, R. S. HIF-1 as a target for drug development. *Nat Rev
Drug Discov* **2**, 803-811 (2003). https://doi.org:10.1038/nrd1199
56   Bárdos, J. I. & Ashcroft, M. Hypoxia-inducible factor-1 and oncogenic signalling.
Bioessays **26**, 262-269 (2004). https://doi.org:10.1002/bies.20002

57 Matsumoto, T. & Claesson-Welsh, L. VEGF receptor signal transduction. *Sci STKE* **2001**,
re21 (2001). https://doi.org:10.1126/stke.2001.112.re21
58 Cross, M. J., Dixelius, J., Matsumoto, T. & Claesson-Welsh, L. VEGF-receptor signal
transduction. *Trends Biochem Sci* **28**, 488-494 (2003). https://doi.org:10.1016/s0968-
0004(03)00193-2
59 Ferrara, N., Gerber, H. P. & LeCouter, J. The biology of VEGF and its receptors. *Nat Med* **9**,
669-676 (2003). https://doi.org:10.1038/nm0603-669
60 Chatterjee, O. *et al.* An overview of the oxytocin-oxytocin receptor signaling network. *J*
Cell Commun Signal **10**, 355-360 (2016). https://doi.org:10.1007/s12079-016-0353-7
61 Siwen Wu, K. W., Tengfei Dou, Sisi Yuan, Dong-Dong Wu, Changrong Ge, Junjing
Jia,Zhengchang Su. High-quality genome assembly of a white eared pheasant individual
and related functional and genetics data resources. *BioRxiv* (2023).
https://doi.org:https://www.biorxiv.org/content/10.1101/2023.11.09.566452v1
62 Alexander, D. H., Novembre, J. & Lange, K. Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res* **19**, 1655-1664 (2009).
https://doi.org:10.1101/gr.094052.109
63 Malinsky, M., Matschiner, M. & Svardal, H. Dsuite - Fast D-statistics and related
admixture evidence from VCF files. *Mol Ecol Resour* **21**, 584-595 (2021).
https://doi.org:10.1111/1755-0998.13265
64 Patterson, N. *et al.* Ancient admixture in human history. *Genetics* **192**, 1065-1093 (2012).
https://doi.org:10.1534/genetics.112.145037
65 Wang, Y. M. *et al.* Integrating Genomic and Transcriptomic Data to Reveal Genetic
Mechanisms Underlying Piao Chicken Rumpless Trait. *Genomics Proteomics*
Bioinformatics **19**, 787-799 (2021). https://doi.org:10.1016/j.gpb.2020.06.019
66 Qanbari, S. *et al.* Genetics of adaptation in modern chicken. *PLoS Genet* **15**, e1007989
(2019). https://doi.org:10.1371/journal.pgen.1007989
67 Wu, S. *et al.* Artificial selection footprints in indigenous and commercial chicken
genomes. *BMC Genomics* **25**, 428 (2024). https://doi.org:10.1186/s12864-024-10291-5
68 Wang, M. S. *et al.* Positive selection rather than relaxation of functional constraint drives
the evolution of vision during chicken domestication. *Cell Res* **26**, 556-573 (2016).
https://doi.org:10.1038/cr.2016.44
69 Churchill, G. A. & Doerge, R. W. Empirical threshold values for quantitative trait mapping.
Genetics **138**, 963-971 (1994). https://doi.org:10.1093/genetics/138.3.963
70 Lelli, A. *et al.* Class III myosins shape the auditory hair bundles by limiting microvilli and
stereocilia growth. *J Cell Biol* **212**, 231-244 (2016).
https://doi.org:10.1083/jcb.201509017
71 Josep del Hoyo, A. E., Jordi Sargatal, David A Christie.
https://en.wikipedia.org/wiki/Handbook_of_the_Birds_of_the_World, 1992).
72 Wu, L., Fu, J. & Shen, S. H. SKAP55 coupled with CD45 positively regulates T-cell
receptor-mediated gene transcription. *Mol Cell Biol* **22**, 2673-2686 (2002).
https://doi.org:10.1128/mcb.22.8.2673-2686.2002
73 D'Orléans-Juste, P., Plante, M., Honoré, J. C., Carrier, E. & Labonté, J. Synthesis and
degradation of endothelin-1. *Can J Physiol Pharmacol* **81**, 503-510 (2003).
https://doi.org:10.1139/y03-032

74 Neylon, C. B. Vascular biology of endothelin signal transduction. *Clin Exp Pharmacol*
Physiol **26**, 149-153 (1999). https://doi.org:10.1046/j.1440-1681.1999.03013.x

75 Masaki, T., Miwa, S., Sawamura, T., Ninomiya, H. & Okamoto, Y. Subcellular mechanisms
of endothelin action in vascular system. *Eur J Pharmacol* **375**, 133-138 (1999).
https://doi.org:10.1016/s0014-2999(99)00252-6

76 Graham, A. M. & McCracken, K. G. Convergent evolution on the hypoxia-inducible factor
(HIF) pathway genes EGLN1 and EPAS1 in high-altitude ducks. *Heredity (Edinb)* **122**, 819-
832 (2019). https://doi.org:10.1038/s41437-018-0173-z

77 Martindale, J. L. & Holbrook, N. J. Cellular response to oxidative stress: signaling for
suicide and survival. *J Cell Physiol* **192**, 1-15 (2002). https://doi.org:10.1002/jcp.10119

78 Davis, R. J. Signal transduction by the c-Jun N-terminal kinase. *Biochem Soc Symp* **64**, 1-
12 (1999). https://doi.org:10.1515/9781400865048.1

79 Kyriakis, J. M. & Avruch, J. Mammalian mitogen-activated protein kinase signal
transduction pathways activated by stress and inflammation. *Physiol Rev* **81**, 807-869
(2001). https://doi.org:10.1152/physrev.2001.81.2.807

80 Tibbles, L. A. & Woodgett, J. R. The stress-activated protein kinase pathways. *Cell Mol*
Life Sci **55**, 1230-1254 (1999). https://doi.org:10.1007/s000180050369

81 Leppä, S. & Bohmann, D. Diverse functions of JNK signaling and c-Jun in stress response
and apoptosis. *Oncogene* **18**, 6158-6162 (1999). https://doi.org:10.1038/sj.onc.1203173

82 Myung, P. S., Boerthe, N. J. & Koretzky, G. A. Adapter proteins in lymphocyte antigen-
receptor signaling. *Curr Opin Immunol* **12**, 256-266 (2000).
https://doi.org:10.1016/s0952-7915(00)00085-6

83 Krogsgaard, M., Huppa, J. B., Purbhoo, M. A. & Davis, M. M. Linking molecular and
cellular events in T-cell activation and synapse formation. *Semin Immunol* **15**, 307-315
(2003). https://doi.org:10.1016/j.smim.2003.09.002

84 Kane, L. P., Lin, J. & Weiss, A. Signal transduction by the TCR for antigen. *Curr Opin*
Immunol **12**, 242-249 (2000). https://doi.org:10.1016/s0952-7915(00)00083-2

85 Lukacs, N. W. Role of chemokines in the pathogenesis of asthma. *Nat Rev Immunol* **1**,
108-116 (2001). https://doi.org:10.1038/35100503

86 Vicente-Manzanares, M., Sancho, D., Yáñez-Mó, M. & Sánchez-Madrid, F. The leukocyte
cytoskeleton in cell migration and immune interactions. *Int Rev Cytol* **216**, 233-289
(2002). https://doi.org:10.1016/s0074-7696(02)16007-4

87 Rhie, A. *et al.* Towards complete and error-free genome assemblies of all vertebrate
species. *Nature* **592**, 737-746 (2021). https://doi.org:10.1038/s41586-021-03451-0

88 Zhao, J. *et al.* Transcriptomics reveals the molecular regulation of Chinese medicine
formula on improving bone quality in broiler. *Poult Sci* **102**, 103044 (2023).
https://doi.org:10.1016/j.psj.2023.103044

89 Wu, S. *et al.* Annotations of four high-quality indigenous chicken genomes identify more
than one thousand missing genes in subtelomeric regions and micro-chromosomes with
high G/C contents. *BMC Genomics* **25**, 430 (2024). https://doi.org:10.1186/s12864-024-
10316-z

90 Kapustin, Y., Souvorov, A., Tatusova, T. & Lipman, D. Splign: algorithms for computing
spliced alignments with identification of paralogs. *Biol Direct* **3**, 20 (2008).
https://doi.org:10.1186/1745-6150-3-20

91　Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
357-359 (2012). https://doi.org:10.1038/nmeth.1923
92　Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res* **41**, D590-596 (2013).
https://doi.org:10.1093/nar/gks1219
93　Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21 (2013).
https://doi.org:10.1093/bioinformatics/bts635
94　Grabherr, M. G. *et al.* Full-length transcriptome assembly from RNA-Seq data without a
reference genome. *Nat Biotechnol* **29**, 644-652 (2011). https://doi.org:10.1038/nbt.1883
95　Nawrocki, E. P. & Eddy, S. R. Infernal 1.1: 100-fold faster RNA homology searches.
Bioinformatics **29**, 2933-2935 (2013). https://doi.org:10.1093/bioinformatics/btt509
96　Kalvari, I. *et al.* Rfam 14: expanded coverage of metagenomic, viral and microRNA
families. *Nucleic Acids Res* **49**, D192-d200 (2021). https://doi.org:10.1093/nar/gkaa1047
97　Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics* **25**, 1754-1760 (2009).
https://doi.org:10.1093/bioinformatics/btp324
98　Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078-2079 (2009). https://doi.org:10.1093/bioinformatics/btp352
99　McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res* **20**, 1297-1303 (2010).
https://doi.org:10.1101/gr.107524.110
100　Chen, Z. L. *et al.* A high-speed search engine pLink 2 with systematic evaluation for
proteome-scale identification of cross-linked peptides. *Nat Commun* **10**, 3404 (2019).
https://doi.org:10.1038/s41467-019-11337-z
101　Reynolds, J., Weir, B. S. & Cockerham, C. C. Estimation of the coancestry coefficient: basis
for a short-term genetic distance. *Genetics* **105**, 767-779 (1983).
https://doi.org:10.1093/genetics/105.3.767
102　Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158
(2011). https://doi.org:10.1093/bioinformatics/btr330
103　Narasimhan, V. *et al.* BCFtools/RoH: a hidden Markov model approach for detecting
autozygosity from next-generation sequencing data. *Bioinformatics* **32**, 1749-1751
(2016). https://doi.org:10.1093/bioinformatics/btw044
104　Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next Generation
Sequencing Data. *BMC Bioinformatics* **15**, 356 (2014). https://doi.org:10.1186/s12859-
014-0356-4
944

Table 1: Gene annotation of two *Crossoptilon* species, *C. crossoptilon* (WT) and *C. mantchuricum* (BR), using reference- and RNA-based methods

Breeds	Reference-based			RNA-based		# Total genes	# Total pseudogenes
	# Intact	# Partial	# Pseudogenes	# NT-supported	# Novel		
WT	14,815	750	1,519	751	61	16,377	1,519
BR	14,518	209	1,976	631	52	15,410	1,976
Common	14,057		1,040	422	28	14,507	1,040

Table 2: Numbers of SNPs identified in the *C. auritum* (BL), *C. mantchuricum* (BR) and *C. crossoptilon* (WT) populations

Breeds	# individuals	# Total SNPs
BL	12	10,147,889
BR	41	6,849,438
WT	10	9,004,245
All	63	16,459,558

Table 3: The f3-statistic and associated z-scores for the three choices of (target; source 1, source 2) among the three species

Source 1	Source 2	Target	f3	std. err	Z-score
WT	BR	BL	-0.0586	0.0022	-26.45
WT	BL	BR	1.2340	0.0769	16.04
BR	BL	WT	0.8392	0.0047	180.24

Table 4: Summary of selective sweeps (SS) and discrete selective sweeps (DSS) identified in pairwise comparisons between three *Crossoptilon* species populations

Comparison	BL			BR			WT		
	# SS (# DSS)	# Genes in DSS	# Pathways involved	# SS (# DSS)	# Genes in DSS	# Pathways involved	# SS (# DSS)	# Genes in DSS	# Pathways involved
BL vs. BR	208 (128)	143	48	638 (463)	474	56	NA	NA	NA
BL vs. WT	92 (72)	67	17	NA	NA	NA	274 (181)	167	34
BR vs. WT	NA	NA	NA	306 (184)	196	45	359 (188)	167	36

Figure 1. Heatmaps of expression levels of new genes in tissues of *C. crossoptilon* (WT) and *C. mantchuricum* (BR). **a.** Expression pattern of the NT-supported new genes in 20 tissues of WT. **b.** Expression pattern of the novel genes in 20 tissues of WT. **c.** Expression pattern of the NT-supported new genes in three tissues of BR. **d.** Expression pattern of the novel genes in three tissues in BR. **e.** Expression pattern of the 17 selected new genes in WT measured by RT-qPCR. **f.** Expression pattern of the 17 selected new genes quantified using RNA-seq data in WT. The expression levels of genes are scaled such that 0 represents a gene that was not expressed and 1 represents a gene that has the highest expression level in the tissue (Materials and Methods). Figures a-d and f were based on TPMs, and e was based on RT-qPCR results.

Figure 2. Fixation rate of pseudogenes in the WT and BR populations and distribution of the first pseudogenization alleles along the CDSs of parental genes. **a.** Number of pseudogenization alleles with indicated frequencies in the BR population. **b.** Number of the pseudogenization alleles with indicated frequencies in the WT populations. **c.** Distribution of the first pseudogenization alleles in the BR genome along the CDSs of parental genes. Notably, 24.8%, 60.5%, and 14.7% of the first pseudogenization alleles are in the first 10%, middle 80% and last 10% of the CDS regions. **d.** Distribution of the first pseudogenization alleles in the WT genome along the CDSs of parental genes. Notably, 25.7%, 57.8% and 16.5% of the first pseudogenization alleles are in the first 10%, middle 80% and last 10% of the CDS region.

980
Figure 3. Functional analysis of the unique genes and pseudogene in the WT and BR genomes.
a. Venn diagram showing shared and unique genes (left) and pseudogenes (right) in the two
species. Of note, 658 WT genes are pseudogenized in BR and 173 BR genes are pseudogenized
in WT, and there are a total of 3,357 non redundant unique genes and pseudogenes in the two
genomes. **b.** The unique genes/pseudogenes in the WT and BR genomes that are involved in
pathways related to four functional categories: neuronal functions, cardiovascular functions,
energy metabolism, and immunity. Notably, unique genes in a species often became unique
pseudogenes in the other species.

989
990 **Figure 4.** Geographic distribution and population structure of the WT, BL and BR populations. **a.**
991 Geographic distribution of the three species used in this study. The color of the map indicates
992 altitude level (m). **b.** Principal component analysis of the populations of the three species based
993 on their SNP spectrum. **c.** Ancestral analysis of the individuals of the three species using
994 ADMIXTURE for K=2 and 3.

995
Figure 5. Distribution of the ZF_{ST} and $Z\Delta\pi$ values of each sliding window for the three pairwise
comparisons. **a.** BL VS. BR comparison. **b.** BL VS. WT comparison. **c.** BR VS. WT comparison. **d.**
Multiple alignments of part of CDSs of two genes in DSSs in different species. In subfigures a, b
and c, the top panel represents the distribution of $Z\Delta\pi$ values and the bottom right one
represents the distribution of ZF_{ST} values. Each dot in the bottom left panel represents a sliding
window with its $Z\Delta\pi$ and ZF_{ST} values being its coordinates in the plot. The dashed lines
represent the cut-offs for the ZF_{ST} and $Z\Delta\pi$ values.

1003
Figure 6. Manhattan plots for genome-wide analyses of the evolution of the three species. **a-c.**
Manhattan plots of ZF_{ST} for the three pair-wise comparisons of the species. The blue line
indicates the cut-off value. **d-f.** Manhattan plots of $Z|\Delta\pi|$ for the three pair-wise comparisons
of the species. The blue line indicates the cut-off value. **g-i.** Manhattan plots of Tajima's D
values for each species. **j-l.** Manhattan plots of the ratio of individuals that have long ROH
(>100kb) for each species. In the figures, each dot represents a sliding window. The horizontal
axes are pseudo chromosomes ordered in decreasing size.

1011
Figure 7. Comparison of unique genes and pseudogenes in the WT and BR genomes and genes
in selective sweeps in the three species. **a.** Venn diagram of the set of unique genes and
pseudogenes in the WT and BR genomes (unique set) and the set of genes in selective sweeps
in the three species (genes in SS). **b.** The genes with GO term assignment in selection sweeps in
each species that are involved in the same pathways of the same four functional categories as
are the unique genes and unique pseudogenes with GO term assignment in the WT and BR
genome, although the two sets only share four genes, i.e., *CAMK2G* and *VWF* are
pseudogenized in BR, *PRKCD* is missing in BR, and *PIK3C2G* is pseudogenized in WT (Figure 3b).