The maternal vGluT2 and embryonic mGluR3 signaling relay system controls

offspring wing dimorphism in pea aphid

Yiyang Yuan[1], Yanyan Wang[2,3], Wanwan Ye[2,3], Liqiang Xie[2], Erliang Yuan[2], Huijuan

Guo[2], Shifan Wang[4], Fang Dong[5], Keyan Zhu-Salzman[6], Feng Ge[1]*, Yucheng Sun[2]*

[1]Institute of Plant Protection, Shandong Academy of Agricultural Sciences, Jinan

250100, Shandong Province, China

[2]State Key Laboratory of Integrated Management of Pest Insects and Rodents, Institute

of Zoology, Chinese Academy of Sciences, Beijing 100101, China

[3]School of Life Sciences, Hebei University, Baoding 071002, China

[4]Department of Immunology and Infectious Diseases, Harvard T. H. Chan School of

Public Health, Boston, Massachusetts 02115, USA

[5]Xinjiang Production and Construction Corps Key Laboratory of Special Fruits and

Vegetables Cultivation Physiology and Germplasm Resources Utilization, Agriculture

College of Shihezi University, Shihezi 832003, China

[6]Department of Entomology, Texas A&M University, College Station, TX 77843

[*]Author for correspondence: Feng Ge (Tel: 86-531-66658008, Fax +86-0531-66657914,

email: gef@ioz.ac.cn); Yucheng Sun (Tel: 86-10-64807123, Fax +86-10-64807099,

email: sunyc@ioz.ac.cn).

Short running title: Transgenerational modulation of wing morph in aphid

Abstract

Transgenerational phenotypic plasticity (TPP) refers to the phenomenon that

environmental conditions experienced by one generation can influence the phenotype

of subsequent generations to adapt to the environment without modification of their

DNA sequences. Aphid wing dimorphism is a textbook example of TPP by which a

maternal aphid perceives the environmental cues to decide the wing morph of her

offspring. However, the signaling mechanism from mother to daughter remains unclear.

In this study, we showed that the population density and physical contact caused high

proportion of winged offspring in the pea aphid *Acyrthosiphon pisum*. Its *vesicular*

glutamate transporter 2 (*ApvGluT2*) and *metabotropic glutamate receptor 3*

(*ApmGluR3*) were identified by tissue-specific RNA-seq as differentially expressed

genes in the head and embryo respectively between solitary and more densely housed

maternal aphids. Elevated expression of brain *ApvGluT2* and embryonic *ApmGluR3* led

to increases in the winged proportion. Knockdown of either gene inhibited

phosphorylation of ApFoxO in embryos. Furthermore, EMSA showed that

dephosphorylated ApFoxO directly bound to the promotor of *hedgehog* (*ApHh*), a

morphogen gene for wing development, to repress its transcription in stage 20 embryos,

causing a lower winged proportion. Our results demonstrated that brain *vGluT2* and

embryonic *mGluR3* coordinately relayed the maternal physical contact signals and

control wing development in offspring, showcasing a novel regulatory mechanism

underlying physical contact-dependent, transgenerational wing dimorphism in aphids.

Keywords: Transgenerational phenotypic plasticity, *Acyrthosiphon pisum*, Wing

polymorphism, Glutamate signaling, FoxO

Introduction

The transgenerational phenotypic plasticity (TPP) is a fascinating capacity of individual genotypes, when experiencing different environmental conditions, to produce different phenotypes in subsequent generations (*Bell and Hellmann, 2019*). Since the impact of parental experience is expected to persist across generations, the phenotypic adjustment warranted by parents could be beneficial to offspring (*Guillaume et al., 2016*). This process involves detection of environmental cues by a parent, signal delivery, and signal perception by offspring and subsequent alternative phenotypes (*Bell and Hellmann, 2019*). Parent-to-offspring signaling can be accomplished via mechanisms that are rapid and non-genetic, such as through neurotransmitters, hormones, epigenetic modifications, and microbiota (*Donelan et al., 2020*).

Insect wing polyphenism is an intrinsic characteristic; a single species can produce flightless (wingless or short-wing) and flight-capable (winged or long-wing) morphs depending on the environment (*Brisson, 2010*; *Zera, 1997*; *Zhang et al., 2019*). Numerous originally winged insect species have become flightless because secondary flight loss favors their adaptation to environmentally stable habitats (*Roff 1990*; *Roff, 1994*; *Wagner and Liebherr, 1992*). Aphid wing dimorphism is an iconic example of TPP in that maternal aphids perceive the environmental cues, such as population density, nutrients, natural enemy, or even physical contact, and signal the offspring during the embryonic period, by which alate (winged) or apterous (wingless) daughters are destined before birth (*Braendle et al., 2005*; *Ishikawa and Miura, 2013*). Of the environmental cues, high population density is widely recognized as a biotic stress

reflecting the intraspecific competition within habitat, which could reliably drive the

transition from wingless to winged aphids across generations (*Deem et al., 2024*; *Sng*

and Ackerman, 2020). Since wingless aphids typically have higher fecundity than the

winged morph, switching to the dispersal winged morph from the reproductive morph

alleviates resource competition in offspring (*Deem et al., 2024*). In some strains of pea

aphids, even two maternal aphids housed in a petri dish can efficiently induce a high

proportion (>90%) of winged offspring than singly housed aphids (*Sutherland, 1969*;

Yuan et al., 2023a; *Yuan et al., 2023b*). Physical contact seems to be a trigger for adults

to produce winged offspring (*Sutherland, 1969*). Similarly, the aphid alarm pheromone

can also induce groups of aphids, rather than single individuals, to produce a higher

proportion of winged offspring, suggesting that the singly housed aphids that lack

physical contact cannot initiate wing morph change in offspring despite perception of

the alarm pheromone (*Kunert et al., 2005*).

Recent studies have identified a number of genes and gene networks involved in

the regulation of insect wing polyphenism. In planthoppers, the zinc finger

homeodomain transcription factor (Zfh1) and insulin/insulin-like growth factor

signaling (IIS) pathways coordinately modulate the fate of wing development in a

Forkhead transcription factor subgroup O (FoxO)-dependent manner (*Xu et al., 2015*;

Zhang et al., 2022). Dephosphorylated FoxO suppresses wing development by

inhibiting the transcription of cellular proliferation genes and wing-patterning genes, or

through repressing the transcriptional activity of Rotund, a zinc finger transcription

factor of wing developmental genes (*Chen et al., 2023*; *Lin et al., 2020*; *Xu et al., 2021*;

Yuan et al., 2023a; *Zhang et al., 2021*). However, these mechanisms are within one

generation prior to the critical switching time point at the nymphal stage. In contrast,

the wing morph of some aphids is determined transgenerationally. Ecdysteroid, miR-

9b, dopamine, and IIS signaling in maternal pea aphids are altered by experiencing high

population density, which increases the proportion of winged offspring (*Liu and*

Brisson, 2023; *Shang et al., 2020*; *Vellichirammal et al., 2017*; *Yuan et al., 2023a*). In

offspring, the TOR (the target of rapamycin) and Wnt signaling pathways respectively

control the occurrence of autophagic and apoptotic degradation of wing disc in 1^{st} instar

nymph of pea aphids, resulting in the formation of wingless morph (*Yuan et al., 2023b*;

Zhou et al., 2023). However, the molecules in the brain and embryo that are specifically

stimulated by maternal density and determining offspring wing morph before birth are

largely unknown.

Glutamate is a primary excitatory neurotransmitter. It not only functions in the

signal transduction cascade mediating the parent-offspring communication in the locust,

but is involved in the regulation of its interconvertible phase change, from the

solitarious to gregarious phase in response to high population density (*Yang et al., 2023*).

Maintaining an optimal glutamate concentration at the right time and in the right tissue

is important for proper nerve cell signaling, and embryo and tissue development.

Dysfunctional glutamatergic synaptic transmission has been linked to transgenerational

effects on anxiety, fear, pain sensitization, and embryonic development in animals

(*Jones et al., 2022*; *Liu et al., 2017*; *Tao et al., 2020*; *Van Winkle, 2022*). Glutamate

signaling involves a sophisticated network of metabolic pathways and transport

mechanisms that facilitate glutamate synthesis, release, and clearance, ensuring its

proper function as the primary excitatory neurotransmitter in the central nervous system

(*Featherstone, 2010*). This process begins with glutamate production in the presynaptic

neurons via the enzyme glutaminase, followed by glutamate sequestration into synaptic

vesicles by vesicular glutamate transporters (vGluTs). Upon synaptic activation,

glutamate is exocytosed into the synaptic cleft, where it interacts with ionotropic or

metabotropic glutamate receptors (iGluRs and mGluRs) on the postsynaptic neuron,

triggering a variety of signaling cascades (*Willard and Koochekpour, 2013*).

The glutamate signaling pathway could modify the phosphorylation level of FoxO.

Phosphorylated FoxO no longer binds to promoters and thus loses the suppression of

the network of wing development genes (*Howlett et al., 2008*; *Xu et al., 2015*; *Zhang et

al., 2018*). It therefore is plausible that the maternal glutamate pathway controls the

phosphorylation level the embryonic FoxO, and thereby regulating the developmental

trajectory of wing disc in offspring. In this study, maternal *vesicular glutamate

transporter 2* (*ApvGluT2*) and *metabotropic glutamate receptor 3* (*ApmGluR3*) of pea

aphids were identified by the tissue-specific RNA-seq as differentially expressed genes;

both were induced by physical contact or by high population density. Given the

importance of the glutamate signaling pathway in perceiving the population density and

transducing the signal to offspring, the effects of brain *ApvGluT2* and embryonic

ApmGluR3 were determined on (i) increases in the proportion of winged offspring, and

(ii) FoxO phosphorylation level and downstream wing developmental genes at stage 20

embryo stage, the crucial time point for determination of the wing morph.

Results

Physical contact and high maternal density increased the proportion of winged offspring. To determine the transgenerational effects of maternal density and physical contact on the wing morph in pea aphid offspring, maternal aphids in groups of 1, 2, and 8 were placed in a petri dish respectively, mimicking the solitary, physical contact, and maternal density effects (Figure 1A). Maternal aphids treated with physical contact and high density for 2- and 4-hr had higher proportions of winged offspring than solitary aphids (Figure 1B). Notably, 2- and 4-hrcontacts between two adults resulted in 54.1% and 96.5% winged offspring, similar to the effects of high-density treatment for the same time periods, 86.5% and 94.6% winged offspring, respectively (Figure 1B). Thus, physical contact could be utilized to obtain effective induction of winged offspring and even explain the effect achieved by high population density.

Since data collected from the two-adult contacting for 2-hr treatment exhibited a full range in percentage of winged offspring from 0% to 100%, the locomotor activity of maternal aphids was traced to determine the possible correlation with the proportion of winged offspring. Three locomotor parameters, including movement distance, velocity, and contact frequency of maternal aphids were monitored by the EthoVision system (Noldus) (Figure S1 A). All three maternal locomotor parameters exhibited strong positive correlations with the winged offspring proportion (P value < 0.0001) (Figure S1 B-D). Furthermore, since movement distance and velocity of maternal aphids were not significantly different between solitary and two-aphid contact for 4-hr

treatments, physical contact between maternal aphids could be the key driving factor

for higher proportion of winged offspring (Figure S1 E-F).

Physical contact and high density increased the expression of brain *ApvGluT2* and

embryonic *ApmGluR3* in maternal aphids. RNA-seq was used to determine tissue-

specific gene profiles of solitary maternal aphids and those subjected to 4-hr physical

contact treatment. A total of 563 differential expressed genes (DEGs) were identified in

maternal heads, while 339 genes were differentially expressed in embryos with the

criteria of the absolute value of Log_2 (fold change) > 0.58 and *P* value < 0.05 (Figure

1C). Of which, only 6 DEGs functioning in neurotransmitter signaling were identified,

including *ApvGluT2* (LOC100168565), *ApmGluR3* (LOC107882629), *Glutamate-*

gated chloride channel α (*ApGluCl-α*, LOC100162577), *Octopamine receptor β-1R*

(*ApOctβ1R*,   LOC100166522),   *Glycine   receptor   subunit   α1*   (*ApGLRA1*,

LOC100169196), and *Serotonin receptor 2A* (*ApHTR2A*, LOC100575043) (Figure 1D).

qPCR results further confirmed that only head *ApvGluT2* and embryonic *ApmGluR3*

had higher transcripts in maternal aphids experiencing physical contact than in solitary

aphids (Figure 1E). Genes associated with glutamate signaling components were

annotated, and their expression levels in maternal brain and embryo tissues were listed

(Figure S2). Furthermore, differential expression of *ApvGluT2* and *ApmGluR3* were not

observed in other tissues or in whole body except for gut *ApvGluT2* (Figure 1F-G). To

validate the involvement of *ApvGluT2* and *ApmGluR3* in perceiving high population

density, 15 adults vs. 1 adult were clipped into a plant seedling for 24-hr, and heads and

embryos were then dissected and sampled. RT-qPCR showed that 15-adult treatment

increased *ApvGluT2* transcripts by 64.9% in the head and *ApmGluR3* transcripts by

34.2% in the embryo relative to solitary aphids (Figure 1H). These results suggested

that maternal brain vGluT2 and embryonic mGluR3 were involved in the physical

contact-induced production of winged offspring.

Maternal *ApvGluT2* expressed in antennal lobes of aphid brain favored winged

offspring in a locomotor activity-independent manner. Previous research showed

that neural vGluT2 could modulate animal locomotor activity (*Roccaro-Waldmeyer et*

al., 2018). It was therefore speculated that physical contact-induced *ApvGluT2* in aphid

brain may increase the proportion of winged offspring by enhancing the locomotor

activity of maternal aphids or by cascading the glutamate pathway to regulate the wing

development across generations. To experimentally test these hypotheses,

pharmacological assays were performed on maternal aphids (Figure 2A), followed by

monitoring of the locomotor activities and proportion of winged offspring of aphids

subjected to 2-adults contact for 2-hr (Figure 2A). Application of glutamate agonist

glutamic acid increased winged offspring from 47.6% to 68.1% (Figure 2B).

Conversely, vGluT2 inhibitor Chicago Sky Blue (CSB) reduced the proportion of

winged offspring from 53.3% to 31.2% (Figure 2F). However, either had any effects on

locomotor activities (Figure 2C-E, 2G-I). Fluorescence *in situ* hybridization (FISH)

revealed that *ApvGluT2* was mainly expressed in the neural antennal lobes of the aphid

brain, and the fluorescent signal intensified as the maternal population density or

contact duration increased (Figure 2J-K). Amputating the antennae of maternal aphids

significantly reduced the proportion of winged offspring from 95.2% to 77.0% (Figure

2L). Together, aphids activated brain *ApvGluT2* in response to maternal density and physical contact, yet the resulting higher proportion of winged offspring had little to do with the locomotor activity.

Embryonic *ApmGluR3* relayed the key transgenerational signal involved with wing dimorphism. Because the embryonic period is a critical time point for determination of offspring wing morph, we hypothesized that *ApmGluR3* in the embryo was a key regulator downstream of *ApvGluT2* that mediated maternal-offspring communication. Application of mGluR antagonist LY341495 decreased the winged offspring by 33% (Figure 3A), whereas injection of mGluR agonist (2R, 4R)-4-aminopyrrolidine-2,4-dicarboxylate (APDC) increased the proportion of winged offspring by 40.4% (Figure 3B). Topical application of ds*ApmGluR3-RNA* knocked down the expression of *ApmGluR3* by 62.2% in the embryo, and the percentage of winged offspring dropped from 56.3% to 36.5% (Figure 3C-E). To establish a causal link between brain vGluT2 and embryo mGluR3, ds*ApvGluT2-RNA* and APDC were administered simultaneously (Figure 3F-G). Notably, reduction of winged offspring caused by the knockdown of *ApvGluT2* diminished upon application of APDC, the agonist of embryonic mGluR3 (Figure 3 F-G). FISH assays revealed abundant *ApmGluR3* in stage 20 embryos (Figure 3H). More importantly, the expression of *ApmGluR3* in the wing disc area of the embryo was up-regulated by density and contact treatments (Figure 3H-I). Presumably, ApmGluR3 functioned downstream of ApvGluT2 in the signaling pathway and mediated transgenerational wing dimorphism during embryonic development.

ApmGluR3 promoted phosphorylation of ApFoxO in embryos. Our previous study

indicated that the embryonic transcription factor, ApFoxO is a crucial suppressor of

wing development in pea aphids (*Yuan et al., 2023a*). In planthoppers, phosphorylation

of FoxO in wing bud cells caused its translocation from the nucleus to cytoplasm,

thereby losing its transcriptional repressor function in wing development (*Lin et al.,

2016*; *Xu et al., 2015*). We therefore examined whether ApmGluR3 affected

phosphorylation of *Ap*FoxO in embryos (Figure 4A). Initially, we confirmed that

ApFoxO indeed suppressed winged offspring: knockdown of *ApFoxO* in embryos of

maternal aphids by 28.6% (*P* value < 0.0001; Figure 4B) led to a significant increase

in the percentage of winged offspring, from 53.9% to 61.8% (*P* value = 0.0255; Figure

4C). Amino acid sequence alignment of ApFoxO with its homologs in fruit fly

(DmFoxO) and human (hFoxO4) identified three conserved putative phosphorylation

sites in ApFoxO, T15, S187, and S250 (Figure S3). Akt could further phosphorylate

these sites to sequestrate FoxO in the cytoplasm, thus preventing FoxO factors from

transactivating their target genes (*Calnan and Brunet, 2008*). Our results found that the

phosphorylation of ApFoxO was successfully detected by antibody against

phosphorylated T15, but failed to be detected with antibodies targeting phosphorylated-

S187 and S250 (Figure 4D-F, S4). Physical contact for 4-hr significantly increased

phosphorylated *Ap*FoxO in embryos (by 59.8%) compared to the solitary counterparts

(Figure 4D). Knockdown of *ApmGluR3* or *ApvGluT2* reduced the phosphorylation level

of ApFoxO in embryos by 21.3% and 54.3%, respectively (Figure 4E-F). In addition,

applying the specific Akt inhibitor MK-2206 remarkably reduced the percentage of

winged offspring, from 49.8% to 24.9% (Figure 4G). These results implied that the ApvGluT2-ApmGluR3 cascade increased the phosphorylation of ApFoxO in embryos, promoting the winged offspring.

ApFoxO bound to *ApHh* promotor to suppress its activation. Our previous study found that the transcripts of four wing development genes, including *Hedgehog* (*Hh*), *Vestigial* (*Vg*), *Decapentaplegic* (*Dpp*) and *Mob as tumor suppressor* (*Mats*), were significantly altered by knockdown of *ApFoxO* in the embryos of pea aphids (*Yuan et al., 2023a*). The stage 20 embryo appears to be a critical point in development for an aphid to determine irreversibly its winged or wingless morph (*Ishikawa and Miura, 2013*; *Ogawa and Miura, 2013*). Knockdown of *ApmGluR3* at stage 20 embryo decreased expression of *ApHh* and *ApMats* by 37.7% and 41.5% (Figure 5A). Interestingly, knockdown of *ApFoxO* caused an increase in *ApHh* transcripts, but a decrease in *ApMats* in stage 20 embryos (*Yuan et al., 2023a*). Accordingly, we ruled out *ApMats* and focused on *ApHh* as the suppressing target of ApFoxO. The application of ds*ApHh-RNA* reduced the expression of *ApHh* by 62.4% in stage 20 embryos, and decreased the proportion of winged offspring from 42.5% to 23.5% (Figure 5B-C). Knockdown of *ApFoxO* in stage 20 embryo increased the proportion of winged offspring by 40.4%. Percentage of winged offspring returned to the control level when both *ApFoxO* and *ApHh* were silenced (Figure 5D-E). These findings demonstrated that ApFoxO repressed wing development via inhibiting *ApHh* expression at the 20[th] embryo stage.

Electrophoretic mobility shift assays (EMSA) were then performed to detect

binding of ApFoxO to the promotor of *ApHh* via the FoxO recognition element (*FRE*),

TTGTTTAC. Typically, FoxO interacts with the *FRE*-element through its highly

conserved DNA-binding domain (DBD) consisting of ~100 amino acid residues to

regulate transcription of target genes (*Furuyama et al., 2000*; *Xuan and Zhang, 2005*).

The sequence alignment showed that DBD of *ApFoxO* shared high sequence similarity

to its homologs in brown planthopper *Nilaparvata lugens*, fruit fly *Drosophila*

melanogaster and human *Homo sapiens* (Figure 5F). Two FREs, FRE1 and FRE2, were

identified in the 3-kb segment upstream of the transcription start site of *ApHh* (Figure

5G-H). Recombinant maltose-binding protein (MBP)-DBD$_{ApFoxO}$ was expressed in

Escherichia coli and purified for EMSA (Figure S5). Biotin-labeled probes of FRE2

result in mobility shift in a concentration-dependent manner, while no binding to biotin-

labeled FRE1 was detected (Figure 5G-H). Notably, the intensity of shift bands was

greatly reduced by adding the unlabeled FRE2 probe. No binding occurred when the

cis-element was mutated (Figure 5H), supporting specific interaction between *FRE2*

and ApFoxO.

We further compared the temporal expression patterns of *ApHh* between offspring

collected from solitarious and the 4-hr physical contact treatment across the

developmental stages from stage 20 embryos to the 4th instar nymphs. Little treatment

difference was observed in *ApHh* transcripts in postnatal development stages,

suggesting that the stage 20 embryo was a critical time point for decision-making

regarding the developmental trajectory of the wing disc (Figure 5I). Together, these

findings indicated that embryonic ApFoxO and ApHh were the executors to determine

winged dimorphism in aphids.

Discussion

The phenomenon where a single genotype is expressed in a spectrum of phenotypes,

both within a single generation and across successive ones, is central to comprehension

of how populations adapt to their environments. This adaptability is particularly

complicated in the context of TPP due to its obscure nature of signal transduction

mechanisms   facilitating   parent-offspring   communication.   Although   some

understanding has been attained about within-generation wing plasticity in insects,

much less is known regarding the transgenerational regulatory mechanism. Our

research has demonstrated that maternal brain *ApvGluT2*, mainly expressed in antennal

lobes, and embryonic *ApmGluR3* were specifically up-regulated when aphids

experienced physical contact or high population density. This was followed by an

elevated phosphorylation of *Ap*FoxO in stage 20 embryos, which caused the loss of its

suppression of *ApHh*, as well as a high proportion of winged offspring (Figure 6).

Conversely, lower expression of *ApvGluT2* and *ApmGluR3* in solitary aphids sustained

a higher level of unphosphorylated ApFoxO, which effectively suppressed *ApHh*

expression and favored the production of wingless offspring. These findings showcased

how pea aphids deployed the glutamate signaling pathway in regulating

transgenerational wing dimorphism in response to environmental cues, and highlighted

the importance of Hh as secreted protein governing wing development trajectory in

stage 20 embryos.

High population density is a reliable environmental cue for intense competition or deteriorating habitat conditions for individuals, resulting in a high proportion of winged offspring. The tactile, olfactory, and visual signals are essential cues for aphids to perceive population density and trigger appropriate phenotypic changes (*Topaz et al., 2012*). For instance, physical contact could induce gregarious locusts from a solitary population by releasing aggregation pheromone 4-vinylanisole (*Guo et al., 2020*; *Yang et al., 2023*). In aphids, the increased frequency of physical contact due to increasing population density could also explain wing induction (*Sutherland, 1969*). A "pseudo-crowding" hypothesis has been proposed to explain the predator-induced wing induction in aphids. The perception of predators or the alarm pheromone increases the locomotion activity of aphids, causing higher frequency of physical contact among individuals, similar to what happens when aphids are crowded. Although single individuals could behaviorally react to the alarm pheromone, lack of physical contact renders them unable to trigger wing production (*Kunert et al., 2005*). Consistently, aphids with ablated antennae fail to produce more winged offspring in the presence of predators (*Kunert et al., 2005*). Our previous screening of natural pea aphids gave rise to a strain that was highly sensitive to physical contact in triggering winged offspring, which provided an excellent biotype for dissecting the underlying molecular mechanism (*Yuan et al., 2023b*). Our results found a strong correlation between the locomotor activity of maternal aphids and the proportion of winged offspring in the treatment of two-adult contacting for 2-hr. Presumably, elevated mobility of maternal aphids led to higher frequency of physical contact, in agreement with the pseudo-

crowding hypothesis.

Low titers of serotonin, dopamine and octopamine are found in densely populated

pea aphids (*Vellichirammal et al., 2016*). Of these biogenic amines, only dopamine was

reported to be a suppressor for winged morph production (*Liu and Brisson, 2023*). In

this study, physical contact stimulated higher transcription activity of *ApvGluT2* mainly

located in the antennal lobes of the maternal brain. Blocking the vGluT2 activity

reduced the proportion of winged offspring but did not alter maternal locomotor activity.

It has been shown that vGluTs are essential in sensing mechanical pain in

somatosensory system, dorsal horn and brain (*Seal, 2016*). As evidenced by transiently

expressing *vGluT3* in neurons of mammal deep dorsal horn, it could directly receive

Aβ primary sensory input caused by mechanical stimuli, and relay polysynaptic

transmission onto lamina I pain processing neurons (*Peirs et al., 2015*; *Seal et al., 2009*).

Our phylogenetic analysis and sequence alignment indicated that ApvGluT2 belonged

to animal Clade III in the solute carrier (SLC) 17 subfamily, sharing 21.33%, 22.33%,

and 20.74% amino acid identity with human vGluT1, vGluT2, and vGluT3, respectively,

along with 12 conserved transmembrane regions (Figure S6 A-B). Notably, ApvGluT2

exhibited a significant structural similarity to human vGluT2, as evidenced by minimal

root mean square deviation (RMSD) value of 4.131 Å, suggesting ApvGluT2

maintained relatively conserved function in aphids (Figure S6 C). Most likely, a specific

group of *ApvGluT2*-expressing neurons in antennal lobes of aphids received physical

contact-induced sensory input, then transmitted the synaptic signals to downstream

nerves to modulate the offspring wing morph at their embryonic period.

Considering that the antennal lobes are located at a considerable distance from the

embryos, an important issue is how the brain ApvGluT2 controls the wing development

trajectory in embryos. Our results revealed that the embryonic *ApmGluR3* was up-

regulated by maternal high density and physical contact experiences. Although

ApmGluR3 was expressed throughout the embryo, it could be up-regulated by density

and contact treatments in the area of wing discs. Injection of mGluR agonist rescued

the negative effect of ds*ApvGluT2* on the proportion of winged offspring, suggesting

ApvGluT2 and *ApmGluR3* coordinately regulated the production of wing morph in

offspring triggered by maternal density. Early studies have shown that the vGluT-

mGluR signal axis orchestrates the function of specific neuronal networks involved in

motor coordination, emotions, and cognition (*Ibrahim et al., 2020*). Furthermore, the

glutamatergic neurons in the ventral nerve cord and thoracic abdominal ganglion could

project to the reproductive tract to regulate reproduction (*Castellanos et al., 2013*;

Deshpande et al., 2022; *Rodríguez-Valentín et al., 2006*). Therefore, it is likely that

glutamatergic neurons in antennal lobes of aphids recruited vGluT2-mediated signaling

to activate the glutamatergic neurons in the ventral nerve cord and thoracic abdominal

ganglion, innervating the mGluR3-expressed cells in the wing disc of embryos to

control their wing development.

Although FoxO acts as a major suppressor of the winged morph in both within-

and trans-generational wing polyphenism, the regulatory routes may vary in

developmental stages by targeting different downstream wing development genes.

Some such as *Vg* and *Ubx* that control the wing patterning in wing buds could be

suppressed by FoxO within a generation (*Liu et al., 2020*; *Zhang et al., 2021*). As for

the transgenerational regulation, we showed that physical contact-induced increases of

brain *ApvGluT2* and embryonic *ApmGluR3* enhanced the phosphorylation level of

ApFoxO, as well as the proportion of winged offspring. In contrast, knockdown of

ApvGluT2 and *ApmGluR3*, and the application of Akt antagonist could reduce the

phosphorylation level of ApFoxO, consistent with a previous study, where mGluR

activates PI3K in *Drosophila* larval motor neuron to alter phosphorylation of FoxO

(*Howlett et al., 2008*).

During the differentiation of Drosophila imaginal disc, *Hh* is transcribed in the

posterior compartment of the wing disc and acts as a short-range morphogen that is

secreted anteriorly, establishing a concentration gradient that provides positional

information to wing disc cells and determines the developmental trajectory of the wing

(*Hartl and Scott, 2014*; *Neto-Silva et al., 2009*). *Hh* controls the expression of the long-

range morphogen *Dpp* to regulate the pattern and growth of the wing disc through its

essential downstream component *Cubitus interruptus* (*Bejarano et al., 2007*). In this

study, the embryonic expression of *ApDpp* was not significant for ds*GFP* vs.

ds*ApmGluR3* group and solitary vs. physical contact group, leaving the possibility that

the clean sampling of wing discs of stage 20 embryos instead of the whole body could

reveal the wing gene regulatory network (GRN) in pea aphids. Moreover, knockdown

of *ApmGluR3* did not affect *vestigial*, which was wing specific and necessary for wing

development in insects, suggesting *ApmGluR3* indirectly regulated the wing GRN in

embryos, and the connection between *ApmGluR3* and the wing GRN could be mediated

by other downstream developmental or physiological processes that affected wing

morph development. Furthermore, our data demonstrated that FoxO could directly bind

to the promotor of *ApHh* to suppress its expression in stage 20 embryos. Beyond the

role in wing development, Hh functions as a secreted signaling protein in many other

organs, regulating a range of processes including embryonic development, tissue

homeostasis, organ regeneration, axon guidance, and cellular metabolism (*Briscoe and*

Thérond, 2013). Since the wingless and winged aphids exhibit a range of differences in

morphological, behavioral and life history (*Brisson, 2010*), the different expression of

Hh in stage 20 embryos could be critical to aphid wing-morph differentiation in

response to high population density. Together, we have provided a novel mechanistic

insight into the regulation of TPP and demonstrated, for the first time that the glutamate

signal transduction cascade mediates the transgenerational wing dimorphism in aphids.

Materials and methods

Aphid rearing. The pea aphid *A. pisum* (strain: Ningxia Red) was originally collected

from *Medicago sativa* in Ningxia Province and had been reared in the laboratory for

over 8 years. Nymphs from the same parthenogenetic pea aphid female were reared on

Vicia faba at 18-20°C, with 60% relative humidity and a photoperiod of a 16 hr: 8 hr

(light/dark) cycle. To eliminate the transgenerational effects on offspring morphs,

females were maintained at low density (three per plant) on *V. faba* seedlings for more

than three generations (*Sutherland, 1969*). This strain was used to test the effects of

two-adult physical contact or crowding on the proportion of winged offspring.

Induction of winged offspring. To effectively induce a high proportion of winged offspring, the maternal density and contacting duration were evaluated in petri dishes (35 mm in diameter) as described by 14. Three groups of aphids consisting of 1, 2, or 8 wingless adults, respectively, were placed in a petri dish for 0.5-, 2-, and 4-hr. After treatment, each adult was transferred to a freshly detached *V. faba* leaf kept in petri dishes with 1% agar to reproduce for 24-hr. After newborns reached the fourth instar, the proportion of winged offspring from each adult was calculated.

For density treatment, two groups of 1 and 15 maternal aphids were placed on fava seedlings for 24-hr. This was followed by transferring each adult to a fresh *V. faba* leaf as described above.

Aphid locomotor activity tracking in petri dishes. One or two maternal aphids were placed in a petri dish (diameter 35 mm), and locomotor activities of each aphid within 2- or 4-hr were recorded (25 frames/s) with a video camera. Six or 12 aphids in each treatment were simultaneously recorded per replicate. The valid video data for each treatment were collected and analyzed at least 40 replicates. Videos were analyzed using the EthoVision XT software (v.11.5, Noldus Information Technology) to measure the total distance (moving distance, unit: cm) and the average velocity of movement (move velocity, unit: cm/s), as well as the contact frequency. Physical contact between two maternal aphids was considered to have occurred when their proximity distance was less than 0.5 cm.

RNA-Seq. Maternal aphids housed individually or two per petri dish for 4-hr were used for differential gene expression analyses. The heads and embryos were dissected from

10 and 5 adult aphids per replicate, respectively. Four of such biological replicates were

collected RNA-seq. DEGs were identified with DESeq2 R package (1.20.0) using the

expected number of Fragments Per Kilobase of transcript sequence per Million base

pairs sequenced. The sequencing data were deposited in the Short Read Archive (NCBI)

under accession number PRJNA1017088.

RT-qPCR. Total RNA was extracted using TRIzol reagent (Invitrogen, CA, USA), and

reverse transcribed using the PrimeScriptTM RT reagent Kit with gDNA Eraser (Perfect

Real Time) kit (Takara, Dalian, China). The RT-qPCR reactions were carried out on the

PikoReal 96 Real-Time PCR System (Thermo) using the SuperReal PreMix Color

(SYBR Green) kit (Tiangen, Beijing, China). Three technical replicates were applied

for each biological replicate. The housekeeping gene ribosomal protein L27 (RPL27)

was used as the internal qPCR standard.Gene expression data were analyzed by relative

quantification with the $2^{-\triangle\triangle CT}$ method (*Livak and Schmittgen, 2001*). Specific primers

for each gene were designed using Primer Premier 6 and tested in preliminary

experiments. The oligonucleotides are listed in Table S1.

Pharmacological experiments. Glutamate signaling agonist glutamic acid

(Topscience, Shanghai) and mGluR agonist (2R, 4R)-ADPC (Enzo Life Sciences, New

York) were dissolved in water at concentrations of 33.98 mM and 100 nM, respectively.

vGluT2 antagonist Chicago Sky Blue 6B (CSB) (Topscience, Shanghai), mGluR

antagonist LY341495 (MedChemExpress, New Jersey), and Akt inhibitor MK-2206

(Selleckchem, Houston) were dissolved in DMSO at concentrations of 10 mM, 16.98

mM, and 5 mM, respectively. A volume of 100 nL of each chemical, except for MK2206,

which was 23.6 nL, was delivered into hemolymph from dorsal abdomens using the

Nanoject II microinjection system (Drummond Scientific Company, Broomall, PA,

USA). An equivalent amount of solvent was used as a solvent control.

Fluorescence in situ hybridization. FISH was performed with the technique modified

by Kliot et al. (*2014*). After washing in TBST (TBS with 0.2% Triton-X) for 10 min,

sample section slices were rinsed three times in the hybridization buffer (20 mM Tris-

HCl, pH 8.0, 0.9 M NaCl, 0.01% sodium dodecyl sulfate, and 30% formamide), and

hybridized overnight in the hybridization buffer containing 10 pmol of the fluorescent

RNA probe (conjugated with FAM or Cy5). After 3x rinses in TBST, samples were

mounted in Fluoroshield Mounting Medium with DAPI (Abcam). Sections were

imaged using a Leica M205C confocal microscope (Zeiss, Germany). The fluorescence

intensities were quantified using Leica Application Suite (LAS) software.

RNA interference. Double-stranded RNA (dsRNAs) i.e. ds*ApvGluT2*, ds*ApmGluR3*,

ds*ApFoxO*, ds*ApHh*, and the ds*GFP* control were synthesized by T7 RiboMAX Express

RNAi System (Promega, P1700) according to the manufacturer's protocol. Each (2

μg/μl) was mixed with the nanocarrier in a water solution at a volume ratio of 1:1. A

volume of 1 μL of the solution containing 1 μg nanocarrier/dsRNA complex was

topically applied to the dorsal side of the female abdomen. Embryos from aphids or

heads from ten aphids in each sample were collected 24 h after topical application of

dsRNA for testing silencing efficiency using qPCR. Each treatment contained at least

three biological replicates. The primers used for dsRNA synthesis are listed in Table S2.

Antibody preparation and western blotting analysis. Three polyclonal rabbit

antibodies against phosphorylated T15, S187 and S250 of ApFoxO were prepared by ABclonal using synthetic peptides RARSNT(p)WPLPR (10 to 20 aa), RRRAVS(p)METPK (182 to 192 aa) and RARASS(p)NASS (245 to 254 aa) as the immunogens. Western blotting was performed to detect phosphorylation levels of FoxO in embryos of adult females. The antigen–antibody complexes were visualized using a secondary goat anti-rabbit IgG (H+L)-conjugated horseradish peroxidase (HRP) antibody (LABLEAD).

Electrophoretic mobility shift assays (EMSA). The FoxO DNA-binding domain (DBD) (89 to 178 aa) was cloned into the pMAL-c2x expression vector, which contained an N-terminal maltose-binding protein (MBP), and subsequently expressed in *E. coli* strain Rosetta (DE3). Proteins were purified from cell lysates using the pMAL protein fusion and purification system (New England Biolabs). The MBP protein was also independently expressed in pMAL-c2x as control. EMSA was conducted as described previously (*Hellman and Fried, 2007*). Probes containing the DBD binding sites were labeled with biotin at the 5' end. Unlabeled probe was used as the competitor. The probes were incubated with purified MBP/DBD recombinant protein at room temperature for 30 min, run on a non-denaturing 0.5×TBE 6% polyacrylamide gel for 1h (60 V at 4°C), and transferred onto Biodyne® B nylon membrane (Pall Corporation). Signals were visualized using the ChemiDoc XRS kit (Bio-Rad Laboratories, UAS).

Phylogenetic construction of SLC17 proteins. Protein sequences, 333 from 10 different species, were obtained from UniProt (https://www.uniprot.org/). Sequences were aligned with MAFFT v7.310 (*Katoh et al., 2002*) and automated trimmed with

trimAl v1.4 (*Capella-Gutiérrez et al., 2009*). Phylogenetic analysis was performed

using IQ-TREE v2.0.3 with LG+R9 model and ultrafast bootstrap approximation with

1000 replicates (*Minh et al., 2020*). Phylogenetic tree was visualized in Interactive Tree

Of Life (iTOL) (*Letunic and Bork, 2021*).

Protein Sequence and Structure analysis. The multiple sequence alignment was

performed using Multiple Protein Sequence Alignment (MUSCLE). High confidence

protein structure of *Ap*vGluT2 was predicted by Alphafold v2.3.0 (*Jumper et al., 2021*).

The pairwise structural alignments were performed using PyMOL 2.6.0a0

(*www.pymol.org*).

Statistical Analyses. The GraphPad software was used for statistical analyses. All data

were checked for normality by the Wilk–Shapiro test. Two-tailed unpaired t-test was

used to separate the means of normally distributed data, while Mann–Whitney test was

used to analyze nonparametric data. All western blotting and confocal assays were

repeated independently at least three times with similar results. Pearson's correlations

were calculated to determine the relationship between proportion of winged offspring

and maternal locomotor activities.

Data Availability

All data used in the study are included, and some are presented as supplementary information. All protocols have been described in Materials and Methods or in the references therein. No custom codes or mathematical algorithms were used in this manuscript.

Author contribution

Yiyang Yuan project designing, performing experiments, data analysis and manuscript writing; Yanyan Wang, Wanwan Ye, Liqiang Xie, Fang Dong performing experiments; Keyan Zhu-Salzman manuscript writing and revising, and Erliang Yuan, Huijuan Guo, Shifan Wang manuscript revising; Feng Ge, Yucheng Sun project designing and coordination, manuscript writing.

Declare of interests

We declare no competing interest.

Acknowledgements

We thank Prof. Jinfeng Chen for his support in phylogenetic analysis. This project was supported by the National Key R&D Program of China (no. 2023YFD1400800), the Taishan Scholars Program, the Introducing Top Talent Program of Shandong (2023YSYY-006), the Agricultural Science and Technology Innovation Project of the Shandong Academy of Agricultural Sciences (no. CXGC2023F04), the National Natural Science Foundation of China (nos. 31970453, 32250002 and 32372636), and the State Key Laboratory of Integrated Management of Pest Insects and Rodents (no. IPM2306).

References

Bejarano F, Perez L, Apidianakis Y, Delidakis C, Milan M. 2007. Hedgehog restricts its

expression domain in the Drosophila wing. EMBO reports, **8**: 778-783.

Bell AM, Hellmann JK. 2019. An integrative framework for understanding the

mechanisms and multigenerational consequences of transgenerational plasticity.

Annual Review of Ecology, Evolution, and Systematics **50**: 97–118.

Braendle C, Friebe I, Caillaud MC, Stern DL. 2005. Genetic variation for an aphid wing

polyphenism is genetically linked to a naturally occurring wing polymorphism.

Proceedings. Biological sciences **272**: 657–664.

Brisson JA. 2010. Aphid wing dimorphisms: linking environmental and genetic control

of trait variation. Philosophical Transactions of The Royal Society of London Series B-

Biologic Journal **365**: 605–616.

Briscoe J, Thérond PP. 2013. The mechanisms of Hedgehog signalling and its roles in

development and disease. Nat. Rev. Mol. Cell Biol. **14**: 416–429.

Calnan DR, Brunet A. 2008. The FoxO code. Oncogene **27**: 2276–2288.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics **25**: 1972–

1973.

Castellanos MC, Tang JC, Allan DW. 2013. Female-biased dimorphism underlies a

female-specific role for post-embryonic Ilp7 neurons in Drosophila fertility.

Development **140**: 3915–3926.

Chen SJ, Zhang JL, Ma WJ, Wu HJ, Li Y, Shen XX, Xu HJ. 2023. FoxO and rotund

form a binding complex governing wing polyphenism in planthoppers. iScience **26**: 107182.

Deem KD, Gregory LE, Liu X, Ziabari OS, Brisson JA. 2024. Evolution and molecular mechanisms of wing plasticity in aphids. Current Opinion in Insect Science **61**: 101142.

Deshpande SA, Rohrbach EW, Asuncion JD, Harrigan J, Eamani A, Schlingmann EH, Suto DJ, Lee PT, Schweizer FE, Bellen HJ, Krantz DE. 2022. Regulation of Drosophila oviduct muscle contractility by octopamine. iScience **25**: 104697.

Donelan SC, Hellmann JK, Bell AM, Luttbeg B, Orrock JL, Sheriff MJ, Sih A. 2020. Transgenerational Plasticity in Human-Altered Environments. Trends in Ecology & Evolution **35**: 115–124.

Featherstone DE. 2010. Intercellular glutamate signaling in the nervous system and beyond. ACS chemical neuroscience **1**: 4–12.

Furuyama T, Nakazawa T, Nakano I, Mori N. 2000. Identification of the differential distribution patterns of mRNAs and consensus binding sequences for mouse DAF-16 homologues. Biochemical Journal **349**: 629–634.

Guo X, Yu Q, Chen D, Wei J, Yang P, Yu J, Wang X, Kang L. 2020. 4-Vinylanisole is an aggregation pheromone in locusts. Nature **584**: 584–588.

Guillaume AS, Monro K, Marshall DJ. 2016. Transgenerational plasticity and environmental stress: do paternal effects act as a conduit or a buffer? Functional Ecology **30**: 1175–1184.

Hartl TA, Scott MP. 2014. Wing tips: The wing disc as a platform for studying Hedgehog signaling. Methods **68**: 199–206.

Hellman LM, Fried MG. 2007. Electrophoretic mobility shift assay (EMSA) for

detecting protein-nucleic acid interactions. Nature protocols **2**: 1849–1861.

Howlett E, Lin CC, Lavery W, Stern M. 2008. A PI3-kinase-mediated negative

feedback regulates neuronal excitability. PLoS genetics **4**: e1000277.

Ibrahim KS, Abd-Elrahman KS, El Mestikawy S, Ferguson SS. 2020. Targeting

Vesicular Glutamate Transporter Machinery: Implications on Metabotropic Glutamate

Receptor 5 Signaling and Behavior. Molecular Pharmacology **98**: 314–327.

Ishikawa A, Miura T. 2013. Transduction of high-density signals across generations in

aphid wing polyphenism. Physiological Entomology **38**: 150–156.

Jones SK, McCarthy DM, Vied C, Stanwood GD, Schatschneider C, Bhide PG. 2022.

Transgenerational transmission of aspartame-induced anxiety and changes in

glutamate-GABA signaling and gene expression in the amygdala. Proceedings of the

National Academy of Sciences **119**: e2213120119.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool

K, Bates R, Žídek A, Potapenko A, Bridgland A. 2021. Highly accurate protein structure

prediction with AlphaFold. Nature **596**: 583–589.

Katoh K, Misawa K, Kuma KI, Miyata T. 2002. MAFFT: a novel method for rapid

multiple sequence alignment based on fast Fourier transform. Nucleic Acids Research

30: 3059–3066.

Kliot A, Kontsedalov S, Lebedev G, Brumin M, Cathrin PB, Marubayashi JM, Skaljac

M, Belausov E, Czosnek H, Ghanim M. 2014. Fluorescence in situ hybridizations

(FISH) for the localization of viruses and endosymbiotic bacteria in plant and insect

tissues. Journal of Visualized Experiments **84**: e51030.

Kunert G, Otto S, Röse US, Gershenzon J, Weisser WW. 2005. Alarm pheromone mediates production of winged dispersal morphs in aphids. Ecology letters **8**: 596–603.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. Nucleic Acids Research **49**: W293–W296. 62.

Lin X, Yao Y, Wang B, Emlen DJ, Lavine LC. 2016. Ecological trade-offs between migration and reproduction are mediated by the nutrition-sensitive insulin-signaling pathway. International journal of biological sciences **12**: 607–616.

Lin X, Gao H, Xu Y, Zhang Y, Li Y, Lavine MD, Lavine LC. 2020. Cell cycle progression determines wing morph in the polyphenic insect *Nilaparvata lugens*. iScience **23**: 101040.

Liu F, Li X, Zhao M, Guo M, Han K, Dong X, Zhao J, Cai W, Zhang Q, Hua H. 2020. Ultrabithorax is a key regulator for the dimorphism of wings, a main cause for the outbreak of planthoppers in rice. National Science Review **7**: 1181–1189.

Liu Y, Ge R, Zhao X, Guo R, Huang L, Zhao S, Guan S, Lu W, Cui S, Wang S, Wang JH. 2017. Activity strengths of cortical glutamatergic and GABAergic neurons are correlated with transgenerational inheritance of learning ability. Oncotarget **8**: 112401–112416.

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\triangle\triangle CT}$ Method. Methods **25**: 402–408.

Liu X, Brisson JA. 2023. Dopamine mediates the pea aphid wing plasticity. Biology Letters **19**: 20230024.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, Von Haeseler A,

Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Molecular biology and evolution **37**: 1530–1534.

Neto-Silva RM, Wells BS, Johnston LA. 2009. Mechanisms of growth and homeostasis

in the Drosophila wing. Annual Review of Cell and Developmental **25**: 197–220.

Ogawa K, Miura T. 2013. Two developmental switch points for the wing

polymorphisms in the pea aphid *Acyrthosiphon pisum*. Evodevo **4**: 30.

Peirs C, Williams SP, Zhao X, Walsh CE, Gedeon JY, Cagle NE, Goldring AC, Hioki

H, Liu Z, Marell PS, Seal RP. 2015. Dorsal Horn Circuits for Persistent Mechanical

Pain. Neuron **87**: 797–812.

Roccaro-Waldmeyer DM, Girard F, Milani D, Vannoni E, Prétôt L, Wolfer DP, Celio

MR. 2018. Eliminating the VGlut2-dependent glutamatergic transmission of

parvalbumin-expressing neurons leads to deficits in locomotion and vocalization,

decreased pain sensitivity, and increased dominance. Frontiers in behavioral

neuroscience **12**: 146.

Rodríguez‐Valentín R, López‐González I, Jorquera R, Labarca P, Zurita M,

Reynaud E. 2006. Oviduct contraction in Drosophila is modulated by a neural network

that is both, octopaminergic and glutamatergic. Journal of cellular physiology **209**:

183–198.

Roff DA. 1990. The evolution of flightlessness in insects. Ecological Monographs **60**:

389–421.

Roff DA. 1994. The evolution of flightlessness: Is history important? Ecology and

Evolution **8**: 639–657.

Seal RP, Wang X, Guan Y, Raja SN, Woodbury CJ, Basbaum AI, Edwards RH. 2009.

Injury-induced mechanical hypersensitivity requires C-low threshold

mechanoreceptors. Nature **462**: 651–655.

Seal RP. 2016. Do the distinct synaptic properties of VGLUTs shape pain?

Neurochemistry International **98**: 82–88.

Shang F, Niu J, Ding BY, Zhang W, Wei DD, Wei D, Jiang HB, Wang JJ. 2020. The

miR-9b microRNA mediates dimorphism and development of wing in aphids.

Proceedings of the National Academy of Sciences **117**: 8404–8409.

Sng O, Ackerman JM. 2020. Too many people, women, men? The psychological effects

of population density and sex ratio. Current Opinion in Psychology **32**: 38–42.

Sutherland ORW. 1969. The role of crowding in the production of winged forms by two

strains of the pea aphid, *Acyrthosiphon pisum*. Journal of Insect Physiology **15**: 1385–

1410.

Tao W, Chen C, Wang Y, Zhou W, Jin Y, Mao Y, Wang H, Wang L, Xie W, Zhang X, Li

J. 2020. MeCP2 mediates transgenerational transmission of chronic pain. Progress in

neurobiology **189**: 101790.

Topaz CM, D'Orsogna MR, Edelstein-Keshet L, Bernoff AJ. 2012. Locust dynamics:

behavioral phase change and swarming. PLoS Computational Biology **8**: e1002642.

Van Winkle LJ. 2022. Perspective: Might Maternal Dietary Monosodium Glutamate

(MSG) Consumption Impact Pre- and Peri-Implantation Embryos and Their

Subsequent Development? International Journal of Environmental Research and Public

Health **19**: 13611.

Vellichirammal NN, Madayiputhiya N, Brisson JA. 2016. The genomewide

transcriptional response underlying the pea aphid wing polyphenism. Molecular

ecology **25**: 4146–4160.

Vellichirammal NN, Gupta P, Hall TA, Brisson JA. 2017. Ecdysone signaling underlies

the pea aphid transgenerational wing polyphenism. Proceedings of the National

Academy of Sciences **114**: 1419–1423.

Wagner DL, Liebherr JK. 1992. Flightlessness in insects. Trends in Ecology and

Evolution **7**: 216–220.

Willard SS, Koochekpour S. 2013. Glutamate, glutamate receptors, and downstream

signaling pathways. International journal of biological sciences **9**: 948–959.

Xu N, Wei SF, Xu HJ. 2021. Transcriptome Analysis of the Regulatory Mechanism of

FoxO on Wing Dimorphism in the Brown Planthopper, *Nilaparvata lugens* (Hemiptera:

Delphacidae). Insects **12**: 413.

Xu HJ, Xue J, Lu B, Zhang XC, Zhuo JC, He SF, Ma XF, Jiang YQ, Fan HW, Xu JY,

Ye YX. 2015. Two insulin receptors determine alternative wing morphs in planthoppers.

Nature **519**: 464–467.

Xuan Z, Zhang MQ. 2005. From worm to human: bioinformatics approaches to identify

FOXO target genes. Mechanisms of ageing and development **126**: 209–215.

Yang M, Du B, Xu L, Wang H, Wang Y, Lin K, He G, Kang L. 2023. Glutamate-GABA

imbalance mediated by miR-8-5p and its STTM regulates phase-related behavior of

locusts. Proceedings of the National Academy of Sciences **120**: e2215660120.

Yuan Y, Wang Y, Ye W, Yuan E, Di J, Chen X, Xing Y, Sun Y, Ge F. 2023a. Functional

evaluation of the insulin/insulin-like growth factor signaling pathway in determination

of wing polyphenism in pea aphid. Insect Science **30**: 816–828.

Yuan E, Guo H, Chen W, Du B, Mi Y, Qi Z, Yuan Y, Zhu-Salzman K, Ge F, Sun Y.

2023b. A novel gene REPTOR2 activates the autophagic degradation of wing disc in

pea aphid. eLife **12**: e83023.

Zera AJ, Denno RF. 1997. Physiology and ecology of dispersal polymorphism in insects.

Annual Review of Entomology **42**: 207–230.

Zhang CX, Brisson JA, Xu HJ. 2019. Molecular Mechanisms of Wing Polymorphism

in Insects. Annual Review of Entomology **64**: 297–314.

Zhang B, Gong J, Zhang W, Xiao R, Liu J, Xu XS. 2018. Brain-gut communications

via distinct neuroendocrine signals bidirectionally regulate longevity in *C. elegans*.

Genes & development **32**: 258–270.

Zhang JL, Fu SJ, Chen SJ, Chen HH, Liu YL, Liu XY, Xu HJ. 2021. Vestigial mediates

the effect of insulin signaling pathway on wing-morph switching in planthoppers. PLoS

Genetics **17**: e1009312.

Zhang JL, Chen SJ, Liu XY, Moczek AP, Xu HJ. 2022. The transcription factor Zfh1

acts as a wing-morph switch in planthoppers. Nature Communications **13**: 5670.

Zhou P, Zong X, Yan S, Zhang J, Wang D, Shen J. 2023. The Wnt pathway regulates

wing morph determination in *Acyrthosiphon pisum*. Insect Biochemistry and Molecular

Biology **161**: 104003.

Figure 1 Maternal aphids produced high proportions of winged offspring after experiencing physical contact and high density, with increased expression of head *ApvGluT2* and embryo *ApmGluR3*. (A) The schematic procedure of transgenerational induction of winged offspring. (B) Effects of maternal density and duration on the proportion of winged offspring (n=32). Two-adult contact for 4 hr was sufficient to induce over 90% of winged offspring. Kruskal-Wallis test was performed, and Dunn's test was used to differentiate between the means; different letters indicate significant differences ($P < 0.05$). (C) The volcano plots showing differentially expressed genes (DEGs) in the head and embryo induced by two-adult contacting for 4-hr (n=4). DEGs were selected if the absolute value of Log_2 (Fold change) > 0.58 and $P < 0.05$. (D) The

heat map showing the expression patterns of six DEGs functioning in neurotransmitter

signaling in RNAseq analyses (n=4). (E) qPCR confirmation of increased expression

of *ApvGluT2* and *ApmGluR3* in head and embryo, respectively caused by maternal

physical contact (n=3). (F) The tissue-specific expression of *ApvGluT2* and (G)

ApmGluR3 in two-adult contacting for 4-hr treatment vs. solitary treatment (n=3). (H)

Transcripts of head *ApvGluT2* and embryo *ApmGluR3* were higher in group aphids of

15 adults than those of singly kept aphids on plants, as determined by qPCR (n=3).

Student *t* test: $*P < 0.05$, $**P < 0.01$.

Figure 2 The effect of *ApvGluT2* on the proportion of winged offspring and its

location in aphid brain. (A) Schematic diagram showing the procedure of winged

offspring inspection and behavior monitoring in pharmacological experiments in pea

aphids. (B) Increased percentage of winged offspring by glutamic acid injection, an

agonist of vGluT (n>47). (C) The effect of glutamic acid injection on the movement

distance (n=48), (D) the movement velocity (n = 48), and (E) the contact frequency of

maternal aphids (n=24). The Kolmogorov-Smirnov test was used to compare means.

(F) Decreased percentage of winged offspring by CSB injection, an antagonist of vGluT

(n>42). (G) The effect of CSB injection on the movement distance (n=44), (H) the

movement velocity (n=44), and (I) the contact frequency of maternal aphids (n=22). (J)

mRNA-FISH. *ApvGluT2* was mainly expressed in the neural antennal lobes

(highlighted by white squares) of aphid brain and was induced by high maternal density

and contact treatments. Cy5-conjugated *ApvGluT2* probe was shown in red, and nuclei

were stained with DAPI in blue. The selected neuropils (La, lamina; Me, medulla; LoX,

lobula complex; PB, protocerebral bridge) were outlined by dotted lines. (K) The

relative intensity of *ApvGluT2* in the antennal lobes was quantified by LAS X (n >6).

(L) The proportion of winged offspring was declined by antennae amputation (n=36).

The Mann Whitney non-parametric test was used to compare means of the proportions

of winged offspring. Student *t* test was used to compare means of relative intensity of

ApvGluT2. *n.s.*, not significant, *P < 0.05, **P < 0.01.

Figure 3 Embryonic *ApmGluR3* relayed the transgenerational signal from *ApvGluT2* about the induction of winged offspring. (A) Declined proportion of winged offspring by injection of an mGluR antagonist LY341495 (n=47), in contrast to (B) increased proportion by agonist ADPC (n=39). (C) RNAi experimental design. (D) Knockdown of *ApmGluR3* in the embryo (n=3), and (E) reduced proportion of winged offspring (n=47). (F) RNAi efficiency of *ApvGluT2* in the head (n=4). (G) Effect of ds*ApvGluT2* on the proportion of winged offspring was rescued by ADPC (n=36). (H) The expression of *ApmGluR3* in the stage 20 embryos increased as the contact duration

increased. *ApmGluR3* probe conjugated with Cy5 was in red, and nuclei were stained

with DAPI in blue. Wing discs shown in confocal were highlighted by white circles. (I)

The relative intensity of *ApmGluR3* in the wing discs of stage 20 embryos was

quantified by LAS X (n>9). Mann Whitney non-parametric test was used to compare

the means of the proportion of winged offspring. Student *t* test was used to compare

means of transcript level or the relative intensity of *ApmGluR3*. *P < 0.05, **P < 0.01,

***P < 0.001.

Figure 4 ApvGluT2 and ApmGluR3 promoted phosphorylation of embryonic ApFoxO, facilitating the formation of winged aphids. (A) Hypothetical model of *Ap*mGluR3-regulated phosphorylation of *Ap*FoxO in the embryo that controls the wing dimorphism of aphids. (B) RNAi of *ApFoxO* in embryos (n>5), and (C) its effect on the proportion of winged offspring (n>54). (D) The phosphorylated ApFoxO level in the embryo was enhanced by two-adult contacting for 4-hr treatment(n=3) and (E) reduced by ds*ApmGluR3*, and (F) ds*ApvGluT2* (n=3), as determined by western blotting analyses. (G) The proportion of winged offspring was reduced by Akt antagonist MK2206 that directly targeted FoxO as a substrate (n=42). Mann Whitney non-parametric test was used to compare the means of the proportion of winged offspring. Student *t* test was used to compare means of gene transcript or protein level. *P < 0.05, **P < 0.01, ***P < 0.001.

Figure 5 ApFoxO transcriptionally suppressed _ApHh_ at stage 20 embryo. (A)

Knockdown of _ApmGluR3_ significantly reduced the transcripts of _hedgehog_ (_ApHh_)

and _Mob as tumor suppressor_ (_ApMats_) in stage 20 embryos (n=3). (B) RNAi efficiency

of _ApHh_ in stage 20 embryos (n=5). (C) The effects of ds_ApHh_ on the proportion of

winged offspring (n=42). (D) RNAi efficiency of _ApFoxO_ in stage 20 embryos (n=5).

(E) The proportion of winged offspring was rescued by knocking down of _ApHh_ in

ds*ApFoxO* aphids (n>40). (F) Sequence alignment of DNA binding domains (DBD)

among FoxO homologs from brown planthopper (NlFoxO), *Drosophila* (DmFoxO) and

human (hFoxO). (G) EMSA showing that Maltose-binding protein (MBP)/ DNA-

binding domain (DBD) of ApFoxO fusion failed to bind to *FRE1* (FoxO recognition

element 1), but (H) bound to *FRE2* on the *ApHh* promoter. Lane 1: positive control.

Lane 2: negative control. Lane 3: biotin-labeled *FRE* only. Lane 4: 0.5 μg recombinant

MBP +labeled FRE. Lane 5: 0.5 μg or Lane 6: 5 μg of recombinant MBP/DBD fusion

protein + labeled *FRE*. Lane 7: 0.5 μg recombinant MBP/DBD fusion protein + labeled

FRE + 20 μmol (5 ×) unlabeled *FRE* competitor, or Lane 8: 25 × unlabeled competitor.

Lane 9: 0.5 μg recombinant MBP/DBD fusion protein + 5 × biotin-labeled mutant FRE

(FRE_m). Lane 10: 0.5 μg recombinant MBP/DBD fusion protein + biotin-labeled FRE

+ 5 × biotin end-labeled FRE_m. Biotin-labeled *FRE* was all 4 μmol in each lane. (I)

Two-adult contacting increased *ApHh* transcripts only at stage 20 embryos relative to

solitary aphids (n=5). Mann Whitney non-parametric test was used to compare the

means of the proportion of winged offspring. Student *t* test was used to compare means

of gene transcript level. *n.s.* not significant, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

Figure 6 Illustration of how the ApvGluT2-mGluR3 module transgenerational controls wing morphs in response to physical contact. In solitary aphids, relatively high level of unphosphorylated ApFoxO binds to the *cis*-element of *ApHh* promoter, suppressing its expression at stage 20 embryo. As a result, the prevalence of wingless offspring was observed. Once experienced physical contact, maternal aphids increased the *ApvGluT2* expression in antennal lobes of the brain and *Ap*m*GluR3* in wing discs of the embryo. This is followed by increased phosphorylation of ApFoxO. The phosphorylated ApFoxO is transported from the nucleus to the cytoplasm, thereby releasing the transcriptional suppression of *ApHh*, leading to high proportion of wingless offspring. vGluT2, vesicular glutamate transporter 2; mGluR3, metabotropic glutamate receptor 3; FoxO, Forkhead transcription factor subgroup O; Hh, hedgehog; La, lamina; Me, medulla; Lox, lobula complex; PB, protocerebral bridge; MB,

830   mushroom body; CB, central body; LAL, lateral accessory lobe; AL, antennal lobe; DL,

831   dorsal lobe.