GTP cyclohydrolase II (gch2) **and axanthism in ball pythons: a new vertebrate model for**

pterin-based pigmentation

*Alan Garcia-Elfring, °Heather L. Roffey, *Jaren M. Abergas, *Andrew P. Hendry, and *Rowan D.

H. Barrett

*Department of Biology, McGill University, Montreal, QC, Canada

°Biology Department, Vanier College, Montreal, QC, Canada

Corresponding authors: rowan.barrett@mcgill.ca, alan.garcia-elfring@mail.mcgill.ca

Abstract:

Pterin pigments are responsible for many of the bright colours observed across the animal kingdom. However, unlike melanin, the genetics of pterin-based pigmentation has received relatively little attention in animal colouration studies. Here, we investigate a lineage of axanthic ball pythons (*Python regius*) found in captivity as a model system to study pterin pigmentation in vertebrates. By crowdsourcing shed skin samples from commercial breeders and applying a case-control study design, we utilized whole-genome pool sequencing (pool-seq) and variant annotation. We identified a premature stop codon in the gene *GTP cyclohydrolase II* (*gch2*), which is associated with the axanthic phenotype. GCH2 catalyzes the first rate-limiting step in riboflavin biosynthesis. This study provides the first identification of an axanthism-associated gene in vertebrates and highlights the utility of ball pythons as a model to study pterin-based pigmentation.

INTRODUCTION

Animal pigmentation has long served as a valuable model for understanding genetics, development, and evolution (Cott 1940; Kronforst et al. 2012; Caro 2017; Endler et al. 2017; San-Jose et al. 2017). Vertebrate pigmentation arises from the differential absorption of light by pigments within specialized cells called melanophores (melanocytes in endothermic vertebrates) and xanthophores. All vertebrates produce a brown/black form of melanin derived from the amino acid tyrosine (Wakamatsu and Ito 2021). In addition, poikilothermic vertebrates (fish, amphibians, and reptiles) display bright colouration through yellow/red pigments found in cells called xanthophores. Xanthophores derive their vibrant yellow or red colouration from pterin or carotenoid pigments. While vertebrates acquire carotenoids exclusively from their diet (Maoka 2020), pterin pigments are synthesized de novo from guanosine triphosphate (GTP). Coloration can also be structural, with poikilothermic vertebrates using guanine nanocrystals in iridophore cells to generate structural colouration via the constructive interference of light. Although melanin-based pigmentation has been extensively studied, other pigment classes, such as pterins, have received less attention.

The mouse has been pivotal in studies of melanocyte biology, helping uncover the genes responsible for melanocyte development and patterning: *Kit*, *Pax3*, *Mitf*, *Edn3*, *Ednrb*, and *Sox10* (Steingrímsson et al. 2005); melanin synthesis: *Tyr*, *Tyrp1*, *Dct* (McNamara et al. 2021; Wakamatsu and Ito 2021); and its regulation, such as *Mc1r* (Garcia-Borron et al. 2005). Meanwhile, the zebrafish model has laid the foundation for understanding poikilothermic vertebrate pattern formation (Singh and Nüsslein-Volhard 2015). Studies on zebrafish have

shown that pattern formation arises from local, short- and long-range cell-cell interactions

between chromatophore types (Frohnhöfer et al. 2013). For example, zebrafish mutants lacking

any one of the chromatophores (iridophores, melanophores, and xanthophores) fail to develop

stripes and often develop spots instead. In double mutants lacking two of the three

chromatophore types, the remaining chromatophores uniformly cover the body (Frohnhöfer et

al. 2013; Owen et al. 2020). Gap junction proteins mediate these cell-cell interactions, and

mutations in these genes can result in labyrinthine patterning (Irion et al. 2014). However, it

remains unclear how widely zebrafish cell-cell interactions apply across vertebrate taxa.

First discovered in butterfly wings (Hopkins 1895), pterin-based pigmentation is

ubiquitous in nature but has received less attention compared to melanin-based pigmentation

(Andrade and Carneiro 2021). The pterin pigment synthesis pathway has been largely

elucidated through studies in *Drosophila* (Glassman and Mitchel 1959; Courtright 1967; Ziegler

and Harmsen 1970; Goto and Sugiura 1971; Evans and Howells 1978; Spradling and Rubin 1983;

Reaume et al. 1991; Kim et al. 2013). Notably, *Drosophila*'s first described mutant, "white,"

lacked eye pterin (drosopterin) and ommochrome pigments (Morgan 1910). While melanin is

often linked to adaptive phenotypes, such as cryptic colouration (Barrett et al. 2019; Harris et

al. 2020; Baltazar-Soares et al. 2024), pterin pigments are key in signaling both within and

between species (Kikuchi and Pfennig 2012; Kikuchi et al. 2014; Andrade and Carneiro 2021). In

vertebrates, zebrafish have been the primary model for studying xanthophore biology,

including xanthophore specification from neural crest cells (Kelsh et al. 1996; Odenthal et al.

1996; Parichy et al. 2000; Minchin and Hughes 2008) and pigment synthesis (Ziegler 2003; Lister

2019).

Studying mutants lacking a pigment can offer insights into bright colouration genetics, a

strategy pioneered by Thomas Hunt Morgan with fruit flies. Axanthic animals, which lack yellow

and red pigments but retain melanin, are rare among vertebrates. Only a few axanthic

amphibians and reptiles have been discovered in nature (Bechtel 1991; Jablonski et al. 2014;

Cattaneo 2015; Kolenda et al. 2017; Cavalcante and Bruni 2018; Borteiro et al. 2021;

Schluckebier et al. 2022). Axanthic vertebrate models are also scarce in pigmentation research,

with the axanthic axolotl, first described by Lyerla and Dalton (1971), being a notable exception

(Dunson 1974; Frost et al. 1986; Bukowski et al. 1990; Masselink and Tanaka 2021). Axanthic

axolotls lack both xanthophores and iridophores, meaning axanthism in axolotls affects

chromatophore development rather than pigment synthesis. However, low survivability in

axanthic axolotls presents challenges for research, including breeding programs, population

genetics studies, and comparative analyses.

Here, we use a line of axanthic ball pythons to study the genetics of yellow pigmentation

in reptiles. We hypothesize that, assuming a similar mechanism of vertebrate pattern formation

to zebrafish (Frohnhöfer 2013; Singh and Nüsslein-Volhard 2015), axanthism in ball pythons

results from defects in pterin pigment synthesis (Ziegler 2003), rather than carotenoid

metabolism or xanthophore development. We expect that xanthophore development remains

unaffected, as axanthic ball pythons exhibit wild-type melanin patterning, which suggests intact

cell-cell interactions. Furthermore, we anticipate that yellow pigmentation (or its absence) is

linked to pterin-based pigmentation, rather than carotenoids, since ball pythons are born with

yellow colouration before acquiring carotenoids from their diet. This suggests that pterins, not

carotenoids, underlie the colouration difference.

METHODS

Sample collection, study design, DNA extraction, and sequencing

We crowd-sourced ball python shed skin samples from breeders, including five samples with

the recessive VPI Axanthic colour morph (Table S1). VPI refers to Vida Preciosa International,

the reptile propagation company that first received the wild-caught axanthic snake from

Western Africa and established the phenotype in captivity. All VPI Axanthic (hereafter referred

to as 'axanthic') ball pythons descend from this individual.

Additionally, we collected 38 samples of the recessive 'Clown' phenotype (Table S2),

which we used as a control for identifying candidate mutations linked to the axanthic trait. We

also gathered samples from other colour morphs: 'Spider' (n=26, Table S3), 'Enchi' (n=7, Table

S4), 'Ivory' (n=15, Table S5), and 'Lavender' (n=14, Table S6). These were used as further control

sets because, according to breeder pedigrees, these morphs do not carry any axanthism

variation.

Similarly, the axanthic samples were confirmed to lack any variation associated with the

Clown, Spider, Enchi, Ivory, or Lavender morphs (i.e., inferred to be homozygous for the

reference alleles at those loci). We identified candidate mutations by comparing genetic

variation in the axanthic pool with multiple reference sample sets known, through pedigree

analysis, to not carry the axanthic variant.

The wild-type ball python colour morph displays a black background with tan-coloured

lateral saddles and traces of yellow pigmentation (Figure 1A). Comparing the wild type to the

albino colour morph highlights normal xanthophore pigmentation and patterning: in non-

axanthic individuals, yellow pigment typically overlaps with the tan-coloured saddles and is

absent from the dark melanic regions (Figure 1B). In contrast, axanthic ball pythons lack yellow

pigmentation but retain the wild-type melanin pattern (Figure 1C).

We extracted and sequenced DNA from the shed skins using a standard three-day

phenol-chloroform procedure. DNA quantification and quality checks were performed with an

Infinite® 200 Nanoquant (Tecan Group Ltd., Männedorf, Switzerland). After normalizing the

DNA from each sample, we combined them into pools in equimolar amounts. These pooled

samples (case and controls) were then sent to the McGill Genome Centre for whole-genome

sequencing. Library preparation was performed using a PCR-based method, and sequencing

was carried out with 150 bp paired-end reads on one lane of the Illumina NovaSeq 6000

platform.

Bioinformatics

We used the trim-fastq.pl program from Popoolation (Kofler et al. 2011a) to filter raw reads

based on quality (--quality-threshold 20) and length (--min-length 50). This step ensured that

each read had a minimum quality score of 20 and a minimum length of 50 bp. We then mapped

the filtered reads to a reference genome using the bwa-mem program with default parameters

(Li and Durbin, 2009). Since a ball python reference genome is not available, we aligned the

reads to both the draft Burmese python reference genome *Pmo2.0* (Castoe et al. 2013) and the

more recent chromosome-length Burmese python assembly (*Python molurus bivittatus-*

5.0.2_HiC), available at DNAzoo.org. This chromosome-length assembly, based on Hi-C data

(Dudchenko et al. 2017; Dudchenko et al. 2018), is aligned to the annotated draft genome

(Castoe et al. 2013) and has been used in previous studies to identify colour-related loci

(Garcia-Elfring et al. 2023).

Next, we converted SAM files to BAM format using SAMtools (Li et al. 2009) and

removed reads with a mapping quality below 20 (-q 20). We then generated mpileup and sync

formats using SAMtools and Popoolation2 (Kofler et al. 2011b), respectively, before proceeding

to variant calling. Variants were called and filtered with bcftools (vcfutils.pl varFilter) using a

minimum depth of five (-d 5) and a mapping quality of 20 (-Q 20).

We calculated SNP-specific F_{ST} estimates using fst-sliding.pl from Popoolation2 (Kofler et

al. 2011b). F_{ST} values were generated for SNP variants in VPI Axanthic (n=5) compared to the

control pools. Since the control pools (e.g., Clown, Spider, etc.) lacked axanthic variation, we

expected the derived allele to be fixed in the axanthic samples and the reference allele to be

fixed in the control pool. A threshold F_{ST} value of 1.0 was used to identify candidate mutations.

To confirm the genomic region and candidate mutation of interest, we generated F_{ST} values

between the axanthic samples and additional control pools (Spider, n=26; Enchi, n=7; Ivory,

n=15; Lavender, n=14).

Variant annotation and identification of polymorphisms and fixed differences

We used the software SnpEff (Cingolani et al. 2012) to annotate variants mapped to

both the draft and chromosome-length assemblies. SnpEff utilizes a genome assembly and its

corresponding General Feature Format (GFF3) file to assess the functional impact of variants on

genomic features. For genomic regions showing elevated FST values ("F_{ST} peaks"), we focused

on individual variants with an F_{ST} of 1.0, that mapped to protein-coding regions and were

predicted to affect gene products. These included nonsense mutations and small indels

classified as "high impact," and missense mutations classified as "moderate impact," which

were considered candidate mutations.

To visualize the position and effect of a candidate mutation on the protein, we used the

program Protter (Omasits et al. 2016). We confirmed the candidate variant by PCR amplification

and Sanger sequencing of the locus using samples that were not included in the pooled

libraries. We validated the candidate loci by Sanger sequencing samples with genotypes

inferred from breeder pedigrees. These included inferred heterozygotes (n=10), homozygotes

for the derived axanthism-causing allele (i.e., axanthic individuals, n=6), and inferred

homozygotes for the wild-type allele (n=10).

Primers used to amplify the region surrounding the candidate locus were designed using

Primer3 (Kõressaar and Remm, 2007; Kõressaar et al. 2018). The primers used for PCR

amplification were as follows:

• Forward: 5'- GAAACTTTTGACCAGACTCAGTC

• Reverse: 5'- CTTCAATCACCACAGCCACTCC

Synteny and phylogenetic analysis of gch1 and gch2

We conducted a synteny analysis and constructed a phylogenetic tree for the *gch1* and *gch2*

genes across multiple species: Burmese python (*Python bivittatus*), brown anole (*Anolis sagrei*),

coelacanth (*Latimeria chalumnae*), and the elephant shark (*Callorhinchus milii*). To ensure focus

on conserved regions, we trimmed the gene sequences (555 bp) before alignment. Due to the

evolutionary divergence of these species, we used MAFFT to build the tree(Katoh et al. 2002).

Phylogenetic trees were then built using FastTree (Price et al. 2010), which estimates

maximum-likelihood phylogenies, allowing us to infer the evolutionary relationships of these

genes across species.

RESULTS

Our whole-genome sequencing and variant calling identified 26,449,779 SNPs across all samples

received from ball python breeders. Analysis of allele frequency differences between the

axanthic pool and the control pool (Clown colour morph) demonstrates overall high levels of

background F_{ST}, likely resulting from the low sample size for the axanthic phenotype (n=5). We

find two areas of high differentiation or F_{ST} peaks. One region (~24.8 Mb) on chromosome two

(Figure 2A) and a second peak on chromosome three, corresponding to the region containing

the Clown mutation (Garcia-Elfring et al. *in prep*). Comparing the axanthic data with additional

control pools (Spider, n=26; Enchi, n=7; Ivory, n=15; Lavender, n=14) confirms that the region of

high differentiation on chromosome two is associated with axanthism (Figure S1). Within the

peak on chromosome two, one variant had both high levels of differentiation (F_{ST} = 1) and is

predicted to have a high impact on the gene product. This variant is a nonsense mutation, and

it maps to exon 5 of LOC103050242 (*GTP cyclohydrolase 1-like*) in the draft assembly. The

candidate SNP consists of a C to T nucleotide change (c.520C>T), changing an arginine residue

(CGA) to a TGA stop codon (p.Arg174*). This genetic variant is expected to result in a truncated

protein (Figure 2B). We validated the axanthic variant found in poolseq data by PCR

amplification and Sanger sequencing of the locus of interest in additional individuals (inferred

heterozygotes: n=10; homozygotes for the derived axanthism-causing allele: n=6; inferred

homozygotes for the wild type allele: n=10). All individuals tested had the genotype that was

inferred from pedigrees by breeders.

Based on the annotation of *gch1* in the Burmese python reference genome and the

absence of an annotated *gch2*, we hypothesized that the gene LOC103050242 (*GTP*

cyclohydrolase 1-like) corresponds to the paralog *gch2*. A phylogenetic analysis shows that the

Burmese python gene LOC103050242 is more closely related to the brown anole

LOC132761499 and coelacanth *gch2* than to Burmese python *gch1* (Figure S2). Synteny analysis

further supports the identification of LOC103050242 as *gch2*. In humans, coelacanth, and

python, *gch1* is flanked by *samd4a*, while in coelacanth and python, *gch2* (LOC103050242 in

python) is flanked by *pigf* and *cript* (Figure S3).

DISCUSSION

Captive-bred ball pythons have emerged as a valuable model system for studying pigmentation

biology, offering unique opportunities to explore the genetics of colouration in reptiles (Brown

et al. 2022; Dao et al. 2023; Garcia-Elfring et al. 2023; Lederer et al. 2023). In this study, we

focused on an axanthic line of ball pythons to investigate the genetic basis of yellow

pigmentation in these snakes. While yellow and red hues in vertebrate integument can arise

from dietary carotenoids or endogenous pigments, the carnivorous diet of snakes, combined

with the yellow colouration observed in wildtype ball pythons, led us to hypothesize that

endogenous pigments are the primary source of yellow colouration in this species. Given the

normal melanin pattern seen in axanthic ball pythons, we speculated that any candidate

mutation would likely be found in a gene associated with pigment biosynthesis rather than in

one related to xanthophore cell-fate specification.

Our analysis identified a mutation in *gch2*, a gene known as *ribA* in bacteria, including *E.*

coli, where it was first isolated and extensively studied (Richter et al. 1993; Averianova et al.

2020). *Gch2* plays a critical role in the biosynthesis of riboflavin (vitamin B2) and the production

of flavin nucleotide coenzymes, which are vital for numerous enzymatic reactions (Bacher et al.

2000; Ren et al. 2005). Unlike animals, bacteria can synthesize riboflavin de novo through an

ancestrally conserved pathway, like that found in plants and fungi (Anam et al. 2020;

Averianova et al. 2020; Nasuno et al. 2022; Tian et al. 2022). Bacterial studies have shown that

gch2 catalyzes the conversion of GTP into 2,5-diamino-6-ribosyl-amino-4(3H)-pyrimidinedione

5⬚-phosphate (DARPP), releasing formate and pyrophosphate as byproducts (Averianova et al.

2020; Klein et al. 2023). In animals, although *gch2* can catalyze the GTP to DARPP reaction, its

physiological role is considered non-essential since the riboflavin pathway is not fully

functional.

The precise function of *gch2* in animals remains less clear compared to that of *gch1*

(*GTP cyclohydrolase I*), which is essential for the synthesis of tetrahydrobiopterin (BH4), a

cofactor for enzymes involved in neurotransmitter production (e.g., dopamine, serotonin) and

nitric oxide synthesis. It also serves as a marker for the xanthophore lineage (Thöny et al. 2000;

Ziegler et al. 2000; Pelletier et al. 2001; Nagao et al. 2014; Hamied et al. 2020; Wu et al. 2022).

In humans, mutations in *gch1* lead to severe metabolic and neurological disorders (Nagatsu

Ichinose, 1996; Longo, 2009; Fanet et al. 2021; Himmelreich et al. 2021), such as

phenylketonuria (Trujillano et al. 2017) and DOPA-responsive dystonia (Hirano and Ueno 1999;

Yoshino et al. 2018; Kostić et al. 2020; Bradley et al. 2021). *Gch1* and BH4 are also crucial for

embryonic development, with homozygous mutations being lethal in mice and zebrafish

(Douglas et al. 2015; Larbalestier et al. 2022).

Although *gch2* and *gch1* share functional similarities, such as guanine ring opening,

formate release, and zinc binding, the reactions they catalyze differ in significant ways

(Auerbach et al. 2000; Kaiser et al. 2002; Ren et al. 2005). The role of *gch2* in vertebrate

pigmentation is less well understood, but it has been identified as an early marker for

xanthophores (Parichy et al. 2000; Miyadai et al. 2023). In zebrafish, *gch2* is essential for larval

xanthophore pigmentation but not in adults (Lister, 2019). Conversely, research in flounder

suggests that *gch2* is necessary for adult melanophore and xanthophore differentiation

(Watanabe et al. 2008).

We discovered a nonsense mutation at residue 174 (p.Arg174*), likely resulting in a

non-functional protein, occurring 16 residues upstream of the conserved histidine believed to

be critical for producing DARPP from GTP (Yadav et al. 2020). Our findings indicate that

xanthophore pigmentation in ball pythons arises from the riboflavin biosynthesis pathway,

rather than the biopterin pathway. Additionally, since loss-of-function mutations in *gch1* often

lead to severe developmental defects, the involvement of *gch2*—rather than *gch1*—is

consistent with the absence of pleiotropic effects in axanthic ball pythons.

The extensive phenotypic variation in ball python colour morphs and the wide range of

colour combinations breeders have produced in captivity provide an opportunity to explore and

test hypotheses regarding the mechanisms behind pattern formation. Recent studies suggest

that all ball pythons lack iridophores, one of the three chromatophore types (Tzika 2024). This

makes ball pythons natural analogs to Shd mutant zebrafish, which also lack iridophores

(Frohnhöfer et al., 2013). In addition to axanthism, ball python colour morphs can exhibit

xanthism, which is characterized by an increase in yellow brightness (possibly due to melanin

changes) or in the distribution of yellow pigment (e.g., the 'Enchi' morph). Many ball python

colour morphs that seem to affect melanophore migration (e.g., Tri-Stripe, which shows altered

melanin patterns) display xanthophore pigmentation in regions typically covered by dark

melanin. This non-overlap of black and yellow pigments suggests that xanthophores and

melanophores repel each other, as observed in zebrafish (Frohnhöfer et al., 2013).

In zebrafish double mutants (Shd;Pfe), where two of the three chromatophore types

(e.g., iridophores and xanthophores) are absent, the remaining chromatophore type (e.g.,

melanophores) uniformly covers the body, uninhibited by cell-cell interactions. Similarly, we

hypothesize that the genetic basis of all-black melanic ball python morphs, such as 'Cinnamon'

(Figure 1H), is likely due to mutations in genes essential for xanthophore development, allowing

melanophores to spread across the body. This contrasts with mutations in melanin production

genes, like agouti, which are more common in mammals (e.g., Sasamori et al. 2017; Reissmann

et al. 2020). Further support for a repulsive interaction between xanthophores and

melanophores comes from the Albino form of the Cinnamon morph (Figure 1I). Instead of

revealing an underlying pterin pigment pattern, which might be expected if melanism was

driven by excessive melanin production, the 'Cinnamon Albino' phenotype presents as an

entirely white ball python with no pterin pigmentation (Figure 1I).

Other recessive lines of axanthism exist in captive-bred ball pythons, such as the 'MJ'

line. Notably, VPI Axanthic and other axanthism lines are not allelic with one another, meaning

crosses between different lines result in double heterozygotes with wildtype coloration,

suggesting distinct genetic causes for each axanthism lineage. These various axanthism lines,

when combined with other color variants (Figure 1D), provide valuable models to further

explore the pterin pigment biosynthesis pathway and identify new axanthism-causing genes in

reptiles. Consistent with our findings, we demonstrated in the VPI lineage that pterin pigment

metabolism, rather than carotenoid metabolism, is associated with yellow coloration in ball

pythons.

In summary, our study highlights a significant link between *gch2* and axanthism in ball

pythons, suggesting that the initial stages of the ancestral riboflavin pathway are crucial for

yellow pigmentation. Further research is necessary to identify the specific xanthophore

pigments involved in ball python pigmentation. Nonetheless, our findings enhance the

understanding of the genetics of pigmentation in reptiles.

CONCLUSION

We used ball python colour morphs to study the genetic basis of xanthophore pigmentation in

snakes. We found axanthism is associated with a key gene in the biopterin biosynthesis

pathway, providing the first evidence that pterins, not carotenoids, produce the yellow pigment

observed in ball pythons. To our knowledge, *gch1* is the first axanthism-associated gene

identified in vertebrates. Our results provide new research avenues for the study of axanthism

and xanthophore pigmentation in snakes and highlights the utility of using captive-bred ball

pythons in pigmentation research.

REFERENCES

Anam, K., Nasuno, R. and Takagi, H., 2020. A novel mechanism for nitrosative stress tolerance

dependent on GTP cyclohydrolase II activity involved in riboflavin synthesis of yeast.

Scientific Reports, 10(1), p.6015.

Andrade, P. and Carneiro, M., 2021. Pterin-based pigmentation in animals. Biology Letters,

17(8), p.20210221.

Auerbach, G., Herrmann, A., Bracher, A., Bader, G., Gütlich, M., Fischer, M., Neukamm, M.,

Garrido-Franco, M., Richardson, J., Nar, H. and Huber, R., 2000. Zinc plays a key role in

human and bacterial GTP cyclohydrolase I. Proceedings of the National Academy of

Sciences, 97(25), pp.13567-13572.

Averianova, L.A., Balabanova, L.A., Son, O.M., Podvolotskaya, A.B. and Tekutyeva, L.A., 2020.

Production of vitamin B2 (riboflavin) by microorganisms: an overview. Frontiers in

Bioengineering and Biotechnology, 8, p.570828.

Bacher, A., Eberhardt, S., Fischer, M., Kis, K. and Richter, G., 2000. Biosynthesis of vitamin b2

(riboflavin). Annual review of nutrition, 20(1), pp.153-167.

Baltazar-Soares, M., Karell, P., Wright, D., Nilsson, J.Å. and Brommer, J.E., 2024. Genomic basis

of melanin-associated phenotypes suggests colour-specific environmental adaptations

in tawny owls. *Molecular Ecology*, p.e17247.

Barrett, R.D., Laurent, S., Mallarino, R., Pfeifer, S.P., Xu, C.C., Foll, M., Wakamatsu, K., Duke-

Cohan, J.S., Jensen, J.D. and Hoekstra, H.E., 2019. Linking a mutation to survival in wild

mice. Science, 363(6426), pp.499-504.

Bechtel, H.B., 1991. Inherited colour defects: Comparison between humans and snakes.

International Journal of Dermatology, 30(4), pp.243-246.

Borteiro, C., Diesel Abegg, A., Hirouki Oda, F., Cardozo, D.E., Kolenc, F., Etchandy, I., Bisaiz, I.,

Prigioni, C. and Baldo, J.D., 2021. Aberrant colouration in wild snakes: case study in

Neotropical taxa and a review of terminology.

Bradley, M., Lyons, S., Coleman, S., Vance, R., Molloy, F. and Widdess-Walsh, P., 2021. A Case of

GCH-1 Mutation Dopa-Responsive Dystonia Requiring High Doses of Levodopa for

Treatment. Tremor and Other Hyperkinetic Movements, 11.

Brown, A.R., Comai, K., Mannino, D., McCullough, H., Donekal, Y., Meyers, H.C., Graves, C.W.,

Seidel, H.S. and BIO306W Consortium, 2022. A community-science approach identifies

genetic variants associated with three colour morphs in ball pythons (Python regius).

Plos one, 17(10), p.e0276376.

Bukowski, L., Erickson, K. and Lyerla, T.A., 1990. Characterization of the yellow pigment in the

axanthic mutant of the Mexican axolotl, Ambystoma mexicanum. Pigment Cell Research,

3(3), pp.123-125.

Caro, T., 2017. Wallace on colouration: contemporary perspective and unresolved insights.

Trends in ecology & evolution, 32(1), pp.23-30.

Castoe, T.A., De Koning, A.J., Hall, K.T., Card, D.C., Schield, D.R., Fujita, M.K., Ruggiero, R.P.,

Degner, J.F., Daza, J.M., Gu, W. and Reyes-Velasco, J., 2013. The Burmese python

genome reveals the molecular basis for extreme adaptation in snakes. Proceedings of

the National Academy of Sciences, 110(51), pp.20645-20650.

Cattaneo, A., 2015. The Zamenis longissimus (Laurenti) axanthic phenotype found on the

Castelporziano Presidential Estate: considerations on its morphology, genetic nature

and probable extinction (Serpentes: Colubridae). Rendiconti Lincei, 26, pp.385-389.

Cavalcante, R. and Bruni, G., 2018. Axanthism in Emys orbicularis hellenica (Valenciennes, 1832)

(Testudines: Emydidae) from Piedmont, northern Italy. The Herpetological Bulletin, 146,

pp.36-38.

Cingolani, P., Platts, A., Wang, L.L., Coon, M., Nguyen, T., Wang, L., Land, S.J., Lu, X. and Ruden,

D.M., 2012. A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118;

iso-2; iso-3. Fly, 6(2), pp.80-92.

Cott, H.B., 1940. Adaptive colouration in animals. Methuen & Co (1940)

Courtright, J.B., 1967. Polygenic control of aldehyde oxidase in Drosophila. Genetics, 57(1),

p.25.

Dao, U.M., Lederer, I., Tabor, R.L., Shahid, B., Graves, C.W. and Seidel, H.S., 2022. Leucism and

stripe formation in ball pythons (Python regius) are associated with variants affecting

endothelin signaling. bioRxiv.

Douglas, G., Hale, A.B., Crabtree, M.J., Ryan, B.J., Hansler, A., Watschinger, K., Gross, S.S.,

Lygate, C.A., Alp, N.J. and Channon, K.M., 2015. A requirement for Gch1 and

tetrahydrobiopterin in embryonic development. Developmental biology, 399(1), pp.129-

138.

Dudchenko, O., Batra, S.S., Omer, A.D., Nyquist, S.K., Hoeger, M., Durand, N.C., Shamim, M.S.,

Machol, I., Lander, E.S., Aiden, A.P. and Aiden, E.L., 2017. De novo assembly of the

Aedes aegypti genome using Hi-C yields chromosome-length scaffolds. Science, 356(6333), pp.92-95.

Dudchenko, O., Shamim, M.S., Batra, S.S., Durand, N.C., Musial, N.T., Mostofa, R., Pham, M., Hilaire, B.S., Yao, W., Stamenova, E. and Hoeger, M., 2018. The Juicebox Assembly Tools module facilitates de novo assembly of mammalian genomes with chromosome-length scaffolds for under $1000. BioRxiv, 254797. Preprint.

Dunson, M.K., 1974. Ultrastructure of pigment cells in wild type and colour mutants of the Mexican axolotl. Cell and Tissue Research, 151, pp.259-268.

Endler, J.A. and Mappes, J., 2017. The current and future state of animal colouration research. Philosophical Transactions of the Royal Society B: Biological Sciences, 372(1724), p.20160352.

Evans, B.A. and Howells, A.J., 1978. Control of drosopterin synthesis in Drosophila melanogaster: mutants showing an altered pattern of GTP cyclohydrolase activity during development. Biochemical genetics, 16(1-2), pp.13-26.

Fanet, H., Capuron, L., Castanon, N., Calon, F. and Vancassel, S., 2021. Tetrahydrobioterin (BH4) pathway: from metabolism to neuropsychiatry. Current Neuropharmacology, 19(5), pp.591-609.

Frohnhöfer, H.G., Krauss, J., Maischein, H.M. and Nüsslein-Volhard, C., 2013. Iridophores and their interactions with other chromatophores are required for stripe formation in zebrafish. Development, 140(14), pp.2997-3007.

Frost, S.K., Epp, L.G. and Robinson, S.J., 1986. The pigmentary system of developing axolotls: IV.

An analysis of the axanthic phenotype.

García-Borrón, J.C., Sánchez-Laorden, B.L. and Jiménez-Cervantes, C., 2005. Melanocortin-1

receptor structure and functional regulation. *Pigment cell research*, *18*(6), pp.393-410.

Garcia-Elfring, A., Sabin, C.E., Iouchmanov, A.L., Roffey, H.L., Samudra, S.P., Alcala, A.J., Osman,

R.S., Lauderdale, J.D., Hendry, A.P., Menke, D.B. and Barrett, R.D., 2023. Piebaldism and

chromatophore development in reptiles are linked to the tfec gene. Current Biology,

33(4), pp.755-763.

Glassman, E. and Mitchell, H.K., 1959. Mutants of Drosophila melanogaster deficient in

xanthine dehydrogenase. Genetics, 44(2), p.153.

Goto, M. and Sugiura, K., 1971. Biosynthesis of pteridines in Drosophila melanogaster and Rana

catesbeiana. In Methods in Enzymology (Vol. 18, pp. 746-761). Academic Press.

Hamied, A., Alnedawy, Q., Correia, A., Hacker, C., Ramsdale, M., Hashimoto, H. and Kudoh, T.,

2020. Identification and characterization of highly fluorescent pigment cells in embryos

of the Arabian killifish (Aphanius Dispar). Iscience, 23(11).

Harris, R.B., Irwin, K., Jones, M.R., Laurent, S., Barrett, R.D., Nachman, M.W., Good, J.M.,

Linnen, C.R., Jensen, J.D. and Pfeifer, S.P., 2020. The population genetics of crypsis in

vertebrates: recent insights from mice, hares, and lizards. Heredity, 124(1), pp.1-14.

Soare

Himmelreich, N., Blau, N. and Thöny, B., 2021. Molecular and metabolic bases of

tetrahydrobiopterin (BH4) deficiencies. Molecular Genetics and Metabolism, 133(2),

pp.123-136.

Hirano, M. and Ueno, S., 1999. Mutant GTP cyclohydrolase I in autosomal dominant dystonia

and recessive hyperphenylalaninemia. Neurology, 52(1), pp.182-182.

Hopkins, F.G., 1895. XV. The pigments of the Pieridæ: a contribution to the study of excretory

substances which function in ornament. Philosophical Transactions of the Royal Society

of London.(B.), (186), pp.661-682.

Irion, U., Frohnhöfer, H.G., Krauss, J., Çolak Champollion, T., Maischein, H.M., Geiger-Rudolph,

S., Weiler, C. and Nüsslein-Volhard, C., 2014. Gap junctions composed of connexins 41.8

and 39.4 are essential for colour pattern formation in zebrafish. Elife, 3, p.e05125.

Jablonski, D., Alena, A., Vlček, P. and Jandzik, D., 2014. Axanthism in amphibians: A review and

the first record in the widespread toad of the Bufotes viridis complex (Anura:

Bufonidae). Belgian Journal of Zoology, 144(2).

Kaiser, J., Schramek, N., Eberhardt, S., Püttmer, S., Schuster, M. and Bacher, A., 2002.

Biosynthesis of vitamin B2: An essential zinc ion at the catalytic site of GTP

cyclohydrolase II. European journal of biochemistry, 269(21), pp.5264-5270.

Katoh, K. and Toh, H., 2008. Recent developments in the MAFFT multiple sequence alignment

program. Briefings in bioinformatics, 9(4), pp.286-298.

Kelsh, R.N., Brand, M., Jiang, Y.J., Heisenberg, C.P., Lin, S., Haffter, P., Odenthal, J., Mullins,

M.C., Van Eeden, F.J., Furutani-Seiki, M. and Granato, M., 1996. Zebrafish pigmentation

mutations and the processes of neural crest development. Development, 123(1),

pp.369-389.

Kikuchi, D.W. and Pfennig, D.W., 2012. A Batesian mimic and its model share colour production

mechanisms. Current Zoology, 58(4), pp.658-667.

Kikuchi, D.W., Seymoure, B.M. and Pfennig, D.W., 2014. Mimicry's palette: widespread use of

conserved pigments in the aposematic signals of snakes. Evolution & Development,

16(2), pp.61-67.

Kim, H., Kim, K. and Yim, J., 2013. Biosynthesis of drosopterins, the red eye pigments of

Drosophila melanogaster. IUBMB life, 65(4), pp.334-340.

Klein, V.J., Brito, L.F., Perez-Garcia, F., Brautaset, T. and Irla, M., 2023. Metabolic engineering of

thermophilic Bacillus methanolicus for riboflavin overproduction from methanol.

Microbial Biotechnology, 16(5), pp.1011-1026.

Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R.V., Nolte, V., Futschik, A., Kosiol, C. and

Schlötterer, C., 2011a. PoPoolation: a toolbox for population genetic analysis of next

generation sequencing data from pooled individuals. PloS one, 6(1), p.e15925.

Kofler, R., Pandey, R.V. and Schlötterer, C., 2011b. PoPoolation2: identifying differentiation

between populations using sequencing of pooled DNA samples (Pool-Seq).

Bioinformatics, 27(24), pp.3435-3436.

Kolenda, K., Najbar, B., Najbar, A., Kaczmarek, P., Kaczmarski, M. and Skawiński, T., 2017. Rare

colour aberrations and anomalies of amphibians and reptiles recorded in Poland.

Herpetology Notes, 10, pp.103-109.

Kõressaar, T. and Remm, M., 2007. Enhancements and modifications of primer design program

Primer3. Bioinformatics, 23(10), pp.1289-1291.

Kõressaar, T., Lepamets, M., Kaplinski, L., Raime, K., Andreson, R. and Remm, M., 2018.

Primer3_masker: integrating masking of template sequence with primer design

software. Bioinformatics, 34(11), pp.1937-1938.

Kostić, V.S., Agosta, F., Tomic, A., Sarasso, E., Kresojevic, N., Basaia, S., Svetel, M., Copetti, M.

and Filippi, M., 2020. Brain structural alterations in patients with GCH1 mutations

associated DOPA-responsive dystonia. Journal of Neurology, Neurosurgery & Psychiatry.

Kronforst, M.R., Barsh, G.S., Kopp, A., Mallet, J., Monteiro, A., Mullen, S.P., Protas, M.,

Rosenblum, E.B., Schneider, C.J. and Hoekstra, H.E., 2012. Unraveling the thread of

nature's tapestry: the genetics of diversity and convergence in animal pigmentation.

Pigment cell & melanoma research, 25(4), pp.411-433.

Larbalestier, H., Keatinge, M., Watson, L., White, E., Gowda, S., Wei, W., Koler, K., Semenova,

S.A., Elkin, A.M., Rimmer, N. and Sweeney, S.T., 2022. GCH1 deficiency activates brain

innate immune response and impairs tyrosine hydroxylase homeostasis. Journal of

Neuroscience, 42(4), pp.702-716.

Lederer, I., Shahid, B., Dao, U., Brogdon, A., Byrtus, H., Delva, M., Deva, O., Hatfield, P., Hertz,

M., Justice, J. and Mavor, S., 2023. A frameshift variant in the melanophilin gene is

associated with loss of pigment from shed skin in ball pythons (Python regius).

Micropublication Biology, 2023.

Li, H. and Durbin, R., 2009. Fast and accurate short read alignment with Burrows–Wheeler

transform. bioinformatics, 25(14), pp.1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G. and

Durbin, R., 2009. The sequence alignment/map format and SAMtools. Bioinformatics,

25(16), pp.2078-2079.

Lister, J.A., 2019. Larval but not adult xanthophore pigmentation in zebrafish requires GTP

cyclohydrolase 2 (gch2) function. Pigment Cell & Melanoma Research, 32(5), pp.724-

727.

Longo, N., 2009. Disorders of biopterin metabolism. Journal of inherited metabolic disease, 32,

pp.333-342.

Lyerla, T.A. and Dalton, H.C., 1971. Genetic and developmental characteristics of a new colour

variant, axanthic, in the Mexican axolotl, Ambystoma mexicanum Shaw. Developmental

Biology, 24(1), pp.1-18.

Maoka, T., 2020. Carotenoids as natural functional pigments. Journal of natural medicines,

74(1), pp.1-16.

Masselink, W. and Tanaka, E.M., 2021. Toward whole tissue imaging of axolotl regeneration.

Developmental Dynamics, 250(6), pp.800-806.

McNamara, M.E., Rossi, V., Slater, T.S., Rogers, C.S., Ducrest, A.L., Dubey, S. and Roulin, A.,

2021. Decoding the evolution of melanin in vertebrates. Trends in Ecology & Evolution.

Miyadai, M., Takada, H., Shiraishi, A., Kimura, T., Watakabe, I., Kobayashi, H., Nagao, Y., Naruse,

K., Higashijima, S.I., Shimizu, T. and Kelsh, R.N., 2023. A gene regulatory network

combining Pax3/7, Sox10 and Mitf generates diverse pigment cell types in medaka and

zebrafish. Development, 150(19).

Minchin, J.E. and Hughes, S.M., 2008. Sequential actions of Pax3 and Pax7 drive xanthophore

development in zebrafish neural crest. Developmental biology, 317(2), pp.508-522.

Morgan, T.H., 1910. Sex limited inheritance in Drosophila. Science, 32(812), pp.120-122.

Nagao, Y., Suzuki, T., Shimizu, A., Kimura, T., Seki, R., Adachi, T., Inoue, C., Omae, Y., Kamei, Y.,

Hara, I. and Taniguchi, Y., 2014. Sox5 functions as a fate switch in medaka pigment cell

development. PLoS genetics, 10(4), p.e1004246.

Nagatsu, T. and Ichinose, H., 1996. GTP cyclohydrolase I gene, tetrahydrobiopterin, and tysorine

hydroxylase gene: Their relations to dystonia and parkinsonism. Neurochemical

research, 21, pp.245-250.

Nasuno, R., Suzuki, S., Oiki, S., Hagiwara, D. and Takagi, H., 2022. Identification and Functional

Analysis of GTP Cyclohydrolase II in Candida glabrata in Response to Nitrosative Stress.

Frontiers in Microbiology, 13, p.825121.

Odenthal, J., Rossnagel, K., Haffter, P., Kelsh, R.N., Vogelsang, E., Brand, M., van Eeden, F.J.,

Furutani-Seiki, M., Granato, M., Hammerschmidt, M. and Heisenberg, C.P., 1996.

Mutations affecting xanthophore pigmentation in the zebrafish, Danio rerio.

Development, 123(1), pp.391-398.

Omasits, U., Ahrens, C.H., Müller, S. and Wollscheid, B., 2014. Protter: interactive protein

feature visualization and integration with experimental proteomic data. Bioinformatics,

30(6), pp.884-886.

Owen, J.P., Kelsh, R.N. and Yates, C.A., 2020. A quantitative modelling approach to zebrafish

pigment pattern formation. *Elife*, *9*, p.e52998.

Parichy, D.M., Ransom, D.G., Paw, B., Zon, L.I. and Johnson, S.L., 2000. An orthologue of the kit-

related gene fms is required for development of neural crest-derived xanthophores and

a subpopulation of adult melanocytes in the zebrafish, Danio rerio. Development,

127(14), pp.3031-3044.

Pelletier, I., Bally-Cuif, L. and Ziegler, I., 2001. Cloning and developmental expression of

zebrafish GTP cyclohydrolase I. Mechanisms of development, 109(1), pp.99-103.

Price, M.N., Dehal, P.S. and Arkin, A.P., 2010. FastTree 2–approximately maximum-likelihood

trees for large alignments. *PloS one*, *5*(3), p.e9490.

Reaume, A.G., Knecht, D.A. and Chovnick, A., 1991. The rosy locus in Drosophila melanogaster:

xanthine dehydrogenase and eye pigments. Genetics, 129(4), pp.1099-1109.

Reissmann, M., Ullrich, E., Bergfeld, U. and Ludwig, A., 2024. Agouti-Signaling Protein and

Melanocortin-1-Receptor Mutations Associated with Coat Color Phenotypes in Fallow

Deer (Dama dama). Genes, 15(8), p.1055.

Ren, J., Kotaka, M., Lockyer, M., Lamb, H.K., Hawkins, A.R. and Stammers, D.K., 2005. GTP

cyclohydrolase II structure and mechanism. Journal of Biological Chemistry, 280(44),

pp.36912-36919.

Richter, G., Ritz, H., Katzenmeier, G., Volk, R., Kohnle, A., Lottspeich, F., Allendorf, D. and

Bacher, A., 1993. Biosynthesis of riboflavin: cloning, sequencing, mapping, and

expression of the gene coding for GTP cyclohydrolase II in Escherichia coli. Journal of

bacteriology, 175(13), pp.4045-4051.

San-Jose, L.M. and Roulin, A., 2017. Genomics of colouration in natural animal populations.

Philosophical Transactions of the Royal Society B: Biological Sciences, 372(1724),

p.20160337.

Sasamori, S., Wiewel, A.S., Thomson, V.A., Kobayashi, M., Nakata, K. and Suzuki, H., 2017.

Potential causative mutation for melanism in rats identified in the agouti signaling

protein gene (Asip) of the Rattus rattus species complex on Okinawa Island,

Japan. *Zoological science*, *34*(6), pp.513-522.

Schluckebier, R., Sachs, M. and Vences, M., 2022. Axanthic green toads, Bufotes viridis (Anura:

Bufonidae), from Cologne, Germany. Herpetology Notes, 15, pp.345-348.

Singh, A.P. and Nüsslein-Volhard, C., 2015. Zebrafish stripes as a model for vertebrate colour

pattern formation. *Current Biology*, *25*(2), pp.R81-R92.

Spradling, A.C. and Rubin, G.M., 1983. The effect of chromosomal position on the expression of

the Drosophila xanthine dehydrogenase gene. Cell, 34(1), pp.47-57.

Steingrímsson, E., Copeland, N.G. and Jenkins, N.A., 2006. Mouse coat colour mutations: from

fancy mice to functional genomics. *Developmental Dynamics: An Official Publication of*

the American Association of Anatomists, *235*(9), pp.2401-2411.

Thöny, B., Auerbach, G. and Blau, N., 2000. Tetrahydrobiopterin biosynthesis, regeneration and

functions. Biochemical Journal, 347(1), pp.1-16.

Tian, Q., Wang, G., Ma, X., Shen, Q., Ding, M., Yang, X., Luo, X., Li, R., Wang, Z., Wang, X. and Fu,

Z., 2022. Riboflavin integrates cellular energetics and cell cycle to regulate maize seed

development. Plant Biotechnology Journal, 20(8), pp.1487-1501.

Trujillano, D., Perez, B., González, J., Tornador, C., Navarrete, R., Escaramis, G., Ossowski, S.,

Armengol, L., Cornejo, V., Desviat, L.R. and Ugarte, M., 2014. Accurate molecular

diagnosis of phenylketonuria and tetrahydrobiopterin-deficient hyperphenylalaninemias

using high-throughput targeted sequencing. European Journal of Human Genetics,

22(4), pp.528-534.*f*

Tzika, A.C., 2024. On the role of TFEC in reptilian colouration. Frontiers in Cell and

Developmental Biology, 12, p.1358828.

Wakamatsu, K. and Ito, S., 2021. Melanins in vertebrates. Pigments, pigment cells and pigment

patterns, pp.45-89.

Watanabe, K., Washio, Y., Fujinami, Y., Aritaki, M., Uji, S. and Suzuki, T., 2008. Adult-type

pigment cells, which colour the ocular sides of flounders at metamorphosis, localize as

precursor cells at the proximal parts of the dorsal and anal fins in early larvae.

Development, growth & differentiation, 50(9), pp.731-741.

Wu, S., Huang, J., Li, Y., Zhao, L. and Liu, Z., 2022. Analysis of yellow mutant rainbow trout

transcriptomes at different developmental stages reveals dynamic regulation of skin

pigmentation genes. Scientific Reports, 12(1), p.256.

Yadav, S., Shinde, R.N., Singh, S., Karthikeyan, S. and Singh, B., 2022. Structurally disordered C-

terminal residues of GTP cyclohydrolase II are essential for its enzymatic activity. Journal

of Biomolecular Structure and Dynamics, 40(19), pp.9318-9331.

Yoshino, H., Nishioka, K., Li, Y., Oji, Y., Oyama, G., Hatano, T., Machida, Y., Shimo, Y., Hayashida,

A., Ikeda, A. and Mogushi, K., 2018. GCH1 mutations in dopa-responsive dystonia and

Parkinson's disease. Journal of neurology, 265, pp.1860-1870.

Ziegler, I. and Harmsen, R., 1970. The biology of pteridines in insects. In Advances in insect

physiology (Vol. 6, pp. 139-203). Academic Press.

Ziegler, I., 2003. The pteridine pathway in zebrafish: regulation and specification during the

determination of neural crest cell-fate. Pigment cell research, 16(3), pp.172-182.

Ziegler, I., McDonaldo, T., Hesslinger, C., Pelletier, I. and Boyle, P., 2000. Development of the

pteridine pathway in the zebrafish, Danio rerio. Journal of Biological Chemistry, 275(25),

pp.18926-18932.

Acknowledgments

599 We thank Douglas Menke and Benjamin Wortman for discussions on gene phylogeny and

600 synteny. This work was supported by an NSERC Discovery Grant and Canada Research Chair to

601 RDHB and APH. We thank Mutation Creation, T Dot Exotics, The Ball Room Canada, Ball Python

602 Genetics Canada, and Designing Morphs for supplying samples.

Data accessibility

605 The data that support the findings of this manuscript have been uploaded to the NCBI

606 Sequence Read Archive (SRA) with the accession PRJNA1122856.

Contributions

609 AGE, APH, and RDHB conceived and designed the study. HR collected samples. AGE and JMA

610 performed the molecular work and AGE the bioinformatics and data analyses. HR performed

611 PCR and Sanger sequencing analysis. AGE wrote the manuscript with input from APH, HR, JMA,

612 and RDHB.

Figure 1. Mendelian phenotypes (colour morphs) in ball pythons. Compared to the (A) wild type, (B) the albino colour morph lacks melanin pigment which allows the display of xanthophore (pterin) pigmentation and patterning, occupying the tan-coloured parts of the wild type. (C) The VPI Axanthic is characterized by a lack of yellow pigmentation on a wildtype pattern. (D) Combining colour morphs with axanthism (VPI Axanthic + Pastel + Fire is shown) can create different forms of the colour anomaly. (E) Some colour morphs like Enchi (Albino + Enchi is shown) are associated with changes in xanthophore migration (e.g., xanthism). (F) Tri-Stripe colour morph exhibits altered melanin patterning. (G) The Albino form of Tri-Stripe shows the effects of this colour morph on xanthophore pigmentation. (H) Melanic colour morphs like Cinnamon do not have xanthophore pigmentation, as shown by an (I) all-white snake in the Albino form. Photo credit: worldofballpythons.com

Figure 2. (A) Manhattan plot of F_{ST} values for SNPs mapped to the chromosome-length assembly. Genomic differentiation between VPI axanthic and a reference pool (Clown colour morph) shows a peak on chromosome two housing the axanthism candidate variant on LOC103050242 (*gch2*). (B) Burmese python amino acid sequence of LOC103050242, with the VPI Axanthic locus indicated by an asterisk (left panel). A nonsense mutation the on the 174th residue (p.Arg174*) likely leads to a defective truncated protein (right panel) and axanthism.