Phenotypic and genetic heterogeneity of *Acinetobacter baumannii* in the course of a chronic infection

Authors: Léa Bednarczuk[a,b], Alexandre Chassard[a#], Julie Plantade[a#], Xavier Charpentier[a] and Maria-Halima Laaberki[a,b*].

a. CIRI, Centre International de Recherche en Infectiologie, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, École Normale Supérieure de Lyon, Univ Lyon, 69007, Lyon, France
b. Université de Lyon, VetAgro Sup, 69280 Marcy l'Etoile, France.

#equivalent contributions
*Correspondence to Maria-Halima Laaberki: maria-halima.laaberki@vetagro-sup.fr

Keywords

Abstract

Acinetobacter baumannii is a nosocomial pathogen associated with various infections, including urinary tract infections (UTIs). In the course of an infection, *A. baumannii* is known to rapidly become resistant to antibiotic therapy, but much less is known about possible adaptation without antibiotic pressure. Through a retrospective study, we investigated within-host genetic diversity during a subclinical five-year UTI in an animal-patient after withdrawal of colistin treatment. We conducted whole-genome sequencing and phenotypic assays on seventeen clonally related isolates from the Sequence Type 25 lineage. Phylogenomic analysis revealed their proximity with animal and human strains from the same country suggesting zoonotic transmission (France). In this case study, the clonally related strains presented variations in genome sizes and nucleotide sequences. Over the course of the infection, *A. baumannii* underwent genome reduction through insertion sequence (IS) recombination, phage excision, or plasmid curing. Alongside this global genome reduction, we observed an expansion of IS*17*, initially located on the endogenous large plasmid. Genetic variations were mainly located in biofilm formation and metabolism genes. We observed repeated variation affecting three biofilm genes and two adhesion operons associated with weak biofilm-forming capacity. Conversely, only two metabolic genes were recurrently affected and phenotypic assays indicated a rather stable metabolism profile between the isolates suggesting minor adaptations to its host. Lastly, an overall decreased antibiotic resistance – expected in the absence of antibiotic treatment - contrasted with a conserved colistin resistance due to a *pmrB* mutation among the isolates.

Impact statement

This study brings a new insight on the genome evolution of the opportunistic pathogen *Acinetobacter baumannii* during a five-year urinary tract infection in the absence of antibiotic treatment. It relies on genomic and phenotypic analyses of multiple isolates from the same animal-patient highlighting the necessity of studying bacterial diversity as opposed to solely on single-isolate approaches. Over the course of the infection, we observed a reduction in genome size, resulting however in a limited loss of metabolism flexibility, but we observed recurrent loss in adhesion capacities and antibiotic resistance. However, this study stands out by the

55 unexpected conserved colistin resistance of the study isolates in the absence of
56 treatment.
57
58

Data summary
The newly sequenced genomes are listed in Table S1, these genomes have been
submitted to the NCBI and their BioProject number is PRJNA1100485. The publicly
available genomes of bacteria from human and animal origin used in this study are
listed in Table S2 All the supplementary tables are available in the Supplementary
Material File.
65
66

Introduction

Acinetobacter baumannii is a non-fermentative Gram-negative pathogenic bacterium responsible for nosocomial infections in humans and animals, particularly in intensive care units (1, 2). *A. baumannii* represents a major threat to public health due to the rapid emergence of multidrug-resistant isolates increasingly difficult to eradicate (3). Indeed, few antibiotics remain effective against this pathogen, such as carbapenems, and the carbapenem resistance of clinical isolates may require poorly tolerated antibiotics such as colistin (4). Emergence of carbapenem resistance in *A. baumannii* therefore limits treatment option and led World Health Organization (WHO) to classify carbapenem-resistant strains of *A. baumannii* (CRAB) as a priority for research and development of new drugs (5). This pathogen is also described in animal infections especially in pets with strains belonging to clones common with human medicine with concerning resistance to human restricted antibiotics such as carbapenems (1, 6).

This bacterium has no particular tropism, causing mainly pulmonary infections but also capable of skin, blood, bones, urinary tract infections (7). The existence of extra-hospital reservoirs has long been disputed, but recent studies demonstrate that *A. baumannii* is an ubiquitous bacterium found both in soils and carried by healthy individuals - humans and animals (8–11). This bacterium is therefore capable of thriving in highly contrasted environments, demonstrating a great capacity for adaptation; antibiotic resistance would be just one facet of its versatility applied to the healthcare context.

Several within-host studies of *A. baumannii* infection describe antibiotic resistance emergence in response to treatment due to non-synonymous single nucleotide polymorphism (SNP) or horizontal gene transfer (12, 13). However, little is known about host adaptation, because brief timeframes of infection limiting observation of bacterial evolution within a host. Additionally, these investigations focused on human infections, typically involving extended antibiotic treatments, making it difficult to decipher bacteria's adaptation to the antibiotic from host adaptation. They also involve mostly single-isolate approaches hindering the understanding of within-host diversity. Consequently, there remains a limited understanding of the fate of antibiotic resistance genes and more broadly of gene evolution post-antibiotic exposure and during host colonization. Seminal within-host microevolution studies on *A. baumannii* observed the accumulation of variations in genes involved surface structures such as

motility, adhesion, biofilm formation or cell wall and capsule biogenesis, suggesting a
consistent selective pressure during antibiotic-treated infections but phenotypic
validation are usually missing (14–16). Phenotypically, antibiotic exposure prompts
bacteria to adopt a biofilm-like lifestyle, enhancing their resistance (17). These
biofilms pose challenges for infection treatment and promote recurrence, notably in
urinary tract infections (UTI) (18). Despite the over-representation of pulmonary and
bloodstream infections in *A. baumannii* studies, the urinary tract is a major source of
A. baumannii isolates in human (19). Tissue tropism may result from mobile genetic
elements (MGE) regulation as exemplified by plasmid-driven regulation of
chromosomal genes modifying strain adhesion pattern (6, 20, 21). Therefore, *A.*
baumannii within-host microevolution may involve crosstalk of MGE and core
genome determinant.
In the following case study, a pet infection emerged from a subcutaneous ureteral
bypass placement which was particularly described as a risk factor for development
of *A. baumannii* UTI in cats (22). We used whole-genome sequencing and phenotype
assays on 17 *A. baumannii* strains isolated from the same cat with UTI over a five-
year period without antibiotic treatment. We sought to investigate the fate of antibiotic
resistance and explore evolutionary strategies upon infection.
118

Materials and methods

Strains
A. baumannii strains identified in the Laboratoire Vétérinaire Départemental du
Rhône, France (LVD69) were stored in an archive since 2014. Nine strains originated
from distinct urine samples collected between 2014 and 2019 and referred to as the
longitudinal strains, while all others were derived from a single urine sample collected
later in October 2019, referred to as the synchronic isolates. Samples were obtained
not for the purpose of this study or research, but rather for diagnostic purposes. The
longitudinal strains were initially isolated on a blood agar plate and the synchronic
strains were isolated using three different media: LB (Lysogeny Broth, Sigma) agar,
Mueller-Hinton agar (Sigma) and CHROMagar *Acinetobacter* (CHROMagar). All the
strains used in this study are listed in Supplemental **Table S1**.

131 Metabolic assays

Metabolic assays were conducted using Biolog Gen III Microplates (Hayward, CA)
following the manufacturer's instructions. In brief, the strains were streaked on LB-
agar and incubated over night at 30°C. For each strain, a single colony was
resuspended in inoculating fluid A (IFA, Biolog) using a cotton-tipped Inoculatorz
swab (Biolog) to reach a cell density of 90-98%. Each well of the Gen III MicroPlate,
containing either a specific carbon source or a chemical sensitivity testing agent,
were filled with 100 µL of the bacterial suspensions and incubated at 33°C for 48
hours in the OmniLog incubator/reader measuring the reduction of tetrazolium redox
dye (forming a purple color) due to bacterial growth.

141 Genome sequencing and annotation

Bacteria were grown in liquid LB medium and DNA was purified using Wizard
genomic DNA kit (Promega). DNA quantifications were performed using Qubit™
dsDNA HS Assay Kit (Thermo Fisher Scientific). For each isolate, Oxford Nanopore
and Illumina DNA sequencings were both performed. Nanopore sequencing were
performed using MinIon flow cells [FLO-MIN106] using the SQK-RBK004 barcoding
kit. Illumina sequencing was performed on Illumina HiSeq 2000 with paired 150-base
sequence reads (Novogene). Each read set was assembled individually using
Unicycler (version 0.4.9, 41) and annotated using Prokka (version 1.14.5, 42) on the
Galaxy pipeline using as reference AB5075-UW protein sequences (GeneBank
accession number: CP008706.1). For analysis purposes, plasmid pF14-11 was
annotated using pA297-3 sequence (GeneBank accession number: KU744946)
except for the *tet*R genes that were named after pAB04 (Genebank accession
number: CP012007, 19, 49–52). Putative prophages were predicted using PHASTER
(27). Putative resistance genes were detected using Resfinder (version 4.5.0, 54)
and insertion sequences using ISfinder (29). All genomes are available at NCBI
under BioProject PRJNA1100485 with quality performed by the NCBI platform using
CheckM (v1.2.2) with completeness between 98.46% and contamination ranging
from 0.27% to 0.53%.⯑⯑

161 Construction of bacterial strains

All the oligonucleotides used in this study for genetic modification are listed in
supplementary **TABLE S3**. Gene disruptions were performed using a scarless
genome editing strategy described previously (30). For plasmid curing of strain F14-
11, its Δ*higA-higB::sacB-aac4* derivative was grown overnight in LB (2mL) at 37°C,
the next day, 100μL were spread on M63 with 10% sucrose for 24h growth at 37°C
then purified on the same medium. About 150 sucrose-resistant CFUs were then
patched on LB agar, LB Tetracycline 30μg/mL, LB Apramycin 30μg/mL. Tetracycline
and apramycin resistant clones were screened by PCR and plasmid extraction for
plasmid loss.

Phylogenetic analysis of the study strains
All chromosomes were aligned to the F14-11 chromosome, set as a reference, using
the Gubbins script generate_ska_alignment.py, which creates an alignment using
split k-mer analysis version 2 (SKA2, version 0.3.7, 58). The resulting alignment
measured 4,061,760 bp with 337 variable positions. The strain 17021 was
additionally included to root the tree. This strain is a suitable root as it lies outside the
other study strains while still being phylogenetically close. Maximum likelihood tree
reconstruction was performed with Gubbins (version 3.2.1, 57) relying on RAxML
using the GTRGAMMA model (version 8.2.12, 56) for constructing the phylogeny in
each iteration to mitigate the bias of horizontal sequence transfer mechanisms.
Statistical support for internal branches of the tree was evaluated by bootstrapping
with 10,000 iterations. Tree visualization was conducted using iTOL (version 6.9, 57).

Phylogenetic analysis with other closely related ST25
Given the close genetic relationships among the study strains, only the first isolated
one, F14-11, was used for the selection of closely related strains from other studies.
The topgenome (-t) feature of WhatsGNU was used to identify the ten closest strains
from a database comprising 4,325 genomes of *A. baumannii* available on GenBank.
The database was constructed by downloading all publicly accessible genomes of *A.*
baumannii from NCBI in February 2023 (Genome database, search criteria:
assembly level = {Chromosome, Complete, Contig}, exclude partial, exclude
anomalous).
To expand the diversity of the strains set, the top genomes of strains from the top 10
closest strains of F14-11 as well as four additional ST25 strains (OIFC143, CI79,

Naval18 and 107m) were also included. To prevent erroneous conclusions regarding
geographic proximity, only one strain per study was included. Incomplete sequenced
genomes were excluded. All the genomes were mapped to the F14-11 chromosome
set as a reference using the Gubbins script generate_ska_alignment.py, which
creates an alignment using SKA2 (version 0.3.7, 58). The resulting alignment
measured 4,061,760 bp with 1,511 variable positions. The output was then analyzed
by the Gubbins algorithm relying on RAxML using the GTRGAMMA model (version
8.2.12, 58) for constructing the phylogeny in each iteration to generate an unbiased
phylogeny. The reproducibility of the node positions in the tree topology was
assessed using bootstrap analysis with 10^4 replicates. No root was assigned
because strains known to be outgroups were genetically too distant. Including such
distant outgroups would result in insufficient sequence alignment and therefore
insufficient information for accurate phylogenetic analysis. Finally, the tree was
visualized, incorporating host and isolation location data, using the R package ggtree
(version 3.12.0, 58).

Variant detection analysis
Primary Illumina sequence reads (fastq) of monophyletic isolates were aligned to the
earliest (F14-11), using the tool snippy (Galaxy version 4.6.0, 59). The primary
Illumina reads of the F14-11 strain were also subjected to the analysis as control and
returned no SNPs. Deletions, insertions and rearrangements were identified using
the software package MAUVE (version 2.4.0, 60). The list of the identified genetic
variations is provided in **TABLE S3**.

Statistical analysis
All statistical analyses were performed using RStudio 4.1.2 (www.rstudio.com). The
medians of the biofilm formation levels of the different strains were compared using
the non-parametric Kruskal-Wallis test, followed by a Bonferroni-Holm correction for
pairwise comparisons with strain F14-11. The potential correlation between
chromosome size and the number of months elapsed between isolation dates was
evaluated using Spearman's correlation test. For months with multiple strains
isolated, the mean chromosome size of those strains was used.

Antibiotic susceptibilities testing

Susceptibility of the strains to a panel of antibiotics was evaluated by disc diffusion
on Mueller-Hinton agar (Biorad, France) following the CA-SFM 2013
recommendations
(https://resapath.anses.fr/resapath_uploadfiles/files/Documents/2013_CASFM.pdf).
Inhibition values were interpreted according to CA-SFM 2013 breakpoints for all
antibiotics but aztreonam, for which *Pseudomonas* spp. breakpoints are given. The
strain *Pseudomonas aeruginosa* CIP 7110 was used as control for all the antibiotic
susceptibility testing. MIC experiments were also conducted to evaluate susceptibility
to colistin and piperacillin according to EUCAST guidelines (37).

Biofilm formation assay
Biofilm-forming capacities of the isolates were evaluated using the crystal violet
staining method, with slight modifications to the protocol previously described (38).
Briefly, exponential phase cells were inoculated into 96-well polystyrene plates in
200µL of BM2 minimal medium (39) supplemented with 10mM potassium glutamate
(BM2G) at starting OD_{600} of 0.1 and incubated at 37°C in the absence of light for 48
hours, without shaking. The medium was replaced with fresh BM2G medium after 24
hours of incubation. After 48 hours, the cells in suspension were removed and the
adherent cells were fixed with 3.7% formaldehyde for 10 minutes. The wells were
then washed twice with distilled water and 200µL of 0.1% crystal violet was added to
each well, incubated for 10 minutes, and washed three times with distilled water. To
solubilize the crystal violet, 200µL of 95% ethanol was added and the absorbance
was measured at 590nm with Infinite M200 PRO (TECAN) and Magellan 7.1 SP1
(TECAN).
248

Results

A ST25 monophyletic infection related to human and animal cases
A retrospective study was performed on *A. baumannii* isolates cultured from a single
animal-patient (cat) over a 5-year period (2014 to 2019). This patient-animal
developed a urinary *A. baumannii* infection following the placement of a urinary
invasive medical device. The early isolate was identified as multidrug resistant,
including to human-restricted antibiotic carbapenems, but not to colistin. However,

two colistin treatments failed to clear the infection. Given the limited antibiotic arsenal
authorized in veterinary medicine, and the absence of invalidating clinical signs, no
further antibiotic treatment was attempted. During medical follow-ups, eight strains
(referred as longitudinal strains) were sequentially isolated and stored by the hospital
laboratory from November 2014 to February 2019 (**FIG S1 & TABLE S1**). In addition,
nine strains (referred to as synchronic strains) were subsequently isolated from a
single complete urine sample collected on October 2019.

FIG 1. Phylogenetic proximity of F14-11 to other strains isolated from human and animal
infections worldwide. (A) Unrooted Maximum Likelihood phylogeny constructed using Gubbins
based on RAxML. Tips are colored by geographic origin, labeled with the country's initial, and
shaped by host type. Bootstrap values are shown at nodes. Scale indicates branch length in base
substitutions. (B) World map with countries highlighted according to the specified color scheme.

Whole-genome sequencing and hybrid assembly of the seventeen study strains
revealed clonally related strains belonging to sequence type (ST) 25 clone, ST
associated with UTI in human and animals (6, 20). To identify the closest genomes to
F14-11, the earliest isolate, we used the Similar Genome Finder utility of the
WhatsGNU tool to query an *A. baumannii* genome database retrieved from the NCBI
(5234 genomes, downloaded on February, 17[th] 2023, 23). Most strains were from
human origin as publicly available *A. baumannii* genomes are mostly related to
human medicine. The phylogenetic tree constructed from twenty-three ST25 strains

reveals an apparent correlation between isolation location and phylogenetic proximity
for Eurasian and American distributions. However, both North American strains were
potentially military imports from overseas explaining their clustering with an Iraqi
strain (41) (**FIG 1**). This analysis also indicated a clustering of clones independently
of their host implying zoonotic transfer and highlighting the relevance of investigating
animal infections for human health (42). The closest human strain, 16A1524, was
indeed isolated in France in 2016. Interestingly, the closest strain, 40288, was also
isolated in France from a dog UTI in 2015 (30). F14-11 and 40288 exhibit only 31
SNPs across their chromosomes and 1 SNP between their ~145Kb-length plasmids
p40288 and pF14-11 (**TABLE S2**). The pF14-11 backbone was also detected in 22
of closest genomes (coverage from 65% to 100%, **TABLE S2**). Despite its reduced
size, pF14-11 shares 100% identity with larger plasmids pVB82 (215Kb) and
pHWBA8 (195Kb) carried by close eponymous strains (**FIG S2** and **TABLE S2**). This
size reduction may explain the inability of pF14-11 to undergo conjugation (**FIG S2**).

FIG 2. Phylogenic tree of the study strains and 40288 associated with the distribution of the genetic variations. (A) Left: Maximum Likelihood Phylogeny constructed with Gubbins based on RAxML. The tree was rooted with strain 17021. Scale indicates branch length in base substitutions. Blue nodes indicate bootstrap values >80. Right: Representation of SNPs and microindels in black, insertions in pink and deletions in green along the genomes in comparison to F14-11 (reference genome). (B) Cumulative distribution of the genetic variations along the F14-11 genome with *y*-axis indicating the number of study strains harboring the genetic variation. The genes most prone to variation are shown.

Genome reduction and insertion sequences dynamics

To further investigate within-host strain diversity, we performed whole-genome comparison of the sixteen study strains to the early one F14-11, set as a reference (**FIG S1** & **TABLE S1**). This analysis revealed a low number of SNPs scattered throughout the genome in comparison to F14-11, with a mean of 31 SNPs per genome ranging from 3 SNPs for the closest genomes to 78 SNPs for the most distant one (**TABLE S1**).

Phylogenetic analysis suggests an absence of correlation between isolation date and phylogenetic relationships (**FIG 2**). Notably, longitudinal isolates F18-02 and F19-02 form an independent clade, referred as clade I, but sharing a hypothetical common ancestor with all other strains. In contrast, late synchronic strains isolated in 2019 cluster with strains isolated in 2015 implying the coexistence of different lineages during the infection. A correlation is observed between the chromosome size of strains and the time elapsed (R^2=0.86, P=6.7×10^{-3}, Spearman test) indicating a reduction in size during infection (**TABLE S1**). Genome reduction operates on MGE such as prophages. Indeed, most strains show independent excisions of one to three of the four prophages identified in strain F14-11. Close strains F18-02 and F19-02 lost one and two prophages respectively (**FIG 2**, large deletions highlighted in green and **FIG S3A**, prophage regions 2 and 4). The endogenous plasmid underwent further size reduction with four of the synchronic isolates lack a 12-18 kb plasmid region corresponding to the BREX phage resistance system while three have lost the plasmid entirely (**FIG 2 & S3B and C**) (43).

Insertion sequences (IS) recombination led to chromosome deletions. The most striking evidence of this phenomenon is the deletion of a 36.6kb region, fixed in a subclade of eight strains within clade II, flanked by to identical IS*Aba34* in F14-11 (**FIG 2** and **FIG S3**). In addition, in comparison to F14-11 chromosome, we observed multiple new chromosomal insertions by an IS originated from the plasmid (**FIG S4**). Indeed, IS*17*, an IS*5* family insertion sequence, was absent from the chromosome of the earliest strain, F14-11, but carried by its plasmid pF14-11. While a single instance of IS*17* was found in the chromosomes of strains isolated in 2015, it significantly multiplied in synchronic strains, climaxing with 75 occurrences in strain F19-10-MH12. Interestingly, IS*17* is also present on p40288 of related 40288 strain although retrained to this replicon.
.

Recurrent variation of adhesion and biofilm genes

Over 82% of SNPs and microindels of the strains studied are in coding sequences, with 80% of non-synonymous SNPs. Several of these genes are involved in adhesion and biofilm formation – genes that are also affected by insertion and deletion events (**FIG 2**; *cup* (44), *csu* (45), *pga* (46), *bapA* (47), *algC* (48) genes). All clade II strains exhibit at least one mutation in one of these genes, while clade I strains retain genetic sequences identical to those of F14-11. Strikingly, genes involved in biogenesis of archaic and classical types of chaperone-usher pili involved in adhesion step of biofilm formation (*csu* and *cup* operon respectively) were subjected to IS insertion, deletion and frameshift. Specifically, the *csu* operon is affected by insertions by distinct IS in strains F15-01A (IS*17*) and F15-06 (IS*Aba33*), this latter insertion being conserved in most of synchronic isolates (7 out of 9). Genes of the classical pili Abp1 (formerly Cup), *cupA* and *cupC*, were also affected by large deletions and indel leading to frameshift in most strains of the clade II. Additionally, genes involved in biofilm formation such as *algC* and *pgaABCD* but also *bapA* gene were also disrupted by IS insertion or subjected to deletion events (**FIG 2**).

FIG 3. Biofilm formation by the study strains and the dog UTI strain 40288 associated with their

phylogenetic relationships. Blue nodes indicate bootstrap values >80. Biofilm formation was

assessed by measuring the optical density of crystal violet stained matrix and compared to

environmental strain DSM30011. Error bars are generated from 12 replicas from 4 independent

experiments. Statistically significant differences with F14-11 (Kruskal-Wallis rank-sum tests with

Bonferroni-Holm correction) are highlighted with asterisks: (*) if the p-value < 0.05, (* *) if the p-

value ≤ 0.01, and (* * *) if the p-value ≤ 0.001.

Hence, during the infection, may co-exist strains with varying adhesion and biofilm

abilities. To bring experimental evidence to this hypothesis, biofilm-forming capacities

of the study strains were measured and compared with those of the environmental

DSM30011 strain described as strong biofilm-producer (**FIG 3**) (39). As observed for

human clinical strains, the animal clinical strain F14-11 forms less biofilm than

DSM30011 (3-fold; p-value: 0.00143). However, F14-11 carries a plasmid closely

related to plasmid pAB5 that downregulates biofilm *pga* and adhesion *cup* genes in

strain UPAB1 (20, 21). We therefore tested the hypothesis that pF14-11 may

restraint biofilm production. Yet, pF14-11 curing did not increase adhesion, biofilm

nor PNAG production (**FIG S5**). We also tested if the residual biofilm capacity of F14-

11 could involve type IV pili or *prp* type I pili but fail to identify its adhesion

mechanism (**FIG S5**).

Surprisingly, F16-05 isolate is a strong biofilm producer despite a frameshift in the

cupC gene and a non-synonymous SNP in *pgaB*. However, its adhesion capacity is

indeed greatly affected (**FIG S5**). Moreover, in spite of the genome proximity between

F14-11, F18-02 and F19-02 these latter two strains produce biofilm amount

comparable to DSM30011 (p-value=1) (**FIG 3**). However, none of the SNPs, mainly

found in intergenic region, nor the phage deletions of these strains are common

between F16-05 and the late isolates and could account for this phenotype. One

hypothesis would be that biofilm production is repressed in F14-11 strain by a

chromosomally encoded factor. In contrast, strains from the clade II exhibit

comparable or reduced biofilm production in comparison to F14-11. However,

considering the numerous adverse variations in their adhesion and biofilm genes, a

more drastic reduction in biofilm formation was expected. In addition, F14-11, despite

identical biofilm and adhesion genes (*cup*, *csu*, *pga*, *algC* genes), produces less

biofilm than dog UTI strain 40288 (p-value: 1.3e-05). These results indicate that the

F14-11 strain isolated at an early stage of infection exhibits low biofilm production,

possibly due to repression of biofilm genes by a chromosomally encoded factor.
These genes were subjected to recurrent mutation in subsequently isolated strains.

Limited conserved metabolism modification
Bacterial metabolism versatility is a crucial factor for successful colonization of a new
environment. Similarly to humans, cat's urine composition is highly variable and
complex with thousands of unique compounds (49, 50). The variations occur both
among different individuals and within the same individual over time, under the
influence of environmental factors, particularly the type of diet (49). Interestingly,
genetic variations in study strains mainly occur in metabolic genes (**FIG S8**).
However, only two metabolic entities appear recurrently affected in the isolated
strains: the *pyrG* gene and the *hca* operon. Most strains (13/17) have a substitution
in *pyrG* gene encoding the putative PyrG enzyme catalyzing amination of UTP to
CTP thereby regulating intracellular CTP levels (**FIG 2**) (51). The non-synonymous
substitution (H193R) being in the vicinity of the conserved putative UTP binding site
we hypothesized that it increases the enzyme affinity for its substrate. We also
observed repeated gene variation (for 14 strains out of 17) for aminobenzoate
utilization (*hca* operon, **FIG 2**). Strains presented either a non-synonymous
substitution (G269V) in the putative membrane transporter HcaK, or a stop codon
mutation in the gene encoding its putative repressor HcaR, indicating a possible
increase in aminobenzoate utilization. This family of compounds can be found in
human and animal food, sometime used as food preservative, excreted in urine (52).
Metabolic capacities may also be affected by deletion events. For instance, the
36.6kb deletion beforementioned, encompasses 27 genes including 19 metabolism
genes whereas only 40.8% of F14-11 genes are of this category (1533/3758). This
conserved large deletion suggests a dispensability of associated metabolic functions
for growth in bladder.

FIG 4. Heatmap of AUC_{48h} for growth either in 70 different nutrient sources, or two pH and three
salinity conditions. Amino acids are highlighted in bold blue. The AUC_{48h} parameter was extracted
from kinetic data over 48 h in Biolog Gen III MicroPlate. The highest and lowest AUC_{48h} are
indicated by green and red colors, respectively. The x-axis lists the tested compounds and
conditions, while the y-axis lists the study strains, which are displayed in the order corresponding
to the phylogenetic tree presented in **Figure 2A**. Mean data of two independent experiments are
shown.

We further explored the metabolism capacities of study strains using Biolog
metabolic phenotypic microarray (**FIG 4**). Overall strains demonstrated a poor ability
to grow in carbohydrates but were able to utilize most amino acids as sole nutrient
source. This shared metabolic profile is advantageous in the context of urinary
infection as urine usually lacks carbohydrates whereas amino acids and small
peptides are abundant (49, 50). The metabolic profiles of the different strains were
mostly similar, with the exception of F19-10-MH12 strain whose metabolic capacities
may have been profoundly affected by numerous IS*17* insertions (**FIG 4 & S4**). For
the set of nutrient sources tested, metabolic capacities were inconsistently altered
and could be explained by allelic variation and transposition. For instance, a P51H
substitution likely abrogated GABA permease function resulting in a growth defect in
GABA in strains F19-10-8 and F19-10-CH6. IS*Aba53* insertion upstream an amino
acid transporter and downstream an arginine *N*-succinyltransferase may account for
slower growth on L-arginine, with ~2-fold lower area under the curve over 48 hours
(AUC_{48h}) of strains F19-10-2 and F19-10-9 compared to F14-11. More surprisingly,
seven strains exhibit a more efficient growth with D-malic acid with up to ~5-fold

higher AUC$_{48h}$ compared to F14-11 (F19-10-2, F19-10-9, F19-10-CH3, F19-10-CH6

and F19-10-CH8). This gain of function for utilization of an artificial amino-acid may

reflect an increase efficiency of enzymes from the tricarboxylic acid (TCA) cycle and

the requirement of this metabolism pathway for growth in the urinary tract as

observed for Uropathogenic *E. coli* (53). An alternative explanation would be an

increased metabolization of D-malic acid that is a food preservative authorized in

animal feed (54). This unexpected gain in function, may therefore reflect adaptation

to the host's urine composition.

Overall resistance loss contrast with colistin resistance fixation

FIG 5. Heatmap of antibiotic susceptibility of the study strains associated with their phylogenetic

relationships. Blue nodes indicate bootstrap values >80.. Abbreviations. AMK: Amikacin, GEN:

Gentamicin, TOB: Tobramycin, CIP: Ciprofloxacin, PIP: Piperacillin, TZP: Piperacillin +

tazobactam, TIC: Ticarcillin, TIM: TIC + clavulanic acid, CAZ: Ceftazidime, FEP: Cefepime, IMP:

Imipenem, MEM: Meropenem, ATM: Aztreonam, TET: Tetracycline, COL: colistin. Antibiotic

susceptibilities were either measured using minimal inhibition concentration (MIC) or by the disk

diffusion methods. The antibiotics tested are listed on the x-axis and the study strains on the y-

axis.

The first isolate F14-11 is resistant to penicillins, cephalosporines (but ceftazidime),

fluoroquinolones, aminoglycosides (but amikacin) and carbapenems. Following

colistin treatments, colistin resistance developed in 2015 isolates (**FIG 5**). Colistin

resistance was mainly maintained after cessation of treatment with the exception of

two sensitive isolates (F18-02 and F19-02). Colistin resistant isolates share a

PmrB(P170T) substitution. We ensured that PmrB(P170T) transfer to F14-11

resulted in colistin resistance (MIC>16µg/mL). Contrarily to PmrB(P170L) associated

to a growth defect in laboratory define medium (55), F14-11 derivatives

PmrB(P170T) and parental F14-11 have comparable growth rate (**FIG S6**). Isolation

of colistin-sensitive strains with wild type *pmrB* sequence suggests coexistence of

both resistant and sensitive strains (**FIG 5** ; F18-02 and F19-02).

In contrast to colistin resistance, several antibiotic resistances were lost in the clade

II in comparison to F14-11. Most susceptibilities emerged due to plasmid loss. Early

isolate plasmid pF14-11 harbors genes conferring resistance to tetracycline (*tetA(B)*),

aminoglycosides (*strA/strB, aac(3)-IIa, aac(6')-Ian*) and sulfonamides (*sul2*)

frequently associated with this family of plasmid (42). Plasmid loss expectedly results

in susceptibility to tetracycline and tobramycin as observed for strains F19-10-MH12,

F19-10-2, and F19-10-9 (**FIG 5**). However, plasmid loss alone may not explain

recovered susceptibility to antibiotics as two tobramycin-sensitive strains still carry a

plasmid (**FIG 2 & 5**; F19-10-CH6 and F19-10-8). Their intact resistance gene

sequences led us to hypothesize that plasmid-specific deletion events encompassing

a putative DNA primase encoding gene may lower plasmid copy number and maybe

decrease associated resistance to tobramycin.

Chromosomal resistance carried by mobile genetic elements were also subjected to

curing. F14-11 strain carries two copies of Tn*2006* carrying *bla*$_{OXA-23}$ gene accounting

for its carbapenem resistance. Closely related strain 40288 also bears two Tn*2006*

copies located either in an AbaR4 resistance island inserted in the *comM* gene or

interrupting the *vgrG* gene (30) . Compared to 40288, F14-11 present a

recombination event between Tn*2006* copies leading to a split AbaR4. Further

deletions of one or two Tn*2006* in several strains may explain increased susceptibility

to carbapenems (**FIG 5** ; respectively in F19-10-4, F19-10-CH3, F19-10-CH8 F19-10-

MH6 and F19-10-MH12). Additionally, resistance decreases without antibiotic gene

loss may be related to compromised growth in laboratory medium associated with

large chromosomal deletion events (**FIG 5 & S7**; F19-10-2, F19-10-9 and F19-10-

MH6). Taken together, these results indicate an overall decline in antibiotic

resistance mediated by curing of mobile genetic elements. They also suggest

coexistence of strains with varying resistance levels within a population hence
illustrating within host heteroresistance in the absence of antibiotic pressure.

Discussion

High versality of certain clones of *A. baumannii* may explain their ability to colonize various host and dominate the global epidemiology of this pathogen. In this study, we examined within-host genomic evolution during prolonged urinary infection of a ST25 strain in an animal host. The genetic proximity between this animal strain and human strains strongly favors the hypothesis of a zoonotic transmission, with a potential transmission from a human to animal hosts (a cat for F14-11 strain and a dog for 40288 strain). A relatedness between human and animal isolates of the ST25 lineage was also suggested by a recent study (42). Remarkably, most animal ST25 strains analyzed in this latter study were isolated from urinary infection in greater extent than for human strains suggesting a host-specific tropism. Following transmission, strains may further evolve by mutations, insertions and deletions presumably mediated by MGE-driven intragenomic rearrangements, particularly through IS dynamics known to promote evolution *in vitro* (56, 57). In our study, the single plasmid copy of IS17 was subsequently detected in the chromosomes of strains collected during infection and extensively propagated in the genomes of late isolates, leading to the pseudogenization of accessory genes. Additionally, recombination of IS*Aba34* led to the loss of a large chromosomal fragment encompassing genes mainly involved in metabolism. While IS element expansion and subsequent recombination events are traditionally associated with the restricted lifestyle of endosymbionts and their dependence on the host, genome reduction have recently been linked with pathogenicity (58). These results bring additional evidence of IS dynamics shaping bacterial evolution to adapt to a new environment, as recently described for the plasmid-mediated IS*1* transposition in clinical Enterobacteriacea (59).

Genetic variations are mainly observed in genes associated with biofilm formation and metabolism. Strikingly, we found that two of three type I *pili* structures encoded by these ST25 isolates were profoundly affected by mutations (*csu* and *cup* genes). Interestingly, several studies described recurrent downregulation or mutation affecting of the *csu* operon following infection by human clinical isolates (15, 55, 60). Besides type I *pili*, other biofilm formation genes present genetic variations such as *algC* and *pga* operon. The *cup* and *pga* genes were also described as downregulated by large conjugative plasmid of a urinary strain involving plasmid-encoded transcriptional regulators that directly or indirectly repressed their expression (20).

For the present study urinary strain, curing the related large plasmid did not modify
adhesion pattern suggesting that a strong repression of these genes may also
involve chromosomal determinants. Repression of biofilm and adhesion gene
expression indicates a dispensability for infection and led to their evolution by
mutation. This hypothesis is strongly sustained by the recurrent and varying types of
gene inactivation observed in those loci. Adhesive structures such as *pili* are highly
immunogenic and are even exploited for vaccine development (61). Loss of these
appendages would favor immune escape.

Furthermore, our data suggest a coexistence of strong and poor biofilm-producers
rather than a homogenous population. This coexistence was also described in other
urinary pathogen during a long-term human urinary infection with *Klebsiella*
pneumoniae (62). These observations may indicate heterogeneous occupation of the
urinary tract with spatially distinct niches. Alternatively, strong and poor biofilm-
producers could collaborate, with sufficient extracellular matrix to allow poor biofilm-
producer strains to embed themselves in the biofilm. The metabolic genes have
undergone the strongest evolution within the host during this long-term infection. This
trend is also observed in longitudinal within-host studies conducted on other bacterial
species indicating common survival strategies (63–66). However, strains display an
unexpected stable metabolism profile. This observation may reflect our technical
limitation as only 70 nutrient sources were tested. Nevertheless, the reduced
genomes may be sufficient to efficiently grow on nutrients available in their host
environment, such as amino acids, and even to adapt to this specific medium with as
near neutral pH, low sodium and chloride, and even presence of diet-associated
compound (D-malic acid metabolism). This latter observation brings new
perspectives to the management of patients suffering from *A. baumannii* infection,
and the potential role of host diet in patient care. Our study also present divergences
with previous studies on human infections. Indeed, we observed only few mutations
in translational machinery genes or in genes related to replication, recombination,
and repair (16, 60). However, these evolutions may have been shaped by alternative
antibiotic treatments performed on human patients. Moreover, contrarily to previous
report, colistin resistance was maintained in the study strains. Snitkin *et al.* isolated
colistin-sensitive strains from human patients after removal of this antibiotic (60).
However, this study used a single-isolate approach randomly isolated that may not
reflect the overall population. Alternatively, the PmrB(P170T) described in the

present study may represent a low-cost mutation or even be beneficial in the context
of urine infection. Indeed, recent findings on PmrAB indicate that this two-component
system responds to low pH (5.8) and metal ions (Fe^{2+}, Zn^{2+} and Al^{3+}) (67). Therefore,
the PmrB(P170T) mutation could have been maintained to mitigate a urine
component toxicity.
Our results also unveiled a hidden phenotypic and genetic diversity within a clinical
sample. Rather than to restrict conclusion to a single isolate with defined
characteristics, the advent of new sequencing technology such as single-cell
sequencing should enable a comprehensive understanding of pathogen genomic
dynamics during infection (68).
566
567

568 **Author statements**
569

570 **Conflicts of interest**
571 The authors declare that there are no conflicts of interest.
572

Funding information
This work was supported by the LABEX ECOFECT (ANR-11-LABX-0048) of
Université de Lyon, within the program "Investissements d'Avenir" (ANR-11-IDEX-
0007) operated by the French National Research Agency (ANR) and by the
RESPOND program of the Université de Lyon (UDL). LB was supported by VetAgro
Sup (PhD funding). AC was supported by Fondation pour la Recherche Médicale
(FRM) (PhD funding).
580
581

Acknowledgements
We thank the HORIGENE group members for useful comments throughout the study.
We are grateful to Michel Pépin and Brigitte Chataigner from Laboratoire Vétérinaire
Départemental (VetAgro Sup) for allowing access to *A. baumannii* animal isolates.

We thank Agnese Lupo and Marisa Haenni (Anses Lyon, Unité Antibiorésistance et

Virulence Bactériennes, France) for performing antimicrobial susceptibility tests and

Maxime Bruto (Anses Lyon, UMR Mycoplasmes, France), Lucie Etienne and Carine

Rey (CIRI, France) for fruitful discussions and comments on phylogenetic analysis.

References

1.   Nocera FP, Attili A-R, De Martino L. 2021. Acinetobacter baumannii: Its
Clinical Significance in Human and Veterinary Medicine. 2. Pathogens 10:127.

2.   van der Kolk JH, Endimiani A, Graubner C, Gerber V, Perreten V. 2019.
Acinetobacter in veterinary medicine, with an emphasis on Acinetobacter baumannii.
J Glob Antimicrob Resist 16:59–71.

3.   Sun R, Song P, Walline J, Wang H, Xu Y, Zhu H, Yu X, Xu J. 2020. Morbidity
and mortality risk factors in emergency department patients with Acinetobacter
baumannii bacteremia. World J Emerg Med 11:164–168.

4.   Iacono M, Villa L, Fortini D, Bordoni R, Imperi F, Bonnal RJP, Sicheritz-Ponten
T, De Bellis G, Visca P, Cassone A, Carattoli A. 2008. Whole-genome
pyrosequencing of an epidemic multidrug-resistant Acinetobacter baumannii strain
belonging to the European clone II group. Antimicrob Agents Chemother 52:2616–
2625.

5.   Tacconelli E. 2018. Discovery, research, and development of new antibiotics:
the WHO priority list of antibiotic-resistant bacteria and tuberculosis. The Lancet
Infectious Diseases 18:318–327.

6.   Lupo A, Châtre P, Ponsin C, Saras E, Boulouis H-J, Keck N, Haenni M, Madec
J-Y. 2016. Clonal Spread of Acinetobacter baumannii Sequence Type 25 Carrying
blaOXA-23 in Companion Animals in France. Antimicrobial Agents and
Chemotherapy 61:10.1128/aac.01881-16.

7.   Maragakis LL, Perl TM. 2008. Acinetobacter baumannii: epidemiology,
antimicrobial resistance, and treatment options. Clin Infect Dis 46:1254–1263.

8.   Bordenave Sylvain, Goñi-Urriza María Soledad, Caumette Pierre, Duran
Robert. 2007. Effects of Heavy Fuel Oil on the Bacterial Community Structure of a
Pristine Microbial Mat. Applied and Environmental Microbiology 73:6089–6097.

9.   Sarma PM, Bhattacharya D, Krishnan S, Lal B. 2004. Assessment of intra-
species diversity among strains of Acinetobacter baumannii isolated from sites
contaminated with petroleum hydrocarbons. Can J Microbiol 50:405–414.

10.   Eveillard M, Kempf M, Belmonte O, Pailhoriès H, Joly-Guillou M-L. 2013.
Reservoirs of Acinetobacter baumannii outside the hospital and potential involvement
in emerging human community-acquired infections. Int J Infect Dis 17:e802-805.

11.   Wilharm G, Skiebe E, Łopińska A, Higgins PG, Weber K, Schaudinn C,
Neugebauer C, Görlitz K, Meimers G, Rizova Y, Blaschke U, Heider C, Cuny C,

Drewes S, Heuser E, Jeske K, Jacob J, Ulrich RG, Bochenski M, Kasprzak M, Burda
E, Ciepliński M, Kaługa I, Jankowiak Ł, Aguirre JI, López-García A, Höfle U, Jagiello
Z, Tobółka M, Janic B, Zieliński P, Kamiński M, Frisch J, Siekiera J, Wendel AF,
Brauner P, Jäckel U, Kaatz M, Müller S, Lübke-Becker A, Wieler LH, Wachsmann J
von, Thrukonda L, Helal M, Epping L, Wolf SA, Semmler T, Jerzak L. 2024. On the
ecology of Acinetobacter baumannii – jet stream rider and opportunist by nature.
bioRxiv https://doi.org/10.1101/2024.01.15.572815.

12.   Hawkey J, Ascher DB, Judd LM, Wick RR, Kostoulias X, Cleland H, Spelman
DW, Padiglione A, Peleg AY, Holt KE. 2018. Evolution of carbapenem resistance in
Acinetobacter baumannii during a prolonged infection. Microb Genom 4:e000165.

13.   Kim M, Park J, Park W. 2021. Genomic and phenotypic analyses of multidrug-
resistant Acinetobacter baumannii NCCP 16007 isolated from a patient with a urinary
tract infection. Virulence 12:150–164.

14.   Zhang J, Xie J, Li H, Wang Z, Yin Y, Wang S, Chen H, Wang Q, Wang H.
2022. Genomic and Phenotypic Evolution of Tigecycline-Resistant Acinetobacter
baumannii in Critically Ill Patients. Microbiol Spectr 10:e0159321.

15.   Wright MS, Iovleva A, Jacobs MR, Bonomo RA, Adams MD. 2016. Genome
dynamics of multidrug-resistant Acinetobacter baumannii during infection and
treatment. Genome Medicine 8:26.

16.   Li H, Zhang J, Wang Z, Yin Y, Gao H, Wang R, Jin L, Wang Q, Zhao C, Wang
Z, Wang H. 2021. Evolution of Acinetobacter baumannii in Clinical Bacteremia
Patients. Infect Drug Resist 14:3553–3562.

17.   Penesyan A, Nagy SS, Kjelleberg S, Gillings MR, Paulsen IT. 2019. Rapid
microevolution of biofilm cells in response to antibiotics. npj Biofilms Microbiomes
5:1–14.

18.   Lila ASA, Rajab AAH, Abdallah MH, Rizvi SMD, Moin A, Khafagy E-S, Tabrez
S, Hegazy WAH. 2023. Biofilm Lifestyle in Recurrent Urinary Tract Infections. 1. Life
13:148.

19.   Venanzio GD, Moon KH, Weber BS, Lopez J, Ly PM, Potter RF, Dantas G,
Feldman MF. 2019. Multidrug-resistant plasmids repress chromosomally encoded
T6SS to enable their dissemination. PNAS 116:1378–1383.

20.   Venanzio GD, Flores-Mireles AL, Calix JJ, Haurat MF, Scott NE, Palmer LD,
Potter RF, Hibbing ME, Friedman L, Wang B, Dantas G, Skaar EP, Hultgren SJ,
Feldman MF. 2019. Urinary tract colonization is enhanced by a plasmid that

regulates uropathogenic Acinetobacter baumannii chromosomal genes. Nature
Communications 10:2763.

21.   Benomar S, Di Venanzio G, Feldman MF. 2021. Plasmid-encoded H-NS
controls extracellular matrix composition in a modern Acinetobacter baumannii
urinary isolate. J Bacteriol JB0027721.

22.   Kopecny L, Palm CA, Drobatz KJ, Balsa IM, Culp WTN. 2019. Risk factors for
positive urine cultures in cats with subcutaneous ureteral bypass and ureteral stents
(2010-2016). J Vet Intern Med 33:178–183.

23.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial
genome assemblies from short and long sequencing reads. PLOS Computational
Biology 13:e1005595.

24.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation.
Bioinformatics 30:2068–2069.

25.   Hamidian M, Ambrose SJ, Hall RM. 2016. A large conjugative Acinetobacter
baumannii plasmid carrying the sul2 sulphonamide and strAB streptomycin
resistance genes. Plasmid 87–88:43–50.

26.   Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, Huse
HK, Zurawski DV, Brittnacher MJ, Manoil C. 2015. Resources for Genetic and
Genomic Analysis of Emerging Pathogen Acinetobacter baumannii. Journal of
Bacteriology 197:2027–2035.

27.   Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016.
PHASTER: a better, faster version of the PHAST phage search tool. Nucleic Acids
Res 44:W16–W21.

28.   Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O,
Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance
genes. J Antimicrob Chemother 67:2640–2644.

29.   Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. 2006. ISfinder: the
reference centre for bacterial insertion sequences. Nucleic Acids Res 34:D32-36.

30.   Godeux A-S, Svedholm E, Barreto S, Potron A, Venner S, Charpentier X,
Laaberki M-H. 2022. Interbacterial Transfer of Carbapenem Resistance and Large
Antibiotic Resistance Islands by Natural Transformation in Pathogenic Acinetobacter.
mBio 13:e02631-21.

31.   Derelle R, Wachsmann J von, Mäklin T, Hellewell J, Russell T, Lalvani A,
Chindelevitch L, Croucher NJ, Harris SR, Lees JA. 2024. Seamless, rapid and

accurate analyses of outbreak genomic data using Split K-mer Analysis (SKA).
bioRxiv https://doi.org/10.1101/2024.03.25.586631.

32.   Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and
post-analysis of large phylogenies. Bioinformatics 30:1312–1313.

33.   Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to
the phylogenetic tree display and annotation tool. Nucleic Acids Research gkae268.

34.   Yu G. 2020. Using ggtree to Visualize Data on Tree-Like Structures. Curr
Protoc Bioinformatics 69:e96.

35.   Seemann T. 2015. snippy: fast bacterial variant calling from NGS reads.

36.   Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve: Multiple Alignment
of Conserved Genomic Sequence With Rearrangements. Genome Res 14:1394–
1403.

37.   Brown D, Cantón R, Dubreuil L, Gatermann S, Giske C, MacGowan A,
Martínez-Martínez L, Mouton J, Skov R, Steinbakk M, Walton C, Heuer O, Struelens
MJ, Högberg LD, Kahlmeter G. 2015. Widespread implementation of EUCAST
breakpoints for antibacterial susceptibility testing in Europe. Eurosurveillance
20:21008.

38.   Tomaras AP, Dorsey CW, Edelmann RE, Actis LAY 2003. Attachment to and
biofilm formation on abiotic surfaces by Acinetobacter baumannii: involvement of a
novel chaperone-usher pili assembly system. Microbiology 149:3473–3484.

39.   Repizo GD, Gagné S, Foucault-Grunenwald M-L, Borges V, Charpentier X,
Limansky AS, Gomes JP, Viale AM, Salcedo SP. 2015. Differential Role of the T6SS
in Acinetobacter baumannii Virulence. PLoS One 10:e0138265.

40.   Moustafa AM, Planet PJ. 2020. WhatsGNU: a tool for identifying proteomic
novelty. Genome Biol 21:58.

41.   Hamidian M, Hall RM. 2016. The resistance gene complement of D4, a
multiply antibiotic-resistant ST25 Acinetobacter baumannii isolate, resides in two
genomic islands and a plasmid. Journal of Antimicrobial Chemotherapy 71:1730–
1732.

42.   Lupo A, Valot B, Saras E, Drapeau A, Robert M, Bour M, Haenni M, Plésiat P,
Madec J-Y, Potron A. 2023. Multiple host colonization and differential expansion of
multidrug-resistant ST25-Acinetobacter baumannii clades. Sci Rep 13:21854.

43.   Rafei R, Koong J, Osman M, Al Atrouni A, Hamze M, Hamidian M. 2022.
Analysis of pCl107 a large plasmid carried by an ST25 Acinetobacter baumannii

strain reveals a complex evolutionary history and links to multiple antibiotic
resistance and metabolic pathways. FEMS Microbes 3:xtac027.

44.   Tamadonfar KO, Di Venanzio G, Pinkner JS, Dodson KW, Kalas V,
Zimmerman MI, Bazan Villicana J, Bowman GR, Feldman MF, Hultgren SJ. 2023.
Structure–function correlates of fibrinogen binding by Acinetobacter adhesins critical
in catheter-associated urinary tract infections. Proceedings of the National Academy
of Sciences 120:e2212694120.

45.   Pakharukova N, Tuittila M, Paavilainen S, Malmi H, Parilova O, Teneberg S,
Knight SD, Zavialov AV. 2018. Structural basis for Acinetobacter baumannii biofilm
formation. PNAS 115:5558–5563.

46.   Choi AHK, Slamti L, Avci FY, Pier GB, Maira-Litrán T. 2009. The pgaABCD
locus of Acinetobacter baumannii encodes the production of poly-beta-1-6-N-
acetylglucosamine, which is critical for biofilm formation. J Bacteriol 191:5953–5963.

47.   Loehfelm TW, Luke NR, Campagnari AA. 2008. Identification and
characterization of an Acinetobacter baumannii biofilm-associated protein. J Bacteriol
190:1036–1044.

48.   Sahu PK, Iyer PS, Barage SH, Sonawane KD, Chopade BA. 2014.
Characterization of the *algC* Gene Expression Pattern in the Multidrug Resistant
Acinetobacter baumannii AIIMS 7 and Correlation with Biofilm Development on
Abiotic Surface. The Scientific World Journal 2014:1–14.

49.   Kim Y, Xu W, Barrs V, Beatty J, Kenéz Á. 2022. In-depth characterisation of
the urine metabolome in cats with and without urinary tract diseases. Metabolomics
18:19.

50.   Bouatra S, Aziat F, Mandal R, Guo AC, Wilson MR, Knox C, Bjorndahl TC,
Krishnamurthy R, Saleem F, Liu P, Dame ZT, Poelzer J, Huynh J, Yallou FS,
Psychogios N, Dong E, Bogumil R, Roehring C, Wishart DS. 2013. The Human Urine
Metabolome. PLOS ONE 8:e73076.

51.   Jørgensen CM, Hammer K, Jensen PR, Martinussen J. 2004. Expression of
the pyrG gene determines the pool sizes of CTP and dCTP in Lactococcus lactis. Eur
J Biochem 271:2438–2445.

52.   Cremin P, Kasim-Karakas S, Waterhouse AL. 2001. LC/ES-MS detection of
hydroxycinnamates in human plasma and urine. J Agric Food Chem 49:1747–1750.

53.   Alteri CJ, Smith SN, Mobley HLT. 2009. Fitness of Escherichia coli during

Urinary Tract Infection Requires Gluconeogenesis and the TCA Cycle. PLOS

Pathogens 5:e1000448.

54.   EFSA Panel on Additives and Products or Substances used in Animal Feed

(FEEDAP). 2014. Scientific Opinion on the safety and efficacy of malic acid and a

mixture of sodium and calcium malate when used as technological additives for all

animal species. EFSA Journal 12:3563.

55.   Pournaras S, Poulou A, Dafopoulou K, Chabane YN, Kristo I, Makris D,

Hardouin J, Cosette P, Tsakris A, Dé E. 2014. Growth Retardation, Reduced

Invasiveness, and Impaired Colistin-Mediated Cell Death Associated with Colistin

Resistance Development in Acinetobacter baumannii. Antimicrobial Agents and

Chemotherapy 58:828–832.

56.   Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM. 2017. The

dynamics of molecular evolution over 60,000 generations. Nature 551:45–50.

57.   Schneider D, Lenski RE. 2004. Dynamics of insertion sequence elements

during experimental evolution of bacteria. Res Microbiol 155:319–327.

58.   Murray GGR, Charlesworth J, Miller EL, Casey MJ, Lloyd CT, Gottschalk M,

Tucker AW (Dan), Welch JJ, Weinert LA. 2021. Genome Reduction Is Associated

with Bacterial Pathogenicity across Different Scales of Temporal and Ecological

Divergence. Molecular Biology and Evolution 38:1570–1579.

59.   Sastre-Dominguez J, DelaFuente J, Toribio-Celestino L, Herencias C,

Herrador-Gómez P, Costas C, Hernández-García M, Cantón R, Rodríguez-Beltrán J,

Santos-Lopez A, Millan AS. 2024. Plasmid-encoded insertion sequences promote

rapid   adaptation   in   clinical   enterobacteria.   bioRxiv

https://doi.org/10.1101/2024.03.01.582297.

60.   Snitkin ES, Zelazny AM, Gupta J, NISC Comparative Sequencing Program,

Palmore TN, Murray PR, Segre JA. 2013. Genomic insights into the fate of colistin

resistance and Acinetobacter baumannii during patient treatment. Genome Res

23:1155–1162.

61.   Hevia A, Delgado S, Sánchez B, Margolles A. 2015. Molecular Players

Involved in the Interaction Between Beneficial Bacteria and the Immune System.

Frontiers in Microbiology 6.

62.   Kalu M, Tan K, Algorri M, Jorth P, Wong-Beringer A. 2022. In-Human

Multiyear Evolution of Carbapenem-Resistant Klebsiella pneumoniae Causing

Chronic Colonization and Intermittent Urinary Tract Infections: A Case Study.
mSphere 7:e0019022.

63.   Chaguza C, Senghore M, Bojang E, Gladstone RA, Lo SW, Tientcheu P-E,
Bancroft RE, Worwui A, Foster-Nyarko E, Ceesay F, Okoi C, McGee L, Klugman KP,
Breiman RF, Barer MR, Adegbola RA, Antonio M, Bentley SD, Kwambana-Adams
BA. 2020. Within-host microevolution of Streptococcus pneumoniae is rapid and
adaptive during natural colonisation. Nat Commun 11:3442.

64.   Gabrielaite M, Johansen HK, Molin S, Nielsen FC, Marvig RL. Gene Loss and
Acquisition in Lineages of Pseudomonas aeruginosa Evolving in Cystic Fibrosis
Patient Airways. mBio 11:e02359-20.

65.   Barker CR, Painset A, Swift C, Jenkins C, Godbole G, Maiden MCJ, Dallman
TJ. 2020. Microevolution of Campylobacter jejuni during long-term infection in an
immunocompromised host. Sci Rep 10:10109.

66.   Ley SD, de Vos M, Van Rie A, Warren RM. 2019. Deciphering Within-Host
Microevolution of Mycobacterium tuberculosis through Whole-Genome Sequencing:
the Phenotypic Impact and Way Forward. Microbiology and Molecular Biology
Reviews 83:10.1128/mmbr.00062-18.

67.   Yamada N, Kamoshida G, Shiraishi T, Yamaguchi D, Matsuoka M, Yamauchi
R, Kanda N, Kamioka R, Takemoto N, Morita Y, Fujimuro M, Yokota S-I, Yahiro K.
2024. PmrAB, the two-component system of Acinetobacter baumannii, controls the
phosphoethanolamine modification of lipooligosaccharide in response to metal ions.
J Bacteriol e0043523.

68.   Lan F, Demaree B, Ahmed N, Abate AR. 2017. Single-cell genome
sequencing at ultra-high-throughput with microfluidic droplet barcoding. Nat
Biotechnol 35:640–646.

819