Epidermal cell fusion promotes the transition from an embryonic to a larval

transcriptome in *C. elegans*

Owen H. Funk, Daniel L. Levy*, David S. Fay*

Department of Molecular Biology, University of Wyoming, Laramie, WY 82071, USA

*Corresponding authors:

David S. Fay   Daniel L. Levy

University of Wyoming   University of Wyoming

Department of Molecular Biology   Department of Molecular Biology

1000 E. University Avenue   1000 E. University Avenue

Laramie, WY 82071   Laramie, WY 82071

Phone: 307-343-0385 (cell)   Phone: 307-417-0354

Fax: 307-766-5098   Fax: 307-766-5098

E-mail: davidfay@uwyo.edu   E-mail: dlevy1@uwyo.edu

Abstract

Cell fusion is a fundamental process in the development of many multicellular

organisms but its precise role in gene regulation and differentiation remains largely

unknown. The *Caenorhabditis elegans* epidermis, which is comprised of multiple

syncytial cells in the adult, represents a powerful model for studying cell fusion in the

context of animal development. The largest of these epidermal syncytia, hyp7,

integrates 139 individual nuclei through processive cell fusion mediated by the

fusogenic protein EFF-1. To explore the role of cell fusion on developmental progression

and associated gene expression changes, we conducted transcriptomic analyses of *eff-*

1 fusion-defective *C. elegans* mutants. Our RNAseq findings showed widespread

transcriptomic changes including the enrichment of epidermal genes and molecular

pathways involved in epidermal function during development. Single-molecule

fluorescence *in situ* hybridization further validated the observed altered expression of

mRNA transcripts. Moreover, bioinformatic analysis suggests that fusion may play a key

role in promoting developmental progression within the epidermis. Our results

underscore the significance of cell–cell fusion in shaping transcriptional programs during

development.

Summary Statement

eff-1 meditated cell fusion drives transcriptomic progression from an embryonic to larval

state in the *C. elegans* epidermis, without which developmental progression is delayed.

Key Words

Cell Fusion, EFF-1, *C. elegans* Development, Epidermis, Syncytia, Transcription

Introduction

Multinucleate syncytial cells are evolutionarily conserved and are found throughout the

tree of life in physiologically healthy tissues and in disease states such as cancer (1-6).

The need for a syncytium varies among organisms and tissues; multiple nuclei may be

required to metabolically support larger cells or to facilitate the rapid production of large

amounts of mRNA for developmental processes or in response to extracellular cues (7-

11). There are multiple paths by which multinucleate syncytia arise, including cell–cell

fusion of multiple mononucleate cells (1, 12-15). Cellular fusion necessarily merges the

cytoplasm of two distinct cells, but how mixing of cytoplasmic factors changes the

regulatory environment of individual nuclei and thus affects the transcriptome within a

newly formed syncytium is poorly understood (16-19).

The *C. elegans* epidermis (also termed hypodermis) is an excellent model to address

questions related to the function of syncytial nuclei. *C. elegans* has a well-characterized

and largely invariant cell lineage in which over a third of somatic nuclei ultimately reside

within syncytial cells (14, 20). The adult worm epidermis is composed of multiple

syncytial cells, the largest of which is hyp7, which surrounds most of the adult body

(14). hyp7 forms through progressive rounds of cell fusion, beginning with 23 embryonic

fusion events that create the initial larval hyp7 prior to hatching. This process is

completely dependent on cell–cell fusion mediated by the fusogen EFF-1 and is

followed by the addition of 116 additional nuclei through EFF-1–mediated fusion events

during larval development (21-23).

eff-1 was initially discovered through genetic screens as the lone factor that is both

necessary and sufficient to induce cell fusion in the *C. elegans* epidermis (22, 23).

Mutations in *eff-1* lead to the disruption of embryonic fusion events in the developing

hypodermal syncytium, leaving the hypodermal progenitors unfused and embryonic cell

boundaries intact. *eff-1* mutant worms exhibit variable morphological and developmental

defects, highlighting the importance of cell fusion for proper body morphogenesis (21,

23-25).

Here we explore the role of embryonic cell fusion in gene expression and differentiation

in the developing epidermal syncytium through RNAseq of worms containing reduction-

of-function mutations in EFF-1, which block most epidermal fusion events in the

embryo. Reduced fusion led to an increase in embryo-associated gene transcripts

concurrent with a decreased expression of larval-associated transcripts in genes

expressed specifically in the epidermis. These findings suggest a developmental delay

in the acquisition of epidermal cell fates and a possible role for cell fusion in activating

gene regulatory programs associated with epidermal differentiation, adding to the

growing list of functions for cell fusion and syncytialization.

Results and Discussion

***eff-1*–mediated cell fusion is essential for normal developmental progression and**

morphology in *C. elegans*

The discovery of EFF-1 as the primary fusogen governing most epidermal cell fusions in

C. elegans originated from classical mutagenesis screens (21-23). This included two

mutations—P183L (*hy21*) and S441L (*oj55*)—that exhibit variable penetrance fusion

defects and temperature sensitivity. Based on available data, both mutations appear to

be partial reduction-of-function mutations rather than complete knockouts and would be

expected to contain background mutations as a result of the mutagenesis, as well as

additional differences in strain backgrounds (23). We therefore generated strains

containing an *eff-1* putative null mutation using CRISPR (P37Stop), which places two

stop codons in exon 1 that is present in all *eff-1* isoforms (**Fig. 1A**). As discussed below,

this precise modification may lead to a stronger reduction of function of *eff-1* as

compared with the historical alleles and allows for a cleaner comparison to our wild type

(WT; N2) fusion-competent control.

Consistent with previous reports, *eff-1*(P37Stop) mutants displayed characteristic

morphological defects, including a bulging anterior region, tail spike abnormalities, and

a dumpy (Dpy) phenotype that persisted into adulthood (**Fig. 1B**) (23). To visualize

fusion defects in *eff-1* mutants we imaged apical junctions using AJM-1::GFP and

confirmed that *eff-1(*P37Stop) mutants have an unfused epidermis that persists into

adulthood (**Fig. 1C,D**). We note that the original characterization of the *eff-1* mutants

reported a complete lack of fusion during late embryonic stages but with partial fusion

occurring during larval development at temperatures below 25°C (23).

We next characterized the developmental timing of *eff-1*(P37Stop) and the historical

mutants from hatching to first molt and to reproductive adulthood. Like *eff-1*(P183L) and

Figure 1. *eff-1* **mutant worms display morphological defects and developmental delay.**
A) *eff-1* gene locus with CRISPR edit P37Stop and mutant alleles P183L (*hy21*) and S441L
(*oj55*) indicated. B) Brightfield imaging of L1 and adult N2 and *eff-1* mutants as indicated. Scale
bars: L1, 10μm; adult, 100μm. C) Schematic of L1 hypodermis with *eff-1*–mediated fusion
events marked with dashed green lines, seam-cells in pink and P cells in red. D) Representative
images of AJM-1::GFP fluorescence, a marker of apical cell–cell junctions in the epidermis, in
N2 (WT) and *eff-1* mutant L1 larvae; lateral seam cells are labeled. E) Quantification of
developmental timepoints in N2 and *eff-1* mutants. Time to first molt was measured via the
cessation of pharyngeal pumping. Time to hatch and adulthood was measured by locomotion
and sexual maturity, respectively. n of > 40 per condition and timepoint.

eff-1(S441L), *eff-1*(P37Stop) exhibited developmental delays in the time to first molt and

the time to reach adulthood, although the delays in *eff-1*(P183L) and *eff-1*(P37Stop)

appeared more pronounced than *eff-1*(S441L) (**Fig. 1E**). Additionally, all three *eff-1*

mutants were less synchronous than WT and we observed a low frequency of larval

arrest in the *eff-1*(P183L) and *eff-1*(P37Stop) mutants.

Disruption of cell fusion in embryonic development leads to consistent,

widespread transcriptomic changes in *eff-1* mutants

The transition from embryo to larva is accompanied by widespread transcriptional

rewiring as the newly hatched L1s begin locomotion and feeding and as many tissues

and cell types acquire their terminal post-mitotic fate (26-29). To examine the role of cell

fusion in the embryonic-to-larval transition, we conducted RNAseq on synchronized

early L1-stage WT, *eff-1*(P37Stop), *eff-1*(S441L), and *eff-1*(P183L) L1 animals. For *eff-*

1(P37Stop), differential expression analysis revealed 2046 upregulated and 912

downregulated differentially expressed genes (DEGs) relative to WT with a false-

discovery rate (FDR) cutoff of <0.05 and $\log_2$(fold change) cutoff of >0.5 or <-0.5 (**Fig.**

2A, File S1). Additionally, we found 3534 upregulated and 2558 downregulated DEGs in

eff-1(S441L) mutants, and 6499 up and 6829 down DEGs in *eff-1*(P183L) mutants (**Fig.**

S1A, File S1). The larger number of DEGs observed in the previously isolated mutant

strains is not entirely unexpected given that these strains were generated by random

mutagenesis and that the parental strains used to generate these alleles may differ

somewhat from our N2 reference strain. As such, some of the transcriptomic changes in

these strains are likely attributable to background variations.

Despite potential background differences, we observed a strong overlap between *eff-*

1(P37Stop), *eff-1*(S441L), and *eff-1*(P183L), with 958 shared DEGs that were

differentially regulated in the same direction (293 downregulated and 665 upregulated)

(**Fig. 2B,C**). Furthermore, $\log_2$-transformed fold-change data for the top DEGs from all

three mutant strains also showed high concordance, with top genes shifting in the same

direction regardless of the *eff-1* mutation (**Fig. 2D**). Euclidian distances between

samples also confirmed transcriptome-wide similarities between *eff-1* point mutations

and P37Stop as compared with WT (**Fig. S1B**). Notably, the P183L transcriptome

appears highly divergent, even more so than the P37Stop despite being only a point

mutation. We suspect this is due to additional background mutations introduced in the

original derivation of this strain, further highlighting the utility of a clean, engineered

knockout.

Taken together, these data indicate a shared core transcriptomic response to the

absence of epidermal cell fusion in L1 larvae, regardless of the specific *eff-1* allele or

Figure 2. RNAseq of multiple *eff-1* mutants indicates a common transcriptome response to fusion disruption.

A) Volcano plot of top DEGs (downregulated in blue, upregulated in red) for *eff-1*(P37Stop). B) Overlap of significant (adjusted p-value < .05, log$_2$FC < -0.5 or > 0.5) DEGs from P37Stop, S441L, and P183L. C) MA (minus average; Bland-Altman) plot of *eff-1*(P37Stop) mutants with upregulated (red) and downregulated (blue) DEGs from S441L and P183L relative to WT. DEGs up- and downregulated in one mutant are more often found differentially expressed in the same direction in the other mutant. Right-side inset shows density of colored (shared DEGs) points D) Heatmap of log$_2$-transformed fold changes (FCs) of top 200 DEGs shared between the *eff-1* mutants.

genetic background. The substantial overlap in both the identity of DEGs and the

directionality of their fold changes across all *eff-1* mutant conditions suggests a

conserved transcriptional signature linked directly to the disruption of epidermal

syncytialization. This highlights a fundamental set of gene expression changes that is

consistently triggered by the failure of cells to fuse in the developing epidermis.

Disruption of *eff-1*–mediated fusion alters epidermal cell expression

The hyp7 syncytium has many functions including the production of a new cuticle at

each molting cycle, which occurs via the synthesis and secretion of apical extracellular

matrix (aECM) proteins that include many collagens (30-34). Consistent with a

disruption of these functions, Gene Ontology (GO) enrichment analysis showed a

substantial number of terms associated with epidermal-specific processes, specifically

extracellular proteins involved in molting and cuticle production (e.g., structural

constituent of cuticle, collagen-containing ECM, etc.) (**Fig. 3A**). These findings support

the notion that normal hypodermal differentiation and function are disrupted or

significantly delayed in *eff-1* mutants. Combined with the tissue enrichment analysis

(**Fig. 3C**), these data suggest that the changes in gene expression observed in fusion-

defective worms occur primarily among genes and pathways associated with the

epidermis and epidermal functions relative to other cell and tissue types. To this point,

intersecting all shared DEGs with genes with detectable expression in the L1 epidermis

found 89.6% of DEGs were present, strengthening the claim that most of genes found

to change their expression are present in the larval epidermis.

In addition to the curated set of GO terms on Wormbase, we noted an enrichment in

embryonic precuticle- or "sheath"-associated genes that accompanied the decrease in

L1-specific genes (**Fig. 3B**). The sheath is the first precuticle formed during embryonic

development prior to secretion of the first cuticle (35-38)**.** The presence of precuticle

gene expression at the L1 stage in *eff-1* mutants suggests a delay in the transition from

precuticle to cuticle gene expression programs. This finding, coupled with the observed

downregulation of L1-specific collagen genes, points toward an overall impairment or

Figure 3. The null *eff-1* mutation affects diverse functions.

A) GO term enrichment using topGO with DEGs from *eff-1*(P37Stop) RNAseq. Colors indicate GO term category as shown in the key. Numbers associated with each bar indicate the observed/expected genes for each GO term displayed. B) Heatmap of z-score–transformed mean expression and log_2(FC) values for selected sheath genes in WT and *eff*-1(P37Stop) L1s. C) Tissue enrichment analysis of DEGs based on gold-standard small-scale enrichment

experiments (37). DEGs were intersected with lists of tissue-specific genes, and p-values were
calculated based on hypergeometric significance. D) Representative images of *col-42* and *rhy-1*
smiFISH hybridization in WT and e*ff-1*(S441L) L1s and 3-fold-stage embryos, respectively.
mRNAs were quantified based on the number of fluorescent puncta. RNAseq counts (transcripts
per million; TPM) for WT and *eff-1* mutants for *col-42* mRNA at L1 larvae and rhy-1 mRNA in
embryos. Developmental trajectory of *col-42* and *rhy-1* expression in WT animals taken from
modENCODE expression data (39). Error bars represent standard error; minimum of nine
animals per condition.

delay in the transcriptional wiring necessary for proper L1 epidermal maturation and

function.

To independently validate and visualize changes in the expression of specific DEGs, we

used single-molecule inexpensive fluorescence *in situ* hybridization (smiFISH) of two

DEGs identified in our RNAseq data, *col-42* and *rhy-1*. *col-42* encodes a structural

collagen largely restricted to larvae and adults and was significantly downregulated in

both *eff-1*(S441L) and *eff-1*(P183L) mutants (**Fig. 3D**). *col-42* is expressed during early

larval development, and smiFISH imaging showed almost no detectable signal in *eff-1*

mutants at L1 (**Fig. 3D**). Conversely, *rhy-1* is a transmembrane acyltransferase that is

expressed in the embryonic epidermis (40, 41). Consistent with the transcriptomic data,

the number of *rhy-1* mRNA fluorescent puncta was also elevated in *eff-1* mutants (**Fig.**

3D).

Although large changes in the transcriptomes of *eff-1* mutants were evident, such

changes may be due to direct or indirect effects caused by the disruption of epidermal

cell functions. To distinguish between these possibilities, we looked for overlap between

gold-standard lists of tissue-specific genes from small-scale single-tissue experiments

and the *eff-1* DEGs (42). This analysis showed a significant enrichment among

downregulated DEGs for transcripts associated with the hyp7 syncytium, hypodermis,

and epithelial system, over other non-syncytial tissues (**Fig. 3C**), consistent with a

model in which unfused hyp7 progenitors fail to upregulate genes found in the L1

syncytial hypodermis. This suggests that the observed transcriptomic changes are, in

large part, the direct consequence of an unfused epidermis rather than of a systemic

developmental delay. Collectively, our data support the central finding that fusion plays a

critical role in promoting the timely differentiation and function of the epidermis based on

hallmark changes in gene expression.

***eff-1* mutant transcriptional changes are consistent with a developmental delay**

In addition to shared pathways and functions associated with the DEGs isolated across

eff-1 mutations, we found a striking enrichment in the DEGs associated with the

transition from embryonic to larval development. By examining the developmental

trajectory of DEGs at timepoints throughout the WT worm lifecycle from modENCODE

(39, 43), we found an enrichment in L1-associated genes in the downregulated DEG

set, along with upregulated DEGs showing substantial enrichment for genes normally

expressed in the late embryo (**Fig. 4A, D**). Furthermore, genes that were not

differentially expressed (FDR > 0.05) in the *eff-1* mutants were predominantly those for

which expression levels remain relatively stable across normal development.

Collectively, these observations strongly suggest that the normal transcriptional shift

from late-embryonic to L1 larval gene expression programs is substantially delayed in

worms deficient for EFF-1–mediated fusion.

To further investigate the hypothesis that *eff-1* mutant L1s inappropriately maintain high

expression of embryonic genes while failing to upregulate larval-specific genes, we

performed a linear regression analysis. This analysis used WT expression data across

multiple embryonic and larval timepoints to define gene sets that are developmentally

upregulated or downregulated during the transition from late-embryonic to larval stages.

We then compared the average fold changes of these developmentally dynamic gene

sets to *eff-1*(P37Stop) mutants. This comparison revealed a net increase in the

expression of developmentally downregulated (i.e., typically embryonically enriched)

genes and a corresponding decrease in the expression of developmentally upregulated

(i.e., typically larvally enriched) genes, irrespective of whether individual genes met the

strict criteria for differential expression (**Fig. 4B**). This broader analysis reinforces the

conclusion that the absence of cell fusion leads to a significant delay in the acquisition

of the appropriate L1 transcriptome.

Figure 4. Fusion mutants fail to acquire the L1 epidermal transcriptome.
Normal (WT) developmental trajectories of genes that were upregulated (red) or downregulated (blue) in *eff-1* mutants. Shaded region denotes 95% confidence interval. Fusion-defective transcriptomes show a marked increase in embryonic transcripts with a concurrent downshift in larval-associated transcripts. Dashed line indicates an approximate hatching time of 800 min. B) Genes were binned using a linear regression of modENCODE data to determine transcripts that went up (green), down (purple) or stayed consistent (gray) across normal WT development. Average log$_2$–transformed fold change data of *eff-1*(P37Stop) RNAseq data for each category of genes are shown. Error bars represent standard error, and p-values calculated using Mann-Whitney U test, ****P < .00001. C) Heatmap of z-score normalized DEG expression for genes identified as ELT-3 regulated from the TF2DNA dataset (49). D) Schematic of WT and *eff-1* mutant gene expression profiles across development, showing a lack of transition from the embryonic to larval transcriptome in fusion defective *eff-1* mutants.

The transition from an embryonic to larval transcriptome is facilitated by many transcription factors, including ELT-3, an embryonically enriched GATA transcription factor with roles in the transcriptional activation of epidermis-specific genes and developmental pathways (44-46). Previously, an *elt-3* reporter construct was found to be expressed in only a subset of hypodermal precursor cells, and thus it has been used as a reporter for fusion, or lack thereof, in *eff-1* mutant worms (47). Intriguingly, we found a

large swath of ELT-3 regulated genes identified through the TF2DNA database as

significantly downregulated in *eff-1* mutant worms (**Fig. 4C**). Examination of the ELT-3

chromatin immunoprecipitation sequencing (ChIPseq) signal across all up, down, and

unchanged genes also showed an enrichment of downregulated genes (**Fig. S2**). The

observed downregulation of ELT-3 target genes in *eff-1* mutants provides a potential

mechanistic link between the failure of cell fusion and the observed developmental

delay. Namely, cell fusion may be necessary for the appropriate stage-specific activity of

key developmental regulators such as ELT-3, which may be expressed in only a subset

of cells contributing to epidermal syncytia.

Taken together, our work underscores the critical role of EFF-1–mediated cell fusion in

the developmental progression of *C. elegans* and highlights its importance for the timely

activation of the L1 larval transcriptome. Our findings suggest that the process of fusion

itself may be a key mechanistic trigger promoting the shift from the embryonic to the L1

transcriptional landscape in fusion-driven syncytia such as hyp7. Although these results

highlight the importance of syncytial formation for the proper expression of

developmental genes, precisely how fusion orchestrates transcriptional maturation

remains an open question. Future investigations will be needed to unravel the molecular

mechanisms involved and to identify specific factors that may drive the identity and

transcriptional programs of developing syncytial nuclei.

Materials and Methods

Strains used and maintenance

C. elegans strains were maintained according to standard protocols and were

propagated at 22°C. The two *eff-1* mutant strains used here were WH171 (*eff-1*(*oj55*);

jcIs1 *[ajm-1::GFP + unc-29(+) + rol-6(su1006)]*) and BP76 (*eff-1*(*hy21*); jcIs1 *[ajm-*

1::GFP + unc-29(+) + rol-6(su1006)]). MH1384 (*[kuIs46 (ajm-1::GFP; unc-119(+))]*) was

used to compare WT AJM-1::GFP expression to that in e*ff-1* mutants. The P37Stop

strain as generated using CRISPR via the method described in Ghanta et. al 2021 (48).

Briefly, RNPs loaded with gRNA (GGTGTCTTGGAACAGTGTGG) were injected along

with repair template designed to add two stop codons along with an SpeI recognition

site into the gonads of animals expressing *kuls46*[AJM-1::gfp] and *bqSi640*[*dpy-*

7p::FRT::mCherry::*his-58*::FRT::GFP::*his-58*].

Developmental timing analysis

Time to hatch was determined by plating young adult worms, allowing them to lay eggs

for 2 hours, and then removing the adults and measuring the percentage of unhatched

eggs that remained over time. Time to molt was determined by plating synchronized L1

larvae and watching for cessation of pharyngeal pumping to mark the beginning of the

L1–L2 molt. Time to adulthood was measured by plating synchronized L1 larvae and

noting the presence of embryos in the gonads. All experiments were carried out at 22°C.

Microscopy

354 Confocal images for **Fig. 1** were taken using CellSens on an Olympus Spin-SR,

355 Spinning Disc, Super Resolution System using an inverted Olympus IX83 microscope

356 and Yokogawa W1 spinning disc. smiFISH images were taken of worms fixed in

357 methanol/acetone as described (49). Larvae for **Fig. 1** images were prepared in 10 mM

358 levamisole on 3% agarose pads.

RNA isolation and RNAseq

361 L1s were synchronized by bleaching gravid adults and allowing embryos to arrest in M9

362 buffer. L1s were spun down and lysed for RNA collection via the Direct-zol kit from

363 Zymo. RNA quality was confirmed using an Agilent TapeStation with minimum RINs

364 (RNA Integrity Number) of 8.0. mRNA libraries were constructed using TruSeq Library

365 Prep Kits from Illumina and were sequenced on an Illumina Nextseq P3 flow cell.

Transcriptomic analysis

Paired-end reads were processed as follows. Low-quality reads were filtered using

fastp, and their quality was assessed using fastQC (0.12.1). Reads were aligned using

STAR and pseudoaligned using Salmon to the *ce11* reference genome, duplicates were

removed using picard, and counts were collected using htseq (50-52). Differential

expression was determined using DESeq2 (v. 1.42.1), with counts filtered for >1 TPM

across all samples . DEGs were defined as genes with a Benjamini-Hochberg adjusted

$p < 0.05$ and $\log_2$(fold change) > 0.5 or < -0.5 , **(**processed files in **File S1)**. Heatmaps

were generated using the seaborn clustermap() function using Euclidian distance to

obtain hierarchal linkages and z-score normalization by rows. Intersectional analysis

was done by collecting modENCODE developmental expression data, processed in

Hutter and Soh 2016 (39, 43). Upregulated, downregulated, and unchanged

classifications were determined by performing a linear regression for expression data

across developmental time points and collecting genes with a derived slope of >1

(developmentally upregulated) or <-1 (developmentally downregulated). Genes with a

derived slope from 1 to -1 were categorized as unchanged. Raw and processed data

files can be found through GEO accession: (GSE301850).

GO term analysis

GO term enrichment was carried out using topGO (v. 2.54.0) on ontologies from

WormBase using the "weight01" algorithm, and significance was assessed with a

Fisher's statistic. Genes found to be differentially expressed (FDR < 0.05) in both S441L

and P183L were used as the test set, and the background gene set was specified as

genes with a base mean TPM of >1 in all three transcriptomes.

Tissue enrichment

Gold-standard tissue-specific gene sets were obtained from https://worm.princeton.edu/

(42) and compared to DEGs from the *eff-1* mutant RNAseq data. Fischer's exact test

was used to calculate enrichment significance.

smiFISH

smiFISH was carried out as described (49). Probe sets were designed against *col-42*

and *rhy-1* using Oligostan with FLAP-X extensions (53). The FLAP-X secondary probes

were labeled with Quasar 670 from Biosearch Technologies. Quantification was done in

FIJI by manually drawing designated regions of interest around embryos and L1s of

appropriate age and sufficient fixation, thresholding across experimental replicates, and

using the "analyze particles" option.

ELT-3 ChIPseq analysis

ELT-3–associated genes were obtained from the TF2DNA dataset (54) and compared to our experimentally derived DEGs. L1 ELT-3 ChIPseq data from modENCODE was downloaded as read-depth–normalized bigwig files. Heatmaps were generated with *deeptools* (v 3.5.6) (55). Up- and downregulated gene sets were designated from the *eff-1* P37Stop dataset with the same criteria (adjusted p-value < .05, log2 fold change > 0.5 or < -0.5. Unchanged genes were taken as a random subset of 5000 genes with adjusted p-value > .2.

Acknowledgements

This work was supported by the National Institutes of Health grants GM136236 and GM125091 (to D.S.F.), GM134885 (to D.L.L.). This project was also supported by an Institutional Development Award (IDeA) from the National Institute of General Medical Sciences of the National Institutes of Health under P20GM103432. We thank Amy Fluet for editing this manuscript, the Nishimura lab for guidance and implementation of smiFISH, the Sundaram and Murray labs for insightful conversation and feedback on the results.

Author contributions

Conceptualization, OHF, DLL, DSF; Investigation, OHF; Writing – Original Draft, OHF; Writing – Review & Editing, OHF, DLL, DSF; Funding Acquisition, DLL, DSF; Supervision, DLL, DSF.

Figure S1. RNAseq of *eff-1* mutants shows substantial overlap in gene expression changes.

A) MA plots of *eff-1*(P183L) and *eff-1*(S441L) with DEGs (adjusted p < 0.05) in red for upregulated and blue for downregulated. B) Euclidian distance matrix of regularized log transformed RNAseq counts of *eff-1* mutants and WT controls from multiple sequencing runs.

Figure S2. ELT-3 ChIPseq signal across *eff-1* differentially expressed genes.
L1 ELT-3 ChIPseq profile across DEGs from *eff-1*(P37Stop). ELT-3 ChIPseq signal was derived from modENCODE data and plotted for a region spanning 2 kb upstream and 500 bp downstream from the transcription start site (TSS) of upregulated, downregulated, and unchanged genes.

References

References

1. B. M. Ogle, M. Cascalho, J. L. Platt, Biological implications of cell fusion. *Nat Rev*
Mol Cell Biol **6**, 567-575 (2005).
2. S. J. Calvert *et al.*, Studies of the dynamics of nuclear clustering in human
syncytiotrophoblast. *Reproduction* **151**, 657-671 (2016).
3. P. J. Brooks, M. Glogauer, C. A. McCulloch, An Overview of the Derivation and
Function of Multinucleated Giant Cells and Their Role in Pathologic Processes. *Am J*
Pathol **189**, 1145-1158 (2019).
4. L. I. Larsson, B. Bjerregaard, J. F. Talts, Cell fusions in mammals. *Histochem Cell*
Biol **129**, 551-561 (2008).
5. S. M. Abmayr, G. K. Pavlath, Myoblast fusion: lessons from flies and mice.
Development **139**, 641-656 (2012).
6. M. Roper, C. Ellison, J. W. Taylor, N. L. Glass, Nuclear and genome dynamics in
multinucleate ascomycete fungi. *Curr Biol* **21**, R786-793 (2011).
7. J. J. Dowling, C. C. Weihl, M. J. Spencer, Molecular and cellular basis of genetically
inherited skeletal muscle disorders. *Nat Rev Mol Cell Biol* **22**, 713-732 (2021).
8. M. Cardamone, B. T. Darras, M. M. Ryan, Inherited myopathies and muscular
dystrophies. *Semin Neurol* **28**, 250-259 (2008).
9. Y. Iosilevskii, B. Podbilewicz, Programmed cell fusion in development and
homeostasis. *Curr Top Dev Biol* **144**, 215-244 (2021).
10. T. L. Orr-Weaver, When bigger is better: the role of polyploidy in organogenesis.
Trends Genet **31**, 307-315 (2015).
11. O. Deshpande, J. de-Carvalho, D. V. Vieira, I. A. Telley, Astral microtubule cross-
linking safeguards uniform nuclear distribution in the Drosophila syncytium. *J Cell*
Biol **221** (2022).
12. S. Demin, M. Berdieva, A. Goodkov, Cell-cell fusions and cell-in-cell phenomena in
healthy cells and cancer: Lessons from protists and invertebrates. *Semin Cancer*
Biol **81**, 96-105 (2022).
13. E. H. Chen, E. Grote, W. Mohler, A. Vignery, Cell-cell fusion. *FEBS Lett* **581**, 2181-
2193 (2007).
14. S. Alper, B. Podbilewicz, Cell fusion in Caenorhabditis elegans. *Methods Mol Biol*
475, 53-74 (2008).
15. P. S. Aguilar *et al.*, Genetic basis of cell-cell fusion mechanisms. *Trends Genet* **29**,
427-437 (2013).
16. M. Kim *et al.*, Single-nucleus transcriptomics reveals functional
compartmentalization in syncytial skeletal muscle cells. *Nat Commun* **11**, 6375
(2020).
17. M. J. Petrany *et al.*, Single-nucleus RNA-seq identifies transcriptional heterogeneity
in multinucleated skeletal myofibers. *Nat Commun* **11**, 6374 (2020).
18. K. Williams, K. Yokomori, A. Mortazavi, Heterogeneous Skeletal Muscle Cell and
Nucleus Populations Identified by Single-Cell and Single-Nucleus Resolution
Transcriptome Assays. *Front Genet* **13**, 835099 (2022).

19.   X. Liu *et al.*, Analysis of cell fate from single-cell gene expression profiles in C.
elegans. *Cell* **139**, 623-633 (2009).

20.   B. Podbilewicz, J. G. White, Cell fusions in the developing epithelial of C. elegans.
Dev Biol **161**, 408-424 (1994).

21.   B. Podbilewicz *et al.*, The C. elegans developmental fusogen EFF-1 mediates
homotypic fusion in heterologous cells and in vivo. *Dev Cell* **11**, 471-481 (2006).

22.   G. Shemer *et al.*, EFF-1 is sufficient to initiate and execute tissue-specific cell fusion
in C. elegans. *Curr Biol* **14**, 1587-1591 (2004).

23.   W. A. Mohler *et al.*, The type I membrane protein EFF-1 is essential for
developmental cell fusion. *Dev Cell* **2**, 355-362 (2002).

24.   S. L. Koneru *et al.*, A role for the fusogen eff-1 in epidermal stem cell number
robustness in Caenorhabditis elegans. *Sci Rep* **11**, 9787 (2021).

25.   C. Brabin, P. J. Appleford, A. Woollard, The Caenorhabditis elegans GATA factor ELT-1
works through the cell proliferation regulator BRO-1 and the Fusogen EFF-1 to
maintain the seam stem-like fate. *PLoS Genet* **7**, e1002200 (2011).

26.   J. I. Murray *et al.*, Automated analysis of embryonic gene expression with cellular
resolution in C. elegans. *Nat Methods* **5**, 703-709 (2008).

27.   M. E. Boeck *et al.*, The time-resolved transcriptome of C. elegans. *Genome Res* **26**,
1441-1450 (2016).

28.   J. Cao *et al.*, Comprehensive single-cell transcriptional profiling of a multicellular
organism. *Science* **357**, 661-667 (2017).

29.   J. S. Packer *et al.*, A lineage-resolved molecular atlas of. *Science* **365** (2019).

30.   G. J. Hendriks, D. Gaidatzis, F. Aeschimann, H. Großhans, Extensive oscillatory gene
expression during C. elegans larval development. *Mol Cell* **53**, 380-392 (2014).

31.   M. W. Meeuse *et al.*, Developmental function and state transitions of a gene
expression oscillator in Caenorhabditis elegans. *Mol Syst Biol* **16**, e9498 (2020).

32.   A. D. Chisholm, T. I. Hsiao, The Caenorhabditis elegans epidermis as a model skin. I:
development, patterning, and growth. *Wiley Interdiscip Rev Dev Biol* **1**, 861-878
(2012).

33.   I. L. Johnstone, Cuticle collagen genes. Expression in Caenorhabditis elegans.
Trends Genet **16**, 21-27 (2000).

34.   A. P. Page, I. L. Johnstone, The cuticle. *WormBook* 10.1895/wormbook.1.138.1, 1-15
(2007).

35.   M. Costa, B. W. Draper, J. R. Priess, The role of actin filaments in patterning the
Caenorhabditis elegans cuticle. *Dev Biol* **184**, 373-384 (1997).

36.   J. R. Priess, D. I. Hirsh, Caenorhabditis elegans morphogenesis: the role of the
cytoskeleton in elongation of the embryo. *Dev Biol* **117**, 156-173 (1986).

37.   V. P. Mancuso *et al.*, Extracellular leucine-rich repeat proteins are required to
organize the apical extracellular matrix and maintain epithelial junction integrity in
C. elegans. *Development* **139**, 979-990 (2012).

38.   M. V. Sundaram, N. Pujol, The Caenorhabditis elegans cuticle and precuticle: a
model for studying dynamic apical extracellular matrices in vivo. *Genetics* **227**
(2024).

39.   M. B. Gerstein *et al.*, Integrative analysis of the Caenorhabditis elegans genome by
the modENCODE project. *Science* **330**, 1775-1787 (2010).

40.   C. Shen, Z. Shao, J. A. Powell-Coffman, The Caenorhabditis elegans rhy-1 gene
inhibits HIF-1 hypoxia-inducible factor activity in a negative feedback loop that does
not include vhl-1. *Genetics* **174**, 1205-1214 (2006).

41.   D. K. Ma, R. Vozdek, N. Bhatla, H. R. Horvitz, CYSL-1 interacts with the O2-sensing
hydroxylase EGL-9 to promote H2S-modulated hypoxia-induced behavioral
plasticity in C. elegans. *Neuron* **73**, 925-940 (2012).

42.   R. Kaletsky *et al.*, Transcriptome analysis of adult Caenorhabditis elegans cells
reveals tissue-specific gene and isoform expression. *PLoS Genet* **14**, e1007559
(2018).

43.   H. Hutter, J. Suh, GExplore 1.4: An expanded web interface for queries on. *Worm* **5**,
e1234659 (2016).

44.   J. S. Gilleard, J. D. McGhee, Activation of hypodermal differentiation in the
Caenorhabditis elegans embryo by GATA transcription factors ELT-1 and ELT-3. *Mol*
Cell Biol **21**, 2533-2544 (2001).

45.   G. Broitman-Maduro, S. Sun, T. Kikuchi, M. F. Maduro, The GATA factor ELT-3
specifies endoderm in Caenorhabditis angaria in an ancestral gene network.
Development **149** (2022).

46.   J. S. Gilleard, Y. Shafi, J. D. Barry, J. D. McGhee, ELT-3: A Caenorhabditis elegans
GATA factor expressed in the embryonic epidermis during morphogenesis. *Dev Biol*
208, 265-280 (1999).

47.   J. H. Shinn-Thomas, J. J. del Campo, J. Wang, W. A. Mohler, The EFF-1A Cytoplasmic
Domain Influences Hypodermal Cell Fusions in C. elegans But Is Not Dependent on
14-3-3 Proteins. *PLoS One* **11**, e0146874 (2016).

48.   K. S. Ghanta, T. Ishidate, C. C. Mello, Microinjection for precision genome editing in.
STAR Protoc **2**, 100748 (2021).

49.   D. M. Parker, L. P. Winkenbach, A. Parker, S. Boyson, E. O. Nishimura, Improved
Methods for Single-Molecule Fluorescence In Situ Hybridization and
Immunofluorescence in Caenorhabditis elegans Embryos. *Curr Protoc* **1**, e299
(2021).

50.   A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
(2013).

51.   R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and
bias-aware quantification of transcript expression. *Nat Methods* **14**, 417-419 (2017).

52.   S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).

53.   N. Tsanov *et al.*, smiFISH and FISH-quant - a flexible single RNA detection approach
with super-resolution capability. *Nucleic Acids Res* **44**, e165 (2016).

54.   M. Pujato, F. Kieken, A. A. Skiles, N. Tapinos, A. Fiser, Prediction of DNA binding
motifs from 3D models of transcription factors; identifying TLX3 regulated genes.
Nucleic Acids Res **42**, 13500-13512 (2014).

55.   F. Ramírez *et al.*, deepTools2: a next generation web server for deep-sequencing
data analysis. *Nucleic Acids Res* **44**, W160-165 (2016).