Cyclic di-GMP as an Antitoxin Regulates Bacterial Genome Stability and Antibiotic Persistence in Biofilms

Hebin Liao[1,2,6,7], Xiaodan Yan[1,2,7], Chenyi Wang[1,2], Chun Huang[1,2], Wei Zhang[1,2], Leyi Xiao[1,2], Jun Jiang[5], Yongjia Bao[5], Tao Huang[5], Hanbo Zhang[5], Chunming Guo[5], Yufeng Zhang[1,2,4], Yingying Pu[1,2,3,*]

[1]The State Key Laboratory Breeding Base of Basic Science of Stomatology & Key Laboratory of Oral Biomedicine Ministry of Education, School & Hospital of Stomatology, Medical Research Institute, Wuhan University, Wuhan, 430079, China.

[2]Frontier Science Center for Immunology and Metabolism, Wuhan University, Wuhan, 430079, China.

[3]Department of Immunology, Hubei Province Key Laboratory of Allergy and Immunology, State Key Laboratory of Virology and Medical Research Institute, Wuhan University School of Basic Medical Sciences, Wuhan, 430079, China.

[4]Taikang Center for Life and Medical Sciences, Wuhan University, Wuhan, 430079, China.

[5]Center for Life Sciences, School of Life Sciences, Yunnan University, Kunming, 650500, China.

[6]Translational Medicine Research Center, North Sichuan Medical College, Nanchong, 637000, China.

[7]These authors contributed equally: Hebin Liao, Xiaodan Yan

[*]To whom correspondence should be addressed. Email: yingyingpu@whu.edu.cn

Abstract

Biofilms are complex bacterial communities characterized by a high persister prevalence, which contributes to chronic and relapsing infections. Historically, biofilm persister formation has been linked to constraints imposed by their dense structures. However, we observed an elevated persister frequency accompanying the stage of cell adhesion, marking the onset of biofilm development. Subsequent mechanistic studies uncovered a distinctive type of toxin-antitoxin (TA) module triggered by cell adhesion, which is responsible for this elevation. In this module, the toxin HipH acts as a genotoxic deoxyribonuclease, inducing DNA double strand breaks and genome instability. While the second messenger c-di-GMP functions as the antitoxin, exerting control over HipH expression and activity. The dynamic interplay between c-di-GMP and HipH levels emerges as a crucial determinant governing genome stability and persister generation within biofilms. These findings unveil a unique TA system, where small molecules act as the antitoxin, outlining a biofilm-specific molecular mechanism influencing genome stability and antibiotic persistence, with potential implications for treating biofilm infections.

Keywords

Biofilm/ cyclic di-GMP/ HipH/ Persister/ Toxin-antitoxin system

Introduction

The prevalence of chronic and recurrent microbial infections has been significantly attributed to the formation of biofilms - complex, surface-attached bacterial communities (Costerton et al, 1999; Hall-Stoodley et al, 2004; Lewis, 2005). In biofilms, bacteria exhibit reduced antibiotic sensitivities, a phenomenon occurring independently of the presence of resistance determinants on genetic elements (Ceri et al, 1999). This antibiotic insensitivity is linked to the elevated prevalence of persister cells in biofilms, with rates ranging from 10 to 1000 times higher compared to their planktonic counterparts (Lewis, 2005; Yan & Bassler, 2019). Persister cells, recognized as phenotypic variants in an isogenic population, demonstrate a remarkable ability to withstand antibiotic treatment and, intriguingly, can resuscitate once the antibiotic pressure is alleviated (Balaban et al, 2004; Bigger, 1944; Lewis, 2007; Zhou et al, 2023). These characteristics collectively position persister cells as key contributors to the perpetuation of persistent infections of biofilms. Moreover, while persister cells themselves may not carry genetic mutations associated with resistance, their survival mechanisms create conditions that foster the evolution of antibiotic resistance (Brauner et al, 2016; Fridman et al, 2014; Levin-Reisman et al, 2017).

Despite the high frequency in biofilms and critical roles in both clinical context and evolution processes, the mechanisms underlying biofilm persister formation are the subject of intense debate. The effects induced by high density, including nutrient and oxygen deficiency, and quorum sensing within dense structure of developed biofilms, have long been considered as primary factors leading to persister formation (Ganesan et al, 2016; Nguyen et al, 2011; Yan & Bassler, 2019; Yan et al, 2018). This association originally stems from the hypothesis of limitations in nutrient, oxygen, or antibiotic penetration into the dense structures. Nevertheless, recent challenges to this hypothesis have arisen from studies indicating that the matrix mesh size of biofilms is significantly larger than the size of molecules such as nutrients, oxygen, and antibiotics (Ganesan et al, 2016; Yan & Bassler, 2019; Yan et al, 2018). Notably, most antibiotics do not strongly interact with biofilm matrix components (Spoering & Lewis, 2001). An alternative perspective emerges from observations that cells in mature biofilms exhibit metabolic activity similar to cells in the stationary phase (Nguyen et al, 2011; Walters et al, 2003; Waters et al, 2016), during which the rate of persister cell formation significantly surpasses that of cells in the exponential phase. This phenomenon is likely attributed to nutrient exhaustion in the local environment surrounding mature biofilms (Nguyen et al, 2011; Walters et al, 2003; Waters et al, 2016). Meanwhile, some findings propose the involvement of specific genes in persister cell formation in biofilms (Zhao et al, 2013), many of these observations are derived from research conducted under planktonic (free-swimming) conditions with bulk cells (Yan & Bassler, 2019).

Studying a biofilm during its early developmental stage offers a unique vantage point, minimizing the impact of dense structure or nutrient exhaustion, thus allowing for a controlled exploration of intrinsic factors influencing persister cell formation. Cell-surface adhesion is the critical initiation step in the early stage of biofilm development, encompassing both reversible and irreversible processes (Berne et al, 2018; Tolker-Nielsen, 2015). During reversible adhesion, planktonic bacteria utilize surface appendages such as flagella and pili, fostering a weak attachment through van der Waals forces (Ren et al, 2018). Subsequently, these reversible adhered cells sense environmental cues linked to the surface, activating signaling pathways that lead to the synthesis of cyclic di-GMP (c-di-GMP). Elevated c-di-GMP levels function as a secondary messenger, orchestrating a shift in bacterial behavior by influencing the expression of genes crucial to biofilm development. This signaling cascade propels the transition from reversible to irreversible adhesion, wherein bacteria establish microcolonies on the surface, solidifying a more stable and irreversible connection (Jenal et al, 2017; Valentini & Filloux, 2016).

To explore the intrinsic factors influencing persister cell formation during the early stages of biofilm
development, we employed a static biofilm system for our investigation (Fig. 1A). This system is widely
employed in research to scrutinize the pivotal processes of initial adherence during biofilm formation
(Biswas & Mettlach, 2019; Merritt et al, 2005; Seviour et al, 2021; Sung et al, 2022). In our initial study, we
noted an increased persister frequency coinciding with the step of cell adhesion. This observation prompted
the hypothesis that elevated c-di-GMP levels in individual cells potentially facilitate the formation of
persister cells within biofilms. Further investigation substantiated this hypothesis, demonstrating that the
elevation in c-di-GMP indeed promotes the formation of persister cells. Our detailed analysis uncovered the
role of c-di-GMP as an antitoxin, exerting control over both the expression and genotoxicity of HipH,
thereby regulating genome stability and persister cell formation. These findings propose a biofilm-specific
mechanism for antibiotic persistence. Notably, targeting this toxin-antitoxin system effectively inhibited
drug tolerance in Uropathogenic *Escherichia coli* (UPEC) infections, presenting promising therapeutic
strategies against chronic and relapsing infections.

Results

An increased propensity of persister formation upon cell ahesion

To enhance the clinical relevance of our study, we employed UPEC, the predominant cause of urinary tract infections (UTIs) and responsible for approximately 80% of cases. UPEC has a notable ability to form biofilms on various surfaces, including catheter materials, bladder epithelial cell walls, and within bladder cells (Supplementary Information). In our investigation, we selected the clinical isolate UPEC16. This strain demonstrates a significant phylogenetic similarity to established UPEC reference strains such as CFT073 and UTI89 (Fig. S1A). Moreover, it displays sensitivity to a wide range of antibiotics and possesses the capability for surface adhesion. To culture UPEC16 cells in the static biofilm system, exponential phase cells with an optical density (OD600) of 0.3 were diluted in 10-fold in fresh media, followed by static incubation in microtiter dishes for varying time intervals. Planktonic bacteria were then gently washed away, leaving adhered cells for further investigation. In the persister counting assay, both the remaining adhered cells and the removed planktonic cells were exposed to bactericidal antibiotics (ampicillin, 150 μg/ml, 10-fold of MIC) for varying durations. Subsequently, persister counting was conducted through plating.

Biphasic killing curves revealed distinct patterns. When cells were subjected to the same duration of static culture, the persister rate within the adhered population notably surpassed that observed in the planktonic population (Fig. 1B-D). Additionally, the persister rate within adhered populations exhibited a positive correlation with the prolonged duration of static culture (Fig. 1B-D). In contrast, the persister rate within planktonic populations displayed little variation with an extended duration of static culture (Fig. 1B-D). These observations underscore the crucial role of cell adhesion in promoting persister cell formation during the early stages of biofilm development. This raises questions about established ideas linking persister formation mainly with the dense structure of biofilms, suggesting a need to reconsider the factors that influence persister dynamics during biofilm initiation."

The correlation between heightened c-di-GMP levels and increased persister frequency

The increase in c-di-GMP levels is a distinctive feature of cell-surface adhesion, marking the transition of cells from a planktonic lifestyle to a sessile one (Fernandez et al, 2020; Jones et al, 2015; Valentini & Filloux, 2016). Building on our observations of increased persister rates upon cell adhesion, we assumed that elevated c-di-GMP levels might play a role in promoting persister cell formation. To investigate this hypothesis and discern the relationship between c-di-GMP levels and persister rates, we introduced a ratiometric fluorescent c-di-GMP sensing system (Vrabioiu & Berg, 2022) into the UPEC16 strain. MrkH is a transcription factor derived from *Klebsiella pneumoniae*. The system utilizes the conformational changes that occur in MrkH upon binding to c-di-GMP by strategically inserting it into mVenus[NB] between residues Y145 and N146, resulting in the creation of the biosensor referred to as mVenus[NB]-MrkH. This strategic integration results in a ratiometrically reduced fluorescence signal when the biosensor is bound to varying levels of c-di-GMP. Furthermore, a red fluorescent protein mScarlet-I is co-expressed under the regulation of the *proC* promoter (Davis et al, 2011), enabling the normalization of the system's expression level. Consequently, the inverse of the ratio between mVenus[NB]-MrkH and mScarlet-I (R^{-1}) exhibits a direct correlation with the intracellular level of c-di-GMP, establishing a reliable indicator for monitoring c-di-GMP dynamics in UPEC16 (Fig. S1B).

By monitoring the fluorescent intensity of the system at the single-cell level under a time-lapse
microscope, we observed the dynamic activation of c-di-GMP during the process of cell adhesion. We
imaged subpopulations on the surface at varied intervals. In the initial 30 minutes following static
incubation in microtiter dishes, the adhered subpopulation displayed a sharply peaked R^{-1} distribution with
a low median of 0.64 (Fig. 1B). As the attachment duration increased to 2 hours, the R^{-1} distribution
broadened, accompanied by a higher median of 0.93 (Fig. 1C). Further extension to 3.5 hours resulted in a
continued broadening of the R^{-1} distribution, accompanied by an even higher median of 1.18 (Fig. 1D).
Meanwhile, the R^{-1} distribution of the planktonic subpopulation from the same culture system and time
points remained low, with a median of 0.31 to 0.35, repectively (Fig. S1C-E). These findings replicate the
observed positive correlation between c-di-GMP production and attachment duration. Moreover, the results
suggest a positive association between c-di-GMP levels and the persister rate (Fig. 1B-D).

Microcolonies emerged on the surface of microtiter dishes when the duration of static culture extended to
24 hours (Fig. 1E). We employed confocal laser scanning microscopy to monitor spatial distribution of
c-di-GMP signals within these microcolonies. This investigation revealed a stratified pattern of c-di-GMP
levels across the cross-section of the static biofilms (R^{-1}, Fig. 1E), highlighting the intricate regulatory role
of this signaling molecule in shaping bacterial behavior and biofilm structure. Notably, cells with the
highest c-di-GMP levels were surface attached, predominantly situated at the bottom layers of the biofilm
(Fig. 1E). To further investigate this observation, we isolated the surface attached cells and unattached cells
in the biofilms for persister counting assay. The results demonstrate that this spatial distribution aligned
with a subpopulation that exhibited the highest persister rate (Fig. 1F). Furthermore, we resuspended the
cells within the static biofilms and used FACS to isolate subpopulations with varying levels of c-di-GMP
(Fig. 1G). An antibiotic killing assay revealed that subpopulations with higher concentrations of c-di-GMP
had a higher frequency of persister cells (Fig. 1H). Time-lapse imaging also revealed that cells with high
intracellular c-di-GMP levels survived antibiotic treatment and resuscitated to form microcolonies (Fig. 1I).
These findings collectively suggest that increased levels of cellular c-di-GMP play a crucial role in
promoting persister formation.

Heightened c-di-GMP levels promotes persister formation

To investigate the involvement of c-di-GMP in promoting the transition of persister phenotype in biofilms,
we first constructed a UPEC16 strain with the ability to degrade c-di-GMP in a controlled manner. This was
achieved through the overexpression of PdeH, an efficient c-di-GMP phosphodiesterase in *E. coli*, under
the control of the araBAD promoter (pBAD::*pdeH*) (Pesavento et al, 2008). Inducing PdeH expression
before cell attachment disrupted cell adhesion, validating the efficacy of our system. Initiating with
uninduced cells, we allowed them to adhere to a surface in a microliter dish (Fig. 2A). After a 2-hour
incubation, we observed robust cell adhesion similar to the vector control, reflected in comparable
c-di-GMP R^{-1} medians. Following this initial observation, we replaced the old medium with fresh medium
containing arabinose (0.2%) for PdeH induction. After an additional 30 minutes of incubation, we noted a
substantial decrease in c-di-GMP R^{-1} levels in the PdeH overexpression strain (median of 0.75) compared to
the vector control strain (median of 0.98) (Fig. 2B-C). Crucially, the biphasic killing curve showed
significantly lower persister rates in adhered populations with PdeH overexpression compared to the vector
control strain (Fig. 2D). These findings suggest that c-di-GMP plays a pivotal role in promoting persister
cell formation during cell-surface adhesion.

In parallel, we engineered a UPEC16 strain overexpressing DgcZ, a highly efficient diguanylate cyclase
responsible for synthesizing c-di-GMP, under the araBAD promoter (pBAD::*dgcZ*). In a consistent
approach, we allowed the cells to adhere to a surface in a microliter dish before induction (Fig. 2A). After a
2-hour incubation, we observed cell adhesion comparable to the vector control, reflected in similar
c-di-GMP R^{-1} medians. Subsequently, we replaced the old medium with fresh medium containing arabinose
(0.2%) for DgcZ induction. Following an additional 30 minutes of incubation, we noted a substantial
increase in c-di-GMP R^{-1} levels in the DgcZ overexpression strain (median of 1.72) compared to the vector
control strain (Fig. 2E). Consistently, the biphasic killing curve indicated significantly higher persister rates
in adhered populations with DgcZ overexpression compared to the vector control strain (Fig. 2D). These
findings reinforce the critical role of c-di-GMP in promoting persister cell formation during cell-surface
adhesion.

Toxin protein HipH acts downstream of c-di-GMP signaling in regulating persister formation in
biofilms

Consistent with the observations in adherent cells, planktonic cells with DgcZ overexpression demonstrated
a significant increase in the persister ratio (Fig. 3A). Yet, cells overexpressing DgcZ-mutant (DgcZ-m,
G206A, G207A, E208Q), which lacks diguanylate cyclase activity (Boehm et al, 2009; Jonas et al, 2008),
did not manifest a parallel increase (Fig. 3A). These results reinforce that elevated levels of c-di-GMP play
a key role in promoting persister cell transition. To explore the mechanisms by which elevated c-di-GMP
levels promote bacterial persister formation, we employed a systematic screening approach using the *E. coli*
Keio Knockout Collection (KEIO collection) (Baba et al, 2006). Given that the K-12 strain, the background
strain of the KEIO collection, might not be ideal for cell adhesion and biofilm formation, we initially
introduced the pBAD::*dgcZ* construct into individual mutant strains. This allowed us to establish a
background characterized by increased c-di-GMP levels (Fig. 3B). Subsequently, persister counting assays
were performed by exposing exponentially growing transformed strains to the antibiotic ampicillin at a
concentration of 150 μg/ml for a duration of 3 hours, followed by plating. This experimental design allowed
us to assess the impact of upregulated c-di-GMP on persister cell formation within specific genetic
backgrounds. Results show that knocking out the toxin gene *hipH* significantly reduces the frequency of
persister formation in the presence of high levels of c-di-GMP (Fig. 3C). Meanwhile, the knockout of *hipH*
alone does not lead to a significant alteration in persister frequency during the exponentially growing phase.
These observations suggest that HipH acts downstream of c-di-GMP signaling in the process of persister
formation and antibiotic tolerance. Considering HipH belongs to the family of HipA-like Ser/Thr kinases,
we investigated the impact of knocking out *hipA*. In contrast, knocking out *hipA* had little effect on the
frequency of persister formation with elevated c-di-GMP levels (Fig. 3C). This observation suggests that
HipH plays a distinct role from HipA in inducing persister formation.

Given that c-di-GMP exhibits upregulation in cell-surface adhesion, we conducted a *hipH* knockout in
UPEC16 strain (UPEC16-Δ*hipH*) and evaluated its influence on persister formation in biofilms (Fig. 3D).
In *in vitro* biofilms, we observed a substantial decrease in persister levels following *hipH* knockout,
contrasting with the wild-type strain that consistently displayed notably high levels of persister cells (Fig.
3E). These results suggest that c-di-GMP regulated HipH toxicity plays a crucial role in persister formation
of biofilms. For *in vivo* biofilm formation, we introduced both the wild-type and UPEC16-Δ*hipH* strains,
each at a concentration of 2*10^8 CFU, into the mouse bladders through transurethral catheterization (Fig.
3D). Following 24 hours duration of infection, both strains exhibited comparable levels of intracellular
bacterial communities (IBC) within the epithelial cells lining the bladder lumen (Fig. S2A-B).

Subsequently, the infected mice were sacrificed, and their bladders were isolated for colony counting
assays and antibiotic killing assays. Consistent with the findings from *in vitro* biofilms, the wild-type
UPEC16 strain displayed a notably high rate of persister formation. In contrast, the UPEC16-Δ*hipH* strain
exhibited a substantial reduction in persister cell frequency under the same conditions (Fig. 3F). The impact
of HipH knockout on persister formation is consistently observed in both *in vitro* and *in vivo* settings
associated with biofilms, but not in planktonic populations (Fig. 3C, left panel). These results reinforce
HipH toxicity is under the regulation of c-di-GMP, and emphasize the significance of HipH in shaping the
persistence and antibiotic tolerance characteristics of biofilms.

We observed a significant elevation in *hipH* gene expression upon the overexpression of DgcZ
(pBAD::*dgcZ*) (Fig. 3G-H). Conversely, the strain overexpressing DgcZ-m, which is unable to increase
c-di-GMP levels after induction, exhibit no activation of *hipH* gene expression (Fig. 3G). Furthermore, we
labeled HipH with BFP at the C-terminal end and placed it under the *hipH* native promoter in the UPEC16
strain with c-di-GMP sensing system. Fluorescent imaging revealed a positive correlation between HipH
and c-di-GMP expression at the single-cell level (Fig. S3A), supported by a correlation coefficient of 0.694
(Fig. S3B). Additionally, we engineered an *E. coli* strain in which *hipH* promoter is replaced by the
constitutive promoter p-rhaB (Fig. S3C). In this strain, *hipH* expression remains unaffected by changes in
c-di-GMP levels (Fig. S3D), indicating that regulation occurs at the promoter region. Furthermore, the
persister ratio was comparable to that of the Δ*hipH* strain but significantly lower than that of wild-type
strain under high c-di-GMP levels (Fig. S3E). These results together suggest that c-di-GMP exerts a
transcriptional regulatory role in controlling the expression of HipH.

c-di-GMP and HipH form a toxin-antitoxin module

We then independently overexpressed HipH (pBAD::*hipH*) to observe its impact on cell growth. The
results demonstrated that HipH overexpression substantially inhibit cell growth, with only 9.8±3.4% of
cells retaining growth potential (Fig. 3I). Further analysis of the growth-inhibited cells revealed a
predominant presence of Viable but Non-Culturable (VBNC) cells, constituting 72±5% of the total cell
population, alongside a significant proportion of dead cells, accounting for 17.9±2.3% of the total cells (Fig.
3J). This evidence suggests that HipH functions as a potent toxin capable of inhibiting cell growth or even
inducing cell death. Interestingly, despite the observed decrease in growth potential, the persister rate
remains low as that of wild-type strains (Fig. 3K). This phenomenon is interpreted as cells entering a state
of deep dormancy with reduced resuscitation capabilities (Fig. 3L).

To test the role of c-di-GMP as an antitoxin in alleviating the toxic effect of HipH, we elevated
intracellular c-di-GMP levels by overexpressing DosC, a moderately efficient diguanylate cyclase, or DgcZ,
a highly efficient diguanylate cyclase, in the strain overexpressing HipH. This manipulation resulted in a
2.72-fold and 4.22-fold increase in the mean of c-di-GMP R^{-1} score, respectively (Fig. S3F). Comparing to
the strain overexpressing HipH alone, those strains co-overexpressing HipH and c-di-GMP exhibited a
significantly recovered growth potential (Fig. 3I). Specifically, 36±6.7% of cells retained growth potential
in the strain expressing HipH and DosC, while approximately 85±18% retained growth potential in the
strain expressing HipH and DgcZ. Further analyzing the portion of growth-inhibited cells revealed that the
strain overexpressing HipH and DosC exhibited a decreased rate of VBNC cells (62±6% of the total cell
population) and dead cells (2.1±0.2% of the total cell population) (Fig. 3J). The strain overexpressing HipH
and DgcZ showed an even lower rate of VBNCs (14±17% of the total cell population) and dead cells
(1.5±0.4% of the total cell population) (Fig. 3J). These results support the assumption that c-di-GMP acts as

an effective antitoxin, enabling HipH-toxified cells to recover their growth potential. Consistently,
increased rates of persister cells were observed, with $10^{-2.8\pm0.19}$ in the strain overexpressing HipH and
DosC, and $10^{-2.5\pm0.34}$ in the strain overexpressing HipH and DgcZ (Fig. 3K). These results collectively
indicate that c-di-GMP functions as an antitoxin in alleviating HipH's harmful effects, steering cells from
deep dormancy depth to a shallower dormancy state with enhanced resuscitation capabilities, thereby
promoting persister cell formation (Fig. 3L).

HipH is a deoxyribonuclease inducing genome instability

hipH stands out as an atypical toxin gene due to its lack of a genetically linked antitoxin cognate, existing as
a solitary gene unlike the typical toxin-antitoxin gene pairs found within the same operon (Jurenas et al,
2022; Maeda et al, 2017). Previous studies believed it a homolog of the toxin protein HipA, functioning as
a Ser/Thr protein kinase involved in regulating ribosome assembly and cell metabolism (Gratani et al,
2023). However, our findings challenge this prevailing belief. The truncated mutant of HipH, lacking the
kinase catalytic motif (Δ5, amino acid 340-372), demonstrated a comparable ability to inhibit cell growth
and induced VBNCs compared to the intact HipH (Fig. 4A-C). These results suggest the involvement of
another domain in HipH that is responsible for its toxic activity.

Structure analysis revealed that, in addition to its kinase catalytic motif (amino acid 340-372), HipH also
contains a DNA binding HTH domain (amino acid 11-32), a gly loop and autophosphorylation site (amino
acid 190-215), and an activation loop (amino acid 222-229) (Fig. 4A). To identify the specific domain
responsible for its toxicity, we generated various truncated mutants and evaluated their capacity in
inhibiting cell growth. Our results showed that overexpression of mutants lacking the DNA binding domain
HTH (Δ1 and Δ2) did not inhibit cell growth or induce VBNCs or dead cells (Fig. 4B-C). This indicates that
the DNA binding domain HTH is indispensable for the observed toxicity of HipH. Consistent with these
results, fluorescent microscopy confirmed the nucleoid localization of wild-type HipH (Fig. 4D), indicating
that the primary function of HipH is related to DNA interaction.

We observed the elongated filamentous shape induced by HipH overexpression in cells (Fig. S4A),
which is consistent with previous findings (Gratani et al, 2023). We interpret this as a potential
characteristic associated with DNA damage. Furthermore, considering the results of the truncated mutation
assay (Fig. 4A-C), we hypothesize that HipH plays a role in inducing DNA damage. To investigate this
hypothesis, we first examined whether HipH overexpression induces the SOS response, a bacterial defense
mechanism against DNA damage involving the expression of various genes. qPCR analysis revealed a
significant upregulation of SOS response genes, such as *recA*, *sulA*, and *tisB*, in cells overexpressing HipH
(Fig. S4B). Notably, the near abolition of cell growth potential upon knockout of *recA,* a crucial protein
essential for DNA damage repair, in cells overexpressing HipH further supports the conclusion that HipH
induces DNA damage (Fig. S4C).

To investigate the specific type of DNA damage caused by HipH, we conducted *in vitro* DNA-cleaving
assays using purified HipH. To address its insolubility, we engineered a fusion construct with a high
solubility protein, GFP, at the C-terminal end to facilitate protein purification. The results demonstrated
HipH's capability to cleave supercoiled plasmids, leading to the production of double-strand breaks (DSBs),
as evidenced by prominent smears observed on agarose gels (Fig. 4E-G). Further analysis revealed a
positive correlation between HipH's DNA cleavage capacity and its protein concentration, temperature, and
incubation time (Fig. 4E-G). These findings collectively indicate that HipH cleaves DNA and produces
DSBs. The overexpression of RecG, a key player in the repair of DSBs in DNA (Whitby et al, 1994),

successfully rescued growth deficient cells resulting from HipH overexpression (Fig. S4D), providing
additional evidence that HipH induces DSBs in DNA. These compelling results demonstrate that HipH
functions as a deoxyribonuclease, exerting genotoxic effects on genomic DNA integrity.

c-di-GMP acts as an antitoxin to alleviate HipH genotoxicity

To investigate whether c-di-GMP plays as an antitoxin to mitigate HipH genotoxicity, we first explored the
potential of direct interaction between c-di-GMP and HipH. A biotinylated c-di-GMP pull-down assay (Fig.
5A) (Chambers & Sauer, 2017), conducted from both cell lysate and purified protein, revealed a physical
binding between c-di-GMP and HipH (Fig. 5B-D, Fig. S5A-B), confirming their direct interaction. The
dissociation constant (K_D), calculated based on the concentration of c-di-GMP at which half-maximal
binding of HipH occurs, was determined as 7.6 µM (Fig. 5B, bottom panel). Additionally, c-di-GMP
successfully displaced HipH-GFP-His from HipH-GFP-His/biotin-c-di-GMP complexes, while other
nucleotides such as cAMP and GTP did not exhibit the same effect (Fig. 5E). This underscores the
specificity of the binding between c-di-GMP and HipH.

Subsequently, we introduced c-di-GMP into the HipH DNA-cleavage assay system, and notably, its
addition exerted a significant suppressive effect on the cleavage activity of HipH. This effect was observed
for both relaxed and supercoiled plasmids, as evidenced by a reduction of smears and double-strand break
(DSB) bands on agarose gels (Fig. 5F-I). It is noteworthy that other nucleotides, including cAMP and GTP,
did not demonstrate the same suppressive effect on HipH's cleavage activity (Fig. S5C). Furthermore, it has
been reported that HipB serves as an antitoxin for HipH (Gratani et al, 2023; Maeda et al, 2017). However,
when we tested the potential impact of HipB on HipH's DNase activity by purifying and introducing HipB
into our cleavage system, we found that HipB failed to inhibit HipH's DNase activity (Fig. S5C). These
findings suggest that c-di-GMP specifically interacts with HipH, leading to the inhibition of its DNase
activity.

To monitor DNA breakage in live bacterial cells with varying stoichiometry between HipH and
c-di-GMP, we utilized Gam-GFP, a double-strand DNA end-binding protein of bacteriophage Mu that
forms foci at DSBs in living cells (Shee et al, 2013). Microscopic examination revealed that cells
overexpressing HipH alone displayed Gam-GFP foci in 8.85±1.28% of cases, while only 0.30±0.12% of
wild-type cells demonstrated such foci (Fig. 5J-K). Further investigations showed that elevation of
intracellular c-di-GMP levels resulted in a significant reduction in the percentage of cells with Gam-GFP
foci, with DosC addition decreasing the occurrence to 1.34±0.39% and DgcZ to 0.46±0.19% (Fig. 5J-K).
These *in vitro* and *in vivo* findings conclusively demonstrate that c-di-GMP acts as an antitoxin for HipH,
effectively suppressing its genotoxic effects.

Discussion

In the intricate biofilm dynamics, the origin of persister formation has largely been ascribed to the
constraints imposed by the dense structures, such as limited nutrients, oxygen, and antibiotics penetration,
or quorum sensing. Our investigation reveals that cell adhesion, a pivotal event initiating biofilm
development, emerges as a trigger for persister formation. This concept introduces a layer of complexity
that reshapes our understanding of drug tolerance in biofilms. Beyond the traditional emphasis on dense
biofilm structures, the scope should now encompass single-layer cells on the surface. Such a deviation from

conventional perspectives highlights the necessity for a more nuanced and comprehensive approach to
address the challenges of antibiotic efficacy within biofilm communities.

c-di-GMP emerges as a pivotal secondary messenger in the intricate orchestration of bacterial lifestyles,
playing a key role in mediating cell-surface adhesion and exerting influence over the transition from
planktonic to sessile existence. Our comprehensive exploration, employing a spectrum of techniques,
brings to light a significant revelation: heightened levels of c-di-GMP in cells drive the formation of
persisters. Contrary to conventional understanding, this system unveils a unique function for c-di-GMP—it
acts as a noncanonical antitoxin, which controls both the expression and genotoxicity of toxin protein
HipH. Within this framework, c-di-GMP, a small molecule with just two guanine bases, takes on the role of
an antitoxin, counterbalancing the toxin. This discovery challenges established toxin-antitoxin (TA) system
norms, in which antitoxins typically manifest as proteins or RNA molecules directly encoded from gene
pairs within the same operon of toxin. The unconventional role of c-di-GMP as an antitoxin redefines our
comprehension of the molecular players orchestrating persister formation, adding a layer of complexity to
the regulatory mechanisms governing bacterial adaptation to sessile lifestyle.

HipH emerges as an intriguing toxin within the bacterial realm, standing apart due to its solitary
existence without a genetically linked antitoxin cognate—an uncommon trait in known toxin-antitoxin
systems. Previous studies have identified HipH as a Ser/Thr protein kinase, sharing homology with the
well-characterized toxin HipA. These studies demonstrated its primary protein substrates to be the
ribosomal protein RpmE (L31) and the carbon storage regulator CsrA (Gratani et al, 2023). While the
overproduction of HipH was observed to impact bacterial DNA segregation and cell division, the
underlying mechanism remained unexplained. In this study, we have uncovered that HipH primarily
functions as a potent DNase, providing an explanation for the observed phenomenon. Our findings further
reveal that that HipH, as a potent DNase, induces a deep cell dormancy depth and elevated rates of VBNC
cells and dead cells, rather than persister cells with the capacity of resuscitation. Significantly, our
investigation further sheds light on the regulation of HipH genotoxicity by c-di-GMP in a dual manner.
Firstly, high intracellular levels of c-di-GMP induce HipH expression. Subsequently, c-di-GMP acts as an
antitoxin, mitigating the potent toxicity of HipH. This, in turn, maintains cells with an adequate dormancy
depth, facilitating the formation of persister cells and conferring antibiotic tolerance. It is intriguing to delve
deeper into research exploring the domain-function relationship in HipH. However, our efforts have often
encountered challenges, as mutations can readily induce protein aggregates, impeding thorough analysis.
Achieving a comprehensive understanding of this domain-function relationship necessitates the
development of innovative methodologies and approaches.

Taken together, we reveal a sophisticated regulatory circuit underlying bacterial genome instability and
antibiotic persistence in biofilms. This regulated genome instability can have significant implications for
adaptive potential of bacterial populations, shaping their ability to persist, evolve, and develop resistance
mechanisms. By targeting this c-di-GMP regulated genotoxicity of toxin protein HipH, we successfully
inhibited drug tolerance in UPEC induced biofilms. These findings provide potential therapeutic strategies
for combating chronic and relapsing infections associated with biofilms.

Materials and Methods

Bacterial strains and growth conditions

Bacterial strains used in this study included *Escherichia coli* strains MG1655, BW25113, UPEC16, and

the Keio library Knockout Collection (Supplementary Table S1). UPEC16 was isolated from a urine

sample (#16) obtained from a local urinary tract infection (UTI) patient at the First Affiliated Hospital of

Kunming Medical University, following the guidelines of the Committee on Human Subject Research

and Ethics (2022-L-203 and CHSRE2022030). The genome sequence has been deposited in the National

Genomics Data Center under the accession number "GWHERGI00000000". All strains were cultured in

Luria Broth (LB) medium, with routine growth condition at 37°C and 220 rpm. To support plasmid

maintenance, media were supplemented with the corresponding antibiotics as needed: chloramphenicol

(25 µg/ml), kanamycin (20 µg/ml), gentamicin (20 µg/ml), and ampicillin (100 µg/ml). For inducible

expression experiments, the medium was supplemented with induces such as 0.002%-0.2% arabinose for

the Arabinose-induction system, 1 mM IPTG for the Lac-induction system, and 1 µg/ml

anhydrotetracycline for the Tet-induction system.

Strains construction

Construction of Target DNA Knockout. The generation of target DNA knockout strains involved the use

of λ-red mediated gene replacement (Datsenko & Wanner, 2000). In this procedure, a 500 bp upstream

region of the target DNA sequence, a resistance cassette for substitution (CmR for UPEC16 Δ*hipH*, GmR

for UPEC16 Δ*araD-B*), and a 500 bp downstream region of the target DNA sequence were fused through

fusion PCR. The resulting fusion fragments were then transformed into electrocompetent cells along with

a recombineering helper plasmid, pSim6 (Datta et al, 2006). After a recovery period of 3-5 hours, the

transformed cells were plated on selection plates containing the corresponding antibiotics. the

antibiotic-selected recombinant colonies were subsequently validated through PCR analysis and

sequencing. Finally, the correct recombinants were incubated at 42°C to eliminate the

temperature-sensitive helper plasmid pSim6.

Recombinant Plasmid Construction. Construction of recombinant plasmids was performed using the 2 ×

MultiF Seamless Assembly Mix (ABclonal, RK21020). To tag HipH, *hipH*, together with its native

promoter, was fused with Blue Fluorescent Protein gene *bfp* and inserted into the pBAD backbone by *in*

vitro recombination. For Gam-GFP overexpression, the *gam* gene from bacteriophage Mu was fused with

Green Fluorescent Protein gene *gfp* and inserted into a p15A ori plasmid by *in vitro* recombination. The

gam-gfp was placed under the control of the Tetracycline induction system (Addgene, pFD152, #125546).

Similarly, for *recG* overexpression, the *recG* gene was cloned into a p15A ori plasmid, placed under the

control of the T5 promoter-Lac-induction system. To express HipH-His, mCherry-His, DosC, and DgcZ

(DgcZ-m), these genes were cloned into pBAD, under the control of the Arabinose-induction promoter.

Additionally, for the c-di-GMP sensor system purchased from Addgene (#182291), the plasmid origin

was substituted with the p15A ori.

Static biofilm culture

431   The methodology employed in this study was adapted from the established one (Merritt et al, 2005).

432   Bacterial strains of interest were initially inoculated in a 3-to-5-ml culture in tubes and allowed to grow

433   overnight. Subsequently, the stationary cultures were diluted 1:100 in the desired media and cultured until

434   reaching the exponential phase with an OD600 of 0.3. Next, 2 ml of each diluted culture was transferred

435   into a fresh microtiter dish (Biosharp, Cat#BS-20-GJM), which has not been undergo tissue culture

treatment. In cases where the subsequent experiment involves imaging, it is necessary to ensure that the microtiter dish has a glass bottom. The plate was covered and incubated at optimal growth temperature for the desired amount of time. While the lids for the microtiter dishes are reusable, it is important to clean them with 70% (v/v) ethanol and allow them to air-dry prior to each experiment.

For cultures involving PdeH or DgcZ induction, it is important to statically incubate cells in microtiter dish for 2 hours at 37°C first. This incubation period allows cells adherence to the surface. Subsequently, the inducer arabinose was added to the desired concentration to initiate protein induction with desired duration.

Antibiotic killing assay and persister counting

For cells from static culture, planktonic cells were directly transferred into new tubes and harvested by centrifugation. The remained microliter dishes were gently washed twice by PBS buffer, and adherent cells were harvested by vigorous pipetting. In other cases, the cells were harvested routinely. Subsequently, the harvested cells directly plated on LB plate for colony-forming unit (CFU) counting without antibiotic challenge.

The harvested cells were washed with PBS buffer and subsequently resuspended into fresh LB containing 150 μg/ml ampicillin and incubated for varying durations (0 to 4.5 hours) at 37°C with shaking at 220 rpm. The survived cells after ampicillin challenge were harvested by centrifugation and diluted in PBS, followed by plating on LB plate for overnight culture at 37°C. Persister counting was performed on the next day. Persister ratios were calculated as the ratio of the CFU after antibiotic treatment to the CFU without antibiotics challenge. At least three biological replicates were performed for each experiment. Averages and standard deviations presented are representative of biological replicates.

Cell growth potential assay and VBNC determination.

To evaluate growth potential, bacterial cells were cultured to exponential phase in LB medium to an OD600 of 0.3. For strains harboring an arabinose-induction system for specific protein expression, 0.002% arabinose (w/v) was added to the medium for 3 to 4 hours at 37°C with shaking at 220 rpm. The cells were then plated for colony-forming unit (CFU) counting on LB plates before and after arabinose induction. The cells with growth potential were determined as the ratio of CFUs after arabinose induction to CFUs before arabinose induction. The dead cells ratio was assessed by staining with PI dye. The ratio of VBNC cells was calculated as 100% minus the sum of the ratios of cells with growth potential and dead cells. Averages and standard deviations presented are representative of three biological replicates.

Microscopy

Bright-field and fluorescence imaging. Inverted microscope Nikon (eclipse Ti2) and confocal microscope (Leica, Stellaris 5 WLL) were used in this study. The illumination was provided by different lasers, at wavelengths 405 nm for BFP and Hoechst 33258, 488 nm for GFP and SYTO[TM]24, 488 nm (Nikon) or 515 nm (Leica) for mVenus[NB], 561 nm for propidium iodide (PI) and mScarlet-I, respectively. The fluorescence emission signal was imaged to an sCMOS camera (pco.edge 4.2 bi). Appropriate filter sets were selected for each fluorophore according to their spectrum. Image analysis was done by ImageJ software (Fiji). For c-di-GMP sensor analysis, the ratio of mVenus[NB] to mScarlet-I (R) has a negative

correlation with the concentration of c-di-GMP. Therefore, R^{-1} demonstrates a positive correlation with

the concentration of c-di-GMP.

Cells staining. PI, Hoechst 33258 and SYTO^{TM}24 were added to culture at a final concentration of 1

μg/ml, 10 μg/ml and 10 μM, respectively. The cells were then incubated in the dark at room temperature

for 15 to 30 minutes. Prior to imaging, the cells were washed twice with PBS.

Gam-GFP foci imaging. The method employed in this study was adapted from the established one (Shee et

al, 2013). Cells expressing toxin protein HipH along with DosC or DgcZ under the arabinose-induction

system and expressing Gam-GFP under Tetracycline-induction system were cultured to exponential phase

in LB medium with supplement of adequate antibiotics for plasmid maintenance. Gam-GFP was induced

by 1 μg/ml anhydrotetracycline for 30 minutes and then HipH and DosC (or DgcZ) were induced by 0.2%

(w/v) arabinose for 2.5 h at 37°C and 220 rpm. The cells were washed twice using PBS buffer and then

visualized under a fluorescence microscope. The Gam-GFP foci were analyzed using ImageJ software

from the individual cell intensity histograms. At least four neighboring pixels, each with an individual

fluorescence intensity (f) greater than 1.2 times the mean fluorescence intensity of the cell (a), were

deemed as foci. Meanwhile, the area occupied by these pixels needed to be less than one-fifth of the total

bacterial area.

Time-lapse imaging. For recording antibiotic killing and bacteria resuscitation processes, cells harboring

c-di-GMP sensor in the 24-hour static culture were collected and washed twice with PBS buffer and

imaged on a gel-pad containing 3% low melting temperature agarose in PBS (Pu et al, 2019; Pu et al,

2016). The gel-pad, prepared at the center of the FCS3 chamber as a gel island, was observed under bright

field or epifluorescence illumination. Subsequently, the gel-pad was encircled by LB broth containing 150

μg/ml ampicillin for 6 hours at 35°C. Fresh LB broth was introduced, and the growth medium was refreshed

every 3 hours to allow cells to recover sufficiently.

Fluorescence-activated Cell Sorting FACS analysis

All samples were analyzed using a Beckman CytoFLEX SRT flow cytometer equipped with a 100 μm

nozzle, and normal saline served as the sheath fluid. The c-di-GMP sensor-labeled strain in the 24-hour

static biofilm growth phase was washed and resuspended in sterile PBS. Bacterial cells were identified

based on FSC (forward scatter) and SSC (side scatter) parameters. Cells were sorted into distinct groups

according to their fluorescence intensity (PB450 for BFP, FITC for mVenus^{NB}, ECD for mScarlet-I). The

obtained results were analyzed using FlowJo V10 software (Treestar, Inc.).

Protein purification

All proteins used in this study were purified through affinity chromatography using a His-tag and a nickel

column. Briefly, MG1655 Δ*ara* harboring pBAD::*hipH-GFP*-His (or *GFP*-His) grew to exponential

growth phase in 500 ml LB medium. Induction of HipH-GFP-His (or GFP-His) was carried out by 0.002%

arabinose for 2 hours at 37°C with shaking at 220 rpm. The cells were harvested and resuspended in 25

ml binding buffer (50 mM Tris-HCl, 200 mM NaCl, 10 mM imidazole, 0.5% Triton X-100, pH 8.0)

containing 1 mg/ml lysozyme. The samples were incubated at 37 °C for 30 minutes and then subjected to

ultrasonic disruption.The cell lysate was centrifuged at 12,000 g for 20 minutes at 4°C, and the

supernatant was collected for protein purification. Ni-TED SefinoseTM Resin (Sangon Biotech, C610030)

was utilized for the purification of His-tagged proteins as per the manual. Finally, HipH-GFP-His (or
GFP-His) was stored in elution buffer containing 10% glycerol at -80°C.

RNA extraction and real-time quantitative PCR (RT-qPCR) assay

Bacterial strains were grown to mid-exponential growth phase and then induced by 0.002% arabinose for
2 hours. Cells were harvested and washed by PBS twice. Total RNA extraction was carried out using total
RNA extraction reagent (Vazyme R403, China), following the manufacture's instruction. Subsequently, 1
μg of total RNA was digested with DNase I and reverse transcribed to cDNA using a reverse transcription
kit (Vazyme R223, China). The resulting cDNA was used as template for RT-qPCR (Vazyme Q711,
China). The RT-qPCR assay were analyzed using BIO-RAD CFX Connect Real-Time PCR Detection
Systems. The 16S rRNA gene was selected as the internal control. The primers used in RT-qPCR assay
were listed in supplementary Table S1. All experiments were performed in biological triplicates to ensure
reliability and reproducibility of the results.

Biotin c-di-GMP pull down assay

The pull-down assay was conducted as previously described (Chambers & Sauer, 2017) with some
modifications. In the case of using purified proteins, purified HipH-GFP-His or GFP-His protein was
added into binding mix containing biotin-c-di-GMP (200 pmol/μl, BIOLOG, B098), 0.2 μl EDTA (200
mM), 2 μl 10×buffer (100 mM Tris-HCl, pH 7.5, 500 mM KCl, 10 mM DTT). The reaction mix was
brought to a final volume of 20 μl with the addition of H_2O and then incubated at 25°C for 20 minutes. In
the case of using cell lysate, MG1655 Δ*ara* harboring pBAD::*hipH*-His (or *mCherry*-His) was cultured to
exponential phase with an OD600 of 0.3 in 30 ml LB medium. 0.002% arabinose was then added into the
culture for another 2 h at 37°C with shaking at 220 rpm. The cells were harvested and resuspended in 1.5
ml lysis buffer (TBST containing 1 mg/ml lysozyme, 5 μg/ml DNaseI, 1× protease inhibitor), followed by
incubation at 37°C for 30 minutes and three cycles of freezing and thawing. The cell lysate was
centrifuged at 12,000g for 20 minutes at 4°C, and the supernatant was collected for the pull-down assay.
The specified amount of cell lysate was introduced into a reaction mix containing the indicated amount of
biotin-c-di-GMP, 1 μl EDTA (200 mM), and 10 μl 10× buffer. The reaction mix was brought to a final
volume of 100 μl with the addition of H_2O and then incubated at 25°C for 20 minutes.

10 μl (per sample) Streptavidin (SA) magnetic beads (Thermo Scientific, 88816) was washed twice
using TBST and resuspended in 150 μl TBST. Washed SA beads were added into reaction buffer and
incubated at 25°C for 1 h on a rotary table. Subsequently, the SA beads were washed five times using
TBST. Biotin-c-di-GMP-conjugated proteins on SA beads were eluted with SDS loading buffer and
analyzed by Western blot.

DNA cleavage assay

For DNA cleavage, 500 ng supercoiled plasmid pBR322 obtained from the FastPure Plasmid Mini Kit
(Vazyme, DC201-01) or relaxed plasmid pBR322 (derived from supercoiled pBR322 digested by
Topoisomerase I, Takara, 2240A), along with 6 μl 5× buffer (170 mM Tris-HCl, pH 7.5, 120 mM KCl, 10
mM DTT, 20 mM MgCl, 25 mM spermidine, 25% Glycerol) and 3 μl BSA (1 mg/ml), were added to the
binding mix (as mentioned in Biotin c-di-GMP pull down assay section).

The reaction mix was brought to a final volume of 30 μl with the addition of H_2O and then incubated at
37°C for varying duration. The reactions were halted by adding SDS to a final concentration of 0.2% and
incubated at 37°C for 20 minutes. The resulting product DNA was analyzed by 1% (w/v) agarose gel
electrophoresis.

Mouse urinary tract infection

Cells of strains UPEC16-WT and UPEC16 Δ*hipH* were grew to exponential phase with an OD600 of 0.3,
respectively. The cells were harvested and washed with PBS, followed by resuspension in equal volumes
of PBS. Bladder infection was conducted following previously described procedures with slight
modifications (Pang et al, 2022). 8-week female BALB/c mice were anesthetized using isoflurane. $2*10^8$
CFU of each strain were transurethrally inoculated by use of 25G blunt needle. Two rounds of infection
were performed with 24-hour interval. At 48-hour post-infection, the mice were sacrificed by cervical
dislocation, and the bladders were aseptically removed for following IBC counting assay and persister
counting assay. Animal studies were approved by the Animal Care Ethics Committee of Medical
Research Institute, Wuhan University. We confirm that all experiments conform to the relevant regulatory
standards.

Statistical analysis

577   Statistical analysis was performed in GraphPad Prism 9 software for Windows. Significance was
578   ascertained by two-tailed Student's t test or Two-way ANOVA. Error bars represent the standard
579   deviations of the mean from at least three independent experiments. A value of $P<0.05$ was considered
580   significant. " * " indicate significant differences (*, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; ****, $P<0.0001$).

Data availability

583   This study includes no data deposited in external repositories.

584
585

Acknowledgements

We thank Profs. Hongbin Shu (Wuhan University) and Fan Bai (Peking University) for valuable
discussions. We also thank the members of our laboratory for helpful discussion. This work is supported by
the grants to Y.P. from the National Key R&D Program of China (2021YFC2701602), the Natural Science
Foundation of China (31970089), and Science Fund for Distinguished Young Scholars of Hubei Province
(2022CFA077). This work is also supported by the grants to Y.Z. for the National Science Fund for
Distinguished Young Scholars (82025011) and C.G. from the Natural Science Foundation of Yunnan
Province of China (202001BB050005). We also thank all the staff in the Core Facilities of Medical
Research Institute at Wuhan University and the Core Facilities at School of Life Sciences at Peking
University for their technical support.

Conflict of interest

Authors declare that they have no competing interests.

References
Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M, Wanner
BL, Mori H (2006) Construction of Escherichia coli K-12 in-frame, single-gene knockout
mutants: the Keio collection. *Molecular systems biology* **2:** 2006 0008

604
605 Balaban NQ, Merrin J, Chait R, Kowalik L, Leibler S (2004) Bacterial persistence as a
606 phenotypic switch. *Science* **305:** 1622-1625

607
608 Berne C, Ellison CK, Ducret A, Brun YV (2018) Bacterial adhesion at the single-cell level.
609 *Nature reviews Microbiology* **16:** 616-627

610
611 Bigger J (1944) Treatment of Staphyloeoeeal Infections with Penicillin by Intermittent
612 Sterilisation. *Lancet***:** 497-500

613
614 Biswas I, Mettlach J (2019) A Simple Static Biofilm Assay for Acinetobacter baumannii.
615 *Methods in molecular biology* **1946:** 159-165

616
617 Boehm A, Steiner S, Zaehringer F, Casanova A, Hamburger F, Ritz D, Keck W, Ackermann M,
618 Schirmer T, Jenal U (2009) Second messenger signalling governs Escherichia coli biofilm
619 induction upon ribosomal stress. *Molecular microbiology* **72:** 1500-1516

620
621 Brauner A, Fridman O, Gefen O, Balaban NQ (2016) Distinguishing between resistance,
622 tolerance and persistence to antibiotic treatment. *Nature reviews Microbiology* **14:** 320-330

623
624 Ceri H, Olson ME, Stremick C, Read RR, Morck D, Buret A (1999) The Calgary Biofilm Device:
625 new technology for rapid determination of antibiotic susceptibilities of bacterial biofilms.
626 *Journal of clinical microbiology* **37:** 1771-1776

627
Chambers JR, Sauer K (2017) Detection of Cyclic di-GMP Binding Proteins Utilizing a
Biotinylated Cyclic di-GMP Pull-Down Assay. *Methods in molecular biology* **1657:** 317-329

Costerton JW, Stewart PS, Greenberg EP (1999) Bacterial biofilms: a common cause of
persistent infections. *Science* **284:** 1318-1322

Datsenko KA, Wanner BL (2000) One-step inactivation of chromosomal genes in Escherichia
coli K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United*
States of America **97:** 6640-6645

Datta S, Costantino N, Court DL (2006) A set of recombineering plasmids for gram-negative
bacteria. *Gene* **379:** 109-115

Davis JH, Rubin AJ, Sauer RT (2011) Design, construction and characterization of a set of
insulated bacterial promoters. *Nucleic acids research* **39:** 1131-1141

Fernandez NL, Hsueh BY, Nhu NTQ, Franklin JL, Dufour YS, Waters CM (2020) Vibrio
cholerae adapts to sessile and motile lifestyles by cyclic di-GMP regulation of cell shape.
Proceedings of the National Academy of Sciences of the United States of America **117:**
29046-29054

Fridman O, Goldberg A, Ronin I, Shoresh N, Balaban NQ (2014) Optimization of lag time
underlies antibiotic tolerance in evolved bacterial populations. *Nature* **513:** 418-421

Ganesan M, Knier S, Younger JG, Solomon MJ (2016) Associative and Entanglement
Contributions to the Solution Rheology of a Bacterial Polysaccharide. *Macromolecules* **49:**
8313-8321

Gratani FL, Englert T, Nashier P, Sass P, Czech L, Neumann N, Doello S, Mann P, Blobelt R,
Alberti S, Forchhammer K, Bange G, Hofer K, Macek B (2023) E. coli Toxin YjjJ (HipH) Is a
Ser/Thr Protein Kinase That Impacts Cell Division, Carbon Metabolism, and Ribosome
Assembly. *mSystems* **8:** e0104322

Hall-Stoodley L, Costerton JW, Stoodley P (2004) Bacterial biofilms: from the natural
environment to infectious diseases. *Nature reviews Microbiology* **2:** 95-108

Jenal U, Reinders A, Lori C (2017) Cyclic di-GMP: second messenger extraordinaire. *Nature*
reviews Microbiology **15:** 271-284

Jonas K, Edwards AN, Simm R, Romeo T, Romling U, Melefors O (2008) The RNA binding
protein CsrA controls cyclic di-GMP metabolism by directly regulating the expression of
GGDEF proteins. *Molecular microbiology* **70:** 236-257

Jones CJ, Utada A, Davis KR, Thongsomboon W, Zamorano Sanchez D, Banakar V, Cegelski L,
Wong GC, Yildiz FH (2015) C-di-GMP Regulates Motile to Sessile Transition by Modulating
MshA Pili Biogenesis and Near-Surface Motility Behavior in Vibrio cholerae. *PLoS pathogens*
11: e1005068

Jurenas D, Fraikin N, Goormaghtigh F, Van Melderen L (2022) Biology and evolution of
bacterial toxin-antitoxin systems. *Nature reviews Microbiology* **20:** 335-350
678
679 Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shoresh N, Balaban NQ (2017) Antibiotic
680 tolerance facilitates the evolution of resistance. *Science* **355:** 826-830
681
682 Lewis K (2005) Persister cells and the riddle of biofilm survival. *Biochemistry Biokhimiia* **70:**
683 267-274
684
685 Lewis K (2007) Persister cells, dormancy and infectious disease. *Nature reviews Microbiology* **5:**
686 48-56
687
688 Maeda Y, Lin CY, Ishida Y, Inouye M, Yamaguchi Y, Phadtare S (2017) Characterization of
689 YjjJ toxin of Escherichia coli. *FEMS microbiology letters* **364**
690
691 Merritt JH, Kadouri DE, O'Toole GA (2005) Growing and analyzing static biofilms. *Current*
692 *protocols in microbiology* **Chapter 1:** Unit 1B 1
693
694 Nguyen D, Joshi-Datar A, Lepine F, Bauerle E, Olakanmi O, Beer K, McKay G, Siehnel R,
695 Schafhauser J, Wang Y, Britigan BE, Singh PK (2011) Active starvation responses mediate
696 antibiotic tolerance in biofilms and nutrient-limited bacteria. *Science* **334:** 982-986
697
698 Pang Y, Cheng Z, Zhang S, Li S, Li X, Li X, Zhang X, Li X, Feng Y, Cui H, Chen Z, Liu L, Li
699 Q, Huang J, Zhang M, Zhu S, Wang L, Feng L (2022) Bladder epithelial cell phosphate
700 transporter inhibition protects mice against uropathogenic Escherichia coli infection. *Cell reports*
701 **39:** 110698
702
703 Pesavento C, Becker G, Sommerfeldt N, Possling A, Tschowri N, Mehlis A, Hengge R (2008)
704 Inverse regulatory coordination of motility and curli-mediated adhesion in Escherichia coli.
705 *Genes & development* **22:** 2434-2446
706
707 Pu Y, Li Y, Jin X, Tian T, Ma Q, Zhao Z, Lin SY, Chen Z, Li B, Yao G, Leake MC, Lo CJ, Bai
708 F (2019) ATP-Dependent Dynamic Protein Aggregation Regulates Bacterial Dormancy Depth
709 Critical for Antibiotic Tolerance. *Molecular cell* **73:** 143-156 e144
710
711 Pu Y, Zhao Z, Li Y, Zou J, Ma Q, Zhao Y, Ke Y, Zhu Y, Chen H, Baker MAB, Ge H, Sun Y,
712 Xie XS, Bai F (2016) Enhanced Efflux Activity Facilitates Drug Tolerance in Dormant Bacterial
713 Cells. *Molecular cell* **62:** 284-294
714
715 Ren Y, Wang C, Chen Z, Allan E, van der Mei HC, Busscher HJ (2018) Emergent
716 heterogeneous microenvironments in biofilms: substratum surface heterogeneity and bacterial
717 adhesion force-sensing. *FEMS Microbiol Rev* **42:** 259-272
718
719 Seviour T, Winnerdy FR, Wong LL, Shi X, Mugunthan S, Foo YH, Castaing R, Adav SS,
720 Subramoni S, Kohli GS, Shewan HM, Stokes JR, Rice SA, Phan AT, Kjelleberg S (2021) The
721 biofilm matrix scaffold of Pseudomonas aeruginosa contains G-quadruplex extracellular DNA
722 structures. *NPJ biofilms and microbiomes* **7:** 27

723
724 Shee C, Cox BD, Gu F, Luengas EM, Joshi MC, Chiu LY, Magnan D, Halliday JA, Frisch RL,
725 Gibson JL, Nehring RB, Do HG, Hernandez M, Li L, Herman C, Hastings PJ, Bates D, Harris
726 RS, Miller KM, Rosenberg SM (2013) Engineered proteins detect spontaneous DNA breakage in
727 human and bacterial cells. *eLife* **2:** e01222
728
729 Spoering AL, Lewis K (2001) Biofilms and planktonic cells of Pseudomonas aeruginosa have
730 similar resistance to killing by antimicrobials. *Journal of bacteriology* **183:** 6746-6751
731
732 Sung K, Park M, Chon J, Khan S (2022) Methods to grow and measure in vitro static biofilms.
733 *Encyclopedia of Infection and Immunity* **4:** 408-429
734
735 Tolker-Nielsen T (2015) Biofilm Development. *Microbiology spectrum* **3:** MB-0001-2014
736
737 Valentini M, Filloux A (2016) Biofilms and Cyclic di-GMP (c-di-GMP) Signaling: Lessons from
738 Pseudomonas aeruginosa and Other Bacteria. *The Journal of biological chemistry* **291:**
739 12547-12555
740
741 Vrabioiu AM, Berg HC (2022) Signaling events that occur when cells of Escherichia coli
742 encounter a glass surface. *Proceedings of the National Academy of Sciences of the United States
743 of America* **119**
744
745 Walters MC, 3rd, Roe F, Bugnicourt A, Franklin MJ, Stewart PS (2003) Contributions of
746 antibiotic penetration, oxygen limitation, and low metabolic activity to tolerance of
747 Pseudomonas aeruginosa biofilms to ciprofloxacin and tobramycin. *Antimicrob Agents
748 Chemother* **47:** 317-323
749
750 Waters EM, Rowe SE, O'Gara JP, Conlon BP (2016) Convergence of Staphylococcus aureus
751 Persister and Biofilm Research: Can Biofilms Be Defined as Communities of Adherent Persister
752 Cells? *PLoS pathogens* **12:** e1006012
753
754 Whitby MC, Vincent SD, Lloyd RG (1994) Branch migration of Holliday junctions:
755 identification of RecG protein as a junction specific DNA helicase. *The EMBO journal* **13:**
756 5220-5228
757
758 Yan J, Bassler BL (2019) Surviving as a Community: Antibiotic Tolerance and Persistence in
759 Bacterial Biofilms. *Cell Host Microbe* **26:** 15-21
760
761 Yan J, Moreau A, Khodaparast S, Perazzo A, Feng J, Fei C, Mao S, Mukherjee S, Kosmrlj A,
762 Wingreen NS, Bassler BL, Stone HA (2018) Bacterial Biofilm Material Properties Enable
763 Removal and Transfer by Capillary Peeling. *Advanced materials* **30:** e1804153
764
765 Zhao J, Wang Q, Li M, Heijstra BD, Wang S, Liang Q, Qi Q (2013) Escherichia coli toxin gene
766 hipA affects biofilm formation and DNA release. *Microbiology* **159:** 633-640
767
768 Zhou Y, Liao H, Pei L, Pu Y (2023) Combatting persister cells: The daunting task in
769 post-antibiotics era. *Cell insight* **2:** 100104

770
771

772
773

Figure 1. Cell adhesion promotes elevation of c-di-GMP levels and persister formation.

(A) Schematic overview of the static biofilm system designed to investigate intrinsic factors influencing persister cell formation during the early stages of biofilm development. (B-D) Left panel: the biphasic killing curves of planktonic cell subpopulation and surface-adherent cell subpopulation from static biofilm system at the culture time of 0.5 h (B), 2h (C) or 3.5 h (D), respectively (n=4). Right panel: Frequency distribution of c-di-GMP levels (R^{-1} score) within the surface-adherent subpopulation from the same static biofilm system and time points. Insert fluorescent images provide representative illustrations of c-di-GMP levels in adherent cells, measured using the c-di-GMP sensor system (Fig. S1B) integrated into *E. coli* cells. Scale bar, 2 μm. R^{-1} score was determined using the fluorescent intensity of mVenus^{NB} and mScarlet-I in the system. (E) Spatial heterogeneity in c-di-GMP levels (R^{-1} score) within a 24-hour static biofilm captured through confocal microscopy. (F) The persister ratio in planktonic cell subpopulation and surface-adherent

cell subpopulation collected from the 24-hour static biofilms (ampicillin, 150 μg/ml, 3-hour at 37°C with
shaking at 220 rpm) (n=3). (G-H) The persister ratio (n=3) (H) in High-R^{-1} or Low-R^{-1} subpopulations
sorted from the 24-hour static biofilms (ampicillin, 150 μg/ml, 3-hour at 37°C with shaking at 220 rpm) (G).
(I) Representative time-lapse images of the persister assay using cells with different R^{-1} values. The cells
resuspended from 24-hour a static biofilm were imaged on a gel pad, and subsequently treated with LB
broth containing 150 μg/ml ampicillin, imaged over 6-hour at 35°C and then allowed to resuscitate in fresh
LB. Scale bar, 5/10 μm, as indicated. Error bars represent standard deviations of biological replicates.
Significance was ascertained by two-tailed Student's t test. Statistical significance is denoted as *P <0.05,
**P < 0.01, ***, P<0.001, ****, P<0.0001.

796
797

Figure 2. High c-di-GMP levels promote persister formation.

(A) Schematic overview illustrating the manipulation of c-di-GMP levels in surface-adherent cell subpopulations for persister ratio determination. (B-C, E) Frequency distribution of c-di-GMP levels (R^{-1} score) within the surface-adherent subpopulation from strains overexpressing either PdeH (pBAD::*pdeH*) (B) or DgcZ (pBAD::*dgcZ*) (E), along with a vector control (pBAD::vector) (C). Cells preparation followed the procedure depicted in (A). Insert fluorescent images provide representative illustrations of c-di-GMP levels in adherent cells, measured using the c-di-GMP sensor system (Fig. S1B) integrated into different stains. Scale bar, 2 μm. (D) Biphasic killing curves of the surface-adherent subpopulations from (A) (n=3). Error bars represent standard deviations of biological replicates. Significance was ascertained by two-tailed Student's t test. Statistical significance is denoted as *P <0.05, **P < 0.01, ***P < 0.001.

809
810

Figure. 3. HipH and c-di-GMP a pair of toxin-antitoxin system.

(A) Determination of persister ratios in the strains overexpressing DgcZ or DgcZ-m (G206A, G207A, E208Q), with 0.002% arabinose induction for 2 h in the pBAD plasmid. Persister counting was performed following antibiotic killing (ampicillin, 150 μg/ml, 3-hour at 37°C with shaking at 220 rpm) (n=3). (B) Schematic overview illustrating the screening of downstream targets of c-di-GMP by using *E. coli* Keio Knockout Collection, the screen was conducted in the presence of high levels of c-di-GMP by inducing the expression of DgcZ with 0.002% arabinose for 2 h. (C) Identification of HipH as the downstream target of c-di-GMP in persister formation. Left panel: persister assay performed without high levels of c-di-GMP (pBAD-vector); Right panel: persister assay performed in the presence of high levels of c-di-GMP (pBAD-*dgcZ*) (ampicillin, 150 μg/ml, 3-hour at 37°C with shaking at 220 rpm) (n=9). (D) Schematic overview of clinical UPEC16 isolation, genetic modification, and *in vitro* or *in vivo* biofilm culture. (E) Antibiotic killing (ampicillin, 150 μg/ml, 3-hour at 37°C with shaking at 220 rpm) and persister counting assay for *in vitro* biofilm cells (n=3). (F) Antibiotic killing (ampicillin, 150 μg/ml, 3-hour at 37°C with shaking at 220 rpm) and persister counting assay for *in vivo* IBC cells (n=9 for WT, n=11 for Δ*hipH*). (G) Relative expression level of *hipH* under different levels of cellular c-di-GMP, determined by RT-qPCR and normalized to the levels of 16S rRNA. DgcZ and DgcZ-m were induced by 0.002% arabinose for 2 h in the pBAD plasmid (n=3). (H) Fluorescence intensity of HipH fused to BFP under varying levels of cellular

c-di-GMP. DgcZ was induced by 0.002% arabinose for 2 hours using the pBAD plasmid. The inserted

fluorescent images provide representative illustrations of HipH-BFP. (I-K) Detection of the ratio of cells

with growth potential (I), VBNCs or death cells (J) and persisters (K) in strains overexpressing HipH alone,

or overexpressing HipH and DosC together, or overexpressing HipH and DgcZ together, induced by 0.002%

arabinose for 4 hours. Persister counting performed followed antibiotic killing (ampicillin, 150 μg/ml,

3-hour at 37°C with shaking at 220 rpm) (n=3). (L) Schematic representation of the dormancy depth

controlled by HipH and c-di-GMP, and the relationship with cell state. Error bars represent standard

deviations of biological replicates. Significance was ascertained by two-tailed Student's t test (ACEFGHK)

or two-tailed Two-way ANOVA (I-J). Statistical significance is denoted as $*P <0.05$, $**P < 0.01$, ***,

$P<0.001$; ****, $P<0.0001$.

Figure 4. HipH is a deoxyribonuclease exerting genotoxic effects.

(A) Structure analysis of HipH protein and Schematic representation of truncated mutants illustrating key
structural domains. (B-C) Detection of the ratio of cells with growth potential (B) and VBNCs or death cells
(C) in strains overexpressing different truncated HipH mutants, induced by 0.002% arabinose for 3 h (n=3).
(D) Subcellular localization of HipH. HipH was fused with mScarlet-I and induced by 0.002% arabinose
for 2 hours. The gDNA was stained with Hoechst 33258 for 15 minutes at room temperature. Scale bar, 1
μm. (E-G) Left panel: *in vitro* DNA cleavage assay using 500 ng supercoiled plasmid DNA, (E) with
varying concentrations of HipH-GFP (0, 25, 50, 100, 200 or 400 nM), with purified GFP protein as control;
(F) with different durations of reaction time (15, 30, 45 and 60 minutes); (G) with different reaction
temperature (25, 30, 37 and 42°C). If not specified, 200 nM of HipH-GFP was added to each reaction. "NC",
negative control, indicates supercoiled plasmid without addition of HipH-GFP. Right panel: illustration of
the relationship between the length of DNA fragments and their ratio to total DNA, as indicated in the
corresponding upper panel. SSB, single-strand break; DSB, double strand break. Error bars represent
standard deviations of biological replicates. Significance was ascertained by two-tailed Two-way ANOVA.
Statistical significance is denoted as *P <0.05, **P < 0.01.

857
858

Figure 5. c-di-GMP acts as an antitoxin to repress HipH DNA Cleavage Activity.

(A) Schematic representation of the biotin-c-di-GMP pull-down assay for HipH protein. (B-E)

Identification of the interaction between c-di-GMP and HipH using the biotinylated c-di-GMP pull-down

assay. (B) Upper panel: 1.5 μM purified HipH-GFP-His and biotin-c-di-GMP were added in gradient

concentrations (30, 20, 10, 7.5, 5, 2.5, 0 μM). Bottom panel: the intensity of bound protein, obtained by

analyzing the immunoblot shown in the upper panel using ImageJ, relative to the concentrations of
c-di-GMP used in the pull-down assays. (C) 0.3 μM purified HipH-GFP-His and 20 μM biotin-c-di-GMP or
biotin were added. (D) Purified HipH-GFP-His or GFP-His was added in gradient concentrations (1.5, 0.75,
0.375, 0.19, 0 μM), and 20 μM biotin-c-di-GMP was added. 0.5 μg protein was used as the input sample. (E)
2 mM c-di-GMP, cAMP, or GTP were used to elute HipH-GFP-His from SA beads, as well as
HipH-GFP-His/biotin-c-di-GMP complexes prepared from the pull-down reaction mix containing 1.5 μM
purified HipH-GFP-His and 20 μM biotin-c-di-GMP. (F-I) *In vitro* DNA cleavage assay using 500 ng
supercoiled plasmid DNA (F-G) or relaxed plasmid DNA (H-I). "+" indicated 1.67 mM c-di-GMP and "++"
indicated 5 mM c-di-GMP. 200 nM HipH-GFP-His or GFP-His was used in Cleavage reaction at 37°C for
1 h. (G) and (I) represent the relationship between the length of DNA fragments and their ratio to total DNA,
as shown in B and D, respectively. SSB: single-strand break (relaxed plasmid DNA); DSB: double-strand
break. (J) Statistic analysis of the ratio of cells with Gam-GFP foci in different strains: overexpressing
HipH alone, or HipH with DosC, or HipH with DgcZ (n=3). (K) Representative images of cells expressing
Gam-GFP in the aforementioned different strains. Scale bar, 5 μm. Error bars represent standard deviations
of biological replicates. Significance was ascertained by two-tailed Two-way ANOVA. Statistical
significance is denoted as *P <0.05, **P < 0.01.

881
882

883 **Tables and their legends**

884 **Table S1. Key resources table.**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse Anti-His-tag-mAb	MBL	Cat#D291-3; RRID: AB_10597733
HRP-Goat Anti-Mouse IgG (H+L)	ABclonal	Cat#AS003; RRID: AB_2769851
Strains and plasmids		
Escherichia coli strain MG1655	Yale Genetic Stock Center	CGSC#6300
Escherichia coli strain BW25113	Yale Genetic Stock Center	CGSC#7636
BW25113 KEIO library	Dharmacon (GE life sciences)	Cat#OEC4988
UroPathogenic *Escherichia coli* UPEC16	First Affiliated Hospital of Kunming Medical University	N/A
UPEC16 Δ*hipH*	This paper	N/A
UPEC16 Δ*ara*	This paper	N/A
UPEC16 p15A::c-di-GMP-sensor	This paper	N/A
UPEC16 Δ*ara* pBAD::vector	This paper	N/A
UPEC16 Δ*ara* pBAD::*pdeH*	This paper	N/A
UPEC16 Δ*ara* pBAD::*dgcZ*	This paper	N/A
UPEC16 p15A::c-di-GMP-sensor P-*hipH-bfp*	This paper	N/A
MG1655 p15A::c-di-GMP-sensor	This paper	N/A
MG1655 Δ*ara* pBAD::vector p15A::c-di-GMP-sensor	This paper	N/A
MG1655 Δ*ara* pBAD::*dosC* p15A::c-di-GMP-sensor	This paper	N/A
MG1655 Δ*ara* pBAD::*dgcZ* p15A::c-di-GMP-sensor	This paper	N/A
MG1655 Δ*ara hipH-bfp* pBAD::vector	This paper	N/A
MG1655 Δ*ara hipH-bfp* pBAD::*dgcZ*	This paper	N/A
MG1655 Δ*ara* pBAD::*dgcZ*-m	This paper	N/A
MG1655 Δ*ara* pBAD::*hipH*-His (or *hipH* mutants)	This paper	N/A
MG1655 Δ*ara* pBAD::*hipH*-mScarlet-I	This paper	N/A
MG1655 Δ*ara* pBAD::*gfp*-His	This paper	N/A
MG1655 Δ*ara* pBAD::*hipB*-His	This paper	N/A
MG1655 Δ*ara* pBAD::*mCherry*-His	This paper	N/A
MG1655 Δ*ara* pBAD::Vector, p15A::*gam*-GFP & *hipH*	This paper	N/A
MG1655 Δ*ara* pBAD::*dosC*, p15A::*gam*-GFP & *hipH*	This paper	N/A
MG1655 Δ*ara* pBAD::*dgcZ*, p15A::*gam*-GFP & *hipH*	This paper	N/A
MG1655 Δ*ara* pBAD::*hipH* pT5::vector	This paper	N/A
MG1655 Δ*ara* pBAD::*hipH* pT5::*recG*	This paper	N/A
BW25113 pBAD::*hipH*	This paper	N/A
BW25113 Promoter-rhaB-*hipH* pBAD::*dgcZ*	This paper	N/A
BW25113 Δ*hipH* pBAD::*dgcZ*	This paper	N/A
BW25113 Δ*recA* pBAD::*hipH*	This paper	N/A
Chemicals, peptides, and recombinant proteins		
Cycle di-GMP	APExBIO	Cat#B7839

2'-Biotin-16-c-di-GMP	BIOLOG	Cat#B098
HipH-GFP-His	This paper	N/A
GFP-His	This paper	N/A
Ampicillin	Sangon Biotech	Cat#A610028
Chloramphenicol	Sangon Biotech	Cat#A600118
Gentamicin	Sangon Biotech	Cat#A620217
Kanamycin	Sangon Biotech	Cat#A600286
Arabinose	Sigma	Cat#V900920
IPTG	Sangon Biotech	Cat# A600168
Anhydrotetracycline	MedChemExpress	Cat#HY-118660
qPCR Primers		
recA-QP1 5-CTGCTGATCTTCATCAACCA-3	This paper	N/A
recA-QP2 5-AACAGAGGCGTAGAATTTCAG-3	This paper	N/A
sulA-QP1 5-GGGCTTATCAGTGAAGTTGTC-3	This paper	N/A
sulA-QP2 5-GGCTAATCTGCATTACTTTCGT-3	This paper	N/A
tisB-QP1 5-ATGAACCTGGTGGATATCGC-3	This paper	N/A
tisB-QP2 5-TTACTTCAGGTATTTCAGAACAGC-3	This paper	N/A
hipH-QP1 5-GATTACTGCACAACACCCTG-3	This paper	N/A
hipH-QP2 5-GAAGAACCCACAATTTCTCCTG-3	This paper	N/A
Software and algorithms		
Fiji	Schindelin et al., 2012	https://fiji.sc/; RRID: SCR_002285
FlowJo	Treestar, Inc.	https://www.flowjo.com/

Fig. S1. A c-di-GMP biosensor was used in UPEC16 strain.

(A) The phylogenetic analysis has revealed that UPEC16 shares close genetic relatedness with
well-characterized UPEC reference strains such as CFT073 and UTI89. (B) Schematic overview of
c-di-GMP sensor. The design of c-di-GMP biosensor is based on fluorescent protein mVenus^{NB}, with MrkH
inserted between residues Y145 and N146. Conformational changes in MrkH, induced upon binding to
c-di-GMP, result in reduced fluorescence of the c-di-GMP sensor. A red fluorescent protein mScarlet-I is
co-expressed under the regulation of the *proC* promoter. (C-E) Frequency distribution of c-di-GMP levels
(R^{-1} score) within the planktonic subpopulation from the static biofilm system harboring c-di-GMP
biosensor at 0.5 h (C), 2 h (D), 3.5 h (E).

900
901

Fig. S2. UPEC16 infection of mouse bladder.

(A) Quantification of intracellular bacterial communities (IBC) count per slide within the epithelial cells
lining the bladder lumen infected by UPEC16-WT or UPEC16-Δ*hipH*. (n=6). (B) Histological
examinations of H&E stained UPEC16-WT or UPEC16-Δ*hipH* infected bladder tissue sections.
Representative images of IBC were indicated by red arrows.

908
909

Fig. S3. The correlation between the levels of c-di-GMP and HipH.

(A) Representative fluorescent images showing HipH-BFP fluorescence and c-di-GMP level R^{-1}. UPEC16
strain with the c-di-GMP sensing system and BFP-labeled HipH were cultured statically five-hours in
microtiter dishes. (B) Correlation analysis between HipH-BFP fluorescence values and c-di-GMP level R^{-1}
scores from (A). (C) Schematic representation of the promoter modification of the *hipH* gene. (D) Relative
expression level of *hipH* in the Promoter-*rhaB-hipH* strain under varying levels of cellular c-di-GMP,
determined by RT-qPCR and normalized to the levels of 16S rRNA. DgcZ was induced by 0.002%
arabinose for 2 hours using the pBAD plasmid (n=3). (E) Determination of persister ratios in the strains
overexpressing DgcZ with 0.002% arabinose induction for 2 hours using the pBAD plasmid. Persister
counting was performed following antibiotic treatment (ampicillin, 150 μg/ml, 3 hours at 37°C with
shaking at 220 rpm) (n=3). (F) Determination of relative cellular concentrations of c-di-GMP by c-di-GMP
sensing system in cells overexpressing DosC or DgcZ, with 0.002% arabinose induction for 2 h. Error bars

923 represent standard deviations of biological replicates. Significance was ascertained by two-tailed Student's
924 t test. Statistical significance is denoted as ****, P<0.0001.

Fig. S4. HipH induced genome instability.

(A) The cells length was detected in MG1655 strains which overexpressed HipH induced by 0.002% arabinose for 2 h under a light microscope. (B) The relative mRNA level of some SOS response genes, *recA*, *sulA* and *tisB*, were detected by qPCR in MG1655 strains which overexpressed HipH or HipH-ΔHTH induced by 0.002% arabinose for 3 h. The mRNA level was normalized by 16S rRNA level (n=3). (C) The ratio of cells with growth potential in different strains, WT: overexpressing HipH in BW25113 WT. Δ*recA*: overexpressing HipH in BW25113 Δ*recA* strain (n=3). HipH was induced by 0.002% arabinose for 2 h. (D) The ratio of cells with growth potential were detected in MG1655 strains which overexpressed HipH induced by 0.002% arabinose and overexpressed RecG induced by 1 mM IPTG for 2 h (n=3). Error bars represent standard deviations of biological replicates. Significance was ascertained by two-tailed Student's t test (A-B) or two-tailed Two-way ANOVA (C-D). Statistical significance is denoted as *P <0.05, **P < 0.01. ***, P<0.001; ****, P<0.0001.

940
941

Fig. S5. c-di-GMP binding inhibits HipH DNA Cleavage Activity.

(A) Lysates from cells overexpressing HipH-His or mCherry-His was added in gradient concentrations (0,
15, 30, 60, 120, 240 μg), with biotin-c-di-GMP was added at a constant concentration (8 μM). (B) Lysates
from cells overexpressing HipH-His was added at a constant concentration (240 μg), and biotin-c-di-GMP
was added in gradient concentrations (0, 0.5, 1, 2, 4, 8 μM). (C) Left panel: In vitro DNA cleavage assay
using 500 ng of supercoiled plasmid DNA. A concentration of 200 nM HipH-GFP-His was employed in the
cleavage reaction at 37°C for 1 hour. Subsequently, 5 mM c-di-GMP, 2.5 mM cAMP, 5 mM GTP, or 200
nM HipB-His were added to the cleavage reaction. Right panel: Illustration of the relationship between the
length of DNA fragments and their ratio to total DNA.

A

Cell adherence	Static growth — 2 h
Arabinose induction	Fresh LB with **arabinose** — Static growth 1.5 h
Adherent cell collection	
Persister counting assay	Ampicillin killing

B pBAD::*pdeH*

Frequency vs c-di-GMP level (R⁻¹)

Median R⁻¹=0.75
n=436

C pBAD::vector

Frequency vs c-di-GMP level (R⁻¹)

Median R⁻¹=0.98
n=476

D

Survival ratio (log10) vs killing time (h)

- pBAD::*pdeH*
- pBAD::Vector
- pBAD::*dgcZ*

ns
***
***
**
**
*

E pBAD::*dgcZ*

Frequency vs c-di-GMP level (R⁻¹)

Median R⁻¹=1.72
n=364

A Persister Ratio (log10) — Vector, DgcZ, DgcZ-m; **** ****

B pBAD::dgcZ → KEIO library → Planktonic cells in exponential phase → Persister assay

C Planktonic culture — Persister Ratio (log10); WT, ΔhipA, ΔhipH; pBAD::vector, pBAD::dgcZ; ****

D Bladder, UPEC, Gene knockout, ΔhipH, Static biofilm, Transurethrally infection, 2×10^8 CFU Twice, Washing, Ultrasonic peeling, For persister assay

G Relative mRNA level of hipH — Vector, DgcZ, DgcZ-m; ** ***

H HipH-Bfp — Fluorescence intensity; Vector n=206, DgcZ n=195; Vector, DgcZ (2 μm)

I Cells with growth potential (%) — Vector, HipH, HipH+DosC, HipH+DgcZ; * *

J Ratio of VBNCs or Death cells (%) — VBNC, Death; Vector, HipH, HipH+DosC, HipH+DgcZ; * *

K Persister Ratio (log10) — Vector, HipH, HipH+DosC, HipH+DgcZ; ** *

L Growth, Persister, VBNC, Death; Dormancy depth; WT, HipH, HipH & c-di-GMP

E Biofilm *in vitro* — Persister Ratio (log10); UPEC16-WT, UPEC16-ΔhipH; **

F IBC *in vivo* — Persister Ratio (log10); UPEC16-WT, UPEC16-ΔhipH; *

A

WT — *hipH* gene

Δ1 1-32 : HTH-domain

Δ2 1-135 : N-domain

Δ3 190-215 : gly loop, autophosphorylation site

Δ4 222-229 : activation loop

Δ5 340-372 : kinase catalytic motif

B

Cells with growth potential (%)

Vector, HipH-WT, HipH-Δ1, HipH-Δ2, HipH-Δ3, HipH-Δ4, HipH-Δ5

C

Cells without growth potential (%)

VBNC, Death

Vector, HipH-WT, HipH-Δ1, HipH-Δ2, HipH-Δ3, HipH-Δ4, HipH-Δ5

D

HipH-mScarlet-I   gDNA   Merge

1 μm

E

GFP (nM)   HipH-GFP (nM)

200 0 25 50 100 200 400

Supercoiled   DSB   SSB

Frequency vs DNA fregmengts length (bp)

— 400 nM
— 200 nM
— 100 nM
— 50 nM
— 25 nM

F

Reaction time (min)

NC 15 30 45 60

Supercoiled   DSB   SSB

Frequency vs DNA fregmengts length (bp)

— 15 min
— 30 min
— 45 min
— 60 min

G

Reaction temperature (°C)

NC 25 30 37 42

Supercoiled   DSB   SSB

Frequency vs DNA fregmengts length (bp)

— 25 °C
— 30 °C
— 37 °C
— 42 °C

A

0.003

CFT073
UPEC16
UTI89
536
AI39
CE10
SMS-3-5
EDL933
CB9615
K12_MG1655
BL21DE3
SE11
E24377A

B2
D
E
A
B1

B

Promoter ProC — mScarlet-I — mVenus^NB-MrkH

$$c\text{-di-GMP} \propto R^{-1} = \left(\frac{mVenus^{NB}\text{-MrkH}}{mScarlet\text{-I}}\right) - 1$$

mVenus^NB- MrkH   +   c-di-GMP   =   mVenus^NB-MrkH with c-di-GMP

C

median=0.34 n=137
planktonic cell 0.5 h

Frequency
c-di-GMP level (R^{-1})

D

median=0.35 n=145
planktonic cell 2 h

Frequency
c-di-GMP level (R^{-1})

E

median=0.31 n=71
planktonic cell 3.5 h

Frequency
c-di-GMP level (R^{-1})

A

IBC count per slide

ns

UPEC16-WT   UPEC16-Δ*hipH*

B

50 μm   UPEC16-WT

50 μm   UPEC16-Δ*hipH*

A

c-di-GMP level R⁻¹

HipH-BFP

2 μm

2 μm

B

r = 0.694
n=147
P<0.0001

c-di-GMP sensor R⁻¹

HipH-BFP fluorescence value

C

Promoter-*hipH*   HipH

Promoter-*rhaB*

D

Promoter-*rhaB-hipH*

Relative mRNA level of *hipH*

ns

Vector   DgcZ

E

pBAD::*dgcZ*

Persister Ratio (log10)

***

***

WT   Δ*hipH*   Promoter-*rhaB-hipH*

F

Vector

Fold change=0.87

c-di-GMP sensor R⁻¹

n=135   n=112

0   0.002%

Ara concentration

DosC

Fold change=2.72

c-di-GMP sensor R⁻¹

n=77

n=85

0   0.002%

Ara concentration

DgcZ

Fold change=4.22

c-di-GMP sensor R⁻¹

n=118   n=76

0   0.002%

Ara concentration

A

HipH-His

mCherry-His

Protein

Biotin-c-di-GMP   Input

B

Biotin-c-di-GMP

Protein

HipH-His

C

	1	2	3	4	5	6
pBR322	+	+	+	+	+	+
HipH-GFP	-	+	+	+	+	+
c-di-GMP	-	-	+	-	-	-
cAMP	-	-	-	+	-	-
GTP	-	-	-	-	+	-
HipB	-	-	-	-	-	+