Nitrogen starvation response in hornworts and liverworts provides little evidence for complex priming to the cyanobiont

Yuling Yue[1,2], Gaurav Sablok[3,4], Anna Neubauer[1,2], Jaakko Hyvönen[3], Péter Szövényi[1,2]

1 Department of Systematic and Evolutionary Botany, University of Zurich, Switzerland, 8008, Zollikerstrasse 107

2 Zurich-Basel Plant Science Center, Zurich, Switzerland

3 Finnish Museum of National History (Botany), University of Helsinki, 00100 Helsinki, Pohjoinen Rautatiekatu 13, Finland

4 Institute for Biochemistry and Biology, University of Potsdam, 14476 Potsdam-Golm, Karl-Liebknecht-Str. 24-25, Germany

Yuling Yue: yuling.yue@uzh.ch, 0009-0009-3210-8511

Gaurav Sablok : gaurav.sablok@uni-potsdam.de , 0000-0002-4157-9405

Anna Neubauer: anna.neubauer@uzh.ch, 0000-0002-5102-2491

Jaakko Hyvönen: jaakko.hyvonen@helsinki.fi, 0000-0001-7559-8295

Correspoding author: *Péter Szövényi: peter.szoevenyi@uzh.ch, 0000-0002-0324-4639*

Date of submission: 20.06.2024

Number of figures: 9
Number of tables: 0

Word count: 6055
Supplementary Tables: 24
Supplementary Figures: 10

Runing title: nitrogen starvation and plant-cyanobacteria symbiosis

Highlight: Our results suggest that the most critical step in establishing plant-cyanobacteria interactions using non-host plants is the attraction of the cyanobiont. This finding has significant impact on crop engineering.

37 ***Keywords:*** starvation response, plant-cyanobacteria symbiosis, nitrogen uptake, land plants,
38 evolution, crop plants, engineering

39
40
41

Abstract

Mutualistic plant-microbe symbiotic interactions are thought to have evolved from a loose
association between host plants and microbes when nutrients are limited. Therefore, the
molecular network enabling intimate mutualistic plant-microbe symbioses may have evolved
from a nutrient starvation response shared by all land plants. While the molecular link between
nutrient status and symbiotic interaction is well-established, it remains poorly understood in
some systems. This is especially true for the symbiotic associations between plants and
cyanobacteria.

To test the conservation of the starvation network across land plants as well as to investigate the
link between nutrient starvation and symbiosis initiation in the plant-cyanobacteria symbiosis,
here we explore the transcriptional responses to nutrient starvation in two non-vascular plant
species, a hornwort *Anthoceros agrestis* and a liverwort *Blasia pusilla*, forming plant-
cyanobacteria endophytic symbioses. We observe a deep conservation of the systemic starvation
response across land plants. However, very few if any components of the starvation network
appear to be specific to cyanobacteria hosting plants, providing little evidence for extensive and
specific priming to the cyanobiont. Moreover, we found that some bioactive molecules known to
be important in initiating the plant-mycorrhiza and nodule-forming bacteria symbioses, may also
have a similar role in plant-cyanobacteria symbioses.

Introduction

It is hypothesized that the evolution of intimate mutualistic symbiotic interactions between
plants and microbes occurred in multiple consecutive steps with the very first step entailing the

capability of the host plant to attract microbes and initiate the symbiotic interaction in nutrient
poor environments (Delaux and Schornack, 2021; Isidra-Arellano et al., 2021). Therefore, it has
been proposed that the molecular network enabling and controlling intimate plant-microbe
symbiotic interactions could have evolved from a general starvation response shared by all
embryophytes (Isidra-Arellano *et al.*, 2021; Shi *et al.*, 2021, 2022). Indeed, it has been observed
for decades that the establishment of plant-arbuscular mycorrhizal fungi (AMF) and plant- nodule
forming bacteria symbiotic interactions can only be initiated under nutrient poor conditions.
More specifically, increasing the concentration of phosphorus inhibits mycorrhizal infection of
host plants (Balzergue et al., 2013). Similarly, nodule formation is triggered under nitrogen but
also phosphorus starvation (Lu et al., 2020; Nguyen et al., 2021). This dependence is regulated
by multiple molecular switches linking the genetic networks of the nutrient status of the host
plant with those involved in the initiation and establishment of the symbiotic interaction (Shi et
al., 2021; Jarratt-Barnham et al., 2022). For instance, plants able to form symbiotic interaction
with AMF or nodule-forming bacteria release various compounds to the soil when kept under
nutrient poor conditions. The best studied of these compounds are strigolactones and flavonoids
(Misson et al., 2005; Hernández et al., 2007). Strigolactones are crucial for the initiation of plant-
AMF symbioses by triggering the germination of fungal spores and growth of hyphae (Akiyama
et al., 2005). Similarly, flavonoids released by the plant host are necessary to attract nodule
forming bacteria by the plant host and to initiate the nodule-forming legume-bacteria interaction
(Abdel-Lateif et al., 2012). Furthermore, master regulators of the phosphate/nitrogen starvation
network have been shown to directly regulate many genes including members of the common
symbiosis signaling pathway (CSP) required to prepare the host plant for accepting the symbiont
including a specialized colonization interface and adjusted immune reaction (Shi et al., 2021).
These observations are in line with the hypothesis that an abiotic nutrient starvation response is
potentially conserved across embryophytes and might have been expanded to initiate and
regulate plant-soil microbe interactions. Nevertheless, this hypothesis remains to be tested
because information on the consequences of nutrient starvation is mainly available for vascular
plants but missing for most non-vascular plant lineages.
103
While the molecular link between nutrient status of the host plant and establishment, as well as
maintenance of the symbiotic interaction is well known in the plant-AMF and nodule forming
bacteria symbioses, it is poorly understood in other plant-microbe symbiotic interactions. Some
plants form endophytic, extracellular symbioses with cyanobacteria which are also under the
control of the nutrient status of the host plant (Meeks, 1998; Xu and Wang, 2023; Álvarez et al.,
2023b). On one hand, endophytic plant-cyanobacteria symbioses are similar to the plant-AMF
and nodule forming bacteria symbioses because nitrogen starvation is necessary for the
reconstitution of the symbiotic interaction in vitro as well as to maintain cyanobacterial nitrogen
fixation (Enderlin and Meeks, 1983; Kimura and Nakano, 1990; CAMPBELL and MEEKS, 1992).

More specifically, upon nitrogen starvation the host plant is known to release the so-called
hormogonia inducing factor (HIF), mobilizing and attracting the cyanobiont (Splitt and Risser,
2016; Nishizuka and Hashidoko, 2018; Hashidoko et al., 2019; Alvarenga et al., 2022). This
response resembles that of the intracellular plant-AMF and -nodule forming bacteria symbioses
in which flavonoids and strigolactones are secreted and sensed by the microbial partner. On the
other hand, plant-endophytic cyanobacteria differ from the well-investigated plant-microbe
symbioses (plant-AMF/-nodule-forming bacteria) in two important aspects. While most plant-
endophytic cyanobacteria symbioses are extracellular and the cavities of the host plant
containing the cyanobiont are present even in the absence of the cyanobiont (Meeks, 2003;
Gorelova and Baulina, 2009; Xu and Wang, 2023), plant-AMF and -nodule forming bacteria
symbioses are all intracellular and formation of host plant cellular structures essential for the
colonization of the microbial partner must be induced by nutrient starvation and by contact with
the microbial partner (Oldroyd, 2013; Ma and Chen, 2021). Therefore, nutrient starvation may
induce a less complex genetic network in plant-cyanobacteria systems mainly related to the
production of the HIF and immune system adjustment in contrast to intracellular symbioses
(plant-AMF/-nodule-forming bacteria). Nevertheless, this hypothesis remains to be tested
because the genetic network activated by starvation in plants forming endophytic symbiotic
interaction with cyanobacteria is poorly characterized (Chatterjee et al., 2022).
131
To investigate these questions, we studied the transcriptional response to starvation in two non-
vascular plant species representing ideal model systems of the plant-cyanobacteria symbioses
using time-series RNA-seq data. We starved the hornwort model *Anthoceors agrestis* and the
liverwort *Blasia pusilla*, with an estimated divergence time of 450 My representing two
independently evolved realizations of the plant-cyanobacteria endophytic symbioses (Meeks,
1998; Duggan et al., 2013; Szövényi et al., 2015; Álvarez et al., 2023b). Then we compared gene
expression changes associated with the starvation reaction in these two bryophyte species with
data available for vascular plants to identify convergent and divergent transcriptomic responses
(Fig. 1). We further used this data set to search for transcriptomic responses shared by the two
cyanobacteria symbiosis forming organisms but missing from bryophytes and vascular plants
unable to form symbiotic interaction, or only engaged in symbiosis with AMF and/or nodule-
forming bacteria.
144

Figure 1. Phylogenetic relationship of the species used in this study and their symbiotic partners. Species were selected to represent four groups: species establishing endophytic symbiosis with cyanobacteria, species establishing symbiosis with AMF, species establishing symbiosis with nodule-forming bacteria, and species unable to form symbiosis with any of the three symbiotic partners. Ellipses after each species refer to the type of symbiotic interaction they are involved with.

Materials and Methods

Plant and cyanobiont material

We used the hornwort species *A. agrestis* and the liverwort model species *B. pusilla* to carry out the starvation experiment. The *A. agrestis* isolate was established from a single spore of the BONN isolate using culture conditions and techniques described earlier (Frangedakis et al., 2021). Young gametophytes were grown and vegetatively propagated for 1-2 months on solid KNOP (Frangedakis et al., 2021) medium in Petri dishes prior to initiating liquid cultures. The *B. pusilla* isolate was kindly provided by Elke Dittman (University of Potsdam). *B. pusilla* gametophytes were also precultured in Petri dishes on KNOP medium under the very same conditions described for *A. agrestis*. Prior to the starvation experiment we established liquid cultures for both species by dissecting Petri dish grown gametophytes into 0.5 mm fragments using forceps. We put

gametophyte fragments into 200 ml liquid culture in a 500 ml flask containing BCD (Cove et al.,
2009) and BG11 (Enderlin and Meeks, 1983) medium for *A. agrestis* and *B. pusilla*, respectively.
To enhance plant growth, we kept the liquid cultures on an orbital shaker (150 rpm; SHKE2000-
1CE; Thermo Fisher Scientific, Waltham, Massachusetts, USA) at 23°C under continuous light (17–
20 µmol·s^{-1}·m^{-2}, as measured using an LM2-1137 light meter with a Quantum Q37008 sensor
[LI-COR Biosciences, Lincoln, Nebraska, USA]). Stock cultures of *A. agrestis* were subcultured in
fresh media every two weeks by filtering the tissues using a cell strainer (BioSwisstec Premium
Cell Strainer, 100 µm pore size). As cyanobiont, we used *Nostoc punctiforme* ATCC 29133. We
precultured *N. punctiforme* on solid BG11 medium in Petri dishes for 2-3 weeks and inoculated
50 ml BG11 in a 125 ml flask. These liquid cultures were kept under the same conditions as the
hornwort and the liverwort cultures.
179
180
181 ***Starvation experiment***
182
To start the starvation experiment, we collected two cell strainers (BioSwisstec Premium Cell
Strainer, 100 µm pore size) of plant tissue for each plant species (corresponding to a ~169 mg dry
weight) and transferred them into one 500 ml Erlenmeyer flask, in total we used 12 flasks. Six out
of the 12 flasks contained 200 ml fresh media without combined nitrogen (BCD-N/BG11-N,
"starvation flasks", details are provided in the Table S1), while the other six flasks were filled with
200 ml fresh media with combined nitrogen (BCD/BG11, "control flasks", details are provided in
the Table S1). To monitor plant response to starvation, we sampled thallus tissues from the
starvation flasks after 1,2,3,7,8,9, and 10 days and from the control flasks after 1,2,3,7, and 10
days of inoculation (details see in Table S1). On each date, we collected ~110 mg fresh weight of
plant tissue (corresponding to ~3.5 mg dry weight) in five biological replicates for both species.
199 ***RNA extraction and sequencing***
200
We immediately submerged collected thallus tissues of *A. agrestis* into RNA*later*® (Sigma-Aldrich,
product number: R0901-100ML) and stored them at 4°C overnight. On the next day, we removed
the tissues from the RNA*later* solution, flash-froze them in liquid nitrogen, and stored them until
RNA extraction at -20°C or -80°C in the freezer. For *B. pusilla*, we directly flash-freeze collected

thallus tissues in liquid nitrogen. To extract RNA, we grinded the plant tissue into a fine powder
with a pestle and a mortar in the presence of liquid nitrogen and processed the samples using
the Spectrum™ Plant Total RNA-Kit from Sigma-Aldrich (product number: STRN250), including on-
column DNA digestion using the on-column DNAse kit (Sigma-Aldrich, product number:
DNASE70-1SET). The Agilent 2100 bioanalyzer (Agilent, Santa Clara, CA) and nanodrop were used
to test the integrity, purity, and concentration of the RNA samples. For each sampling point we
kept the three biological replicates with the best RNA qualities. The 42 mRNA libraries for *B.*
pusilla and 30 mRNA libraries for *A. agrestis* were prepared using NEBNext® Ultra Directional RNA
Library Prep Kit and paired end sequencing was performed on Illumina NovaSeq6000 (2 × 150 bp)
by Novogene UK, Cambridge, London.
215
216

Draft assembly of the Blasia pusilla genome

In-vitro grown gametophytic cultures of *B. pusilla* on MS medium (Murashige & Skoog
Medium (MS1), 2008) were used for genomic DNA extraction using the plant-EZ DNA extraction
kit and were sequenced using PacBio-SMRT at the Institute of Biotechnology, University of
Helsinki, Finland. Reads mapped (using BLASR (Chaisson and Tesler, 2012)) to organelle and
cyanobacterial genomes were removed. In addition to the PacBio sequencing, short read
sequencing was done to support hybrid genome assembly and polishing. Short reads were
trimmed using Fastp (Chen et al., 2018) and assembled together with the filtered PacBio reads
using the hybrid genome assembler Masurca (Zimin et al., 2013). The hybrid genome assembly
was polished with cleaned short reads using Pilon (Walker et al., 2014) with a minimum read
depth of 10. Finally, we used blobtools v.1.1.1 (Laetsch and Blaxter, 2017), the NCBI nr database
and the average coverage of Illumina reads for each scaffold to remove scaffolds of contaminant
origin. After visual assessment we kept scaffolds with a taxonomic assignment of viridiplantae or
streptophyte. Scaffolds with other taxonomic assignments were discarded. Completeness and
redundancy of our assembled genome was assessed by searching translated peptide sequences
of the reconstructed gene set against the hidden Markov model of Benchmarking Universal
Single-Copy Orthologs BUSCO V.5 (Seppey et al., 2019) using the Viridiplantae and Embryophyta
datasets retrieved from OrthoDB v11 (Kuznetsov et al., 2023).

Creating a comprehensive transcript dataset for Blasia pusilla

We combined all RNA-seq data and the draft genome sequence to create reference transcripts
for *B. pusilla* which was used later for gene expression estimation. We generated a genome-
guided as well as a *de novo* TRINITY transcriptome assembly using all collected RNA-seq reads (42

RNA libraries) (Grabherr et al., 2011). We used the PASA pipeline (Haas et al., 2008) to combine
the *de novo* and genome-guided assemblies into a non-redundant set of transcripts and putative
genes. Using this approach, we created a transcript to gene mapping file and a non-redundant
set of transcripts. We ran all transcripts through Transdecoder
(https://github.com/TransDecoder/TransDecoder) to obtain their best ORF and peptide
translation. To reduce the number of potential transcripts and gene models, we discarded all
putative gene ids without at least one complete ORF prediction. To identify potential
contaminants in this filtered transcript file, we selected the longest ORF for each putative gene
and searched them against the Eggnog database in TRAPID 2.0 (Bucchini et al., 2021) and
assessed their taxonomic assignment. The initial run indicated that most of the contamination
concerned yeast. To remove these transcripts, we searched all transcripts against the full
transcriptome of *Saccharomyces cerevisisae* S288C (assembly R64) transcripts using blastn
version 2.12.0+ (Altschul et al., 1990) and discarded all transcripts passing the following filtering
criterium: -evalue ≤ 0.0001, similarity value ≥ 90%, and query coverage ≥ 80%. (Yeast
transcriptomics data was downloaded from here:
https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/146/045/GCF_000146045.2_R64/GCF_0001
46045.2_R64_rna.fna.gz) After that, we ran a further taxonomic assignment using TRAPID 2.0
and removed another smaller set of genes mapping to Viruses, Bacteria, Archeae, Opisthokonta,
and other Eukaryota. The final assembly had 22635 genes represented by 156239 transcripts.
262
The quality of the assembled transcripts was evaluated using three methods: (1) by assessing the
read content of the transcriptome; (2) by calculating the 'gene' Contig Nx Statistic; and (3) the
completeness of the transcript dataset. To assess the read content of the transcript assembly,
we counted the number of overall aligned read pairs to our assembled transcripts using Bowtie2
(Langmead and Salzberg, 2012). We computed the conventional Nx length statistic using the
Trinity toolkit utilities (TrinityStats.pl). Finally, we assessed the completeness of the transcript
dataset by BUSCO V.5 (Seppey et al., 2019) using both the Viridiplantae and Embryophyta
references retrieved from OrthoDB v11 (Kuznetsov et al., 2023).
271
272
273 ***Expression quantification and time-series analysis***
274
We applied the transcript quantification method in the mapping-based mode implemented in
Salmon 1.4.0 (Patro et al., 2017), the other parameters were set to default. As reference we used
the *B. pusilla* transcriptome set (assembled in this study), and the *A. agrestis* BONN cDNA
sequences (Li et al., 2020). After running Salmon, we imported the quantification files into R with
the tximport package (Soneson et al., 2015). The median of ratios method of normalization was
used to normalize expression estimates. Prior to further statistical analyses, we removed

transcripts from the data set for which the sum of the normalized expression counts over all
samples was < 10. We summarized gene expression values at the gene level for both species.
283
To gain insight into the overall qualitative change of gene expression associated with starvation
in *B. pusilla* and *A. agrestis*, we carried out principal component analysis (PCA) (Brand, 2013) on
normalized and standardized RNA-seq expression data. To identify genes significantly responding
to nitrogen starvation in *A. agrestis* and *B. pusilla* plants, we employed the approach provided in
maSigPro version 1.66.0 (Nueda et al., 2014), an algorithm modeling gene expression time series
data with GLMs (general linear models). We chose the backward selection algorithm with an
alpha value of 0.05, a Q-value threshold of 0.05 and a polynomial degree of 3. The expression
data comprised RNAseq data of seven and five time points for *B. pusilla* and *A. agrestis*
respectively. We clustered the significant (Q ≤ 0.05) DEGs into different profiles using the "hclust"
cluster method in maSigPro and the Mclust option estimating the optimal k, choosing
k.mclust=TRUE. Finally, we have tested different numbers of k to obtain a robust clustering of
expression profiles by comparing the clusters at each step of k with the previous ones using the
command "see.genes (get$sig.genes, k=...).".
297
298 ***Comparative transcriptomic analysis***
299
To compare the gene set reacting for nitrogen starvation in *A. agrestis* and *B. pusilla* and that of
vascular plants, we created orthogroups using Orthofinder v2.5.5 (Emms & Kelly, 2019) with
default options. We included symbiotic and and non-symbiotic bryophyte and vascular plant
species covering a range of taxonomic diversity increasing the accuracy of the orthofinder
algorithm (full species list is provided in Table S2). Furthermore, we collected transcriptomic data
in response to starvation for bryophytes and vascular plants able or unable to establish symbiotic
interaction with AM fungi, nodule-forming bacteria, cyanobacteria, or a combination of these
partners. More specifically, we obtained data for non-symbiotic species (*Marchantia*
polymorpha, Arabidopsis thaliana), species involved in interaction with AMF (*Marchantia*
paleacea, Triticum aestivum, Solanum lycopersicon, Brachypodium distachyon, Medicago sativa)
or nodule-forming bacteria (*Medicago sativa*). We retrieved the up and downregulated gene lists
for all these species but *M. polymorpha* from the supplementary material of the original
publications (see Table S3). For *M. polymorpha* we used DESeq2 (Love et al., 2014) method to
test for differential gene expression (details are provided in the Table S4). Finally, expression
information was added to the orthofinder results and orthogroups with convergent or divergent
expression responses were identified. For *A. agrestis* and *B. pusilla*, up and downregulated genes
were identified in the maSigPro analyses explained above. For the other species *M. paleacea, A.*
thaliana, B. distachyon we retrieved the up and downregulated set of genes as they were given
in the corresponding publications. For species such as *T. aestivum*, *M. sativa* and *S. lycopersium*

in which transcriptomic response was recorded in multiple tissue types, we did the following: we
selected the most significant differential expressed genes among all the tissues tested. If data for
different tissues were available and the reaction of genes was tissue-dependent, we selected the
greatest absolute log2 fold change value.
323
324
325   ***Functional enrichment***
326
To functionally identify and characterize the broad biological pathways and processes involved
in nitrogen starvation, we obtained GO and KEGG annotations for the *A. agrestis* and *B. pusilla*
gene set using Trapid 2.0 (see Table S5-S8). We carried out gene ontology and KEGG enrichment
analysis of up and downregulated genes using the Parent-Child Algorithm and a Fisher's exact
test implemented in the TopGO package 2.46.0 (Alexa, 2006). We also carried out manual
annotation of genes showing differential expression using the Plaza 5.0 database
(https://bioinformatics.psb.ugent.be/plaza/). We retrieved transcription factor annotations and
genes involved in hormone synthesis from the hornwort genome paper (Li et al., 2020).
Furthermore, we also processed the full proteome of both species using the PlantTFBD website
(https://planttfdb.gao-lab.org/)(Tian et al., 2020) to assign genes to transcription factor (TF)
families (see Table S9).
343   ## Results
344
345   ***Draft genome assembly of Blasia pusilla***
346
To aid our analyses, we generated a draft genome assembly for the *B. pusilla* with a combination
of short and long-read data (see Methods section). After removing potential contaminants with
Blobtools, our assembly consisted of 1456 scaffolds/contigs with a total length of ca. 347 Mbp
(see details in the Table S10 and the final genome assembly on Figshare) which fits to the genome
size estimated by previous c-value analyses (Renzaglia et al., 1995). Our results showed that the
blobtools filtered draft genome assembly was of good quality containing 71%/79.7%
(Embryophyta/Viridiplantae dataset) of the BUSCO genes. These BUSCO values are close to the
figures obtained for high-quality genome assemblies generated using long-read technologies
(*Marchanthia polymorpha*, *Physcomitrella patens*, *Pleurozium schreberi,* and *Funaria*
hygrometrica) (Bowman et al., 2017; Lang et al., 2018; Pederson et al., 2019; Kirbis et al., 2020).

Furthermore, over 96% of our RNA-seq reads could be mapped to the draft genome implying that
the coding part of the genome is well covered (Supplementary File 1).
359

Comprehensive transcriptomic data set

Initial annotation of the draft genome using braker1 and RNA-seq evidence resulted in a
considerable proportion of fragmented gene models. Furthermore, we observed a large number
of highly expressed gene models that were present in the transcriptome assemblies but missing
from the genome-based braker annotation. Therefore, we decided to carry out a genome-guided
as well as a *de novo* transcriptome assembly and combined their results. After contaminant
filtering (eg. yeast Viruses, Bacteria, Archeae, Opisthokonta, and other Eukaryota) and removing
gene models lacking at least one complete CDS sequence, the final assembly consisted of 22636
genes that were represented by 156239 transcripts. Completeness of the contamination filtered
gene set (longest ORF per gene) was superior to that obtained by conventional annotation of the
genome (see values above 71% and 79.7%) and captured 85.1% and 97.7% of the Embryophyta
and Viridiplantae BUSCO gene set, respectively (The final transcriptome assembly is available on
Figshare). The transcriptome assembly was of high quality based on all three statistics assessed
(Supplementary File 2).

Nitrogen starvation induces transcriptional reprogramming in both bryophytes investigated

Visualization of gene expression changes under nitrogen starvation over the time course of the
experiment (standardized and centralized PCA) indicated considerable transcriptional
reprogramming in both species (Fig. 2). More specifically, control and starvation samples formed
two well distinguishable groups in both species after seven days. While transcriptional
reprogramming occurred in both species, the temporal reactions to starvation differed
considerably. In the hornwort (*A. agrestis*) the difference between control and starvation
samples became considerable for the last two sampling points of the experiment. By contrast, in
the liverwort (*B. pusilla*) global transcriptional changes were already detectible after a single day
of nitrogen starvation. Quantitatively, we found that 20737 *A. agrestis* gene models were
expressed in at least one of the nitrogen starvation samples, which is about 80.5% of the 25837
currently annotated gene set (Li et al., 2020) (Table S11, S12). Similarly, of the total 22636 gene
models 22331 (98.6%) showed expression during nitrogen starvation in *B. pusilla* (Table S13, S14).
Our statistical analysis indicated (MaSiPro Q≤0.05) that a similar number of genes, 1522 (7.34%
of the expressed genes) and 1023 (4.58fig. 1% of the expressed genes), reacted to nitrogen
starvation in *A. agrestis* and *B. pusilla*, respectively (Fig. 3 and Fig. 4).

Figure 2. Principal component analysis (PCA) of overall transcriptomic response to nitrogen-
starvation in *A. agrestis* (36 samples) and B. *pusilla* (48 samples)*.* Each data point (triangle/dot)
refers to a sample. Case (solid dot): samples starved of combined nitrogen; Control (solid
triangle): grown in full media (see materials and methods); TP: sampling days after starvation.

402
403

Figure 3. Expression profile of genes experiencing significant response to nitrogen starvation in
A. agrestis. Each plot corresponds to a cluster of genes showing similar expression profiles over
the time-course. Dots correspond to the average normalized expression values over all genes of
a cluster. The three dots for each time point represent the three biological replicates. Gene
expressions in the nitrogen starved (in red, case) and control samples (in green, control) are
shown. Solid lines connect median expression values per time point. Dashed lines are fitted
regression curves with a maximum polynomial degree of 3.

Days after starvation

412
413
Figure 4. Expression profile of genes experiencing significant response to nitrogen starvation in
B. pusilla. Each plot corresponds to a cluster of genes showing similar expression profiles over
the time-course. Dots correspond to the average normalized expression values over all genes of
a cluster. The three dots for each time point represent the three biological replicates. Gene
expression in the nitrogen starved (in red, case) and control (in green, control) samples are
shown. Solid lines connect median expression values per time point. Dashed lines are fitted
regression curves with a maximum polynomial degree of 3.

A. agrestis and *B. pusilla* **show similar transcriptional responses to nitrogen starvation**

To learn more about the dynamics of gene expression change over the time course, we identified
genes showing similar temporal expression dynamics. We ran K-means clustering using a range
of K values (1-9) to find a subjective optimal K value. After visual inspection of group expression
profiles, we decided to use K=4 which provided a good balance between cluster distinctness and
within-cluster uniformity (see results for K=2-9 in Fig. S1-S2). We did so for the two species
separately.

Anthoceros agrestis

In the hornwort *A. agrestis*, we found four major groups of genes showing well distinguishable
expression dynamics (see Table S12) and the enriched GO terms for all four clusters are displayed
in Table S15, Fig. S3-S6 and the enriched go terms for up/downregulated genes are shown in Fig.
5. Fig. 5A shows the enriched GO terms for upregulated genes of cluster 2 and Fig. 5B for clusters

1,3 and 4. 657 genes (cluster 2) showed gradually increasing expression under nitrogen starvation
while their expression remained constant during the whole-time course in the control plants. We
refer to this group as **(I) early acting upregulated genes**. GO, KEGG, and manual annotation
indicated that this group was strongly enriched for genes involved in transmembrane transport
both at the plasma membrane and that of the vacuole including amino-acid transporters and
various ion transporters (including ammonium, nitrate/nitrite, potassium, sulfate, iron
transporters). We further found that the group of **early acting upregulated genes** was enriched
for lipid synthesis related genes. Besides that, we observed the upregulation of genes related to
plant hormonal activity. In particular, we found that genes involved in the synthesis of gibberellic
acid precursors (*KO, KAO, CPS/KS*), and strigolactones (*CCD7, CCD8*), modifying the concentration
and effect of cytokinin (cytokinin receptor), signaling ABA (*NCED, abi5, bZIP*) and jasmonate
action (*JAZ/Tify*) or involved with auxin concentration change were upregulated (*AUX1/LAX*). In
parallel with this we also observed that genes involved in protecting the plant from oxidative
damage were highly upregulated (flavonoid synthesis, phenylpropanoid pathway, polyphenol
oxidases, alcohol dehydrogenases, glutathione). Nitrogen starvation also led to strong changes
in carbohydrate metabolism including the upregulation of genes synthesizing trehalose and
catalyzing some reactions of glycolysis and gluconeogenesis. Starvation further led to the
induction of enzymes catalyzing the release of nitrogen from amino acids and enabling the
assimilation of ammonium (glutamine synthase). The upregulation of various Ca-dependent
kinases suggested the involvement of calcium signaling in the response to nitrogen starvation. In
addition to that, we observed the expression of enzymes involved in cell wall remodeling. Finally,
the early reacting gene set also contained several transcription factor (TF) genes that are likely
orchestrating the molecular response (*GAPR_G2*-like [*NIN*-like], *MYB*-related, *bHLH*, *HD-III-ZIP*,
WRKY, RAM1 and other *GRAS, NAC, PHD, C2H2, bZIP, ARID, SET, RWP-RK, OFP, JAZ/TIFY, C2C2-*
Dof).
463
A second set of genes (325 genes, cluster 4) experienced downregulation immediately after
starvation, while their expression values were only slightly downregulated in the control samples.
We refer to these genes as **(II) early acting downregulated genes**. These sets of genes are
primarily enriched for pathways related to ribosome assembly/genesis, mRNA and rRNA
processes plus amino acid synthesis. That is, genes responsible for the assembly of ribosomes
and protein synthesis are immediately blocked by nitrogen starvation. We also found some TF
genes being downregulated (*AP2/EREBP, GNAT, AS2/LOB, C2H2, HD-ZIP*).
471
Genes of cluster 3 were also downregulated but their expression decrease started only after the
$3^{rd} - 4^{th}$ day of starvation. We refer to these genes as **(III) later acting downregulated genes**.
This gene set was primarily enriched for photosynthesis-related genes, such as photosynthetic
complexes (PSI, PSII) and genes responsible for chlorophyll synthesis suggesting that blocking of

photosynthetic processes occurs somewhat later during nitrogen starvation. Some genes related
to hormonal response and/or transcription factors were also members of this group like cytokinin
oxidases (*CKX*), a *YUCCA* gene, ABA and cytokinin synthesis genes (*ABA4, IPT*), *PLATZ, C3H*.
479
480   Finally, genes of cluster 1 showed relatively rapid but weak downregulation during the first three
481   days of starvation and their expression only slightly recovered after seven to ten days.
482   Functionally, this cluster showed similarity to the **later acting downregulated genes** (cluster 3)
483   because it contained genes involved with ribosome biogenesis, but also in protein synthesis,
484   protein translation, proteasome building, and protein folding.

486   A: Enriched GO terms of upregulated genes (cluster2)

GO Molecular Functions

B: Enriched GO terms of downregulated genes (cluster 1, 3 and 4)

Figure 5. GO enrichment analysis (Molecular function ontology, P value < 0.05, Fisher's exact test)
of genes up/downregulated under nitrogen starvation in *A. agrestis*. A: enriched GO terms of
upregulated genes from cluster 2. B: enriched GO terms of downregulated genes from cluster 1,
3 and 4.

Blasia pusilla

The temporal dynamic of *B. pusilla* genes reacting to nitrogen starvation was similar to those of
A. agrestis ([Fig. 6A a](#)nd B, and see Table S14, GO enrichment results in Table S16 and Fig. S7-S10).
Nevertheless, genes of cluster 2 showed a more gradual expression increase and reached their
maximum expression value after 10 days of starvation, while genes of cluster 4 peaked already
by the third day. We further refer to these two clusters as **gradually** and **rapidly reacting**
upregulated genes. The **gradually reacting** group contained genes increasing nitrogen and
sulfate uptake from the environment and their assimilation within the cell (glutamine synthase,
ammonium, nitrate/nitrite and oligopeptide transporters). We also detected that this group
contained several hormonal response genes related to stress response such as auxin (*SAUR*), *ABA*
(*ABF*), gibberellin precursors (*KAO*) and cytokinin (*CYP735A*). Several genes related to cell
signaling and protection against oxidative damage were upregulated such as phosphatidyl
inositol and flavonoid synthesis genes, polyphenol oxidases and peroxidases. There was an
indication of intensified signaling, such as increased expression of various kinases including Ca-
dependent kinases suggesting that starvation reaction may be linked to Ca signaling. The
increased activity of histone acetylase genes suggests that epigenetic reprogramming is taking

place and may contribute to transcriptional reorganization. Finally, we also found that multiple
TF genes were also part of this group (*AS2/LOB, bHLH, AP2/EREBP*).

The group of **rapidly reacting upregulated** genes contained genes with similar functions. We
observed multiple genes involved with nitrogen uptake, such as ammonium and nitrite/nitrate
transporters. Besides the previously mentioned transporters, we also detected the immediate
upregulation of sulfate transporters. An exclusive set of genes not detected in the gradually
upregulated group were those involved with cell wall modifications/remodeling. Similarly,
multiple mRNA, protein and amino acid synthesis-related genes were present in the rapidly
reacting but not in the gradually upregulated group of genes. Finally, we also found a handful of
TF genes that reacted rapidly to nitrogen starvation (*AP2/EREBP, Trihelix, bHLH, TRAF, NAC, MYB-*
related, ZF-HD, bZIP).

Downregulated genes were also distributed in two clusters. Genes of cluster 1 were rapidly
downregulated after nitrogen starvation, reaching their minimum expression level already on
day 3. These genes are referred to as **early acting downregulated genes**. We found that multiple
photosystem I/II and chlorophyll synthesis-related genes occurred in this group. Similarly, genes
involved with ribosome biogenesis and translation were present as well as genes involved with
signaling, auxin and gibberellin response, DNA replication and cell wall remodeling. In contrast to
A. agrestis, we found that multiple TF genes were also part of this group (*GRAS, NAC, bHLH, GATA,*
AP2/EREBP, C2H2, SET, bZIP, WRKY, HD-ZIP). Therefore, besides some TF genes, this cluster of
early downregulated genes was functionally like cluster 2 of *A. agrestis*. While photosynthesis
related genes reacted somewhat later in *A. agrestis* they did so earlier in *B. pusilla*.

Finally, genes of cluster 3 were slightly downregulated and showed relatively low expression in
the nitrogen starved samples while regained expression in the control samples. Genes of this
cluster are functionally overlapping with those found in the early downregulated group. For
instance, we found genes involved in nitrogen preservation such as amino acid hydrolysis and
amino acid synthesis. Similarly, some photosynthesis related genes were also present.
Furthermore, auxin efflux carriers, signaling protein kinases and some cell wall synthesis genes
were also present. Finally, we also found some TF genes to be downregulated (*NAC, C2H2, bHLH,*
AS2/LOB).

545   A: Enriched GO terms of upregulated genes (cluster 2 and 4)

548   B: Enriched GO terms of downregulated genes (cluster 1 and 3)

549   **Figure 6.** GO enrichment analysis (Molecular function ontology, P value < 0.05, Fisher's exact

550   test) of genes upregulated (genes of cluster 2 and 4) during nitrogen starvation in *B. pusilla*. A:

551   enriched GO terms of upregulated genes from cluster 2 and 4. B: enriched GO terms of

552   downregulated genes from cluster 1 and 3.

Evolutionary and functional conservation of the nitrogen starvation response between A.
agrestis and B. pusilla

We hypothesized that part of the starvation response is evolutionary conserved between the liverwort and hornwort models used and could have been present in their most recent common ancestor (the common ancestor of bryophytes). We found 53 orthogroups showing shared upregulation under nitrogen starvation both in *B. pusilla* and *A. agrestis* (see Table S17, S18). GO, KEGG enrichment analysis and manual annotation of genes implies that this set of genes is enriched for plasma membrane transport processes aiding the uptake of various ions, nitrogen fixation and organic material transport. Another set of orthogroups implied that hormonal responses are also shared. We found that bZIP transcription factors (TFs) (*ABF*), cytokinin-modifying enzymes, gibberellin precursor synthesis genes, and auxin action related genes were upregulated in both species. A third set of orthogroups suggested that response of the major signal transduction and signaling cascades to nitrogen starvation is also conserved between the two species. For instance, orthogroups containing leucine-rich repeat protein kinases, protein kinases (nodule lectin) and calcium-dependent kinases show shared upregulation in both species during nitrogen starvation. We also found orthogroups containing the enzyme involved in trehalose synthesis (trehalose-6-phosphate synthase) upregulated in both species producing the essential and potent signaling molecule as well as important osmotic protectant within plant cells, trehalose. A fourth set of orthogroups suggests that nitrogen recycling from nucleotides or from chlorophyll is also evolutionary conserved between the two species. We found that orthogroups containing genes with function in obtaining nitrogen from purine nucleotides (amidase (allantoate deiminase (*allC*))) or orthogroups containing homolog of the stay-green gene that degrade chlorophyll are upregulated in both species. Finally, several shared upregulated orthogroups contained genes with functions either related to pathogen defense, or to protection from oxidative damage. This involved enzymes of the phenylpropanoid pathway, various peroxidases, and subtilisin-like proteases.

The 26 orthogroups exhibiting shared downregulation showed similar properties to what we have found in the separate analysis of the two species (see Table S17, S19). Downregulated genes were involved in transcription and translation processes which appeared to be blocked during nitrogen starvation. Similarly, chlorophyll synthesis and cell-wall expansion related genes and various metabolic processes were downregulated.

While the systemic response to nitrogen starvation was largely conserved between the model hornwort and the model liverwort, we also observed some notable differences. Our comprehensive TF annotation revealed that upregulation of various TF families was either specific to the hornwort, or to the liverwort. We found that some orthogroups containing *HD-*

592   *ZIP*, *WRKY*, and *TIFY* TFs were only upregulated in the hornwort, while *AS2/LOB, AP2/EREBP,*
593   *Trihelix* and *TRAF* were only in the liverwort (Fig. 7). Similarly, orthogroups containing strigolacton
594   precursor enzymes and some lipid biosynthesis pathway genes were only upregulated in the
595   hornwort model.
596
597

598
599
600   **Figure 7.** Venn diagram illustrating upregulated transcription factor families (TFs) in the liverwort
601   (*B. pusilla)* and the hornwort (*A. agrestis)*. The intersection refers to shared upregulation.
602
603
604   ***Conservation of the response to nitrogen starvation among bryophytes and across land plants***
605
We hypothesized that all land plants deeply share a starvation response which was further
extended with symbiont and lineage-specific modules to regulate symbiotic interactions
between plants and their microbes. Both the hornwort *A. agrestis* and the liverwort *B. pusilla* can
establish intimate symbiotic interaction with cyanobacteria while such interactions are not
observed in other liverworts such as *M. polymorpha* and *M. paleacea*. Therefore, we reasoned
that all four liverworts are expected to share a common response to nutrient starvation which

has been extended with further modules in the hornwort (*A. agrestis*) and liverwort (*B. pusilla*) to enable a more intimate interaction with cyanobacteria. We identified 21 orthogroups showing shared upregulation in *A. agrestis* and *B. pusilla* during nutrient starvation but no response in *M. paleacea* and *M. polymorpha* (See Table S20-S23). These 21 orthogroups contained genes with putative functions including beta-glucanases, molybdate transporters, aldehyde dehydrogenases, trehalose synthetases, and urea transporters. We also found that several orthogroups containing TFs showed shared upregulation among others, bZIPs (*ABF*).

We further reasoned that if upregulation of these orthogroups is required to specifically attract cyanobacteria, they should not be upregulated in vascular plants establishing symbiotic interactions with either AMF or nodule-forming bacteria but not with cyanobacteria. Nevertheless, we found that most of the 21 orthogroups were also upregulated in one or more vascular plants not able to establish intimate interaction with cyanobacteria ([Fig. 8](#)).

Figure 8. Selected orthogroups upregulated under nutrient starvation (either nitrogen or phosphate starvation) a wide range of symbiotic and non-symbiotic vascular and non-vascular plant species (cyanobacteria-symbiosis, AM fungi symbiosis, nodule-forming bacteria symbiosis, and non-symbiotic species). Black: no gene is assigned to this orthogroup, green: orthogroup contains genes but none of them are upregulated, yellow: at least 1 gene assigned to the orthogroup is upregulated in response to starvation.

636
637
Discussion
639

It has been proposed that the regulatory network responsible for initiating plant-microbe
symbiotic interactions has evolved from a conserved, ancient nutrient starvation response that
was already present in the common ancestor of all land plants (Isidra-Arellano et al., 2021; Paries
and Gutjahr, 2023). Later this network has further recruited and extended to genes specifically
regulating the intimate interaction between plants and AM fungi, nodule-forming bacteria and
potentially cyanobacteria (Ma and Chen, 2021; Delaux and Schornack, 2021; Puginier et al., 2022;
Shi et al., 2022). Here we investigated this hypothesis by exploring gene expression changes
during nitrogen starvation in the two major lineages of bryophytes, hornworts and liverworts,
able to establish intimate symbiotic interaction with cyanobacteria (Meeks, 1998; Álvarez et al.,
2023a). By conducting comparative analyses with symbiotic and non-symbiotic vascular and non-
vascular plants, we found that the systemic gene expression response to nitrogen/phosphate
starvation is similar in bryophytes and vascular plants. Additionally, examination of the dataset
allowed us to identify several groups of compounds that may serve as hormogonia-inducing
factors (HIF), while providing little evidence for an extensive set of genes involved in priming the
host plant for the cyanobiont. These findings allow us to propose a hypothetical description of
the starvation response and its convergence and divergence in the hornwort and liverwort
models, which is detailed below (Fig. 9).

657
658

Systemic response to nitrogen starvation appears to be conserved across bryophytes and
vascular plants

Response to nitrogen starvation in vascular plants involves sensing nitrogen levels, transduction
of the signal into the cell, and adjusting gene expression and metabolism to preserve metabolic
balance (Varala et al., 2018; Gaudinier et al., 2018; Wang et al., 2019; Ueda and Yanagisawa,
2019; Ueda et al., 2020). We observed conservation in the response of *A. agrestis* and *B. pusilla*
as well as with vascular plants in most of these steps.

Sensing and signaling of low nitrogen environment
Low nitrogen levels are sensed by *NRT* genes (nitrate sensor) which work together with calcium
channels and activate calcium signaling pathways inducing transcription factors as well as further
genes involved in nitrogen uptake, nitrogen metabolism and auxin regulation (Mascia et al., 2019;
Dong et al., 2022; Fang et al., 2022; Adavi and Sathee, 2023). We observed that major
components of the calcium signaling pathway (calmodulins, calcium-dependent protein kinases,

serine-threonine protein kinases) were induced both in *A. agrestis* and *B. pusilla* during starvation. This suggests that calcium signaling is a common theme between nitrogen starvation in bryophytes and vascular plants. Calcium signaling is also central to transmit systematic response to various stresses (Dong et al., 2022). Whether calcium signaling involves calcium spiking or other mechanisms is unknown.

Nitrogen Uptake, Transport, assimilation, and Related Modulators

In vascular plants nitrogen limitation rapidly activates nitrogen uptake systems enabling its transport from the environment to the cell (Fan et al., 2017; Aluko et al., 2023). Nitrate (*NRT*), ammonium (*AMT*) transporters are upregulated to increase nitrogen uptake and assimilation (Rui et al., 2022; Aluko et al., 2023). In vascular plants, activation of nitrogen uptake and assimilation mainly occurs in the root. While roots are not present in non-vascular plants, rhizoid plays an analogous role (Jones and Dolan, 2012; Szövényi et al., 2019). We found that nitrate and ammonium transporter homologs were upregulated in the *B. pusilla* and *A. agretis* systems like in vascular plants. Specifically, an ammonium transporter *AMT2* and a nitrate transporter *NRT2.2* homolog were upregulated both in *A. agrestis* and *B. pusilla*. Amino acid and peptide transporters were also upregulated that likely help the uptake of amino acids/peptides from the environment. In vascular plants nitrogen is mainly available and taken up as NO_3^-, and NH_4^+ (Masclaux Daubresse et al., 2010). NO_3^- is then reduced to NO_2^- by the nitrate reductase (*NR*) and then transported to the chloroplast together with NH_4^+. NO_2^- is reduced to NH_4^+ by the nitrite reductase and together with the transported NH_4^+ fed into the GS/GOGAT cycle to be assimilated into glutamate which serves as a precursor of multiple metabolites (Dubey et al., 2014). When vascular plants are starved of nitrogen, protein reserves of the chloroplast are proteolyzed and NH_4^+ and amino acids are recycled into glutamine by GS. We found that nitrogen starvation led to the upregulation of the *GS* and not the *GOGAT* enzyme in both species *B. pusilla* and *A. agrestis* suggesting recycling of protein resources to maintain nitrogen balance of the cell. This implies the presence of a physiological response conserved between vascular plants and bryophytes in preserving nitrogen balance.

Besides nitrogen transporters, our transcriptome analyses also suggested that ABC transporters are induced under nitrogen starvation (both *ABCG* and *ABCB* transporters). *ABCG* transporters transport lipids and or (iso)flavonoids as well as phytohormones. Even though it has been shown that cyanobacteria receive carbohydrates (at least some of it in the form of sucrose) from its plant partner (Kaplan and Peters,1998), the necessary transporters of either symbiont have not been identified. We found that *SWEET13* homologs were upregulated in *A. agrestis* under nitrogen starvation conditions likely transporting sugars towards the cyanobiont.

Hormonal response

In flowering plants, nitrogen and phosphate starvation cause hormone synthesis
activation/repression. Hormonal changes lead to both physiological and morphological
adaptations increasing nutrient usage. Auxin synthesis and transport are well known to be
activated under low nitrogen conditions to change root shoot allocation (Sultana et al., 2020).
The most obvious effect of auxin is to trigger root hair growth under nitrogen and phosphate
limitation (López-Bucio et al., 2002). Auxin response is apparent in both species data sets (*GH3,*
AUX1) and likely trigger rhizoid development as was shown in the liverwort *M. polymorpha* (Rico-
Reséndiz et al., 2020; Rico-Resendiz et al., 2021; Suzuki et al., 2021). Because auxin is necessary
for normal organogenesis in *M. polymorpha*, it is possible that it also plays a role in the
developmental process of the cavities containing the cyanobacterial cells in both *B. pusilla* and
A. agrestis (Suzuki et al., 2023).

Similarly, a low nitrogen environment is known to promote ABA accumulation which will affect
lateral root formation, nitrate sensing and transport in vascular plants (Ondzighi-Assoume et al.,
2016; Wang et al., 2018; Lee et al., 2020). Overall, ABA helps to increase tolerance to low nitrogen
conditions in general. We assume that ABA has a similar stress alleviating effect in the hornwort
and the liverwort, however its effect awaits further scrutiny.

The role of cytokinin in nitrogen starvation of vascular plants is poorly explored but phosphate
starvation reduces cytokinin levels in *A. thaliana* thereby affecting the shoot to root ratio and
enabling root hair growth (Kiba et al., 2011; Lv et al., 2021). Similarly, it was found that cytokinin
oxidase genes, decreasing the amount of active cytokinin, are upregulated during phosphate
starvation in *M. polymorpha* (Rico-Resendiz et al., 2020). Furthermore, cytokinin is known to
promote rhizoid development in *M. polymorpha* like its effect on root hair growth in flowering
plants and may have a similar function in *A. agrestis* and *B. pusilla* (Rico-Reséndiz et al., 2020;
Rico-Resendiz et al., 2021).

GA levels increase in vascular plants under nitrogen starvation, and it likely triggers stress
tolerance against low nitrogen conditions. GA receptors and canonical gibberellins are not
present in bryophytes (Guillory and Bonhomme, 2021), but hormonal diterpenoids regulate
various aspects of development in mosses as well as in liverworts. For instance, *DELLA* proteins
promote protection against oxidative stress response and coordinate growth across various
environmental inputs in the liverwort *M. polymorpha* (Hernández-García et al., 2021). Our results
suggest that this response is also likely conserved between liverworts, hornworts, mosses and
vascular plants.

Jasmonates are also upregulated under defense in flowering plants and are likely to regulate
sugar, amino acid synthesis and NH_4^+ uptake (Sun et al., 2020; Lv et al., 2021). By contrast, in the
hornwort model some *JAZ* homologs were upregulated which are known to repress jasmonate
synthesis in liverworts and promote cell growth and reproductive fitness (Monte et al., 2019).
Furthermore, reduced jasmonate synthesis leads to repressed defense responses. While the role
of *JAZ* upregulation during nitrogen starvation in the hornwort is currently unclear, it is possible
that it decreases plant defenses against pathogens and this way triggers the priming of the host
plant to the cyanobiont (Monte et al., 2019).
755
Together with jasmonates, salicylic acid (SA) also plays an important role in the defense against
pathogens in vascular plants (Ding and Ding, 2020). SA has been proposed to be involved in the
signaling between host plant and the cyanobiont in the *Azolla-Nostoc* symbiosis (de Vries et al.,
2018, 2023). While most biosynthetic enzymes of SA are present in the hornwort genome, we
found none of them upregulated during nitrogen starvation. Therefore, our data do not support
the findings observed in the *Azolla-Nostoc* symbiosis.
762
763 **A.**   **B.**
764

765
766
Figure 9. Systemic response to nitrogen starvation and potential priming of the host to the
cyanobiont. A.) Up- and downregulated biological processes under nitrogen starvation shared by
B. pusilla and *A. agrestis*. B.) Biological processes/genes potentially contribute to priming the
plant host to the cyanobiont. Transporters are shown on light gray background, transcription
factors on dark gray background in white letters.
Transcription factors
The nitrogen starvation induced calcium signaling pathway converges on various transcription
factor genes that act as master regulators of the nitrogen starvation response. In vascular plants
bZIP transcription factors are usually upregulated and transmit ABA response, increase nitrogen

uptake and regulate root branching (Zhang et al., 2015; Yang et al., 2019; Ma et al., 2020). *WRKY*
transcription factors are also induced and increase nitrogen stress tolerance (Curci et al., 2017;
Wang et al., 2019). Upregulation of *MYB* and MYB-related transcription factors increase nitrogen
influx, development, cell cycle and morphology (*NIGT*) (Curci et al., 2017; Wang et al., 2019),
while CCAAT-binding TFs (such as *TaNFYA)* are important regulators of nitrate transporters (Qu
et al., 2015; He et al., 2015). *C2H2* TFs are known to trigger low-nitrogen adaptation and *Dof* TFs
can induce ammonium transporters as well as the synthesis of carbonic scaffolds to be used for
nitrogen assimilation (Yanagisawa et al., 2004; Noguero et al., 2013; Chen et al., 2017).
784
We found that multiple members of these TF families were also upregulated in either one or both
species investigated. For instance, *bZIP* TFs of the ABA responsive element binding factor (*ABF*),
MYB-related, *bHLH* and *NAC* TFs were upregulated during nitrogen starvation in both species.
Homologs of these TFs were also upregulated under nitrogen starvation in *M. polymorpha* and
phosphate starvation in *M. paleacea* (Radhakrishnan et al., 2020; Rico-Reséndiz et al., 2020).
Therefore, these TFs appear to represent a conserved subset of master regulators that could have
been already present in the common ancestor of most bryophytes and some of them may have
master regulatory functions even shared with vascular plants.
793
NAC TFs are also often upregulated under nitrogen starvation and affect root growth, branching
and nitrate uptake (Vidal *et al.*, 2013, 2014; He *et al.*, 2015). Stress induced *NAC* TFs have recently
also been found to be induced under nitrogen starvation in rice (Qi et al., 2023). Furthermore, it
was also shown that the *OsNAC1* gene directly induced the expression of multiple nitrate
transporters (*NRT* genes) and led to higher nitrogen use (NUE) efficiency, biomass and grain yield
(Tang et al., 2019; Li et al., 2020a; Qi et al., 2023). Therefore, we hypothesize that the connection
between nitrogen starvation induced *NAC* TFs and *NRT* genes is likely to be conserved among
liverworts, hornworts and vascular plants. Future work should reveal to what extent this
conserved systemic regulatory module is linked to further modules potentially enabling
colonization of the host plant by the cyanobiont.
804
MYB-related TFs, especially phosphate response 1 (*PHR1*), is a master regulator of the phosphate
starvation response in flowering plants as well as in *M. polymorpha* (Rouached et al., 2010; Rico-
Reséndiz et al., 2020; Isidra-Arellano et al., 2021; Tan et al., 2023; Paries and Gutjahr, 2023).
Furthermore, *PHR1* is essential in priming the host plant for plant AMF and rhizobia symbiosis by
regulating plant immunity and a diverse set of genes necessary for the initiation and
establishment of the symbiotic interaction (Isidra-Arellano et al., 2021; Paries and Gutjahr, 2023).
Importantly, our analysis indicates that the upregulated Myb-related transcription factors are
homologous to the *PHR1* gene of *A. thaliana* and *M. polymorpha*. Therefore, it is likely that the
upregulation of *PHR1* homologs and their master regulatory function in nitrogen/phosphate

starvation are shared between the hornwort, the liverwort models (and other liverworts) and
vascular plants. How much the regulated gene set differs between flowering plants and our two
model species needs further investigation.
817
Multiple *bHLH* TFs are known to be upregulated under various stresses in flowering plants
including phosphate and nitrogen-starvation (Yang et al., 2016; Subudhi et al., 2020). Some of
these genes regulate root growth while some others influence phosphate/nitrogen starvation
response genes (Crombez et al., 2019; He et al., 2021). Phylogenetic analysis shows that the *bHLH*
gene identified in the hornwort and liverwort models forms a monophyletic group with *Azolla*
filiculoides genes. Furthermore, it is embedded in a large clade containing only bryophyte *bHLH*s.
Therefore, this gene could potentially be important for the cyano-hornwort/liverwort symbiotic
interaction.
826
While some TFs reacted similarly in the hornwort and liverwort species (*A. agrestis* and *B. pusilla*)
investigated, many TF family genes showed divergent expression patterns whose potential
function remains to be investigated. For instance, *AS2/LOB* TFs are known to repress nitrate
starvation response genes in *A. thaliana*, while they were upregulated only in our model liverwort
B. pusilla (Rubin et al., 2009; Dash et al., 2015). Similarly, *AP2/EREPB* TFs are known to be induced
in nitrogen starved flowering plants (Yang et al., 2015), but they were only upregulated in the
liverwort model studied. Finally, the role of *Trihelix* and *TRAF* TFs in nitrogen starvation response
of *B. pusilla* is unclear.
835
Intriguingly, we found many more TF genes to be upregulated under nitrogen starvation in the
hornwort than in the liverwort. Some of them are well-known to show similar responses in
flowering plants including, *HD-ZIP III* (Lin et al., 2022), *WRKY* (Wang et al., 2019), *MYB* (Zhang et
al., 2012; Wang et al., 2019, 2020b), *C2H2* (SUN et al., 2012; Chen et al., 2017), *C2C2-Dof*
(Yanagisawa et al., 2004; Curci et al., 2017), *GARP* (Medici et al., 2015) and *RWP_RK* (Chardin et
al., 2014; Wu et al., 2020) TFs. It is unclear why these TFs are not responsive in the liverwort *B.*
pusilla although these genes are mostly present in the genome of liverwort *B. pusilla.* We
speculate that the difference between the hornwort and the liverwort model may be partly due
to their divergent habit/growth form and thallus organization as well as their ability to engage in
symbiotic interaction with AMF (see below).
846
847
848   ***Nitrogen starvation and symbiotic interaction***
849
It has been known for a long time that nutrient starvation (Pi/N) is a prerequisite for the
establishment of plant-AMF and plant-rhizobia symbioses (Ma and Chen, 2021; Das et al., 2022).

In these interactions master regulators of the phosphate (*PHR*) or nitrogen starvation syndrome
(*NIN*) directly regulate genes involved in pre-colonization signaling of the symbiotic interaction
with AMF or rhizobia, respectively (Diédhiou and Diouf, 2018; Luo et al., 2023). As the very first
step of symbiosis initiation, these master regulators activate the synthesis and secretion of
special chemical signals, strigolactones in AMF and flavonoids in nodule-forming symbiosis, which
trigger the growth and colonization potential of the microbes. As a response to the chemical
signals, the microbes release signaling molecules perceived by host plant receptors and activate
precolonization signaling (Oldroyd, 2013; Zipfel and Oldroyd, 2017; Oldroyd and Leyser, 2020;
Jhu and Oldroyd, 2023).

Similarly, plant-cyanobacteria symbiotic interactions are also suppressed in the presence of
sufficient amounts of combined nitrogen (Meeks, 1998; Adams and Duggan, 2008; Álvarez et al.,
2023b). Starved host plants release the so-called hormogonia inducing factor (HIF) triggering the
conversion of vegetative cyanobacterial filaments to motile hormogonia that can colonize the
host plant (Campbell and Meeks, 1989; Bergman et al., 1996; Meeks, 2003; Adams, 2005).
Nevertheless, the exact chemical composition of the HIF is poorly understood and appears to
vary across species. Previous studies found that monosaccharides (arabinose, glucose),
diacylglycerols, or even various pigments likely function as HIFs (Rasmussen, 1994; Nilsson et al.,
2006; Adams and Duggan, 2012; Liaimer et al., 2015; Hashidoko et al., 2019; Alvarenga et al.,
2022; Carrell et al., 2022). Intriguingly, our transcriptomic analysis suggests that genes
responsible for the synthesis of some of these compounds are upregulated under nitrogen
starvation both in the hornwort and in the liverwort system investigated. Furthermore, they are
rarely upregulated in species forming no symbiotic interaction with cyanobacteria.

For instance, we found that genes involved in flavonoid synthesis as well as *ABCG* transporter
homologs known to transport isoflavonoids are upregulated in both the liverwort and the
hornwort systems. This suggests that flavonoid type molecules can potentially function as HIF
components. This is in line with previous findings that flavonoids can affect the activity of a gene
cluster involved in hormogonia formation of the cyanobiont (Cohen and Yamasaki, 2000).
Hormogonia are the motile forms of cyanobacteria necessary for the colonization of the host
plant. Similarly, we observed the upregulation of trehalose-synthetase genes. Trehalose was
recently proposed to potentially be transported from the host plant to the cyanobiont and may
have HIF-like activity in peat mosses (Carrell et al., 2022). Therefore, both compounds should be
investigated for their HIF activity in the future.

In the plant-AMF and nodule-forming bacteria symbioses starvation is not only necessary to
attract and mobilize the microbial partner but it is also essential to prime the host for the
interaction by adjusting its physiology as well as creating specialized cellular structures guiding

microbial colonization (Isidra-Arellano et al., 2021; Shi et al., 2021; Das et al., 2022). While
specialized cellular structures (cavities) hosting the microbes are performed and do not need to
be induced in the hornwort- and liverwort-cyanobacteria symbioses, it is unclear how much
further priming is necessary to accept the cyanobiont. Collectively, our data suggests that some
genes induced by nitrogen starvation may be involved in priming, but an extended network of
genes is absent. Therefore, it is possible that no extensive priming is necessary besides
chemotactic attraction and HIF-mediated mobilization of the cyanobiont. Alternatively, the
absence of an extensive shared priming network may be a consequence of the independent
evolution of symbiotic interaction regulation in hornworts and liverworts.

While the gene network involved in priming is unlikely to be extensive, our data provides some
candidate genes whose expression is triggered by nitrogen starvation and could potentially be
important in priming the plant host for symbiotic interaction. This is in line with the observations
that starvation is not only necessary to attract and mobilize the cyanobiont but it is also essential
to maintain symbiotic interaction (CAMPBELL and MEEKS, 1992). For instance, it is known that
blocking the upregulation of nitrate transporter genes in the plant-nodule-forming bacteria
interaction will inhibit the establishment of the symbiotic interaction (Wang et al., 2020a).
Therefore, it is possible that the upregulation and proper localization of various transporters in
the hornwort and liverwort during nitrogen starvation helps priming the host plant for the
symbiotic interaction. In the hornwort and the liverwort, we also found sugar transporters
(including *SWEET*) to be upregulated. This is in line with the observation that photosynthates are
provided by the host plant to the cyanobiont (de Vries and de Vries, 2022). *Nostoc* with
dysfunctional sugar transporters will not establish symbiotic interaction (Ekman et al., 2013), and
that host plant cavities containing the cyanobiont are filled up with polysaccharide containing
mucilage. Therefore, it is possible that upregulation of sugar transporters is necessary for the
initiation and establishment of the symbiotic interaction by attracting, feeding and retaining the
cyanobiont. Finally, we also detected alcohol dehydrogenases to be mainly upregulated in the
hornwort and liverwort models during nitrogen starvation but not in species without intimate
interaction with cyanobacteria. Alcohol dehydrogenase overexpression is often observed in
pathogen colonization and was shown to support biotrophy (Pathuri et al., 2011). Therefore,
overexpression of such enzymes may be important in priming the host plant for symbiotic
interaction.

Several nitrogen-starvation induced genes in A. agrestis are likely involved in priming the host
plant for AMF symbiosis

While the above-mentioned transcriptional responses were mainly shared between the
hornwort and the liverwort, a large number were specific to only one of the model species
investigated. Intriguingly, more genes and TFs were upregulated in the hornwort model and
many of these were shared with species exhibiting symbiotic interaction with AMF. We speculate
that the hornwort-specific upregulation of these genes reflects the ability of *A. agrestis* to form
symbiotic interactions with AMF.

NIN-like TFs (*NLP* genes, *RPW-RK* transcription factors) are especially interesting in this respect
because they are master regulators of the nitrogen-starvation response regulating many other
genes (Chardin et al., 2014). We found that multiple *RPW-RK* transcription factors homologous
to the NIN-like proteins of *M. polymorpha* (Mu and Luo, 2019; Hsin et al., 2021) were significantly
upregulated but only in *A. agrestis*. Therefore, our data suggests that the sensing and signal
transduction of low nitrogen environments in *A. agrestis* likely involve Ca^{2+} signaling and NIN-like
proteins in a similar way to what is described for vascular plants (Mu and Luo, 2019).

The upregulation of a *RAM1* homolog in the hornwort model during nitrogen starvation is also
intriguing. Expression of *RAM1* together with other *GRAS* TFs is known to be induced by
phosphate starvation in vascular plants establishing AMF symbiosis and is essential in inducing
various genes required to form the infection thread and the interface between the host plant
and the fungal partner (Gobbato et al., 2012; Xue et al., 2015). *RAM1* is also essential for lipid
provisioning (Luginbuehl et al., 2017). *RAM1* is a member of the common symbiosis signaling
pathway (CSP) genes with a complete set of homologs present in the hornwort genome (Li et al.,
2020b). By contrast, most homologs of the CSP genes are absent from the *B. pusilla* genome.
Furthermore, we found that no *GRAS* TFs were upregulated under nitrogen starvation in *B.*
pusilla. This suggests that *GRAS* TFs may not be necessary to prime the host plant for the
cyanobiont in the hornwort and liverwort model systems but could be important in priming the
hornwort for AMF colonization. This is further supported by the upregulation of lipid synthesis
genes in the hornwort *A. agrestis* but not in the liverwort *B. pusilla* (Rich et al., 2021).

Another prominent example is the upregulation of genes involved in the biosynthesis of
strigolactone precursors during nitrogen starvation which was also recently observed by other
studies (Radhakrishnan et al., 2020; Rich et al., 2021). We found that nitrogen starvation
positively affects the expression of *D27* and the gene encoding the enzyme catalyzing subsequent
steps of strigolactone biosynthesis – *CCD7.* A new form of strigolactone has recently been shown
to affect AMF colonization in the liverwort *M. paleacea* (Kodama et al., 2022). It was also
observed that some of the key strigolactone synthesis genes are missing from the genome of *M.*
polymorpha, a sister species of *M. paleacea*, which is not able to establish symbiotic interaction
with AMF. Therefore, the secretion of strigolactones into the environment appears to be

necessary for AMF interaction. When comparing the hornwort and the liverwort proteomes, we
made similar observations and found that genes crucial for the biosynthesis of strigolactones
were missing from the *B. pusilla* but were present in the *A. agrestis* genome. Therefore, we
hypothesize that strigolactones are unlikely to be necessary to establish the plant-cyanobiont
interaction. While our data and previous observations support our hypothesis, the potential role
of strigolactones and *GRAS* TFs in the hornwort-cyanobacteria interaction cannot be fully ruled
out and warrants further scrutiny.
972
973
974 ***Overall conclusions***
975
Our study also has some important applied implications for engineering plant-cyanobacteria
symbiotic interactions. The observation that almost no orthogroups were specifically
upregulated in the cyanobacteria host plants suggests that little host plant priming is needed to
accept the cyanobiont. Therefore, the attraction and mobilization of the cyanobiont by the HIF
may be the most critical step in establishing symbiotic interaction. Our data suggest that beyond
that, very little if any priming is necessary in the host plant to accept the cyanobiont. Therefore,
identifying compounds with HIF activity and applying them under natural conditions may be key
to attracting cyanobacteria by non-host plants.
984
985
986 **Supplementary data** (available at Figshare https://figshare.com/s/0acebe58f3dee5065469)
987
Table S1. Design of the starvation experiment (*A. agrestis* and *B. pusilla*) and the number of
replicates used for RNA sequencing.
Table S2. Proteomes used in the orthofinder analysis.
Table S3. RNA-seq data used for comparative transcriptomic analyses.
Table S4. Genes showing significant (padj<0.05) response (up and down-regulation) over the
course of the nitrogen starvation experiment for *M. polymorpha*.
Table S5. GO annotation of *A. agrestis* genes including the molecular function, biological
process and cellular localization ontologies.
Table S6. Kegg annotation of *A. agrestis* genes.
Table S7. GO annotation of *B. pusilla* genes.
Table S8. KEGG annotation of *B. pusilla* genes.
Table S9. Predicted transcription factors and genes involved in hormone synthesis and signaling
(*B. pusilla* and *A. agrestis*).
Table S10. Blobtools contamination filtering of the *B. pusilla* draft genome assembly.

Table S11 Normalized expression values for *A. agrestis* over the course of the time-series
experiment.
Table S12. Results of the maSigpro analysis for *A. agrestis*.
Table S13. Normalized expression values for *B. pusilla* over the course of the time-series
experiment.
Table S14. Results of the maSigpro analysis for *B.pusilla*.
Table S15. GO-enrichement analyses results for *A. agrestis* (all four clusters).
Table S16. GO enrichment analysis results for *B. pusilla* (in all four clusters).
Table S17. Up- and down-regulated genes and their orthogroup assignment in *A. agrestis* and *B.*
pusilla.
Table S18. Upregulated genes in the *B. pusilla* and *A. agrestis*.
Table S19. Genes downregulaten in the *B. pusilla* and *A. agrestis*.
Table S20. Upregulated genes in the vascular and nonvascular species investigated.
Table S21. Genes upregulated in the selected nonvascular and vascular plants under
nitrogen/phosphate starvation.
Table S22. Downregulated genes in the vascular and nonvascular species investigated.
Table S23. Genes downregulated in the selected nonvascular and vascular plants under
nitrogen/phosphate starvation
1020
1021
1022   In Supplementary_Figures
1023
Fig. S1. Expression clusters (maSigpro analysis) for *A. agrestis* at kcluster=2-9.
Fig. S2. Expression clusters (maSigpro analysis) for *B. pusilla* at kcluster=2-9.

Fig. S3. Functional enrichment analyses of cluster 1 under nitrogen starvation in *A. agrestis.*

Fig. S4. Functional enrichment analyses of cluster 2 under nitrogen starvation in *A. agrestis.*

Fig. S5 Functional enrichment analyses of cluster 3 under nitrogen starvation in *A. agrestis.*.

Fig. S6 Functional enrichment analyses of cluster 4 under nitrogen starvation in *A. agrestis.*

Fig. S7 Functional enrichment analyses of cluster 1 under nitrogen starvation in *B. pusilla.*

Fig. S8 Functional enrichment analyses of cluster 2 under nitrogen starvation in *B. pusilla*.

Fig. S9 Functional enrichment analyses of cluster 3 under nitrogen starvation in *B. pusilla*.

Fig. S10 Functional enrichment analyses of cluster 4 under nitrogen starvation in *B. pusilla*.

1034

1035   Supplementary_file_1 Description, statistics and quality assessment of the draft genome
1036   assembly.

1037   Supplementary_file_2 Description and quality assessment of the transcriptomic data set used to
1038   estimate gene expression for *B. pusilla*.

1039

1040
1041
1042
1043   **Author Contribution**

1044   PS and YY conceptualized the study. YY, GS and AN generated the primary sequence data. YY,
1045   PS and GS carried out the draft genome assembly. YY and PS carried out *de novo*-transcriptomic
1046   assembly, conducted detailed transcriptomic and comparative transcriptomic analyses. YY and
1047   PS wrote the manuscript. All co-authors revised and approved the final manuscript.
1048
1049
1050   **Conflict of Interest**
1051
1052   No conflict of interest declared.

1053

Funding Statement

This study was financially supported by grants of the Swiss National Science Foundation (grant
nos. 131726, 160004, 184826 and 212509 to PS), a pilot grant of the University Research Priority
Program "Evolution in Action" of the University of Zurich to PS and YY, a Georges and Antoine
Claraz Foundation grant to YY and PS. This project was also carried out in the framework of MAdLand
(https://madland.science, DFG priority programme 2237, PS-1111/1 to PS, RE 1697/19-1 and 20-1 to SAR).
Academy of Finland grant 295595 to JH enabled genome sequencing of Blasia pusilla by GS.

1061

Data Availability

Raw DNA and RNA sequencing data used in this publication was submitted to NCBI Short Read Archive (SRA) under the BioProject ID PRJNA1099914 (SRA submission SUB14374705, SUB14439959, SUB14439962). The transcriptome reference and the draft genome assemblies are available on Figshare (will be replaced by DOI once published). Supplementary tables are also available on Figshare (will be replaced by DOI once published).

Acknowledgments

We are grateful to Elke Dittmann for providing *Blasia pusilla* and cyanobacteria cultures. We are thankful to Jessy Zhang for assistance in sequencing at the Novogene UK. We are also thankful for the S3IT team and the ScienceCloud infrastructure at the University of Zurich for providing computational resources. JH and G thank CSC IT center for science (Espoo, Finland) for computational resources and advise in Finland. We also appreciate early discussions about the topic with John Bowman and Charles A. Delwiche.

References

Abdel-Lateif, K., Bogusz, D., & Hocher, V. (2012). The role of flavonoids in the establishment of plant roots endosymbioses with arbuscular mycorrhiza fungi, rhizobia and Frankia bacteria. *Plant signaling & behavior*, *7*(6), 636–641.

Akiyama, K., Matsuzaki, K., & Hayashi, H. (2005). Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi. *Nature*, *435*(7043), 824–827.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of molecular biology*, *215*(3), 403–410.

Adams, D.G. (2002). Cyanobacteria in Symbiosis with Hornworts and Liverworts. In: Rai, A.N., Bergman, B., Rasmussen, U. (eds) Cyanobacteria in Symbiosis. Springer, Dordrecht. pp. 117-135.

Adams, D. G., & Duggan, P. S. (2008). Cyanobacteria-bryophyte symbioses. *Journal of experimental botany*, *59*(5), 1047–1058.

Adams, D.G., Duggan, P.S. (2012). Signalling in Cyanobacteria–Plant Symbioses. In: Perotto, S., Baluška, F. (eds) Signaling and Communication in Plant Symbiosis. Signaling and Communication in Plants, vol 11. Springer, Berlin, Heidelberg. pp 93-121.

Adavi SB, Sathee L. Calcium regulates primary nitrate response associated gene transcription in a time- and dose-dependent manner. Protoplasma. 2024 Mar;261(2):257-269.

Aluko, O. O., Kant, S., Adedire, O. M., Li, C., Yuan, G., Liu, H., & Wang, Q. (2023). Unlocking the potentials of nitrate transporters at improving plant nitrogen use efficiency. *Frontiers in plant science*, *14*, 1074839.

Alvarenga, D. O., Elmdam, I. V., Timm, A. B., & Rousk, K. (2023). Chemical Stimulation of Heterocyte Differentiation by the Feather Moss Hylocomium splendens: a Potential New Step in Plant-Cyanobacteria Symbioses. *Microbial ecology*, *86*(1), 419–430.

Álvarez, C., Jiménez-Ríos, L., Iniesta-Pallarés, M., Jurado-Flores, A., Molina-Heredia, F. P., Ng, C. K. Y., & Mariscal, V. (2023). Symbiosis between cyanobacteria and plants: from molecular studies to agronomic applications. *Journal of experimental botany*, *74*(19), 6145–6157.

Baerenfaller, K., Grossmann, J., Grobei, M. A., Hull, R., Hirsch-Hoffmann, M., Yalovsky, S., Zimmermann, P., Grossniklaus, U., Gruissem, W., & Baginsky, S. (2008). Genome-scale proteomics reveals Arabidopsis thaliana gene models and proteome dynamics. *Science (New York, N.Y.)*, *320*(5878), 938–941.

Balzergue, C., Chabaud, M., Barker, D. G., Bécard, G., & Rochange, S. F. (2013). High phosphate reduces host ability to develop arbuscular mycorrhizal symbiosis without affecting root calcium spiking responses to the fungus. *Frontiers in plant science*, *4*, 426.

Bergman, B., Matveyev, A.V., & Rasmussen, U. (1996). Chemical signalling in cyanobacterial-plant symbioses. *Trends in Plant Science, 1*, 191-197.

Bucchini, F., Del Cortona, A., Kreft, Ł., Botzki, A., Van Bel, M., & Vandepoele, K. (2021). TRAPID 2.0: a web application for taxonomic and functional analysis of de novo transcriptomes. *Nucleic acids research*, *49*(17), e101.

Chaisson, M. J., & Tesler, G. (2012). Mapping single molecule sequencing reads using basic local alignment with successive refinement (BLASR): application and theory. *BMC bioinformatics*, *13*, 238.

Campbell, E. L., & Meeks, J. C. (1989). Characteristics of Hormogonia Formation by Symbiotic Nostoc spp. in Response to the Presence of Anthoceros punctatus or Its Extracellular Products. *Applied and environmental microbiology*, *55*(1), 125–131.

Campbell, E. L., & Meeks, J. C. (1992). Evidence for Plant-Mediated Regulation of Nitrogenase Expression in the Anthoceros-Nostoc Symbiotic Association. Journal of General Microbiology, 138, 473–480.

Carrell, A. A., Veličković, D., Lawrence, T. J., Bowen, B. P., Louie, K. B., Carper, D. L., Chu, R. K., Mitchell, H. D., Orr, G., Markillie, L. M., Jawdy, S. S., Grimwood, J., Shaw, A. J., Schmutz, J., Northen, T. R., Anderton, C. R., Pelletier, D. A., & Weston, D. J. (2022). Novel metabolic interactions and environmental conditions mediate the boreal peatmoss-cyanobacteria mutualism. *The ISME journal*, *16*(4), 1074–1085.

Chardin, C., Girin, T., Roudier, F., Meyer, C., & Krapp, A. (2014). The plant RWP-RK transcription factors: key regulators of nitrogen responses and of gametophyte development. *Journal of experimental botany*, *65*(19), 5577–5587.

Chatterjee, P., Schafran, P., Li, F. W., & Meeks, J. C. (2022). *Nostoc* Talks Back: Temporal Patterns of Differential Gene Expression During Establishment of *Anthoceros-Nostoc* Symbiosis. *Molecular plant-microbe interactions : MPMI*, *35*(10), 917–932.

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics (Oxford, England)*, *34*(17), i884–i890.

Cohen, M. F., & Yamasaki, H. (2000). Flavonoid-induced expression of a symbiosis-related gene in the cyanobacterium Nostoc punctiforme. *Journal of bacteriology*, *182*(16), 4644–4646.

Cove, D. J., Perroud, P. F., Charron, A. J., McDaniel, S. F., Khandelwal, A., & Quatrano, R. S. (2009). Culturing the moss Physcomitrella patens. *Cold Spring Harbor protocols*, *2009*(2), pdb.prot5136.

Crombez, H., Motte, H., & Beeckman, T. (2019). Tackling Plant Phosphate Starvation by the Roots. *Developmental cell*, *48*(5), 599–615.

Curci, P. L., Aiese Cigliano, R., Zuluaga, D. L., Janni, M., Sanseverino, W., & Sonnante, G. (2017). Transcriptomic response of durum wheat to nitrogen starvation. *Scientific reports*, *7*(1), 1176.

Das, D., Paries, M., Hobecker, K., Gigl, M., Dawid, C., Lam, H. M., Zhang, J., Chen, M., & Gutjahr, C. (2022). PHOSPHATE STARVATION RESPONSE transcription factors enable arbuscular mycorrhiza symbiosis. *Nature communications*, *13*(1), 477.

Dash, M., Yordanov, Y. S., Georgieva, T., Kumari, S., Wei, H., & Busov, V. (2015). A systems biology approach identifies new regulators of poplar root development under low nitrogen. *The Plant journal : for cell and molecular biology*, *84*(2), 335–346.

Delaux, P. M., & Schornack, S. (2021). Plant evolution driven by interactions with symbiotic and pathogenic microbes. *Science (New York, N.Y.)*, *371*(6531), eaba6605.

Diédhiou, I., & Diouf, D. (2018). Transcription factors network in root endosymbiosis establishment and development. *World journal of microbiology & biotechnology*, *34*(3), 37.

Ding, P., & Ding, Y. (2020). Stories of Salicylic Acid: A Plant Defense Hormone. *Trends in plant science*, *25*(6), 549–565.

Dong, Q., Wallrad, L., Almutairi, B. O., & Kudla, J. (2022). Ca^{2+} signaling in plant responses to abiotic stresses. *Journal of integrative plant biology*, *64*(2), 287–300.

Dubey, R.S., Srivastava, R.K., & Pessarakli, M. (2014). Physiological Mechanisms of Nitrogen Absorption and Assimilation in Plants under Stressful Conditions. *Handbook of Plant and Crop Physiology*.n.pag

Duggan, P. S., Thiel, T., & Adams, D. G. (2013). Symbiosis Between the Cyanobacterium Nostoc and the Liverwort Blasia Requires a CheR-type MCP Methyltransferase. Symbiosis, 59, 111–120.

Ekman, M., Picossi, S., Campbell, E. L., Meeks, J. C., & Flores, E. (2013). A Nostoc punctiforme sugar transporter necessary to establish a Cyanobacterium-plant symbiosis. *Plant physiology*, *161*(4), 1984–1992.

Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome biology*, *20*(1), 238.

Enderlin, C. S., & Meeks, J. C. (1983). Pure culture and reconstitution of the Anthoceros-Nostoc symbiotic association. *Planta*, *158*(2), 157–165.

Fan, X., Naz, M., Fan, X., Xuan, W., Miller, A. J., & Xu, G. (2017). Plant nitrate transporters: from gene function to application. *Journal of experimental botany*, *68*(10), 2463–2475.

Fang, Y., Qin, X., Liao, Q., Du, R., Luo, X., Zhou, Q., Li, Z., Chen, H., Jin, W., Yuan, Y., Sun, P., Zhang, R., Zhang, J., Wang, L., Cheng, S., Yang, X., Yan, Y., Zhang, X., Zhang, Z., Bai, S., … Yan, J. (2022). The genome of homosporous maidenhair fern sheds light on the euphyllophyte evolution and defences. *Nature plants*, *8*(9), 1024–1037.

Frangedakis, E., Waller, M., Nishiyama, T., Tsukaya, H., Xu, X., Yue, Y., Tjahjadi, M., Gunadi, A., Van Eck, J., Li, F. W., Szövényi, P., & Sakakibara, K. (2021). An Agrobacterium-mediated stable transformation technique for the hornwort model Anthoceros agrestis. *The New phytologist*, *232*(3), 1488–1505.

Gaudinier, A., Rodriguez-Medina, J., Zhang, L., Olson, A., Liseron-Monfils, C., Bågman, A. M., Foret, J., Abbitt, S., Tang, M., Li, B., Runcie, D. E., Kliebenstein, D. J., Shen, B., Frank, M. J., Ware,

D., & Brady, S. M. (2018). Transcriptional regulation of nitrogen-associated metabolism and growth. *Nature*, *563*(7730), 259–264.

Gobbato, E., Marsh, J. F., Vernié, T., Wang, E., Maillet, F., Kim, J., Miller, J. B., Sun, J., Bano, S. A., Ratet, P., Mysore, K. S., Dénarié, J., Schultze, M., & Oldroyd, G. E. (2012). A GRAS-type transcription factor with a specific function in mycorrhizal signaling. *Current biology : CB*, *22*(23), 2236–2241.

Gorelova, O.A., Baulina, O.I. (2009). Ultrastructure of cyanobacterium *Nostoc* sp. f. *Blasia* cell forms in persisting populations. *Microbiology* **78**, 609–617

Guillory, A., & Bonhomme, S. (2021). Phytohormone biosynthesis and signaling pathways of mosses. *Plant molecular biology*, *107*(4-5), 245–277.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., … Regev, A. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature biotechnology*, *29*(7), 644–652.

Haas, B. J., Salzberg, S. L., Zhu, W., Pertea, M., Allen, J. E., Orvis, J., White, O., Buell, C. R., & Wortman, J. R. (2008). Automated eukaryotic gene structure annotation using EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome biology*, *9*(1), R7.

Hashidoko, Y., Nishizuka, H., Tanaka, M., Murata, K., Murai, Y., & Hashimoto, M. (2019). Isolation and characterization of 1-palmitoyl-2-linoleoyl-sn-glycerol as a hormogonium-inducing factor (HIF) from the coralloid roots of Cycas revoluta (Cycadaceae). *Scientific reports*, *9*(1), 4751.

He, Q., Lu, H., Guo, H., Wang, Y., Zhao, P., Li, Y., Wang, F., Xu, J., Mo, X., & Mao, C. (2021). OsbHLH6 interacts with OsSPX4 and regulates the phosphate starvation response in rice. *The Plant journal : for cell and molecular biology*, *105*(3), 649–667.

He, X., Qu, B., Li, W., Zhao, X., Teng, W., Ma, W., Ren, Y., Li, B., Li, Z., & Tong, Y. (2015). The Nitrate-Inducible NAC Transcription Factor TaNAC2-5A Controls Nitrate Response and Increases Wheat Yield. *Plant physiology*, *169*(3), 1991–2005.

Hernández-García, J., Sun, R., Serrano-Mislata, A., Inoue, K., Vargas-Chávez, C., Esteve-Bruna, D., Arbona, V., Yamaoka, S., Nishihama, R., Kohchi, T., & Blázquez, M. A. (2021). Coordination between growth and stress responses by DELLA in the liverwort Marchantia polymorpha. *Current biology : CB*, *31*(16), 3678–3686.e11.

Hernández, G., Ramírez, M., Valdés-López, O., Tesfaye, M., Graham, M. A., Czechowski, T., Schlereth, A., Wandrey, M., Erban, A., Cheung, F., Wu, H. C., Lara, M., Town, C. D., Kopka, J., Udvardi, M. K., & Vance, C. P. (2007). Phosphorus stress in common bean: root transcript and metabolic responses. *Plant physiology*, *144*(2), 752–767.

Hsin, K. T., Yang, T. J., Lee, Y. H., & Cheng, Y. S. (2021). Phylogenetic and Structural Analysis of NIN-Like Proteins With a Type I/II PB1 Domain That Regulates Oligomerization for Nitrate Response. *Frontiers in plant science*, *12*, 672035.

Isidra-Arellano, M. C., Delaux, P. M., & Valdés-López, O. (2021). The Phosphate Starvation Response System: Its Role in the Regulation of Plant-Microbe Interactions. *Plant & cell physiology*, *62*(3), 392–400.

Jarratt-Barnham, E., Zarrabian, D., & Oldroyd, G. E. D. (2022). Symbiotic regulation: How plants seek salvation in starvation. *Current biology : CB*, *32*(1), R46–R48.

Jhu, M. Y., & Oldroyd, G. E. D. (2023). Dancing to a different tune, can we switch from chemical to biological nitrogen fixation for sustainable food security?. *PLoS biology*, *21*(3), e3001982.

Jones, V. A., & Dolan, L. (2012). The evolution of root hairs and rhizoids. *Annals of botany*, *110*(2), 205–212.

Kuznetsov, D., Tegenfeldt, F., Manni, M., Seppey, M., Berkeley, M., Kriventseva, E. V., & Zdobnov, E. M. (2023). OrthoDB v11: annotation of orthologs in the widest sampling of organismal diversity. *Nucleic acids research*, *51*(D1), D445–D451.

Kiba, T., Kudo, T., Kojima, M., & Sakakibara, H. (2011). Hormonal control of nitrogen acquisition: roles of auxin, abscisic acid, and cytokinin. *Journal of experimental botany*, *62*(4), 1399–1409.

Kimura, J., & Nakano, T. (1990). Reconstitution of a Blasia-Nostoc symbiotic association under axenic conditions. Nova Hedwigia, 50, 191–200.

Kodama, K., Rich, M. K., Yoda, A., Shimazaki, S., Xie, X., Akiyama, K., Mizuno, Y., Komatsu, A., Luo, Y., Suzuki, H., Kameoka, H., Libourel, C., Keller, J., Sakakibara, K., Nishiyama, T., Nakagawa, T., Mashiguchi, K., Uchida, K., Yoneyama, K., Tanaka, Y., … Kyozuka, J. (2022). An ancestral function of strigolactones as symbiotic rhizosphere signals. *Nature communications*, *13*(1), 3974.

Kolmogorov, M., Bickhart, D. M., Behsaz, B., Gurevich, A., Rayko, M., Shin, S. B., Kuhn, K., Yuan, J., Polevikov, E., Smith, T. P. L., & Pevzner, P. A. (2020). metaFlye: scalable long-read metagenome assembly using repeat graphs. *Nature methods*, *17*(11), 1103–1110.

Konishi, M., & Yanagisawa, S. (2013). Arabidopsis NIN-like transcription factors have a central role in nitrate signalling. *Nature communications*, *4*, 1617.

Kumar, A., Batra, R., Gahlaut, V., Gautam, T., Kumar, S., Sharma, M., Tyagi, S., Singh, K. P., Balyan, H. S., Pandey, R., & Gupta, P. K. (2018). Genome-wide identification and characterization of gene family for RWP-RK transcription factors in wheat (Triticum aestivum L.). *PloS one*, *13*(12), e0208409.

Laetsch DR and Blaxter ML. BlobTools: Interrogation of genome assemblies [version 1; peer review: 2 approved with reservations]. *F1000Research* 2017, **6**:1287

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature methods*, *9*(4), 357–359.

Lee, W. J., Truong, H. A., Trịnh, C. S., Kim, J. H., Lee, S., Hong, S. W., & Lee, H. (2020). NITROGEN RESPONSE DEFICIENCY 1-mediated CHL1 induction contributes to optimized growth performance during altered nitrate availability in Arabidopsis. *The Plant journal : for cell and molecular biology*, *104*(5), 1382–1398.

Li, W., He, X., Chen, Y., Jing, Y., Shen, C., Yang, J., Teng, W., Zhao, X., Hu, W., Hu, M., Li, H., Miller, A. J., & Tong, Y. (2020). A wheat transcription factor positively sets seed vigour by regulating the grain nitrate signal. *The New phytologist*, *225*(4), 1667–1680.

Li, F. W., Nishiyama, T., Waller, M., Frangedakis, E., Keller, J., Li, Z., Fernandez-Pozo, N., Barker, M. S., Bennett, T., Blázquez, M. A., Cheng, S., Cuming, A. C., de Vries, J., de Vries, S., Delaux, P. M., Diop, I. S., Harrison, C. J., Hauser, D., Hernández-García, J., Kirbis, A., … Szövényi, P. (2020). Anthoceros genomes illuminate the origin of land plants and the unique biology of hornworts. *Nature plants*, *6*(3), 259–272.

Liaimer, A., Helfrich, E. J., Hinrichs, K., Guljamow, A., Ishida, K., Hertweck, C., & Dittmann, E. (2015). Nostopeptolide plays a governing role during cellular differentiation of the symbiotic cyanobacterium Nostoc punctiforme. *Proceedings of the National Academy of Sciences of the United States of America*, *112*(6), 1862–1867.

Lin, Y., Chu, S., Xu, X., Han, X., Huang, H., Tong, Z., & Zhang, J. (2022). Identification of Nitrogen Starvation-Responsive miRNAs to Reveal the miRNA-Mediated Regulatory Network in *Betula luminifera*. *Frontiers in genetics*, *13*, 957505.

López-Bucio, J., Hernández-Abreu, E., Sánchez-Calderón, L., Nieto-Jacobo, M. F., Simpson, J., & Herrera-Estrella, L. (2002). Phosphate availability alters architecture and causes changes in hormone sensitivity in the Arabidopsis root system. *Plant physiology*, *129*(1), 244–256.

Lu, M., Cheng, Z., Zhang, X. M., Huang, P., Fan, C., Yu, G., Chen, F., Xu, K., Chen, Q., Miao, Y., Han, Y., Feng, X., Liu, L., & Fu, Y. F. (2020). Spatial Divergence of *PHR-PHT1* Modules Maintains Phosphorus Homeostasis in Soybean Nodules. *Plant physiology*, *184*(1), 236–250.

Luginbuehl, L. H., Menard, G. N., Kurup, S., Van Erp, H., Radhakrishnan, G. V., Breakspear, A., Oldroyd, G. E. D., & Eastmond, P. J. (2017). Fatty acids in arbuscular mycorrhizal fungi are synthesized by the host plant. *Science (New York, N.Y.)*, *356*(6343), 1175–1178.

Luo, Z., Wang, J., Li, F., Lu, Y., Fang, Z., Fu, M., Mysore, K. S., Wen, J., Gong, J., Murray, J. D., & Xie, F. (2023). The small peptide CEP1 and the NIN-like protein NLP1 regulate NRT2.1 to mediate root nodule formation across nitrate concentrations. *The Plant cell*, *35*(2), 776–794.

Lv, X., Zhang, Y., Hu, L., Zhang, Y., Zhang, B., Xia, H., Du, W., Fan, S., & Kong, L. (2020). Low-Nitrogen Stress Stimulates Lateral Root Initiation and Nitrogen Assimilation in Wheat: Roles of Phytohormone Signaling. *Journal of Plant Growth Regulation, 40*, 436-450.

Laughlin, M. J., McGaughey, D. S., Crews, J. R., Chao, N. J., Rizzieri, D., Ross, M., Gockerman, J., Cirrincione, C., Berry, D., Mills, L., Defusco, P., LeGrand, S., Peters, W. P., & Vredenburgh, J. J.

(1998). Secondary myelodysplasia and acute leukemia in breast cancer patients after autologous bone marrow transplant. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*, *16*(3), 1008–1012.

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome biology*, *15*(12), 550.

Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone reads using repeat graphs. *Nature biotechnology*, *37*(5), 540–546.

Ma, Y., & Chen, R. (2021). Nitrogen and Phosphorus Signaling and Transport During Legume-Rhizobium Symbiosis. *Frontiers in plant science*, *12*, 683601.

Ma, N., Dong, L., Lü, W., Lü, J., Meng, Q., & Liu, P. (2020). Transcriptome Analysis of Maize Seedling Roots in Response to Nitrogen-, Phosphorus-, and Potassium Deficiency. Plant and Soil, 447, 637–658.

Mascia, M., Sega, D., Zamboni, A., & Varanini, Z. (2019). Nitrogen Starvation Differentially Influences Transcriptional and Uptake Rate Profiles in Roots of Two Maize Inbred Lines with Different NUE. *International journal of molecular sciences*, *20*(19), 4856.

Masclaux-Daubresse, C., Daniel-Vedele, F., Dechorgnat, J., Chardon, F., Gaufichon, L., & Suzuki, A. (2010). Nitrogen uptake, assimilation and remobilization in plants: challenges for sustainable and productive agriculture. *Annals of botany*, *105*(7), 1141–1157.

Medici, A., Marshall-Colon, A., Ronzier, E., Szponarski, W., Wang, R., Gojon, A., Crawford, N. M., Ruffel, S., Coruzzi, G. M., & Krouk, G. (2015). AtNIGT1/HRS1 integrates nitrate and phosphate signals at the Arabidopsis root tip. *Nature communications*, *6*, 6274.

Meeks, J. C. (1998). Symbiosis Between Nitrogen-Fixing Cyanobacteria and Plants. BioScience, 48, 266–276.

Meeks, J. C. (2003). Symbiotic Interactions Between Nostoc punctiforme, a Multicellular Cyanobacterium, and the Hornwort Anthoceros punctatus. Symbiosis, 35, 55–71.

Misson, J., Raghothama, K. G., Jain, A., Jouhet, J., Block, M. A., Bligny, R., Ortet, P., Creff, A., Somerville, S., Rolland, N., Doumas, P., Nacry, P., Herrera-Estrella, L., Nussaume, L., & Thibaud, M. C. (2005). A genome-wide transcriptional analysis using Arabidopsis thaliana Affymetrix gene chips determined plant responses to phosphate deprivation. *Proceedings of the National Academy of Sciences of the United States of America*, *102*(33), 11934–11939.

Monte, I., Franco-Zorrilla, J. M., García-Casado, G., Zamarreño, A. M., García-Mina, J. M., Nishihama, R., Kohchi, T., & Solano, R. (2019). A Single JAZ Repressor Controls the Jasmonate Pathway in Marchantia polymorpha. *Molecular plant*, *12*(2), 185–198.

Mu, X., & Luo, J. (2019). Evolutionary analyses of NIN-like proteins in plants and their roles in nitrate signaling. *Cellular and molecular life sciences : CMLS*, *76*(19), 3753–3764.

Nguyen, N. N. T., Clua, J., Vetal, P. V., Vuarambon, D. J., De Bellis, D., Pervent, M., Lepetit, M., Udvardi, M., Valentine, A. J., & Poirier, Y. (2021). PHO1 family members transport phosphate from infected nodule cells to bacteroids in Medicago truncatula. *Plant physiology*, *185*(1), 196–209.

Nilsson, M., Rasmussen, U., & Bergman, B. (2006). Cyanobacterial chemotaxis to extracts of host and nonhost plants. *FEMS microbiology ecology*, *55*(3), 382–390.

Nishizuka, H., & Hashidoko, Y. (2018). Comparison of Nostocean hormogonium induction and its motility on solid plates between agar and gellan gum at varying gel matrix concentrations. *Bioscience, biotechnology, and biochemistry*, *82*(3), 525–531.

Noguero, M., Atif, R. M., Ochatt, S., & Thompson, R. D. (2013). The role of the DNA-binding One Zinc Finger (DOF) transcription factor family in plants. *Plant science : an international journal of experimental plant biology*, *209*, 32–45.

Nueda, M. J., Tarazona, S., & Conesa, A. (2014). Next maSigPro: updating maSigPro bioconductor package for RNA-seq time series. *Bioinformatics (Oxford, England)*, *30*(18), 2598–2602.

Oldroyd G. E. (2013). Speak, friend, and enter: signalling systems that promote beneficial symbiotic associations in plants. *Nature reviews. Microbiology*, *11*(4), 252–263.

Oldroyd, G. E. D., & Leyser, O. (2020). A plant's diet, surviving in a variable nutrient environment. *Science (New York, N.Y.)*, *368*(6486), eaba0196.

Ondzighi-Assoume, C. A., Chakraborty, S., & Harris, J. M. (2016). Environmental Nitrate Stimulates Abscisic Acid Accumulation in Arabidopsis Root Tips by Releasing It from Inactive Stores. *The Plant cell*, *28*(3), 729–745.

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast and bias-aware quantification of transcript expression. *Nature methods*, *14*(4), 417–419.

Paries, M., & Gutjahr, C. (2023). The good, the bad, and the phosphate: regulation of beneficial and detrimental plant-microbe interactions by the plant phosphate status. *The New phytologist*, *239*(1), 29–46.

Pathuri, I. P., Reitberger, I. E., Hückelhoven, R., & Proels, R. K. (2011). Alcohol dehydrogenase 1 of barley modulates susceptibility to the parasitic fungus Blumeria graminis f.sp. hordei. *Journal of experimental botany*, *62*(10), 3449–3457.

Puginier, C., Keller, J., & Delaux, P. M. (2022). Plant-microbe interactions that have impacted plant terrestrializations. *Plant physiology*, *190*(1), 72–84.

Pederson, E. R. A., Warshan, D., & Rasmussen, U. (2019). Genome Sequencing of *Pleurozium schreberi*: The Assembled and Annotated Draft Genome of a Pleurocarpous Feather Moss. *G3 (Bethesda, Md.)*, *9*(9), 2791–2797.

Qi 杞金芳, J., Yu 郁露, L., Ding 丁静丽, J., Ji 姬晨晨, C., Wang 汪社亮, S., Wang 王创, C., Ding 丁广大, G., Shi 石磊, L., Xu 徐芳森, F., & Cai 蔡红梅, H. (2023). Transcription factor OsSNAC1 positively regulates nitrate transporter gene expression in rice. *Plant physiology*, *192*(4), 2923–2942.

Qu, B., He, X., Wang, J., Zhao, Y., Teng, W., Shao, A., Zhao, X., Ma, W., Wang, J., Li, B., Li, Z., & Tong, Y. (2015). A wheat CCAAT box-binding transcription factor increases the grain yield of wheat with less fertilizer input. *Plant physiology*, *167*(2), 411–423.

Radhakrishnan, G. V., Keller, J., Rich, M. K., Vernié, T., Mbadinga Mbadinga, D. L., Vigneron, N., Cottret, L., Clemente, H. S., Libourel, C., Cheema, J., Linde, A. M., Eklund, D. M., Cheng, S., Wong, G. K. S., Lagercrantz, U., Li, F. W., Oldroyd, G. E. D., & Delaux, P. M. (2020). An ancestral signalling pathway is conserved in intracellular symbioses-forming plant lineages. *Nature plants*, *6*(3), 280–289.

Rasmussen, U. (1994). Early Communication in the Gunnera - Nostoc Symbiosis: Plant-Induced Cell Differentiation and Protein Synthesis in the Cyanobacterium. Molecular Plant-Microbe Interactions, 7, 696.

Renzaglia, K. S., Rasch, E. M., & Pike, L. M. (1995). Estimates of Nuclear DNA Content in Bryophyte Sperm Cells: Phylogenetic Considerations. American Journal of Botany, 82, 18.

Rich, M. K., Vigneron, N., Libourel, C., Keller, J., Xue, L., Hajheidari, M., Radhakrishnan, G. V., Le Ru, A., Diop, S. I., Potente, G., Conti, E., Duijsings, D., Batut, A., Le Faouder, P., Kodama, K., Kyozuka, J., Sallet, E., Bécard, G., Rodriguez-Franco, M., Ott, T., … Delaux, P. M. (2021). Lipid exchanges drove the evolution of mutualism during plant terrestrialization. *Science (New York, N.Y.)*, *372*(6544), 864–868.

Rico-Resendiz, F.; Diaz-Santana, Z.H.U.; Dipp-Alvarez, M.; Cruz-Hernandez, A.; Bowman, J.L.; Herrera-Estrella, L.; Ishizaki, K.; Arteaga-Vazquez, M.A.; Cruz-Ramirez, A. Phosphate Starvation Triggers Transcriptional Changes in the Biosynthesis and Signaling Pathways of Phytohormones in Marchantia polymorpha . Biol. Life Sci. Forum 2021, 4, 89.

Rouached, H., Arpat, A. B., & Poirier, Y. (2010). Regulation of phosphate starvation responses in plants: signaling players and cross-talks. *Molecular plant*, *3*(2), 288–299.

Rubin, G., Tohge, T., Matsuda, F., Saito, K., & Scheible, W. R. (2009). Members of the LBD family of transcription factors repress anthocyanin synthesis and affect additional nitrogen responses in Arabidopsis. *The Plant cell*, *21*(11), 3567–3584.

Rui, W., Mao, Z., & Li, Z. (2022). The Roles of Phosphorus and Nitrogen Nutrient Transporters in the Arbuscular Mycorrhizal Symbiosis. *International journal of molecular sciences*, *23*(19), 11027.

Shi, J., Zhao, B., Jin, R., Hou, L., Zhang, X., Dai, H., Yu, N., & Wang, E. (2022). A phosphate starvation response-regulated receptor-like kinase, OsADK1, is required for mycorrhizal symbiosis and phosphate starvation responses. *The New phytologist*, *236*(6), 2282–2293.

Shi, J., Zhao, B., Zheng, S., Zhang, X., Wang, X., Dong, W., Xie, Q., Wang, G., Xiao, Y., Chen, F., Yu, N., & Wang, E. (2021). A phosphate starvation response-centered network regulates mycorrhizal symbiosis. *Cell*, *184*(22), 5527–5540.e18.

Soneson, C., Love, M. I., & Robinson, M. D. (2015). Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences. *F1000Research*, *4*, 1521.

Splitt, S. D., & Risser, D. D. (2016). The non-metabolizable sucrose analog sucralose is a potent inhibitor of hormogonium differentiation in the filamentous cyanobacterium Nostoc punctiforme. *Archives of microbiology*, *198*(2), 137–147.

Subudhi, P. K., Garcia, R. S., Coronejo, S., & Tapia, R. (2020). Comparative Transcriptomics of Rice Genotypes with Contrasting Responses to Nitrogen Stress Reveals Genes Influencing Nitrogen Uptake through the Regulation of Root Architecture. *International journal of molecular sciences*, *21*(16), 5759.

Sultana, N., Islam, S., Juhasz, A., Yang, R., She, M., Alhabbar, Z., Zhang, J., & Ma, W. (2020). Transcriptomic Study for Identification of Major Nitrogen Stress Responsive Genes in Australian Bread Wheat Cultivars. *Frontiers in genetics*, *11*, 583785.

Sun, L., Di, D. W., Li, G., Li, Y., Kronzucker, H. J., & Shi, W. (2020). Transcriptome analysis of rice (Oryza sativa L.) in response to ammonium resupply reveals the involvement of phytohormone signaling and the transcription factor OsJAZ9 in reprogramming of nitrogen uptake and metabolism. *Journal of plant physiology*, *246-247*, 153137.

Sun, Z., Ding, C., Li, X., & Xiao, K. (2012). Molecular Characterization and Expression Analysis of TaZFP15, a C2H2-Type Zinc Finger Transcription Factor Gene in Wheat (Triticum aestivum L.). Journal of Integrative Agriculture, 11, 31–42.

Suzuki, H., Kato, H., Iwano, M., Nishihama, R., & Kohchi, T. (2023). Auxin signaling is essential for organogenesis but not for cell survival in the liverwort Marchantia polymorpha. *The Plant cell*, *35*(3), 1058–1075.

Suzuki, H., Kohchi, T., & Nishihama, R. (2021). Auxin Biology in Bryophyta: A Simple Platform with Versatile Functions. *Cold Spring Harbor perspectives in biology*, *13*(3), a040055.

Szövényi, P., Frangedakis, E., Ricca, M., Quandt, D., Wicke, S., & Langdale, J. A. (2015). Establishment of Anthoceros agrestis as a model species for studying the biology of hornworts. *BMC plant biology*, *15*, 98.

Szövényi, P., Waller, M., & Kirbis, A. (2019). Evolution of the plant body plan. *Current topics in developmental biology*, *131*, 1–34.

Tan, Q. W., Lim, P. K., Chen, Z., Pasha, A., Provart, N., Arend, M., Nikoloski, Z., & Mutwil, M. (2023). Cross-stress gene expression atlas of Marchantia polymorpha reveals the hierarchy and regulatory principles of abiotic stress responses. *Nature communications*, *14*(1), 986.

Tang, W., Ye, J., Yao, X., Zhao, P., Xuan, W., Tian, Y., Zhang, Y., Xu, S., An, H., Chen, G., Yu, J., Wu, W., Ge, Y., Liu, X., Li, J., Zhang, H., Zhao, Y., Yang, B., Jiang, X., Peng, C., … Wan, J. (2019).

Genome-wide associated study identifies NAC42-activated nitrate transporter conferring high nitrogen use efficiency in rice. *Nature communications*, *10*(1), 5279.

Tian, F., Yang, D. C., Meng, Y. Q., Jin, J., & Gao, G. (2020). PlantRegMap: charting functional regulatory maps in plants. *Nucleic acids research*, *48*(D1), D1104–D1113.

Ueda, Y., Ohtsuki, N., Kadota, K., Tezuka, A., Nagano, A. J., Kadowaki, T., Kim, Y., Miyao, M., & Yanagisawa, S. (2020). Gene regulatory network and its constituent transcription factors that control nitrogen-deficiency responses in rice. *The New phytologist*, *227*(5), 1434–1452.

Ueda, Y., & Yanagisawa, S. (2019). Perception, transduction, and integration of nitrogen and phosphorus nutritional signals in the transcriptional regulatory network in plants. *Journal of experimental botany*, *70*(15), 3709–3717.

Varala, K., Marshall-Colón, A., Cirrone, J., Brooks, M. D., Pasquino, A. V., Léran, S., Mittal, S., Rock, T. M., Edwards, M. B., Kim, G. J., Ruffel, S., McCombie, W. R., Shasha, D., & Coruzzi, G. M. (2018). Temporal transcriptional logic of dynamic regulatory networks underlying nitrogen signaling and use in plants. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(25), 6494–6499.

Vidal, E. A., Álvarez, J. M., & Gutiérrez, R. A. (2014). Nitrate regulation of AFB3 and NAC4 gene expression in Arabidopsis roots depends on NRT1.1 nitrate transport function. *Plant signaling & behavior*, *9*(3), e28501.

Vidal, E. A., Moyano, T. C., Riveras, E., Contreras-López, O., & Gutiérrez, R. A. (2013). Systems approaches map regulatory networks downstream of the auxin receptor AFB3 in the nitrate response of Arabidopsis thaliana roots. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(31), 12840–12845.

de Vries, S., Herrfurth, C., Li, F. W., Feussner, I., & de Vries, J. (2023). An ancient route towards salicylic acid and its implications for the perpetual Trichormus-Azolla symbiosis. *Plant, cell & environment*, *46*(9), 2884–2908.

de Vries, S., & de Vries, J. (2022). Evolutionary genomic insights into cyanobacterial symbioses in plants. *Quantitative plant biology*, *3*, e16.

de Vries, S., de Vries, J., Teschke, H., von Dahlen, J. K., Rose, L. E., & Gould, S. B. (2018). Jasmonic and salicylic acid response in the fern Azolla filiculoides and its cyanobiont. *Plant, cell & environment*, *41*(11), 2530–2548.

Wang, Y. Y., Cheng, Y. H., Chen, K. E., & Tsay, Y. F. (2018). Nitrate Transport, Signaling, and Use Efficiency. *Annual review of plant biology*, *69*, 85–122.

Wang, Q., Huang, Y., Ren, Z., Zhang, X., Ren, J., Su, J., Zhang, C., Tian, J., Yu, Y., Gao, G. F., Li, L., & Kong, Z. (2020). Transfer cells mediate nitrate uptake to control root nodule symbiosis. *Nature plants*, *6*(7), 800–808.

Wang, J., Song, K., Sun, L., Qin, Q., Sun, Y., Pan, J., & Xue, Y. (2019). Morphological and Transcriptome Analysis of Wheat Seedlings Response to Low Nitrogen Stress. *Plants (Basel, Switzerland)*, *8*(4), 98.

Wang, D., Xu, T., Yin, Z., Wu, W., Geng, H., Li, L., Yang, M., Cai, H., & Lian, X. (2020). Overexpression of OsMYB305 in Rice Enhances the Nitrogen Uptake Under Low-Nitrogen Condition. *Frontiers in plant science*, *11*, 369.

Wu, Z., Liu, H., Huang, W., Yi, L., Qin, E., Yang, T., Wang, J., & Qin, R. (2020). Genome-Wide Identification, Characterization, and Regulation of RWP-RK Gene Family in the Nitrogen-Fixing Clade. *Plants (Basel, Switzerland)*, *9*(9), 1178.

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C. A., Zeng, Q., Wortman, J., Young, S. K., & Earl, A. M. (2014). Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PloS one*, *9*(11), e112963.

Xu, P., & Wang, E. (2023). Diversity and regulation of symbiotic nitrogen fixation in plants. *Current biology : CB*, *33*(11), R543–R559.

Xue, L., Cui, H., Buer, B., Vijayakumar, V., Delaux, P. M., Junkermann, S., & Bucher, M. (2015). Network of GRAS transcription factors involved in the control of arbuscule development in Lotus japonicus. *Plant physiology*, *167*(3), 854–871.

Xiao, C. L., Chen, Y., Xie, S. Q., Chen, K. N., Wang, Y., Han, Y., Luo, F., & Xie, Z. (2017). MECAT: fast mapping, error correction, and de novo assembly for single-molecule sequencing reads. *Nature methods*, *14*(11), 1072–1074.

Yanagisawa, S., Akiyama, A., Kisaka, H., Uchimiya, H., & Miwa, T. (2004). Metabolic engineering with Dof1 transcription factor in plants: Improved nitrogen assimilation and growth under low-nitrogen conditions. *Proceedings of the National Academy of Sciences of the United States of America*, *101*(20), 7833–7838.

Yang, T., Hao, L., Yao, S., Zhao, Y., Lu, W., & Xiao, K. (2016). TabHLH1, a bHLH-type transcription factor gene in wheat, improves plant tolerance to Pi and N deprivation via regulation of nutrient transporter gene transcription and ROS homeostasis. *Plant physiology and biochemistry : PPB*, *104*, 99–113.

Yang, J., Wang, M., Li, W., He, X., Teng, W., Ma, W., Zhao, X., Hu, M., Li, H., Zhang, Y., & Tong, Y. (2019). Reducing expression of a nitrate-responsive bZIP transcription factor increases grain yield and N use in wheat. *Plant biotechnology journal*, *17*(9), 1823–1833.

Yang, W., Yoon, J., Choi, H., Fan, Y., Chen, R., & An, G. (2015). Transcriptome analysis of nitrogen-starvation-responsive genes in rice. *BMC plant biology*, *15*, 31.

Zhang, L., Zhang, L., Xia, C., Zhao, G., Liu, J., Jia, J., & Kong, X. (2015). A novel wheat bZIP transcription factor, TabZIP60, confers multiple abiotic stress tolerances in transgenic Arabidopsis. *Physiologia plantarum*, *153*(4), 538–554.

Zhang, L., Zhao, G., Jia, J., Liu, X., & Kong, X. (2012). Molecular characterization of 60 isolated wheat MYB genes and analysis of their expression during abiotic stress. *Journal of experimental botany*, *63*(1), 203–214.

Zipfel, C., & Oldroyd, G. E. (2017). Plant signalling in symbiosis and immunity. *Nature*, *543*(7645), 328–336.

Zimin, A. V., Marçais, G., Puiu, D., Roberts, M., Salzberg, S. L., & Yorke, J. A. (2013). The MaSuRCA genome assembler. *Bioinformatics (Oxford, England)*, *29*(21), 2669–2677.