Causal network perturbation analysis identifies known and novel type-2 diabetes driver genes

Yue Zhao[1*], Hash Brown Taha[2*], Ansarullah[3*], Kailiang Qian[2], Parveen Kumar[1], J. Matthew Mahoney[3],

Hao He[1], Joshy George[1, §], Sheng Li[1,2, 3,4,§]

[1]The Jackson Laboratory for Genomic Medicine, Farmington, CT, USA.

[2]Department of Cancer Biology, Keck School of Medicine, Norris Comprehensive Cancer Center,

University of Southern California, Los Angeles, CA, USA

[3]Center for Biometric Analysis, The Jackson Laboratory, Bar Harbor, ME, USA

[4]The Jackson Laboratory for Mouse Genetics, Bar Harbor, ME, USA.

[5]Department of Genetics and Genome Sciences, University of Connecticut School of Medicine,

Farmington, CT, USA.

[*]*These authors contributed equally to this work and are co-first authors.*

[§]Correspondence should be addressed to

Dr. Joshy George: joshy.george@jax.org;

Dr. Sheng Li : sli68423@usc.edu

Keywords: ssNPA, meta-cell, type-2 diabetes, β-cell, causal network perturbation

Abstract

The molecular pathogenesis of type 2 diabetes mellitus (T2DM) involves environmental and genetic factors that remain poorly understood. Increased insulin resistance leads to pancreatic β-islet cell exhaustion and failure, leading to more severe T2DM. As such, targeting β-cell dysfunction is an attractive pathway for T2DM treatment. Single-cell RNA sequencing can help identify differentially expressed genes (DEGs) that may contribute to disease-associated pathophysiology, but cannot capture the full spectrum of disease pathophysiology, as some genes that are not differentially expressed may still play important roles in disease progression or mechanisms. As such, we investigated single-cell gene expression changes in β-cells from healthy (C57BL/6J) and diabetic (NZO/HlLtJ) mice fed with normal or high-fat, high-sugar diet (HFHS) across the spectrum of T2DM development (prediabetic, mild or severely diabetic) using an innovative integration of the causal network perturbation assessment (ssNPA) framework with meta-cell transcriptome analysis to identify driver genes for T2DM. By generating a reference causal network and conducting *in silico* perturbations, we identified three classes of genes implicated in T2DM pathophysiology: (1) DEGs that were not perturbed (e.g, *Igf2bp2* strongly linked to β cell dysfunction and glucose regulation), (2) perturbed non-DEGs (e.g., *Glp1r* a therapeutic target of modern diabetes drugs) that were perturbed, and (3) DEGs that were perturbed (e.g., *Pdx1* a master β cell transcription factor), and validated some of the targets using the KOMP database. Together, these results show that each analytic layer captures complementary mechanisms underlying T2DM severity, and their synthesis provides a more complete view of the disease pathophysiology.

Introduction

Type 2 diabetes mellitus (T2DM) comprises more than 90% of all diabetes cases, impacting approximately 6.8% (537 million individuals) of the global population in the year 2021[1,2]. T2DM is caused by a complex interplay between environmental and genetic factors. However, while some of the major environmental factors (i.e., diet and physical activity) are known, the genetic bases of T2DM remain poorly understood. The endocrine portion of the pancreas is constituted by highly specialized hormone-secreting entities known as islets of Langerhans. The islets comprise α, β, delta, PP, and ghrelin cells that secrete glucagon, insulin, somatostatin, pancreatic polypeptide, and ghrelin, respectively[3]. The onset of full-blown T2DM occurs when β cells suffer from failure and exhaustion leading to the loss of their capacity to secrete appropriate amounts of insulin in response to elevated blood glucose levels as a result of insulin resistance in peripheral tissues.

Current research insights on T2DM are limited due to the variability in environmental and genetic factors among diverse populations. Additionally, there is a notable gap in long-term clinical trials that comprehensively assess the efficacy of emerging treatments across different stages of the disease. In recent years, single-cell RNA sequencing (scRNA-seq) has proven critical for investigating comprehensive gene expression profiles, revealing the presence of heterogeneous gene expression patterns, even within the same cell-types[4]. Furthermore, diverse phenotypes of pancreatic β cells have been observed within a single islet[5-7]. The application of scRNA-seq has significantly contributed to our understanding of β-cell maturation, β-cell heterogeneity, β-cell failure, and β-cell function in both healthy and diseased states. However, scRNA cannot capture the full spectrum of disease pathophysiology, as some genes that are not differentially expressed may still play important roles in disease progression or mechanisms. Additionally, no study to date has delineated the gene changes associated with clinical transitions across T2DM's spectrum by comparing prediabetic, mildly, and severely diabetic stages.

To answer this gap, we used two mouse lines (C57BL/6J and NZO/HILTJ) that were kept on a standard or high-fat high-sugar (HFHS) diet regimen to create the following phenotypes: healthy/non-diabetic (C56BL/6J on standard diet), prediabetic (C56BL/6J on HFHS diet), mildly diabetic (NZO/HILTJ on standard diet) or severely diabetic (NZO/HlLtJ on HFHS diet). At the end of the treatment, we performed scRNA on isolated β-cells to make a comparison between healthy and prediabetic states;

prediabetic and severely diabetic states and mildly diabetic and severely diabetic states (**Figure 1A-B**).

Additionally, since conventional differential expression analyses do not effectively detect certain type of

disruptions in gene networks, we   performed a causal network perturbation assessment (ssNPA)

framework[8] in combination with a meta-cell transcriptome analysis (meta-ssNPA) (**Figure 1C**). By

generating a reference causal network and conducting *in silico* perturbations, we identified three classes

of genes implicated in T2DM pathophysiology: (1) non-perturbed DEGs, (2) perturbed non-DEGs and

(3) perturbed DEGs, and validated our candidates using the Knockout Mouse Phenotyping (KOMP)

Project database.

Our findings reveal that integrating differential expression, perturbation, and shared DEG-perturbed

analyses provides a comprehensive understanding of T2DM severity mechanisms. The DEG-only layer

highlighted genes such as *Igf2bp2*[9] that regulate insulin signaling and glucose metabolism, while the

perturbed-only layer identified *Glp1r* across all three comparisons, a clinically relevant therapeutic

target modulating β cell function[10]. The perturbed-DEG layer captured core regulatory genes like

Pdx1[11], essential for insulin synthesis and secretion. Together, these complementary layers illustrate

how distinct molecular processes converge to drive β cell dysfunction and metabolic imbalance in

T2DM.

Material & Methods

Animal studies and islet isolation

In strict adherence to the standards set by the Association for Assessment and Accreditation of Laboratory Animal Care, our facility at The Jackson Laboratory has upheld the care and treatment of mice. We acquired male and female mice from two distinct strains: C57BL/6J (B6; RRID:IMSR_JAX:000664), and NZO/HlLtJ (NZO; RRID:IMSR_JAX:002105), starting at the age of four weeks. The mice were provided with two types of diets from Research Diets: a high-fat, high-sucrose diet (HFHS, comprising 44% kcal from fat and 1360 kcal from sucrose; formula D19070208) and a control diet (10% kcal from fat and devoid of sucrose; formula D19072203), both containing equal fiber content. The diets were given *ad libitum* starting from the age of six weeks. At the age of fifteen weeks, islet isolation was performed, and mice were euthanized through cervical dislocation and the common bile duct at the Sphincter of Oddi was clamped. Collagenase solution (three mL of a solution containing collagenase P (5 units/ml) and DNaseI (1mg/ml) in Hank's Balanced Salt Solution (HBSS)) was inserted into the bile duct proximal to the final bifurcation leading to the liver to inflate the pancreas. We then removed the pancreas for digestion by the collagenase solution at 37°C for 40 minutes. Post-digestion, samples were agitated for ten seconds, diluted it with 10 mL of HBSS, and centrifuged for 3 minutes at 300 RPM. After two washes with HBSS, the pellet was resuspended in 5 mL HBSS, and handpicked islets were collected using a clean petri dish containing HBSS. The islets were then transferred to 24-well plates with 1 mL of the warmed media (containing RPMI 1640, 10% FBS, glutamine, and HEPES) and incubated overnight at 37°C. After overnight incubation islets were centrifuged and the supernatant was discarded. Finally, islets were resuspended in a 1-2 mL of preheated (37°C) StemPro Accutase dissociation solution (Fisher Scientific, cat # A1110501). The cell suspension was gently pipetted for 30 seconds to facilitate the dissociation of islets until the media appeared translucent and there were no visible clumps, usually within 2-5 minutes. Following dissociation, 2-3 mL of RPMI complete medium was added, and the cell suspension was filtered through a 20 μm strainer. The cells were then centrifuged at 230 RCF for 3 minutes, the supernatant was removed, and the cells were resuspended in 2-3 mL of RPMI complete medium.

Preprocessing of Single Cell RNA-Seq

We processed raw fastq reads from Illumina sequencing for scRNA-Seq by aligning them to the mouse reference genome (mm10/GRCm38) using 10X Cell Ranger version 6.1.1 with standard settings. To demultiplex the strain of origin from samples containing mixed strains, we employed demuxlet version 2 (https://github.com/statgen/popscle)[12], utilizing known genomic variations from VCF files obtained from the Sanger Mouse Genomes Project[13]. We focused on sites that were biallelic and varied among our two strains (B6, and NZO). We ran demuxlet on scRNA-seq data with specific parameters "--α 0.0 -- α 0.5 --tag-group CB --tag-UMI UB --field GT", accounting for all cells identified by Cell Ranger.

Quality Control and Filtering of Single Cell RNA-Seq Data

Post-processing, we applied quality control measures to the feature count matrices generated by Cell Ranger. Single cells with fewer than 500 (islet) genes, more than 20% (islet) mitochondrial transcripts, or over 50% ribosomal transcripts were excluded. Additionally, genes not detected in at least three single cells per sequenced library were also omitted. To correct for potential ambient RNA contamination in islet samples, we utilized decontX from the celda V1.10.0 package, adhering to developer guidelines on GitHub. The SCDS V1.10.0 package was deployed to eliminate cell doublets, using default settings.

Clustering and Identification of Cell Types in Single Cell RNA-Seq

Following the filtration and quality control, single cells/nuclei underwent normalization and were clustered using Seurat version 4.1, with batch variations across libraries corrected by harmony version 0.1.0. Gene expression data from single cells were normalized based on library size and log-transformed. Dimensionality was reduced using principal component analysis (PCA) on the 2,500 most variable genes, and these principal components (PCs) underwent batch correction using harmony. The batch-corrected PCs were then used for Louvain-based clustering, with the resolution parameter adjusted between 0.1 and 1 according to the dataset specifics.

We identified differential marker genes for various groups via the Wilcoxon rank-sum test or MAST within the Seurat package, employing an false discovery rate (FDR) cutoff of 0.1 and a fold change

cutoff of 1.5. Cluster-specific genes aided in assigning cell types. Independently, cell type assignment

was also performed unbiasedly using the SingleR package, comparing each single-cell transcriptome to

reference transcriptome profiles of known cell types. For formal differential gene expression tests, we

created pseudobulks within cell types by aggregating gene expression counts across all cells of a given

type from a single mouse, applying DESeq2 on the pseudobulks. This approach using negative binomial

modeling with pseudobulks derived from single-cell transcriptomic data is noted for its efficacy and

accuracy in differential expression testing in single-cell contexts[14]. Low-expression genes have been

filtered out the OGFSC algorithm[15].

Refine the existing Single Sample Network Perturbation Assessment (ssNPA) method

There are many limitations that still exist for the current ssNPA framework[8]. First, the framework is not

well suited to handle single cell data. The single cell data have sparsity issues where a lot of genes have

zero expressions in many cells. This violates the assumption of the linear model ssNPA is using and thus

hinders accurate prediction performance. We tackle this problem by a simple meta-cell idea[16]. We

randomly pick several cells from the same cell type and average the gene expressions as a new meta-cell

sample. This approach maintains the relative level of all gene expression in the cell (high expression

genes are still higher and lower expression genes are still lower in the meta-cells) while removing the

zeros. Secondly, the current ssNPA framework uses a simple criterion to filter out Perturbed genes: as

long as the average Perturbance Score is higher in the test group than the control group, the gene is

annotated as a Perturbed Gene. This approach does not consider the random noise in the Perturbance

Score, thus induces false positive results. We refined this process by applying Wilcoxon test[17] onto test

and control group. Perturbance Score and use a FDR < 0.05 as the threshold to filter the Perturbed Genes

where statistical significance is introduced in the comparison. We performed one sided Wilcoxon test

with the alternative hypothesis that the control group Perturbance Score $<$ test group Perturbance Score.

Workflow and experimental design to detect perturbed genes via meta-cell local reference causal

network

Based on the discussion above, we propose the following meta-ssNPA workflow (**Figure 1**): the

scRNA-seq data from various sample groups underwent standard processing using a single-cell pipeline

to identify distinct cell types. Subsequently, meta-cell transcriptome data was generated for each of these

identified cell types. Using the control group samples as a reference dataset, perturbation scores were

computed for all genes within a given gene network, employing the ssNPA framework. Perturbance

scores were calculated by comparing the network predictions derived from the reference network and

the test group data, with the expectation of distinguishing between the test and control groups. Then, we

visualize all samples perturbance score with t-distributed stochastic neighbor embedding (t-SNE) and

compare them against real group ID (test or control). After verifying that the test and control group are

separated well in visualization indicating the Perturbance Scores can be used to distinguish the two

groups. The Wilcoxon tests[17] were applied to filter out genes exhibiting significant perturbations, and

finally, pathway analysis was conducted for further interpretation of the findings. The genes with a

significant adjusted p-value (< 0.05) in the Wilcoxon test between test and control perturbance score is

defined as Perturbed Genes.

We combined a meta-cell transcriptome analysis with the ssNPA framework to identify the genes that

are perturbed in the test condition (HFHS diet) compared to that of the control using a causal network[8].

The causal network is computed using a Fast Greedy Equivalent Search (FGES) algorithm[18] on a

directed acyclic graph (DAG), where the nodes are genes, and the directed edges are the causal

relationships between the genes. The FGES algorithm iteratively adds an edge if the addition decreases

the Bayesian Information Criteria (BIC), where the BIC is calculated based on a linear model where the

expression of a gene is predicted by fitting a linear model with the expression levels of its parents,

spouses, and children (Markov Blanket) in the current network as the predictors. The single-cell gene

expression data from the mouse given the standard diet is used to construct the causal network. The

difference between the predicted and actual values (i.e., perturbation score) is used to identify genes

essential for T2DM pathogenesis. Since, unlike the standard differential expression analysis, this method

considers the interactions between the genes and their neighbors, several genes identified in our analysis

could not have been identified otherwise. Importantly, the expression level of perturbed genes can be

similar between test and control groups, but the interactions between them and their neighbor genes

vary.

DEG analysis

To identify DEGs associated with diet, we applied the Wilcoxon test on scRNA-seq expression data

between control and test group for β cells using the same code in[17]. Genes with an FDR < 0.05 and

overall >were considered DEGs. Venn diagrams were generated to visualize the overlap between

perturbed genes and DEGs.

Pathway Analysis assessment: perturbed and differentially expressed genes

218 Gene set enrichment analysis of KEGG mouse pathways was conducted using the hypergeometric test.

219 This analysis assessed both perturbed genes and DEGs. The pathways were ranked based on their p-

220 values, and the top 10 pathways with the lowest p-values were selected.

Validation using the Knockout Mouse Project Database

The KOMP database, accessible at the International Mouse Phenotyping Consortium website, serves as

a valuable resource for researchers, offering comprehensive phenotypic data and genetic insights on a

wide array of knockout mouse models. This platform facilitates the understanding of gene function and

disease mechanisms and is instrumental in advancing the study of human diseases, including the

identification of potential therapeutic targets.https://www.mousephenotype.org/understand/start-using-

the-impc/impc-data-generation.

Results

Given that scRNA-seq may fail to capture all disease-relevant genes, we focused our analysis on

comparing and contrasting three gene categories: non-perturbed DEGs, perturbed non-DEGs, and

perturbed DEGs, to evaluate their respective relevance to T2DM pathophysiology across its

pathophysiological continuum to demonstrate that all three are essential to capture the full molecular

spectrum of the disease. We selected the NZO/HlLtJ mouse strain as the mild and severe diabetic model

as it displays signs of metabolic syndrome symptoms including morbid obesity, fasting hyperglycemia,

hyperinsulinemia, insulin resistance, and hypercholesterolemia, and is a well-established mildly diabetic

model[19-21].

Differentially expressed and perturbed β cells genes between prediabetic and healthy states

To identify prediabetic-state-specific perturbed genes, we performed ssNPA analysis using scRNA-seq

of β cells isolated from healthy (C57BL/6J, standard diet) and prediabetic (C57BL/6J, HFHS diet) mice.

In total, β cells exhibited 573 DEGs and 1,389 perturbed genes (**Figure 2A-B**). Of these, 289 genes were

shared between DEGs and perturbed signatures (including 235 up-regulated and 54 down-regulated

perturbed DEGs). We also identified 1,100 perturbed-only genes (perturbed but not DEGs) and 2,267

genes that were neither DEGs nor perturbed.

Among the non-perturbed DEGs, we identified several genes related to T2DM pathophysiology

including *Igf2bp2, Irs2, Hk2, Ldha, Apoe, Klf2, Il1r2, Oas1, Oas2, Oas3, Rbp4, Slc2a3, Dusp4, Adm,*

and Inhba. Irs2 and Igf2bp2 are well-established T2DM susceptibility genes that regulate insulin and

growth factor signaling *Irs2* acts as a major mediator of insulin and IGF signaling, ensuring proper β cell

survival and glucose responsiveness[22,23], whereas *Igf2bp2* stabilizes key metabolic transcripts that

influence glucose homeostasis[9]. *Hk2* and *Ldha* are important glycolytic enzymes that help maintain

cellular energy balance in β cells by regulating glucose phosphorylation and lactate production[24]. *Slc2a3*

encodes a glucose transporter that enhances glucose uptake during periods of metabolic stress[25], while

Rbp4 is an adipokine associated with insulin resistance and altered lipid metabolism in obesity and

diabetes[26].

Other genes in this group play significant roles in inflammation, stress response, and cellular

remodeling. *Il1r2* acts as a decoy receptor that modulates interleukin-1–mediated inflammation in

pancreatic β cells[27]. *Oas1*, *Oas2*, and *Oas3* are interferon-stimulated genes that may reflect immune

activation contributing to β cell stress[28]. *Dusp4*[29] *and Klf2*[30] regulate transcriptional, and kinase

pathways linked to oxidative and endoplasmic reticulum stress responses. *Apoe*[31] *and Inhba*[32] influence

lipid handling and transforming growth factor signaling, both of which contribute to β cell

dedifferentiation and senescence. *Adm* encodes adrenomedullin, a vasoprotective and cytoprotective

peptide that supports islet perfusion and survival under metabolic load[33]. Together, these DEGs

represent interconnected metabolic, inflammatory, and stress-adaptive mechanisms underlying the

transition from β cell compensation to dysfunction in T2DM.

Among the perturbed non DEGs, we identified several genes with strong relevance to β cell function and

T2DM pathophysiology, including *Glp1r, Kcnj11, Abcc8, Slc2a2, Foxo1, Txnip, Wfs1, Ptpn1* and *Dpp4.*

Glp1r encodes the glucagon-like peptide 1 receptor, which mediates incretin signaling to enhance

glucose-stimulated insulin secretion and promote β cell proliferation[10]. *Igf1r* regulates growth and

survival signaling in β cells[34]. *Kcnj11* and *Abcc8* encode the subunits of the ATP sensitive potassium

channel complex that is essential for glucose stimulated insulin secretion. Both have been associated

with T2DM[35,36]. *Slc2a2* encodes GLUT2, the major glucose transporter that allows β cells to sense
circulating glucose levels[37]. *Foxo1* integrates insulin and oxidative stress signaling and is key for
maintaining β cell identity and proliferation[37]. *Txnip* connects oxidative stress to glucose toxicity by
inhibiting thioredoxin[38], while *Wfs1* protects the endoplasmic reticulum from stress and supports insulin
folding and secretion[39]. *Ptpn1* encodes protein tyrosine phosphatase 1B, which downregulates insulin
receptor signaling[40], and *Dpp4* modulates incretin degradation, impacting insulin release[41].

Additional perturbed non-DEGs such as *Pcsk1, Prlr, Rictor, Gcgr, Ptger* and *Nfe2l.,* also play important
roles in β cell metabolism and signaling. *Pcsk1* is involved in proinsulin processing[42], while *Prlr*
(prolactin receptor) supports β cell growth and functional adaptation during metabolic stress[43]. *Rictor*, a
component of the mTORC2 complex, influences insulin signaling and cytoskeletal organization[44].
Gcgr[45] and *Ptger3*[46] mediate glucagon and prostaglandin signaling, both of which regulate calcium and
cyclic AMP pathways that control insulin secretion. *Nfe2l2* encodes NRF2, a transcription factor that
drives antioxidant defense and maintains redox homeostasis[47]. Together, these perturbed non-DEGs
indicate early disruption of metabolic, oxidative, and signaling networks that compromise β cell
adaptability before major transcriptional changes occur in diabetes progression.

Among the perturbed DEGs, we identified several genes closely linked to β cell signaling, metabolism,
and stress adaptation, including *Pdx1, Ffar1, Gipr, Pik3r1, Pdk1, Slc2a1, Gcg, Ucn3, G6pc2, and*
Gdf15. Pdx1 is a master transcription factor for pancreatic β cell identity and insulin gene expression[11].
Ffar1[48] and *Gipr* encode receptors that sense free fatty acids and incretins, respectively, enhancing
glucose stimulated insulin secretion[49]. *Pik3r1*[50] and *Pdk1*[51] are core components of insulin signaling that
regulate β cell survival and glucose metabolism. *Slc2a1* facilitates basal glucose uptake and
compensatory glucose transport under stress[52]. *Gcg* encodes glucagon[53], and *Ucn3* is a β cell secretory
product that fine tunes paracrine signaling with α cells and is associated with T2DM development[54].
G6pc2 participates in glucose cycling within β cells, influencing fasting glucose regulation[55], while
Gdf15 acts as a stress responsive cytokine linked to metabolic adaptation and protection against
glucolipotoxicity[56].

Differentially expressed and perturbed β cells genes between severely and prediabetic states

To identify diabetes-state-specific perturbed genes, we performed ssNPA analysis using scRNA-seq of β

cells isolated from severely diabetic mice (NZO, HFHS diet) and prediabetic mice (C57BL/6J, standard

diet). In total, β cells exhibited 1,289 DEGs (344 up-regulated and 945 down-regulated) and 413

perturbed genes (**Figure 3A-B**). Among these, 281 genes were shared between DEG and perturbed

signatures (170 up-regulated perturbed DEGs and 111 down-regulated perturbed DEGs). We also

identified 132 perturbed-only genes and 1,205 genes that were neither DEGs nor perturbed.

Among these non-perturbed DEGs, we identified similar and additional several key regulators of β cell

metabolism, insulin signaling, and nutrient sensing, including *Pdx1, Slc2a2, Irs2, Lepr, Fgf21, Ffar1,*

Pdk1, Pdk4, Pck2, and *Hk2. Lepr*[57] and *Fgf21*[58] reflect endocrine crosstalk between adipose tissue, liver,

and pancreatic islets, coordinating systemic energy balance and glucose homeostasis, while *Pck2*

participates in gluconeogenic and cataplerotic flux, ensuring metabolic energy flexibility[59]. Additional

genes such *as Gipr, Sst, Insig1, Cd36, Apoe, Rbp4, Ceacam1, Hhex, Maf,* and *Slc2a1* contribute to the

integrated regulation of insulin secretion, lipid metabolism, and stress response. *Sst* encodes

somatostatin, a paracrine inhibitor that modulates islet hormone release[60]. *Insig1* regulates lipid

biosynthesis and endoplasmic reticulum homeostasis[61]. *Cd36*[62], *Apoe*[31], and *Rbp4*[26] are involved in lipid

and retinol transport, influencing insulin sensitivity and β cell lipid metabolism. *Ceacam1* promotes

insulin clearance and maintains insulin receptor recycling[63], while *Hhex*[64] and *Maf*[65] are transcription

factors required for islet cell differentiation and function. Together, these genes form a tightly

coordinated network that integrates glucose and lipid metabolism, insulin signaling, and intercellular

communication to maintain β cell function and systemic glucose homeostasis.

Among the perturbed non-DEGs, several are critical for maintaining β cell identity, stress tolerance, and

insulin signaling, including *Nkx6.1, Glp1r, Wfs1, Igf1r, Nfe2l2, Cebpb, Atf6, Relb, Bace2*, *Ppp1r15a* and

Cstb. *Nkx6.1* is a core transcription factor required for β cell differentiation and insulin gene

transcription, ensuring mature β cell function[66,67]. *Cebpb* and *Atf6* contribute to stress-adaptive

transcriptional programs, particularly during ER stress and inflammatory signaling[68,69]. *Relb* functions in

the noncanonical NF-κB pathway, mediating inflammatory tolerance[70], while *Bace2* modulates

proinsulin processing and insulin granule maturation[71]. *Ppp1r15a*, also known as GPR43 is a key

regulator of the integrated stress response that promotes recovery from ER stress through

dephosphorylation of eIF2α[72]. *Cstb,* a potential exerkine[73], regulates lysosomal protease activity,

protecting cells from proteotoxic stress and may contribute to T2DM[74]. Collectively, these genes

represent a cohesive stress-adaptive and survival network that safeguards β cell function in the face of

metabolic overload and inflammatory stress characteristic of diabetes progression.

Among the perturbed DEGs, several are central regulators of β cell differentiation, glucose sensing, and

insulin secretion, including *Mafa, Neurod1, Pcsk1, Slc30a8, Ptpn1, Ucp2, Ddit3, Socs3, Abca1*, and

C2cd4b. *Mafa* and *Neurod1* are key transcription factors essential for maintaining mature β cell identity

and activating insulin gene expression. In particular, *MAFA*-associated missense mutations have been

implicated in familial insulinomatosis and diabetes. *Slc30a8* encodes the zinc transporter ZnT8, crucial

for insulin crystallization and storage within secretory granules and its mutations are linked with

T2DM[75]. *Ptpn1* negatively regulates insulin receptor signaling, and its overactivity contributes to insulin

resistance and β cell stress[76]. *Ucp2* modulates mitochondrial coupling and reactive oxygen species

generation, fine-tuning insulin secretion efficiency[77]. *Ddit3* (CHOP) is a marker of ER stress and a key

mediator of β cell apoptosis during chronic metabolic overload[78]. *Socs3* suppresses cytokine and insulin

signaling under inflammatory conditions[79], while *Abca1* regulates cholesterol efflux, preserving

membrane integrity and insulin granule function[80]. *C2cd4b* has been genetically linked to T2DM

susceptibility, possibly influencing calcium-mediated signaling in β cells[81].

Differentially expressed and perturbed β cells genes between severely and mildly diabetic sages

To identify diabetes-state–specific perturbed genes, we performed ssNPA analysis using scRNA-seq of

β cells isolated from severely diabetic mice (NZO, HFHS diet) and midly diabetic mice (NZO, standard

diet). In total, β cells exhibited 1,293 DEGs (1,371 down-regulated and 162 up-regulated) and 966

perturbed genes (**Figure 4A-B**). Among these, 637 genes were shared between DEG and perturbed

signatures (566 down-regulated perturbed DEGs and 71 up-regulated perturbed DEGs). We also

identified 329 perturbed-only genes (perturbed but not differentially expressed) and 1,946 genes that

were neither perturbed nor differentially expressed.

Among these genes, several are central to insulin signaling, glucose metabolism, and energy regulation,

including *Pparg*, *Ppargc1a*, *Irs2*, *Slc2a2*, *Lepr*, *Dpp4*, *Rbp4*, *Camk1d*, *Pdk1*, and *Gcg*. *Pparg* and

Ppargc1a coordinate lipid and glucose metabolism, promoting insulin sensitivity and mitochondrial

biogenesis in peripheral tissues[82]. *Camk1d* regulates calcium-dependent signaling pathways tied to

insulin secretion and is associated with increased risk for T2DM[83]. Additional genes such as *Hk1*, *Ldha*,

Ffar4, *Ffar2*, *C2cd4a*, *Sec16b*, *Ksr2*, *Cdk4*, *Adm*, and *Abcc9* contribute to the metabolic flexibility and

energy homeostasis of β cells. *Hk1* initiates glycolysis by phosphorylating glucose and its disruption is

implicated in T2DM[84]. *C2cd4a* has been genetically linked to T2DM and likely regulates calcium-

mediated exocytosis in β cells[85]. *Sec16b* is involved in endoplasmic reticulum export and secretory

pathway maintenance and has been recently identified as a key regulator of glucose homeostasis[86]. *Ksr2*

acts as a scaffold protein regulating AMPK and ERK signaling, linking cellular energy status with

metabolic output and is involved in T2DM pathogenesis[87]. *Cdk4* promotes β cell proliferation and

adaptive expansion during insulin resistance[88]. *Abcc9* encodes the SUR2 subunit of ATP-sensitive

potassium channels, influencing membrane excitability and insulin release, and along with *Abcc8,*

contribute to T2DM[89]. Collectively, these genes form a network that integrates glucose and lipid

metabolism, hormonal signaling, and adaptive stress responses that are fundamental to β cell

compensation in diabetes.

Among the perturbed non-DEGs, several are key regulators of pancreatic lineage specification,

mitochondrial metabolism, and insulin signaling, including *Mt.nd1*, *Glp1r*, *Nkx2.2*, *Foxa2*, *Ndufaf2*,

Atp1b1 and *Fosl2* . *Mt.nd1* encodes a mitochondrial NADH dehydrogenase subunit essential for

oxidative phosphorylation and ATP generation, supporting the high energy demands of insulin

secretion[90]. *Nkx2.2* and *Foxa2* are core transcription factors governing pancreatic development, β cell

differentiation, and maintenance of glucose-sensing machinery[91,92]. *Ndufaf2* participates in

mitochondrial complex I assembly, ensuring efficient electron transport and cellular respiration[93].

Atp1b1 encodes a subunit of Na□/K□-ATPase, which maintains membrane potential and ion gradients

critical for insulin vesicle exocytosis[94]. *Fosl2* is part of the AP-1 transcriptional complex that modulates

cellular growth and metabolic gene expression[95].

Among the perturbed DEGs, several play critical roles in mitochondrial metabolism, β cell

differentiation, and glucose utilization, including *Mt.nd2*, *Nkx6.1*, *H2.t22*, *Mt.atp8* and *Pappa2*. *Mt.nd2*

and *Mt.atp8* encode mitochondrial components of the electron transport chain, essential for ATP

production required for insulin exocytosis[96,97]. *H2.t2 is* an MHC-related genes, potentially linking

immune modulation with metabolic stress responses in β cells[98]. *Pappa2* modulates local IGF signaling

by cleaving IGF-binding proteins, enhancing insulin sensitivity and growth factor bioavailability[99].

Comparative overlap of differentially expressed and perturbed genes across the diabetes
continuum

To further illustrate that DEGs and perturbed genes capture both unique and shared molecular signatures

across the T2DM continuum, we compared the number of overlapping genes across all three conditions

and between each condition (**Figure 5A–F,** see **Supplementary Tables 1–4** for the full list of genes).

We identified 63 non-perturbed DEGs shared across all three comparisons (**Figure 5B**), including *Irs2*.

Irs2 plays a central role in insulin signaling, transmitting signals from the insulin and IGF-1 receptors to

downstream pathways like PI3K-Akt that regulate glucose uptake, β-cell survival, and hepatic glucose

production. Impaired *Irs2* expression or phosphorylation is a well-established contributor to insulin

resistance and β-cell failure, two key hallmarks of T2DM. Mouse models lacking *Irs2* develop severe

diabetes due to defective β-cell compensation, confirming its causal role in maintaining glucose

homeostasis[100].

We identified 12 perturbed-only genes shared across all three comparisons (**Figure 5D**), including

Glp1r, a clinically relevant gene encoding the glucagon-like peptide-1 receptor. Glp1r is a key

therapeutic target in T2DM, as it mediates insulin secretion and is the basis for GLP-1 receptor agonist

drugs used to improve glycemic control and promote weight loss[10]. Interestingly, despite its strong

association with T2DM, Glp1r was not detected as a DEG in the scRNA-seq data, suggesting that

network-level perturbation analysis can uncover functionally important regulatory changes that are not

apparent from expression-level differences alone.

We identified 16 perturbed DEGs shared across all three comparisons (**Figure 5F**). These included

Serping1, which encodes C1 esterase inhibitor, a key regulator of the complement and coagulation

cascades, which have been implicated in metabolic inflammation, endothelial dysfunction, and insulin

resistance in T2DM. Consistent with prior microarray analyses[101], *Serping1* was identified among the

core genes enriched in pathways related to pancreatic secretion and immune activation, suggesting that

its dysregulation may contribute to β-cell stress and impaired insulin signaling.

Together, these findings demonstrate that integrating DEG and perturbation analyses provides a more

comprehensive view of T2DM pathophysiology. While DEGs highlight transcriptional shifts directly

measurable at the expression level, perturbation-based analysis captures regulatory and signaling

alterations that may occur without major expression changes. The overlap and divergence between these

layers reveal distinct yet complementary molecular mechanisms across the T2DM continuum.

Validation of perturbed genes using the KOMP database

To corroborate the perturbed genes identified with the meta-ssNPA analysis, we employed the KOMP

database and generated visual representations for the genes previously linked to T2DM. The *in vivo*

mouse reference data for validation of genes was generated by the Knock-out Mouse Project-KOMP

(www.mousephenotype.org)[102,103]. The perturbed genes selected for validation were based on their

response to a glucose tolerance test, the standard method to diagnose diabetic or obese phenotypes in

both preclinical and clinical research. We selectively chose three genes that demonstrated improved

glucose tolerance/intolerance based on their area under curve profiles (AUC) and have been associated

with T2DM: *Glp1r, Abcc8, and Kcnj11*.

Across all three knockout models, glucose tolerance tests revealed significant impairments in glucose

clearance compared to wild-type controls, consistent with diabetic phenotypes (**Figure 6A-C**). These

findings validate the computationally predicted perturbations, as all three genes are established

regulators of pancreatic β-cell excitability and insulin secretion. Loss of *Glp1r* impairs incretin-mediated

insulin release[10], while *Abcc8* and *Kcnj11* encode subunits of the ATP-sensitive potassium (K_ATP)

channel essential for insulin granule exocytosis[104-106].

Discussion

Several studies have utilized gene expression data to construct causal networks[107-109]. Additionally,

researchers have identified gene features that exhibit strong predictive power for specific phenotypes[110-

112]. In this context, the innovative approach of ssNPA assesses how the gene network of a set of control

samples is perturbed when presented with a new query sample. The underlying rationale of ssNPA is

based on the notion that, in many diseases, an observed phenotype may arise from alterations in different components of the 'healthy' gene network. The ssNPA framework offers several distinct advantages over traditional approaches. Firstly, it enables the inference of topological relationships among genes within the causal network providing valuable insights into the interconnections and regulatory dynamics among genes. Secondly, the data-driven nature of the network allows for the discovery of novel information enhancing our understanding of the complex mechanisms underlying the disease. Thirdly, ssNPA does not rely on prior pathway knowledge, making it a valuable tool for investigating gene perturbations in a hypothesis-free manner. In comparison, prior studies on T2DM primarily relied on DEG analysis and genome-wide association studies (GWAS), which lack the aforementioned advantages of the ssNPA framework.

T2DM is a chronic metabolic disorder characterized by heterogeneity and polygenic traits. The genetic bases of T2DM are poorly understood, highlighting the need of approaches that can help investigate the genes and associated signaling pathways contributing to its onset. To this end, we constructed a transcriptional network that explored perturbed genes, hub genes, and associated pathways and performed validation using the KOMP database. The ssNPA enabled the identification of several genes already established in the diabetes context while also unveiling novel genes previously unrecognized in relation to diabetes. In our comparison between C57BL/6J mice on a regular chow diet and those on a HFHS diet, the analysis highlighted a significant number of perturbed genes. This observation is particularly noteworthy as there is a limited report of genes during the *prediabetic* state, emphasizing the importance of these findings in understanding new-onset diabetes/prediabetes. The key findings in this comparison included genes, such as *Glp1r* and *Abcc8*, which we subsequently validated using the KOMP database.

SsNPA is a sophisticated method designed to assess perturbations in gene networks at the level of individual samples. Instead of focusing on isolated genes, ssNPA delves into the broader landscape of interconnected gene networks. The methodology begins by inferring a global gene network, utilizing causal graph learning derived from a set of reference samples. Upon the introduction of a new sample, ssNPA calculates the degree of deviation of this sample from the established reference network at every gene point. This approach furnishes in-depth information regarding the topology of network perturbations. By generating a perturbation feature vector, i.e. perturbance score, ssNPA allows for the

classification or clustering of samples, which can be instrumental in distinguishing between cell types,

disease subtypes, or any other biological distinctions under investigation. This tool provides an

advanced perspective on how various perturbations, such as environmental changes, drug treatments, or

genetic mutations, affect the broader dynamics of gene networks. SsNPA and differential gene

expression analysis, while both employed in the domain of genomics, serve distinctly different

analytical purposes. Differential gene expression analysis aims to identify individual genes that exhibit

statistically significant differences in expression between two or more conditions, such as healthy versus

diseased states. Its output is often a list of upregulated or downregulated genes. In contrast, ssNPA

focuses on assessing gene network perturbations in individual samples. Instead of concentrating on the

behavior of single genes, ssNPA evaluates how perturbations, such as mutations or drug treatments,

influence entire gene networks or pathways. It provides insights into deviations from a reference

network, shedding light on both the magnitude and topology of network perturbations. While differential

gene expression gives a granular view of specific genes' behavior, ssNPA offers a holistic perspective on

how interconnected gene networks respond to various conditions.

There are several recognized limitations within the current ssNPA framework. Primarily, the framework

exhibits suboptimal performance when applied to single-cell data. Such data often present sparsity

challenges, characterized by numerous genes that register zero expressions across a multitude of cells.

This phenomenon disrupts the linear model's foundational assumption upon which ssNPA operates,

resulting in diminished prediction accuracy. A proposed solution to this challenge, introduced by Baran

and Bercovich[16], involves the innovative 'meta-cell' concept. By randomly selecting multiple cells of

identical cell types and computing the mean gene expressions, a novel meta-cell sample is generated.

This methodology retains the relative gene expression hierarchy within the cells, ensuring genes with

higher expressions remain dominant, and those with lower expressions continue to be subdued in the

meta-cells, while concurrently mitigating the zero-expression issue. A secondary concern pertains to the

criteria ssNPA employs to discern Perturbed genes. In its original design, a gene is classified as

"Perturbed" provided its average Perturbance Score exceeds that of the control group. This method,

however, overlooks the potential influence of random noise on the Perturbance Score, which can

inadvertently result in False Positive outcomes. To enhance precision, we integrated a statistical

approach, employing the Wilcoxon test as elucidated by Li et al.[17]. By contrasting the Perturbance

Scores of both test and control groups and setting a FDR threshold of less than 0.05, we established a

more rigorous criterion for identifying perturbed genes. For this analysis, a one-sided Wilcoxon test was

executed, operating under the alternative hypothesis that the control group's Perturbance Score is inferior

to that of the test group. The KOMP validation process is designed to be broad and encompasses

multiple cell types, rather than being specific to any single cell type. This approach allows for a more

generalizable understanding of gene function across different biological contexts.

Conclusion

In conclusion, our study employed meta-ssNPA, an innovative combination of meta-cell transcriptome

analysis with the ssNPA framework, to investigate gene perturbations in various conditions. We

identified genes that are perturbed in the context of T2DM and metabolic disorders, shedding light on

potential therapeutic targets. The analysis revealed DEGs and perturbed genes in C57BL/6J mice fed on

a HFHS diet compared to those on a standard diet. Pathway enrichment analysis highlighted the

involvement of these genes in critical metabolic pathways. Moreover, we identified known T2DM genes

such as *Glp1r*, *Igf1r*, and *Zbtb20*. which have previously been linked to metabolic traits and insulin

regulation. Novel genes such as *Bpifc*, *Itga11*, *P2rx2*, *Tnip1* and *Khl32* are also discovered by the

analysis and validated through KOMP database. To validate our findings, we leveraged the KOMP

databasex knockout and characterize all protein-coding genes in the mouse genome, which corroborated

the identified perturbed genes obtained from the meta-ssNPA analysis. Additionally, we discovered

novel genes that exhibited perturbations despite not being DEGs and not directly associated with T2DM.

Figure Legend

Figure 1. Overview of the meta-ssNPA framework and experimental design. A) Experimental setup

showing comparisons across the T2DM continuum. C57BL/6J mice on a standard diet were compared to

those on a high-fat, high-sucrose (HFHS) diet to model healthy versus prediabetic states, while

NZO/HlLtJ mice on standard and HFHS diets represented severe versus prediabetic and severe versus

mild diabetic conditions, respectively. **B)** Schematic representation of the single-cell network

perturbation analysis (ssNPA) pipeline. Single-cell RNA sequencing data from islets of control and

HFHS-fed mice were processed to construct cell-type-specific networks, compute perturbation scores,

and identify gene-specific causal relationships using the FGES algorithm. **C)** The meta-ssNPA

framework integrates reference and disease single-cell transcriptomes into a unified meta-cell network,

quantifies gene perturbation scores, and performs differential analyses to identify perturbed genes and

their causal subnetworks. Selected genes were further validated using the Knockout Mouse Phenotyping

Program (KOMP) database.

Figure 2. Differentially expressed and perturbed genes in β-cells from healthy and prediabetic

mice. A) Volcano plot showing the relationship between gene expression changes (log☐ fold change)

and statistical significance (−log☐☐FDR) in β-cells from C57BL/6J mice fed a standard versus a high-

fat, high-sucrose (HFHS) diet. Triangles denote perturbed genes, while circles represent non-perturbed

genes; red and green indicate up- and downregulated DEGs, respectively. Several key metabolic

regulators such as *Pdx1*, *Slc2a2*, *Foxo1*, and *Gcg* exhibit significant expression or perturbation

differences between groups. **B)** Venn diagram depicting the overlap between DEGs (left) and perturbed-

only genes (right). Of the total identified genes, 289 were shared between both categories, while 284 and

1,100 were unique to DEGs and perturbed-only sets, respectively, indicating that perturbation analysis

reveals additional functionally altered genes beyond those identified by expression changes alone.

Figure 3. Differentially expressed and perturbed genes in β-cells from prediabetic and severely

diabetic mice. A) Volcano plot illustrating the relationship between gene expression changes (log☐ fold

change) and statistical significance (−log☐☐FDR) in β-cells from NZO/HILtJ mice fed a high-fat, high-

sucrose (HFHS) diet compared to those on a standard diet. Triangles represent perturbed genes, while

circles denote non-perturbed genes; red and green indicate up- and downregulated DEGs, respectively.

Several critical β-cell and insulin signaling regulators, including *Glp1r*, *Slc2a2*, *Irs2*, *Wfs1*, and

Neurod1, showed significant expression or perturbation differences between groups. **B)** Venn diagram

showing the overlap between DEGs (left) and perturbed-only genes (right). Of the total identified genes,

281 were shared between both categories, while 1,008 and 132 were unique to DEGs and perturbed-only

sets, respectively. This highlights that perturbation analysis uncovers additional network-level

alterations beyond transcriptional changes detectable through differential expression alone.

Figure 4. Differentially expressed and perturbed genes in β-cells from mild and severe diabetic

mice. A) Volcano plot showing the relationship between gene expression changes (log☐ fold change)

and statistical significance (−log☐☐FDR) in β-cells from NZO/HILtJ mice with mild versus severe

diabetes. Triangles denote perturbed genes, while circles represent non-perturbed genes; red and green

indicate up- and downregulated DEGs, respectively. Several metabolic and β-cell transcriptional

regulators, including *Pparg*, *Foxa2*, *Slc2a2*, *Lepr*, and *Gcg*, exhibited marked differences in expression

or perturbation between disease stages. **B)** Venn diagram showing the overlap between DEGs (left) and

perturbed-only genes (right). A total of 637 genes were shared between both categories, while 656 and

329 were unique to DEGs and perturbed-only sets, respectively. These findings suggest that as diabetes

severity increases, both transcriptional dysregulation and network-level perturbations become more

extensive, reflecting progressive β-cell dysfunction.

Figure 5. Overlap of DEGs and perturbed genes across stages of the T2DM continuum. Venn

diagrams show the overlap in differentially expressed and perturbed genes identified across three

comparisons: prediabetes versus healthy (red), severe versus prediabetes (green), and severe versus mild

diabetes (blue). **A)** Overlap of all DEGs across conditions, revealing both shared and stage-specific

transcriptional changes. **B)** Overlap of DEGs only, indicating a limited set of consistently altered genes

across all comparisons. **C)** Overlap of all perturbed genes, showing broader network-level disruptions

shared between disease stages. **D)** Overlap of perturbed-only genes, highlighting 12 genes shared across

all three conditions that were not captured by differential expression alone. **E)** Overlap between DEGs

and perturbed genes, demonstrating partial convergence between expression-level and network-level

changes. **F)** Overlap of DEGs or perturbed genes, showing the combined molecular landscape across the

T2DM continuum. Collectively, these results illustrate that DEG and perturbation analyses capture both

distinct and overlapping molecular features underlying progressive T2DM pathophysiology.

Figure 6. In vivo validation of perturbed genes using the Knockout Mouse Phenotyping (KOMP)

database. Glucose tolerance test (GTT) data from knockout (KO) mice were analyzed to validate genes

identified as perturbed in β-cells. **(A–C)** Time-course plots show blood glucose levels following glucose

administration in *Glp1r*, *Abcc8*, and *Kcnj11* KO mice (orange) compared with wild-type controls (blue).

KO mice exhibited significantly impaired glucose clearance, reflected by elevated glucose levels over

time. **(A–C, right)** Box plots of the area under the glucose response curve (AUC) further confirm

reduced glucose tolerance in KO mice relative to controls, with highly significant Wilcoxon p-values.

These findings corroborate the role of *Glp1r*, *Abcc8*, and *Kcnj11* as key regulators of β-cell function and

glucose homeostasis.

Acknowledgments

The authors thank the Knock-out Mouse Project (KOMP) and members of Li Lab for discussions. The authors thank The Jackson Laboratory Computational Sciences and Research IT teams for technical support. Research reported in this publication was partially supported by The Jackson Laboratory Cube Initiatives. S. Li was supported by startup funds from The Jackson Laboratory and by the National Cancer Institute of the National Institutes of Health under Award Number P30CA034196. S. Li is also supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM133562, by the National Human Genomic Research Institute of the National Institutes of Health under Award Number U01HG013175, by the National Cancer Institute of the National Institutes of Health under Award Number U01CA271830 and U01CA271830-03S1, by the NIH Common Fund under Award Number U54AG079753, and by the National Institute of Aging of the National Institutes of Health under Award Number R56AG071766.

Zheng, Y., Ley, S. H. & Hu, F. B. Global aetiology and epidemiology of type 2 diabetes mellitus and its complications. *Nat Rev Endocrinol* **14**, 88-98 (2018). https://doi.org:10.1038/nrendo.2017.151

Sun, H. *et al.* IDF Diabetes Atlas: Global, regional and country-level diabetes prevalence estimates for 2021 and projections for 2045. *Diabetes Res Clin Pract* **183**, 109119 (2022). https://doi.org:10.1016/j.diabres.2021.109119

Da Silva Xavier, G. The Cells of the Islets of Langerhans. *J Clin Med* **7** (2018). https://doi.org:10.3390/jcm7030054

Li, S. *et al.* Advancing biological understanding of cellular senescence with computational multiomics. *Nat Genet* **57**, 2381-2394 (2025). https://doi.org:10.1038/s41588-025-02314-y

Hrovatin, K. *et al.* Delineating mouse beta-cell identity during lifetime and in diabetes with a single cell atlas. *Nat Metab* **5**, 1615-1637 (2023). https://doi.org:10.1038/s42255-023-00876-x

Camunas-Soler, J. *et al.* Patch-Seq Links Single-Cell Transcriptomes to Human Islet Dysfunction in Diabetes. *Cell Metab* **31**, 1017-1031 e1014 (2020). https://doi.org:10.1016/j.cmet.2020.04.005

Bakhti, M., Bottcher, A. & Lickert, H. Modelling the endocrine pancreas in health and disease. *Nat Rev Endocrinol* **15**, 155-171 (2019). https://doi.org:10.1038/s41574-018-0132-z

Buschur, K. L., Chikina, M. & Benos, P. V. Causal network perturbations for instance-specific analysis of single cell and disease samples. *Bioinformatics* **36**, 2515-2521 (2020). https://doi.org:10.1093/bioinformatics/btz949

Wang, J., Chen, L. & Qiang, P. The role of IGF2BP2, an m6A reader gene, in human metabolic diseases and cancers. *Cancer Cell Int* **21**, 99 (2021). https://doi.org:10.1186/s12935-021-01799-x

Zheng, Z. *et al.* Glucagon-like peptide-1 receptor: mechanisms and advances in therapy. *Signal Transduct Target Ther* **9**, 234 (2024). https://doi.org:10.1038/s41392-024-01931-z

Fujimoto, K. & Polonsky, K. S. Pdx1 and other factors that regulate pancreatic beta-cell survival. *Diabetes Obes Metab* **11 Suppl 4**, 30-37 (2009). https://doi.org:10.1111/j.1463-1326.2009.01121.x

12 Kang, H. M. *et al.* Multiplexed droplet single-cell RNA-sequencing using natural genetic
variation. *Nat Biotechnol* **36**, 89-94 (2018). https://doi.org:10.1038/nbt.4042
13 Keane, T. M. *et al.* Mouse genomic variation and its effect on phenotypes and gene regulation.
Nature **477**, 289-294 (2011). https://doi.org:10.1038/nature10413
14 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014). https://doi.org:10.1186/s13059-014-
0550-8
15 Hao, J., Cao, W., Huang, J., Zou, X. & Han, Z. G. Optimal Gene Filtering for Single-Cell data
(OGFSC)-a gene filtering algorithm for single-cell RNA-seq data. *Bioinformatics* **35**, 2602-2609
(2019). https://doi.org:10.1093/bioinformatics/bty1016
16 Baran, Y. *et al.* MetaCell: analysis of single-cell RNA-seq data using K-nn graph partitions.
Genome Biol **20**, 206 (2019). https://doi.org:10.1186/s13059-019-1812-2
17 Li, Y., Ge, X., Peng, F., Li, W. & Li, J. J. Exaggerated false positives by popular differential
expression methods when analyzing human population samples. *Genome Biol* **23**, 79 (2022).
https://doi.org:10.1186/s13059-022-02648-4
18 Ramsey, J., Glymour, M., Sanchez-Romero, R. & Glymour, C. A million variables and more: the
Fast Greedy Equivalence Search algorithm for learning high-dimensional graphical causal
models, with an application to functional magnetic resonance images. *Int J Data Sci Anal* **3**, 121-
129 (2017). https://doi.org:10.1007/s41060-016-0032-z
19 Leiter, E. H. *et al.* NIDDM genes in mice: deleterious synergism by both parental genomes
contributes to diabetogenic thresholds. *Diabetes* **47**, 1287-1295 (1998).
https://doi.org:10.2337/diab.47.8.1287
20 Ortlepp, J. R. *et al.* A metabolic syndrome of hypertension, hyperinsulinaemia and
hypercholesterolaemia in the New Zealand obese mouse. *Eur J Clin Invest* **30**, 195-202 (2000).
https://doi.org:10.1046/j.1365-2362.2000.00611.x
21 Reifsnyder, P. C., Churchill, G. & Leiter, E. H. Maternal environment and genotype interact to
establish diabesity in mice. *Genome Res* **10**, 1568-1578 (2000).
https://doi.org:10.1101/gr.147000
22 Withers, D. J. *et al.* Disruption of IRS-2 causes type 2 diabetes in mice. *Nature* **391**, 900-904
(1998). https://doi.org:10.1038/36116
23 Brady, M. J. IRS2 takes center stage in the development of type 2 diabetes. *J Clin Invest* **114**,
886-888 (2004). https://doi.org:10.1172/JCI23108
24 Rabbani, N., Xue, M. & Thornalley, P. J. Hexokinase-2-Linked Glycolytic Overload and
Unscheduled Glycolysis-Driver of Insulin Resistance and Development of Vascular
Complications of Diabetes. *Int J Mol Sci* **23** (2022). https://doi.org:10.3390/ijms23042165
25 Ziegler, G. C., Almos, P., McNeill, R. V., Jansch, C. & Lesch, K. P. Cellular effects and clinical
implications of SLC2A3 copy number variation. *J Cell Physiol* **235**, 9021-9036 (2020).
https://doi.org:10.1002/jcp.29753
26 Fan, J. & Hu, J. Retinol binding protein 4 and type 2 diabetes: from insulin resistance to
pancreatic beta-cell function. *Endocrine* **85**, 1020-1034 (2024). https://doi.org:10.1007/s12020-
024-03777-5
27 Boni-Schnetzler, M. *et al.* beta Cell-Specific Deletion of the IL-1 Receptor Antagonist Impairs
beta Cell Proliferation and Insulin Secretion. *Cell Rep* **22**, 1774-1786 (2018).
https://doi.org:10.1016/j.celrep.2018.01.063

28 Hovanessian, A. G. & Justesen, J. The human 2'-5'oligoadenylate synthetase family: unique
interferon-inducible enzymes catalyzing 2'-5' instead of 3'-5' phosphodiester bond formation.
Biochimie **89**, 779-788 (2007). https://doi.org:10.1016/j.biochi.2007.02.003
29 Denhez, B. *et al.* Diabetes-Induced DUSP4 Reduction Promotes Podocyte Dysfunction and
Progression of Diabetic Nephropathy. *Diabetes* **68**, 1026-1039 (2019).
https://doi.org:10.2337/db18-0837
30 Zhong, F. *et al.* Reduced Kruppel-like factor 2 expression may aggravate the endothelial injury
of diabetic nephropathy. *Kidney Int* **87**, 382-395 (2015). https://doi.org:10.1038/ki.2014.286
31 Martins, I. J. *et al.* Apolipoprotein E, cholesterol metabolism, diabetes, and the convergence of
risk factors for Alzheimer's disease and cardiovascular disease. *Mol Psychiatry* **11**, 721-736
(2006). https://doi.org:10.1038/sj.mp.4001854
32 Brown, M. L. & Schneyer, A. A Decade Later: Revisiting the TGFbeta Family's Role in
Diabetes. *Trends Endocrinol Metab* **32**, 36-47 (2021). https://doi.org:10.1016/j.tem.2020.11.006
33 Teich-Alasia, S. A study of the vascularisation of pedicle flaps using disulphine blue. *Br J Plast*
Surg **24**, 282-288 (1971). https://doi.org:10.1016/s0007-1226(71)80071-1
34 Gardner, E. J. *et al.* Damaging missense variants in IGF1R implicate a role for IGF-1 resistance
in the etiology of type 2 diabetes. *Cell Genom* **2**, None (2022).
https://doi.org:10.1016/j.xgen.2022.100208
35 Haghvirdizadeh, P. *et al.* KCNJ11: Genetic Polymorphisms and Risk of Diabetes Mellitus. *J*
Diabetes Res **2015**, 908152 (2015). https://doi.org:10.1155/2015/908152
36 Li, M., Han, X. & Ji, L. Clinical and Genetic Characteristics of ABCC8 Nonneonatal Diabetes
Mellitus: A Systematic Review. *J Diabetes Res* **2021**, 9479268 (2021).
https://doi.org:10.1155/2021/9479268
37 Sansbury, F. H. *et al.* SLC2A2 mutations can cause neonatal diabetes, suggesting GLUT2 may
have a role in human insulin secretion. *Diabetologia* **55**, 2381-2385 (2012).
https://doi.org:10.1007/s00125-012-2595-0
38 Choi, E. H. & Park, S. J. TXNIP: A key protein in the cellular stress response pathway and a
potential therapeutic target. *Exp Mol Med* **55**, 1348-1356 (2023). https://doi.org:10.1038/s12276-
023-01019-8
39 Chapla, A. *et al.* WFS1 Gene-associated Diabetes Phenotypes and Identification of a Founder
Mutation in Southern India. *J Clin Endocrinol Metab* **107**, 1328-1336 (2022).
https://doi.org:10.1210/clinem/dgac002
40 Coronell-Tovar, A. *et al.* Protein tyrosine phosphatase 1B (PTP1B) function, structure, and
inhibition strategies to develop antidiabetic drugs. *FEBS Lett* **598**, 1811-1838 (2024).
https://doi.org:10.1002/1873-3468.14901
41 Saini, K., Sharma, S. & Khan, Y. DPP-4 inhibitors for treating T2DM - hype or hope? an
analysis based on the current literature. *Front Mol Biosci* **10**, 1130625 (2023).
https://doi.org:10.3389/fmolb.2023.1130625
42 Ramos-Molina, B., Martin, M. G. & Lindberg, I. PCSK1 Variants and Human Obesity. *Prog*
Mol Biol Transl Sci **140**, 47-74 (2016). https://doi.org:10.1016/bs.pmbts.2015.12.001
43 Banerjee, R. R. *et al.* Gestational Diabetes Mellitus From Inactivation of Prolactin Receptor and
MafB in Islet beta-Cells. *Diabetes* **65**, 2331-2341 (2016). https://doi.org:10.2337/db15-1527
44 Kumar, A. *et al.* Fat cell-specific ablation of rictor in mice impairs insulin-regulated fat cell and
whole-body glucose and lipid metabolism. *Diabetes* **59**, 1397-1406 (2010).
https://doi.org:10.2337/db09-1061

45 Jia, Y., Liu, Y., Feng, L., Sun, S. & Sun, G. Role of Glucagon and Its Receptor in the
Pathogenesis of Diabetes. *Front Endocrinol (Lausanne)* **13**, 928016 (2022).
https://doi.org/10.3389/fendo.2022.928016
46 Yu, Y., Jia, Y. Y., Wang, M., Mu, L. & Li, H. J. PTGER3 and MMP-2 play potential roles in
diabetic nephropathy via competing endogenous RNA mechanisms. *BMC Nephrol* **22**, 27 (2021).
https://doi.org/10.1186/s12882-020-02194-w
47 Dodson, M. *et al.* NRF2 and Diabetes: The Good, the Bad, and the Complex. *Diabetes* **71**, 2463-
2476 (2022). https://doi.org/10.2337/db22-0623
48 Li, Z., Xu, X., Huang, W. & Qian, H. Free Fatty Acid Receptor 1 (FFAR1) as an Emerging
Therapeutic Target for Type 2 Diabetes Mellitus: Recent Progress and Prevailing Challenges.
Med Res Rev **38**, 381-425 (2018). https://doi.org/10.1002/med.21441
49 Campbell, J. E. *et al.* GIPR/GLP-1R dual agonist therapies for diabetes and weight loss-
chemistry, physiology, and clinical applications. *Cell Metab* **35**, 1519-1529 (2023).
https://doi.org/10.1016/j.cmet.2023.07.010
50 Tsay, A. & Wang, J. C. The Role of PIK3R1 in Metabolic Function and Insulin Sensitivity. *Int J*
Mol Sci **24** (2023). https://doi.org/10.3390/ijms241612665
51 Hashimoto, N. *et al.* Ablation of PDK1 in pancreatic beta cells induces diabetes as a result of
loss of beta cell mass. *Nat Genet* **38**, 589-593 (2006). https://doi.org/10.1038/ng1774
52 Stefanidis, I. *et al.* The contribution of genetic variants of SLC2A1 gene in T2DM and T2DM-
nephropathy: association study and meta-analysis. *Ren Fail* **40**, 561-576 (2018).
https://doi.org/10.1080/0886022X.2018.1496931
53 Zhao, S. *et al.* A GLP-1/glucagon (GCG)/CCK(2) receptors tri-agonist provides new therapy for
obesity and diabetes. *Br J Pharmacol* **179**, 4360-4377 (2022). https://doi.org/10.1111/bph.15860
54 Alarslan, P. *et al.* Increased urocortin 3 levels are associated with the risk of having type 2
diabetes mellitus. *J Diabetes* **12**, 474-482 (2020). https://doi.org/10.1111/1753-0407.13020
55 Bahl, V. *et al.* G6PC2 controls glucagon secretion by defining the set point for glucose in
pancreatic alpha cells. *Sci Transl Med* **17**, eadi6148 (2025).
https://doi.org/10.1126/scitranslmed.adi6148
56 Eddy, A. C. & Trask, A. J. Growth differentiation factor-15 and its role in diabetes and
cardiovascular disease. *Cytokine Growth Factor Rev* **57**, 11-18 (2021).
https://doi.org/10.1016/j.cytogfr.2020.11.002
57 Do, O. H., Gunton, J. E., Gaisano, H. Y. & Thorn, P. Changes in beta cell function occur in
prediabetes and early disease in the Lepr (db) mouse model of diabetes. *Diabetologia* **59**, 1222-
1230 (2016). https://doi.org/10.1007/s00125-016-3942-3
58 Fisher, F. M. & Maratos-Flier, E. Understanding the Physiology of FGF21. *Annu Rev Physiol* **78**,
223-241 (2016). https://doi.org/10.1146/annurev-physiol-021115-105339
59 Beale, E. G., Harvey, B. J. & Forest, C. PCK1 and PCK2 as candidate diabetes and obesity
genes. *Cell Biochem Biophys* **48**, 89-95 (2007). https://doi.org/10.1007/s12013-007-0025-6
60 Hernandez, C., Simo-Servat, O. & Simo, R. Somatostatin and diabetic retinopathy: current
concepts and new therapeutic perspectives. *Endocrine* **46**, 209-214 (2014).
https://doi.org/10.1007/s12020-014-0232-z
61 Ouyang, S., Mo, Z., Sun, S., Yin, K. & Lv, Y. Emerging role of Insig-1 in lipid metabolism and
lipid disorders. *Clin Chim Acta* **508**, 206-212 (2020). https://doi.org/10.1016/j.cca.2020.05.042
62 Puchalowicz, K. & Rac, M. E. The Multifunctionality of CD36 in Diabetes Mellitus and Its
Complications-Update in Pathogenesis, Treatment and Monitoring. *Cells* **9** (2020).
https://doi.org/10.3390/cells9081877

63 Najjar, S. M. Regulation of insulin action by CEACAM1. *Trends Endocrinol Metab* **13**, 240-245
(2002). https://doi.org:10.1016/s1043-2760(02)00608-2
64 Zhang, J., McKenna, L. B., Bogue, C. W. & Kaestner, K. H. The diabetes gene Hhex maintains
delta-cell differentiation and islet function. *Genes Dev* **28**, 829-834 (2014).
https://doi.org:10.1101/gad.235499.113
65 Butler, A. E. *et al.* Beta cell nuclear musculoaponeurotic fibrosarcoma oncogene family A
(MafA) is deficient in type 2 diabetes. *Diabetologia* **55**, 2985-2988 (2012).
https://doi.org:10.1007/s00125-012-2666-2
66 Aigha, II & Abdelalim, E. M. NKX6.1 transcription factor: a crucial regulator of pancreatic beta
cell development, identity, and proliferation. *Stem Cell Res Ther* **11**, 459 (2020).
https://doi.org:10.1186/s13287-020-01977-0
67 Taylor, B. L., Liu, F. F. & Sander, M. Nkx6.1 is essential for maintaining the functional state of
pancreatic beta cells. *Cell Rep* **4**, 1262-1275 (2013). https://doi.org:10.1016/j.celrep.2013.08.010
68 Matsuda, T. *et al.* Ablation of C/EBPbeta alleviates ER stress and pancreatic beta cell failure
through the GRP78 chaperone in mice. *J Clin Invest* **120**, 115-126 (2010).
https://doi.org:10.1172/JCI39721
69 Chu, W. S. *et al.* Activating transcription factor 6 (ATF6) sequence polymorphisms in type 2
diabetes and pre-diabetic traits. *Diabetes* **56**, 856-862 (2007). https://doi.org:10.2337/db06-1305
70 Frier, B. C., Noble, E. G. & Locke, M. Diabetes-induced atrophy is associated with a muscle-
specific alteration in NF-kappaB activation and expression. *Cell Stress Chaperones* **13**, 287-296
(2008). https://doi.org:10.1007/s12192-008-0062-0
71 Alcarraz-Vizan, G. *et al.* BACE2 suppression promotes beta-cell survival and function in a
model of type 2 diabetes induced by human islet amyloid polypeptide overexpression. *Cell Mol*
Life Sci **74**, 2827-2838 (2017). https://doi.org:10.1007/s00018-017-2505-1
72 McNelis, J. C. *et al.* GPR43 Potentiates beta-Cell Function in Obesity. *Diabetes* **64**, 3203-3217
(2015). https://doi.org:10.2337/db14-1938
73 Taha, H. B. Lactate: a potential brain-targeting exerkine. *J Physiol* (2024).
https://doi.org:10.1113/JP287745
74 Liu, C. *et al.* Cathepsin B deteriorates diabetic cardiomyopathy induced by streptozotocin via
promoting NLRP3-mediated pyroptosis. *Mol Ther Nucleic Acids* **30**, 198-207 (2022).
https://doi.org:10.1016/j.omtn.2022.09.019
75 Rutter, G. A. & Chimienti, F. SLC30A8 mutations in type 2 diabetes. *Diabetologia* **58**, 31-36
(2015). https://doi.org:10.1007/s00125-014-3405-7
76 Bowden, D. W. Association of the PTPN1 gene with type 2 diabetes and insulin resistance.
Discov Med **4**, 427-432 (2004).
77 Liu, J., Li, J., Li, W. J. & Wang, C. M. The role of uncoupling proteins in diabetes mellitus. *J*
Diabetes Res **2013**, 585897 (2013). https://doi.org:10.1155/2013/585897
78 Yong, J. *et al.* Chop/Ddit3 depletion in beta cells alleviates ER stress and corrects hepatic
steatosis in mice. *Sci Transl Med* **13** (2021). https://doi.org:10.1126/scitranslmed.aba9796
79 Jorgensen, S. B. *et al.* Deletion of skeletal muscle SOCS3 prevents insulin resistance in obesity.
Diabetes **62**, 56-64 (2013). https://doi.org:10.2337/db12-0443
80 He, Y. *et al.* Diabetes Impairs Cellular Cholesterol Efflux From ABCA1 to Small HDL Particles.
Circ Res **127**, 1198-1210 (2020). https://doi.org:10.1161/CIRCRESAHA.120.317178
81 Kycia, I. *et al.* A Common Type 2 Diabetes Risk Variant Potentiates Activity of an
Evolutionarily Conserved Islet Stretch Enhancer and Increases C2CD4A and C2CD4B
Expression. *Am J Hum Genet* **102**, 620-635 (2018). https://doi.org:10.1016/j.ajhg.2018.02.020

82 Cataldi, S., Costa, V., Ciccodicola, A. & Aprile, M. PPARgamma and Diabetes: Beyond the
Genome and Towards Personalized Medicine. *Curr Diab Rep* **21**, 18 (2021).
https://doi.org:10.1007/s11892-021-01385-5

83 Fromont, C. *et al.* Discovery of Highly Selective Inhibitors of Calmodulin-Dependent Kinases
That Restore Insulin Sensitivity in the Diet-Induced Obesity in Vivo Mouse Model. *J Med Chem*
63, 6784-6801 (2020). https://doi.org:10.1021/acs.jmedchem.9b01803

84 Sundaram, S. M., Doughty, L. A. & Sereda, M. W. Location matters: hexokinase 1 in glucose
metabolism and inflammation. *Trends Endocrinol Metab* **33**, 665-667 (2022).
https://doi.org:10.1016/j.tem.2022.07.005

85 Kuo, T. *et al.* Identification of C2CD4A as a human diabetes susceptibility gene with a role in
beta cell insulin secretion. *Proc Natl Acad Sci U S A* **116**, 20033-20042 (2019).
https://doi.org:10.1073/pnas.1904311116

86 Zhang, R. X. *et al.* Identification of SEC16B as a novel regulator of glucose homeostasis.
Diabetologia **68**, 2247-2262 (2025). https://doi.org:10.1007/s00125-025-06501-8

87 Pearce, L. R. *et al.* KSR2 mutations are associated with obesity, insulin resistance, and impaired
cellular fuel oxidation. *Cell* **155**, 765-777 (2013). https://doi.org:10.1016/j.cell.2013.09.058

88 Stamateris, R. E. *et al.* Noncanonical CDK4 signaling rescues diabetes in a mouse model by
promoting beta cell differentiation. *J Clin Invest* **133** (2023). https://doi.org:10.1172/JCI166490

89 Bryan, J. *et al.* ABCC8 and ABCC9: ABC transporters that regulate K+ channels. *Pflugers Arch*
453, 703-718 (2007). https://doi.org:10.1007/s00424-006-0116-z

90 Yu, P., Yu, D. M., Liu, D. M., Wang, K. & Tang, X. Z. Relationship between mutations of
mitochondrial DNA ND1 gene and type 2 diabetes. *Chin Med J (Engl)* **117**, 985-989 (2004).

91 Doyle, M. J. & Sussel, L. Nkx2.2 regulates beta-cell function in the mature islet. *Diabetes* **56**,
1999-2007 (2007). https://doi.org:10.2337/db06-1766

92 Lantz, K. A. *et al.* Foxa2 regulates multiple pathways of insulin secretion. *J Clin Invest* **114**, 512-
520 (2004). https://doi.org:10.1172/JCI21149

93 Abu Hanna, F. *et al.* Lack of mitochondrial complex I assembly factor NDUFAF2 results in a
distinctive infantile-onset brainstem neurodegenerative disease with early lethality. *Orphanet J*
Rare Dis **19**, 92 (2024). https://doi.org:10.1186/s13023-024-03094-0

94 Hasbani, N. R. *et al.* Type 2 Diabetes Modifies the Association of CAD Genomic Risk Variants
With Subclinical Atherosclerosis. *Circ Genom Precis Med* **16**, e004176 (2023).
https://doi.org:10.1161/CIRCGEN.123.004176

95 He, X. *et al.* FOSL2 activates TGF-beta1-mediated GLUT1/mTOR signaling to promote diabetic
kidney disease. *J Diabetes Investig* **16**, 187-203 (2025). https://doi.org:10.1111/jdi.14360

96 Chen, J., Gusdon, A. M., Piganelli, J., Leiter, E. H. & Mathews, C. E. mt-Nd2(a) Modifies
resistance against autoimmune type 1 diabetes in NOD mice at the level of the pancreatic beta-
cell. *Diabetes* **60**, 355-359 (2011). https://doi.org:10.2337/db10-1241

97 Kytovuori, L. *et al.* A novel mutation m.8561C>G in MT-ATP6/8 causing a mitochondrial
syndrome with ataxia, peripheral neuropathy, diabetes mellitus, and hypergonadotropic
hypogonadism. *J Neurol* **263**, 2188-2195 (2016). https://doi.org:10.1007/s00415-016-8249-2

98 Keller, M. P. & Attie, A. D. Physiological insights gained from gene expression analysis in
obesity and diabetes. *Annu Rev Nutr* **30**, 341-364 (2010).
https://doi.org:10.1146/annurev.nutr.012809.104747

99 Lopez-Gambero, A. J. *et al.* Alteration of IGF-1 bioavailability due to PAPPA2 deficiency leads
to sex-specific metabolic disturbances. *Metabolism* **171**, 156355 (2025).
https://doi.org:10.1016/j.metabol.2025.156355

100   White, M. F. IRS proteins and the common path to diabetes. *Am J Physiol Endocrinol Metab*
877   **283**, E413-422 (2002). https://doi.org/10.1152/ajpendo.00514.2001
878 101   Ding, L., Fan, L., Xu, X., Fu, J. & Xue, Y. Identification of core genes and pathways in type 2
879   diabetes mellitus by bioinformatics analysis. *Mol Med Rep* **20**, 2597-2608 (2019).
880   https://doi.org/10.3892/mmr.2019.10522
881 102   Groza, T. *et al.* The International Mouse Phenotyping Consortium: comprehensive knockout
882   phenotyping underpinning the study of human disease. *Nucleic Acids Res* **51**, D1038-D1045
883   (2023). https://doi.org/10.1093/nar/gkac972
884 103   Dickinson, M. E. *et al.* High-throughput discovery of novel developmental phenotypes. *Nature*
885   **537**, 508-514 (2016). https://doi.org/10.1038/nature19356
886 104   Stancill, J. S. *et al.* Chronic beta-Cell Depolarization Impairs beta-Cell Identity by Disrupting a
887   Network of Ca(2+)-Regulated Genes. *Diabetes* **66**, 2175-2187 (2017).
888   https://doi.org/10.2337/db16-1355
889 105   Osipovich, A. B., Stancill, J. S., Cartailler, J. P., Dudek, K. D. & Magnuson, M. A.
890   Excitotoxicity and Overnutrition Additively Impair Metabolic Function and Identity of
891   Pancreatic beta-Cells. *Diabetes* **69**, 1476-1491 (2020). https://doi.org/10.2337/db19-1145
892 106   Remedi, M. S. & Nichols, C. G. Hyperinsulinism and diabetes: genetic dissection of beta cell
893   metabolism-excitation coupling in mice. *Cell Metab* **10**, 442-453 (2009).
894   https://doi.org/10.1016/j.cmet.2009.10.011
895 107   Friedman, N. Inferring cellular networks using probabilistic graphical models. *Science* **303**, 799-
896   805 (2004). https://doi.org/10.1126/science.1094068
897 108   Sachs, K., Perez, O., Pe'er, D., Lauffenburger, D. A. & Nolan, G. P. Causal protein-signaling
898   networks derived from multiparameter single-cell data. *Science* **308**, 523-529 (2005).
899   https://doi.org/10.1126/science.1105809
900 109   Sedgewick, A. J., Shi, I., Donovan, R. M. & Benos, P. V. Learning mixed graphical models with
901   separate sparsity parameters and stability-based model selection. *BMC Bioinformatics* **17 Suppl**
902   **5**, 175 (2016). https://doi.org/10.1186/s12859-016-1039-0
903 110   Huang, G. T., Tsamardinos, I., Raghu, V., Kaminski, N. & Benos, P. V. T-ReCS: stable selection
904   of dynamically formed groups of features with application to prediction of clinical outcomes.
905   *Pac Symp Biocomput* **20**, 431-442 (2015).
906 111   Sedgewick, A. J. *et al.* Mixed graphical models for integrative causal analysis with application to
907   chronic lung disease diagnosis and prognosis. *Bioinformatics* **35**, 1204-1212 (2019).
908   https://doi.org/10.1093/bioinformatics/bty769
909 112   Raghu, V. K., Poon, A. & Benos, P. V. Evaluation of Causal Structure Learning Methods on
910   Mixed Data Types. *Proc Mach Learn Res* **92**, 48-65 (2018).
911

A

standard diet

Healthy vs. Prediabetes

C57BL/6J ♂ ♀

High Fat High Sucrose diet

Severe diabetes vs. Prediabetes

standard diet

NZO/HILTJ ♂ ♀

Severe vs. Mild diabetes

High Fat High Sucrose diet

B

C

Perturbed genes and their causal network.