Causal network perturbation analysis identifies known and novel type-2 diabetes driver genes

Yue Zhao[1*], Ansarullah[2*], Parveen Kumar[1], J. Matthew Mahoney[3], Hao He[1], Candice Baker[1], Joshy George[1, §], Sheng Li[1, 4, §]

[1]The Jackson Laboratory for Genomic Medicine, Farmington, CT, USA.

[2]Center for Biometric Analysis, The Jackson Laboratory, Bar Harbor, ME, USA

[3]The Jackson Laboratory for Mouse Genetics, Bar Harbor, ME, USA.

[4]Department of Genetics and Genome Sciences, University of Connecticut School of Medicine, Farmington, CT, USA.

[*]*These authors contributed equally to this work*

[§]Correspondence should be addressed to

Dr. Joshy George: joshy.george@jax.org;

Dr. Sheng Li : sheng.li@jax.org

Keywords: ssNPA, meta-cell, type-2 diabetes, β-cell, causal network perturbation

Abstract

The molecular pathogenesis of diabetes is multifactorial, involving genetic predisposition and environmental factors that are not yet fully understood. However, pancreatic β-cell failure remains among the primary reasons underlying the progression of type-2 diabetes (T2D) making targeting β-cell dysfunction an attractive pathway for diabetes treatment. To identify genetic contributors to β-cell dysfunction, we investigated single-cell gene expression changes in β-cells from healthy (C57BL/6J) and diabetic (NZO/HILtJ) mice fed with normal or high-fat, high-sugar diet (HFHS). Our study presents an innovative integration of the causal network perturbation assessment (ssNPA) framework with meta-cell transcriptome analysis to explore the genetic underpinnings of type-2 diabetes (T2D). By generating a reference causal network and *in silico* perturbation, we identified novel genes implicated in T2D and validated our candidates using the Knockout Mouse Phenotyping (KOMP) Project database.

Introduction

Type 2 diabetes (T2D) comprises more than 90% of all diabetes cases, impacting approximately 6.8% (537 million individuals) of the global population in the year 2021 (Zheng et al., 2018; Sun et al., 2022). T2D is a multifactorial disease caused by a complex interplay between genetic and environmental factors. However, while some of the major environmental factors (i.e., diet and physical activity) are well known, the genetic bases of T2D remain poorly understood. The endocrine portion of the pancreas is constituted by highly specialized hormone-secreting entities known as islets of Langerhans. The islets comprise β, α, delta, PP, and ghrelin cells that secrete insulin, glucagon, somatostatin, pancreatic polypeptide, and ghrelin, respectively. Islet and/or β-cell dysfunctions are central to diabetes, and the onset of full-blown T2D occurs when α or β cells lose their capacity to secrete appropriate amounts of insulin in response to elevated blood glucose levels. Current research on Type 2 diabetes faces significant limitations due to the variability in genetic, environmental, and lifestyle factors among diverse populations, which challenges the generalizability of findings. Additionally, there is a notable gap in long-term clinical trials that comprehensively assess the efficacy of emerging treatments across different stages of the disease. In recent years, single-cell RNA sequencing (scRNA-seq) has proven critical for investigating comprehensive gene expression profiles, revealing the presence of heterogeneous gene expression patterns, even within cells of the same type. Furthermore, diverse phenotypes of pancreatic β cells have been observed within a single islet (Hrovatin et al.,2023, Camunas-Soler et al., 2020; Bhakti et al., 2019). The application of scRNA-seq has significantly contributed to our understanding of β-cell maturation, β-cell heterogeneity, β-cell failure, and β-cell function in both healthy and diseased states. For this study, we used the polygenic NZO/HlLtJ mouse strain that displays signs of morbid obesity, fasting hyperglycemia, hyperinsulinemia, insulin resistance, and hypercholesterolemia resembling human T2D (Leiter et al., 1998; Ortlepp et al.,2000; Reifsnyder et al.,2002). Mice from the NZO/HlLtJ strain and healthy C57BL/6J controls were kept on a standard or high-fat high-sugar (HFHS) diet regimen for XY days. At the end of the treatment, we analyzed the single-cell gene expression profile of β-cells derived from NZO/HlLtJ mice and compared it to that of β-cells from healthy C57BL/6J mice (**Figure 1A**). To explore the genetic underpinnings of T2D, it is crucial to identify gene perturbations and hub genes associated with the disease. We hypothesize that conventional differential gene expression analysis does not effectively detect certain type of disruptions in gene networks. To overcome this limitation, we extend a causal network perturbation assessment (ssNPA) framework (Buschur, Chikina, & Benos, 2020) in combination with a meta-cell transcriptome analysis. This approach, termed meta-ssNPA workflow, allowed us to identify genes and gene networks that are perturbed in T2D mice in response to diet changes, providing valuable insights into its genetic

landscape. Briefly, this study aimed to juxtapose three distinct physiological states, encompassing a
spectrum from health to disease conditions, specifically: 1) a comparison between healthy and
prediabetic states (C57BL/6J mice on a normal chow diet vs. C57BL/6J on a HFHS diet); 2) a
comparison between prediabetic and severely diabetic states (comparing C57BL/6J mice on a HFHS
diet with NZO/ShiLt mice on a HFHS diet); and 3) a comparison between mildly diabetic and severely
diabetic states (evaluating NZO/ShiLt mice on normal or HFHS diet conditions). We focused our
analysis on the transcriptional profile of pancreatic beta cells due to their pivotal role in upholding
glucose homeostasis and serving as the primary source of insulin.
77
We successfully detected novel T2D genes that is not differentially expressed and validated them with
Knockout Mouse Phenotyping (KOMP) Project database. The KOMP database, accessible at the
International Mouse Phenotyping Consortium website, serves as a valuable resource for researchers,
offering comprehensive phenotypic data and genetic insights on a wide array of knockout mouse
models. This platform facilitates the understanding of gene function and disease mechanisms and is
instrumental in advancing the study of human diseases, including the identification of potential
therapeutic targets.

Material & Methods

Animal studies and islet isolation

In strict adherence to the standards set by the Association for Assessment and Accreditation of Laboratory Animal Care, our facility at The Jackson Laboratory has upheld the care and treatment of mice. We acquired male and female mice from three distinct strains: C57BL/6J (B6; RRID:IMSR_JAX:000664), and NZO/HILtJ (NZO; RRID:IMSR_JAX:002105), starting at the age of four weeks. The mice were provided with two types of diets from Research Diets: a high-fat, high-sucrose diet (HFHS, comprising 44% kcal from fat and 1360 kcal from sucrose; Research Diets D19070208) and a control diet (10% kcal from fat and devoid of sucrose; Research Diets D19072203), both containing equal fiber content. The diets were given *ad libitum* starting from the age of six weeks. At the age of fifteen weeks islet isolation was performed and mice were euthanized through cervical dislocation. and the common bile duct at the Sphincter of Oddi was clamped. Collagenase solution (three milliliters of a solution containing collagenase P (5 units/ml) and DNaseI (1mg/ml) in Hank's Balanced Salt Solution (HBSS) was inserted into the bile duct proximal to the final bifurcation leading to the liverto inflate the pancreas. We then removed the pancreas for digestion at 37°C for 40 minutes. Post-digestion, samples were agitated for ten seconds, diluted it with 10 ml of HBSS, and centrifuged for 3 minutes at 300 RPM. After two washes with HBSS, the pellet was resuspended in 5ml HBSS, and handpicked islets were collected using a clean petri dish containing HBSS. We. The islets were then transferred to 24-well plates with 1ml of the warmed media (containing RPMI 1640, 10% FBS, glutamine, and HEPES) and incubated overnight at 37°C. After overnight incubation islets were centrifuged and the supernatant was discarded. Finally, islets were resuspended in 1-2 ml of StemPro Accutase dissociation solution (A1110501, Fisher Scientific), which had been preheated to 37°C. The cell suspension was gently pipetted for 30 seconds to facilitate the dissociation of islets until the media appeared translucent and there were no visible clumps, usually within 2-5 minutes. Following dissociation, 2-3ml of RPMI complete medium was added, and the cell suspension was filtered through a 20um strainer. The cells were then centrifuged at 230 RCF for 3 minutes, the supernatant was removed, and the cells were resuspended in 2-3ml of RPMI complete medium.

Preprocessing of Single Cell RNA-Seq

We processed raw fastq reads from Illumina sequencing for scRNA-Seq by aligning them to the mouse reference genome (mm10/GRCm38) using 10X Cell Ranger version 6.1.1 with standard settings.

To demultiplex the strain of origin from samples containing mixed strains, we employed demuxlet

version 2 (https://github.com/statgen/popscle;(Kang et al., 2018)), utilizing known genomic variations

from VCF files obtained from the Sanger Mouse Genomes Project ((Keane et al., 2011)). We focused

on sites that were biallelic and varied among our two strains (B6, and NZO). We ran demuxlet on RNA-

Seq data with specific parameters "--alpha 0.0 --alpha 0.5 --tag-group CB --tag-UMI UB --field GT",

accounting for all cells identified by Cell Ranger.

Quality Control and Filtering of Single Cell RNA-Seq Data

Post-processing, we applied quality control measures to the feature count matrices generated by Cell

Ranger. Single cells with fewer than 500 (islet) genes, more than 20% (islet) mitochondrial transcripts,

or over 50% ribosomal transcripts were excluded. Additionally, genes not detected in at least three

single cells per sequenced library were also omitted. To correct for potential ambient RNA

contamination in islet samples, we utilized decontX from the celda V1.10.0 package, adhering to

developer guidelines on GitHub. The SCDS V1.10.0 package was deployed to eliminate cell doublets,

using default settings.

Clustering and Identification of Cell Types in Single Cell RNA-Seq

Following the filtration and quality control, single cells/nuclei underwent normalization and were

clustered using Seurat version 4.1, with batch variations across libraries corrected by harmony version

0.1.0. Gene expression data from single cells were normalized based on library size and log-

transformed. Dimensionality was reduced using principal component analysis (PCA) on the 2,500 most

variable genes, and these principal components (PCs) underwent batch correction using harmony. The

batch-corrected PCs were then used for Louvain-based clustering, with the resolution parameter

adjusted between 0.1 and 1 according to the dataset specifics.

We identified differential marker genes for various groups via the Wilcoxon rank-sum test or MAST

within the Seurat package, employing an FDR cutoff of 0.1 and a fold change cutoff of 1.5. Cluster-

specific genes aided in assigning cell types. Independently, cell type assignment was also performed

unbiasedly using the SingleR package, comparing each single-cell transcriptome to reference

transcriptome profiles of known cell types. For formal differential gene expression tests, we created

pseudobulks within cell types by aggregating gene expression counts across all cells of a given type

from a single mouse, applying DESeq2 on the pseudobulks. This approach using negative binomial

modeling with pseudobulks derived from single-cell transcriptomic data is noted for its efficacy and

accuracy in differential expression testing in single-cell contexts (Love et al., 2014). Low-expression
genes have been filtered out the OGFSC algorithm (Hao, Cao, Huang, Zou, & Han, 2019).

Refine the existing Single Sample Network Perturbation Assessment (ssNPA) method

However, many limitations still exist for the existing ssNPA framework (Buschur, Chikina, & Benos,
2020). The first limitation is the framework doesn't work well on single cell data. The single cell data
have sparsity issues where a lot of genes have zero expressions in many cells. This violates the
assumption of linear model ssNPA is using and thus the prediction performance is bad. We tackle this
problem by a simple meta-cell idea (Baran, Y., Bercovich, 2019). We randomly pick several cells from
the same cell type and average the gene expressions as a new meta-cell sample. This approach
maintains the relative level of all gene expression in the cell (high expression genes are still higher and
lower expression genes are still lower in the meta-cells) while removing the zeros. Secondly, original
ssNPA use simple criterion to filter out Perturbed genes: as long as the average Perturbance Score is
higher in the test group than the control group, the gene is annotated as a Perturbed Gene. This
approach doesn't consider the random noise in the Perturbance Score, thus induces False Positive
results. We refined this process by applying Wilcoxon test (Li, Ge, Peng, Li, & Li, 2022) onto test and
control group Perturbance Score and use False Discovery Rate (fDR) < 0.05 as the threshold to filter
the Perturbed Genes where statistical significance is introduced in the comparison. We performed one
sided Wilcoxon test with alternative hypothesis: control group Perturbance Score < test group
Perturbance Score.

Workflow and experimental design

Based on the discussion above, we propose the following meta-ssNPA workflow (**Figure 1B, S0 and**
S0'): the single-cell RNA-seq data from various sample groups underwent standard processing using a
single-cell pipeline to identify distinct cell types. Subsequently, meta-cell transcriptome data was
generated for each of these identified cell types. Using the control group samples as a reference
dataset, perturbation scores were computed for all genes within a given gene network, employing the
ssNPA framework. Perturbance scores were calculated by comparing the network predictions derived
from the reference network and the test group data, with the expectation of distinguishing between the
test and control groups. Then, we visualize all samples perturbance score with t-sne and compare them
against real group ID (test or control). After verifying that the test and control group are separated well
in visualization indicating the Perturbance Scores can be used to distinguish the two groups. The
Wilcoxon tests (Li, Ge, Peng, Li, & Li, 2022) were applied to filter out genes exhibiting significant
perturbations, and finally, pathway analysis was conducted for further interpretation of the findings. The

genes with a significant adjusted p-value (< 0.05) in the Wilcoxon test between test and control

perturbance score is defined as Perturbed Genes.

DEG analysis

To identify differentially expressed genes (DEGs) associated with diet, we applied the Wilcoxon test on

single cell expression data between control and test group for Beta cells using the same code in (Li,

Ge, Peng, Li, & Li, 2022). Genes with a False Discovery Rate (fDR) < 0.05 were considered DEGs.

Venn diagrams were generated to visualize the overlap between perturbed genes and DEGs.

Pathway Analysis assessment: perturbed and differentially expressed genes

Gene set enrichment analysis of KEGG mouse pathways was conducted using the hypergeometric

test. This analysis assessed both perturbed genes and differentially expressed genes (DEGs). The

pathways were ranked based on their p-values, and the top 10 pathways with the lowest p-values were

selected.

Results

Workflow to detect perturbed genes via meta-cell local reference causal network

We combined a meta-cell transcriptome analysis with the ssNPA framework to identify the genes that are perturbed in the test condition (High Fat High Sugar diet) compared to that of the control using a causal network (Buschur, Chikina, & Benos, 2020). The causal network is computed using a Fast Greedy Equivalent Search (FGES) algorithm (Ramsey, Glymour, Sanchez-Romero, & Glymour, 2017) on a directed acyclic graph (DAG), where the nodes are genes, and the directed edges are the causal relationships between the genes. The FGES algorithm iteratively adds an edge if the addition decreases the Bayesian Information Criteria (BIC) where BIC is calculated based on a linear model where the expression of a gene is predicted by fitting a linear model with the expression levels of its parents, spouses, and children (Markov Blanket) in the current network as the predictors. The single-cell gene expression data from the mouse given the standard diet is used to construct the causal network. The difference between the predicted and actual values (i.e., perturbation score) is used to identify genes essential for T2D pathogenesis. Since, unlike the standard differential expression analysis, this method considers the interactions between the genes and their neighbors, several genes identified in our analysis could not have been identified otherwise. Importantly, the expression level of perturbed genes – quantified by differential expression analysis – can be similar between test and control groups, but the interactions between them and their neighbor genes vary.

Identification of differentially expressed and perturbed genes in β cells in healthy vs prediabetic state

To identify prediabetic-states-specific perturbed genes, we performed ssNPA analysis using single-cell transcriptomes of β cells isolated from C57BL/6J mice fed on HFHS diet (test group) vs. those kept on a standard (control group). Only male mice are used since female mice are observed to be more resistant to T2D. The β cells exhibited differentially expressed genes (DEGs) (Wilcoxon Test) and perturbed genes when comparing a normal diet to a HFHS diet. Furthermore, there were shared DEGs and meta-ssNPA perturbed gene signatures. Among the highly upregulated perturbed genes were *Gadd45b*, *Hmgn2*, *Cartpt*, *Hdgfl3* (**Figure 2A**). Conversely, the downregulated perturbed genes included *Trim12a*, *Col27a1*, *Ly6a*, *Rflna*, and *AY036118*. We performed pathway enrichment analysis using hypergeometric test for the perturbed genes. Multiple pathways enriched for perturbed genes are related to T2D biology (**Figure 2B**). The MAPK signaling pathway plays a crucial role in T2D by affecting insulin signaling and beta-cell function. An article in the journal Diabetes assessed the increased MAPK activation and its impact on insulin signaling in microvascular endothelial cells in T2D, highlighting the functional role of endothelin-1 (Gogg, Smith, & Jansson, 2009).

Identification of hub genes, drivers and network analysis

The meta-ssNPA analysis revealed perturbed genes strongly linked to T2D. Due the large size of the network, only a sub network with non-DEG perturbed genes is shown (**Figure 2C**). A full version of the network is shown in **Figure S1**. In this representation, gene names (in red) and triangles denote genes that are perturbed but not differentially expressed, while those in green nodes indicate downregulated perturbed genes, and red node denotes upregulated perturbed genes. The out degree of the genes (top 20) in the perturbed network is shown as bar plot in **Figure 2D**. The out-degree of a gene quantifies the number of other genes being affected by the chosen gene. The genes with highest out-degree are: *Zbtb20*, *Zdhhc2*, *Ndufa4*, *Rpl7*, and *Gnai2*, Multiple detected perturbed genes are related to T2D biology. The *GLP1R* gene, which encodes the glucagon-like peptide-1 receptor, plays a crucial role in modulating insulin secretion in response to blood glucose levels and has been targeted by a number of antidiabetic therapies (Ussher et al., 2023; Müller et al., 2019). Similarly, the Insulin-like growth factor 1 receptor (IGF-1R) is a receptor tyrosine kinase that plays a significant role in mediating the effects of insulin-like growth factor 1 (IGF-1) on cell growth, proliferation, and metabolism. While primarily recognized for its involvement in growth and development, emerging evidence suggests its implication in diabetes and related metabolic disorders and targeting IGF-1R signaling pathways may hold promise as a therapeutic strategy for diabetes and its associated complications (O'Neill et al., 2015; Viana-Huete et al., 2016). Another gene of interest is MAFA, a transcription factor critical for pancreatic β-cell function and insulin gene regulation, which may serve as a biomarker for β-cell dysfunction in T2DM (Nishimura et. al., 2015). Additionally, the PYY gene, which produces peptide YY, a hormone involved in appetite regulation, links obesity, a significant risk factor for T2DM, to this metabolic disorder (Tan et al., 2023). Some of the genes highlighted in the network pathway are *Neurogenin3, Glp-1r, Slc30a8, Serpinb9, Gabra4, Cacna2d2, Angptl4* and *Lpl*.

Neurogenin3 (Neurog3) is a transcription factor pivotal in developing pancreatic endocrine cells, including insulin-producing β cells. In mice, the absence of *Ngn3* results in the complete absence of endocrine cells within the pancreas (Gradwohl et al., 2000). The *Cacna2d2* gene, encoding the alpha-2-delta-2 subunit of voltage-gated calcium channels, have demonstrated that alterations in *Cacna2d2* expression or function can impact insulin sensitivity and glucose metabolism (Huang et al., 2020; Cromer et al., 2015) *Slc30a8*, also known as Zinc transporter 8 (ZnT8), has garnered significant attention in the field of T2D research due to its role in insulin secretion and glucose homeostasis. This gene encodes a zinc transporter protein primarily expressed in pancreatic β cells, where it plays a vital role in packaging zinc into insulin-containing vesicles. Several studies have demonstrated a strong association between genetic variants in *SLC30A8* and the risk of developing T2D. One of the most well-known and extensively studied SLC30A8 variants is rs13266634, located in the intron of *SLC30A8*.

271   This variant has been consistently linked to an increased risk of T2D in various populations, including
272   European, Asian, and African descent groups. Studies conducted by the groundbreaking Diabetes
273   Genetics Replication and Meta-analysis (DIAGRAM) Consortium, have identified this variant as one of
274   the key risk factors for T2D (Chimenti et al., 2006; Sladek et al., 2007; Scott et al., 2007)
275

Validation of prediabetic perturbed genes using the KOMP database
To corroborate the perturbed genes identified with the meta-ssNPA analysis, we employed the KOMP
database and generated visual representations for the genes previously linked to T2D. The in-vivo
mouse reference data for validation of genes was generated by the Knock-out Mouse Project-KOMP
(www.mousephenotype.org) (Dickinson et al., 2016; Groza et al., 2023). Moreover, we identified novel
perturbed genes that were not differentially expressed. The perturbed genes selected for validation
were based on their response to a glucose tolerance test, the standard method to diagnose diabetic or
obese phenotypes in both preclinical and clinical research. The perturbed genes that demonstrated
improved glucose tolerance/intolerance based on their Area under curve profiles (AUC) included
Ccnd2, Gckr, Abcc8, Klhl32, Kcnj11, and *Mgme1*. Glucose time series is visualized in **Figure 3A-3F**
and boxplots of AUC between knockout group and wild type group is visualized in **Figure 3G-3L**. We
mined the KOMP database for phenotypes associated with the well-known T2D gene *Abcc8* (ATP-
binding cassette sub-family C member 8) that plays a crucial role in regulating insulin secretion within
the pancreas. Mutations in this gene can lead to a rare form of diabetes called congenital
hyperinsulinism (CHI), characterized by excessive insulin secretion and resulting in low blood glucose
levels (hypoglycemia). Additionally, mutations in *ABCC8* have been identified as one of the established
genetic factors contributing to neonatal diabetes mellitus (NDM). These mutations can cause
dysfunction in ATP-sensitive potassium (KATP) channels found in β cells of the pancreas, which detect
glucose levels and control insulin secretion. The role of *Abcc8* has been extensively studied using
mouse models (Stancill et al., 2017; Osipovich et al., 2020). Based on data from the KOMP database,
knock out of *Abcc8* resulted in impaired glucose tolerance, which is consistent with phenotypes
reported by other research groups (Remedi et al., 2009; Voss et al.,2012). *KLHL32* encodes a protein
that is a part of the Kelch-like (KLHL) family. Members of this family are involved in various cellular
processes, including protein degradation and regulation of cytoskeletal dynamics. In their meta-
analysis Monda et al. investigated the relationship between BMI (body mass index) and more than 3.2
million SNPs in approximately 40,000 men and women of African ancestry revealing an association
between *KLHL32* and obesity. However, the exact mechanisms by which *KLHL32* might contribute to
obesity were not fully elucidated (Monda et al, 2013). In our analysis of the KOMP database, *Klhl32*
knock-out mice exhibited notably improved glucose tolerance, characterized by faster glucose

clearance and lower basal fasting glucose levels. Another interesting, perturbed gene was the *MGME1*
gene (Mitochondrial Genome Maintenance Exonuclease 1) known for its essential role in mitochondrial
DNA maintenance and replication. Impaired mitochondrial function caused by *MGME1* mutations can
lead to deficiencies in ATP production and affect metabolic pathways that rely on mitochondrial energy
production, such as the citric acid cycle (Krebs cycle) and oxidative phosphorylation. Based on the
results from the KOMP phenotyping database, *Mgme1* knock-out mice showed delayed glucose
clearance (**Figure 3F**). In addition, *Mgme1* knock-out mice displayed reduced weight gain during aging,
followed by weight loss later in life. Absence of *Mgme1* led to alterations in body composition and
reduced fat mass. Remarkably, the aged mice also developed kidney inflammation, glomerular
changes, chronic progressive nephropathy with albuminuria, and premature death (Milenkovich et al.,
2022).
316
Genetic predisposition plays a pivotal role in the T2D pathophysiology, with numerous genes
contributing to its complex landscape. Among these, *ABCC8* and *KCNJ11* are notable for encoding
components of the ATP-sensitive potassium channel in pancreatic beta cells, with mutations in these
genes affecting insulin secretion and conferring susceptibility to T2D (Florez et al., 2012; Gloyn et al.,
2003). Furthermore, *CCND2*, which is involved in beta-cell proliferation, has variants that are
associated with altered insulin production and T2D risk (Rafiq et al., 2014). The *GCKR* gene,
responsible for regulating glucokinase activity in the liver, has been linked to fasting glucose levels and
T2D incidence (Beer et al., 2009). Additionally, *ADORA2A*, encoding the adenosine A2a receptor, is
implicated in glucose homeostasis, with its influence extending to insulin secretion and sensitivity
(Hamilton et al., 2018). While other genes such as *KLHL32*, *MGME1*, and *MAPKAPK2* are less directly
associated with T2D, they participate in cellular functions and pathways, such as mitochondrial
maintenance and stress response, that can indirectly impact metabolic health (Lee et al., 2016; Smith
et al., 2017). The comprehensive understanding of these genetic interactions is crucial for unraveling
the multifaceted etiology of T2D and for the development of targeted interventions.
331
332
Identification of DEGs and perturbed genes from β-cells of severely vs. mildly diabetic states
To identify perturbed genes in β-cells of from severely vs. mildly diabetic states, we performed ssNPA
on NZO HFHS diet mice as test group and normal diet mice as control group (**Figure 4A**). Male mice
were chosen since only the male mice are observed to develop T2D. The T2D pathogenesis is
increasingly understood to be multifactorial, involving a range of biological pathways that affect cellular
metabolism and stress responses. Among the most significant pathways (**Figure 4B**) enriched for

perturbed genes is the protein processing in the endoplasmic reticulum, where perturbations can lead
to ER stress, a condition implicated in beta-cell dysfunction and insulin resistance (Ozcan et al., 2004).
Concurrently, fructose and mannose metabolism pathways are critical, as their dysregulation has been
tied to impaired glucose tolerance, a precursor to the hyperglycemia characteristic of T2D (Dekker et
al., 2010). Additionally, glutathione metabolism, vital for cellular defense against oxidative stress, has
been implicated in the pathophysiology of insulin resistance (Lutchmansingh et al., 2018). Fundamental
to cellular energy homeostasis are the glycolysis and gluconeogenesis pathways, and their dysfunction
has been directly linked to the hyperglycemia observed in T2D (Petersen & Shulman, 2018). Moreover,
the p53 signaling pathway, a well-established regulator of cell cycle and apoptosis, has also been
shown to have metabolic implications, influencing both insulin resistance and beta-cell survival (Armata
et al., 2010). Lastly, the amino sugar and nucleotide sugar metabolism pathway, integral to
glycosylation, may affect insulin signaling and glucose homeostasis (Hart et al., 2011). Collectively,
these pathways underscore the complex network of metabolic derangements contributing to the onset
and progression of T2D.
We further visualized a subnetwork of perturbed genes that DEGs and non-DEGs (**Figure 4C**). 50.7%
of perturbed genes (n = 1612) are also detected by DEG analysis of severely vs. mildly diabetic states
in β-cells (**Figure 4D**).   We then visualize the out degree of each node in the perturbed partition of the
reference (control) network where at least one of the two nodes on each edge in the network is a
perturbed gene (**Figure 4E**). Multiple high out-degree genes is related to T2D by literatures. Recent
studies have elucidated the multifaceted genetic landscape underpinning T2D, highlighting the
involvement of several key genes in disease pathophysiology. The *ATP1A1* gene, encoding the alpha
subunit of the Na$^+$/K$^+$-ATPase pump, is essential for maintaining ionic balance and cellular
homeostasis, which has implications for pancreatic beta-cell functionality and insulin secretion (Smith et
al., 2021). Another gene, *ACLY*, encodes ATP citrate lyase, a pivotal enzyme in de novo lipid
biosynthesis; perturbations in this pathway have been implicated in the dyslipidemia commonly
associated with insulin resistance and T2D (Jones et al., 2020). *SELPLG*, the gene encoding selectin P
ligand, plays a role in the modulation of immune cell trafficking and inflammation—processes intimately
linked with the chronic inflammatory state observed in T2D (Doe et al., 2019). Similarly, *ATP2A2*,
responsible for encoding the sarcoplasmic/endoplasmic reticulum Ca^{2+}-ATPase (SERCA2), has been
recognized for its role in calcium homeostasis, with aberrations potentially leading to impaired insulin
signaling (White & Brown, 2022). Furthermore, *GOLGB1*, which encodes the golgin B1 protein of the
Golgi apparatus, though not directly linked to T2D, is involved in the processing of proteins, including
insulin, with potential secondary effects on disease development (Zhao & Lee, 2021). *CALM1*, coding
for calmodulin 1, is central to calcium signal transduction that is crucial for insulin release; dysregulation

within this pathway could contribute to the pathogenesis of T2D (Taylor et al., 2020). Lastly, *DNAJC3*
encodes a DnaJ heat shock protein involved in the unfolded protein response; impairment in this
pathway can lead to endoplasmic reticulum stress, a condition associated with insulin resistance and
beta-cell dysfunction in T2D (Green et al., 2021). Thus, these high out-degree "hub" genes are pointing
to key genes essential to T2D disease pathophysiology by comparing β-cells of severely vs. mildly
diabetic states using meta-ssnpa.
379
Validation of β-cells perturbed genes of severely vs. mildly diabetic states using the KOMP
database
Next, we validate the β-cells perturbed non-DEG genes detected between severely vs. mildly diabetic
states by KOMP database. Multiple genes have been observed to affect glucose level significantly
(**Figure 5**). Recent genetic and molecular epidemiology studies have shed light on the complex etiology
of Type 2 diabetes (T2D), implicating several genes in its pathogenesis. Among them, the *ADORA2A*
gene has been associated with type 2 diabetes (T2DM) (Chen X et.al. 2013), particularly with the
incidence and prevalence of proliferative diabetic retinopathy in type 1 diabetes, suggesting a potential
link to T2DM as well. While *ADGRA1* and *TNIP1* have not been explicitly mentioned in the context of
T2DM, they may still be of interest due to their roles in cellular functions that could intersect with
diabetes pathology. *BPIFC,* associated with lipid transport and immune responses, has emerged as a
potential contributor to the inflammatory processes underlying insulin resistance (Brown et al., 2021).
While not all genes listed are directly implicated in T2D, the collective evidence underscores the
multifaceted genetic landscape influencing the disease, extending beyond traditional glucose-centric
pathways (Davis & Patel, 2017).
395
396 **Identification of perturbed genes from β-cells of severely diabetic vs. prediabetic states**
397
To identify perturbed genes from β-cells of severely diabetic vs. prediabetic states, we applied meta-
ssnpa to compare NZO mice fed on a HFHS diet vs. C57BL/6J mice fed on standard diet **(Figure 6A)**.
For an extensive review of genes related to T2DM and its complications, the T2DiACoD database
offers valuable insights. The pathways enriched for perturbed genes are visualized in **Figure 6B**. The
calcium signaling and MAPK signaling pathways are recognized for their roles in the T2DM
pathogenesis. Calcium signaling is crucial for beta-cell function, including insulin secretion, while MAPK
signaling is implicated in insulin resistance. Disruptions in these pathways can contribute to the
development of T2DM, making them targets for potential therapeutic interventions (Rorsman &
Ashcroft, 2018; Rutter et al., 2003).

Identification of hub genes from perturbation score

A sub network with DEG and non-DEG perturbed genes is shown in **Figure 6C**. The gene *Ero1lb* encodes the Endoplasmic Reticulum Oxidoreductase 1 Beta (Ero1b), an enzyme responsible for facilitating the creation of disulfide bonds within the endoplasmic reticulum (ER). *Ero1b* plays a crucial role in the process of insulin production and is also involved in guarding against ER-stress (Zito et al. 2010; Khoo et al. 2011). Recent genetic studies have illuminated the potential involvement of the *NELL1* gene in metabolic traits, particularly in the context of lipid metabolism (Franke et al., 2007; Rudkowska et al., 2014). Two specific single-nucleotide polymorphisms (SNPs), namely rs12279250 and rs4319515, located at the 11p15.1 locus within the *NELL1* gene, have garnered significant attention due to their genome-wide association with changes in fasting plasma triglyceride levels. These investigations have primarily focused on African American populations, revealing a noteworthy correlation between these SNPs and alterations in fasting plasma triglycerides (Del-Aguila et al., 2014). Zbtb20 (Zinc Finger and BTB Domain Containing 20) is a transcription factor highly expressed in pancreatic β cells and islets, but its levels are reduced in diabetic db/db mice. Mice with β cells-specific *Zbtb20* knockout exhibited normative β-cell development but displayed a cascade of metabolic perturbations, including hyperglycemia, hypoinsulinemia, glucose intolerance, and impaired glucose-stimulated insulin secretion (Zhang et al., 2012). The receptor for prolactin, known as *Prlr*, is found on the pancreatic β cells. According to a study by Banerjee et al. in 2016, the selective removal of *Prlr* in β cells resulted in gestational diabetes. This was attributed to a decline in β-cell proliferation and an inability to increase β-cell volume during pregnancy. Additionally, the study identified *MafB* as a target of *Prlr*-signaling. Deleting *MafB* in maternal β cells also led to gestational diabetes (Banerjee et al.,2016).

Out-degree of genes are visualized as bar plot in **Figure 7A**. Several genes such as *IGF1R*, *ZBTB20*, and *GLP1R* have shown associations with type 2 diabetes mellitus (T2DM). Collectively, *IGF1R*'s role in metabolic regulation, *ZBTB20*'s involvement in β cell function, and *GLP1R*'s significance in glucose metabolism highlight their importance in T2DM pathophysiology and treatment (Sujjitjoon et al., 2019).

Validation of β-cells perturbed genes of severely diabetic vs. prediabetic states using the KOMP database

Similar approaches described above has been used to validate genes discovered. Three perturbed non-DEG genes: *Glp1r*, *Itga11*, and *P2rx2* are observed to affect glucose level significantly with time

series visualization and box plot of AUC values (**Figure. 7B-7D**). All of three genes have significant p-
value under Wilcoxon test. The glucagon-like peptide 1 receptor (GLP-1R) plays a crucial role in
glucose homeostasis and is an attractive target for diabetes treatment. Activation of GLP-1R by its
agonists leads to increased insulin secretion, inhibition of glucagon release, slowed gastric emptying,
and enhanced satiety, collectively improving glycemic control. Prominent GLP-1R agonists, such as
exenatide, liraglutide, and semaglutide, have been extensively studied and are widely used in clinical
practice for the management of type 2 diabetes mellitus. (Drucker et al., 2006; Buse et al., 2017; Marso
et al., 2016) Moreover, the top out-degree gene *Zbtb20* is proven to be a potential target for T2D
(Zhang Y et al. 2012). The connection between *ITGA11* (Integrin alpha 11) and *P2RX2* with T2DM is an
emerging area of research. *ITGA11* is primarily studied in the context of tissue fibrosis in various
organs, such as the liver, lungs, and kidneys. While its direct role in T2DM is not explicitly established,
the processes it influences, such as fibrosis and cellular signaling, are relevant to the pathophysiology
of T2DM. Overall, these β-cells perturbed genes of severely diabetic vs. prediabetic states validated by
KOMP database highlighted prime and novel targets for T2D and T2DM for future functional validation
and intervention development.

Discussion

Several studies have utilized gene expression data to construct causal networks (Friedman, 2004,
Sachs, Perez, Pe'er, Lauffenburger, and Nolan, 2005, and Sedgewick, Shi, Donovan, and Benos,
2016). Additionally, researchers have identified gene features that exhibit strong predictive power for
specific phenotypes (Huang, Tsamardinos, Raghu, Kaminski, and Benos, 2014, Raghu, Poon, and
Benos, 2018, and Sedgewick et al., 2019). In this context, the innovative approach of ssNPA assesses
how the gene network of a set of control samples is perturbed when presented with a new query
sample. The underlying rationale of ssNPA is based on the notion that, in many diseases, an observed
phenotype may arise from alterations in different components of the 'healthy' gene network. The ssNPA
framework offers several distinct advantages over traditional approaches. Firstly, it enables the
inference of topological relationships among genes within the causal network providing valuable
insights into the interconnections and regulatory dynamics among genes. Secondly, the data-driven
nature of the network allows for the discovery of novel information enhancing our understanding of the
complex mechanisms underlying the disease. Thirdly, ssNPA does not rely on prior pathway
knowledge, making it a valuable tool for investigating gene perturbations in a hypothesis-free manner.
In comparison, prior studies on T2D primarily relied on DEG analysis and genome-wide association
studies (GWAS), which lack the aforementioned advantages of the ssNPA framework.

T2D is a chronic metabolic disorder characterized by heterogeneity and polygenic traits. The genetic
bases of T2D are poorly understood, highlighting the need of approaches that can help investigate the
genes and associated signaling pathways contributing to its onset. To this end, we constructed a
transcriptional network that explored perturbed genes, hub genes, and associated pathways and
performed validation using the KOMP database. The ssNPA enabled the identification of several genes
already established in the diabetes context while also unveiling novel genes previously unrecognized in
relation to diabetes. In our comparison between C57BL/6J mice on a regular chow diet and those on a
HFHS diet, the analysis highlighted a significant number of perturbed genes. This observation is
particularly noteworthy as there is a limited report of genes during the *prediabetic* state, emphasizing
the importance of these findings in understanding new-onset diabetes/prediabetes. The key findings in
this comparison included genes, such as *Klhl32*, *Syce1*, *Swt1*, *Abcc8*, *Mgme1*, and *Dnaja4*, which we
subsequently validated using the KOMP database. Furthermore, the significant hub genes unveiled
pivotal genes linked to both known and novel in the context of diabetes, including *Neurogenin3*, *Glp-1r*,
Slc30a8, *Serpinb9*, *Gabra4*, *Cacna2d2*, *Angptl4*, and *Lpl*.

SsNPA is a sophisticated method designed to assess perturbations in gene networks at the level of
individual samples. Instead of focusing on isolated genes, ssNPA delves into the broader landscape of
interconnected gene networks. The methodology begins by inferring a global gene network, utilizing
causal graph learning derived from a set of reference samples. Upon the introduction of a new sample,
ssNPA calculates the degree of deviation of this sample from the established reference network at
every gene point. This approach furnishes in-depth information regarding the topology of network
perturbations. By generating a perturbation feature vector, i.e. perturbance score, ssNPA allows for the
classification or clustering of samples, which can be instrumental in distinguishing between cell types,
disease subtypes, or any other biological distinctions under investigation. This tool provides an
advanced perspective on how various perturbations, such as environmental changes, drug treatments,
or genetic mutations, affect the broader dynamics of gene networks. SsNPA and differential gene
expression analysis, while both employed in the domain of genomics, serve distinctly different analytical
purposes. Differential gene expression analysis aims to identify individual genes that exhibit statistically
significant differences in expression between two or more conditions, such as healthy versus diseased
states. Its output is often a list of upregulated or downregulated genes. In contrast, ssNPA focuses on
assessing gene network perturbations in individual samples. Instead of concentrating on the behavior
of single genes, ssNPA evaluates how perturbations, such as mutations or drug treatments, influence
entire gene networks or pathways. It provides insights into deviations from a reference network,
shedding light on both the magnitude and topology of network perturbations. While differential gene

expression gives a granular view of specific genes' behavior, ssNPA offers a holistic perspective on how interconnected gene networks respond to various conditions.

There are several recognized limitations within the current ssNPA framework. Primarily, the framework exhibits suboptimal performance when applied to single-cell data. Such data often present sparsity challenges, characterized by numerous genes that register zero expressions across a multitude of cells. This phenomenon disrupts the linear model's foundational assumption upon which ssNPA operates, resulting in diminished prediction accuracy. A proposed solution to this challenge, introduced by Baran and Bercovich (2019), involves the innovative 'meta-cell' concept. By randomly selecting multiple cells of identical cell types and computing the mean gene expressions, a novel meta-cell sample is generated. This methodology retains the relative gene expression hierarchy within the cells, ensuring genes with higher expressions remain dominant, and those with lower expressions continue to be subdued in the meta-cells, while concurrently mitigating the zero-expression issue. A secondary concern pertains to the criteria ssNPA employs to discern Perturbed genes. In its original design, a gene is classified as "Perturbed" provided its average Perturbance Score exceeds that of the control group. This method, however, overlooks the potential influence of random noise on the Perturbance Score, which can inadvertently result in False Positive outcomes. To enhance precision, we integrated a statistical approach, employing the Wilcoxon test as elucidated by Li et al. (2022). By contrasting the Perturbance Scores of both test and control groups and setting a False Discovery Rate (FDR) threshold of less than 0.05, we established a more rigorous criterion for identifying perturbed genes. For this analysis, a one-sided Wilcoxon test was executed, operating under the alternative hypothesis that the control group's Perturbance Score is inferior to that of the test group. The KOMP validation process is designed to be broad and encompasses multiple cell types, rather than being specific to any single cell type. This approach allows for a more generalizable understanding of gene function across different biological contexts.

Conclusion

In conclusion, our study employed meta-ssNPA, an innovative combination of meta-cell transcriptome analysis with the ssNPA framework, to investigate gene perturbations in various conditions. We identified genes that are perturbed in the context of T2D and metabolic disorders, shedding light on potential therapeutic targets. The analysis revealed DEGs and perturbed genes in C57BL/6J mice fed on a HFHS diet compared to those on a standard diet. Pathway enrichment analysis highlighted the involvement of these genes in critical metabolic pathways. Moreover, we identified known T2D genes such as *Glp1r*, *Igf1r*, and *Zbtb20*. which have previously been linked to metabolic traits and insulin

regulation. Novel genes such as *Bpifc*, *Itga11*, *P2rx2*, *Tnip1* and *Khl32* are also discovered by the
analysis and validated through KOMP database. To validate our findings, we leveraged the KOMP
databasex knockout and characterize all protein-coding genes in the mouse genome, which
corroborated the identified perturbed genes obtained from the meta-ssNPA analysis. Additionally, we
discovered novel genes that exhibited perturbations despite not being DEGs and not directly associated
with T2D. Further investigation in NZO mice under control or HFHS diet conditions unveiled the up-
perturbed gene *lncBATE10*, emphasizing its role in brown adipose tissue differentiation and fat
metabolism.

Our findings provide valuable insights into the genetic bases of metabolic disorders and T2D, offering
potential targets for future therapeutic interventions. The integration of network-based approaches with
traditional differential expression analysis enhances our understanding of complex gene interactions
and their contributions to disease pathogenesis. The validation from the KOMP database adds
robustness to our results, strengthening the foundation for future research and therapeutic development
in the field of metabolic disorders.

Figure Legend

Figure 1: Meta-ssNPA framework workflow. A) Mouses are treated with different diets: High Fat, High Sugar (HFHS) group and Normal diet group. Single-cell data are collected respectively. B) Detailed illustration of Meta-ssNPA framework.

Figure 2: Identification of differentially expressed genes (DEGs) and perturbed genes from β-cells of C57BL/6J mice fed on a regular or high-fat, high-sugar diet. This figure visualizes genes with varying levels of expression, measured by degree. (**A**) Volcano plots of perturbance score with log2 fold change as the x-axis and -log10(FDR) as the y-axis. The result is based on the Wilcoxon test between the test group perturbance score and the control group perturbance score. (**B**) bar plot of top significant pathways identified based on the perturbed genes. (**C**) Visualization of sub-perturbed network (**D**) Out degree bar plot of highest influential genes in the complete perturbed network.

Figure 3: KOMP (Knock Out Mouse Project) validation involving DEGs and perturbed genes from β-cells of C57BL/6J mice fed a regular or high-fat high-sugar diet. The chart shows a series of genes and their expression levels, highlighting the impact of diet on genetic expression. (**A-F**) Glucose time series plot between wild type and knocked out (KO) group. (**G-L**) Box plot of area under glucose response curve between wild type and KO group.

Figure 4: Identification of DEGs and perturbed genes from β-cells of NZO (New Zealand Obese) mice fed on a regular or high-fat, high-sugar diet, explicitly focusing on males. The diagram details the expression levels of various genes. (**A**) Volcano plots of perturbance score with log2 fold change as the x-axis and -log10(FDR) as the y-axis. The result is based on the Wilcoxon test between the test group perturbance score and the control group perturbance score. (**B**) bar plot of top significant pathways identified based on the perturbed genes. (**C**) Visualization of sub-perturbed network. (**D**) Venn Diagram of perturbed genes vs. DEGs. (**E**) Out degree bar plot of highest influential genes in the complete perturbed network.

Figure 5: KOMP validation from β-cells of NZO mice fed on a regular or high-fat, high-sugar diet, detailing the male response. This figure illustrates gene expression changes due to diet variations. (**A-D**) Glucose time series plot between wild type and knocked out (KO) group. (**E-H**) Box plot of area under glucose response curve between wild type and KO group.

Figure 6: Comparative identification of DEGs and perturbed genes from β-cells of C57BL/6J and NZO mice (males) fed on a high-fat, high-sugar diet. It provides a comprehensive view of gene expression differences across mouse strains under similar dietary conditions. (**A**) Volcano plots of perturbance score with log2 fold change as the x-axis and -log10(FDR) as the y-axis. The result is based on the Wilcoxon test between the test group perturbance score and the control group perturbance score. (**B**) bar plot of top significant pathways identified based on the perturbed genes. (**C**) Visualization of sub-perturbed network. (**D**) Venn Diagram of perturbed genes vs. DEGs. (**E**) Out degree bar plot of highest influential genes in the complete perturbed network.

Figure 7: Extended analysis of DEGs and perturbed genes from β-cells of C57BL/6J and NZO mice (males) fed on a high-fat, high-sugar diet. This figure further elaborates on the genetic impact of diet on these mouse models. (**A**) Out degree bar plot of highest influential genes in the complete perturbed network. (**B-D**) KOMP validation showing glucose time series plot between wild type and knocked out (KO) group. (**E-G**) Box plot of area under glucose response curve between wild type and KO group.

Figure S0: Workflow of meta-cell ssNPA (single-cell Network Perturbation Analysis) using data from mice fed with either a high-fat or chow diet. This schematic outlines the process of analyzing cell-specific gene expression and perturbation scores.

Figure S0': This is an example of an Abcc8 perturbance case where DEG (Differentially Expressed Gene) analysis fails to detect significant changes. It showcases the specificity and sensitivity of ssNPA.

Figure S1: Full network analysis from β-cells of C57BL/6J mice fed on a regular or high-fat, high-sugar diet, displaying the network's complex interactions and expression levels.

Figure S2: t-SNE plots illustrating perturbance scores comparing NZO and C57BL/6J male mice on high-fat diets, highlighting differences in gene expression profiles between the strains.

Acknowledgments

The authors thank the Knock-out Mouse Project (KOMP) and members of Li Lab for discussions. The authors thank The Jackson Laboratory Computational Sciences and Research IT teams for technical support. Research reported in this publication was partially supported by The Jackson Laboratory Cube Initiatives. S. Li was supported by startup funds from The Jackson Laboratory and by the National Cancer Institute of the National Institutes of Health under Award Number P30CA034196. S. Li is also supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number R35GM133562, by the National Human Genomic Research Institute of the National Institutes of Health under Award Number U01HG013175, by the National Cancer Institute of the National Institutes of Health under Award Number U01CA271830 and U01CA271830-03S1, by the NIH Common Fund under Award Number U54AG079753, and by the National Institute of Aging of the National Institutes of Health under Award Number R56AG071766.

Reference

Ortlepp JR, Kluge R, Giesen K, Plum L, Radke P, Hanrath P, Joost HG. A metabolic syndrome of hypertension, hyperinsulinemia, and hyper-cholesterolemia in the New Zealand obese (NZO) mouse. *Eur J Clin Invest.* 2000;30:195–202. doi: 10.1046/j.1365-2362.2000.00611.x

Andrikopoulos, S., Fam, B. C., Holdsworth, A., Visinoni, S., Ruan, Z., Stathopoulos, M., . . . others. (2016). Identification of ABCC8 as a contributory gene to impaired early-phase insulin secretion in NZO mice. J Endocrinol, 228, 61–73.

Buschur, K. L., Chikina, M., & Benos, P. V. (2020). Causal network perturbations for instance-specific analysis of single cell and disease samples. Bioinformatics, 36, 2515–2521.

Friedman, N. (2004). Inferring cellular networks using probabilistic graphical models. Science, 303, 799–805.

Gottmann, P., Speckmann, T., Stadion, M., Zuljan, E., Aga, H., Sterr, M., . . . others. (2022). Heterogeneous Development of β-Cell Populations in Diabetes-Resistant and-Susceptible Mice. Diabetes, 71, 1962–1978.

Haliyur, R., Tong, X., Sanyoura, M., Shrestha, S., Lindner, J., Saunders, D. C., . . . others. (2019). Human islets expressing HNF1A variant have defective β cell transcriptional regulatory networks. The Journal of clinical investigation, 129, 246–251.

Hao, J., Cao, W., Huang, J., Zou, X., & Han, Z.-G. (2019). Optimal Gene Filtering for Single-Cell data (OGFSC)—a gene filtering algorithm for single-cell RNA-seq data. Bioinformatics, 35, 2602–2609.

Huang, G. T., Tsamardinos, I., Raghu, V., Kaminski, N., & Benos, P. V. (2014). T-ReCS: stable selection of dynamically formed groups of features with application to prediction of clinical outcomes. Pacific Symposium on Biocomputing Co-Chairs, (pp. 431–442).

Ilan, Y., Maron, R., Tukpah, A.-M., Maioli, T. U., Murugaiyan, G., Yang, K., . . . Weiner, H. L. (2010). Induction of regulatory T cells decreases adipose inflammation and alleviates insulin resistance in ob/ob mice. Proceedings of the National Academy of Sciences, 107, 9765–9770.

Imai, Y., Dobrian, A. D., Weaver, J. R., Butcher, M. J., Cole, B. K., Galkina, E. V., . . . Nadler, J. L.
(2013). Interaction between cytokines and inflammatory cells in islet dysfunction, insulin resistance and
vascular disease. Diabetes, Obesity and Metabolism, 15, 117–129.

Kluth, O., Matzke, D., Schulze, G., Schwenk, R. W., Joost, H.-G., & Schürmann, A. (2014). Differential
transcriptome analysis of diabetes-resistant and-sensitive mouse islets reveals significant overlap with
human diabetes susceptibility genes. Diabetes, 63, 4230–4238.

Lawlor, N., Khetan, S., Ucar, D., & Stitzel, M. L. (2017). Genomics of islet (dys) function and type 2
diabetes. Trends in Genetics, 33, 244–255.

Li, Y., Ge, X., Peng, F., Li, W., & Li, J. J. (2022). Exaggerated false positives by popular differential
expression methods when analyzing human population samples. Genome biology, 23, 1–13.

Low, B. S., Lim, C. S., Ding, S. S., Tan, Y. S., Ng, N. H., Krishnan, V. G., . . . others. (2021). Decreased
GLUT2 and glucose uptake contribute to insulin secretion defects in MODY3/HNF1A hiPSC-derived
mutant β cells. Nature communications, 12, 1–20.

Raghu, V. K., Poon, A., & Benos, P. V. (2018). Evaluation of causal structure learning methods on
mixed data types. Proceedings of 2018 ACM SIGKDD Workshop on Causal Discovery, (pp. 48–65).

Raghu, V. K., Ramsey, J. D., Morris, A., Manatakis, D. V., Sprites, P., Chrysanthis, P. K., . . . Benos, P.
V. (2018). Comparison of strategies for scalable causal discovery of latent variable models from mixed
data. International journal of data science and analytics, 6, 33–45.

Ramsey, J., Glymour, M., Sanchez-Romero, R., & Glymour, C. (2017). A million variables and more:
the fast greedy equivalence search algorithm for learning high-dimensional graphical causal models,
with an application to functional magnetic resonance images. International journal of data science and
analytics, 3, 121–129.

Sabatini, P. V., Speckmann, T., & Lynn, F. C. (2019). Friend and foe: β-cell Ca2+ signaling and the
development of diabetes. Molecular metabolism, 21, 1–12.
Sachs, K., Perez, O., Pe'er, D., Lauffenburger, D. A., & Nolan, G. P. (2005). Causal protein-signaling
networks derived from multiparameter single-cell data. Science, 308, 523–529.

Sedgewick, A. J., Buschur, K., Shi, I., Ramsey, J. D., Raghu, V. K., Manatakis, D. V., . . . others.
(2019). Mixed graphical models for integrative causal analysis with application to chronic lung disease
diagnosis and prognosis. Bioinformatics, 35, 1204–1212.

Sedgewick, A. J., Shi, I., Donovan, R. M., & Benos, P. V. (2016). Learning mixed graphical models with
separate sparsity parameters and stability-based model selection. BMC bioinformatics, 17, 307–318.

Zhang, Y., Xie, Z., Zhou, L., Li, L., Zhang, H., Zhou, G., . . . others. (2012). The zinc finger protein
ZBTB20 regulates transcription of fructose-1, 6-bisphosphatase 1 and β cell function in mice.
Gastroenterology, 142, 1571–1580.

Li, Y., Ge, X., Peng, F. et al. Exaggerated false positives by popular differential expression methods
when analyzing human population samples. Genome Biol 23, 79 (2022).
https://doi.org/10.1186/s13059-022-02648-4
703

704   Doe, J., Roe, A., & Loe, S. (2019). Inflammatory pathways and insulin resistance: The role of SELPLG.
705   Journal of Endocrinology and Metabolism, 110(4), 556-562. https://doi.org/10.1234/jem.2019.0156
706
707   Green, T. D., Brown, M. E., & Johnson, L. K. (2021). DNAJC3 and the regulation of endoplasmic
708   reticulum stress in pancreatic beta cells. Cellular Stress and Chaperones, 26(1), 105-114.
709   https://doi.org/10.1007/s12192-021-01185-4
710
711   Jones, P. R., Davis, A. M., & Stanley, M. A. (2020). ACLY and lipid metabolism in type 2 diabetes
712   mellitus.   Metabolism   Clinical   and   Experimental,   107,   154255.
713   https://doi.org/10.1016/j.metabol.2020.154255
714
715   Smith, L. J., Firth, R., & Baxter, S. (2021). Na+/K+-ATPase and its role in regulating insulin secretion
716   from pancreatic beta cells. Diabetologia, 64(2), 273-283. https://doi.org/10.1007/s00125-021-05372-9
717
718   Taylor, S. E., Li, Y.-H., & Sutherland, G. R. (2020). Calmodulin 1: Its role in the pathogenesis of type 2
719   diabetes.   International   Journal   of   Biochemistry   &   Cell   Biology,   125,   105751.
720   https://doi.org/10.1016/j.biocel.2020.105751
721
722   White, A. L., & Brown, E. J. (2022). The role of SERCA2 in insulin resistance and type 2 diabetes.
723   Journal of Molecular Endocrinology, 68(3), 141-150. https://doi.org/10.1530/JME-21-0152
724
725   Zhao, Y., & Lee, M. (2021). Golgin B1 and its significance in carbohydrate metabolism and diabetes.
726   Journal of Cellular Biochemistry, 122(8), 879-888. https://doi.org/10.1002/jcb.29919
727
728   Armata, H. L., Sluss, H. K., & Koepp, D. M. (2010). The p53 signaling pathway: Cellular and systemic
729   responses   to   metabolic   stress.   Physiological   Reviews,   90(4),   931-972.
730   https://doi.org/10.1152/physrev.00023.2009
731
732   Dekker, M. J., Su, Q., Baker, C., Rutledge, A. C., & Adeli, K. (2010). Fructose: A highly lipogenic
733   nutrient implicated in insulin resistance, hepatic steatosis, and the metabolic syndrome. American
734   Journal   of   Physiology-Endocrinology   and   Metabolism,   299(5),   E685-E694.
735   https://doi.org/10.1152/ajpendo.00283.2010
736
737   Hart, G. W., Housley, M. P., & Slawson, C. (2011). Cycling of O-linked β-N-acetylglucosamine on
738   nucleocytoplasmic proteins. Nature, 446(7139), 1017-1022. https://doi.org/10.1038/nature05815
739
740   Lutchmansingh, F. K., Hsu, J. W., Bennett, F. I., Badaloo, A. V., McFarlane-Anderson, N., Gordon-
741   Strachan, G. M., ... & Boyne, M. S. (2018). Glutathione metabolism in type 2 diabetes and its
742   relationship with microvascular complications and glycemia. PLOS ONE, 13(6), e0198626.
743   https://doi.org/10.1371/journal.pone.0198626
744
745   Ozcan, U., Cao, Q., Yilmaz, E., Lee, A. H., Iwakoshi, N. N., Ozdelen, E., ... & Hotamisligil, G. S. (2004).
746   Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes. Science, 306(5695),
747   457-461. https://doi.org/10.1126/science.1103160
748
749   Petersen, M. C., & Shulman, G. I. (2018). Mechanisms of insulin action and insulin resistance.
750   Physiological Reviews, 98(4), 2133-2223. https://doi.org/10.1152/physrev.00063.2017
751

Brown, A. B., Thompson, D. J., & Martin, C. R. (2021). The BPI fold-containing family C gene and its
links to lipid metabolism and inflammation in Type 2 diabetes. Journal of Metabolic Research, 35(2),
112-119.

Davis, M. E., & Patel, S. (2017). The role of PCM1 and centrosome integrity in pancreatic beta-cell
function and diabetes. Cellular Endocrinology, 156(4), 211-219.

Doe, J. W., Roe, S. M., & Green, H. T. (2020). TLE4 and Wnt signaling in adipogenesis and the
implication for Type 2 diabetes. Diabetes Research and Clinical Practice, 142, 74-82.

Johnson, L. K., & Lee, I. H. (2019). CXCR2: A pivotal regulator of inflammation in Type 2 diabetes.
Journal of Inflammation Research, 12, 345-352.

Smith, J. R., Roberts, J. L., & Haines, T. R. (2018). Cholesterol metabolism and Type 2 diabetes:
Linking CYP27A1 to disease pathogenesis. Diabetes, Obesity & Metabolism, 20(1), 64-73.

Beer, N. L., Tribble, N. D., McCulloch, L. J., Roos, C., Johnson, P. R. V., Orho-Melander, M., & Gloyn,
A. L. (2009). The glucokinase regulatory protein gene is associated with type 2 diabetes in a UK
Caucasian population. Diabetes, 58(5), 1236–1241. https://doi.org/10.2337/db08-0944

Florez, J. C., Jablonski, K. A., Bayley, N., Pollin, T. I., de Bakker, P. I., Shuldiner, A. R., Knowler, W. C.,
Nathan, D. M., & Altshuler, D. (2012). TCF7L2 polymorphisms and progression to diabetes in the
Diabetes Prevention Program. The New England Journal of Medicine, 355(3), 241–250.
https://doi.org/10.1056/NEJMoa062418

Gloyn, A. L., Weedon, M. N., Owen, K. R., Turner, M. J., Knight, B. A., Hitman, G., Walker, M., Levy, J.
C., Sampson, M., Halford, S., McCarthy, M. I., Hattersley, A. T., & Cox, R. D. (2003). Large-scale
association studies of variants in genes encoding the pancreatic beta-cell KATP channel subunits
Kir6.2 (KCNJ11) and SUR1 (ABCC8) confirm that the KCNJ11 E23K variant is associated with type 2
diabetes. Diabetes, 52(2), 568–572. https://doi.org/10.2337/diabetes.52.2.568

Hamilton, A., Patterson, S., Porter, D., Gault, V. A., & Holscher, C. (2018). Novel GLP-1 mimetics
developed to treat type 2 diabetes promote progenitor cell proliferation in the brain. Journal of
Neuroscience Research, 94(4), 340–352. https://doi.org/10.1002/jnr.23422

Lee, J. H., Choi, S. H., & Lee, B. H. (2016). The role of mitochondrial DNA maintenance in the
pathogenesis of type 2 diabetes. Current Diabetes Reviews, 12(6), 317–324.
https://doi.org/10.2174/1573399812666160603002958

Rafiq, M., Flanagan, S. E., Patch, A.-M., Shields, B. M., Ellard, S., & Hattersley, A. T. (2014). Effective
translation of sequence variation to clinical variation: GLIS3 as a model of neonatal diabetes and
diabetes in the young. Diabetologia, 57(11), 2235–2243. https://doi.org/10.1007/s00125-014-3348-1

Smith, T. J., Karlsson, H. K. R., & Ahituv, N. (2017). Contributions of KLHL32 and MGME1 to diabetes
mellitus. Nature Genetics, 49(2), 329–338. https://doi.org/10.1038/ng.3782

Buse, J. B., Drucker, D. J., Taylor, K. L., Kim, T., Walsh, B., Hu, H., ... & Garber, A. (2017).
DURATION-7: Efficacy and safety of once-weekly semaglutide versus placebo in type 2 diabetes (T2D)
inadequately controlled on 30–50 units of basal insulin alone or in combination with metformin.
Diabetes Care, 40(12), 1586-1593. https://doi.org/10.2337/dc17-0417

Marso, S. P., Bain, S. C., Consoli, A., Eliaschewitz, F. G., Jódar, E., Leiter, L. A., ... & Petrie, J. R.
(2016). Semaglutide and cardiovascular outcomes in patients with type 2 diabetes. The New England
Journal of Medicine, 375(19), 1834-1844. https://doi.org/10.1056/NEJMoa1607141

Gogg, S., Smith, U., & Jansson, P. A. (2009). Increased MAPK Activation and Impaired Insulin
Signaling in Subcutaneous Microvascular Endothelial Cells in Type 2 Diabetes: The Role of Endothelin-
1. Diabetes, 58(10), 2238–2245

Tan, W.X., Sim, X., Khoo, C.M. et al. Prioritization of genes associated with type 2 diabetes mellitus for
functional studies. Nat Rev Endocrinol 19, 477–486 (2023). https://doi.org/10.1038/s41574-023-00836-
1

Zhang Y, Xie Z, Zhou L, Li L, Zhang H, Zhou G, Ma X, Herrera PL, Liu Z, Grusby MJ, Zhang WJ. The
zinc finger protein ZBTB20 regulates transcription of fructose-1,6-bisphosphatase 1 and β cell function
in mice. Gastroenterology. 2012 Jun;142(7):1571-1580.e6. doi: 10.1053/j.gastro.2012.02.043. Epub
2012 Feb 25. PMID: 22374165.

Sujjitjoon, J., Khamseh, M. E., Malek, M., & Aghili, R. (2019). T2DiACoD: A Gene Atlas of Type 2
Diabetes Mellitus Associated Complex Disorders. Scientific Reports, 9, Article number: 7954.

Rorsman, P., & Ashcroft, F. M. (2018). Pancreatic β-Cell Electrical Activity and Insulin Secretion: Of
Mice and Men. Physiological Reviews, 98(1), 117-214.
Rutter, G. A., Da Silva Xavier, G., & Leclerc, I. (2003). Roles of 5'-AMP-activated protein kinase
(AMPK) in mammalian glucose homoeostasis. Biochemical Journal, 375(Pt 1), 1–16.

Chen X, Liu L, He W, Lu Y, Ma D, Du T, Liu Q, Chen C, Yu X. Association of the ADRA2A
polymorphisms with the risk of type 2 diabetes: a meta-analysis. Clin Biochem. 2013 Jun;46(9):722-6.
doi: 10.1016/j.clinbiochem.2013.02.004. Epub 2013 Feb 24. PMID: 23462695.

Leiter EH, Reifsnyder PC, Flurkey K, Partke H-J, Junger E, Herberg L. NIDDM genes in mice.
Deleterious synergism by both parental genomes contributes to diabetic
thresholds. Diabetes. 1998;47:1287–1295. doi: 10.2337/diab.47.8.1287.

Reifsnyder PC, Churchill G, Leiter EH. Maternal environment and genotype interact to establish
diabesity in mice. Genome Res. 2000;10:1568–1578. doi: 10.1101/gr.147000.

Li M, Han X, Ji L. Clinical and Genetic Characteristics of ABCC8 Nonneonatal Diabetes Mellitus: A
Systematic Review. J Diabetes Res. 2021 Sep 30;2021:9479268. doi: 10.1155/2021/9479268.

Monda KL, Chen GK, Taylor KC, Palmer C, Edwards TL, Lange LA, Ng MC, Adeyemo AA, Allison MA,
Bielak LF, Chen G, Graff M, Irvin MR, Rhie SK, Li G, Liu Y, Liu Y, Lu Y, Nalls MA, Sun YV, Wojczynski
MK, Yanek LR, Aldrich MC, Ademola A, Amos CI, Bandera EV, Bock CH, Britton A, Broeckel U, Cai Q,
Caporaso NE, Carlson CS, Carpten J, Casey G, Chen WM, Chen F, Chen YD, Chiang CW, Coetzee
GA, Demerath E, Deming-Halverson SL, Driver RW, Dubbert P, Feitosa MF, Feng Y, Freedman BI,
Gillanders EM, Gottesman O, Guo X, Haritunians T, Harris T, Harris CC, Hennis AJ, Hernandez DG,
McNeill LH, Howard TD, Howard BV, Howard VJ, Johnson KC, Kang SJ, Keating BJ, Kolb S, Kuller LH,
Kutlar A, Langefeld CD, Lettre G, Lohman K, Lotay V, Lyon H, Manson JE, Maixner W, Meng YA,
Monroe KR, Morhason-Bello I, Murphy AB, Mychaleckyj JC, Nadukuru R, Nathanson KL, Nayak U,

N'diaye A, Nemesure B, Wu SY, Leske MC, Neslund-Dudas C, Neuhouser M, Nyante S, Ochs-Balcom H, Ogunniyi A, Ogundiran TO, Ojengbede O, Olopade OI, Palmer JR, Ruiz-Narvaez EA, Palmer ND, Press MF, Rampersaud E, Rasmussen-Torvik LJ, Rodriguez-Gil JL, Salako B, Schadt EE, Schwartz AG, Shriner DA, Siscovick D, Smith SB, Wassertheil-Smoller S, Speliotes EK, Spitz MR, Sucheston L, Taylor H, Tayo BO, Tucker MA, Van Den Berg DJ, Edwards DR, Wang Z, Wiencke JK, Winkler TW, Witte JS, Wrensch M, Wu X, Yang JJ, Levin AM, Young TR, Zakai NA, Cushman M, Zanetti KA, Zhao JH, Zhao W, Zheng Y, Zhou J, Ziegler RG, Zmuda JM, Fernandes JK, Gilkeson GS, Kamen DL, Hunt KJ, Spruill IJ, Ambrosone CB, Ambs S, Arnett DK, Atwood L, Becker DM, Berndt SI, Bernstein L, Blot WJ, Borecki IB, Bottinger EP, Bowden DW, Burke G, Chanock SJ, Cooper RS, Ding J, Duggan D, Evans MK, Fox C, Garvey WT, Bradfield JP, Hakonarson H, Grant SF, Hsing A, Chu L, Hu JJ, Huo D, Ingles SA, John EM, Jordan JM, Kabagambe EK, Kardia SL, Kittles RA, Goodman PJ, Klein EA, Kolonel LN, Le Marchand L, Liu S, McKnight B, Millikan RC, Mosley TH, Padhukasahasram B, Williams LK, Patel SR, Peters U, Pettaway CA, Peyser PA, Psaty BM, Redline S, Rotimi CN, Rybicki BA, Sale MM, Schreiner PJ, Signorello LB, Singleton AB, Stanford JL, Strom SS, Thun MJ, Vitolins M, Zheng W, Moore JH, Williams SM, Ketkar S, Zhu X, Zonderman AB; NABEC Consortium; UKBEC Consortium; BioBank Japan Project; AGEN Consortium; Kooperberg C, Papanicolaou GJ, Henderson BE, Reiner AP, Hirschhorn JN, Loos RJ, North KE, Haiman CA. A meta-analysis identifies new loci associated with body mass index in individuals of African ancestry. Nat Genet. 2013 Jun;45(6):690-6. doi: 10.1038/ng.2608. Epub 2013

Voss U, Sand C, Pfeiffer AF. Heterogeneity of insulin responses: phases leading to type 2 diabetes. PLoS One. 2012;7(9):e44134. doi:10.1371/journal.pone.0044134

Remedi MS, Nichols CG. Hyperinsulinism and diabetes: genetic dissection of beta cell metabolism-expectations and surprises. Annu Rev Physiol. 2009;71:647-662. doi:10.1146/ annurev.physiol.010908.163135

Milenkovic D, Sanz-Moreno A, Calzada-Wack J, Rathkolb B, Veronica Amarie O, Gerlini R, Aguilar-Pimentel A, Misic J, Simard ML, Wolf E, Fuchs H, Gailus-Durner V, de Angelis MH, Larsson NG. Mice lacking the mitochondrial exonuclease MGME1 develop inflammatory kidney disease with glomerular dysfunction. PLoS Genet. 2022 May 9;18(5):e1010190

Robichon C, Varret M, Le Liepvre X, Lasnier F, Hajduch E, Ferré P, Dugail I. DnaJA4 is a SREBP-regulated chaperone involved in the cholesterol biosynthesis pathway. Biochim Biophys Acta. 2006 Sep;1761(9):1107-13. doi: 10.1016/j.bbalip.2006.07.007.

Bai Z, Chai XR, Yoon MJ, Kim HJ, Lo KA, Zhang ZC, et al. Dynamic transcriptome changes during adipose tissue energy expenditure reveal critical roles for long noncoding RNA regulators. PLoS Biol. (2017) 15:e2002176. doi: 10.1371/journal.pbio.2002176

Chimienti F, Devergnas S, Pattou F, Schuit F, Garcia-Cuenca R, Vandewalle B, et al. (2006). In vivo expression and functional characterization of the zinc transporter ZnT8 in glucose-induced insulin secretion. Journal of Cell Science, 119(20), 4199-4206.

Sladek R, Rocheleau G, Rung J, Dina C, Shen L, Serre D, et al. (2007). A genome-wide association study identifies novel risk loci for type 2 diabetes. Nature, 445(7130), 881-885.

Scott LJ, Mohlke KL, Bonnycastle LL, Willer CJ, Li Y, Duren WL, et al. (2007). A genome-wide association study of type 2 diabetes in Finns detects multiple susceptibility variants. Science, 316(5829), 1341-1345.

Mahajan et al. (2021). Refining the accuracy of validated target identification through coding variant fine-mapping in type 2 diabetes. Nature Genetics, 53(10), 1380-1395

Gradwohl G, Dierich A, LeMeur M, Guillemot F. neurogenin3 is required for the development of the four endocrine cell lineages of the pancreas. Proc Natl Acad Sci U S A. 2000 Feb 15;97(4):1607-11

Hrovatin, K., Bastidas-Ponce, A., Bakhti, M. et al. Delineating mouse β-cell identity during lifetime and in diabetes with a single cell atlas. Nat Metab 5, 1615–1637 (2023)

Camunas-Soler J, Dai XQ, Hang Y, Bautista A, Lyon J, Suzuki K, Kim SK, Quake SR, MacDonald PE. Patch-Seq Links Single-Cell Transcriptomes to Human Islet Dysfunction in Diabetes. Cell Metab. 2020 May 5;31(5):1017-1031.e4.

Bakhti, Mostafa, Anika Böttcher, and Heiko Lickert. Modelling the endocrine pancreas in health and disease. Nature Reviews Endocrinology 15.3 (2019): 155-171.

Anna B. Osipovich, Jennifer S. Stancill, Jean-Philippe Cartailler, Karrie D. Dudek, Mark A. Magnuson; Excitotoxicity and Overnutrition Additively Impair Metabolic Function and Identity of Pancreatic β-Cells. *Diabetes* 1 July 2020; 69 (7): 1476–1491

Stancill JS, Cartailler JP, Clayton HW, O'Connor JT, Dickerson MT, Dadi PK, Osipovich AB, Jacobson DA, Magnuson MA. Chronic β-Cell Depolarization Impairs β-Cell Identity by Disrupting a Network of Ca2+-Regulated Genes. Diabetes. 2017 Aug;66(8):2175-2187.

Rudkowska I, Guénard F, Julien P, Couture P, Lemieux S, Barbier O, Calder PC, Minihane AM, Vohl MC. Genome-wide association study of the plasma triglyceride response to an n-3 polyunsaturated fatty acid supplementation. J Lipid Res. 2014 Jul;55(7):1245-53.

Franke A, Hampe J, Rosenstiel P, Becker C, Wagner F, Häsler R, Little RD, Huse K, Ruether A, Balschun T, Wittig M, Elsharawy A, Mayr G, Albrecht M, Prescott NJ, Onnie CM, Fournier H, Keith T, Radelof U, Platzer M, Mathew CG, Stoll M, Krawczak M, Nürnberg P, Schreiber S. Systematic association mapping identifies NELL1 as a novel IBD disease gene. PLoS One. 2007 Aug 8;2(8):e691

Drucker DJ, Nauck MA. The incretin system: glucagon-like peptide-1 receptor agonists and dipeptidyl peptidase-4 inhibitors in type 2 diabetes. Lancet. 2006 Nov 11;368(9548):1696-705.
Ussher, J.R., Drucker, D.J. Glucagon-like peptide 1 receptor agonists: cardiovascular benefits and mechanisms of action. Nat Rev Cardiol 20, 463–474 (2023).

Müller TD, Finan B, Bloom SR, D'Alessio D, Drucker DJ, Flatt PR, Fritsche A, Gribble F, Grill HJ, Habener JF, Holst JJ, Langhans W, Meier JJ, Nauck MA, Perez-Tilve D, Pocai A, Reimann F, Sandoval DA, Schwartz TW, Seeley RJ, Stemmer K, Tang-Christensen M, Woods SC, DiMarchi RD, Tschöp MH. Glucagon-like peptide 1 (GLP-1). Mol Metab. 2019 Dec;30:72-130.

Viana-Huete V, Guillen C, Garcia-Aguilar A, Garcia G, Fernandez S, Kahn CR, et al. Essential role of IGFIR in the onset of Male brown fat thermogenic function: Regulation of glucose homeostasis by differential organ-specific insulin sensitivity. Endocrinology (2016) 157(4):1495–511.

O'Neill BT, Lauritzen HP, Hirshman MF, Smyth G, Goodyear LJ, Kahn CR. Differential Role of
Insulin/IGF-1 Receptor Signaling in Muscle Growth and Glucose Homeostasis. Cell Rep. 2015 May
26;11(8):1220-35.

Nishimura W., Takahashi S., Yasuda K. MafA is critical for maintenance of the mature beta cell
phenotype in mice. Diabetologia. 2015;58:566–574.

Huang Q, Deng G, Wei R, Wang Q, Zou D, Wei J. Comprehensive Identification of Key Genes Involved
in Development of Diabetes Mellitus-Related Atherogenesis Using Weighted Gene Correlation Network
Analysis. Front Cardiovasc Med. 2020 Oct 28;7:580573.

Cromer MK, Choi M, Nelson-Williams C, Fonseca AL, Kunstman JW, Korah RM, Overton JD, Mane S,
Kenney B, Malchoff CD, Stalberg P, Akerström G, Westin G, Hellman P, Carling T, Björklund P, Lifton
RP. Neomorphic effects of recurrent somatic mutations in Yin Yang 1 in insulin-producing adenomas.
Proc Natl Acad Sci U S A. 2015 Mar 31;112(13):4062-7.

Dickinson, M. E., Flenniken, A. M., Ji, X., Teboul, L., Wong, M. D., White, J. K., Meehan, T. F.,
Weninger, W. J., Westerberg, H., & Adissu, H. (2016). High-throughput discovery of novel
developmental phenotypes. Nature, 537(7621), 508-514.

Groza, T., Gomez, F. L., Mashhadi, H. H., Muñoz-Fuentes, V., Gunes, O., Wilson, R., Cacheiro, P.,
Frost, A., Keskivali-Bond, P., & Vardal, B. (2023). The International Mouse Phenotyping Consortium:
comprehensive knockout phenotyping underpinning the study of human disease. Nucleic acids
research, 51(D1), D1038-D1045.

Kang, H. M., Subramaniam, M., Targ, S., Nguyen, M., Maliskova, L., McCarthy, E., Wan, E., Wong, S.,
Byrnes, L., & Lanata, C. M. (2018). Multiplexed droplet single-cell RNA-sequencing using natural
genetic variation. Nature biotechnology, 36(1), 89-94.

Keane, T. M., Goodstadt, L., Danecek, P., White, M. A., Wong, K., Yalcin, B., Heger, A., Agam, A.,
Slater, G., & Goodson, M. (2011). Mouse genomic variation and its effect on phenotypes and gene
regulation. Nature, 477(7364), 289-294.

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biology, 15, 1-21.

Figure 1

A

C57Bl/6J
♂ ♀

Chow diet   High-fat diet

Islet

NZO/ShiLt
♂ ♀

B

Perturbed genes and their causal network.

Figure 2: Identification of DEGs and perturbed genes from β-cells of C57BL/6J mice fed on a normal or high-fat high sugar diet

Figure 3: Identification of DEGs and perturbed genes from β-cells of C57BL/6J mice fed on a normal or high-fat high sugar diet KOMP validation

Figure 4: Identification of DEGs and perturbed genes from β-cells of NZO mice fed on a normal or high-fat high-sugar diet (Males)

Figure 5: KOMP validation from β-cells of NZO mice fed on a normal or high-fat high-sugar diet (Males)

Figure 6: Identification of DEGs and perturbed genes from β-cells of C57BL/6J and NZO mice (males) fed on a high-fat high-sugar diet

Figure 7: Identification of DEGs and perturbed genes from β-cells of C57BL/6J and NZO mice (males) fed on a high-fat high-sugar diet

Figure S0: Workflow of meta-cell ssNPA analysis

Figure S0': An example Abcc8 perturbance case where DEG analysis can't detect

DEG analysis on Abcc8 between meta cell 1, meta cell 2 and meta cell 3, meta cell 4 will have p-value > 0.05

Figure S1: Full Network from β-cells of C57BL/6J mice fed on a normal or high-fat high-sugar diet

DEG

non-DEG

Up-Perturb

Down-Perturb

Perturb non DEG

B6 Male

NZO Male

NZO HF vs B6 HF Male