Title: Classification of psychedelics and psychoactive drugs based on brain-wide imaging of

cellular c-Fos expression

Authors: Farid Aboharb[1,2,#], Pasha A. Davoudian[1,3,4,#], Ling-Xiao Shao[1,5], Clara Liao[1,3], Gillian

N. Rzepka[1], Cassandra Wojtasiewicz[1], Mark Dibbs[5], Jocelyne Rondeau[5], Alexander M.

Sherwood[6], Alfred P. Kaye[5,7,8], Alex C. Kwan[1,9,*]

Affiliations:

[1]Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, 14853, USA

[2]Weill Cornell Medicine/Rockefeller/Sloan-Kettering Tri-Institutional MD/PhD Program, New

York, NY, 10021, USA

[3]Interdepartmental Neuroscience Program, Yale University School of Medicine, New Haven, CT,

06511, USA

[4]Medical Scientist Training Program, Yale University School of Medicine, New Haven, CT,

06511, USA

[5]Department of Psychiatry, Yale University School of Medicine, New Haven, CT, 06511, USA

[6]Usona Institute, Fitchburg, WI, 53711, USA

[7]Clinical Neurosciences Division, VA National Center for PTSD, West Haven, CT, 06477, USA

[8]Wu Tsai Institute, Yale University, New Haven, CT, 06511, USA

[9]Department of Psychiatry, Weill Cornell Medicine, New York, NY, 10065, USA

[#]These authors contributed equally to the work

[*]Correspondence to Alex Kwan, Ph.D., Room 111 Weill Hall, 526 Campus Road, Ithaca, NY,

14853, United States; E-mail: alex.kwan@cornell.edu

Keywords: Psilocybin, ketamine, MDMA, antidepressant, entactogen, drug discovery,

immediate early gene, neural plasticity

ABSTRACT

Psilocybin, ketamine, and MDMA are psychoactive compounds that exert behavioral effects with distinguishable but also overlapping features. The growing interest in using these compounds as therapeutics necessitates preclinical assays that can accurately screen psychedelics and related analogs. We posit that a promising approach may be to measure drug action on markers of neural plasticity in native brain tissues. We therefore developed a pipeline for drug classification using light sheet fluorescence microscopy of immediate early gene expression at cellular resolution followed by machine learning. We tested male and female mice with a panel of drugs, including psilocybin, ketamine, 5-MeO-DMT, 6-fluoro-DET, MDMA, acute fluoxetine, chronic fluoxetine, and vehicle. In one-versus-rest classification, the exact drug was identified with 67% accuracy, significantly above the chance level of 12.5%. In one-versus-one classifications, psilocybin was discriminated from 5-MeO-DMT, ketamine, MDMA, or acute fluoxetine with >95% accuracy. We used Shapley additive explanation to pinpoint the brain regions driving the machine learning predictions. Our results support a novel approach for characterizing and validating psychoactive drugs with psychedelic properties.

INTRODUCTION

Psychedelics include classic serotonergic psychedelics, such as psilocybin and 5-methoxy-*N,N*-dimethyltryptamine (5-MeO-DMT), and related psychoactive compounds, such as ketamine and 3,4-methylenedioxymethamphetamine (MDMA). These compounds have recently gained widespread interest as potential therapeutics for neuropsychiatric disorders[1, 2]. Psilocybin with psychological support is under active investigation as a treatment for major depressive disorder and treatment-resistant depression[3, 4, 5, 6, 7]. Subanesthetic ketamine has long been studied for its efficacy for treating depression[8, 9, 10] and post-traumatic stress disorder (PTSD)[11]. The research efforts culminated in the approval of esketamine nasal spray by the FDA in the United States for treatment-resistant depression[12, 13]. Finally, MDMA-assisted psychotherapy has undergone phase III clinical trials for the treatment of moderate to severe PSTD[14, 15]. The clinical relevance has sparked intense interest in understanding the shared and distinct aspects of these compounds' mechanisms of action.

Beyond the known psychedelics, there is also growing excitement for synthesizing novel psychedelic-inspired analogs that can be new chemical entities for therapeutics[16, 17, 18]. Ideally, the novel compounds would retain therapeutic effects while improving pharmacokinetics, minimizing perceptual effects, and eliminating cardiovascular risks. A major roadblock in this

pursuit lies in developing screens that can filter thousands of psychedelic-inspired analogs to a

manageable list of the most promising compounds for further in-depth characterizations.

Currently, most screens operate at the molecular or behavioral level. At the molecular level,

candidate compounds can be docked *in silico* with the structure of the 5-HT$_{2A}$ receptor, followed

by biochemical measurements of receptor engagement and activation of downstream G-protein

and beta-arrestin pathways. This target-based approach has yielded exciting leads[19, 20, 21, 22], but

assumes that the 5-HT$_{2A}$ receptor is the key mediator of the therapeutic effect, which has not

been proven conclusively. At the behavioral level, candidate compounds may be tested in

animals for defined phenotypes. Simple characterizations such as changes in animal movement

patterns may be automated to increase throughput and accuracy[23, 24]. However, more complex

behavioral assays relevant for depression suffer from limitations including poor construct validity

and weak predictive power for drug efficacy in humans[25].

The development of a new screening method may complement current molecular and

behavioral approaches to accelerate preclinical drug discovery. Classic psychedelics and

ketamine share the ability to enhance neural plasticity in the brain[26], as evidenced by the rapid

and persistent growth of dendritic spines in the rodent medial frontal cortex after a single dose

of ketamine[27, 28], psilocybin[29], and related serotonergic receptor agonists[30, 31, 32, 33]. A promising

approach may thus focus on quantifying indicators of neural plasticity in native brain tissues. To

this end, immediate early genes are activated in a cell in response to increased firing activity or

an external stimulus[34]. The immediate early genes are a key part of neural plasticity, because

they enable neurons to adapt to stimuli by regulating gene expression, which is crucial for

protein synthesis that are needed for synaptic modifications and learning[35, 36]. Taking classic

psychedelics as an example, drug administration induces robust increases in the expression of

immediate early genes[37, 38], including c-Fos, that can be detected starting in as few as 30

minutes in multiple brain regions[39, 40]. More recently, technological advances in tissue clearing,

light sheet fluorescence microscopy, and automated detection of nuclei have enabled high-

throughput mapping of the expression of immediate early genes such as c-Fos in the whole

mouse brain[41, 42]. We and others have applied this method to characterize the impact of

psilocybin and ketamine[43, 44, 45], joining a rapidly growing number of studies using brain-wide

imaging of fluorescence signals to study drugs[46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57]. Although these early

studies have provided valuable biological insights, only one or two drugs were typically included

in each study thus far. Developing the method as a drug screen requires evaluating its feasibility

and accuracy on a larger panel of compounds.

In this study, we measured brain-wide c-Fos expression in male and female mice for 8 drug

conditions, including a variety of psychedelics, related psychoactive compounds, and vehicle

control. We developed a pipeline for analysis and classification based on explainable machine

learning, determining performance in one-versus-rest and one-versus-one classification tasks.

We implemented Shapley additive explanation to interpret the machine learning models to

identify the brain regions driving the classifications. Collectively the results demonstrate brain-

wide imaging of immediate early gene expression as a promising approach for preclinical drug

discovery.

RESULTS

Psychedelics and related drugs in the study

For this study, we evaluated 8 drug conditions: psilocybin (PSI, 1 mg/kg, i.p., single dose),

ketamine (KET, 10 mg/kg, i.p., single dose), *5*-methoxy-*N,N*-dimethyltryptamine (5-MeO-DMT or

5MEO, 20 mg/kg, i.p., single dose), 6-fluoro-*N,N*-diethyltryptamine (6-fluoro-DET or 6-F-DET, 20

mg/kg, i.p., single dose), 3,4-methylenedioxymethamphetamine (MDMA, 7.8 mg/kg, i.p., single

dose), acute fluoxetine (A-SSRI, 10 mg/kg, i.p., single dose), chronic fluoxetine (C-SSRI, 10

mg/kg, i.p., one dose every day for 14 days), and saline vehicle (SAL, 10 mL/kg, i.p., single

dose) (**Fig. 1a**).

Fig. 1. Imaging brain-wide c-Fos expression at cellular resolution following drug administration. a. Chemical structures for the 8 conditions included in this study: psilocybin (PSI), ketamine (KET), 5-MeO-DMT (5MEO), 6-fluoro-DMT (6-F-DET), MDMA, acute fluoxetine (A-SSRI), chronic fluoxetine (C-SSRI, daily for 14 days), and saline vehicle (SAL). **b.** Time course of head-twitch response following the administration of 5-MeO-DMT, psilocybin, 6-fluoro-DET, or saline vehicle. Line, mean. Shading, 95% confidence interval based on 1000 bootstraps. N = 3 males and 3 females for each drug, except 4 males and 3 females for saline. **c.** Box plot of the total number of head twitches detected within a 2-hour period after drug administration. Wilcoxon rank-sum test. *, $P < 0.05$, **, $P < 0.01$. **d.** Experimental timeline. **e.** Box plot of the total number of c-Fos+ cells in the brain for each drug condition. Cross, female individual. Circle, male individual. N = 64 mice, including 4 males and 4 females for each drug. **f.** An example of the fluorescence images of c-Fos+ cells in the mouse brain for a psilocybin-treated mouse acquired by light sheet fluorescence microscopy. Inset, magnified view of the dorsal anterior cingulate cortex. For **b** and **c**, the psilocybin and saline vehicle data had been shown in a prior study[33].

We elected to investigate these compounds for several reasons. Psilocybin is a classic

psychedelic that acts on the 5-HT$_{2A}$ receptor. Psilocybin stands at the forefront of ongoing late-

stage clinical trials evaluating psychedelics' efficacy for treating depression[3, 4, 5, 6, 7]. Ketamine is

primarily a NMDA receptor antagonist[58]. Despite the distinct molecular targets, ketamine and

psilocybin have similarities in their plasticity-promoting action and behavioral effects[59, 60], making

ketamine an intriguing compound to contrast with psilocybin. The doses and route of

administration for psilocybin and ketamine were chosen based on prior studies showing

behavioral effects in mice[29, 61].

5-MeO-DMT is a classic serotonergic psychedelic in the same tryptamine chemical class as

psilocybin[16]. There is clinical interest in evaluating 5-MeO-DMT as a treatment for depression[62,

63]. At a dose of 20 mg/kg in mice, 5-MeO-DMT induces head-twitch response and evokes

structural rewiring in the mouse medial frontal cortex[33]. Compared to psilocybin, 5-MeO-DMT is

shorter-acting and has higher affinity for the 5-HT_{1A} receptor than for the 5-HT_{2A} receptor. Thus

5-MeO-DMT serves as a useful case of another tryptamine psychedelic with distinct

pharmacokinetics and receptor target profile. 6-fluoro-DET is also a tryptamine like psilocybin

and 5-MeO-DMT. Although bioavailable in the brain and a 5-HT_{2A} receptor agonist[64, 65], 6-fluoro-

DET induces autonomic effects without causing perceptual changes in humans[66]. Therefore, it

has been used as an active, non-hallucinogenic control in a clinical study[67]. Concordantly, 6-

fluoro-DET provided ineffective as a substitute compound for rats trained to discriminate LSD or

2,5-dimethoxy-4-iodoamphetamine (known as DOI)[64, 68]. To corroborate these prior results, we

measured the effect of 6-fluoro-DET on head-twitch response in mice using magnetic ear tags

for automated detection of head movements. Our results showed that, unlike 1 mg/kg psilocybin

and 20 mg/kg 5-MeO-DMT which elicited robust head-twitch responses[33], mice administered

with 20 mg/kg 6-fluoro-DET were not statistically different from controls (**Fig. 1b, c**). Our study

adds to other recent studies[20, 21] that included 6-fluoro-DET as a non-hallucinogenic tryptamine

for comparison. The dose of 6-fluoro-DET was chosen to match the dose of 5-MeO-DMT.

MDMA is different from psilocybin: it is a member of the phenethylamine chemical class and has

distinct pro-social and euphoric qualities[69]. MDMA can act on monoamine transporters to

enhance release and inhibit reuptake of neuromodulators including serotonin, thus it has been

characterized as an entactogen rather than a classic psychedelic[70]. MDMA holds clinical

relevance particularly for PTSD[14, 15]. We selected a dose of 7.8 mg/kg for MDMA based on prior

work showing that this dose facilitates fear extinction learning in mice[71]. Fluoxetine is a

commonly prescribed antidepressant that is a selective serotonin reuptake inhibitor (SSRI).

Clinical interest lies in understanding the relative efficacies of SSRIs versus psilocybin[4] and

whether ketamine or psilocybin is suitable for treatment-resistant depression[5, 12, 13]. SSRIs

require chronic administration to exert therapeutic effects, therefore likely engage a mechanism

of action distinct than that of psilocybin and ketamine. For these reasons, we included acute and

chronic fluoxetine for this study. We chose a dose of 10 mg/kg, which was used for acute and

chronic administration of fluoxetine in mice previously[72, 73]. Control animals received a single

injection of saline vehicle.

Light sheet fluorescence imaging of cellular c-Fos expression

For each of the 8 drugs, we tested 4 male and 4 female C57BL/6J mice, totaling 64 animals for

the entire data set. Brains were collected 2 hours after the administration of the single dose or 2

hours after the administration of the last dose for the chronic fluoxetine condition (**Fig. 1d**). The

2-hour interval was chosen assuming drug penetrance to the brain by 0.5 hours and peak c-Fos

expression after an additional 1.5 hours[74]. Brains were processed for tissue clearing and c-Fos

immunohistochemistry (see **Methods**). Light sheet fluorescence microscopy was used to image

each brain at a resolution of 1.8 μm per pixel in the x- and y-axis and at 4 μm intervals in the z-

axis, which allowed for sampling of all cells in the entire brain without any gap. The images were

analyzed using neural nets for automated detection of fluorescent puncta corresponding to c-

Fos+ cells (see **Methods**). The number of c-Fos+ cells detected in each brain for each condition

is presented in **Figure 1e**. An example image collected from a mouse administered with

psilocybin is shown in **Figure 1f**.

To investigate the regional distribution of c-Fos+ cells, we aligned the images of each brain to

the Allen Brain Atlas and segmented the images into summary structures based on the Allen

Mouse Brain Common Coordinate Framework[75] (see **Methods**; **Supplementary Table 1**). The

number of c-Fos+ cells in each brain region for all animals is provided in **Supplementary Table**

2. To visualize the entire data set, we normalized the c-Fos+ cell count in each brain region by

the total number of c-Fos+ cells of each brain and by the spatial volume of the brain region.

Figure 2 is a heatmap of the resulting c-Fos+ cell density for all the samples. We observed that

c-Fos+ cell density was generally high in the isocortex, olfactory area, hippocampal area,

striatum and pallidum, and thalamus, whereas expression was lower in the midbrain and

hindbrain, and cerebellum. The c-Fos expression pattern can be compared to the regional

transcript expression of various receptors (**Fig. S1**). There were individual differences across

samples from the same drug, but also notable contrasts across different drugs. This begets

questions such as: How does the individual variability compare with the differences across

drugs? How well can whole-brain c-Fos maps be used to discriminate the different drugs?

Fig. 2. c-Fos+ cell density listed by brain region for all samples by drugs. The c-Fos+ cell density was defined as the c-Fos+ cell count in each brain region divided by the total number of c-Fos+ cells in each brain and the spatial volume of the brain region. The pixels in the heatmap are positioned by brain region (row) and animal grouped by drug (column). The intensity of the pixel is pseudo-colored by the value of the c-Fos+ cell density. The brain regions including acronyms and other details are provided in **Supplementary Table 1**.

Machine learning pipeline for classifying drugs based on brain-wide c-Fos distribution

To answer these questions, we developed a pipeline for quantitative comparison of the brain-

wide c-Fos expression data between different drug conditions. We posited that different

compounds may elicit distinct regional distribution of cellular c-Fos expression that can serve as

fingerprints for classifying drugs. The pipeline starts with a matrix of c-Fos+ cell counts for

different brain regions from different samples (first panel, **Fig. 3a**). This matrix of c-Fos+ cell

counts undergoes preprocessing, starting with normalization (dividing the c-Fos+ cell count in

each region by the total c-Fos+ cell count of the brain) (second panel, **Fig. 3a**). Normalization is

important because there may be batch effects across samples. The data were then processed

to scale the input data to a standard range such that the values across brain regions are more

comparable and amenable to fitting machine learning models (second panel, **Fig. 3a**), using

Yeo-Johnson transformation (monotonic transformation of data using a power function) and

robust scaling (median subtraction and interquartile range scaling). We will herein refer to the

values after this preprocessing step as the c-Fos scores.

Fig. 3. A machine learning pipeline for drug prediction and performance of one-versus-rest classification.
a. The pipeline consisted of three steps. First, c-Fos+ cell counts for each brain region undergo normalization, Yeo-Johnson transformation, and robust scaling, into c-Fos scores. Second, the Boruta procedure is used to select the set of informative brain regions. Third, c-Fos scores from this set of brain regions were used to fit a ridge logistic regression model. For each iteration, 75% of the data in each drug condition were used for region selection and training through the three steps, and the remaining 25% of the data were withheld initially, but then processed and tested with the ridge logistic regression model. The entire process was iterated using different splits of the data for

Next, we adapted the Boruta feature selection procedure[19] to determine which brain regions to

include for model fitting and testing (third panel, **Fig. 3a**). The Boruta procedure is a

permutation-based method for determining feature importance. It starts by creating "shadow

features": for example, if the data contains 48 c-Fos scores for brain region 1 for various

conditions, then the corresponding shadow feature will be those same 48 c-Fos scores with

scrambled drug labels. Shadow variants were created for all brain regions to create the

expanded Boruta dataset. A random forest classifier was built using this Boruta dataset to

determine a feature-importance value for each brain region. If a brain region has a higher

feature-importance value than the largest feature-importance value from shadow features, then

brain region 1 is a "hit". This permutation process is iterated 100 times. Given that each brain

region can achieve only one of two outcomes (hit or no hit) in each iteration, the distribution of

outcomes across all iterations is a binomial distribution, and a brain region is included by the

statistical criterion of exceeding 95th percentile of the binomial distribution. Why Boruta? We

used the Boruta procedure in lieu of including all brain regions, because many regions likely

contribute little or nothing towards differential drug action and their inclusion in the model would

increase noise and lead to overfitting. A distinctive advantage of Boruta is that brain regions do

not compete with each other, but rather with the shadows. As a result, the number of brain

regions selected by Boruta is not pre-determined but instead dictated by the data as needed.

For the last step, the c-Fos scores from the selected brain regions are used to construct a ridge

logistic regression model (fourth panel, **Fig. 3a**). The entire pipeline is evaluated using 4-fold

splits, where 75% of the data in each drug condition was used to train and fit the model, while

the remaining 25% of the data is used to test the model. Importantly, we emphasize that we

used only the training data to optimize the preprocessing parameters, run feature selection, and

construct regression model. The same optimized preprocessing parameters and selected

features were then later applied for the test data, ensuring no data leakage. The splits were

repeated 100 times to evaluate the prediction accuracy of the pipeline.

One-versus-rest classification shows drug prediction accuracy well above chance

We performed a linear discriminant analysis on the c-Fos scores of all 64 samples, just after the

preprocessing step. We plotted the data for the top two linear discriminants (**Fig. 3b**). This

visualization clearly shows that the differences in c-Fos scores across drugs are more separable

than the differences in c-Fos scores across samples within the same drug condition. Drugs that

alter the serotonergic tone via different mechanisms of action are positioned differently along the

first linear discriminant. By contrast, 5-MeO-DMT, 6-fluoro-DET, and psilocybin are separable

along the second linear discriminant.

We first tested the pipeline with the entire data set and asked the models to predict the exact

drug condition. The confusion matrix shows how the predicted drug labels compared with the

true drug labels (**Fig. 3c**). Because there were 8 conditions, the chance-level accuracy was

12.5% (1 out of 8). We found that the model was the most accurate at identifying the MDMA,

chronic fluoxetine, and 5-MeO-DMT samples, with 100%, 89%, and 81% accuracy respectively.

Performance for other conditions were lower, yielding an overall mean accuracy of 67% for all

drugs. Performance was the lowest for saline and acute fluoxetine at 38% and 47%

respectively. Our interpretation for the low-performance conditions is that tradeoffs must be

made to solve this 8-way classification problem. The machine learning model uses the cross-

entropy loss function, which seeks to maximize the probability of labeling training data correctly

across the entire training set, rather than drawing boundaries in a one-vs-rest fashion. In this

global approach, individual decision boundaries may be placed in a way which under performs

on one label, such as saline, while leading to a greater improvement on others. In other words,

the model was fitted with the goal of maximizing the overall mean classification accuracy, which

was not necessarily the most ideal for distinguishing any one specific condition such as saline.

Nevertheless, the mean accuracy of 67% was still substantially higher than chance level of

12.5%.

Confusion matrices are calculated based on a single decision threshold, which may exaggerate

true positive rate for one drug type at the expense of more false positives for another drug type.

To understand our model performance from a different perspective, we plotted precision-recall

curves (**Fig. 3d**). These curves consider performance across all possible decision thresholds

and summarize the results in terms of precision (true positives relative to false positives) and

recall (true positives relative to false negative). The perfect classifier would have an area under

the precision-recall curve (precision-recall AUC) of 1. Across all drugs, the pipeline yielded a

mean precision-recall AUC value of 0.75. This is well above the theoretical chance-level of

0.125 for 1 out of 8 drugs and the empirical chance-level of 0.12 calculated with shuffled data.

The performance based on precision-recall AUC for predicting different drugs corresponds in

rank order to the accuracy in the confusion matrix. Overall, these results provide evidence that

brain-wide c-Fos maps can be leveraged to identify the exact drug administered out of a panel

of related psychoactive compounds.

A likely use case for the pipeline is to determine how a novel chemical entity may be positioned

in the pharmacological space based on the c-Fos expression pattern. To simulate this scenario,

we performed a leave-one-drug-out analysis, in which we trained a model using 7 conditions

(psilocybin, ketamine, 5-MeO-DMT, MDMA, acute fluoxetine, chronic fluoxetine, and saline), but

then tested it on all conditions including 6-fluoro-DET. We found that 6-fluoro-DET was most

frequently classified as psilocybin at 44% chance but could also be detected as saline at 29%

chance (**Fig. S2**), which is in general agreement with 6-fluoro-DET being a non-hallucinogenic

5-HT$_{2A}$ receptor agonist.

One-versus-one classification suggests a small list of brain regions drives drug

prediction

We reasoned that one-versus-one classification, where the machine learning pipeline solves a

binary problem of deciding between two drugs (**Fig. 4a**), may provide deeper insights into the

factors that distinguish specific drug classes. Given the prominence of psilocybin in clinical trials

and drug discovery, we were particularly interested in comparisons between psilocybin and

other conditions that differ in serotonergic receptor affinities (5-MeO-DMT), mechanism of action

(MDMA, acute fluoxetine, ketamine), or hallucinogenic potency (6-fluoro-DET). We trained the

same machine learning pipeline using subsets of data involving only two or three drugs. The

binary classifiers achieved near-perfect accuracy reflected by precision-recall AUC values at or

exceeding 0.90, with the notable exception of psilocybin versus 6-fluoro-DET which had a

precision-recall AUC of 0.59 (**Fig. 4b**). The difficulty in discerning between a classic

serotonergic psychedelic and the non-hallucinogenic 5-HT$_{2A}$ receptor agonist extended beyond

psilocybin: 5-MeO-DMT versus 6-fluoro-DET as well as psilocybin and 5-MeO-DMT versus 6-

fluoro-DET also yielded modest precision-recall AUC values at 0.80 and 0.57 respectively,

relative to chance level of 0.5 for one-versus-one classifications. These results suggest that

brain-wide cellular c-Fos expression is effective at discriminating between exemplars from

different drug classes, such as a classic psychedelic versus an entactogen, a classic

psychedelic versus a dissociative, and a classic psychedelic versus SSRI. It also effectively

distinguishes between the two classic psychedelics psilocybin and 5-MeO-DMT. However, the

prediction is less reliable for the specific problem of predicting a non-hallucinogenic 5-HT$_{2A}$

receptor agonist relative to a classic psychedelic.

Fig. 4. **Performance of one-versus-one classification. a.** Schematic illustrating the one-versus-one classification problem. **b.** The mean area under the precision-recall curve across all splits for different binary classifiers. Dark gray, real data. Light gray, shuffled data. **c.** The number of brain regions selected via the Boruta procedure for inclusion in the regression model. **d.** Heatmaps showing the fraction of splits when a cortical (left) or thalamic (right) region was included in the regression model. The regions are sorted based on usage in all classifiers. Regions that were included in <75% of the splits across all conditions are not shown.

As mentioned, a feature of the Boruta procedure is that a different number of regions may be

included depending on the data and the desired classification. Indeed, there were differences in

the brain regions chosen for the various drug prediction problems and different training and

testing splits of the same data (**Fig. 4c**). Most classifiers relied on <35 brain regions for drug

prediction, except for the two comparisons involving MDMA which included around 40 - 70 brain

regions. Furthermore, we plotted how often various cortical and thalamic regions were selected

by the machine learning models (**Fig. 4d**). Regions such as retrosplenial areas (RSPd, RSPv),

somatosensory areas (SSp-m, SSp-tr, SSp-ll), and lateral networks (VISC, AId) were included

often, but different classifiers relied on them to different extents. We will explore the importance

of specific brain regions quantitatively in the next section using Shapley additive explanation.

Many thalamic regions were consistently included in comparisons involving MDMA, which

contributed to the higher total number of brain regions used by classifiers when MDMA was

involved. Overall, the results suggest that one-versus-one drug classifications based on brain-

wide c-Fos expression is highly accurate, with the machine learning models only needing data

from a small number of brain regions to produce the prediction.

Using Shapley additive explanation to highlight key brain regions driving drug prediction

A brain region selected by Boruta in the pipeline suggests that it is informative, yet it does not

communicate the importance of its contribution to the final prediction. To better understand how

the c-Fos scores in individual brain regions contribute to decisions in one-versus-one drug

classifications we used Shapley additive explanation (SHAP) (**Fig. 5a**). SHAP uses a game-

theoretical approach to determine how the brain regions contribute to driving the machine

learning regression model from a starting base value to the final output value for decision[21]. To

illustrate, we present the force plot of two test brain samples in one of our cross-validation splits

(**Fig. 5b**). The top half of the plot shows the c-Fos scores in selected brain regions for the

sample of psilocybin and their additive contributions to the decision. In this instance, regions

such as posteromedial visual area (VISpm, c-Fos score = 0.44) and lateral habenula (LH, c-Fos

score = -0.78) were among the drivers leading to an overall positive SHAP value to predict

psilocybin. The posteromedial visual area is located between the primary visual cortex and

retrosplenial cortex[76] and has been suggested to mediate visual information between the

neighboring regions[77]. Lateral habenula neurons had spiking activity associated with

undesirable outcomes[78, 79], which is consistent with their posited role in mediating depression-

related symptoms[80] and contributing to antidepressant response[81]. Intriguingly, another driver

was the parafascicular nucleus (PF, c-Fos score = -1.74), which is implicated in arousal and

head movements[82]. By contrast, the c-Fos scores in the same set of selected brain regions

sums to an overall negative SHAP value for the 5-MeO-DMT sample, providing the basis for the

correct prediction in this case. Across all splits tested for the psilocybin-versus-5-MeO-DMT

comparison, we identified regions that were included in >75% of the machine learning models,

and then ranked these regions by mean SHAP value difference, which highlight the brain

regions most responsible for driving the classification (**Fig. 5c, d**).

Fig. 5. Shapley additive explanation for identifying brain regions driving the prediction of 5-MeO-DMT from psilocybin. a. Diagram illustrating the concept behind SHAP values. The ridge regression model is akin to a black box that takes c-Fos scores as inputs to produce a prediction. SHAP values can be computed to quantitatively assess how strong and in what direction the c-Fos score of each brain region contributes to the prediction. **b**. Example force plots for a psilocybin sample and a 5-MeO-DMT sample from one split, illustrating how actual c-Fos scores of brain regions add to shift the model's output from the base value to the final value. **c**. Plot relating a region's c-Fos scores to the SHAP values across individual splits of the 100 iterations for the 5-MeO-DMT-versus-psilocybin classification. Brain regions were shown only if they were used by >=75% of the splits and listed in rank order by the absolute value of the mean difference in SHAP values between the two drug conditions. The values in parentheses are the absolute value of the mean difference in SHAP values between the two drug conditions. **d**. Visualization of the brain regions included in **c**, color coded according to the compound which evoked higher c-Fos score in the region.

We also analyzed other one-versus-one classification problems using Shapley additive

explanation. For MDMA versus psilocybin, there was a longer list including 32 brains regions

that were used in at least 75% of the cross-validation splits (**Fig. 6a, b**). Half of these regions

(16/32) were in the thalamus. Given the larger number of regions in each model, the SHAP

value differences tended to be smaller, because there is redundancy in the information provided

by the regions.

Fig. 6. Brain regions driving the prediction of MDMA, ketamine, or fluoxetine from psilocybin. a, b. Similar to Fig. 5c, d for MDMA-versus psilocybin classification. **c, d.** Similar to Fig. 5c, d for ketamine-versus psilocybin classification. **e, f.** Similar to Fig. 5c, d for acute fluoxetine-versus psilocybin classification.

For ketamine versus psilocybin, the top 5 regions that were consistently included in >96% of the

cross-validation splits and had the highest SHAP value differences were the visceral area

(VISC), gustatory area (GU), dorsal agranular insular area (AId), xiphoid thalamic nucleus (Xi),

and nucleus of reuniens (RE) (**Fig. 6c, d**). VISC and GU have direct connections to AId, all of

which are part of the lateral subnetworks of the mouse neocortex[83, 84]. The mouse insular cortex

contains various cell types that express an abundance of 5-HT_{2A} and 5-HT_{1A} receptors[85], which

may predispose it to stronger activation by psilocybin. Indeed, the higher c-Fos scores in these

lateral cortical regions informed the model to predict psilocybin. Of note, the insular cortex is

considered a core region in the mouse homolog of the salience network[86, 87], which has been

implicated in mood regulation and depression in humans[88]. Xi and RE are part of the midline

thalamus, which receives visual inputs to mediate behavioral responses to threat[89].

Interestingly, higher c-Fos scores in these midline thalamic regions are routinely used by the
machine learning models to predict ketamine.
373
Finally, we also plotted SHAP value differences comparing acute fluoxetine and psilocybin (**Fig.**
6e, f). Here, the strongest differences were detected by in regions involved in somatosensation
and motor control, including cortical somatosensory regions (SSs, SSp-m), primary motor cortex
(MOp), substantia nigra (SNr), and caudoputamen (CP). These effects may relate to the
previously noted effects of psychedelic on the integration of tactile sensory inputs[90]. Other
implicated regions are the interpeduncular nucleus (IPN) and medial mammillary nucleus (MM),
which are deep midbrain regions that are component of the limbic midbrain circuitry with long-
range connections to habenula, amygdala, and hippocampus.
382
DISCUSSION
In this study, we evaluated the possibility of using whole-brain imaging of cellular c-Fos
expression for drug classification. We developed a machine learning pipeline with key features
including adapting the statistical Boruta procedure to select informative brain regions and using
Shapley additive explanation to identify features that drive the classifications. We tested the
approach using 64 mice that were administered with a panel of psychedelics and related
psychoactive drugs. The results demonstrated high accuracy in various one-versus-rest and
one-versus-one classification problems, supporting the utility of the approach for preclinical drug
discovery. For dissemination, the data and code are available at a public repository.
392
Immunohistochemistry can be influenced by factors such as fixation method, incubation time,
antibody quality, and antigen retrieval techniques. Consequently, the c-Fos antibody staining
can differ from sample to sample. Here, the issue of inter-sample variability was mitigated by not
using the absolute c-Fos+ cell counts for analysis, but instead using the proportional distribution
in each brain region by dividing c-Fos+ cell counts in each region by the total count in each
brain. For instance, if the entire brain was stained poorly and the total c-Fos+ cell count is low,
the proportion distribution should remain unchanged. This normalization step is possible when
whole-brain data is acquired via light sheet fluorescence microscopy. Experimentally, the
variation in antibody staining is also reduced because active electrotransport methods were
used for immunolabeling. Although the normalization step is expected to help with inter-sample
variability, we note that the 64 samples were processed for imaging over 3 batches (details are
provided in **Methods**), and some differences may arise from batch effects.

On average, only a small number of brain regions (~25 brain regions, except for the two

comparisons involving MDMA which included ~50 brain regions) out of the >300 summary

structures in the brain were included in the machine learning models. From our prior study

comparing psilocybin and ketamine[43], we know that both compounds induce increases in c-

Fos+ expression in numerous brain regions including dorsal and ventral anterior cingulate

cortex (ACAd, ACAv), prelimbic area (PL), primary visual cortex (VISp), retrosplenial cortex

(RSP), mediodorsal thalamus (MD), locus coeruleus (LC), lateral habenula (LH), claustrum

(CLA), basolateral amygdala (BLA), and central amygdala (CEA). These brain regions are likely

important for drug action, but shared targets of ketamine and psilocybin are not helpful for

distinguishing the compounds. By design, the machine learning pipeline emphasizes brain

regions with c-Fos expression changes that can discriminate between drug conditions, for which

we found a short list of brain regions.

We anticipate the pipeline to be useful for classifying new chemical entities. For instance, when

a novel psychedelic-inspired compound is synthesized, we may predict its action in the brain by

its position in the linear discriminant axes (**Fig. 3b**) and the proximity to existing drug labels

(**Fig. 3c**). We simulated how such a scenario could work by fitting the pipeline with 7

compounds and testing 6-fluoro-DET as if the classifier has never seen it previously (**Fig. S2**).

For the full panel of drugs tested, we show that the exact drug could be identified with mean

accuracy of 67%, significantly above the chance level of 12.5%. It is instructive to ask how the

pipeline's performance compared with other approaches to classify drugs. For humans,

psilocybin, ketamine, and MDMA exert comparable acute behavioral effects in metrics such as

experience of unity, oceanic boundlessness, and changed meaning of percepts[69]. However,

MDMA preferentially induce blissful state, whereas ketamine evokes disembodiment and

psilocybin induces elementary imagery and audio-visual synesthesia[69, 91]. In one study, human

participants were asked to guess the administered drug, choosing between mescaline (500 mg

and 300 mg), LSD, and psilocybin[92]. The accuracy for identifying the correct drug ranged from

48% to 58% during the session and 69% to 81% after the study. For animals, there has been

recent progress in capturing videos of freely moving mice and analyzing their motion using

unsupervised machine learning methods. One study used motion sequencing method to

investigate a larger panel of 30 psychoactive compounds and doses from a wide range of drug

classes including benzodiazepines, antidepressants, antipsychotics, and stimulants (but not

psychedelics and the compounds tested in the current study) to show a F1 precision-recall

score of 0.62[23]. Our pipeline based on brain-wide cellular c-Fos expression and machine

learning therefore performed at a level comparable to earlier methods based on human and

animal behaviors.

As with any analysis pipeline, there are methodological choices that can affect the outcome,

which can plague the interpretation as demonstrated in the field of neuroimaging[93]. Our

codebase is available online for anyone to freely use, adapt, and test. We used a statistical

method with the Boruta algorithm, rather than a strict threshold, for region selection. We were

careful about data leakage, using only the training data for parameter optimization and feature

selection, such that the prediction accuracy for test data would not be inflated. We implemented

Shapley additive explanation to decipher the factors driving the decisions, which is a general

approach that should find great utility in neuroscience[94], and has already seen applications in

behavioral classification[95] and spike waveform analyses[96]. There are areas of improvement for

the pipeline. While we opted for the simplicity of treating each brain region on its own, regions

may have correlated responses to drug administration. There may be biological reasons, such

as anatomical proximity or synaptic connectivity, for clustering brain regions prior to region

selection, which may outperform our procedure. Network analyses may be used to explore

potential correlated responses to drugs. Furthermore, the pipeline will benefit from testing a

larger range of compounds including enantiomers, other drug classes, and different doses. The

drugs may be administered in conjunction with a receptor antagonist and a stress or behavioral

manipulation, which will all lead to a richer and more refined picture of the 'drug space'. Finally,

c-Fos is one immediate early gene. It is well characterized as an activity-dependent gene and

has the advantage of nuclear labeling that permits automated detection. However, there are

other immediate early genes and plasticity-related biomarkers that can provide complementary

information.

Here we only demonstrated moderate throughput by performing the whole-brain imaging

approach for a sample size of 64 brains. This falls short of other current screening methods,

which typically involve hundreds of conditions including more compounds, different doses, and

additions of antagonists for competitive assays. For whole-brain imaging, the main issue was

cost, which precluded us from testing at a larger scale. At the moment, the drug injection and

tissue extraction steps are straightforward. The cell counting procedure is mostly automated.

However, the cost per brain is high due to tissue processing and imaging, which may drop in the

future because of the rapid advances in brain clearing methods[97] and the development of

inexpensive light sheet fluorescence microscopes[98, 99]. Thus, there is hope that whole-brain

imaging can become a practical method for screening drugs within the next several years.

In summary, there is intense interest in using psychedelics for the treatment of neuropsychiatric

disorders. Progress hinges on knowing more about existing psychedelics and finding new

psychedelic-inspired drugs with improved characteristics. However, there is currently a paucity

of reliable methods to screen psychedelics and related analogs. Here we developed and

characterized an approach based on whole-brain imaging of cellular c-Fos expression. We

demonstrated high prediction accuracy for drug classifications using a machine learning

pipeline. We expect this and other neuroscience-based approaches to play an important role for

accelerating the preclinical development of psychiatric drugs.

Acknowledgments

Psilocybin, 5-MeO-DMT, and 6-fluoro-DET were provided by Usona Institute's Investigational

Drug & Material Supply Program; the Usona Institute IDMSP is supported by Alexander

Sherwood, Robert Kargbo, and Kristi Kaylo in Madison, WI. This work was supported by NIH

grants R01MH121848 (A.C.K.), R01MH128217 (A.C.K.), R01MH137047 (A.C.K.); One Mind –

COMPASS Rising Star Award (A.C.K.); Cornell Engineering M.D.-M.Eng. program (F.A.); NIH

training grants T32GM007205 (P.A.D.), T32NS041228 (C.L.); NIH fellowship F30DA059437

(P.A.D.); Source Research Foundation student grant (P.A.D.); VA National Center for PTSD

(A.P.K.); Department of Defense HT9425-23-1-0458 (A.P.K.); K08MH122733 (A.P.K.); this work

was funded in part by the State of Connecticut, Department of Mental Health and Addiction

Services, but this publication does not express the views of the Department of Mental Health

and Addiction Services or the State of Connecticut. The views and opinions expressed are

those of the authors.

Contributions

F.A., P.A.D., and A.C.K planned the study. P.A.D., L.X.S., and C.L. performed experiments.

G.N.R. and C.W. assisted with tissue processing and imaging. P.A.D. and M.D. measured

head-twitch responses. F.A. and P.A.D. analyzed the data, with input from A.C.K. on the

pipeline. J.R., A.M.S., and A.P.K. contributed reagents. F.A. and A.C.K. drafted the manuscript.

All authors reviewed the manuscript before submission.

Competing interests

A.C.K. has been a scientific advisor or consultant for Boehringer Ingelheim, Empyrean

Neuroscience, Freedom Biosciences, and Psylo. A.C.K. has received research support from

Intra-Cellular Therapies. A.P.K has received research support from Transcend Therapeutics

and Freedom Biosciences. A.P.K. has a provisional patent application related to psychedelics.

The other authors report no financial relationships with commercial interests.

Data availability

Data and code associated with the study will be available on https://github.com/Kwan-Lab.

METHODS

Animals. We used adult, 8-week-old male and female C57BL/6J mice (#00064, The Jackson

Laboratory). Tissues were collected and imaged in three batches. The first batch performed in

August 2021 included 2 males and 2 females for psilocybin (1 mg/kg, i.p.), 2 males and 2

females for ketamine (10 mg/kg, i.p.), and 2 males and 2 females for saline (10 mL/kg, i.p.).

Data from these mice were included in a previous study[12]. The second batch performed in May

2022 included 2 males and 2 females for psilocybin (1 mg/kg, i.p.), 2 males and 2 females for

saline (10 mL/kg, i.p.), 4 males and 4 females for 5-MeO-DMT (20 mg/kg, i.p.), 4 males and 4

females for 6-fluoro-DET (20 mg/kg, i.p.), 4 males and 4 females for acute fluoxetine (10 mg/kg,

i.p.), 4 males and 4 females for chronic fluoxetine (10 mg/kg, i.p.; daily for 14 days). The third

batch performed in December 2022 included 4 males and 4 females for MDMA (7.8 mg/kg, i.p.)

and 2 males and 2 females for ketamine (10 mg/kg, i.p.). All animals were housed and handled

according to protocols approved by the Institutional Animal Care and Use Committee (IACUC)

at Yale University and Cornell University. Tissue collection for all batches was done at Yale

University, except for ketamine in the third batch that was done at Cornell University. For all

batches, the brain samples were shipped for clearing and imaging at LifeCanvas Technologies

(Cambridge, MA).

Drugs. Psilocybin, 5-MeO-DMT succinate, and 6-fluoro-DET solids were obtained from Usona

Institute's Investigational Drug & Material Supply Program. We used the succinate salt form of

5-MeO-DMT[100] (at equivalent amount to freebase 5-MeO-DMT) because it can be dissolved in

saline. Ketamine hydrochloride injection vial (055853, Henry Schein; or Dechra), fluoxetine

hydrochloride solid (F132, Millipore-Sigma), 3,4-MDMA hydrochloride (13971, Cayman

Chemical), and saline (NDC: 0409-4888-03, Hospira) were purchased from supply vendors.

Psilocybin, 5-MeO-DMT succinate, 6-fluoro-DET, MDMA, and fluoxetine were prepared by

dissolving powders into saline. Ketamine was prepared by diluting from the injection vial. For

ketamine, 5-MeO-DMT succinate, 6-fluoro-DET, MDMA, and acute fluoxetine, the working

solutions were prepared fresh on the day of experiment. For psilocybin, a stock solution was

made and then the working solution was made from stock solution, with both solutions prepared

within 1 month from the day of experiment. For chronic fluoxetine, the working solution was

prepared on the first day of administration and then kept in 4°C and used for the remainder of

the chronic treatment.

Tissue collection and imaging. All the samples underwent the same tissue collection and

imaging protocols. Two hours following the single-dose injection or injection of the last dose for

chronic fluoxetine, mice were deeply anesthetized with isoflurane and transcardially perfused

with phosphate buffered saline (P4417, Sigma-Aldrich) followed by paraformaldehyde (PFA, 4%

in PBS). Brains were fixed in 4% PFA for 24 hours at 4°C, after which they were transferred to

0.1% sodium azide in PBS for storage until clearing. The SHIELD protocol was used to process

the whole mouse brains. A stochastic electrotransport device[101] was used to clear samples for 4

days at 42°C, followed by active immunolabeling using eFLASH technology integrating

electrotransport[101] and SWITCH[102]. Each brain sample was stained with 3.5 μg of rabbit anti-c-

Fos monoclonal antibody (Abcam, #ab214672), followed by 10 μg of mouse anti-NeuN

monoclonal antibody (Encor Biotechnology, #MCA-1B7) and then by fluorescently conjugated

secondaries in 1:2 primary:secondary molar ratios (Jackson ImmunoResearch). Following

active labeling, refractive index matching (n = 1.52) was done through incubation in EasyIndex

(LifeCanvas Technologies). Samples were then imaged at 3.6× magnification with a SmartSPIM

light sheet fluorescence microscope (LifeCanvas Technologies) at a resolution of 1.8 μm/pixel

for XY sampling with 4 μm step size for Z sampling over the entire brain. Imaging was done

blinded to treatment conditions.

Atlas registration and cell counting. Fluorescence images were tile-corrected, de-striped, and

registered to the Allen Brain Atlas using an automated process. For each brain, the image from

the NeuN channel was registered to 8-20 atlas-aligned reference samples using

SimpleElastix[103], which implemented successive rigid, affine, and b-spline warping algorithms.

The final atlas alignment value for each sample was determined by taking the average

alignment generated across intermediate reference samples. Cell detection was automated by

using a custom convolutional neural networked designed using the TensorFlow python package.

First, a U-Net-based detection network was used to locate fluorescent puncta corresponding to

c-Fos-immunolabeled cells. Second, a ResNet-based network was used to filter putative cells to

arrive at a final list of cell locations. Each cell location was projected onto the Allen Brain Atlas to

identify its anatomical region. We segmented the brain into 316 summary structures based on

the Allen Mouse Brain Common Coordinate Framework[75]. We omitted the 'fiber tracts' summary

structure in the analysis to focus on grey matter structures. Counts were then generated on a

per-region basis for each sample.

Batch effect correction. We observed differences in the total number of c-Fos+ cells in

psilocybin samples across batch 1 and 2, saline samples across batch 1 and 2, and ketamine

samples across batch 1 and 3. Batch effects are common and, in this study, may arise from

differences in antibody quality, microscope condition, and/or subtle changes in the automated

cell counting procedure. To correct for these differences, a scaling factor was calculated for the

psilocybin, ketamine, and saline conditions individually. This factor was calculated by taking the

mean total c-Fos+ cell counts of the batch 2 (psilocybin, saline) or 3 (ketamine) mice belonging

to the same drug condition and dividing by mean total c-Fos+ cell counts of the batch 1

(psilocybin, saline, ketamine) mice belonging to the same drug condition. The factor was 2.78

for psilocybin, 4.94 for ketamine, and 3.11 for saline. These factors were applied to the per-

region c-Fos+ cell count data in batch 1 to shift the c-Fos+ cell counts to be more comparable to

the later batches. All analyses were performed after the batch effect correction. We emphasize

that this batch correction step should not affect the machine learning analysis pipeline described

below. This is because the first step of the pipeline is to divide per-region count by total count in

each brain, meaning that the absolute values of the cell count should have minimal influence on

model fits but instead it is the relative values of the cell count (e.g., proportion of c-Fos+ cell

residing in one brain region over another brain region in a sample) that mattered for analysis

and prediction.

Head-twitch response. Head movements were recorded using a magnetic ear tag system as

described in detail previously[33]. Briefly, an ear tag consisted of a neodymium magnet (N45, 3

mm diameter, 0.5 mm thick, #D1005-10, SuperMagnetMan) that was adhered to an aluminum

ear tag (La Pias #56780, Stoelting) with cyanoacrylate glue (Super Glue Ultra Gel Control,

$1739050, Loctite). The neodymium magnet was coated with a nitrocellulose marker (#7056,

ColorTone) and dried for >2 h, which helped to reduce ear irritation for the mice. This magnetic

ear tag was applied to the mouse's ear using an ear tag applicator (#56791, Stoelting). For

measurement, the animal was put inside a plastic cube (4" x 4" x 4"). A spool of enameled

cooper wire (30 AWG) was used to wind around the cube like a solenoid, with the ends of the

wire connected to a current-to-voltage preamplifier (PP444, Pyle) where the voltage was

captured with a computer via a data acquisition device (USB-6001, National Instruments). Each

mouse was recorded using one cube. Up to four cubes could be used to record from four mice

at once inside a soundproof chamber. Data acquisition and analysis were done using custom

software written in MATLAB (Mathworks). The voltage signal was sent through a 70 – 110 Hz

bandpass filter because head twitch response had a characteristic ~90 Hz frequency. The

filtered signal was then processed for peak detection to identify individual head-twitch events. A

protocol including parts list for the setup and the MATLAB code is available at https://

github.com/Kwan-Lab/HTR.

Machine learning pipeline – preprocessing. The analysis pipeline used the Python package

sci-kit learn (Version 1.2.1)[104]. The first step of the pipeline was preprocessing, which entails

three steps: normalization, transformation, and scaling. For normalization, we divided each

region's c-Fos+ cell count by the total c-Fos+ cell count across all summary structures used.

This was done to mitigate influence of batch effects across samples. For transformation, each

brain region's normalized c-Fos+ cell counts across different drug conditions were transformed

using Yeo-Johnson power transformation[105]. The Yeo-Johnson transformation is a generalized

form of the Box-Cox transformation. The transformation leads to data values that more closely

approximate a Gaussian distribution. The Yeo-Johnson transformation was implemented in

scikit-learn: *PowerTransformer(method='yeo-johnson', standardize=False)*. The Yeo-Johnson

transformation is parameterized by one variable, lambda. The optimal lambda parameter was

calculated for each brain region independently using maximum likelihood estimation to optimize

for normality. For scaling, for each brain region, the *RobustScaler* module in scikit-learn was

used to subtract the median value and scales values by the range of the 25^{th} to 75^{th} percentile

(quartile scaling). We decided to do this, rather than subtracting mean value and standard-

deviation scaling, because it is less sensitive to outliers. The c-Fos+ cell counts of each brain

region after undergoing the normalization, transformation, and scaling steps are referred to as

the c-Fos scores. To visualize the data, we performed dimensionality reduction on c-Fos scores

across all samples using scikit-learn's *LinearDiscriminantAnalysis* function and plotted the top

two linear discriminants (**Fig. 3b**).

Machine learning pipeline – region selection. Based on Allen Institute definition of summary structures, the brain was divided into 315 regions (316 summary structure and then 'fiber tracts' removed). We were concerned that a model involving c-Fos scores from 315 regions may be overfitting due to our limited sample size of 64 brains. Many regions are likely not informative and only contribute noise to the machine learning models. Therefore, we implemented a method to filter out features (i.e., the brain regions) which were not informative for distinguishing the desired drug conditions. Region selection was carried out using the Boruta algorithm, as implemented in the BorutaPy package[106]. The Boruta algorithm is an 'all relevant features' selection method which seeks to identify all the features with information relevant to a task. This was done by creating scrambled versions of each feature, which are called shadow features, and appending them to the original data set. This expanded data set was then used to fit a random forest classifier, as implemented in scikit-learn. We used the BorutaPy package to automatically select the number of trees for the *RandomForestClassifier()* module based on the size of the feature set. Following this, a threshold was established based on the highest feature importance amongst shadow features. Features exceeding this threshold were considered 'hits' and recorded. This procedure was repeated 100 times. The distribution across these 100 iterations created a binomial distribution. The BorutaPy package rejected features based on the cumulative distribution function of a binomial distribution where p = 0.5, alpha = 0.05, and n = number of hits. Features (i.e., brain regions) that were not rejected by this criterion were the feature included for the next stage of the pipeline.

Machine learning pipeline – classification. We used the c-Fos scores of the selected brain regions to fit a ridge regression model (L2 normalized logistic regression). The regularization parameter C is a hyperparameter used to modulate the penalty strength. Given the interconnected nature of the exact feature set and hyperparameter, as well as our desire to eventually merge results across many cross-validation splits of the data, we opted to fix this parameter to its default value of 1. The 'multinomial' setting was used to generalize from binary classification to multi-class classification.

Cross validation to determine prediction accuracy. The data were evaluated using the aforementioned pipeline using 4-fold splits, where 75% of the data (i.e., 6 brain samples) in each drug condition was used to train and fit the model, while the remaining 25% of the data (i.e., 2 brain samples) was used to test the model. Importantly, preprocessing parameters (e.g., lambda in Yeo-Johnson transformation) and feature selection (brain regions to be included)

were chosen using only the training data to ensure no data leakage. Nevertheless, after those

stages were fixed, the test data would undergo the same preprocessing and feature selection

steps before being inputted into the ridge regression model to generate the prediction of the

drug condition. We performed 100 iterations, each time using a randomized splits for each drug

condition, generated by scikit-learn's *StratifiedShuffleSplit()* function. Combining the outcome

across the 100 iterations, the predicted classifications were used to generate a mean confusion

matrix (**Fig. 3c**). The probabilities assigned to each label for each test data point were combined

to create a composite precision recall curve, generated using scikit-learn's

precision_recall_curve function (**Fig. 3d, Fig. 4b**). The scikit-learn's *auc* function was used to

calculate the area under the curve for each composite precision recall curve (legend of **Fig. 3d**).

We used numpy's random seeds and state objects (numpy.random.RandomState()) to generate

reproducible results. The cross validation splitting function was seeded with an integer, per

scikit-learn's recommendations. Remaining random states were set using a random state

object. A null distribution for area under the precision recall curve was established by shuffling

labels during each cross validation split prior to model fitting and label prediction (**Fig. 4b**).

Shapley additive explanation. SHAP values were generated by the *LinearExplainer* object

from the SHAP package, which accepted test data points and the fit model. We set the feature

perturbation parameter of the *LinearExplainer* to '*correlation_dependent*'. SHAP values were

generated in part by breaking dependencies across features and testing the influence of

perturbations on individual features. This ran the risk of creating unrealistic feature

combinations, because many brain regions which would normally change in lockstep may be

changed individually by the algorithm to infer feature importance, which would lead to inflated

feature importance scores[107]. By using the "correlation dependent" intervention, additional

measures were taken to address correlations in the feature space and credit was distributed

more appropriately. The SHAP values for each test data point were combined across the data

splits from the 100 iterations to arrive at composite SHAP summary plots (**Figs. 5c, 6a, 6c, 6e**).

We determined which brain regions were included in >=75% of the cross-validation splits of the

data (**Figs. 4c, 4d**). Regions meeting this criterion were visualized using the brainrender

package[108] (**Figs. 5d, 6b, 6d, 6f**).

Correlation with gene expression

We calculated the Pearson's correlation coefficient between drug-induced change in c-Fos+ cell

density (by subtracting drug value by saline value and then divide by saline value) and

expression levels for 19,143 genes as gathered from the Allen Brain Institute repository in the

mouse neocortex[109, 110], on a region-by-region basis. These correlation coefficients from all

genes produces a distribution. Genes of interest (e.g., *Htr2a*) were then chosen to be compared

against the distribution to determine a percentile score (*data.rank(pct = True)*). Specifically, we

assessed genes encoding serotonin receptors, glutamate receptors, and neurotransmitter

transporters. We performed this analysis for 4 drugs: psilocybin, 5-MeO-DMT, 6-fluoro-DET, and

MDMA

Leave-one-drug-out analysis

The fitting of the pipeline (*pipelineObj.fit*) was performed on a reduced dataset of cFos scores,

excluding all samples in the 6-fluoro-DET condition. That is, for each split, training data were c-

Fos+ cell count from 75% of the samples from 7 conditions (psilocybin, ketamine, 5-MeO-DMT,

MDMA, acute fluoxetine, chronic fluoxetine, and saline). The test data consist of c-Fos+ cell

count from the remaining 25% of the samples from those 7 conditions and 25% of the samples

drawn from the left-out condition of 6-fluoro-DET. For linear discriminant analysis, the full

dataset was transformed (*pipelineObj.transform*) and plotted using multiple calls to the seaborn

scatterplot function (*sns.scatterplot*).

REFERENCES

1.   Kelmendi B, Kaye AP, Pittenger C, Kwan AC. Psychedelics. *Curr Biol* **32**, R63-R67 (2022).
730

731   2.   Vollenweider FX, Preller KH. Psychedelic drugs: neurobiology and potential for treatment
732   of psychiatric disorders. *Nat Rev Neurosci* **21**, 611-624 (2020).
733

734   3.   Davis AK*, et al.* Effects of Psilocybin-Assisted Therapy on Major Depressive Disorder: A
735   Randomized Clinical Trial. *JAMA Psychiatry* **78**, 481-489 (2021).
736

737   4.   Carhart-Harris R*, et al.* Trial of Psilocybin versus Escitalopram for Depression. *N Engl J*
738   *Med* **384**, 1402-1411 (2021).
739

740   5.   Goodwin GM*, et al.* Single-Dose Psilocybin for a Treatment-Resistant Episode of Major
741   Depression. *N Engl J Med* **387**, 1637-1648 (2022).
742

743   6.   Raison CL*, et al.* Single-Dose Psilocybin Treatment for Major Depressive Disorder: A
744   Randomized Clinical Trial. *JAMA* **330**, 843-853 (2023).
745

746   7.   von Rotz R*, et al.* Single-dose psilocybin-assisted therapy in major depressive disorder: A
747   placebo-controlled, double-blind, randomised clinical trial. *EClinicalMedicine* **56**, 101809
748   (2023).
749

750   8.   Berman RM*, et al.* Antidepressant effects of ketamine in depressed patients. *Biol*
751   *Psychiatry* **47**, 351-354 (2000).
752

753   9.   Murrough JW*, et al.* Antidepressant efficacy of ketamine in treatment-resistant major
754   depression: a two-site randomized controlled trial. *Am J Psychiatry* **170**, 1134-1142
755   (2013).
756

757   10.   Zarate CA, Jr.*, et al.* A randomized trial of an N-methyl-D-aspartate antagonist in
758   treatment-resistant major depression. *Arch Gen Psychiatry* **63**, 856-864 (2006).
759

760   11.   Feder A*, et al.* Efficacy of intravenous ketamine for treatment of chronic posttraumatic
761   stress disorder: a randomized clinical trial. *JAMA Psychiatry* **71**, 681-688 (2014).
762

763   12.   Popova V*, et al.* Efficacy and Safety of Flexibly Dosed Esketamine Nasal Spray Combined
764   With a Newly Initiated Oral Antidepressant in Treatment-Resistant Depression: A
765   Randomized Double-Blind Active-Controlled Study. *Am J Psychiatry* **176**, 428-438 (2019).
766

767   13.   Daly EJ*, et al.* Efficacy and Safety of Intranasal Esketamine Adjunctive to Oral
768   Antidepressant Therapy in Treatment-Resistant Depression: A Randomized Clinical Trial.
769   *JAMA Psychiatry* **75**, 139-148 (2018).
770

771   14.   Mitchell JM*, et al.* MDMA-assisted therapy for severe PTSD: a randomized, double-blind,
772   placebo-controlled phase 3 study. *Nat Med* **27**, 1025-1033 (2021).
773

774   15.   Mitchell JM*, et al.* MDMA-assisted therapy for moderate to severe PTSD: a randomized,
775   placebo-controlled phase 3 trial. *Nat Med* **29**, 2473-2480 (2023).
776

777   16.   Kwan AC, Olson DE, Preller KH, Roth BL. The neural basis of psychedelic action. *Nat*
778   *Neurosci* **25**, 1407-1419 (2022).
779

780   17.   McClure-Begley TD, Roth BL. The promises and perils of psychedelic pharmacology for
781   psychiatry. *Nat Rev Drug Discov* **21**, 463-473 (2022).
782

783   18.   Olson DE. Psychoplastogens: A Promising Class of Plasticity-Promoting
784   Neurotherapeutics. *J Exp Neurosci* **12**, 1179069518800508 (2018).
785

786   19.   Kaplan AL*, et al.* Bespoke library docking for 5-HT2A receptor agonists with
787   antidepressant activity. *Nature*, (2022).
788

789   20.   Wallach J, *et al.* Identification of 5-HT(2A) receptor signaling pathways associated with
790   psychedelic potential. *Nat Commun* **14**, 8221 (2023).
791

792   21.   Dong C, *et al.* Psychedelic-inspired drug discovery using an engineered biosensor. *Cell*
793   **184**, 2779-2792 e2718 (2021).
794

795   22.   Cao D, *et al.* Structure-based discovery of nonhallucinogenic psychedelic analogs.
796   *Science* **375**, 403-411 (2022).
797

798   23.   Wiltschko AB, *et al.* Revealing the structure of pharmacobehavioral space through
799   motion sequencing. *Nat Neurosci* **23**, 1433-1443 (2020).
800

801   24.   Alexandrov V, Brunner D, Hanania T, Leahy E. High-throughput analysis of behavior for
802   drug discovery. *Eur J Pharmacol* **750**, 82-89 (2015).
803

804   25.   Nestler EJ, Hyman SE. Animal models of neuropsychiatric disorders. *Nat Neurosci* **13**,
805   1161-1169 (2010).
806

807   26.   Liao C, Dua AN, Wojtasiewicz C, Liston C, Kwan AC. Structural neural plasticity evoked by
808   rapid-acting antidepressant interventions. *Nat Rev Neurosci*, in press (2024).
809

810   27.   Li N, *et al.* mTOR-dependent synapse formation underlies the rapid antidepressant
811   effects of NMDA antagonists. *Science* **329**, 959-964 (2010).
812

813   28.   Phoumthipphavong V, Barthas F, Hassett S, Kwan AC. Longitudinal Effects of Ketamine on
814   Dendritic Architecture In Vivo in the Mouse Medial Frontal Cortex. *eNeuro* **3**,
815   ENEURO.0133-0115.2016 (2016).
816

817   29.   Shao LX, *et al.* Psilocybin induces rapid and persistent growth of dendritic spines in
818   frontal cortex in vivo. *Neuron* **109**, 2535-2544 e2534 (2021).
819

820   30.   de la Fuente Revenga M, *et al.* Prolonged epigenomic and synaptic plasticity alterations
821   following single exposure to a psychedelic in mice. *Cell Rep* **37**, 109836 (2021).
822

823   31.   Cameron LP, *et al.* A non-hallucinogenic psychedelic analogue with therapeutic
824   potential. *Nature* **589**, 474-479 (2021).
825

826   32.   Lu J, *et al.* An analog of psychedelics restores functional neural circuits disrupted by
827   unpredictable stress. *Mol Psychiatry* **26**, 6237-6252 (2021).
828

829   33.   Jefferson SJ, *et al.* 5-MeO-DMT modifies innate behaviors and promotes structural
830   neural plasticity in mice. *Neuropsychopharmacology*, Online ahead of print (2023).
831

832 34. Sheng M, Greenberg ME. The regulation and function of c-fos and other immediate early
833   genes in the nervous system. *Neuron* **4**, 477-485 (1990).
834

835 35. Yap EL, Greenberg ME. Activity-Regulated Transcription: Bridging the Gap between
836   Neural Activity and Behavior. *Neuron* **100**, 330-348 (2018).
837

838 36. Ma H*, et al.* Excitation–transcription coupling, neuronal gene expression and synaptic
839   plasticity. *Nature Reviews Neuroscience* **24**, 672-692 (2023).
840

841 37. Gonzalez-Maeso J*, et al.* Transcriptome fingerprints distinguish hallucinogenic and
842   nonhallucinogenic 5-hydroxytryptamine 2A receptor agonist effects in mouse
843   somatosensory cortex. *J Neurosci* **23**, 8836-8843 (2003).
844

845 38. Nichols CD, Sanders-Bush E. A single dose of lysergic acid diethylamide influences gene
846   expression patterns within the mammalian brain. *Neuropsychopharmacology* **26**, 634-
847   642 (2002).
848

849 39. Leslie RA, Moorman JM, Coulson A, Grahame-Smith DG. Serotonin2/1C receptor
850   activation causes a localized expression of the immediate-early gene c-fos in rat brain:
851   evidence for involvement of dorsal raphe nucleus projection fibres. *Neuroscience* **53**,
852   457-463 (1993).
853

854 40. Grieco SF*, et al.* Psychedelics and Neural Plasticity: Therapeutic Implications. *J Neurosci*
855   **42**, 8439-8449 (2022).
856

857 41. Renier N*, et al.* Mapping of Brain Activity by Automated Volume Analysis of Immediate
858   Early Genes. *Cell* **165**, 1789-1802 (2016).
859

860 42. Kim Y*, et al.* Mapping social behavior-induced brain activation at cellular resolution in
861   the mouse. *Cell Rep* **10**, 292-305 (2015).
862

863 43. Davoudian PA, Shao LX, Kwan AC. Shared and Distinct Brain Regions Targeted for
864   Immediate Early Gene Expression by Ketamine and Psilocybin. *ACS Chem Neurosci* **14**,
865   468-480 (2023).
866

867 44. Rijsketic DR*, et al.* UNRAVELing the synergistic effects of psilocybin and environment on
868   brain-wide immediate early gene expression in mice. *Neuropsychopharmacology* **48**,
869   1798-1807 (2023).
870

871 45. Datta MS*, et al.* Whole-brain mapping reveals the divergent impact of ketamine on the
872   dopamine system. *Cell Rep* **42**, 113491 (2023).
873

874 46. Bijoch L*, et al.* Whole-brain tracking of cocaine and sugar rewards processing. *Transl*
875   *Psychiatry* **13**, 20 (2023).

47. Kimbrough A, Kallupi M, Smith LC, Simpson S, Collazo A, George O. Characterization of the Brain Functional Architecture of Psychostimulant Withdrawal Using Single-Cell Whole-Brain Imaging. *eNeuro* **8**, (2021).

48. Carrette LLG, Kimbrough A, Davoudian PA, Kwan AC, Collazo A, George O. Hyperconnectivity of Two Separate Long-Range Cholinergic Systems Contributes to the Reorganization of the Brain Functional Connectivity during Nicotine Withdrawal in Male Mice. *eNeuro* **10**, (2023).

49. Azevedo H, Ferreira M, Mascarello A, Osten P, Guimaraes CRW. Brain-wide mapping of c-fos expression in the single prolonged stress model and the effects of pretreatment with ACH-000029 or prazosin. *Neurobiol Stress* **13**, 100226 (2020).

50. Cruces-Solis H, Nissen W, Ferger B, Arban R. Whole-brain signatures of functional connectivity after bidirectional modulation of the dopaminergic system in mice. *Neuropharmacology* **178**, 108246 (2020).

51. Hansen HH, *et al.* Whole-brain activation signatures of weight-lowering drugs. *Mol Metab* **47**, 101171 (2021).

52. Keyes PC, *et al.* Orchestrating Opiate-Associated Memories in Thalamic Circuits. *Neuron* **107**, 1113-1123 e1114 (2020).

53. Roland AV, *et al.* Alcohol Dependence Modifies Brain Networks Activated During Withdrawal and Reaccess: A c-Fos-Based Analysis in Mice. *Biol Psychiatry* **94**, 393-404 (2023).

54. Skovbjerg G, *et al.* Whole-brain mapping of amylin-induced neuronal activity in receptor activity-modifying protein 1/3 knockout mice. *Eur J Neurosci*, (2021).

55. Skovbjerg G, *et al.* Uncovering CNS access of lipidated exendin-4 analogues by quantitative whole-brain 3D light sheet imaging. *Neuropharmacology* **238**, 109637 (2023).

56. Stefaniuk M, *et al.* Global brain c-Fos profiling reveals major functional brain networks rearrangements after alcohol reexposure. *Neurobiol Dis* **178**, 106006 (2023).

57. Tan B, Browne CJ, Nöbauer T, Vaziri A, Friedman JM, Nestler EJ. Drugs of abuse hijack a mesolimbic pathway that processes homeostatic need. *Science* **384**, (2024).

58. Zanos P, *et al.* Ketamine and Ketamine Metabolite Pharmacology: Insights into Therapeutic Mechanisms. *Pharmacol Rev* **70**, 621-660 (2018).

920 59. Savalia NK, Shao LX, Kwan AC. A Dendrite-Focused Framework for Understanding the
921   Actions of Ketamine and Psychedelics. *Trends Neurosci* **44**, 260-275 (2021).
922

923 60. Aleksandrova LR, Phillips AG. Neuroplasticity as a convergent mechanism of ketamine
924   and classical psychedelics. *Trends Pharmacol Sci* **42**, 929-942 (2021).
925

926 61. Ali F*, et al.* Ketamine disinhibits dendrites and enhances calcium signals in prefrontal
927   dendritic spines. *Nat Commun* **11**, 72 (2020).
928

929 62. Reckweg JT*, et al.* The clinical pharmacology and potential therapeutic applications of 5-
930   methoxy-N,N-dimethyltryptamine (5-MeO-DMT). *J Neurochem* **162**, 128-146 (2022).
931

932 63. Davis AK, So S, Lancelotta R, Barsuglia JP, Griffiths RR. 5-methoxy-N,N-
933   dimethyltryptamine (5-MeO-DMT) used in a naturalistic group setting is associated with
934   unintended improvements in depression and anxiety. *Am J Drug Alcohol Abuse* **45**, 161-
935   169 (2019).
936

937 64. Blair JB*, et al.* Effect of ring fluorination on the pharmacology of hallucinogenic
938   tryptamines. *J Med Chem* **43**, 4701-4710 (2000).
939

940 65. Rabin RA, Regina M, Doat M, Winter JC. 5-HT2A receptor-stimulated phosphoinositide
941   hydrolysis in the stimulus effects of hallucinogens. *Pharmacol Biochem Behav* **72**, 29-37
942   (2002).
943

944 66. Kalir A, Szara S. Synthesis and Pharmacological Activity of Fluorinated Tryptamine
945   Derivatives. *J Med Chem* **6**, 716-719 (1963).
946

947 67. Faillace LA, Vourlekis A, Szara S. Clinical evaluation of some hallucinogenic tryptamine
948   derivatives. *J Nerv Ment Dis* **145**, 306-313 (1967).
949

950 68. Helsley S, Fiorella D, Rabin RA, Winter JC. A comparison of N,N-dimethyltryptamine,
951   harmaline, and selected congeners in rats trained with LSD as a discriminative stimulus.
952   *Prog Neuropsychopharmacol Biol Psychiatry* **22**, 649-663 (1998).
953

954 69. Studerus E, Gamma A, Kometer M, Vollenweider FX. Prediction of psilocybin response in
955   healthy volunteers. *PLoS One* **7**, e30800 (2012).
956

957 70. Nichols DE. Differences between the Mechanism of Action of Mdma, Mbdb, and the
958   Classic Hallucinogens - Identification of a New Therapeutic Class - Entactogens. *J
959   Psychoactive Drugs* **18**, 305-313 (1986).
960

961 71. Young MB, Andero R, Ressler KJ, Howell LL. 3,4-Methylenedioxymethamphetamine
962   facilitates fear extinction learning. *Transl Psychiatry* **5**, e634 (2015).
963

964 72. Dulawa SC, Holick KA, Gundersen B, Hen R. Effects of chronic fluoxetine in animal models
965   of anxiety and depression. *Neuropsychopharmacology* **29**, 1321-1330 (2004).
966

967 73. Simard S*, et al.* Fibroblast growth factor 2 is necessary for the antidepressant effects of
968   fluoxetine. *PLoS One* **13**, e0204980 (2018).
969

970 74. Morgan JI, Cohen DR, Hempstead JL, Curran T. Mapping patterns of c-fos expression in
971   the central nervous system after seizure. *Science* **237**, 192-197 (1987).
972

973 75. Wang Q*, et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference
974   Atlas. *Cell* **181**, 936-953 e920 (2020).
975

976 76. Wang Q, Burkhalter A. Area map of mouse visual cortex. *J Comp Neurol* **502**, 339-357
977   (2007).
978

979 77. Roth MM, Helmchen F, Kampa BM. Distinct functional properties of primary and
980   posteromedial visual area of mouse neocortex. *J Neurosci* **32**, 9716-9726 (2012).
981

982 78. Matsumoto M, Hikosaka O. Lateral habenula as a source of negative reward signals in
983   dopamine neurons. *Nature* **447**, 1111-1115 (2007).
984

985 79. Matsumoto M, Hikosaka O. Representation of negative motivational value in the primate
986   lateral habenula. *Nat Neurosci* **12**, 77-84 (2009).
987

988 80. Li B*, et al.* Synaptic potentiation onto habenula neurons in the learned helplessness
989   model of depression. *Nature* **470**, 535-539 (2011).
990

991 81. Yang Y*, et al.* Ketamine blocks bursting in the lateral habenula to rapidly relieve
992   depression. *Nature* **554**, 317-322 (2018).
993

994 82. Fallon IP*, et al.* The role of the parafascicular thalamic nucleus in action initiation and
995   steering. *Curr Biol* **33**, 2941-2951 e2944 (2023).
996

997 83. Zingg B*, et al.* Neural networks of the mouse neocortex. *Cell* **156**, 1096-1111 (2014).
998

999 84. Harris JA*, et al.* Hierarchical organization of cortical and thalamic connectivity. *Nature*
1000   **575**, 195-202 (2019).
1001

1002 85. Ju A, Fernandez-Arroyo B, Wu Y, Jacky D, Beyeler A. Expression of serotonin 1A and 2A
1003   receptors in molecular- and projection-defined neurons of the mouse insular cortex.
1004   *Molecular Brain* **13**, (2020).
1005

1006 86. Gozzi A, Schwarz AJ. Large-scale functional connectivity networks in the rodent brain.
1007   *NeuroImage* **127**, 496-509 (2016).

87. Mandino F, *et al.* A triple-network organization for the mouse brain. *Molecular Psychiatry* **27**, 865-872 (2021).

88. Lynch CJ, *et al.* Frontostriatal salience network expansion in individuals in depression. *Nature*, (2024).

89. Salay LD, Ishiko N, Huberman AD. A midline thalamic circuit determines reactions to visual threat. *Nature* **557**, 183-189 (2018).

90. Duerler P, *et al.* Psilocybin Induces Aberrant Prediction Error Processing of Tactile Mismatch Responses—A Simultaneous EEG–FMRI Study. *Cerebral Cortex* **32**, 186-196 (2022).

91. Vollenweider FX, Kometer M. The neurobiology of psychedelic drugs: implications for the treatment of mood disorders. *Nat Rev Neurosci* **11**, 642-651 (2010).

92. Ley L, *et al.* Comparative acute effects of mescaline, lysergic acid diethylamide, and psilocybin in a randomized, double-blind, placebo-controlled cross-over study in healthy participants. *Neuropsychopharmacology* **48**, 1659-1667 (2023).

93. Botvinik-Nezer R, *et al.* Variability in the analysis of a single neuroimaging dataset by many teams. *Nature* **582**, 84-88 (2020).

94. Goodwin NL, Nilsson SRO, Choong JJ, Golden SA. Toward the explainability, transparency, and universality of machine learning for behavioral classification in neuroscience. *Curr Opin Neurobiol* **73**, 102544 (2022).

95. Goodwin NL, *et al.* Simple Behavioral Analysis (SimBA) as a platform for explainable machine learning in behavioral neuroscience. *Nature Neuroscience* **27**, 1411-1424 (2024).

96. Lee EK, *et al.* Non-linear dimensionality reduction on extracellular waveforms reveals cell type diversity in premotor cortex. *eLife* **10**, (2021).

97. Lai HM, *et al.* Antibody stabilization for thermally accelerated deep immunostaining. *Nature Methods* **19**, 1137-1146 (2022).

98. Vladimirov N, *et al.* Benchtop mesoSPIM: a next-generation open-source light-sheet microscope for cleared samples. *Nature Communications* **15**, (2024).

99. Chen Y, *et al.* Low-cost and scalable projected light-sheet microscopy for the high-resolution imaging of cleared tissue and living samples. *Nature Biomedical Engineering* **8**, 1109-1123 (2024).

1052

100.   Sherwood AM, Claveau R, Lancelotta R, Kaylo KW, Lenoch K. Synthesis and Characterization of 5-MeO-DMT Succinate for Clinical Use. *ACS Omega* **5**, 32067-32075 (2020).

101.   Kim SY*, et al.* Stochastic electrotransport selectively enhances the transport of highly electromobile molecules. *Proc Natl Acad Sci U S A* **112**, E6274-6283 (2015).

102.   Murray E*, et al.* Simple, Scalable Proteomic Imaging for High-Dimensional Profiling of Intact Systems. *Cell* **163**, 1500-1514 (2015).

103.   Marstal K, Berendsen F, Staring M, Klein S. SimpleElastix: A User-Friendly, Multi-lingual Library for Medical Image Registration. In: *2016 IEEE Conference on Computer Vision and Pattern Recognition Workshops (CVPRW)*) (2016).

104.   Pedregosa F*, et al.* Scikit-learn: Machine Learning in Python. *JMLR* **12**, 2825-2830 (2011).

105.   Yeo IK, Johnson RA. A new family of power transformations to improve normality or symmetry. *Biometrika* **87**, 954-959 (2000).

106.   Kursa MB, Rudnicki WR. Feature Selection with the Boruta Package. *Journal of Statistical Software* **36**, 1-13 (2010).

107.   Mase M, Owen AB, Seiler B. Explaining black box decisions by Shapley cohort refinement. *arxiv*, (2020).

108.   Claudi F, Tyson AL, Petrucco L, Margrie TW, Portugues R, Branco T. Visualizing anatomically registered data with brainrender. *Elife* **10**, (2021).

109.   Lein ES*, et al.* Genome-wide atlas of gene expression in the adult mouse brain. *Nature* **445**, 168-176 (2007).

110.   Fulcher BD, Murray JD, Zerbi V, Wang XJ. Multimodal gradients across mouse cortex. *Proc Natl Acad Sci U S A* **116**, 4689-4695 (2019).