Comparison of conservation strategies for California Channel Island Oak (*Quercus tomentella*) using climate suitability predicted from genomic data

Alayna Mead[1,2], Sorel Fitz-Gibbon[1], John Knapp[3], Victoria Sork[1,4]

[1] Department of Ecology and Evolutionary Biology, University of California, Los Angeles
[2] Department of Ecosystem Science and Management, Pennsylvania State University
[3] The Nature Conservancy, California Chapter
[4] Institute of the Environment and Sustainability, University of California, Los Angeles

Abstract

Management strategies such as assisted gene flow could increase resilience to climate change, but knowledge of the evolutionary history and genetic structure of species are needed to determine the risks and benefits of different strategies. *Quercus tomentella*, or Island Oak, is a rare oak restricted to six of the California Islands in California, USA and Baja California, Mexico. Previous work has shown that Island Oaks on each island are genetically differentiated from each other, but less is known about the extent to which island populations are adapted to their local climate conditions. We performed whole-genome sequencing on Island Oak individuals, as well as *Q. chrysolepis*, a closely related species that hybridizes with Island Oak (127 trees in total), to characterize genetic structure and introgression across its range and assess the relationship between genomic variation and climate. We assessed three potential management strategies, with different trade-offs between conserving historic genetic structure and enabling populations to survive changing climates: the *status quo* approach, ecosystem preservation approach which conserves the trees and their associated biodiversity, and species preservation approach which conserves the genetic variation of the species. We compare the impact of the three approaches on predicted maladaptation to climate using Gradient Forest. We also introduce a climate suitability index to identify optimal pairs of seed sources and planting sites for approaches involving assisted gene flow. We found one island (Santa Rosa) that could benefit from the ecosystem preservation approach and also serve as a species preservation site. Overall, we find that both the ecosystem preservation and species preservation approaches performed better than the *status quo* approach. The species preservation approach would produce slightly better-adapted populations to predicted climate change, but without the benefits offered by the ecosystem preservation approach. Choosing between these three approaches will depend on the conservation goals.

Introduction

Climate change may be proceeding too quickly for many populations to adapt to new
conditions, particularly long-lived and foundational species such as forest trees, threatening
not only species but the entire ecosystem (Aitken et al., 2008). However, tree species often
contain abundant genetic variation that may enable them to adapt to novel climates, which
could be a tool for management and conservation. Populations are frequently adapted to local
environments that vary across the species range (Leimu & Fischer, 2008; Sexton et al., 2014;
Sork, 2017; Wang & Bradburd, 2014), and this heterogeneity can enable adaptation to novel
environments. In natural populations, gene flow can introduce novel genetic variation into a
population, including alleles that are adaptive under particular environments (Savolainen et al.,
2007). If these alleles are adaptive in the new environment and selection is strong enough, they
will increase in frequency, enabling the population to be better adapted to environmental
conditions. If adaptation is too slow to match the pace of climate change, as may be the case
for species with long generation times, one potential management strategy is to intentionally
introduce genetic variation that is likely to be adaptive under future conditions. This strategy,
assisted gene flow (AGF), involves transferring genetic material (such as seeds or individuals)
among populations within a species range (Aitken & Bemmels, 2016; Aitken & Whitlock, 2013;
Browne et al., 2019).

Genomic information can help determine the risks and benefits of AGF for a particular species.
The risks of AGF may include outbreeding depression, disruption of local adaptation, and the
introduction of alleles that are deleterious in the new location (Grummer et al., 2022). A
species' genome-wide genetic structure can be used to understand historic patterns of gene
flow among populations and evaluate whether these risks are likely, and putatively adaptive
genetic variation can provide information about specific AGF strategies such as seed sourcing
recommendations (Shryock et al., 2021; Yu et al., 2022). By considering both types of genomic
information, it is possible to compare the predicted outcomes of management strategies such as
assisted gene flow and make management decisions aligning with conservation goals.

Landscape genomics analyses can be used to understand patterns of putatively adaptive genetic variation and identify conservation strategies. Landscape genomics identifies genetic variation associated with landscape characteristics, such as climate and geography, and can determine which factors are most important in explaining genetic variation (Fitzpatrick & Keller, 2015; Rellstab et al., 2015; Sork et al., 2013). These genomic-climate relationships can be used to calculate the genomic offset, a metric of the difference between two climates (at a different location, time period, or both) while scaling climate variables by their importance in explaining the spatial patterns of putatively adaptive alleles and taking into account non-linear relationships with allele frequencies (Fitzpatrick & Keller, 2015). The genomic offset can be calculated for a given site under historic and future climates to predict which parts of a species range are at greatest risk of maladaptation, and it can also be calculated using historic climates and future climates across a species range to identify optimal pairs of seeds sources and planting sites (Gougherty et al., 2021; Lachmuth et al., 2024).

Forest trees have life history characteristics that make them likely to be good candidates for AGF: they often experience high gene flow followed by strong selection on seedlings, enabling local adaptation, similar to the processes that would result from AGF. High levels of gene flow are facilitated by their predominantly outcrossing mating system and long-distance pollen dispersal by wind (Petit & Hampe, 2006). Despite the resulting lack of strong genetic structure, local adaptation is common within forest trees, likely due to the high genetic diversity of a large panmictic species combined with high seedling mortality introducing strong selection against non-locally adapted alleles (Savolainen et al., 2007, 2013; Sork, 2016, 2017; Sork et al., 2013). Strong selection pressure against deleterious alleles, whether introduced through natural or human-assisted gene flow, should reduce the chances that they will be introduced into the gene pool.

Endangered species with restricted ranges and fragmented populations are often the focus of conservation efforts to maintain or restore populations, particularly under changing climates. Yet, it is unclear whether such tree species experience evolutionary processes similar to widespread tree species, making AGF a viable strategy, or whether populations should be

conserved *in situ* without AGF. Generally, landscape genomic studies on trees have found that

genetic differences among populations increase with geographic distance (a pattern of isolation

by distance, or IBD), but that environmental variables also play a role in determining genetic

variation across the range, finding candidate genetic variants associated with climate

(Fitzpatrick & Keller, 2015; Gugger et al., 2021; Jia et al., 2020; Martins et al., 2018). In

comparison to widespread tree species, rare species occurring on islands may be comprised of

disjunct populations with limited gene flow (Di Santo et al., 2022; Gugger et al., 2018), and

island endemic plant species may sometimes have lower genetic diversity than mainland

congeners (Hamabata et al., 2019), resulting in a decreased ability to adapt to changing climatic

conditions. However, in many species, island populations have greater diversity than mainland

populations (García-Verdugo et al., 2015) and some studies on island species have found signs

of local adaptation. In Hawaiian koa, precipitation was strongly associated with genetic

variation both across and within islands (Gugger et al., 2018). In Torrey pine, the disjunct

island and mainland populations show morphological differences in a common garden that are

consistent with adaptation to the contrasting climates (Hamilton et al., 2017). Fine-scale local

adaptation has also been described in bird species occupying islands (Cheek et al., 2022;

Gamboa et al., 2022; Langin et al., 2015). These studies show that local adaptation may be

possible in species with fragmented ranges. Given the potential vulnerability of species with

disjunct populations under climate change, AGF could be a viable conservation strategy if

populations contain sufficient adaptive genetic variation.

Quercus tomentella, or Island Oak, is a rare relictual island tree which is endemic to six of the

California Islands off the western coast of the United States (the Channel Islands) and Mexico

(Isla Guadalupe) (Figure 1A). This species is categorized as endangered by the IUCN (Beckman

& Jerome, 2017), and its threats include browsing from non-native herbivores, erosion, changes

in hydrology, and climate change (Beckman et al., 2019; Beckman & Jerome, 2017); and on

Santa Rosa Island, the loss of a native disperser, the Island Scrub-Jay (Delaney & Wayne, 2005;

Morrison et al., 2011). Island Oak forms groves that co-occur with island endemic plant species

and varieties, forming unique ecosystems that support animal species such as the Island Scrub-

Jay (Pesendorfer et al., 2018; Sawyer et al., 2009). Based on microsatellite markers, *Q. tomentella*

is genetically differentiated across islands and has lower genetic diversity than its more

widespread relative *Q. chrysolepis* (Ashley et al., 2018). Additionally, it can propagate clonally,

which may contribute to its lower genetic diversity. However, to understand how selection has

shaped adaptive genetic variation within and across islands, we used whole genome sequence

data, which will often reveal different spatial patterns than the neutral markers ofcalifornia

microsatellites (Fitzpatrick & Keller, 2015; Gugger et al., 2021; Martins et al., 2018), providing a

better understanding of adaptive potential. While Island Oak currently has a restricted and

fragmented range, it was once widespread throughout mainland California during the Miocene

and Pliocene (Figure 1A), then became restricted to the coast, and eventually the California

Islands as the mainland climate became less temperate (Axelrod, 1967; Muller, 1965). During

glacial periods with lower sea levels, the Northern Channel Islands were connected as a single

island, called Santa Rosae, as recently as 11,000 years ago (Kennett et al., 2008; Reeder-Myers et

al., 2015). During this time, there may have been more gene flow among Santa Rosae Island

populations. It is also possible that gene flow occurred through translocation by native peoples,

as acorns are an important food source, although no direct evidence exists (Rick et al., 2019).

Hybridization may be a source of genetic variation: *Q. tomentella* hybridizes with *Q.*

chrysolepis, a widespread species on mainland California that is also present on five of the

islands (Ashley et al., 2018), and genomic and fossil data support a history of co-occurrence and

ancient introgression between the two species (Axelrod, 1944b, 1944a; Ortego et al., 2018).

In this study, we used a landscape genomic analysis of whole-genome sequencing data

collected from the entire species range of Island Oak to assess the best strategy for preserving

the it and its associated ecosystems under future climate change scenarios on the California

Islands (Figure 1 B, C). Before assessing different management approaches, we need to first

characterize the processes structuring genetic variation in *Q. tomentella*, such as gene flow,

hybridization, and local adaptation. We documented the genetic structure of *Q. tomentella*

along with *Q. chrysolepis* and their hybrids to describe the historic patterns of divergence and

gene flow. Then, we implemented a landscape genomics analysis to identify associations

between genetic variation and climate and identify putatively adaptive SNPs. We compared three management strategies using the genomic offset as a proxy for maladaptation. We also calculated the climate suitability index, a new comparison metric based on the genomic offset that measures the similarity of climates scaled by their genetic importance. Two of the strategies considered here involve the use of assisted gene flow (AGF), in which oak seeds or genotypes are transplanted other locations within the species range (Aitken & Bemmels, 2016; Aitken & Whitlock, 2013). While a combination of ecological and restoration management strategies will likely be needed to conserve this species, here we focus on the potential contribution of AGF to their success.

[1] *Status quo* **approach**: The simplest approach, and the one currently used, is to protect existing populations and augment oak groves by planting acorns from the same island or watershed. This approach maintains the historic genetic population structure, but may not enable populations to survive climate change.

[2] **Ecosystem preservation approach**: To maintain existing oak groves and re-establish previous ones so that Island Oak ecosystems and their associated biodiversity are sustained, restoration projects could use AGF to create a viable oak population under future climates by introducing genotypes that are likely to be best-adapted to future climate conditions at a planting location.

[3] **Species preservation approach**: When a species is threatened throughout its native range, it is sometimes necessary to focus on strategies that will preserve the species, either *in situ* or *ex situ. In situ* sites could include existing oak groves, and/or they could be sites on the islands where they are not present, but have historically existed. Alternatively, species preservation could be accomplished through *ex situ* planting sites on the mainland (Rosenberger et al., 2022; Westwood et al., 2021). As *ex situ* preservation typically cultivates trees in botanical gardens and arboreta, targeting sites with optimal climates is less important than for wild populations, so we focus here on *in situ* preservation of Island Oaks in their native range.

Figure 1. A) Map showing location of the six islands sampled, relation to the mainland, and direction of the California Current. Locations of fossil sites of *Q. declinata*, the fossil equivalent of *Q. tomentella*, from the Miocene (Axelrod, 1939) and Pliocene (Axelrod, 1944b) are marked on mainland California. Isla Cedros to the southeast is shown where *Q. cedroensis* occurs. Map credit: Irina Koroleva, The Nature Conservancy. B and C) Predicted shifts in climate at each sampled location based on max temperature of the warmest month (BIO5) and precipitation of the warmest quarter (BIO12) for two future climate models, CNRM (B) and MIROC (C) under Representative Concentration Pathway (RCP) 4.5, in which emissions peak in 2040 then decline. Closed circles indicate historic climate for a given location, and arrowheads indicate the projected future climate for the same location.

Methods

Collections

Leaf samples from *Q. tomentella* individuals were collected across the six California Islands encompassing the species range. Because *Q. tomentella* hybridizes with *Q. chrysolepis*, we also collected co-occurring *Q. chrysolepis* and putative hybrid individuals from the islands, and *Q. chrysolepis* individuals from across northern and southern mainland California (southernmost in San Bernardino county, northernmost in Siskiyou county, Table S1). We attempted to avoid collecting multiple samples from the same clone by collecting from only one stem within the same grove or cluster of stems, with the exception of Anacapa Island where only one grove of trees exists. Leaf samples were either dried on silica or were placed on ice while transported to the Sork lab, then were frozen at -80° C. In total, based on species identifications in the field, we sequenced and analyzed data for 107 *Q. tomentella* individuals, 17 *Q. chrysolepis* individuals (8 from mainland California and 9 from the islands), and 3 putative hybrids.

DNA extraction and sequencing

Leaves were hand-ground on liquid nitrogen using a mortar and pestle, and approximately 50 mg of tissue was used for DNA extraction. DNA was extracted from leaves using a modified version of the Qiagen DNeasy Plant Mini Kit protocol. First, to remove polyphenols, a prewash step was performed twice. 1 mL of prewash buffer was added to ground leaf tissue, ground in a bead mill for 20 seconds, centrifuged for 10 minutes at 10,000 RPM, and the supernatant was discarded. Prewash buffer consisted of (per sample) 100 ul Tris, 100 ul EDTA, 200 ul 5 M NaCl, 600 ul molecular grade water, and 0.01 g PVP. Following the prewash, the Qiagen protocol was followed. For the silica-dried leaf samples (from Guadalupe and San Clemente Islands) we were unable to extract sufficient amounts of DNA, so extraction was performed by the CCGP mini-core using the Macherey-Nagel NucleoMag Plant kit, with the following modifications: PVP and Proteinase K were added during the digest, and an additional ethanol wash was added before elution. Extracted DNA was sent to UC Davis DNA Technologies and Expression

Analysis Cores for library preparation using a custom SeqWell kit. Whole-genome sequencing

was performed on a NovaSeq 6000 using 150 bp, paired-end sequencing.

Filtering and variant calling

Adapters were trimmed from raw reads using Trim Galore, and reads with a length less than 20

bp were removed. Reads were not trimmed based on quality scores during this step. Reads were

aligned to the *Q. lobata* genome (Sork et al., 2022), rather than the Island Oak genome (Mead et

al., 2024), because the genetic similarity was high (85.8% identity, Mead et al., 2024) and

annotations were available. Alignment used BWA-MEM, with 'markShorterSplits' and

'readGroupHeaderLine' options enabled. Duplicate reads were marked and removed using

GATK MarkDuplicates. Variants were called using GATK HaplotypeCaller with the 'emit-ref-

confidence' option set to 'GVCF'. Variants were hard-filtered using GATK VariantFiltration,

with SNPs and indels filtered separately. For SNPs, we removed variants with quality by depth

(QD) <2, quality (QUAL) <30, mapping quality (MQ) < 40, phred-scaled strand bias (FS) >60,

symmetric odds ratio strand bias (SOR) >3, mapping quality rank sum (MQRankSum) < -12.5,

and read position rank sum (ReadPosRankSum) < -8. We removed indels with QD<2, FS>200,

QUAL<30, and ReadPosRankSum< -20. Repetitive regions of the genome were removed using

vcftools based on the reference genome.

From this set of high-quality variants, we selected biallelic SNPs with high coverage across all

samples for further analysis. Using bcftools (version 1.15.1, Danecek et al., 2021), we selected

only biallelic SNPs, and removed SNPs with a mean depth across all samples <5, set individual

genotypes with depth <5 to missing, removed SNPs with a minor allele frequency <0.01, and

removed SNPs with ≥90% missingness across all individuals. The resulting filtered VCF file was

converted to BED file format using PLINK (version 1.90b6.26, Chang et al., 2015), and variants

in linkage disequilibrium (LD) were pruned using a window size of 50 variants, a window shift

value of 10 variants, and an R^2 threshold of 0.1, which identifies variant pairs with a correlation

>0.1 within the given window and prunes them until no correlated pairs remain. Unless

otherwise noted, analyses were run on this filtered and LD-pruned dataset of 585,298 SNPs. For

analyses requiring no missing data, SNPs were imputed by assigning missing individuals the most common SNP (total missingness in the filtered dataset was 5%).

Climate data

Climate variables were extracted for each locality from WorldClim (version 2.1, historic climate data for 1970-2000) at 30-second resolution. Of the five Channel Islands with Island Oaks present, the Northern Channel Islands (Santa Rosa, Santa Cruz, and Anacapa) are cooler and receive more precipitation than the Southern Channel Islands (Santa Catalina and San Clemente, Figure 1B and C). Precipitation seasonality varies across an east/west gradient in the Northern Channel Islands, with the western part of the range experiencing less precipitation seasonality and more summer rainfall. Guadalupe Island, off the coast of Baja California, Mexico, has warmer winter temperatures and less precipitation, but has less yearly variation in temperature and precipitation, and receives more precipitation during the dry season than the Channel Islands. We also extracted future predictions of climate at our sample sites using downscaled CMIP5 climate models from WorldClim, projected to 2050 under two RCP scenarios: 4.5 (with emissions peaking around 2040) and 8.5 (rising emissions through the 21st century, IPCC 2014). We choose two climate models with different outcomes for Southern California, CNRM-CM5 (a warmer/wetter scenario) and MIROC-ESM (a hotter/drier scenario) (Underwood et al., 2019).

Genetic Structure

Divergence among islands was calculated using Weir and Cockerham's F_{ST} in the R package heirfstat 0.5-11 (Goudet & Jombart, 2022). To perform a principal components analysis (PCA) on the imputed SNP set, we used the R package vegan (version 2.6-2, Oksanen et al., 2019). Because Guadalupe Island trees were divergent from the other populations (see results), we also ran a PCA with those samples excluded to better visualize the genetic structure among the remaining islands. ADMIXTURE (version 1.3.0, Alexander et al. 2009) was used to estimate the ancestry of individuals and characterize genetic structure across the species range. For the

entire dataset, we ran ADMIXTURE for K values 1-10, where K is the number of hypothetical

ancestral populations. We also ran ADMIXTURE on subsets of the data to investigate fine-scale

genetic structure for the following sets: all samples except Guadalupe Island (K=1-10) and

individuals from Santa Cruz and Santa Rosa islands alone (K=1-5). Because Guadalupe Island

populations appear deeply diverged from the other Island Oak populations and are unlikely to

experience contemporary gene flow, we ran further analyses of adaptive genetic variation

separately for the five northernmost islands (hereafter, Channel Islands) and for Guadalupe

Island.

To assess the effects of geography and climate on genetic variation in the Channel Islands

populations, we used a partial redundancy analysis (RDA) in the vegan package, which

partitioned the genetic variance into proportions that could be statistically explained by

climate, by geographical location (via latitude and longitude), by each of these factors alone

while controlling for the other, and by their joint influence. We tested the significance of each

explanatory factor using the `anova.cca` function with 99 permutations.

A redundancy analysis was also used to identify candidate SNPs that are associated with

climate variables. We reduced the climate and environmental variables to a subset with

pairwise correlations below 0.75: BIO5 (Max Temperature of Warmest Month), BIO6 (Min

Temperature of Coldest Month), BIO15 (Precipitation Seasonality), BIO18 (Precipitation of

Warmest Quarter), BIO19 (Precipitation of Coldest Quarter), and elevation. A preliminary

analysis found that candidate SNPs were strongly influenced by the five individuals from

Anacapa, which are from one small grove and had high genetic similarity to each other. Many

of the resulting candidate SNPs were associated with BIO5, which is lowest at the Anacapa site.

In further analysis with Gradient Forest (see below), including the Anacapa samples appeared

to result in overfitting of the model, with unsampled regions that had similar climate to

Anacapa clustering separately from all other regions. Since these trees are relatively isolated, it

is difficult to determine whether this association is the result of natural selection on

temperature or due to inbreeding within this population, so we selected the candidate SNPs

using an RDA on the four larger Channel Islands (Santa Rosa, Santa Cruz, Catalina, and San

Clemente), all of which had samples collected in multiple regions within the island. We

replicated the analysis to identify candidate SNPs on Guadalupe Island, which is genetically

diverged from the other islands but has relatively high genetic diversity and climatic variation,

potentially allowing for local adaptation within the island. Following Forester et al. (2018), we

identified candidate SNPs as those that were strongly associated with the multivariate climate

space. We considered candidate SNPs to be those that were outliers (more than 4 standard

deviations) on the first three PC axes. We did not correct for population structure because the

partial RDA showed a minor influence of geography on genetic variation after controlling for

climate (see Results, Table 2), and because correcting for population structure in an RDA

results in reduced power and increased false positive rates in systems with low population

structure (Forester et al., 2018).

Predicting climate suitability

Gradient Forest (GF, version 0.1-32, Ellis et al., 2012) was used to identify nonlinear

relationships between SNPs and environmental factors, and to predict turnover of genomic

composition across the range of Island Oak. We ran GF on three sets of the imputed SNP data:

1) all SNPs to predict background genomic turnover for the Channel Islands, 2) candidate SNPs

identified from the redundancy analysis for the Channel Islands to predict putatively adaptive

turnover for the Channel Islands, and 3) candidate SNPs identified from Guadalupe Island to

predict genomic turnover within Guadalupe Island and across the entire species range.

Geographic location cannot be included as a predictor in GF analysis, so PCNM axes were

calculated as a proxy for geography using the `pcnm` function in the vegan package. To

calculate the PCNM axes, we set the truncation distance separately for datasets because they

encompassed different areas. For the two analyses on the Channel Islands, a truncation

distance of 150 km was used in calculating PCNM axes to maintain connections between

populations on the Northern and Southern Channel Islands and avoid statistical artifacts

resulting from our clustered sampling design. As recommended by Fitzpatrick & Keller (2015),

we included the first half of the axes with positive eigenvectors (denoting a positive spatial

correlation) in the analysis, in this case 5 PCNM axes. For Guadalupe, we used a truncation

distance of 430 km to account for the larger geographic area and retained 3 PCNM axes. We

also included the same climate variables that were used in the redundancy analysis as

explanatory variables. The `gradientForest` command was run with the following

parameters: number of trees = 500, correlation threshold = 0.5, and max level = $\log_2$(0.368 ×

number of samples/2), following Fitzpatrick & Keller (2015).

We predicted the genetic composition and turnover across the range using the relationships

between climate and allele frequency (in this case, a value of 0, 1, or 2 indicating the number of

copies of a variant present in an individual tree) to scale climate variables by their genetic

importance and predict genomic composition at non-sampled locations from their climates.

The first three predicted axes were mapped to RGB color values which were combined for each

cell and mapped, as in Fitzpatrick & Keller (2015). We did this color assignment for the three

sets of SNPs. For the Channel Islands, we also mapped the difference in genetic turnover

between the genome-wide SNP dataset and the candidate SNPs using the Procrustes residuals.

Using the Gradient Forest predictions of genomic turnover, we calculated the difference

between current and future climates using the genomic-scaled climates variables, or the

genomic offset. We calculated the genomic offset for each population under our three

conservation scenarios: 1) the *status quo* approach to conservation, in which populations are

maintained at their present locations, 2) ecosystem preservation, in which non-local, pre-

adapted genotypes are transplanted to a site, and 3) species preservation, in which genotypes

are conserved by transplanting them to optimal sites. These are similar to the local, reverse,

and forward offset (respectively) calculated in Gougherty et al. (2021), but here we explicitly

test different conservation scenarios for each population. For scenario 3, we calculated offset

on two sets of locations: only the sites where trees were sampled in this study, and all possible

sites on the islands. Because offset is predicted from gridded climate data, these "populations"

are not defined biologically, but are groups of trees within the same climate grid cell that are

likely to have similar adaptive genetic variation based on our gradient forest model. We also

converted offset into a relative climate suitability by dividing it by the maximum offset

calculated in the model, so that a value of 1 represents climate that is identical to climate of

origin, and 0 represents the least suitable site.

Results

Genetic Structure

F_{ST} values among island pairs and mainland *Q. chrysolepis* ranged from near 0 to 0.05 (Table 1).

Within the island populations, the population from Guadalupe Island was the most divergent

from other populations (average pairwise F_{ST} = 0.032), followed by Santa Rosa and Anacapa

(average pairwise F_{ST} = 0.021 and 0.020, respectively). In fact, the average divergence between

Guadalupe Island populations and the Channel Islands populations was similar to that of the

divergence between mainland *Q. chrysolepis* and the island populations primarily consisting of

Q. tomentella. PCA and ADMIXTURE plots also supported divergence of Guadalupe

populations from other *Q. tomentella* samples in California (Figures 2 and 3).

Table 1. Pairwise F_{ST} (Weir and Cockerham's) across all island samples and mainland *Q. chrysolepis* samples.

	Mainland (Qchr)	Santa Rosa	Santa Cruz	Anacapa	Santa Catalina	San Clemente	Guadalupe	Avg	Avg, mainland excluded
Mainland (Qchr)	-	0.049	0.026	0.044	0.012	0.02	0.028	0.030	
Santa Rosa	0.049	-	0.009	0.023	0.019	0.011	0.042	0.026	0.021
Santa Cruz	0.026	0.009	-	0.01	0.009	0.001	0.027	0.014	0.011
Anacapa	0.044	0.023	0.01	-	0.011	0.013	0.045	0.024	0.020
Santa Catalina	0.012	0.019	0.009	0.011	-	0.003	0.026	0.013	0.014
San Clemente	0.02	0.011	0.001	0.013	0.003	-	0.021	0.012	0.010
Guadalupe	0.028	0.042	0.027	0.045	0.026	0.021	-	0.032	0.032

Figure 2. PCAs of genetic variation across SNPs for axes 1-3 for all samples (top) and with Guadalupe Island excluded (bottom). Color of points indicates their collection location, and shape indicates the species as identified in the field (although collectors often noted they were unsure of the species identity).

Figure 3. ADMIXTURE results when the number of ancestral groups (K) was set to values 2-5. Each color indicates a group, and each bar shows the ancestry proportions of each group for an individual tree. Colored bars at the bottom indicate the field species ID (note that collectors on each island were different, and collectors often commented that they were unsure of IDs).

We found evidence of widespread introgression between the two species. Mainland *Q.*
chrysolepis samples clustered separately from the island individuals in the PCA, but island
individuals that were identified in the field as *Q. chrysolepis* or possible hybrids primarily
cluster within the island samples rather than with mainland *Q. chrysolepis* (Figure 2).
ADMIXTURE plots show that *Q. chrysolepis* or hybrid individuals on the Northern Channel
Islands have mixed ancestry, with portions from mainland *Q. chrysolepis* and from the northern
Q. tomentella groups (Figure 3). Trees from the Southern Channel Islands had more *Q.*
chrysolepis ancestry than those from the Northern Channel Islands (Figure 3). Because these
results suggest a history of introgression between the two species on the islands rather than
occasional hybridization events producing F_1 offspring, and because hybridization can be a
source of genetic variation for natural selection (Suarez-Gonzalez et al., 2018), we decided to
include the hybrid individuals and those that morphologically appeared to be *Q. chrysolepis*
from the islands in further analyses of putatively adaptive genetic variation.

ADMIXTURE results for the full dataset, for the dataset with Guadalupe sample excluded, and
for the dataset with only the Northern Channel Islands (Santa Cruz and Santa Rosa) all had the
lowest cross-validation error for K=1 (Figure S1), meaning that in the best model, all
individuals come from a single ancestral population. We plotted admixture proportions for
multiple K values to visualize genetic structure (Figure 3). The plot for K = 2 separates
mainland *Q. chrysolepis* and Guadalupe Island as one ancestry group and the Northern Channel
Islands as another, with admixture between the two groups occurring in hybrid individuals and
in the Southern Channel Islands. For K = 3, Guadalupe Island forms a separate ancestry group.
Across multiple K values, trees from the Southern Channel Islands were more admixed, with
ancestry proportions from both mainland *Q. chrysolepis* and the Northern Channel Island
populations. The plot for K=5 suggests that the gene flow San Clemente primarily receives
from the Northern Channel Islands comes from one region of Santa Cruz, located on the
southwestern part of the island (purple group in Figure 3). Running ADMIXTURE on a subset
dataset including only two of the Northern Channel Islands, Santa Rosa and Santa Cruz,
revealed fine-scale genetic structure. For example, one tree on the southwestern ridge of Santa

Cruz shared some ancestry with Santa Rosa samples, suggesting that gene flow has occurred

from Santa Rosa to Santa Cruz populations (Figure S2).

Adaptive genetic variation

Multiple redundancy analysis models were tested in order to partition genetic variance into

proportions explained by climate, geography, their joint influence, and each factor alone

(controlling for the effects of the other factor). All models significantly explained genetic

variation ($P = 0.01$ for all), indicating that both geography and climate are factors shaping

genetic variation (Table 2). However, climate explained a greater proportion of variance than

geography: in the full model, geography and climate together explained 3.8% of the genetic

variation; with climate alone explaining 2.3%, geography alone explaining 0.9%, and their joint

influence explaining 0.6%. As a proportion of total explainable variance, climate alone

explained 61.6%, geography alone explained 22.5%, and their joint influence explained 15.9%.

The redundancy analysis identified 560 candidate climate-associated SNPs that were outliers in

the Channel Islands dataset and 346 in the Guadalupe dataset. For each candidate SNP, we

determined which climate variable was most strongly correlated. For the Channel Islands,

there were 174 SNPs correlated most strongly with BIO19, 155 with elevation, 88 with BIO5, 73

with BIO18, 47 with BIO15, and 23 with BIO6. For Guadalupe, 204 SNPs were most strongly

correlated with BIO6, 129 with BIO5, 9 with elevation, 2 with BIO15, and 2 with BIO19.

Table 2. Partitioning of genetic variance among Channel Island samples among climate and geography (latitude and longitude) when controlling for the other, as well as their joint influence that cannot be disentangled. First column includes the total amount of variation explained, and second column is as a proportion of variance explained by the full model (0.038). *P*-values were calculated from 99 permutations, except for the joint influence, which is calculated from the total variance explained that cannot be partitioned by climate and geography alone and is not testable using permutations.

Model	Adjusted R^2	Proportion of explainable variance	*P*
full model (climate + geography)	0.038	1	0.01 **
Climate alone	0.023	0.616	0.01 **
Geography alone	0.009	0.225	0.01 **
Joint influence of climate and geography	0.006	0.159	

When Gradient Forest was used to analyze genome-wide SNPs in the Channel Islands samples,

the five PCNM axes describing geographic distance had greater importance values than the

climate variables (Figure S3), suggesting a spatial influence on genetic variation. However, for

the candidate SNPs, important variables included both PCNM axes and environmental

variables (with PCNM3, BIO6, and elevation having the highest R^2 weighted importance values

for the Channel Islands; and BIO5, BIO19, and elevation having the highest values for

Guadalupe) (Figure S3). As expected, overall importance values were also higher when

considering only the candidate SNPs, indicating stronger associations. The relationship

between allelic variation and climate, or genomic turnover, was mapped to visualize predicted

spatial patterns of climate-associated genetic variation for the Channel Islands (Figure 4) and

for Guadalupe (Figure 5). The Southern Channel Islands were more similar to each other, and

the Northern Channel Islands showed variation across an east-west gradient associated with

precipitation seasonality (BIO15) and precipitation of the warmest quarter (BIO18). When

predicting turnover using candidate SNPs from identified from Guadalupe, there was more

turnover within the island than across the Channel Islands (Figure 5), consistent with the

genetic differentiation of Island Oaks on Guadalupe from the other islands, as well as

putatively adaptive genetic differentiation within the island.

Genomic offsets and conservation strategies

Catalina Island has the greatest genomic offset under 2050 climate (CNRM model, RCP 4.5)
than the other islands, followed by Anacapa and eastern Santa Cruz (Figure 4). These results
were consistent across the two climate models and the two RCP scenarios tested (Figure S9).
Within Guadalupe, populations closer to the coast have a higher offset (Figure 5). Offset values
calculated for the two datasets (Channel Islands and Guadalupe) are not comparable to each
other, so we cannot determine whether the risk for Guadalupe is higher or lower than for the
Channel Islands.

Both conservation strategies implementing AGF reduced the genomic offset compared to the
status quo (Figure 6). For most sampled sites, the genomic offset was reduced most with the
species preservation approach compared to the ecosystem preservation approach. Offsets were
slightly reduced when all parts of the islands were included as possible planting sites, rather
than only the sites with sampled oak populations (Figure 6), indicating that some future
optimal sites may be outside of the range sampled in this study. Optimal planting sites were
similar for all populations and included sites that will be coolest under future climates: Santa
Rosa followed by San Clemente (although under CNRM 8.5, San Clemente has higher
suitability than Santa Rosa – Supplemental Table S1). Because Santa Rosa populations are
already located in the part of the species range that will have future climate conditions that are
most similar to current conditions for the species, they benefit more from the ecosystem
preservation approach than the species preservation approach, unlike other populations.

Figure 4. A) Maps of genetic turnover predicted by gradient forest for candidate SNPs on each of the

Channel Islands. Climate variables are scaled by their importance in predicting genetic variation in the gradient forest model and their relationship with allelic turnover as shown in Figure S4, then the genomic composition of a cell is predicted based on its climate. The first three PCs of the genomic composition are mapped to RGB values, so regions with similar colors are expected to have similar genomic composition. Points indicate sample locations. B) PCA of scaled climate values for each cell, with colors matching each grid cell in A. C) Map of the genetic offset, or predicted difference in the current genetic composition and that which would be ideal under future climate conditions. Darker red indicates a greater likelihood of future maladaptation. Results are shown for the CNRM 4.5 scenario for the year 2050 (offsets under other models and scenarios are in Figure S9).

Figure 5. A) Maps of genetic turnover predicted by gradient forest for candidate SNPs on Guadalupe Island. Catalina Island is included as a comparison, but all other Channel Islands were similar, clustering in the dark

blue/purple regions with lower BIO5 (max temperature of the warmest month). Climate variables are scaled by their importance in predicting genetic variation in the gradient forest model, then the genomic composition of a cell is predicted based on its climate. The first three PCs of the genomic composition are mapped to RGB values, so regions with similar colors are expected to have similar genomic composition. Points indicate sample locations. B) Map of the genetic offset, or predicted difference in the current genetic composition and that which would be ideal under future climate conditions. Darker red indicates a greater likelihood of future maladaptation. Results are shown for the CNRM 4.5 scenario for the year 2050; offsets under other models and scenarios are shown in Figure S10. Colors only indicate offset within Guadalupe and are not comparable to offset in other islands in Figure 4. C) PCA of scaled climate values for each cell, with colors matching each grid cell in A.

Figure 6. Comparison of genomic offsets under four management scenarios for the Channel Islands (A) and Guadalupe Island (B). Higher offsets represent greater maladaptation to future climate. Offsets were calculated for each grid cell that included sampled trees, referred to as "populations," and are sorted by island. The *status quo* strategy indicates offsets for each population at its current site. In the ecosystem preservation strategy, the offset is given for a scenario in which the site is planted with the non-local population predicted to be best-adapted to its 2050 conditions. In the species preservation strategy, the offset is given for the population on the island if it were transplanted to another site with the most optimal 2050 conditions, either when including the sampled sites or all sites on the islands. For A and B, offset values are calculated separately, with transplantation only within the Channel Islands or Guadalupe Island, and the offsets are not comparable between the two plots.

Figure 7. Climate suitability in 2050 when matching a sampled source site (columns) to a planting site (rows). Suitability was calculated for each grid cell where trees were collected for this study, and nearby grid cells with similar values were averaged for clarity. Planting sites (rows) are sorted by average suitability, with the optimal sites at the top. Columns are sorted geographically (roughly north to south and west to east). Under the ecosystem preservation approach, optimal source populations for restoring a given site can be selected by looking at climate suitability within a row. Under the species preservation approach, optimal planting sites for a given population can be selected by looking at climate suitability within a column.

Discussion

Our genomic analyses reveal great potential for using seed sources to manage, restore, and expand Island Oak ecosystems and to preserve the species, despite the fact that currently all localities are predicted to be maladapted under future climates. These findings point to the importance of using assisted gene flow for populations on many of the islands to maximize their potential for survival under future climate environments. In this study, we use the combined evidence from landscape genomics and genome-wide genetic structure to compare the potential benefits of several conservation strategies.

Genetic structure

Information about a species' genetic structure can be used to inform seed sourcing decisions, balancing the goals of climate adaptation and preserving historic genetic structure and gene flow patterns. Our results show genetic differentiation across islands, but F_{ST} values are relatively low for an island species (Gugger et al., 2018; Villa-Machío et al., 2020, but see Di Santo et al., 2022), indicating weak population structure among islands (Table 1). Guadalupe Island populations, which are at a geographic distance from the Channel Islands populations that make pollen transfer extremely unlikely, are also the most genetically divergent from other island populations (average pairwise F_{ST} = 0.032, Table 1). On the Channel Islands, Santa Rosa populations appear to be the most genetically distinct, while other islands are less differentiated from each other and show signs of admixture (Table 1, Figure 3). The patterns of genetic structure found here may result from asymmetric gene flow shaped by typical wind patterns during spring flowering (Figure 1A). In the Channel Islands, winds are primarily from the northwest, so trees on Santa Rosa Island are unlikely to receive pollen from other islands, except under rare spring Santa Ana wind conditions, which normally occur in Fall when oaks are not producing pollen. However, Santa Cruz and Anacapa may receive pollen from Santa Rosa, and San Clemente and Catalina may receive pollen from all three northern islands, explaining the increased admixture observed in these individuals (Figure 3). These results suggest that the Northern Channel Islands are unlikely to receive alleles from the warmer

Southern islands, reducing the pool of alleles that could be adaptive under warmer future

climates. Similar to our results, previously analyzed microsatellite data found isolation of

Guadalupe and greater admixture in the Catalina and San Clemente populations (Ashley et al.,

2018), but our ADMIXTURE results show more distinct ancestries between Santa Rosa and the

islands to its east, possibly because whole-genome datasets detect the impact of adaptative

variation more than do neutral markers. Patterns of differentiation among islands were also

similar to our results, but are not directly comparable because different markers and

differentiation statistics were used (Ashley et al., 2018).

Introgression with *Q. chrysolepis* appears to be widespread, except on Santa Rosa and Anacapa,

and is likely to be another important factor shaping genetic structure in Island Oak. Admixture

between the two species is particularly evident in the Southern Channel Islands, but also

present within some individuals in the Northern Channel Islands. Island trees that were

identified as *Q. chrysolepis* based on field morphology clustered more closely with *Q. tomentella*

individuals than with the mainland *Q. chrysolepis* individuals, suggesting a long history of

introgression between the two species on these islands (Figure 2). Our genetic results suggest

that *Q. chrysolepis* may not be present on Santa Rosa or Anacapa, where they have been

previously reported. The isolated Guadalupe Island population, however, appears to be

somewhat genetically similar to the mainland *Q. chrysolepis;* having similar values along PC1

(Figure 2) and forming the same ancestry group for K=2 (Figure 3). Previous work has hinted at

a complex evolutionary history in these two species. Ortego et al. (2018) suggested that *Q.*

chrysolepis is a non-monophyletic species split into a northern and southern lineage, and that

Q. tomentella split from the southern lineage of *Q. chrysolepis* in the late Pliocene or early

Pleistocene. If *Q. tomentella* originated in southwest North America, Guadalupe Island may

have been colonized early in the species' evolutionary history, before its range moved

northward. Colonization of the Channel Islands may have occurred later, as California shifted

toward a more Mediterranean climate and *Q. tomentella* became restricted to the coast

(Axelrod, 1967; Muller, 1965). Additionally, it is possible that the Guadalupe population has

experienced introgression with *Q. cedrocensis*, which we were unable to sample but which is

present on Cedros Island to the southeast of Guadalupe Island (Figure 1A), as well as on the

mainland in southern San Diego County and northern Baja California. In the future,

demographic modeling that includes the Guadalupe Island populations could be used to clarify

the timing of divergence between Guadalupe and the other islands, and the timing of

introgression with *Q. chrysolepis* or other oak species in section *Protobalanus.* While

hybridization is sometimes considered a conservation concern, here it seems to be a part of a

long evolutionary history of introgression between the two species, as is common in oaks

(Kremer & Hipp, 2020; O'Donnell et al., 2021), and could be the source of novel genetic

variation.

Adaptive genetic variation

To test for local adaptation to climate in Island Oak populations, we used a redundancy

analysis to partition genetic variation into proportion explained by climate and geography

separately. Climate accounted for most of the explainable genetic variation among the Channel

Island populations when controlling for the effects of geography (Table 2). Thus, natural

selection is likely affecting the evolution of these island populations more than neutral

differences resulting from genetic drift and limited gene flow within or among islands. Climate

variables that were important in the redundancy analysis and Gradient Forest analysis varied,

suggesting that elevation, temperature, and precipitation variables are important in local

adaptation, without a strong influence of any one factor, unlike some tree species for which

genetic turnover is most associated with precipitation variables (Gugger et al., 2018; Martins et

al., 2018). The moderating impact of ocean temperatures and the fog on all of the islands may

make these variable indistinguishable.

Our findings indicate that populations within the same island may be locally adapted to

different conditions. On Santa Cruz Island, climate-associated genetic variation differed across

an east/west gradient (Figure 4A). Candidate SNPs and climate-associated SNPs produced

different predictions for the rate of genomic change across climate gradients (genomic

turnover) on eastern Santa Cruz Island (Figure S7), suggesting that gene flow occurs between

eastern and western Santa Cruz populations, but that selection results in differentiation of
climate-associated alleles between the regions. Local adaptation could occur as a result of the
precipitation gradient in the Northern Channel Islands, in which eastern regions have less
summer precipitation and greater precipitation seasonality. This trend indicates that local
adaptation may be possible at small scales and in the presence of gene flow, even within
relatively small island populations, as found previously (Cheek et al., 2022; Gamboa et al., 2022;
Gugger et al., 2018; Hamilton et al., 2017; Langin et al., 2015). In contrast, populations on
Catalina and San Clemente show less turnover of adaptive alleles within islands. Candidate
SNPs from Guadalupe showed very little variation or turnover on the Channel Islands (Figure
5A), suggesting that local adaptation among the Guadalupe populations results from variation
in SNPs that are less variable across the Channel Islands. Overall, genomic turnover within
both Guadalupe and Santa Cruz islands indicates fine-scale local adaptation, consistent with
predictions of local adaptation for forest trees (Sork, 2016), and indicating that AGF strategies
using existing genetic variation are viable.

Genetic offset and maladaptation

Genetic offset is a measure of vulnerability to climate change, based on associations between
genomic structure and future climates. Our results predict that, without intervention,
populations on Catalina will be most maladapted to future climates in comparison to the other
Channel Islands populations, followed by eastern Santa Cruz populations (Figure 4C). If gene
flow naturally occurs from the Northern to Southern Channel Islands, as suggested by neutral
genetic structure, both Catalina and San Clemente may currently receive northern alleles that
are maladaptive at hotter sites and could become more maladaptive as climate change
progresses. Catalina is projected to experience a greater increase in temperature than San
Clemente (Figure 1B), resulting in a larger genomic offset. Within Guadalupe, populations at
lower elevations closer to the coast are predicted to have a higher genomic offset. While all
populations are likely to experience some degree of maladaptation to future climates,
populations with a greater genomic offset could be prioritized in conservation efforts. We

cannot compare offsets among Guadalupe and the other islands, so this analysis is most useful

in identifying localities within the island that are most at risk. Lotterhos (2024) cautions

against interpreting genomic offset as a metric of maladaptation when calculated from

genotype-environment associations because it is possible that future climates could be more

beneficial. However, given the absence of Island Oak from warmer and drier microclimates on

the islands, we believe future climates are unlikely to be beneficial for this species.

Evaluation of conservation strategies

For current and future restoration projects taking place on several islands, landscape genomic

findings can inform decisions about what seed sources should be used to maximize success for

plantings. However, different conservation strategies may be preferred depending on

management priorities and policies, and conservation decisions must take those values into

account. If the priority is to maintain "natural" patterns of genetic variation, seeds should only

be introduced among populations that have historically experienced gene flow. However, to

conserve existing Island Oak ecosystems and the species that depend on them, it may be

necessary to introduce alleles that are "pre-adapted" to future climates through assisted gene

flow (Aitken & Bemmels, 2016; Aitken & Whitlock, 2013). By investigating genetic structure

and determining climate suitability for all possible seed sources and planting sites, it is possible

to determine the risks and benefits of different strategies. Here, we compare the predicted

genomic offset and climate suitability under three different management strategies: (1)

conservation of local oak populations without AGF (*status quo*) which may include the use of

local acorns in restoration projects, (2) ecosystem preservation – conserving existing Island

Oak ecosystems by introducing non-local populations that are likely to have the greatest

fitness under future conditions, or (3) species preservation – conserving the species and its

genetic variation in at least part of its native range by planting populations at sites with

optimal future climate.

Since all populations are predicted to be maladapted to future conditions (Figure 4C and 5B),

maintaining historic patterns of gene flow may not effectively conserve populations. Because

future climates are projected to be far outside the current climate space occupied by island oak

(Figure S8), there may not be existing genetic variation within the species that will be well-

adapted to future climates. As a result, the risk of maladaptation is reduced most by preserving

populations at sites that will have climate most similar to that of current Island Oak

populations, as in the species preservation approach (Figure 6). Using this conservation

approach, the optimal planting sites for all populations are those with the coolest projected

future climate – Santa Rosa Island, followed by San Clemente Island (Figure 7). These

locations could be used to preserve genotypes from across the species range. However, this

approach would focus on conserving Island Oak in only a portion of its native range,

potentially neglecting Island Oak ecosystems in regions most affected by climate change.

While the ecosystem preservation approach does not reduce genomic offset as much as the

species preservation approach, it is an improvement over the *status quo* (Figure 6), preserving

existing oak groves by introducing genotypes that are likely to be best-adapted to their future

conditions. The ideal outcome would conserve all existing island oak ecosystems; for this

reason we suggest prioritizing ecosystem-preservation, while also conserving genetic diversity

as a bet-hedge to prevent extinction and loss of genetic variation.

Climate suitability can be used to compare possible seed sources for a given planting site while

taking into account other goals, such as conserving the species' genetic structure. For example,

the ideal seed sources for Santa Rosa's future climate are populations on Catalina and San

Clemente, the warmer Southern Channel Islands, which have experienced more admixture

with *Q. chrysolepis*. While AGF may introduce beneficial alleles, it may also disrupt the natural

genetic structure of populations on Santa Rosa island, which appear to be relatively isolated

and have no evidence of admixture with *Q. chrysolepis*. However, the climate suitability of

nearby central Santa Cruz Island populations is nearly as high as those for the Southern

Channel Islands (Figure 7). As the two islands were connected as recently as 11,000 years ago

(Kennett et al., 2008; Reeder-Myers et al., 2015), using central Santa Cruz as a seed source for

Santa Rosa could represent a more moderate AGF approach that could still introduce beneficial

alleles. Using climate suitability, managers can identify a subset of optimal seed sources, then

select those that best maintain historical evolutionary structure for planting.

For Guadalupe Island, the optimal seed source was a region with high BIO6 (minimum
temperature of the coldest month) and the optimal planting site was the westernmost part of
the island sampled (Figure 5, Table S1). In preserving Guadalupe Island genotypes, predicted
maladaptation was minimized when planting within the island, even when the cooler Channel
Islands were included as possible planting sites (Table S1). The Guadalupe Island Oaks appear
to be differentiated from the Channel Islands at both neutral and putatively adaptive SNPs, and
are unlikely to experience contemporary gene flow from these populations. Because of their
divergence, it is possible that hybridization between Guadalupe and Channel Islands
populations could result in inbreeding depression. Therefore, the Guadalupe island oaks are
probably best managed as a separate lineage at present. However, including Guadalupe
populations in future studies, including common garden experiments, could help determine
whether they contain unique genetic variation could be beneficial under hotter climates when
introduced to the Channel Islands.

The ecosystem- and species-preservation approaches could be complementary and used in
combination with other conservation strategies. Because Santa Rosa Island will have cooler
future climates than other islands it could be an ideal planting site for other populations.
Moreover, because the native Santa Rosa populations are likely adapted to a cooler, more
moderate climate, they could benefit from the introduction of genetic variation that is likely to
be beneficial under climate change. In addition to maintaining genetic diversity within the
native range, *ex situ* collections in botanic gardens and arboreta can be used to conserve the
genetic diversity of species with seeds that cannot be stored through traditional seed banking,
such as oaks (Rosenberger et al., 2022; Westwood et al., 2021). These living collections could be
used as tissue or seed sources for restoration, or for replanting when a natural population is
lost in events such such as wildfire, landslide, or pathogen outbreaks. Lastly, AGF is not likely
to be a viable strategy for populations that are predicted to be most maladapted to future
temperatures and are also from the warmest part of the species range, such as Catalina Island.
For such populations, we propose that supplementary ecological studies should be undertaken

to identify seed sources (for example, specific maternal trees) that may be better adapted to

future climate conditions. Additionally, a genome-informed breeding approach could be used

to predict which genotypes are likely to have the greatest fitness under future environments by

associating fitness with genomic variation (Browne et al., 2019).

Limitations

Landscape genomics tools can be a powerful tool for identifying strategies to conserve species

under changing climates; however, our recommendations depend on the assumption that

populations are locally adapted to modern climate conditions. Because *Q. tomentella* is a

paleoendemic that experienced range contraction and likely survives on the California Islands

because of their more moderate climates, it is possible that populations are already maladapted

to present climate conditions, as found for California populations of valley oak (*Q. lobata*)

(Browne et al., 2019), meaning we underestimate maladaptation here. Furthermore, studies that

have tested whether genomic offset can predict growth in a common garden have found mixed

results (Fitzpatrick et al., 2021; Lind et al., 2023) so genetic and climate data may not be

sufficient to predict optimal climates. Additionally, our predictions define optimal climate as

the mean values over 30 years for a 30-second square grid cell. In reality, trees experience

climatic variation across seasons, years, and microclimates (such as those created by the north-

facing canyons that Island Oaks often inhabit) which are not represented by this climate data.

Common garden experiments could validate our predictions of ideal population-climate

matches prior to implementing assisted gene flow. For example, populations from across the

species range could be planted at a restoration site and monitored over 5-10 years to determine

which have the highest fitness at the site, and could be removed before they reach reproductive

age if introducing non-local alleles is a concern. Santa Rosa Island would be an ideal location

for such an experiment, as it is the optimal planting site for all populations, and local

populations could potentially benefit from the introduction of warmer-adapted populations.

Nonetheless, our findings suggest that island populations are likely to be maladapted to future

climates and that AGF can reduce the predicted maladaptation, so genomic-informed seed

sourcing should be considered as a possible management strategy to conserve and restore

Island Oak ecosystems.

Conclusions

Assisted gene flow could reduce maladaptation of Island Oak to future climates relative to the
restoring populations with local seeds. Conserving the genetic diversity of populations on
islands that are projected to have less dramatic climate shifts, in this case, Santa Rosa and San
Clemente, would conserve the species in parts of its native range, and introducing pre-adapted
genotypes to populations could help preserve existing island oak ecosystems. These two AGF
strategies could be combined together: for example, by establishing restorations at ideal sites
that will conserve rangewide genetic diversity while also introducing beneficial alleles to
existing oak populations that would otherwise not be well-adapted to future climates. This
study identified one locality of major concern (Catalina Island), which may require further
ecological and genomic studies to identify seed sources within the island that may be better
adapted to future climate conditions. While Island Oak is likely to be threatened by climate
change, we show that genomic tools can compare among possible conservation strategies that
may allow these unique ecosystems persist.

Data Availability

659 Sequencing data will be uploaded to NCBI upon manuscript acceptance. Locality and climate
660 data for collected samples are available as Table S1.

661 Analysis scripts will be made available on Github and archived at Zenodo upon manuscript
662 acceptance. R markdown output files are included as supplementary information.

Acknowledgements

This work took place on the ancestral lands of the Chumash (Northern Channel Islands) and
the Tongva (Southern Channel Islands), and we acknowledge them as the traditional caretakers
of the land.

We thank those who helped us with fieldwork and collecting (in alphabetical order): Elizabeth

Becker, James Cabrera, Lauren Dennhardt, Adam Fontenot, Robert Kaaret, Seth Kauppinen,

Reena Lam, Betty Lee, Zhizhong Li, Sergio A. Luvianos-Colín, Bryan Munson, Scott O'Donnell,

John Randall, Joseph Retelskyj, and Kristi Smith. Kari Merrill and Dan Oliveira helped perform

DNA extractions. Sequencing was performed by the DNA Technologies and Expression

Analysis Cores at the UC Davis Genome Center, supported by NIH Shared Instrumentation

Grant 1S10OD010786-01. A special note of appreciation to Luciana Luna for acquiring the

federal permits, coordinating collections, and transporting samples from Guadalupe Island.

Thank you to Jim Howard, Annie Little, David Mazurkiewicz, and Cameron Williams for

supporting collections within the Channel Islands National Park, to Scott Morrison for

supporting this project, and to Jen Baker for her support on the Santa Cruz Island Preserve.

This work was supported and funded by both The Nature Conservancy and the California

Conservation Genomics Project (awarded to University of California by the State of California,

State Budget Act of 2019 [UC Award ID RSI-19-690224]). AM was supported by these sources

and by NSF-PRFB-2209410.

References

Aitken, S. N., & Bemmels, J. B. (2016). Time to get moving: Assisted gene flow of forest trees. *Evolutionary Applications*, *9*(1), 271–290. https://doi.org/10.1111/eva.12293

Aitken, S. N., & Whitlock, M. C. (2013). Assisted gene flow to facilitate local adaptation to climate change. *Annu. Rev. Ecol. Evol. Syst*, *44*, 367–388. https://doi.org/10.1146/annurev-ecolsys-110512-135747

Aitken, S. N., Yeaman, S., Holliday, J. A., Wang, T., & Curtis-McLane, S. (2008). Adaptation, migration or extirpation: Climate change outcomes for tree populations. *Evolutionary Applications*, *1*(1), 95–111. https://doi.org/10.1111/j.1752-4571.2007.00013.x

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, *19*(9), 1655–1664. https://doi.org/10.1101/gr.094052.109

Ashley, M. V., Backs, J. R., Kindsvater, L., & Abraham, S. T. (2018). Genetic variation and structure in an endemic island oak, *Quercus tomentella*, and mainland canyon oak, *Quercus chrysolepis*. *International Journal of Plant Sciences*, *179*(2), 151–161. https://doi.org/10.1086/696023

Axelrod, D. I. (1939). *A miocene flora from the western border of the Mohave desert.* The Carnegie Institution of Washington.

Axelrod, D. I. (1944a). The Mulholland Flora. In R. W. Chaney (Ed.), *Pliocene floras of California and Oregon* (pp. 103–162). The Carnegie Institution of Washington.

Axelrod, D. I. (1944b). The Sonoma Flora. In R. W. Chaney (Ed.), *Pliocene floras of California and Oregon* (pp. 167–218). The Carnegie Institution of Washington.

Axelrod, D. I. (1967). Geologic history of the Californian insular flora. *1st Symposium on the Biology of the California Islands*, 267–314.

Beckman, E., & Jerome, D. (2017). Quercus tomentella. IUCN Red List of Threatened Species. https://www.iucnredlist.org/species/30959/2799049

Beckman, E., Meyer, A., Denvir, A., Gill, D., Man, G., Pivorunas, D., Shaw, K., & Westwood, M. (2019). *Conservation Gap Analysis of Native U.S. Oaks.* The Morton Arboretum.

Browne, L., Wright, J. W., Fitz-Gibbon, S., Gugger, P. F., & Sork, V. L. (2019). Adaptational lag to temperature in valley oak (*Quercus lobata*) can be mitigated by genome-informed assisted gene flow. *Proceedings of the National Academy of Sciences*, *116*(50), 25179–25185. https://doi.org/10.1073/pnas.1908771116

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-generation PLINK: Rising to the challenge of larger and richer datasets. *GigaScience*, *4*(1), s13742-015-0047–0048. https://doi.org/10.1186/s13742-015-0047-8

Cheek, R. G., Forester, B. R., Salerno, P. E., Trumbo, D. R., Langin, K. M., Chen, N., Scott Sillett, T., Morrison, S. A., Ghalambor, C. K., & Chris Funk, W. (2022). Habitat-linked genetic variation supports microgeographic adaptive divergence in an island-endemic bird species. *Molecular Ecology*, *31*(10), 2830–2846. https://doi.org/10.1111/mec.16438

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008

Delaney, K. S., & Wayne, R. K. (2005). Adaptive Units for Conservation: Population Distinction and Historic Extinctions in the Island Scrub-Jay. *Conservation Biology*, *19*(2), 523–533. https://doi.org/10.1111/j.1523-1739.2005.00424.x

Di Santo, L. N., Hoban, S., Parchman, T. L., Wright, J. W., & Hamilton, J. A. (2022). Reduced representation sequencing to understand the evolutionary history of Torrey pine (*Pinus torreyana* parry) with implications for rare species conservation. *Molecular Ecology*, *31*(18), 4622–4639. https://doi.org/10.1111/mec.16615

Ellis, N., Smith, S. J., & Pitcher, C. R. (2012). Gradient Forests: Calculating importance gradients on physical predictors. *Ecology*, *93*, 156–168.

Fitzpatrick, M. C., Chhatre, V. E., Soolanayakanahally, R. Y., & Keller, S. R. (2021). Experimental support for genomic prediction of climate maladaptation using the machine learning approach Gradient Forests. *Molecular Ecology Resources*, *21*(8), 2749–2765. https://doi.org/10.1111/1755-0998.13374

Fitzpatrick, M. C., & Keller, S. R. (2015). Ecological genomics meets community-level modelling of biodiversity: Mapping the genomic landscape of current and future environmental adaptation. *Ecology Letters*, *18*(1), 1–16. https://doi.org/10.1111/ele.12376

Forester, B. R., Lasky, J. R., Wagner, H. H., & Urban, D. L. (2018). Comparing methods for detecting multilocus adaptation with multivariate genotype–environment associations. *Molecular Ecology*, *27*(9), 2215–2233. https://doi.org/10.1111/mec.14584

Gamboa, M. P., Ghalambor, C. K., Scott Sillett, T., Morrison, S. A., & Chris Funk, W. (2022). Adaptive divergence in bill morphology and other thermoregulatory traits is facilitated by restricted gene flow in song sparrows on the California Channel Islands. *Molecular Ecology*, *31*(2), 603–619. https://doi.org/10.1111/mec.16253

García-Verdugo, C., Sajeva, M., La Mantia, T., Harrouni, C., Msanda, F., & Caujapé-Castells, J. (2015). Do island plant populations really have lower genetic variation than mainland populations? Effects of selection and distribution range on genetic diversity estimates. *Molecular Ecology*, *24*(4), 726–741. https://doi.org/10.1111/mec.13060

Goudet, J., & Jombart, T. (2022). *hierfstat: Estimation and Tests of Hierarchical F-Statistics* (0.5-11). https://CRAN.R-project.org/package=hierfstat

Gougherty, A. V., Keller, S. R., & Fitzpatrick, M. C. (2021). Maladaptation, migration and extirpation fuel climate change risk in a forest tree species. *Nature Climate Change*, *11*(2), 166–171. https://doi.org/10.1038/s41558-020-00968-6

Grummer, J. A., Booker, T. R., Matthey-Doret, R., Nietlisbach, P., Thomaz, A. T., & Whitlock, M. C. (2022). The immediate costs and long-term benefits of assisted gene flow in large populations. *Conservation Biology*, *36*(4), e13911. https://doi.org/10.1111/cobi.13911

Gugger, P. F., Fitz-Gibbon, S. T., Albarrán-Lara, A., Wright, J. W., & Sork, V. L. (2021). Landscape genomics of *Quercus lobata* reveals genes involved in local climate adaptation at multiple spatial scales. *Molecular Ecology*, *30*(2), 406–423.

https://doi.org/10.1111/mec.15731

Gugger, P. F., Liang, C. T., Sork, V. L., Hodgskiss, P., & Wright, J. W. (2018). Applying landscape genomic tools to forest management and restoration of Hawaiian koa (*Acacia koa*) in a changing environment. *Evolutionary Applications*, *11*(2), 231–242. https://doi.org/10.1111/eva.12534

Hamabata, T., Kinoshita, G., Kurita, K., Cao, P.-L., Ito, M., Murata, J., Komaki, Y., Isagi, Y., & Makino, T. (2019). Endangered island endemic plants have vulnerable genomes. *Communications Biology*, *2*(1), 1. https://doi.org/10.1038/s42003-019-0490-7

Hamilton, J. A., Royauté, R., Wright, J. W., Hodgskiss, P., & Ledig, F. T. (2017). Genetic conservation and management of the California endemic, Torrey pine (*Pinus torreyana* Parry): Implications of genetic rescue in a genetically depauperate species. *Ecology and Evolution*, *7*(18), 7370–7381. https://doi.org/10.1002/ece3.3306

IPCC. (2014). *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* https://www.ipcc.ch/report/ar5/syr/

Jia, K.-H., Zhao, W., Maier, P. A., Hu, X.-G., Jin, Y., Zhou, S.-S., Jiao, S.-Q., El-Kassaby, Y. A., Wang, T., Wang, X.-R., & Mao, J.-F. (2020). Landscape genomics predicts climate change-related genetic offset for the widespread *Platycladus orientalis* (Cupressaceae). *Evolutionary Applications*, *13*(4), 665–676. https://doi.org/10.1111/eva.12891

Kennett, D. J., Kennett, J. P., West, G. J., Erlandson, J. M., Johnson, J. R., Hendy, I. L., West, A., Culleton, B. J., Jones, T. L., & Stafford, T. W. (2008). Wildfire and abrupt ecosystem disruption on California's Northern Channel Islands at the Ållerød–Younger Dryas boundary (13.0–12.9ka). *Quaternary Science Reviews*, *27*(27), 2530–2545. https://doi.org/10.1016/j.quascirev.2008.09.006

Kremer, A., & Hipp, A. L. (2020). Oaks: An evolutionary success story. *New Phytologist*, *226*(4), 987–1011. https://doi.org/10.1111/nph.16274

Lachmuth, S., Capblancq, T., Prakash, A., Keller, S. R., & Fitzpatrick, M. C. (2024). Novel genomic offset metrics integrate local adaptation into habitat suitability forecasts and inform assisted migration. *Ecological Monographs*, *94*(1), e1593. https://doi.org/10.1002/ecm.1593

Langin, K. M., Sillett, T. S., Funk, W. C., Morrison, S. A., Desrosiers, M. A., & Ghalambor, C. K. (2015). Islands within an island: Repeated adaptive divergence in a single population. *Evolution*, *69*(3), 653–665. https://doi.org/10.1111/evo.12610

Leimu, R., & Fischer, M. (2008). A meta-analysis of local adaptation in plants. *PLOS ONE*, *3*(12), e4010. https://doi.org/10.1371/journal.pone.0004010

Lind, B. M., Candido-Ribeiro, R., Singh, P., Lu, M., Vidakovic, D. O., Booker, T. R., Whitlock, M. C., Yeaman, S., Isabel, N., & Aitken, S. N. (2023). How useful is genomic data for predicting maladaptation to future climate? *BioRxiv.* https://doi.org/10.1101/2023.02.10.528022

Lotterhos, K. E. (2024). Interpretation issues with "genomic vulnerability" arise from conceptual issues in local adaptation and maladaptation. *Evolution Letters*, qrae004.

https://doi.org/10.1093/evlett/qrae004

Martins, K., Gugger, P. F., Llanderal-Mendoza, J., González-Rodríguez, A., Fitz-Gibbon, S. T., Zhao, J.-L., Rodríguez-Correa, H., Oyama, K., & Sork, V. L. (2018). Landscape genomics provides evidence of climate-associated genetic variation in Mexican populations of *Quercus rugosa*. *Evolutionary Applications*, *11*(10), 1842–1858. https://doi.org/10.1111/eva.12684

Mead, A., Fitz-Gibbon, S. T., Escalona, M., Beraut, E., Sacco, S., Marimuthu, M. P. A., Nguyen, O., & Sork, V. L. (2024). The genome assembly of Island Oak (*Quercus tomentella*), a relictual island tree species. *Journal of Heredity*, esae002. https://doi.org/10.1093/jhered/esae002

Morrison, S. A., Sillett, T. S., Ghalambor, C. K., Fitzpatrick, J. W., Graber, D. M., Bakker, V. J., Bowman, R., Collins, C. T., Collins, P. W., Delaney, K. S., Doak, D. F., Koenig, W. D., Laughrin, L., Lieberman, A. A., Marzluff, J. M., Reynolds, M. D., Scott, J. M., Stallcup, J. A., Vickers, W., & Boyce, W. M. (2011). Proactive Conservation Management of an Island-endemic Bird Species in the Face of Global Change. *BioScience*, *61*(12), 1013–1021. https://doi.org/10.1525/bio.2011.61.12.11

Muller, C. H. (1965). Relictual origins of insular endemics in *Quercus*. *1st Symposium on the Biology of the California Islands*.

O'Donnell, S. T., Fitz-Gibbon, S. T., & Sork, V. L. (2021). Ancient Introgression Between Distantly Related White Oaks (*Quercus* sect. *Quercus*) Shows Evidence of Climate-Associated Asymmetric Gene Exchange. *Journal of Heredity*, *112*(7), 663–670. https://doi.org/10.1093/jhered/esab053

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2019). *vegan: Community Ecology Package* (2.5-5). https://CRAN.R-project.org/package=vegan

Ortego, J., Gugger, P. F., & Sork, V. L. (2018). Genomic data reveal cryptic lineage diversification and introgression in Californian golden cup oaks (section *Protobalanus*). *New Phytologist*, *218*(2), 804–818. https://doi.org/10.1111/nph.14951

Pesendorfer, M. B., Baker, C. M., Stringer, M., McDonald-Madden, E., Bode, M., McEachern, A. K., Morrison, S. A., & Sillett, T. S. (2018). Oak habitat recovery on California's largest islands: Scenarios for the role of corvid seed dispersal. *Journal of Applied Ecology*, *55*(3), 1185–1194. https://doi.org/10.1111/1365-2664.13041

Petit, R. J., & Hampe, A. (2006). Some evolutionary consequences of being a tree. *Annual Review of Ecology, Evolution, and Systematics*, *37*, 187–214.

Reeder-Myers, L., Erlandson, J. M., Muhs, D. R., & Rick, T. C. (2015). Sea level, paleogeography, and archeology on California's Northern Channel islands. *Quaternary Research*, *83*(2), 263–272. https://doi.org/10.1016/j.yqres.2015.01.002

Rellstab, C., Gugerli, F., Eckert, A. J., Hancock, A. M., & Holderegger, R. (2015). A practical guide to environmental association analysis in landscape genomics. *Molecular Ecology*, *24*(17), 4348–4370. https://doi.org/10.1111/mec.13322

Rick, T. C., Hofman, C. A., & Reeder-Myers, L. A. (2019). Why Translocate? Evaluating the Evidence and Reasons for Ancient Human Introductions of Wildlife to California's Islands. In K. M. Gill, M. Fauvelle, & J. M. Erlandson (Eds.), *An Archaeology of Abundance: Reevaluating the Marginality of California's Islands* (pp. 248–272). University Press of Florida.

Rosenberger, K., Schumacher, E., Brown, A., & Hoban, S. (2022). Species-tailored sampling guidelines remain an efficient method to conserve genetic diversity ex situ: A study on threatened oaks. *Biological Conservation, 275*, 109755. https://doi.org/10.1016/j.biocon.2022.109755

Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation. *Nature Reviews Genetics, 14*(11), 807–820. https://doi.org/10.1038/nrg3522

Savolainen, O., Pyhäjärvi, T., & Knürr, T. (2007). Gene Flow and Local Adaptation in Trees. *Annual Review of Ecology, Evolution, and Systematics, 38*(1), 595–619. https://doi.org/10.1146/annurev.ecolsys.38.091206.095646

Sawyer, J. O., Keeler-Wolf, T., & Evens, J. M. (2009). CNPS Alliance: *Quercus tomentella—Lyonothamnus floribundus*. In *A Manual of California Vegetation, Second Edition* (p. 1300). California Native Plant Society. https://vegetation.cnps.org/alliance/523

Sexton, J. P., Hangartner, S. B., & Hoffmann, A. A. (2014). Genetic Isolation by Environment or Distance: Which Pattern of Gene Flow Is Most Common? *Evolution, 68*(1), 1–15. https://doi.org/10.1111/evo.12258

Shryock, D. F., Washburn, L. K., DeFalco, L. A., & Esque, T. C. (2021). Harnessing landscape genomics to identify future climate resilient genotypes in a desert annual. *Molecular Ecology, 30*(3), 698–717. https://doi.org/10.1111/mec.15672

Sork, V. L. (2016). Gene flow and natural selection shape spatial patterns of genes in tree populations: Implications for evolutionary processes and applications. *Evolutionary Applications, 9*(1), 291–310. https://doi.org/10.1111/eva.12316

Sork, V. L. (2017). Genomic studies of local adaptation in natural plant populations. *Journal of Heredity, 109*(1), 3–15. https://doi.org/10.1093/jhered/esx091

Sork, V. L., Aitken, S. N., Dyer, R. J., Eckert, A. J., Legendre, P., & Neale, D. B. (2013). Putting the landscape into the genomics of trees: Approaches for understanding local adaptation and population responses to changing climate. *Tree Genetics and Genomes, 9*(4), 901–911. https://doi.org/10.1007/s11295-013-0596-x

Sork, V. L., Cokus, S. J., Fitz-Gibbon, S. T., Zimin, A. V., Puiu, D., Garcia, J. A., Gugger, P. F., Henriquez, C. L., Zhen, Y., Lohmueller, K. E., Pellegrini, M., & Salzberg, S. L. (2022). High-quality genome and methylomes illustrate features underlying evolutionary success of oaks. *Nature Communications, 13*(1), 1. https://doi.org/10.1038/s41467-022-29584-y

Suarez-Gonzalez, A., Lexer, C., & Cronk, Q. C. B. (2018). Adaptive introgression: A plant perspective. *Biology Letters, 14*(3), 20170688. https://doi.org/10.1098/rsbl.2017.0688

Underwood, E. C., Hollander, A. D., Safford, H. D., Kim, J. B., Srivastava, L., & Drapek, R. J. (2019). The impacts of climate change on ecosystem services in southern California.

Ecosystem Services, *39*, 101008. https://doi.org/10.1016/j.ecoser.2019.101008

Villa-Machío, I., Fernández de Castro, A. G., Fuertes-Aguilar, J., & Nieto Feliner, G. (2020). Colonization history of the Canary Islands endemic Lavatera acerifolia, (Malvaceae) unveiled with genotyping-by-sequencing data and niche modelling. *Journal of Biogeography*, *47*(4), 993–1005. https://doi.org/10.1111/jbi.13808

Wang, I. J., & Bradburd, G. S. (2014). Isolation by environment. *Molecular Ecology*, *23*(23), 5649–5662. https://doi.org/10.1111/mec.12938

Westwood, M., Cavender, N., Meyer, A., & Smith, P. (2021). Botanic garden solutions to the plant extinction crisis. *PLANTS, PEOPLE, PLANET*, *3*(1), 22–32. https://doi.org/10.1002/ppp3.10134

Yu, Y., Aitken, S. N., Rieseberg, L. H., & Wang, T. (2022). Using landscape genomics to delineate seed and breeding zones for lodgepole pine. *New Phytologist*, *235*(4), 1653–1664. https://doi.org/10.1111/nph.18223