1  **The importance of small island populations for the long term survival of endangered large-**
2  **bodied insular mammals**
3

Sabhrina Gita Aninta[1*#], Rosie Drinkwater[1,2#], Alberto Carmagnini[2], Nicolas J. Deere[3], Dwi Sendi
Priyono[4], Noviar Andayani[5], Nurul L. Winarni[6], Jatna Supriatna[5], Matteo Fumagalli[1], Greger
Larson[7], Peter H.A. Galbusera[8], Alastair Macdonald[9], Deborah Greer[1], Kusdiantoro Mohamad[10],
Wahono Esthi Prasetyaningtyas[10], Abdul Haris Mustari[11], John Lewis Williams[12], Ross Barnett[13],
Darren Shaw[9], Gono Semiadi[14], James Burton[15], David Seaman[16], Maria Voigt[16], Matthew
Struebig[16], Selina Brace[17], Stephen Rossiter[1], Laurent Frantz[1,2*]
10

[1]School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK
[2]Palaeogenomics group, Faculty of Veterinary Medicine, Ludwig Maximilian University Munich
[3]Durrell Institute of Conservation and Ecology (DICE), University of Kent, Canterbury, CT2 7NR,
UK
[4]Department of Tropical Biology, Faculty of Biology, Universitas Gadjah Mada, Sleman, 55281,
Special Region of Yogyakarta, Indonesia.
[5]Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Indonesia,
Depok, 16424, Indonesia
[6]Research Center for Climate Change, University of Indonesia, Depok, Indonesia
[7]The Palaeogenomics and Bio-Archaeology Research Network, Research Laboratory for
Archaeology and History of Art, University of Oxford, Oxford, UK
[8]Antwerp Zoo Centre for Research and Conservation, Royal Zoological Society of Antwerp,
Antwerp, Belgium
[9]Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Midlothian EH25 9RG
[10]School of Veterinary Medicine and Biomedical Sciences, IPB University, Bogor, Indonesia
[11]Department of Forest Resources Conservation, Faculty of Forestry and Environment, IPB
University, Bogor, Indonesia
[12]Department of Animal Science, Food and Nutrition, Università Cattolica del Sacro Cuore, via
Emilia Parmense, 84, 29122 Piacenza, Italy
[13]Durham Evolution and Ancient DNA, Department of Archaeology, University of Durham, Durham,
United Kingdom
[14]Research Centre for Biosystematics and Evolution, National Research and Innovation Agency
(BRIN), Cibinong 16911, Indonesia
[15]IUCN SSC Asian Wild Cattle Specialist Group
[16]Durrell Institute of Conservation and Ecology (DICE), University of Kent, Canterbury, CT2 7NR
[17]Department of Earth Sciences, Natural History Museum, London, England, United Kingdom,
SW7 5BD, UK
38

39  [#] Equally contributed
40  *corresponding authors: sagitaninta@gmail.com; r.drinkwater@lmu.de; laurent.frantz@lmu.de;
41

**Summary**
Small island populations, particularly those of large vertebrates, are vulnerable to extinction due to
environmental stochasticity and genetic drift. Here, we combined analyses of 113 newly generated
~10x genomes of anoa (*Bubalus* spp.) and babirusa (*Babyrousa* spp.) with habitat suitability
models to assess the evolutionary trajectories of small and large island populations within the
Wallacea archipelago in Indonesia. Our analyses show that, compared to closely related species,
populations on the largest island of Sulawesi exhibit low levels of heterozygosity, high levels of
inbreeding, and a high proportion of deleterious alleles. In contrast, although populations on

smaller islands such as Buton and Togean exhibit lower heterozygosity and higher inbreeding than
in larger islands, they also possess fewer deleterious mutations. Analyses of site frequency spectra
analyses indicate this likely the result of stronger, long-term purging acting in population from
smaller islands. Combined with species distribution models, our results show that populations on
small islands have persisted over a long time frame in high quality habitat while populations in the
larger island have recently suffered from low habitat quality as a result of heavy exploitation of
natural resources. In the case of anoa and babirusa, preservation of small island populations
should focus more on improving habitat quality in small islands and avoid translocating individuals
from the larger island. In the face of ongoing land-use change on the Sulawesi mainland, we
suggest that offshore islands are increasingly crucial for the long-term survival of Wallacea's iconic
and indigenous mammals.
61
**Results and Discussion**
Throughout the Quaternary, populations of large vertebrates on islands have experienced higher
rates of extinction compared to their mainland counterparts[1], likely reflecting the effects of genetic
drift that operate in small populations of slow-reproducing species, and the associated negative
consequences for fitness and adaptability[2,3]. In more recent years, these long-term extinction risks
have been further exacerbated by human activities, including deforestation, mining and hunting[4].
68
Anthropogenic pressures can vary widely among islands. On the Wallacea archipelago (Figure 1),
a global centre of endemism, the exploitation of timber[5] and mineral resources[6] has occurred more
intensively on the large island of Sulawesi than on the smaller offshore islands[7]. As a result, while
small island populations are more likely to go extinct over geological time scale, the relatively low
human impact on small island ecosystems makes them attractive locations for species
conservation programs.
75
Small island populations, however, can suffer from inbreeding depression due to low genetic
diversity[8]. One solution to improve their viability is to translocate individuals from other populations
(e.g. from larger islands where anthropogenic pressures are more acute) to increase genetic
diversity[9]. Aside from increasing numbers of individuals, translocation can lead to genetic rescue, a
process through which the average fitness of individuals is improved by increasing heterozygosity
and decreasing inbreeding[10].
82
Despite the proposed benefit of genetic rescue, both recent empirical work[11] and simulations[12]
suggest that it could lead to long term adverse effects through the introduction of recessive
deleterious mutations if the source population is larger than the population to be rescued. In fact,
recent work has shown that in the case of historically small populations that have had time to purge
highly deleterious mutations, translocation can lead to counterproductive outcomes[12]. It is therefore
crucial to conduct thorough genetic evaluations prior to establishing onerous translocation
interventions[13].
90
The two endangered forest-dependent mammalian genera anoa (*Bubalus depressicornis* and
*Bubalus quarlesi*) and babirusa (*Babyrousa* spp.) expanded across the Wallacea archipelago
approximately 2 Mya[14] . These taxa now occur on islands of different sizes (from Togean,
~230km$^2$, to Sulawesi ~180,000 km$^2$) where they are subject to contrasting levels of anthropogenic
activity[7]. As a result,these taxa are potential targets for translocation and genetic rescue. Their
evolutionary history also means they are excellent models to understand how long term (i.e.
geological timescales) and recent disturbances (i.e. anthropogenic) affect microevolutionary
processes in large-bodied insular species.

To assess the effect that long term isolation and recent human activities have had on the genetic
variation of anoa and babirusa, we combined analyses of newly generated genomic data (113
genomes; ~10x coverage; Table S1) sampled across the larger island of Sulawesi and smaller
islands of Buton and Togean with habitat suitability models. Our results provide guidance for
conservation efforts, demonstrating that small island populations have persisted over a long time
frame in high quality habitats, while their counterparts on larger islands have experienced
diminished habitat quality as a result of heavy exploitation of natural resources.

We aligned anoa and babirusa short-read sequences to the water buffalo (*Bubalus bubalis*) and
babirusa reference genome respectively for all downstream analysis, except for genomic load
analyses for which we used the better annotated reference genome of cattle (*Bos taurus*) and pig
(*Sus scrofa*) (Supplementary Information). Principal component analysis (Figure 1C & D), and
phylogenetic trees (Figure S2) based on nuclear single nucleotide polymorphisms (SNPs),
indicated the existence of four, well supported, lineages of anoa (North Sulawesi, East Central,
Southeast Sulawesi and Buton) and three lineages of babirusa (North Sulawesi, Southeast
Sulawesi and Togean) that map clearly onto the geography of Sulawesi and nearby islands. These
results are largely in agreement with previous findings based on mitochondrial and microsatellite
loci[14]. The phylogeographic structure defined here formed the basis of the population definition
used in all subsequent analyses (Figure 1).

**Populations on smaller island are less genetically diverse than on mainland Sulawesi**
We computed genome-wide Watterson's $\theta$ using ROHan[16]. We found that $\theta$ on the small islands of
Togean (babirusa: mean=0.29 x $10^{-3}$, SE=0.02 x $10^{-3}$) and Buton (anoa: mean=0.58 x $10^{-3}$,
SE=0.02 x $10^{-3}$) was ~2 to ~6 fold lower than in populations from mainland Sulawesi (babirusa:
mean=1.85 x $10^{-3}$, SE=0.1 x $10^{-3}$; anoa: mean=1.33 x $10^{-3}$, SE=0.1 x $10^{-3}$), indicating that small
island populations possess less genetic diversity than their mainland counterparts (Figure 2A). On
the large island of Sulawesi, however, we found that babirusa from the Southeast Sulawesi
possessed lower $\theta$ (mean=1.17 x $10^{-3}$, SE=0.04 x $10^{-3}$) compared to those found on the Northern
peninsula (mean=2.17 x $10^{-3}$, SE=0.06 x $10^{-3}$). Low $\theta$ in Southeast Sulawesi babirusa could be the
result of a recent bottleneck.

We assessed whether any of these populations showed signs of recent bottlenecks by analysing
Runs of Homozygosity (ROH) using ROHan (Supplementary Information). Compared to anoa from
mainland Sulawesi, the genomes of individuals from the small island of Buton contained more
ROH segments (1-267, Figure 2B), consistent with higher inbreeding levels (Figure 2C). Overall,
this indicates that anoa populations from small islands are more inbred than those found on
mainland Sulawesi.

For babirusa, levels of inbreeding (measured by the fraction of the genome in ROH) are highest in
individuals in Southeast (mainland) Sulawesi followed by Togean (Figure 2B & C). This analysis
also indicates that babirusa from both regions are more inbred than most other species of suids
including endangered taxa such as the pygmy hog (*Porcula salvania*), with the exception of the
critically endangered Javanese warty pig (*Sus verrucosus*) (Figure 2B). High levels of ROHs in
Southeast babirusa (Figure 2C) indicate that this population likely underwent a recent population
bottleneck.

To address whether this bottleneck in Southeast Sulawesi is consistent with a recent decrease of
suitable habitat in the area, we constructed an ensemble species distribution model[17] based on

bioclimatic variables and forest cover[5,15] (Supplementary Information). Our models, for both anoa
and babirusa, indicate that there is proportionally more highly suitable habitat (i.e. top 20% of all
suitable habitat; see Supplementary Text) on smaller islands (i.e. Togean and Buton) compared to
mainland Sulawesi (~1.5 fold more highly suitable area for anoa and ~1.6 for babirusa relative to
North Sulawesi; Figure 1, Table S7). We also found that, for both species, habitat suitability was
highly correlated with their protection status, where habitats with higher protection were also found
to be more suitable (Supplementary Information, Figure S4).

On the mainland, the availability of suitable habitat was higher in the North compared to the
Southeast for both taxa. This trend is consistent with recent deforestation rates (between 2000-
2017), which were higher in Southeast Sulawesi compared to the North[18], suggesting that
deforestation has led to the reduced habitat availability detected by our models. These data,
combined with previously reported high rates of poaching[19], imply that population bottlenecks in
babirusa in Southeast Sulawesi have been driven by relatively recent anthropogenic disturbances,
rather than longer term evolutionary processes.

For both babirusa and anoa, we found that differences in levels of genome-wide $\theta$ (including ROH)
and $\theta$ outside of ROH in small island populations were much less pronounced than in Southeast
Sulawesi babirusa (Figure 2B). In Southeast Sulawesi babirusa, $\theta$ computed genome-wide (i.e.
including ROHs) was significantly lower than when computed excluding ROHs (p<0.001; Figure
2B). This pattern implies that babirusa living in Southeast Sulawesi were part of a historically large
population which underwent a bottleneck. Babirusa and anoa living on small islands (Togean and
Buton), however, likely form part of a historically small, yet stable, population that inhabits a highly
suitable habitat, as demonstrated by our species distribution model (Figure 1, Figure S4).
Altogether, this suggests that although they are not as genetically diverse as their mainland
counterparts, small island populations are likely to have been stable over geological time scales,
and potentially since their expansion ~2Mya[14].

**More deleterious alleles segregate on Sulawesi than on small islands**
Recent studies indicated that other historically small populations (*e.g.* Channel Island foxes[20],
Iberian lynx[21], and Indian tigers[22]) have experienced long histories of purging strongly deleterious
recessive variants. To evaluate how small island populations purge deleterious mutations
compared to mainland populations, we computed genomic load in individual genomes, using three
conservation scores: SIFT, PhyloP and phastCons (Supplementary Information). Genomic load
was computed at one-to-one pig-cow ortholog loci (Supplementary Information), to allow cross
species comparison. We first computed a load score by summing conservation scores, weighted
by genotype probability for derived alleles found at homozygous states across the genome (see
Supplementary Information). This homozygous load represents the minimum impact of deleterious
alleles on fitness (i.e. assumes that all deleterious mutations are recessive). We also computed the
load by summing the impact of alleles found at both homozygous and heterozygous states (total
load) across the genome. The total load represents the maximum impact of deleterious alleles on
fitness (i.e. assumes that all deleterious mutations are dominant).

Both homozygous and total load were significantly lower in anoa than in babirusa (p < 0.001; See
Supplementary Information) (Figure 3A & B). Mean total load, computed using all the individuals
from the same population, was also significantly higher in large populations found on mainland
Sulawesi than in smaller island populations such as Togian (babirusa) and Buton (anoa; Figure 3C,
Table S1). This result indicates that deleterious mutations are more abundant in larger, mainland,

populations than in smaller, island populations, despite the fact that larger populations are
expected to purge deleterious mutations more efficiently[12].
198
Mean homozygous load, however, was significantly higher in small island populations than in
larger populations (Figure 3D). Altogether these results indicate that large populations possess
more deleterious recessive alleles overall, most of which are found in heterozygous state and
therefore hidden from selection. Thus, although large populations possess more deleterious
alleles, the overall fitness impact of deleterious mutations is likely to be higher in smaller island
populations which possess more deleterious alleles in homozygous state.
205
**206 Small island populations efficiently purge deleterious alleles**
Deleterious alleles are potentially more likely to be found in homozygous state in individuals from
small island populations due to their lower levels of heterozygosity (Figure 2B). This could lead to
the difference between the total and homozygous load observed in small and large populations
(Figure 3C & D). To address this, we built unfolded site frequency spectra (SFS) using ANGSD[23],
based on alleles that have been assigned different impact ratings by the Variant Effect Predictor
(VEP)[24], *i.e.* low, modifier, moderate, and high (Figure 4, Supplementary Information). Comparing
SFS provides the opportunity to assess the impact of purifying selection on the full frequency range
of deleterious alleles in a population, not just those in heterozygous and homozygous states.
215
These SFS show an excess of low frequency deleterious alleles in mainland populations that have
not experienced population bottleneck(s) such as North Sulawesi (both babirusa and anoa) and
Southeast Sulawesi (anoa) (Figure 4A & B). This pattern is consistent with the effect of purging
reducing the frequency of deleterious alleles in the population. The persistence of deleterious
alleles at low frequency in these large, highly heterozygous, populations is likely due to their
recessive nature which means they are less likely to be exposed to selection.
222
In contrast, we found no noticeable excess of deleterious mutations at low frequency in Southeast
Sulawesi babirusa, likely due to a weaker effect of purging and lower heterozygosity than in the
North, in this population that experienced a recent bottleneck. Small island populations, particularly
Togean babirusa, however, possessed many fixed deleterious mutations, but much fewer
deleterious mutations at low frequency than in large populations. The high degree of fixed
deleterious alleles is likely the result of strong genetic drift leading to higher rate of fixation.
Reduced heterozygosity in these island populations likely explains the near absence of low-
frequency deleterious mutations, a result of long-term purging where recessive alleles are exposed
to selection. This is in contrast with large populations, in which deleterious alleles persist at low
frequency despite purging.
233
We then quantitatively compared the shape of these SFS across different impact categories using
a composite likelihood approach adapted from Nielsen *et al*[25]. Large differences of composite
likelihood values between SFS built using alleles with different levels of impact indicate substantial
differences between the shape of SFS. As the low impact alleles are less likely to be affected by
selection, we used the SNPs in this category as the expected SFS. This analysis shows that the
shape of the SFS of high and low impact mutations are substantially different in small island
populations, while very similar in mainland populations (Figure 4C & D).
241
The shape of the SFS across different impact categories in the small island populations suggests
that although they are smaller than their mainland counterparts, purging is having a stronger
distorting effect on the SFS of more impactful alleles. This is likely because recessive deleterious

alleles are more likely to be exposed to selection in a less heterozygous population. Over time, this
will lead to a reduction in frequency for the most deleterious alleles in small populations, consistent
with our total load scores (Figure 3A) and previous simulation studies[12].

248
**Conclusions**
Our analyses show that anoa and babirusa populations on small islands have remained sufficiently
stable to efficiently purge deleterious alleles. These populations also occupy high-quality habitats,
often within protected areas (e.g. 45% of the Togean islands are protected as a national park).
Small island ecosystems may thus offer a long-term solution to preserve these species, away from
areas such as Southeast Sulawesi where natural resources are more heavily exploited.

255
In contrast, we show that populations on the larger Island of Sulawesi, such as in Southeast
Sulawesi, occupy lower quality habitat due to a higher degree of anthropogenic disturbance, and
show signs of strong bottlenecks and an accumulation of deleterious alleles. These findings imply
that recent anthropogenic disturbances on larger islands may be reshaping the extinction dynamics
previously observed during the Quaternary, a period during which populations on smaller islands
have been generally presumed to be more vulnerable. Our results suggest a reversal of this trend,
indicating that human activities may be rendering larger island populations more susceptible to
extinction than their smaller counterparts.

264
The introduction of individuals from the mainland could be a solution to increase population size
and improve genetic diversity on small islands with high habitat quality and low anthropogenic
pressures. Our genomic data, however, indicate that individuals from mainland populations
possess more deleterious mutations, which, if translocated to smaller, less heterozygous, island
populations could result in fitness decline and increased risk of extinction[12]. We therefore suggest
that, in the case of babirusa and anoa, conservation efforts should focus on maintaining forest
habitat for these species on smaller islands, without the need for a possibly counterproductive, and
onerous, translocation programme. Altogether, our study demonstrates the benefit of combining
genomic information with species distribution modelling to help prediction of future anthropogenic
threats inform species conservation planning in island systems.

**Acknowledgement**
This research was funded by the NERC-Ristekdikti Newton Wallacea joint research programme
(grant number NE/S007067/1; Ristekdikti Grant No: NKB-1799/UN2.R3.1/HKP.05.00/2019
"Biodiversity, Environmental Change, and Land-use Policy in Sulawesi and Maluku"). We thank the
Indonesian Ministry of Environment and Forestry (KLHK), Sulawesi's Provincial Forestry
Departments (BKSDA), and Indonesia's National Research and Innovation Agency (BRIN, formerly
LIPI) for granting research and export permits. This work is a continuation of research under the
permit from Ministry of Forestry no 188/Kpts-IV/2001 and 2918/Kpts-IV/2002, now extended under
permit no. SK.18/KSDAE/SET-3/KSA-2/I/2024. This research was supported by computational and
data resources provided by the Leibniz Supercomputing Centre (LRZ; www.lrz.de). We thank
Athena Syarifa for her support during the start of the project, and Gabriel Renaud for assistance in
ROH analysis. SGA was supported by QMUL and GCRF. RD is supported by NERC and Humboldt
Foundation. MV, NJD and MJS were supported by a Leverhulme Research Leadership grant
Awarded to MJS. SB, SR, and LF were supported by NERC Newton Fund Wallacea. JB was
supported by University of Edinburgh, Rufford Grant, and Stichting Dierentuin Helpen Grant. PG
acknowledges the structural support of the Flemish government.

292
**Author contributions**

S.G.A., R.D., S.R., S.B., L.F.: Conceptualization; S.G.A., R.D., N.J.D., M.V., D.F., D.S., R.B., A.M.,
G.L., P.H.A.G., J.B., M.S.: Data Curation; S.G.A., R.D., A.C., L.F.: Formal Analysis; N.L.W., N.A.,
J.S., M.S., S.R., S.B., L.F.: Funding Acquisition; S.G.A., R.D., K.M., W.E.P., A.H.M, J.L.W., J.B.:
Investigation; S.G.A., R.D., A.C., N.J.D., M.V., M.F., D.S., D.G., M.S.: Methodology; S.G.A., R.D.,
D.S.P., N.L.W.. J.B., N.A., J.S., M.S., S.B., S.R., L.F.: Project Administration; M.S., D.S., K.M.,
W.E.P., J.L.W., R.B., A.M., G.L., P.H.A.G., J.B., M.S., L.F: Resources; S.G.A., R.D., A.C., N.J.D.,
M.F., D.G., L.F.: Software; S.G.A., R.D., S.B., S.R., L.F.: Supervision; S.G.A., R.D., A.C., L.F.:
Validation; S.G.A., R.D., L.F.: Visualization; S.G.A., R.D., S.R., S.B., L.F.: Writing - original draft; all
authors contribute to review and edit part of writing.
303
304 **Declaration of interests**
305 The authors declare no competing interests.
306
307 **Supplemental information**
308 **Document S1.** Figures S1–S4 and Table S1–S3
309 **Supplementary Data.** Excel file containing additional data (**Data S1-S3**)

**References**

1. Rozzi, R., Lomolino, M.V., van der Geer, A.A.E., Silvestro, D., Lyons, S.K., Bover, P., Alcover, J.A., Benítez-López, A., Tsai, C.-H., Fujita, M., et al. (2023). Dwarfism and gigantism drive human-mediated extinctions on islands. Science 379, 1054–1059.

2. Lynch, M., Conery, J., and Burger, R. (1995). Mutation Accumulation and the Extinction of Small Populations. Am. Nat.146 489-518.

3. Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., Chiari, Y., Dernat, R., Duret, L., Faivre, N., et al. (2014). Comparative population genomics in animals uncovers the determinants of genetic diversity. Nature 515, 261–263.

4. Russell, J.C., and Kueffer, C. (2019). Island biodiversity in the Anthropocene. Annu. Rev. Environ. Resour. 44, 31–60.

5. Voigt, M., Supriatna, J., Deere, N.J., Kastanya, A., Mitchell, S.L., Rosa, I.M.D., Santika, T., Siregar, R., Tasirin, J.S., Widyanto, A., et al. (2021). Emerging threats from deforestation and forest fragmentation in the Wallacea centre of endemism. Environ. Res. Lett. 16, 094048.

6. Exploring Energy and Mineral Resources of Indonesia, https://geoportal.esdm.go.id/minerba/.

7. Struebig, M.J., Aninta, S.G., Beger, M., Bani, A., Barus, H., Brace, S., Davies, Z.G., Brauwer, M.D., Diele, K., Djakiman, C., et al. (2022). Safeguarding imperilled biodiversity and evolutionary processes in the Wallacea center of endemism. BioScience 72, 1118–1130.

8. Frankham, R. (2008). Inbreeding and extinction: Island populations. Conserv. Biol. 12, 665–675.

9. Scott, P.A., Allison, L.J., Field, K.J., Averill-Murray, R.C., and Shaffer, H.B. (2020). Individual heterozygosity predicts translocation success in threatened desert tortoises. Science 370, 1086–1089.

10. Bell, D.A., Robinson, Z.L., Funk, W.C., Fitzpatrick, S.W., Allendorf, F.W., Tallmon, D.A., and Whiteley, A.R. (2019). The exciting potential and remaining uncertainties of genetic rescue. Trends Ecol. Evol. 34, 1070–1079.

11. Hedrick, P.W., Robinson, J.A., Peterson, R.O., and Vucetich, J.A. (2019). Genetics and extinction and the example of Isle Royale wolves. Anim. Conserv. *22*, 302–309.

12. Kyriazis, C.C., Wayne, R.K., and Lohmueller, K.E. (2021). Strongly deleterious mutations are a primary determinant of extinction risk due to inbreeding depression. Evol. Lett. 5, 33–47.

13. Bertola, L.D., Miller, S.M., Williams, V.L., Naude, V.N., Coals, P., Dures, S.G., Henschel, P., Chege, M., Sogbohossou, E.A., Ndiaye, A., et al. (2022). Genetic guidelines for translocations: Maintaining intraspecific diversity in the lion (*Panthera leo*). Evol. Appl. 15, 22–39.

14. Frantz, L.A.F., Rudzinski, A., Nugraha, A.M.S., Evin, A., Burton, J., Hulme-Beaman, A., Linderholm, A., Barnett, R., Vega, R., Irving-Pease, E.K., et al. (2018). Synchronous diversification of Sulawesi's iconic artiodactyls driven by recent geological events. Proc. Biol. Sci. 285, 20172566.

15. Hansen, M.C., Potapov, P.V., Moore, R., Hancher, M., Turubanova, S.A., Tyukavina, A., Thau, D., Stehman, S.V., Goetz, S.J., Loveland, T.R. and Kommareddy, A., 2013. High-resolution global maps of 21st-century forest cover change. Science 342, 850-853.

16. Renaud, G., Hanghøj, K., Korneliussen, T. S., Willerslev, E., and Orlando, L. (2019). Joint estimates of heterozygosity and runs of homozygosity for modern and ancient samples Genetics, 212, 587–614.

17. Schmitt, S., Pouteau, R., Justeau, D., De Boissieu, F. and Birnbaum, P. (2017) ssdm: An r package to predict distribution of species richness and composition based on stacked species distribution models. Methods Ecol. Evol. 8, 1795-1803.

18. Supriatna, J., Shekelle, M., Fuad, H.A.H., Winarni, N.L., Dwiyahreni, A.A., Farid, M., Mariati, S., Margules, C., Prakoso, B., and Zakaria, Z. (2020). Deforestation on the Indonesian island of Sulawesi and the loss of primate habitat. Glob. Ecol. Conserv. 24, e01205.

19. Rejeki, I.S. (2018). Wildlife conservation strategy: an assessment of wildlife hunting activities in Sulawesi. (Doctoral dissertation, IPB (Bogor Agricultural University)).

20. Robinson, J.A., Brown, C., Kim, B.Y., Lohmueller, K.E., and Wayne, R.K. (2018). Purging of strongly deleterious mutations explains long-term persistence and absence of inbreeding depression in island foxes. Curr. Biol. 28, 3487–3494.

21. Kleinman-Ruiz, D., Lucena-Perez, M., Villanueva, B., Fernández, J., Saveljev, A.P., Ratkiewicz, M., Schmidt, K., Galtier, N., García-Dorado, A., and Godoy, J.A. (2022). Purging

of deleterious burden in the endangered Iberian lynx. Proc. Natl. Acad. Sci. 119,
e2110614119.
22. Khan, A., Patel, K., Shukla, H., Viswanathan, A., van der Valk, T., Borthakur, U., Nigam, P.,
Zachariah, A., Jhala, Y.V., Kardos, M., et al. (2021). Genomic evidence for inbreeding
depression and purging of deleterious genetic variation in Indian tigers. Proc. Natl. Acad. Sci.
118, e2023018118.
23. Korneliussen, T.S., Albrechtsen, A. and Nielsen, R. (2014). ANGSD: Analysis of Next
Generation Sequencing Data. BMC Bioinform. 15, 1-13
24. McLaren, W., Gil, L., Hunt, S.E., Riat, H.S., Ritchie, G.R.S., Thormann, A., Flicek, P., and
Cunningham, F. (2016). The Ensembl Variant Effect Predictor. Genome Biol. 17, 1-14.
25. Nielsen, R., Williamson, S., Kim, Y., Hubisz, M.J., Clark, A.G., and Bustamante, C. (2005).
Genomic scans for selective sweeps using SNP data. Genome Res. 15, 1566–1575.

**Figure 1. Population structure and habitat suitability**. **(A)** anoa (copyright: chester zoo) and **(B)**
babirusa (copyright: S.L. Mitchell). **(C)** Principal component analysis (PCA) based on SNPS from
genomes (>5x) anoa (53 individuals; 1,053,534 SNPs) and **(D)** babirusa (37 individuals; 1,011,533
SNPs). **(E)** Zone of endemisms defined from Frantz *et al*[14] **(F)** Location of the samples of anoa **(G)**
babirusa, coloured based on regions of endemism. **(H)** Map of Sulawesi (including Buton and
Togean islands) with 2018 forest cover (Global Forest Change repository, v1.6, Hansen *et al*[15],
processed in Voight *et al*[5]) in green. **(I)** Ensemble distribution models constrained by forest cover,
categorised by the habitat suitability score quantile (class one = least suitable habitat, class five =
most suitable habitat) for anoa and **(J)** babirusa.
387

388    **Figure 2. Population-level genetic diversity and runs of homozygosity (ROH). (A)** Genome-
389    wide Watterson's $\theta$ (i.e. including ROH; square) and excluding ROH segments (triangle), **(B)** Sum
390    of size of ROH segments across the genome (NROH) and proportion of ROH across the genome
391    (FROH) **(C)** distribution of ROH segment length **(D)** Map of Sulawesi with different regions
392    considered in the analysis**.**
393

394    **Figure 3. Genomic load in babirusa, anoa and closely related species computed in**
395    **individual genomes using SIFT scores. (A)** Comparison of total load of anoa and babirusa to
396    their close relatives for alleles found at both heterozygous and homozygous states and **(B)** only
397    homozygous. **(C)** Total load within anoa and babirusa populations at both heterozygous and
398    homozygous states and **(D)** only homozygous.
399

**Figure 4. Site frequency spectra (SFS) of deleterious alleles across babirusa and anoa**
**populations. (A)** Site frequency spectra of alleles with different impact categories rating (low,
modifier, moderate, high) across populations of anoa and **(B)** babirusa. **(C)** Mean composite
likelihood (CL) values in anoa and **(B)** babirusa. Composite likelihood (CL) values represent
differences between the shape of the SFS computed using low impact mutation and the SFS built
using moderate, modifier and high impact alleles. Mean CL was computed using 1000 bootstrap
replicates and normalised using CL obtained from the SFS of low impact mutation (see
Supplementary Information).

Habitat class

| one | two | three | four | five |
|-----|-----|-------|------|------|
| <20% | 20-40% | 40-60% | 60-80% | 80-100% |