The importance of small island populations for the long-term survival of endangered large-bodied insular mammals.

Sabhrina Gita Aninta[1*#], Rosie Drinkwater[1,2*#], Alberto Carmagnini[2], Nicolas J. Deere[3], Dwi Sendi Priyono[4], Noviar Andayani[5], Nurul L. Winarni[6], Jatna Supriatna[5], Matteo Fumagalli[1], Greger Larson[7], Peter H.A. Galbusera[8], Alastair Macdonald[9], Deborah Greer[1], Kusdiantoro Mohamad[10], Wahono Esthi Prasetyaningtyas[10], Abdul Haris Mustari[11], John Lewis Williams[12], Ross Barnett[13], Darren Shaw[9], Gono Semiadi[14], James Burton[15], David J. I. Seaman[16], Maria Voigt[16], Matthew Struebig[16], Selina Brace[17], Stephen Rossiter[1], Laurent Frantz[1,2*].

[1]School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK
[2]Palaeogenomics group, Faculty of Veterinary Medicine, Ludwig Maximilian University Munich
[3]Durrell Institute of Conservation and Ecology (DICE), University of Kent, Canterbury, CT2 7NR, UK
[4]Department of Tropical Biology, Faculty of Biology, Universitas Gadjah Mada, Sleman, 55281, Special Region of Yogyakarta, Indonesia.
[5]Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Depok, 16424, Indonesia
[6]Research Center for Climate Change, University of Indonesia, Depok, Indonesia
[7]The Palaeogenomics and Bio-Archaeology Research Network, Research Laboratory for Archaeology and History of Art, University of Oxford, Oxford, UK
[8]Antwerp Zoo Centre for Research and Conservation, Royal Zoological Society of Antwerp, Antwerp, Belgium
[9]Royal (Dick) School of Veterinary Studies, The University of Edinburgh, Midlothian EH25 9RG
[10]School of Veterinary Medicine and Biomedical Sciences, IPB University, Bogor, Indonesia
[11]Department of Forest Resources Conservation, Faculty of Forestry and Environment, IPB University, Bogor, Indonesia
[12]Department of Animal Science, Food and Nutrition, Università Cattolica del Sacro Cuore, via Emilia Parmense, 84, 29122 Piacenza, Italy
[13]Durham Evolution and Ancient DNA, Department of Archaeology, University of Durham, Durham, United Kingdom
[14]Research Centre for Biosystematics and Evolution, National Research and Innovation Agency (BRIN), Cibinong 16911, Indonesia
[15]IUCN SSC Asian Wild Cattle Specialist Group
[16]Durrell Institute of Conservation and Ecology (DICE), University of Kent, Canterbury, CT2 7NR
[17]Department of Earth Sciences, Natural History Museum, London, England, United Kingdom, SW7 5BD, UK.
[#] Equally contributed

Corresponding authors: Sabhrina Gita Aninta, Rosie Drinkwater, Laurent Frantz.

E-mail: sagitaninta@gmail.com; r.drinkwater@lmu.de; laurent.frantz@lmu.de

Author Contributions: S.G.A., R.D., S.R., S.B., L.F.: Conceptualization; S.G.A., R.D., N.J.D., M.V., D.F., D.S., R.B., A.M., G.L., P.H.A.G., J.B., M.S.: Data Curation; S.G.A., R.D., A.C., L.F.: Formal Analysis; N.L.W., N.A., J.S., M.S., S.R., S.B., L.F.: Funding Acquisition; S.G.A., R.D., K.M., W.E.P., A.H.M, J.L.W., J.B.: Investigation; S.G.A., R.D., A.C., N.J.D., M.V., M.F., D.J.I.S., D.G., M.S.: Methodology; S.G.A., R.D., D.S.P., N.L.W.. J.B., N.A., J.S., M.S., S.B., S.R., L.F.: Project Administration; M.S., D.S., K.M., W.E.P., J.L.W., R.B., A.M., G.L., P.H.A.G., J.B., M.S., L.F: Resources; S.G.A., R.D., A.C., N.J.D., M.F., D.G., L.F.:

Software; S.G.A., R.D., S.B., S.R., L.F.: Supervision; S.G.A., R.D., A.C., L.F.: Validation;
S.G.A., R.D., L.F.: Visualization; S.G.A., R.D., S.R., S.B., L.F.: Writing - original draft; all
authors contribute to review and edit part of writing.

Competing Interest Statement: The authors declare no competing interests.

Keywords: conservation genomics, endemic species, genetic load, inbreeding, habitat
suitability modelling.

This PDF file includes:

Main Text
Figures 1 to 4
58

Significance statement
Within tropical archipelagos, such as the Wallacea biodiversity hotspot, larger islands
experience greater resource exploitation compared to smaller ones, highlighting the potential
of smaller islands as refuges for conservation. To investigate the genetic health of populations
on small islands, we used genomic, occurrence, and environmental data from a system of
replicated populations of anoa and babirusa across islands of varying sizes. In contrast to
larger islands like Sulawesi, our results demonstrate that smaller offshore islands not only
provide higher-quality habitats but also support populations that have efficiently purged
harmful mutations. Thus, despite their known vulnerability over geological time-frames, small
island populations can provide long term insurance against human-driven extinction and
conservation efforts should prioritise habitat management over translocations.
70
Abstract
Island populations of large vertebrates have experienced higher extinction rates than
mainland populations over long timescales due to demographic stochasticity, genetic drift and
inbreeding. Conversely, small island populations often experience relatively less
anthropogenic habitat degradation than populations on larger islands, making them potential
targets for conservation interventions such as genetic rescue to improve their overall genetic
diversity. Here we determine the consequences and conservation implications of long-term
isolation and recent human activities on genetic diversity of island populations of two forest-
dependent mammals endemic to the Wallacea archipelago: the anoa (*Bubalus* spp.) and
babirusa (*Babyrousa* spp.). Using genomic analyses and habitat suitability models, we show
that, compared to closely related species, populations on mainland Sulawesi exhibit low
heterozygosity, high inbreeding, and a high proportion of deleterious alleles. In contrast,
populations on smaller islands possess fewer deleterious mutations despite exhibiting lower
heterozygosity and higher inbreeding. Site frequency spectra indicate that these patterns
reflect stronger, long-term purging in smaller island populations. Our results thus suggest that
conservation efforts should focus on protecting small island habitats and avoiding
translocations from mainland populations. This study highlights the crucial role of small
offshore islands for the long-term survival of Wallacea's iconic and indigenous mammals in
the face of development on the mainland.

90
Introduction
Throughout the Quaternary period, island populations of large vertebrates have experienced
higher rates of extinction than their mainland counterparts[1]. This pattern can be attributed to
demographic stochasticity, as well as the strong effects of genetic drift that operate in small
populations of slow-reproducing species, and the associated negative consequences for
fitness and adaptability[2,3].
97
In the past century, these long-term threats to island populations have been further
compounded by human activities such as urbanisation, the introduction of invasive species,
land conversion for agriculture, mining, and hunting[4]. These anthropogenic pressures often
vary across islands, with smaller and more isolated populations typically facing less intense
exploitation compared to larger, more accessible ones[5,6]. Consequently, while populations on

small islands may be more prone to extinction over geological timescales, they can be less
affected by habitat degradation.
105
Among the strategies available to conservation managers of small populations, one of the
most valuable is genetic rescue. In this approach, individuals from nearby, larger populations
are translocated and introduced into a population to increase genetic diversity, thereby
reducing inbreeding depression, mitigating the impact of recessive deleterious alleles on
fitness, and potentially introducing beneficial alleles[7-9]. Although this procedure has achieved
notable successes, recent empirical studies[10] and simulations[11] suggest that it may not
always be suitable. While relocating individuals from larger populations can introduce
beneficial alleles and reduce inbreeding, it also risks introducing harmful recessive alleles,
potentially causing long-term negative consequences. This highlights the need for careful
genetic management[12], especially in historically small populations where natural selection has
likely eliminated many harmful alleles, but which still face fitness challenges due to inbreeding
and low genetic diversity.
118
The island archipelago of Wallacea in Indonesia provides a natural laboratory in which to
determine the combined effects of long-term isolation and recent human activity on island
population genetic structure and viability. Due to its long geological history of isolation,
Wallacea is a global hotspot for endemism. In recent years, it has emerged as a frontier
region for development in Indonesia, and its ecosystems and unique species face increasing
pressures from deforestation and mineral extraction[5,6,13]. To date, human activities have been
concentrated in the more accessible lowland areas on Sulawesi, whereas some smaller
offshore islands have remained less disturbed.
127
Here we determine the consequences and conservation implications of long-term isolation
and recent human activities on the genomic diversity of island populations of two forest-
dependent mammals, the anoa (*Bubalus depressicornis* and *B. quarlesi*) and babirusa
(*Babyrousa* spp.) (Figure 1). Both taxa that are endemic to Wallacea and are broadly co-
distributed on islands of different sizes across the archipelago. Previous work suggests they
underwent range expansions approximately 2 Mya[14], however, current populations are highly
fragmented following several decades of population declines and local extinctions due to
habitat loss and hunting.
136
As endangered and endemic mammals, the anoa and babirusa are the focus of ex-situ
breeding programmes and viewed as potential candidates for conservation programmes
aimed at increasing the size and genetic diversity of wild populations. Through a combination
of population genomic analyses and habitat suitability models, we demonstrate that small
island populations have persisted over long timeframes in high-quality habitats. In contrast,
some populations on the larger island (Sulawesi; Figure 1) have suffered from reduced
habitat quality due to intensive resource exploitation. Thus, conservation strategies for anoa
and babirusa should focus on preserving high-quality habitats on small islands and avoid
translocating individuals from degraded mainland populations. Our findings underscore the
critical role of small offshore islands in ensuring the long-term survival of Wallacea's iconic
mammals amidst ongoing land-use changes on the Sulawesi mainland.
148
Results and Discussion
We generated short-read genome sequence data from samples of anoa and babirusa
collected from across their respective ranges in Wallacea. To determine broad population
genetic structure, we mapped short reads of each taxon to a reference genome of a
conspecific, or, in the case of the anoa, the water buffalo (*Bubalus bubalis*). Using datasets of
3,762,790 unlinked SNPs for anoa, and 3,920,000 unlinked SNPs for babirusa, we performed
principal component analysis (Figure 1C), phylogenetic reconstructions, and admixture
analyses (Figure S2). These analyses revealed concordant patterns of population structure
across both taxa (Figure 1C), in which three distinct lineages were identified corresponding to
individuals from North Sulawesi, Southeast Sulawesi, and the small offshore islands of Buton
(anoa) or Togean (babirusa). This pattern of population structure, which broadly supports
previous findings based on mitochondrial and microsatellite data[14], was then used to define
populations in subsequent analyses.
162

Populations on smaller island are less genetically diverse than on mainland Sulawesi
To determine the consequences of long-term isolation, we quantified levels of genetic
diversity in populations on Sulawesi and smaller nearby islands (Buton and Togean). To do
so, we computed genome-wide Watterson's θ using ROHan[16], an unbiased estimator of
heterozygosity in a single diploid individual under the infinite sites model[17]. We found that θ
on the small islands of Togean (babirusa: mean=0.29 x 10^{-3}, SE=0.02 x 10^{-3}) and Buton
(anoa: mean=0.58 x 10^{-3}, SE=0.02 x 10^{-3}) was ~2 to ~6 fold lower than in populations from
mainland Sulawesi (babirusa: mean=1.85 x 10^{-3}, SE=0.1 x 10^{-3}; anoa: mean=1.33 x 10^{-3},
SE=0.1 x 10^{-3}), indicating that small island populations contain less genetic diversity than
populations on the larger island of Sulawesi (Figure 2A). On Sulawesi, however, we found
that babirusa from the Southeast Sulawesi possessed lower θ (mean=1.17 x 10^{-3}, SE=0.04 x
10^{-3}) compared to those found on the Northern peninsula (mean=2.17 x 10^{-3}, SE=0.06 x 10^{-3}).
175
We then tested whether anoa and babirusa populations are characterised by more variable
levels of genetic diversity compared to related species that do not occur on islands of varying
size. To address this, we compared levels of genetic variability (Watterson's θ) and
inbreeding (ROHs) among anoa and babirusa populations to those observed in 19 closely
related taxa (52 individuals in total), including highly endangered species such as the pygmy
hog (*Porcula salvianus*), the Javan warty pig (*Sus verrucosus*) and banteng (*Bos javanicus*).
Levels of genetic diversity (Watterson's θ) were on average lower in anoa than in other bovid
species (Figure 2A). The same was observed in babirusa, which aside from the North
Sulawesi population, possessed lower levels of genetic diversity than most other suids. Lower
levels of genetic diversity in the Wallacean endemics is likely the result of smaller effective
population sizes resulting from their island habitats and/or recent bottleneck(s).
187
To assess whether populations showed signs of recent bottlenecks, we analysed runs of
homozygosity (ROHs) using ROHan (Supplementary Information). Analyses of ROHs showed
that inbreeding levels and genetic diversity were more variable among anoa and babirusa
populations than in different species (Figure 2B). Small island populations of anoa (Buton)
and babirusa (Togean) possessed higher levels of inbreeding than most other species (Figure
2B). Buton contained longer ROHs than those sampled on mainland Sulawesi, consistent with
higher inbreeding levels on small offshore islands (Figure 2C).
195
Among suids, however, we found that Southeast Sulawesi babirusa possessed more ROHs
and had a larger portion of their genome in ROH than any other population of babirusa and
any other suid species (Figure 2B). Interestingly, despite this high level of inbreeding,
Southeast Sulawesi babirusa possessed average levels of genetic diversity when compared
to other suid species (Figure 2A). Their level of genetic diversity, however, was significantly
higher when excluding ROHs (p<0.001; Figure 2B). The significant change in genetic diversity
when excluding ROHs implies that babirusa living in Southeast Sulawesi were part of a
historically large population which underwent a recent bottleneck.
204
Similar levels of genetic diversity within and outside ROHs in anoa and babirusa living on
small islands (Togean and Buton) indicate they form part of a historically small, yet stable
population that inhabits a highly suitable habitat, as demonstrated by our species distribution
model (Figure 1D, Figure S4, Table S2). Altogether, this suggests that although they are not
as genetically diverse as their large island counterparts, small island populations are likely to
have been stable over geological time scales, and potentially since their expansion ~2Mya[14].
211
Impact of anthropogenic disturbances on demography
To address whether the dramatic bottleneck in Southeast Sulawesi babirusa is consistent with
a recent decrease of suitable habitat in the area, we constructed an ensemble species
distribution model[18] based on bioclimatic variables and forest cover[5,15] (Supplementary
Information). Our models, for both anoa and babirusa, indicate that there is proportionally
more suitable habitat (i.e. top 20% of all suitable habitat; see Supplementary Text) on smaller
islands (i.e. Togean and Buton) compared to mainland Sulawesi (~1.5 fold more highly
suitable area for anoa and ~1.6 for babirusa relative to North Sulawesi; Figure 1D, Table S3).
220
For both species, we related the distribution of suitable habitat to different land use classes
and found considerable overlap with areas that have protected status (Figure S4). This

highlights the effectiveness of the protected area network for conserving habitat for
Wallacea's endemic mammals. On the mainland, the availability of suitable habitat was higher
in the North compared to the Southeast for both taxa (Table S3). This trend is consistent with
recent deforestation rates (between 2000-2017), which were higher in Southeast Sulawesi
compared to the North[19], suggesting that deforestation has led to the reduced habitat
availability detected by our models. These data, combined with previously reported high rates
of poaching[20], imply that population bottlenecks in babirusa in Southeast Sulawesi have been
driven by relatively recent anthropogenic disturbances, rather than longer term evolutionary
processes.
232
More deleterious alleles segregate on Sulawesi than on small islands
Recent studies of several large-bodied mammals indicate that small populations with long
histories of isolation (*e.g.* Channel Island foxes[21], Iberian lynx[22], Indian tigers[23], mammoths[24])
show evidence of accumulation of mildly deleterious alleles, while naturally purging strongly
deleterious recessive variants. To evaluate how small island populations of anoa and
babirusa purge deleterious alleles compared to their mainland populations, we computed
genetic load in individual genomes, using three conservation scores: SIFT, PhyloP and
phastCons (Supplementary Information). To allow cross-species comparisons, we estimated
genetic load for loci that show one-to-one orthology between the pig and cow genomes
(Supplementary Information). We first computed a load score by summing conservation
scores, weighted by genotype probability for derived alleles found at homozygous states
across the genome (see Supplementary Methods). This homozygous load represents the
minimum impact of deleterious alleles on fitness (i.e. assumes that all deleterious alleles are
recessive). We also computed the load by summing the impact of alleles found at both
homozygous and heterozygous states (total load) across the genome. The total load
represents the maximum impact of deleterious alleles on fitness (i.e. assumes that all
deleterious alleles are dominant).
250
Both total and homozygous load were significantly lower in anoa than in babirusa (p<0.001;
see Supplementary Materials) (Figure 3A & B). Mean total load, calculated for all individuals
from the same population, was also significantly higher in large populations found on
mainland Sulawesi than in smaller island populations such as Togean (babirusa) and Buton
(anoa; Figure 3C, Table S1). This result indicates that deleterious alleles are more abundant
in larger, mainland, populations than in smaller, island populations.
257
Mean homozygous load, however, was significantly higher in small island populations than in
larger island populations (Figure 3D). Altogether these results indicate that large populations
possess more deleterious recessive alleles overall, most of which are found in heterozygous
state and therefore hidden from selection. Thus, although large populations possess more
deleterious alleles, the overall fitness impact of deleterious alleles is likely to be higher in
smaller island populations which possess more deleterious alleles in homozygous state.
264
Small island populations efficiently purge deleterious alleles
The difference between the total and homozygous load observed in small and large
populations could be due to either lower levels of heterozygosity, or the lower efficiency of
purifying selection in small populations (Figure 2B). To assess the efficiency of purifying
selection across different populations, we first built unfolded site frequency spectra (SFS)
using ANGSD[25], based on alleles that have been assigned different impact ratings by the
Variant Effect Predictor (VEP)[26], *i.e.* low, modifier, moderate, and high (Figure 4,
Supplementary Information). Comparing SFS provides the opportunity to assess the impact of
purifying selection on the full frequency range of deleterious alleles in a population, not just
those in heterozygous and homozygous states.
275
These SFS show an excess of low frequency deleterious alleles in mainland populations that
have not experienced bottleneck(s), such as populations of anoa in Southeast Sulawesi
(Figure 4A & B) and of both taxa in North Sulawesi. This pattern is consistent with the effect
of purging reducing the frequency of deleterious alleles in a population[21-23]. The persistence of
deleterious alleles at low frequency in these large, highly heterozygous, populations is likely
due to their recessive nature which means they are less likely to be exposed to selection than
in smaller, less heterozygous, populations

283
In contrast, we found fewer deleterious alleles at low frequency in the recently bottlenecked
Southeast Sulawesi babirusa, compared to in the North Sulawesi population. This is likely due
to a weaker effect of purging, and stronger drift in Southeast Sulawesi babirusa than in the
North Sulawesi. Both small island populations, however, possessed many fixed deleterious
alleles, but had fewer deleterious alleles at low frequency than in large populations and in the
recently bottlenecked Southeast babirusa population. The high degree of fixed deleterious
alleles is likely to be the result of strong genetic drift in small populations. Reduced
heterozygosity in these island populations likely explains the near absence of low-frequency
deleterious alleles, a result of long-term purging where recessive alleles are exposed to
selection. This is in contrast with large populations, in which deleterious alleles persist at low
frequency despite purging.
295
To quantitatively compare the shape of the SFS across four impact ratings, we used a
composite likelihood (CL) approach adapted from Nielsen *et al*[27]. Large differences in CL
values across impact ratings point to contrasting shapes of the SFS. As the low impact alleles
are less likely to be affected by selection compared to the other three impact ratings (modifier,
moderate, high), we used the SNPs in this rating as the expected SFS. For each impact rating
we obtained 1000 CL values across bootstrap replicates (see supplementary methods), which
were normalised using the mean of the bootstrap value of the low impact CL. In large island
populations, normalised CL distributions (across bootstrap replicates) of the three higher
impact ratings were centred around one (Figure 4A, B, D, E). This indicates that the SFS of
the higher impact ratings were not quantitatively different from that of the SFS built using low
impact ratings. In contrast, in small island populations, the mean CL across the three higher
impact ratings was higher than one, indicating that the shape of the SFS of the higher impact
alleles are quantitatively different from the shape of the SFS built using low impact alleles
(Figure 4C & D).
310
These differences in the shape of the SFS across different impact ratings in the small island
populations suggests that although they are smaller than their mainland counterparts, purging
is having a stronger distorting effect on the SFS of more impactful alleles. This could be the
result of recessive deleterious alleles that are more likely to be exposed to selection in a less
heterozygous population. Over time, this will lead to a reduction in frequency for the most
deleterious alleles in small populations, consistent with our total load scores (Figure 3A) and
previous simulation studies[11].
318
Conclusions
Our analyses show that anoa and babirusa populations on small islands have remained
sufficiently stable to efficiently purge deleterious alleles. These populations also occupy high-
quality habitats, often within protected areas (e.g. 45% of the Togean islands are protected as
a national park). Small island ecosystems may thus offer a long-term solution to preserve
these species. In contrast, we show that some populations on the larger island of Sulawesi,
such as in the Southeast, occupy lower quality habitat due to a higher degree of
anthropogenic disturbance, and show signs of strong bottlenecks and weaker purifying
selection. These findings imply that recent anthropogenic disturbances on larger islands may
be reshaping the extinction dynamics during the Quaternary, a period during which
populations on smaller islands have been generally presumed to be more vulnerable. Our
results suggest a reversal of this trend, which when combined with a greater human impact
could make large island populations more susceptible to extinction than their smaller island
counterparts.
333
The introduction of individuals from the mainland could be a solution to increase genetic
diversity on small islands with high habitat quality and low anthropogenic pressures. Our
genomic data, however, indicate that individuals from mainland populations possess more
deleterious alleles, which, if translocated to smaller, less heterozygous, island populations
could result in fitness decline and increased risk of extinction[10]. We therefore suggest that,
unless the population size of babirusa and anoa drops dramatically on smaller islands,
conservation efforts should focus on maintaining forest habitat, without the need for a possibly
counterproductive, and onerous, translocation programme. Translocations, however, could
also become useful in this case if the degree of fixed deleterious alleles, which is higher in

smaller island populations (Figure 4), starts having a strong impact on fitness, as purifying
cannot remove these alleles once they are fixed, or to enhance immune function by
introducing new alleles. Altogether, our study demonstrates the benefit of combining genomic
information with species distribution modelling to help predict future anthropogenic threats
and inform species conservation planning for island systems.
348
Materials and Methods
Samples used for this study were obtained from a previous study[14] totalling to 67 anoa (29
from North Sulawesi, 15 from Southeast Sulawesi, and 23 from Buton) and 46 babirusa (24
from North Sulawesi, 14 from Southeast Sulawesi, and 8 from Togean). Detailed descriptions
of the samples used for this study are presented in SI Appendix. DNA was extracted either
from hair follicles from tails or tissue scraps from skeletal remains using the DNeasy Blood
and Tissue kit (Qiagen) with the final extract eluted in 100 µL of TE buffer. Double indexed
standard illumina libraries were built by Novogene (in 2020) or Macrogen (in 2021) (Data S1).
Libraries were pooled equimolarly and sequenced on a Illumina Novaseq S4 platform (150 bp
PE). To compare with other taxa that are closely related, we downloaded 25 genomes from
European Nucleotide Archive (ENA) and 26 genomes from Sequence Read Archive (SRA)
comprising 8 species of Suidae and 10 species of Bovidae (Data S2) that were also
sequenced using Illumina (150 bp PE). Each paired end fastq files sample was trimmed with
AdapterRemoval[28] and aligned to the using the BWA MEM[29] to a closely related reference
genome, i.e. water buffalo and babirusa, and distantly related reference genome, i.e. cow and
pig, for anoa and babirusa, respectively, constructing each to a set of close relative alignment
and distant relative alignment (see SI Text). Other than the population structure and genetic
load analyses that use the distant relative alignment to get gene annotation information, all
downstream analyses were conducted using the close relative alignment. Only genomes with
mean reads depth of at least 5x were analysed. Habitat suitability models were generated
using anoa and babirusa occurrence data and environmental covariates (see SI text).
Ensemble distribution models[30] were generated and analysed in R[31] and QGIS[32]. Further
details for computational analyses used in this study are provided in SI Appendix.
372

Acknowledgement
This research was funded by the NERC-Ristekdikti Newton Wallacea joint research
programme (grant number NE/S007067/1; Ristekdikti Grant No: NKB-
1799/UN2.R3.1/HKP.05.00/2019 "Biodiversity, Environmental Change, and Land-use Policy
in Sulawesi and Maluku"). We thank the Indonesian Ministry of Environment and Forestry
(KLHK), Sulawesi's Provincial Forestry Departments (BKSDA), and Indonesia's National
Research and Innovation Agency (BRIN, formerly LIPI) for granting research and export
permits. This work is a continuation of research under the permit from Ministry of Forestry no
188/Kpts-IV/2001 and 2918/Kpts-IV/2002, now extended under permit no.
SK.18/KSDAE/SET-3/KSA-2/I/2024. This research was supported by computational and data
resources provided by the Leibniz Supercomputing Centre (LRZ; www.lrz.de). We thank
Athena Syarifa for her support during the start of the project, and Gabriel Renaud for
assistance in ROH analysis. SGA was supported by QMUL and GCRF. RD is supported by
NERC and Humboldt Foundation. MV, NJD and MJS were supported by a Leverhulme
Research Leadership Grant Awarded to MJS. SB, SR, and LF were supported by NERC
Newton Fund Wallacea. JB was supported by University of Edinburgh, Rufford Grant, and
Stichting Dierentuin Helpen Grant. PG acknowledges the structural support of the Flemish
government.
391
392 **Data sharing**
393 All sequences used in this study are stored in European Nucleotide Archive accession
394 ERS10527733-ERS10527854. All codes used for analyses in this paper is openly available in
395 https://github.com/sagitaninta/Wallacea (see Supplementary Information)
396

References
1. R. Rozzi *et al.*, Dwarfism and gigantism drive human-mediated extinctions on islands.
Science **379**, 1054–1059 (2023).
2. M. Lynch, J. Conery, R. Burger, Mutation accumulation and the extinction of small
populations. *Am. Nat.* **146**, 489–518 (1995).

3. J. Romiguier *et al.*, Comparative population genomics in animals uncovers the
determinants of genetic diversity. *Nature* **515**, 261–263 (2014).
4. J.C. Russell, C. Kueffer, Island biodiversity in the Anthropocene. *Annu. Rev. Environ.*
Resour. **44**, 31–60 (2019).
5. M. Voigt, et al., Emerging threats from deforestation and forest fragmentation in the
Wallacea centre of endemism. *Environ. Res. Lett.* **16**, 094048 (2021).
6. M. J. Struebig, et al., Safeguarding Imperiled Biodiversity and Evolutionary Processes in
the Wallacea Center of Endemism. *BioScience* **72**, 1118–1130 (2022).
7. R. Frankham, Genetic rescue of small inbred populations: meta-analysis reveals large
and consistent benefits of gene flow. *Mol. Ecol.* **24**, 2610–2618 (2015).
8. P.A. Scott, L.J. Allison, K.J. Field, R.C. Averill-Murray, H.B Shaffer, Individual
heterozygosity predicts translocation success in threatened desert tortoises. *Science*
370, 1086–1089 (2020).
9. K. Ralls, P. Sunnucks, R. C. Lacy, R. Frankham, Genetic rescue: A critique of the
evidence supports maximizing genetic diversity rather than minimizing the introduction of
putatively harmful genetic variation. *Biological Conservation* **251**,108784 (2020).
10. P.W. Hedrick, J.A. Robinson, R.O. Peterson, J.A. Vucetich, Genetics and extinction and
the example of Isle Royale wolves. *Anim. Conserv.* **22**, 302–309 (2019).
11. C.C. Kyriazis, R.K. Wayne, K.E. Lohmueller, Strongly deleterious mutations are a
primary determinant of extinction risk due to inbreeding depression. *Evol. Lett.* **5**, 33–47
(2021).
12. L.D. Bertola *et al.*, Genetic guidelines for translocations: Maintaining intraspecific
diversity in the lion (Panthera leo). *Evol. Appl.* **15**, 22–39 (2022).
13. Exploring Energy and Mineral Resources of Indonesia,
https://geoportal.esdm.go.id/minerba/
14. L.A.F. Frantz *et al.*, Synchronous diversification of Sulawesi's iconic artiodactyls driven
by recent geological events. *Proc. Biol. Sci.* **285**, 20172566 (2018).
15. M.C. Hansen *et al.*, High-resolution global maps of 21st-century forest cover change.
Science **342**, 850–853 (2013).
16. G. Renaud, K. Hanghøj, T.S. Korneliussen, E. Willerslev, L. Orlando, Joint estimates of
heterozygosity and runs of homozygosity for modern and ancient samples. *Genetics* **212**,
587–614 (2019).
17. Watterson G A, On the number of segregating sites in genetical models without
recombination. *Theor. Popul. Biol.* **7**: 256–276. (1975)
18. S. Schmitt, R. Pouteau, D. Justeau, F. De Boissieu, P. Birnbaum, ssdm: An R package
to predict distribution of species richness and composition based on stacked species
distribution models. *Methods Ecol. Evol.* **8**, 1795–1803 (2017).
19. J. Supriatna *et al.*, Deforestation on the Indonesian island of Sulawesi and the loss of
primate habitat. *Glob. Ecol. Conserv.* **24**, e01205 (2020).
20. I.S. Rejeki, Wildlife conservation strategy: An assessment of wildlife hunting activities in
Sulawesi. (Doctoral dissertation, IPB (Bogor Agricultural University), 2018).
21. J.A. Robinson, C. Brown, B.Y. Kim, K.E. Lohmueller, R.K. Wayne, Purging of strongly
deleterious mutations explains long-term persistence and absence of inbreeding
depression in island foxes. *Curr. Biol.* **28**, 3487–3494 (2018).
22. D. Kleinman-Ruiz *et al.*, Purging of deleterious burden in the endangered Iberian lynx.
Proc. Natl. Acad. Sci. **119**, e2110614119 (2022).
23. A. Khan *et al.,* Genomic evidence for inbreeding depression and purging of deleterious
genetic variation in Indian tigers. *Proc. Natl. Acad. Sci.* **118**, e2023018118 (2021).
24. M. Dehasque. *et al.*, Temporal dynamics of woolly mammoth genome erosion prior to
extinction, *Cell.* **187**, 3531-3540.e13 (2024).
25. T.S. Korneliussen, A. Albrechtsen, R. Nielsen, ANGSD: Analysis of next generation
sequencing data. *BMC Bioinform.* **15**, 1–13 (2014).
26. W. McLaren *et al.*, The Ensembl variant effect predictor. *Genome Biol.* **17**, 1–14 (2016).
27. R. Nielsen *et al.*, Genomic scans for selective sweeps using SNP data. *Genome Res.*
15, 1566–1575 (2005).
28. Schubert, M., Lindgreen, S., & Orlando, L. (2016). AdapterRemoval v2: rapid adapter
trimming, identification, and read merging. *BMC Res. Notes*, **9**, 1-7.
29. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM,
Preprint at arXiv 1303.3997 (2013).

30. S. Schmitt, R. Pouteau, D. Justeau, F. De Boissieu, P. Birnbaum, ssdm: An R package to
462   predict distribution of species richness and composition based on stacked species
463   distribution models. *Methods Ecol. Evol.* **8**, 1795–180 (2017).
464   31. R Core Team, R: A Language and Environment for Statistical Computing, Vienna, Austria:
465   R Foundation for Statistical Computing, available at: https://www.r-project.org/ (2024).
466   32. QGIS.org (2021). QGIS Geographic Information System. QGIS Association.
467   http://www.qgis.org.

470
Figure 1. Population structure and habitat suitability. **(A)** Photographs of anoa (top, credit: Chester Zoo) and babirusa (bottom, photo credit: S.L. Mitchell). **(B)** Locations of samples for anoa (left), babirusa (right) coloured based on population structure (Figure S2). **(C)** Zones of endemism (right) based on Frantz *et al*[14] (left) Principal component analysis (PCA) based on SNPs from genomes (>5x) for anoa (middle, 53 individuals; 1,053,534 SNPs) and babirusa (right, 37 individuals; 1,011,533 SNPs), **(D)** Sulawesi, and offshore islands, showing the 2018 forest cover (Global Forest Change repository, v1.6, Hansen *et al*[15], processed in Voight *et al*[11]) in green used to constrain the ensemble distribution models of habitat suitability for anoa (middle) and babirusa (right) categorised by the habitat suitability score quantile (class one = least suitable habitat, class five = most suitable habitat.

Figure 2. Population-level genetic diversity and runs of homozygosity (ROHs).
(A) Genome-wide Watterson's θ computed across the whole genome (triangle) and
excluding ROH (square) in anoa, babirusa, and their respective close relatives, **(B)**
Sum of size of ROH segments across the genome (NROH) and proportion of ROH
across the genome (FROH) **(C)** distribution of ROH segment length.

490
Figure 3. Genetic load in babirusa, anoa and closely related species. (A) Total
load computed in individual genomes using SIFT scores for alleles found at both
heterozygous and homozygous states in anoa, babirusa, and their close relatives.
(B) Homozygous load computed in individual genomes using SIFT scores for alleles
found at homozygous states in anoa, babirusa, and their close relatives. **(C)** Inter-
island comparison of total load and **(D)** homozygous load within the populations of
anoa and babirusa.
498
499

501
Figure 4. Site frequency spectra (SFS) of deleterious alleles across anoa and babirusa populations. Site frequency spectra of alleles with four different impact ratings (low, modifier, moderate, high) across populations of anoa **(A-C)** and babirusa **(D-F)** with a corresponding 1000 bootstrap of composite likelihood (CL) values within each SFS graph representing differences between the shape of the SFS across the four impact ratings using low impact rating as the expected likelihood.