Strength of low-frequency EEG phase entrainment to external stimuli is associated with fluctuations in the brain's internal state

Verónica Mäki-Marttunen[1], Alexandra Velinov[1], Sander Nieuwenhuis[1]

[1]Cognitive Psychology Unit, Faculty of Social Sciences, Leiden University, Wassenaarseweg 52 2333 AK, Leiden, Netherlands.

Abstract

The brain attends to environmental rhythms by aligning the phase of internal oscillations. However, the factors underlying fluctuations in the strength of this phase entrainment remain largely unknown. In the present study we examined whether the strength of low-frequency EEG phase entrainment to rhythmic stimulus sequences varied with pupil size and posterior alpha-band power, thought to reflect arousal level and excitability of posterior cortical brain areas, respectively. We recorded pupil size and scalp EEG while participants carried out an intermodal selective attention task, in which they were instructed to attend to a rhythmic sequence of visual or auditory stimuli and ignore the other perceptual modality. As expected, intertrial phase coherence (ITC), a measure of entrainment strength, was larger for the task-relevant than for the task-irrelevant modality. Across the experiment, pupil size and posterior alpha power were strongly linked with each other, and ITC tracked pupil size: with increasing pupil size, entrainment was higher to the task-relevant stimulus sequence and lower to the task-irrelevant stimulus sequence. Exploratory analyses showed that a temporal relation between ITC and posterior alpha power emerged in the time periods around pupil peaks and pupil troughs. These results indicate that endogenous sources contribute distinctly to the fluctuations of EEG phase entrainment.

Significance statement

Fluctuations in cortical state powerfully shape the perception of external stimuli. Understanding the physiological signatures of cortical state fluctuations is crucial to understand how the brain selectively attends and switches between internal and external content. Here we studied how two signatures of

attentional state, pupil-linked arousal and power in the alpha band, shape the entrainment of brain activity to low-frequency rhythmic stimuli. Our results disclose common and dissociable influences of these signatures at slow time scales. Furthermore, measuring and including pupil size as a covariate in statistical models can help increase statistical power in studies focusing on EEG phase entrainment. Our study provides new evidence on a direct influence of cortical state on the perception of rhythmic stimuli.

Introduction

How we perceive the continuous stream of external sensory stimulation is highly modulated by the internal state of the brain (Matthewson et al., 2012), but the interaction between external and internal mechanisms of attention is still poorly understood. Rhythmic stimuli entrain endogenous oscillatory activity in the brain to the temporal structure of the stimulus sequence (Van Rullen et al. (2014), Ten Oever et al., 2017; Zoefel et al., 2018; Bauer et al., 2018). This *neural entrainment* results in the alignment of the high-excitability oscillatory phase with the predicted onset of the rhythmic stimuli (Lakatos et al., 2008; Stefanics et al., 2010), and provides an advantage for the processing of natural stimuli that are inherently rhythmical, such as speech, music and repetitive motion (Henry and Obleser, 2012; Lakatos et al. 2019). Although phase alignment of oscillations to external rhythms constitutes an important principle of neural dynamics and attention, little is known about the factors that make the strength of phase entrainment fluctuate over time.

Here, we investigated the impact on low-frequency phase entrainment of two physiological measures known to be highly variable across time: pupil-linked arousal and posterior alpha-band power. Attention during tasks requiring sustained performance commonly fluctuates due to variations in the internal state of vigilance or arousal (Unsworth and Robinson, 2017). Pupil size has long been used to measure variations in arousal level in humans (Bradshaw, 1967). Very low or very high levels of vigilance, as characterized by lapses of attention or task-unrelated thoughts, often occur during periods of small and/or large (pre-trial) baseline pupil diameter (van den Brink et al., 2016; Unsworth and Robinson, 2016; Madore et al., 2020). Whether and how arousal modulates the strength of neural entrainment, which is an externally directed process, is still unknown.

Changes in posterior alpha-band power, as measured with electroencephalography (EEG), are thought to reflect the excitability of posterior cortical brain areas that process external (sensory) input (Romei et al., 2008; Samaha et al., 2016). Increased alpha power is suggested to facilitate internally directed attention rather than attention to external stimuli (Cooper et al., 2003; Klimesch et al., 200; Benedek et al., 2014). Lapses of attention and task-unrelated thoughts are associated with increases in pre-trial

posterior alpha power (O'Connell et al., 2009; Compton et al., 2019; Jin et al., 2019; Madore et al., 2020, Groot et al., 2021). Similarly, a study in nonhuman primates found that periods with high alpha power were characterized by attentional lapses and strongly diminished low-frequency entrainment of neuronal oscillations to task-relevant rhythmic stimulus sequences (Lakatos et al., 2016). These results suggest that alpha power is an index of internal state and would be expected to anti-correlate with the strength of entrainment to a task-relevant sequence of stimuli.

Although baseline pupil size and posterior alpha have both been associated with attentional state, their interrelation may not be straightforward. Alpha power as a measure of inattention, as reviewed above, would be expected to correlate negatively with pupil size. That is, low arousal would be related to distraction and an increase in the "idling" alpha. However, studies on the relationship between pupil size and posterior alpha power at rest or in pre-trial periods have found a positive relation and/or an inverted-U relationship (Hong et al., 2014; van Kempen et al., 2019; Ceh et al., 2020; Podvalny et al., 2021; Pfeffer et al., 2022, but see Waschke et al., 2019). Therefore, these two signals may reflect brain states that are at least partly dissociable, and we expected that they would distinctly influence neural entrainment.

In this study, we aimed to test whether and how low-frequency phase entrainment is susceptible to modulations in internal state, as indexed by pupil-linked arousal and posterior alpha power. We simultaneously recorded pupil size and scalp EEG to investigate the relationships between pupil-linked arousal, cortical alpha oscillations and low-frequency phase entrainment during an intermodal attention task. In the task, rhythmic streams of visual and auditory stimuli were concurrently presented. Participants had to detect infrequent targets in the cued modality and ignore the other modality. A small difference in repetition rate (visual, 1.1 Hz, auditory, 1.4 Hz) allowed us to separately assess entrainment to the task-relevant and task-irrelevant stimulus sequences. We hypothesized that transient changes in the strength of entrainment to the attended modality would be related to fluctuations in the arousal level of the participants. We also expected that internal neural processes signaled by alpha power would

modulate entrainment level. This study brings novel insights into the interrelation between external and internal attentional mechanisms of the brain.

Methods

Participants

Thirty-seven students recruited at the author's institution took part in the experimental study. One participant was excluded because the recording was interrupted. EEG data from four participants were excluded from analysis due to excessive artifacts (> 50% of epochs rejected) in the Attend Visual condition. One participant was excluded because of low accuracy (hit rate < 0.5) in the Attend Auditory condition. Five out of the first 15 participants were excluded because they reported an unforeseen visual illusion, where they perceived the afterimage of the green fixation stimulus having a similar color as the magenta targets, probably caused by a complementary afterimage effect (Manzotti, 2017). As a precaution to avoid this illusion, after the first 15 participants we changed the color stimulus, as reported below. Performance did not differ between either set of stimuli for the participants included in the final study (hit rate, first round: 0.77 ± 0.05, second round: 0.79 ± 0.07, T = 0.86, p = 0.398; incorrect responses, first round: 32 ± 30, second round: 38 ± 38, T = 0.41, p = 0.685). The final sample consisted of 26 participants (mean age: 23, age range: 18-28; 22 female). Study exclusion criteria included any psychiatric disorders and wearing contact lenses. The experimental protocol was reviewed and approved by the ethics committee of the Authors' University. Participants signed a consent form prior to participating and received a compensation of 7.5 euros/hour or course credits.

Task and stimuli

Participants performed an intermodal selective attention task (**Fig. 1**), based on that used by Lakatos et al. (2016) to study phase entrainment in monkeys. Participants were presented with simultaneous streams of visual and auditory stimuli and had to respond as quickly as possible to rare target stimuli in the cued modality by pressing the "B" key on a keyboard, and to otherwise withhold responding. The response deadline was 1 second after target onset. The task consisted of ten blocks, each lasting

approximately five minutes. The participants were prompted at the beginning of each block to attend to the visual stimuli (Attend Visual condition) or to the auditory stimuli (Attend Auditory condition), and ignore the other modality. Half of the blocks required the participant to attend to the visual modality, the other half to the auditory modality, in a quasi-random order (A1-V1-A2-A3-V2-A4-V3-V4-A5-V5).

Figure 1. Multimodal task. a) Toy example of a sequence of trials of the intermodal selective attention task. Rectangles represent trials of the auditory modality (tones, represented in blue) and the visual modality (gabor patches, represented in orange). Auditory and visual stimuli were presented simultaneously at slightly different frequencies (auditory: 1.4 Hz, visual: 1.1 Hz). Participants had to attend to only one modality per block, and respond to the target stimuli of that modality (represented in lighter color) with a button press. In this example, the participant responds to the target stimulus in the visual modality. b) Illustration of the calculation of ITC over auditory (ITC_A) and visual (ITC_W) trials in 8-seconds time windows (W1, W2, W3, etc) and 2-seconds steps.

In each block 605 stimuli were presented: 265 visual stimuli and 340 auditory stimuli. In each stream of stimuli, 15% of the stimuli were targets and 85% were standards. The auditory stimuli consisted of short beeps with a 460-Hz pitch for the standards and a 476-Hz pitch for the targets. The visual stimuli consisted of colored RGB Gabor patches (orange standards: RGB 229 133 126, magenta targets: RGB 229 90 172, green fixation point for participants #1-15: RGB 151 199 62; purple fixation point for participants #16-37: RGB 180 137 205) with a 10-degree orientation and a frequency of 0.1 cycles/pixel, created using an online Gabor-patch generator (https://www.cogsci.nl/gabor-generator). Visual stimuli

were made isoluminant with the grey screen background (RGB 167 167 167) using the SHINEcolor toolbox (Dal Ben, 2023). The visual and auditory stimuli were presented at average frequencies of 1.1 Hz and 1.4 Hz, respectively. All stimuli were presented for 50 ms. The fixation point was on the screen in the interval between consecutive visual stimuli. The average interstimulus interval was 387 ms, with a range from 0 (i.e., auditory and visual stimuli presented simultaneously) to 727 ms. To lower a possible contribution of stimulus-evoked EEG responses to our measure of intertrial phase coherence, we added to the interstimulus interval some decorrelated jitter, drawn on each trial from the values [-66, -33, 0, 33, 66 ms]. Entrainment occurs at the average (i.e., expected) frequency of presentation of the stimuli, and thus is independent of the stimulus presentation itself (Besle et al., 2011).

Procedure

Participants were instructed to sit as still as possible throughout the blocks and to always look at the center of the screen, also in the Attend Auditory condition. Their chins were positioned in a chin rest, with their eyes at approximately 73 cm from a monitor with a refresh rate of 60 Hz. For the auditory stimuli custom in-ear air-tube headphones were used to prevent electrical artifacts in the EEG recordings. Participants first performed two practice blocks, one for each condition, which each lasted 2 minutes and contained a total of 256 visual and auditory stimuli. After the practice blocks, the eye-tracker was calibrated and the participants performed the task in a dimly lit room. The task was controlled using Eprime 3.0 with the appropriate Tobii extensions installed. In between blocks, participants could take breaks and decide when to start the following block. Lighting conditions and screen luminance were kept constant across the whole session. Participants took between 60 and 70 minutes to complete the experiment.

Pupillometry

The pupil diameter of both eyes was recorded using a Tobii Pro X3-120 eye-tracker with a sampling rate of 40 Hz. The calibration was performed at the beginning of the task using a 5-point fixation procedure.

The Tobii device automatically marked periods when no pupil signal was recorded (e.g., blinks) and custom code was compiled to save accurate timing data that could then be used to create eye-tracking epochs and segmentations for the behavioral data. Raw pupil data were preprocessed using the PhysioData Toolbox v0.5.0 (Kret and Sjak-Shie, 2019) using the standard pipeline of the Pupil Diameter Analyzer module. One participant was excluded from the pupil analyses due to excessive data loss during preprocessing (> 50% of recording). An exponential decay in pupil size was observed at the beginning of the blocks for most participants. Since this would affect our measures of pupil-linked arousal, we fitted an exponential decay curve to the pupil time series using the MATLAB R2018 (MathWorks, Inc., Natick, MA) functions *fittype* and *fit*, and regressed out this component from the pupil data. For the time windows analyses, the continuous pupil signal was cut into 8-second windows (2-second step size) and then averaged across each window. The resulting time series was *z*-scored separately for each block to make sure that differences between trials in pupil size reflected local fluctuations in pupil size instead of global time-on task effects (Hopstaken et al., 2015).

EEG acquisition and preprocessing

EEG data were recorded using a BioSemi ActiveTwo system with 32 active Ag-AgCl electrodes, at a sampling rate of 512 Hz. The electrodes were arranged according to the international 10-20 system. In addition to the 32 main electrodes, two reference electrodes were positioned on the mastoids, two facial electrodes above and below the left eye and two electrodes on the temples. Raw EEG data were preprocessed in EEGLAB (Delorme and Makeig 2004). The signal was re-referenced to the mastoids and high-pass filtered (0.5 Hz). Bad channels were removed, with the rejection threshold for channel correlation set to the default of 0.8 and the artifact subspace reconstruction parameter set to 10. Missing channels were interpolated from neighboring sites. Then the data were split into Attend Auditory and Attend Visual conditions and epoched into 3-second segments centered on stimulus onset. Bad epochs were rejected automatically (threshold criteria: 150 uV) and an independent component analysis (infomax algorithm with *extended* option to extract sub-Gaussian sources) was performed to identify and remove noise components manually (e.g., eye artifacts, muscle artifacts). Preprocessing

was done using the Neuroscience Gateway (https://www.nsgportal.org, Sivagnanam et al., 2013, Martínez-Cancino et al., 2021).

<u>Intertrial phase coherence and prestimulus alpha power</u>

The cleaned EEG data were used to calculate measures of phase entrainment and alpha power. We first extracted spectral power and phase dynamics using Morlet wavelet decomposition from 0.5 to 30 Hz with 30 logarithmically spaced steps. To calculate the entrainment of EEG oscillations to the stimulus rhythm, we took the 3-second segments mentioned above and computed for each 8-second time window the intertrial phase coherence (ITC), a measure of the consistency across trials of the phase angle at the time of stimulus onset (Tallon-Baudry et al., 1996, **Fig. 1b**). This was done separately for the visual and auditory stimulus streams. For these analyses, we averaged data from electrodes displaying the largest ITC for each modality: AF3, AF4, F3, Fz, F4, FC1 and FC2 for auditory trials and C3, Cz, C4, CP1 and CP2 for visual trials.

For the analysis of prestimulus alpha power, we extracted for each epoch the data from -500 to 0 msec before the stimulus and applied a fast Fourier transform to extract the power density spectrum. We then calculated the area under the curve for the alpha band (8-13 Hz). **Figure 3-1** shows the posterior distribution of prestimulus alpha power. We averaged the data of posterior electrodes (P7, P3, Pz, O1, Oz, O2, P4 and P8), obtaining thus one prestimulus alpha value for each stimulus. For the time window analyses, we computed the average alpha power value across all visual and (separately) auditory stimuli of which the onset latency fell within the 8-second window. The resulting time series were *z*-scored separately for each block.

The analyses described above eventually yielded one pupil-size value, two ITC values and two alpha power values for each 8-second block: one for the task-relevant stimulus sequence (visual trials in the Attend Visual condition; auditory trials in the Attend Auditory condition) and one for the task-irrelevant stimulus sequence (visual trials in the Attend Auditory condition; auditory trials in the Attend Visual condition).

<u>Temporal analysis around pupil peaks and pupil troughs</u>

We analyzed the variables ITC and prestimulus alpha power in segments of 10 windows around peaks and troughs of the pupil timecourse. The normalized pupil size timecourse of each participant was thresholded at +1 standard deviation to detect the peaks and -1 standard deviation to detect the troughs. For this, the function *findpeaks* in MATLAB was employed, with a minimum distance between consecutive peaks and consecutive troughs of 10 time windows. Then, the ITC and alpha power data of those windows were extracted for further analysis. To statistically compare cross-correlations between the time-windowed variables around pupil peaks and troughs, we used cluster-based permutation testing, with 1000 iterations and a significance level of $\alpha = 0.01$ for the individual time points and $\alpha = 0.05$ for the significant clusters. All the temporal analyses were performed on the trials of the task-relevant modality, and data were collapsed across modalities.

Statistics

Hit rate was defined as the proportion of targets in the cued modality that were detected in time (i.e., button press within 1 second after target onset). Any response that was not a hit was counted as an incorrect response. Average ITC values were compared with repeated-measures ANOVA with cued modality (Attend Visual, Attend Auditory) and trial type (visual, auditory) as within-subject factors. To examine the relationships between pupil size, ITC, prestimulus alpha power and task performance (hit rate, number of incorrect responses), we binned the 8-second time windows in 10 equally populated bins on the basis of one variable, and examined the effects of bin (1-10), task relevance (task-relevant vs. task-irrelevant stimulus sequence) and trial type (visual, auditory) on one of the other variables using linear regression analysis. In the analyses of task performance we included in the model both a linear and quadratic effect of bin. Statistics were performed using MATLAB.

Results

Behavioral results

Participants showed better performance in the Attend Auditory condition than in the Attend Visual condition (**Fig. 2a**). The hit rate was significantly higher ($F(1, 495) = 6.78$, $p = 0.009$), whereas the number of incorrect responses did not differ between conditions ($F(1, 475) = -0.88$, $p = 0.11$). We also inspected the relation between pupil size and task performance. Pupil bin had no linear and quadratic effects on hit rate ($F(1, 495) = 1.79$, $p = 0.18$; $F(1, 495) = 2.52$, $p = 0.11$). In contrast, pupil bin had both a linear and a quadratic effect on the number of incorrect responses ($F(1, 475) = 22.65$, $p < 0.001$; $F(1, 475) = 30.90$, $p < 0.001$). More incorrect responses were made at intermediate levels of pupil size (**Fig. 2a**).

Figure 2. Behavioral results. Hit rate and number of incorrect responses as a function of pupil bin (a) and prestimulus alpha bin (b). Colours correspond to the different task-relevant modalities (Aud: auditory; Vis: visual).

Prestimulus alpha power showed a qualitatively similar relationship with task performance. Alpha bin had no linear and quadratic effects on hit rate ($F(1, 515) = 0.04$, $p = 0.62$; $F(1, 515) = 0.08$, $p = 0.76$). In

contrast, alpha bin had both a linear and a quadratic effect on the number of incorrect responses (F(1, 495) = 15.678p < 0.001; F(1, 495) = 19.135p < 0.001). More incorrect responses were made at intermediate levels of prestimulus alpha power (**Fig. 2b**).

Low-frequency phase entrainment to simultaneous trains of auditory and visual stimuli

Figure 3a shows the ITC spectra for the various conditions. The top row shows strong ITC at the frequency corresponding to the task-relevant stimulus sequence (1.4 Hz for Attend Auditory, 1.1 Hz for Attend Visual), suggesting entrainment of EEG oscillations to the task-relevant stimulus rhythm. The corresponding ITC topography was frontal for the auditory stimulus sequence and central for the visual stimulus sequence (**Fig. 3b**). The ITC associated with the task-relevant stimulus sequence was larger for visual trials than for auditory trials (F(1, 100) = 28.67, p < 0.001; **Fig. 3c**). The bottom row of **Figure 3a** suggests that EEG oscillations also became entrained to the task-irrelevant stimulus sequences, although significantly less than to the task-relevant stimulus sequences, (F(1, 100) = 8.94, p = 0.003; **Fig. 3c**) and with more distributed topographies (**Fig. 3b**).

Figure 3. Low-frequency phase entrainment. a) ITC time-frequency spectra, averaged across all electrodes. White horizontal lines indicate the average presentation frequency of auditory (top left) and visual (top right) stimuli. b) Topographies of ITC strength at the peak frequency and the time of stimulus onset. c) Average ITC at the driving frequency as a function of task relevance (task-relevant vs. task-irrelevant stimulus sequence) and trial type.

Figure 3-1. Topoplots of pre-stimulus alpha power distribution for task-relevant trials.

Relation between pupil size, ITC, and prestimulus alpha power

We next inspected the paired associations between pupil size, ITC, and pre-stimulus alpha power. ITC did not show a main effect of pupil bin (**Fig. 4a**; $F(1, 992) = 0.12$, $p = 0.73$). However, there was a significant interaction between bin and task relevance ($F(1, 992) = 6.31$, $p = 0.01$) where the general pattern suggests that with increasing pupil size the low-frequency oscillations became more entrained to the task-relevant stimulus sequence and less entrained to the task-irrelevant stimulus sequence.

When looking at prestimulus alpha power, we found a highly significant main effect of pupil bin on alpha power (**Fig. 4b**; $F(1, 992) = 91.43$, $p < 0.001$). Prestimulus alpha power directly increased with pupil size, in line with a previous study (Podvalny et al., 2021). ITC did not show any significant effect of alpha power bin (**Fig. 4c**; $F(1, 1032) = 2.99$, $p = 0.08$), or interaction with task relevance ($F(1, 1032) = 0.55$, $p = 0.46$).

Figure 4. Paired relation between pupil size, ITC and alpha power. Relation between (a) pupil size and ITC; (b) pupil size and prestimulus alpha power; and (c) prestimulus alpha power and ITC. Different colours correspond to the task-relevant and task-irrelevant modality. Note that the distinction between the task-relevant and task-irrelevant modality is only of interest for ITC, since pupil values are the average over time windows including both task-relevant and task-irrelevant stimuli. There were no significant main effects and interaction effects of trial type (visual, auditory) so we averaged across those levels.

Arousal-related modulation of the temporal relation between ITC and alpha power

As an exploratory analysis, we asked whether pupil-linked arousal may modulate the temporal relation between entrainment and alpha power. Specifically, we inspected entrainment and alpha power associated with trials of the task-relevant modality in periods centered around pupil peaks and troughs (**Fig. 5a**). We found that pupil peaks were preceded by an increase in entrainment, but this effect was transient and small (**Fig. 5b**). More interestingly, a large increase in pre-stimulus alpha was observed prior to pupil peaks, and a large decrease prior to pupil troughs (**Fig. 5c**). This indicates that changes in the electrophysiological signatures occurred prior to the changes in pupil-linked arousal. To investigate whether the changes in entrainment and alpha power associated with changes in arousal were linked, we calculated the cross-correlation between these signals around pupil peaks and troughs. We found that the temporal relation between entrainment and alpha power was strongly modulated around zero-crossings of the pupil derivative (**Fig. 5d**): there was a significant positive correlation between ITC and alpha power in time windows around pupil peaks and a significant

negative correlation in time windows around pupil troughs. Interestingly, around pupil troughs there was a positive correlation between ITC and alpha power at negative lags (-6 and -7, i.e., window starting approximately 14 seconds before the pupil event). This indicates that alpha power and ITC covary during peaks of arousal, but they are temporally shifted around dips of arousal. Taken together, the results suggest that changes in ITC and alpha power are temporally related, and that this relation is strongly modulated by arousal.

Figure 5. Temporal relation between ITC and pre-stimulus alpha power around pupil peaks and pupil troughs. a) Example of one participant's time course of pupil size and detected peaks and troughs marked with yellow and purple triangles, respectively; b) and c) timecourse of ITC and alpha power, respectively, around pupil peaks and

troughs; d) cross-correlogram between ITC and alpha power associated with trials of the task-relevant modality: around pupil peaks (purple) and around pupil troughs (yellow). Negative lags correspond to ITC leading alpha power, while positive lags correspond to alpha power leading ITC. Horizontal lines above and below the time courses indicate significant difference from zero based on permutation testing ($p < 0.05$).

Discussion

The goal of the current study was to unravel how variations in the brain's internal state contribute to fluctuations in neural and behavioral measures of attention at slow timescales. We examined the impact of varying levels of pupil-linked arousal and alpha power on low-frequency phase entrainment. We found that increases in pupil size coincided with more alignment of low-frequency oscillations to the task-relevant stimulus sequence and less alignment to the task-irrelevant stimulus sequence. Furthermore, we found that alpha power and ITC showed a significant correlation, but only in the time windows around pupil peaks and troughs, with the sign of the correlation depending strongly on whether pupil-linked arousal was high or low. These results suggest that variability in internal state modulates how the brain responds to external rhythmic stimulation, and that arousal is a key factor coordinating the oscillatory mechanisms of the brain.

Facilitatory effect of arousal on low-frequency entrainment

A key finding in our study was that increases in pupil size coincided with more entrained low-frequency oscillations to the task-relevant stimulus sequence, while entrainment to the task-irrelevant stimulus sequence decreased. Previous research has found that diminished phase entrainment is associated with lapses in attention and response accuracy (van den Brink et al., 2014; Lakatos et al., 2016). Our finding suggests that transient decreases in pupil-linked arousal may be one factor underlying these observations. Studies in animals suggest that pupil dilation is associated with changes in neuronal excitability and population activity (Reimer et al., 2014; McGinley et al., 2015a; Vinck et al., 2015), tuning the cortex into an active state (McGinley et al. 2015b). In addition, increased arousal is often associated with better task performance at low-to-intermediate arousal levels, while task performance decreases at very high levels of pupil-linked arousal, giving rise to an inverted-U relation between pupil-linked

arousal and performance (van den Brink et al., 2016; Waschke et al., 2019; Beerendonk et al., 2023). Interestingly, Figure 4a shows decreased task-relevant ITC in the time windows with the largest baseline pupil size (bin 10). Our findings suggest that pupil-linked arousal, by shaping the brain's internal state, affects how the brain processes task-relevant rhythmic stimuli.

<u>Arousal-dependent relation between pre-stimulus alpha power and low-frequency entrainment</u>

A second key finding was an absence of a systematic relationship between posterior alpha power and phase entrainment. This was surprising, given that both variables showed significant associations with pupil size. Indeed, alpha power and pupil size showed a strong positive relationship, a similar relationship with task performance (i.e., negative quadratic relationship with number of incorrect responses), as well as a clear temporal relationship (**Fig. 5c**): alpha power peaked before pupil peaks and reached a minimum before pupil troughs—a similar pattern as in Montefusco et al. (2022). This apparent dissociation between pupil size (correlated with entrainment) and posterior alpha power (uncorrelated with entrainment) could be explained by dissociable effects of these variables on perception.

Our exploratory analyses of the temporal relation between pre-stimulus alpha power, ITC and pupil peaks/troughs shed some light on this dissociation: alpha power and ITC showed a significant cross-correlation, but only in the time windows around pupil peaks and troughs, with the sign of the correlation depending strongly on whether pupil-linked arousal was high or low. Interestingly, we found a significant cross-correlation between alpha power and entrainment preceding pupil troughs (at lags corresponding to time windows starting 14 seconds before), suggesting that alpha power lagged with respect to entrainment in the order of 10 seconds. This resembles the finding of Lakatos and colleagues (2016) of a slow anti-correlation in nonhuman primates between alpha power and ITC with a period of around 16 seconds. Intracranial measurements in humans also showed a modulation of alpha power by the phase of delta entrainment (Gomez-Ramirez et al., 2011). We can only speculate why the coupling between entrainment and alpha power was shifted in time at the different extremes of arousal. One possibility is that the peaking of entrainment and alpha power and their correlation around pupil peaks

reflected activity in brainstem arousal centers, which also drove the peaking of pupil size. Indeed, activity in the locus coeruleus occurs shortly before pupil dilation (Joshi et al., 2016), pupil-linked arousal has generalized effects on the brain that are evident at slower time scales (Shine et al., 2016). On the other hand, the counter-relation around pupil throughs may have reflected an inattention mechanism within the alpha band that emerged at low arousal. Indeed, it has been proposed that top-down and primary sensory circuits may contribute to the alpha band with opposing effects (Bollimunta et al., 2008). During low arousal, cortical effects on alpha power may signal basic sensory processes, which are dissociated from pupil-linked arousal effects (Pilipenko & Samaha, 2023), and present a slow modulation also during rest (~17-seconds period; Novak et al. 1992). Given that the results in humans and non-human primates suggest the translational value of the findings discussed here, future research should replicate and further disentangle the relation between alpha power and ITC during periods of large and small pupil-linked arousal. In terms of mechanisms of selective attention, we speculate that arousal is related to higher entrainment and cognitive contributions to alpha power (Sadaghiani and Kleinschmidt, 2016; Clayton et al., 2018). A slow counter-modulation between sensory alpha power and entrainment, rather than being sustained, may emerge during periods of low arousal.

Limitations

From a methodological perspective, it is hard to distinguish between true entrainment of ongoing neural oscillations and the consistent phase alignment caused by a sequence of stimulus-evoked potentials (Haegens and Golumbic 2018; van Diepen and Mazaheri, 2018; Helfrich et al., 2019). Nevertheless, mounting evidence shows that phase coherence persists during no-stimulus periods, suggesting that it is at least in part caused by neural entrainment (Zoefel et al., 2018; Bouwer et al., 2023), and intracranial studies in humans support the presence of wide-spread alignment of oscillations to rhythmic stimuli, not restricted to sensory areas (Gomez-Ramirez et al., 2011; Besle et al., 2011).

Previous studies have found within-trial phase-amplitude coupling between the delta phase of entrained oscillatory activity and alpha power over posterior brain regions (Wostmann et al., 2016; Wilson and Foxe, 2020). Here we were interested in fluctuations at slow time scales, similar to Lakatos

et al. (2016), because they likely represent variation in the brain's internal states. To compute the uniformity of phase angles across trials we used 8-second sliding time windows. The length of these time windows did not allow for sub-second temporal resolution, limiting our ability to study within-trial relationships between entrainment and alpha power at various levels of arousal. Taken together, our results and previous studies suggest that internally and externally-driven oscillatory mechanisms operate at different time scales.

Conclusions

Neural entrainment is an important instrument of selective attention (Schroeder and Lakatos, 2009). Here we reported delta-phase entrainment to both the visual and the auditory stimulus sequences, with stronger entrainment to the task-relevant modality than to the task-irrelevant modality. This is consistent with previous studies using bimodal stimulation (Besle et al., 2011; Lakatos et al., 2016). Most importantly, we found a significant linear relationship between pupil size and the degree to which entrainment tracked the task-relevant instead of the task-irrelevant stimulus sequence, indicating a facilitating effect of arousal on the perception of rhythmic stimulus sequences. We also found a coupling between alpha power and entrainment around pupil peaks and pupil troughs. Our results give clear indications of a differential modulation of brain oscillatory processes by arousal.

References

Bauer AKR, Van Ede F, Quinn AJ, Nobre AC (2021) Rhythmic modulation of visual perception by continuous rhythmic auditory stimulation. *Journal of Neuroscience* 41(33):7065-7075.

Beerendonk L, Mejías JF, Nuiten SA, de Gee JW, Fahrenfort JJ, van Gaal S (2023) A disinhibitory circuit mechanism explains a general principle of peak performance during mid-level arousal. *bioRxiv* 2023.07.28.550956.

Benedek M, Schickel RJ, Jauk E, Fink A, Neubauer AC (2014) Alpha power increases in right parietal cortex reflects focused internal attention. *Neuropsychologia* 56:393-400.

Besle J, Schevon CA, Mehta AD, Lakatos P, Goodman RR, McKhann GM, Emerson RG, Schroeder CE (2011) Tuning of the human neocortex to the temporal dynamics of attended events. *J Neurosci* 31:3176-85.

Bollimunta A, Chen Y, Schroeder, C. E., Ding M (2008) Neuronal mechanisms of cortical alpha oscillations in awake-behaving macaques. *Journal of Neuroscience* 28(40):9976-9988.

Bouwer FL, Fahrenfort JJ, Millard SK, Kloosterman NA, Slagter HA (2023) A silent disco: Differential effects of beat-based and pattern-based temporal expectations on persistent entrainment of low-frequency neural oscillations. J Cogn Neurosci 35:990-1020.

Bradshaw J (1967) Pupil size as a measure of arousal during information processing. *Nature*216(5114): 515-516.

Ceh SM, Annerer-Walcher S, Körner C, Rominger C, Kober SE, Fink A, Benedek M (2020) Neurophysiological indicators of internal attention: An electroencephalography-eye-tracking coregistration study. *Brain Behav* 10:e01790.

Clayton MS, Yeung N, Cohen Kadosh R (2018) The many characters of visual alpha oscillations. *European Journal of Neuroscience* 48(7):2498-2508.

Compton RJ, Gearinger D, Wild H (2019) The wandering mind oscillates: EEG alpha power is enhanced during moments of mind-wandering. *Cogn Affect Behav Neurosci* 19:1184-1191.

Cooper NR, Croft RJ, Dominey SJ, Burgess AP, Gruzelier JH (2003) Paradox lost? Exploring the role of alpha oscillations during externally vs. internally directed attention and the implications for idling and inhibition hypotheses. *Int J Psychophysiol* 47:65-74.

Dal Ben R (2023) SHINE_color: controlling low-level properties of colorful images. *MethodX* 11:102377. https://doi.org/10.1016/j.mex.2023.102377

Delorme A, Makeig S (2004) EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *J Neurosci Methods* 134:9-21.

Gomez-Ramirez M, Kelly SP, Molholm S, Sehatpour P, Schwartz TH, Foxe JJ (2011) Oscillatory sensory

selection mechanisms during intersensory attention to rhythmic auditory and visual inputs: a

human electrocorticographic investigation. *Journal of Neuroscience* 31(50),:18556-18567.

Groot JM, Boayue NM, Csifcsák G, Boekel W, Huster R, Forstmann BU, Mittner M (2021) Probing the

neural signature of mind wandering with simultaneous fMRI-EEG and pupillometry. *Neuroimage*

224:117412.

Haegens S, Golumbic E Z (2018) Rhythmic facilitation of sensory processing: A critical review.

Neuroscience & Biobehavioral Reviews 86,:150-165.

Helfrich RF, Breska A, Knight RT (2019). Neural entrainment and network resonance in support of top-
down guided attention. *Current opinion in psychology*, *29*, 82-89.

Henry MJ, Obleser J (2012) Frequency modulation entrains slow neural oscillations and optimizes

human listening behavior. *Proc Natl Acad Sci* 109:20095-100.

Hong L, Walz JM, Sajda P (2014) Your eyes give you away: prestimulus changes in pupil diameter

correlate with poststimulus task-related EEG dynamics. *PLOS ONE* 9:e91321.

Hopstaken JF, van der Linden D, Bakker AB, Kompier MA (2015) A multifaceted investigation of the link

between mental fatigue and task disengagement. *Psychophysiology* 52:305-15.

Jin CY, Borst JP, van Vugt MK (2019) Predicting task-general mind-wandering with EEG. *Cogn Affect*

Behav Neurosci 19:1059-1073.

Joshi S, Li Y, Kalwani R M, Gold JI (2016) Relationships between pupil diameter and neuronal activity in

the locus coeruleus, colliculi, and cingulate cortex. *Neuron* 89(1),:221-234.

Klimesch W, Sauseng P, Hanslmayr S (2007) EEG alpha oscillations: the inhibition-timing hypothesis.

Brain Res Rev 53:63-88.

Kret ME, Sjak-Shie EE (2019) Preprocessing pupil size data: Guidelines and code. *Behav Res Methods*

51:1336-1342.

Lakatos P, Barczak A, Neymotin SA, McGinnis T, Ross D, Javitt DC, O'Connell MN (2016) Global

dynamics of selective attention and its lapses in primary auditory cortex. *Nat Neurosci* 19:1707-

1717.

Lakatos P, Karmos G, Mehta AD, Ulbert I, Schroeder CE (2008) Entrainment of neuronal oscillations as a mechanism of attentional selection. Science 320:110–113. doi: 10.1126/science.1154735

Lakatos P, Gross J, Thut G (2019) A new unifying account of the roles of neuronal entrainment. *Current Biology* 29(18):R890-R905.

Madore KP, Khazenzon AM, Backes CW, Jiang J, Uncapher MR, Norcia AM, Wagner AD (2020) Memory failure predicted by attention lapsing and media multitasking. *Nature* 587:87-91.

Manzotti R (2017) A perception-based model of complementary afterimages. *SAGE Open* 7(1).

Martínez-Cancino R, Delorme A, Truong D, Artoni F, Kreutz-Delgado K, Sivagnanam S, Yoshimoto K, Majumdar A, Makeig S (2021) The open EEGLAB portal Interface: High-Performance computing with EEGLAB. *NeuroImage*, 224: 116778

Matthewson et al. (2012) Making waves in the stream of consciousness: entraining oscillations in EEG alpha and fluctuations in visual awareness with rhythmic visual stimulation. *Journal of cognitive neuroscience* 24(12):2321-2333.

McGinley MJ, David SV, McCormick DA (2015a) Cortical membrane potential signature of optimal states for sensory signal detection. *Neuron* 87(1):179-192.

McGinley MJ, Vinck M, Reimer J, Batista-Brito R, Zagha E, Cadwell CR, Tolias AS, Cardin JA, McCormick DA. (2015b) Waking state: Rapid variations modulate neural and behavioral responses. *Neuron* 87:1143-1161.

Montefusco-Siegmund R, Schwalm M, Rosales Jubal E, Devia C, Egaña JI, Maldonado PE (2022) Alpha EEG activity and pupil diameter Coupling during inactive wakefulness in humans. *eNeuro* 9:ENEURO.0060-21.2022.

Novak P, Lepicovska V, Dostalek C (1992) Periodic amplitude modulation of EEG. *Neuroscience letters* 136(2):213-215.

O'Connell RG, Dockree PM, Robertson IH, Bellgrove MA, Foxe JJ, Kelly SP (2009) Uncovering the neural signature of lapsing attention: electrophysiological signals predict errors up to 20 s before they occur. *J Neurosci* 29:8604-11.

Oken BS, Salinsky MC, Elsas S (2006) Vigilance, alertness, or sustained attention: physiological basis and measurement. *Clinical neurophysiology* 117(9):1885-1901.

Pilipenko A, Samaha J (2023) Double dissociation of spontaneous alpha-band activity and pupil-linked arousal on additive and multiplicative perceptual gain. *bioRxiv.*

Pfeffer T, Keitel C, Kluger DS, Keitel A, Russmann A, Thut G, Donner TH, Gross J (2022) Coupling of pupil- and neuronal population dynamics reveals diverse influences of arousal on cortical processing. *eLife* 11:e71890.

Podvalny E, King LE, He BJ (2021) Spectral signature and behavioral consequence of spontaneous shifts of pupil-linked arousal in human. *eLife* 10:e68265.

Reimer J, Froudarakis E, Cadwell CR, Yatsenko D, Denfield GH, Tolias AS (2014) Pupil fluctuations track fast switching of cortical states during quiet wakefulness. *Neuron* 84(2):355-362.

Romei et al. (2008) Spontaneous fluctuations in posterior α-band EEG activity reflect variability in excitability of human visual areas. *Cereb Cortex* 18:2010–2018

Sadaghiani S, Kleinschmidt A (2016) Brain networks and α-oscillations: structural and functional foundations of cognitive control. *Trends in cognitive sciences* 20(11):805-817.

Samaha J, Gosseries O, Postle BR (2017) Distinct oscillatory frequencies underlie excitability of human occipital and parietal cortex. *Journal of Neuroscience* 37(11):2824-2833.

Schroeder CE, Lakatos P (2009) Low-frequency neuronal oscillations as instruments of sensory selection. *Trends Neurosci* 32:9-18.

Shine JM, Bissett PG, Bell PT, Koyejo O, Balsters JH, Gorgolewski KJ, Poldrack RA (2016) The dynamics of functional brain networks: integrated network states during cognitive task performance. *Neuron* 92(2):544-554.

Sivagnanam S, Majumdar A, Yoshimoto K, Astakhov V, Bandrowski A, Martone ME, and Carnevale NT (2013) Introducing the Neuroscience Gateway, IWSG, volume 993 of CEUR Workshop Proceedings, CEUR-WS.org.

Stefanics G, Hangya B, Hernádi I, Winkler I, Lakatos P, Ulbert I (2010) Phase entrainment of human delta oscillations can mediate the effects of expectation on reaction speed. *Journal of Neuroscience* 30(41):13578-13585.

Tallon-Baudry C, Bertrand O, Delpuech C, Pernier J. (1996) Stimulus specificity of phase-locked and non-phase-locked 40 Hz visual responses in human. *J Neurosci* 16, 4240–4249.

Ten Oever et al. (2017) Low-frequency cortical oscillations entrain to subthreshold rhythmic auditory stimuli. *Journal of Neuroscience* 37(19):4903-4912.

Unsworth N, Robison MK (2016) Pupillary correlates of lapses of sustained attention. *Cogn Affect Behav Neurosci* 16:601-15.

Unsworth N, Robison MK (2017) The importance of arousal for variation in working memory capacity and attention control: A latent variable pupillometry study. *Journal of Experimental Psychology: Learning, Memory, and Cognition* 43(12):1962.

van den Brink RL, Murphy PR, Nieuwenhuis S (2016) Pupil diameter tracks lapses of attention. *PLOS ONE* 11:e0165274.

van den Brink RL, Wynn S, Nieuwenhuis S (2014) Post-error slowing as a consequence of disturbed low-frequency oscillatory phase entrainment. *J Neurosci* 34:11096-11105.

van Diepen RM, Mazaheri A (2018). The caveats of observing inter-trial phase-coherence in cognitive neuroscience. *Scientific reports*, *8*(1), 2990.

van Kempen J, Loughnane GM, Newman DP, Kelly SP, Thiele A, O'Connell RG, Bellgrove MA (2019) Behavioural and neural signatures of perceptual decision-making are modulated by pupil-linked arousal. *eLife* 8:e42541.

Van Rullen et al. (2014) On the cyclic nature of perception in vision versus audition. *Philosophical transactions of the Royal Society B: Biological sciences* 369(1641):20130214.

Vinck M, Batista-Brito R, Knoblich U, Cardin JA (2015) Arousal and locomotion make distinct contributions to cortical activity patterns and visual encoding. *Neuron* 86(3):740-754.

Waschke L, Tune S, Obleser J (2019) Local cortical desynchronization and pupil-linked arousal differentially shape brain states for optimal sensory performance. *eLife* 8:e51501.

Wilson TJ, Foxe JJ (2020) Cross-frequency coupling of alpha oscillatory power to the entrainment

   rhythm of a spatially attended input stream. *Cogn Neurosci* 11:71-91.

Wöstmann M, Herrmann B, Maess B, Obleser J (2016) Spatiotemporal dynamics of auditory attention

   synchronize with speech. *Proc Natl Acad Sci* 113:3873-8.

Zoefel B, Ten Oever S, Sack AT (2018) The involvement of endogenous neural oscillations in the

   processing of rhythmic input: More than a regular repetition of evoked neural responses. *Front*

   Neurosci 12:95. doi: 10.3389/fnins.2018.00095