Antibodies against medically relevant arthropod-borne viruses in the ubiquitous African rodent *Mastomys natalensis*

Short title: Antibodies against arboviruses in *Mastomys natalensis* from Tanzania

Wim De Kesel [1, 2*], Bram Vanden Broecke [1, 3], Benny Borremans [1, 4], Léa Fourchault [5], Elisabeth Willems [2], Ann Ceulemans [2, 6], Christopher Sabuni [7], Apia Massawe [7], Rhodes H. Makundi [7], Herwig Leirs [1], Martine Peeters [8], Erik Verheyen [1, 5], Sophie Gryseels [1, 5], Joachim Mariën [1, 6] and Kevin K. Ariën [2, 9*]

[1] Evolutionary Ecology Group, Department of Biology, Faculty of Science, University of Antwerp, Antwerp, Belgium

[2] Virology Unit, Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

[3] Terrestrial Ecology Unit, Department of Biology, Ghent University, Ghent, Belgium

[4] Wildlife Health Ecology Research Organization, San Diego, United States of America

[5] OD Taxonomy & Phylogeny, Royal Belgian Institute of Natural Sciences, Brussels, Belgium

[6] Virus Ecology Unit, Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

[7] Institute of Pest Management, Sokoine University of Agriculture, Morogoro, Tanzania

[8] TransVIHMI, University of Montpellier, Institut de Recherche pour le Développement (IRD), INSERM, Montpellier, France

[9] Department of Biomedical sciences, Faculty of Pharmaceutical, Biomedical and Veterinary Sciences, University of Antwerp, Antwerp, Belgium

* Corresponding authors

E-mail: wim.dekesel@uantwerpen.be (WDK); karien@itg.be (KKA)

Abstract

Over the past decades, the number of arthropod-borne virus (arbovirus) outbreaks has increased worldwide. Knowledge regarding the sylvatic cycle (i.e., non-human hosts/environment) of arboviruses is limited, particularly in Africa, and the main hosts for virus maintenance are unknown. Previous studies have shown the presence of antibodies against certain arboviruses (i.e., chikungunya-, dengue- and zika virus) in African non-human primates and bats. We hypothesize that small mammals, specifically rodents, may function as amplifying hosts in anthropogenic environments. The detection of RNA of most arboviruses is complicated by the virus's short viremic period within their hosts. An alternative to determine arbovirus hosts is by detecting antibodies, which can persist several months. We developed a high-throughput multiplex immunoassay to detect antibodies against 15 medically relevant arboviruses. We used this assay to assess almost 1,300 blood samples of the multimammate mouse, *Mastomys natalensis* from Tanzania. In 24% of the samples, we detected antibodies against at least one of the tested arboviruses, with high seroprevalences of antibodies reacting against dengue virus serotype one (7.6%) and two (8.4%) and chikungunya virus (6%). Seroprevalence was higher in females and increased with age, which could be explained by inherent immunity and behavioral differences between sexes and the increased chance of exposure to an arbovirus with age. We evaluated whether antibodies against multiple arboviruses co-occur more often than randomly and found that this may be true for some members of the *Flaviviridae* and *Togaviridae*. In conclusion, the development of an assay against a wide diversity of medically relevant arboviruses enabled the analysis of a large sample collection of one of the most abundant African small mammals. Our findings suggest a role in the transmission of multiple arboviruses by this ubiquitous rodent and

47 provide a solid foundation for future molecular screening to elucidate the role in the arbovirus

48 transmission cycle.

Author summary

One of the main causes of zoonotic related human morbidity and mortality is the

transmission of arthropod-borne viruses such as dengue virus, Yellow Fever virus, and

chikungunya virus. These viruses cannot only infect humans but also livestock, pets, and

wildlife, though our understanding of their non-human hosts remains limited. Rodents are

thought to be an interesting host for these viruses because they can be abundant, often live

near humans and some are already known to be viral hosts. However, research has focused

on non-human primates, neglecting other potential hosts. To address this gap, we have

developed a high-throughput antibody test to screen rodent blood against 15 different

arboviruses. Our findings reveal that a proportion of *Mastomys natalensis*, a common African

rodent species, carry antibodies that (cross-)react against these viruses. We hypothesize that

immunologically naïve juveniles may drive transmission, particularly during population

outbreaks. These outbreaks coincide with environmental conditions that are favorable for

mosquitoes, the vectors of these viruses. Thus, increasing the risk of spillover to humans,

livestock, and wildlife. Understanding the role of rodents in arbovirus transmission dynamics

is crucial for mitigating zoonotic disease risks.

Introduction

The African continent harbors a diverse array of infectious diseases with profound impacts on public health, economic development, and general well-being [1,2]. Diseases caused by arthropod-borne viruses, collectively known as arboviruses, are a growing threat for Africa and the rest of the world especially in relation to climate and environmental changes [3,4]. Arboviruses are a polyphyletic clade that includes several viral families, of which the most important are *Flaviviridae, Togaviridae, Bunyaviridae,* and *Reoviridae* [5]. Some well-known arboviruses, notorious for their devastating effects on human health, are dengue virus, Yellow Fever virus, Zika virus, and chikungunya virus. Mosquitoes, ticks, sandflies, and midges are the primary vectors responsible for arbovirus transmission as they engage in hematophagy. These vectors do not only affect humans and livestock, but also a wide range of wildlife hosts [6–8]. Indeed, while for some arboviruses morbidity and mortality can be high in humans, similar impacts have been detected in other animals by arboviruses such as Rift Valley Fever virus in goats and sheep, West Nile virus in birds and horses and Japanese Encephalitis virus in birds and pigs [4,5,9–11]. The (re-)emergence of arboviruses appears, at least partially, to be caused by the increased urbanization and global connectivity, natural genetic evolution of viruses, and adaptations of the vectors to changing climate and environments [11,12]. Emerging arboviruses pose a threat for humans, livestock, as well as wildlife, therefore it needs to be approached from a One health perspective (i.e., including human, animal, and environment health) [13,14]. Nevertheless, our knowledge about the extent to which wild animals can serve as sylvatic hosts for human-infecting arboviruses and the natural diversity of arboviruses remains insufficient. This significantly limits our

understanding of arbovirus transmission dynamics, which is required to develop effective

control measurements.

For decades, efforts have been made to identify natural reservoirs of arboviruses to

monitor, prevent, and control sources of infection that pose a threat to human health [15–

17]. Several studies have proposed non-human primates as significant potential reservoirs for

arboviruses, as they have found arbovirus antibodies and viral RNA in this animal group [18–

20]. However, other animal groups such as small mammals have often been neglected as

arbovirus hosts [21]. Sporadic reports of arboviruses in small mammal species suggest that a

more comprehensive investigation of their potential role as a host is needed [22–24].

Rodents have a number of characteristics that could make them potentially important

hosts for several pathogens, including arboviruses [23]. Particularly the high species diversity,

the fact that many species can reach high population abundances and turnover rates. The risk

of pathogen spillover to humans increases with the role of some rodents as a pest species,

due to their proximity to humans [25,26]. A notable example of such a pest species is the

ubiquitous rodent *Mastomys natalensis*, commonly known as the multimammate mouse. This

species inhabits many regions of sub-Saharan Africa, with a preference for crop fields, fallow

land, and typically occurring within or at the fringes of urban settlements [27,28]. In east

Africa, and especially in Tanzania, the reproductive cycle of *M. natalensis* is strongly

correlated with seasonal rainfall which leads to strong seasonal fluctuations in density (20 -

500 individuals/hectare), and occasionally even severe population outbreaks (>1000

individuals/hectare) [29–32]. This has large ecological and societal impacts due to crop

damage and influences seasonal transmission dynamics of different pathogens [33–35]. The

multimammate mouse is a known host for several zoonotic pathogens such as *Lassa*

mammarenavirus, *Yersinia pestis*, *Leptospira interrogans*, *Leishmania major* as well as

different ecto- and endoparasites [31,36–52]. No studies have investigated or reported on arboviruses in *M. natalensis*, except Diagne et al. (2019) who have detected Usutu virus RNA in *M. natalensis*. However, other studies have reported on sporadic arbovirus detections in other rodent species in sub-Saharan Africa [22,24,53,54]. These findings, along with the ecology of *M. natalensis* (i.e., high abundance during population outbreaks, proximity to humans, and its status as a proven pathogen host) may suggest that this species plays a role in the natural transmission cycle of arboviruses. Consequently, *M. natalensis* could thus pose a risk to humans in east Africa, particularly in Tanzania as an amplifying host.

The human population in Tanzania has experienced several outbreaks of chikungunya virus, Rift Valley fever virus, West Nile virus, and dengue virus in the past decades [55–59]. Due to the symptomatic similarities between arbovirus and malaria infections, which has a prevalence of around 20% in Tanzania, it is probable that arbovirus cases are underreported [60,61]. While these studies confirm that the local human population is indeed exposed to arboviruses, the specific dynamics of arbovirus transmission in this region remains unclear.

The goal of this study was to investigate the potential of wild *M. natalensis* to serve as a host for arboviruses in their natural environment. To achieve this, we first developed a multiplex immune assay to detect immunoglobulin G (IgG) antibodies against 15 different arboviruses causing disease in humans and subsequently conducted a comprehensive screening of almost 1,300 blood samples obtained from *M. natalensis* from Morogoro, Tanzania.

Materials and Methods

Sample origin

The samples used in this study were collected during previous published and unpublished studies conducted by the University of Antwerp (UAntwerp) and Pest Management Center of the Sokoine University of Agriculture on *M. natalensis* in Morogoro, Tanzania, between 2010 and 2019 [31,62,63] (Fig 1). In this study, the samples were divided in two screening sessions. The first session consisted of approximately 500 dried blood spot (DBS) samples, from wild captured mice that were used in infection and behavioral experiments in six different years (i.e., 2010, 2011, 2015, 2017, 2018, and 2019) with an average of 80 samples per year. The second session consisted of 800 DBS samples from mice involved in capture mark recapture experiments in 2017 and 2019. All samples were randomly selected from the studies regardless of individual characteristics or trapping period.

During these studies, *M. natalensis* were live caught using Sherman traps (H.B. Sherman Traps, Tallahassee, USA) in a heterogeneous landscape (e.g., woodlands, maize fields, and fallow land) on the premises of the Sokoine University of Agriculture in Morogoro, Tanzania. Blood was collected from the retro-orbital plexus using a 50μL hematocrit capillary tube and preserved on filter paper (Serobuvard; LDA 22; Zoopole, France). The filter paper was dried for 12 hours at room temperature and archived at -20°C in envelopes with desiccant. Additional data related to characteristics such as sex, reproductive status, weight, and body measurements were recorded. More detailed Information pertaining to the trapping procedures and sampling methodology can be found in the primary research documents associated with these studies [31,62–64].

Fig 1. African continent with a focus on Tanzania. Samples were collected in the city of Morogoro (red triangle) which is located in the Morogoro region.

Analysis set up and protocol.

To assess the presence of arbovirus antibodies in DBS against a panel of arboviruses, we first developed a multiplex immune assay using Luminex technology [18,65]. Recombinant virus-derived proteins (Table 1) were covalently coupled to carboxyl-functionalized fluorescent magnetic beads (1-3μg/1.25*10^6 beads) (Luminex Corp. MagPlex® Microspheres; Bio-Rad; Temse, Belgium) employing the BioPlex amine coupling kit (Ref.: 171406001; Bio-Rad; Temse, Belgium) following the manufacturer's instructions.

To obtain positive control samples we inoculated captive *M. natalensis* individuals (age: 5-12 months) from our breeding colony at UAntwerp with recombinant virus-derived proteins (Table 1) [66–68]. We subcutaneously injected 4μg of the respective virus protein and 1μL of vaccine adjuvant (Quil-A® adjuvant; InvivoGen; Toulouse, France), dissolved in autoclaved phosphate buffered saline (PBS) to achieve a final volume of 1mL. This inoculum was evenly divided, with 0.5mL administered into the scruff and 0.5mL into the hindlimb of the animal, using a 25-gauge, 12.5mm needle and a 0.5mL syringe. This inoculation was duplicated for each viral protein (i.e., performed in two mice) and repeated twice for each

mouse (i.e., inoculation on day 0 and day 20). We collected blood, according to the same method as in the previous studies, every 10 days from day zero until day 30, at day 30 we also collected whole blood from which serum was extracted. Serum from day 30 from individuals were the antibody response increased over time were considered as positive samples. Day 30 had the highest antibody titer in our tests and is also a time point at which IgG antibody development is anticipated to have reached a peak [69,70].

Table 1. Recombinant arbovirus proteins for the inoculation of captive *Mastomys natalensis*.

Viral family	Virus	Protein (reference)	Supplier
Bunyaviridae	Rift Valley Fever virus (RVFV)	Nucleoprotein (REC31640)	The native antigen company (Kidlington, United Kingdom)
Flaviviridae	Yellow Fever virus (YFV)	Nonstructural protein 1 (YFV-NS1)	The native antigen company (Kidlington, United Kingdom)
Flaviviridae	Zika virus (ZIKV)	Nonstructural protein 1 (40544-V07H)	Interchim (Montluçon Cedex, France)
Flaviviridae	Dengue virus serotype 1 (DENV1)	Nonstructural protein 1 (DEN-004)	Prospecbio (Rehovot, Israel)
Flaviviridae	Dengue virus serotype 2 (DENV2)	Nonstructural protein 1 (PIP048A)	BioRad (Temse, Belgium)
Flaviviridae	Dengue virus serotype 3 (DENV3)	Nonstructural protein 1 (DENV3-NS1)	The native antigen company (Kidlington, United Kingdom)
Flaviviridae	Dengue virus serotype 4 (DENV4)	Nonstructural protein 1 (DENV4-NS1)	The native antigen company (Kidlington, United Kingdom)
Flaviviridae	Usutu virus (USUV)	Nonstructural protein 1 (Ab218552)	The native antigen company (Kidlington, United Kingdom)
Flaviviridae	West Nile virus (WNV)	Nonstructural protein 1 (40346-V07H)	Sinobiological (Eschborn, Germany)
Flaviviridae	Tick-borne Encephalitis virus (TBEV)	Nonstructural protein 1 (TBEV-NS1)	The native antigen company (Kidlington, United Kingdom)
Flaviviridae	Wesselsbron virus (WSLV)	Nonstructural protein 1 (REC31698)	The native antigen company (Kidlington, United Kingdom)
Nairoviridae	Crimean Congo Hemorrhagic Fever virus (CCHFV)	Nucleoprotein (REC31639)	The native antigen company (Kidlington, United Kingdom)
Togaviridae	Chikungunya virus (CHIKV)	Envelope protein 2 (CHI-003)	Prospecbio (Rehovot, Israel)
Togaviridae	Mayaro virus (MAYV)	Envelope protein 2 (REC31644)	The native antigen company (Kidlington, United Kingdom)
Togaviridae	O'nyong nyong virus (ONNV)	Envelope protein 2 (B4TG40)	Interchim (Montluçon Cedex, France)

Screening was done in 96 flat-bottom well plates, each plate contained DBS samples

of 80 wild *M. natalensis*, two background controls, eight negative controls and a six step

dilution series (1:200 – 1:625,000) of a positive pool sample. Each well in the plate contained

50µL of the corresponding sample type. The samples of the wild *M. natalensis* were acquired

by placing a punched-out DBS (round, 0.5 cm diameter) in 200 µL of dilution buffer (1% bovine

serum albumin, 0.2% Tween-20, 5% fetal calf serum, 45% distilled water, 50% Hypertonic PBS

{0.08% NaH_2PO_4, 0.25% Na2HPO$_4$, 8.8% NaCl}). One single DBS punch corresponds to

approximately 10µL of blood [71]. The punched DBS were left to elute overnight, in a 1.5mL

Eppendorf tube, maintained at a temperature of 4°C on a plate shaker. This elution was

considered a 1:100 dilution and was diluted, with dilution buffer, to 1:200 prior to loading in

the 96 well plate. This dilution gave the best signal to noise ratio in our preliminary tests and

are in line with previous studies [18,72]. The background control was reading buffer (1%

bovine serum albumin, 0.05% NaN$_3$, 100% phosphate buffered saline). The eight negative

controls were four DBS, treated the same as the wild *M. natalensis* DBS, and four serum

samples in a 1:200 dilution. All negative controls originated from the breeding colony at

UAntwerp. The positive pool sample was compiled from serum from the positive individuals

acquired through the arbovirus protein inoculation experiment. Serum from 15 positive

individuals (i.e., one for each arbovirus antigen) was pooled to create the positive pool

sample, each individual serum had a final dilution in the pool of 1:200.

In each well of the 96 well plate, 25µL of bead mixture was added. The bead mixture

consisted out of ~1000 protein-coated beads per arbovirus antigen suspended in reading

buffer. The bead mixture of the first screening session did not contain ONNV beads.

Plates, containing 50µL of sample and 25µL of bead mixture per well, were incubated

for one hour at room temperature, in the dark and on a plate shaker (Heidolph Titrimax 100;

VWR; Leuven, Belgium) at 400rpm/min. After incubation, plates underwent washing with

dilution buffer using an automated plate washer (Tecan Hydroflex plate washer; Tecan

Benelux; Mechelen, Belgium). Subsequently, we added 50µL Biotin anti-mouse IgG (4µg/mL)

(Sigma-Aldrich B7022; Merck Life Science; Hoeilaart, Belgium) to each well and incubated for

40 minutes. After another round of washing, we added 50µL of Streptavidin-R-phycoerythrin

(1µg/mL) (10655783; Fisher Scientific; Brussel, Belgium) to each well, followed by a 10-minute

incubation. The last wash step used reading buffer, and the final bead pellet was resuspended

in 150µL of reading buffer. Beads were read on a Bio-Plex 200 System (Bio-Rad; Temse,

Belgium). Results were quantified as the median fluorescent intensity (MFI) based on a

minimum of 100 beads per antigen.

Data analysis and statistics

All data preparation, analysis and statistical procedures were conducted using R

Statistical Software (R version 4.3.3) [73]. We used the body weight of the wild-caught *M.*

natalensis individuals at the time of sample collection as a rough proxy for age, which we

subdivided into three categories based on the 1/3 quantiles of weight; juvenile (5 - 26.7g),

subadult (>26.7 - 42g) and adult (>42 - 91g). These weight classes coincide to the expected

sexual maturity, with sexual maturity estimated to occur between 30 - 40g [29,32].

To control for variation between different assay plates and testing days, the MFI

results were transformed to relative antibody units using the positive dilution series as a

standard curve. The MFI result of the positive control starting dilution (i.e., 1:200) was

equalized to 3,125 units and each following dilution step was adjusted proportionally (i.e., the

final dilution step 1:625,000 corresponded to 1 unit). The results of the two sessions were

combined by linear alignment adjustment. This alignment was based on 86 duplicate samples

encompassing the measurable range, allowing the adjustment of the results from the first

session.

Finally, each sample was categorized as a binary value (i.e., 1= positive, 0= negative)

for each of the tested arboviruses. This was done based on whether the unit value exceeded

the mean cutoff value for that specific arbovirus antigen. Five cutoff values were determined

for each arbovirus antigen: I) the mean plus three times the standard deviation of the negative

controls (i.e. *'NegCtrl'*) [18,65]; the change-point analysis, using R package *'changepoint'*

(version: 2.2.4), calculated at most one changepoint based on the II) mean (i.e. *'CHP.m'*), III)

variance (i.e. *'CHP.v'*) and IV) a combination of mean and variance (i.e. *'CHP.mv'*) of wild-

caught samples [74,75] and V) the maximum value of an average antibody curve (i.e. *'Recap'*).

This curve was based on wild-caught individuals that were recaptured at least three times and

showed seroconversion. Seroconversion of an individual was considered when the

individual's maximum unit value was at least four-fold the minimum unit value. This four-fold

increase is a standard seroconversion confirmation measure in human antibody studies [76].

An average antibody curve, with days as the explanatory variable, was created for each

antigen by aligning the maximum unit value of each recaptured seroconverted individual to

the same day. The binary results were used to calculate the seroprevalence for each arbovirus

along with a 95% confidence interval (CI), using the *'binom.exact'* from the package *'binom'*

(version: 1.1.1.1) [77]. The seroprevalence according to the different cutoff methods was

compared to the seroprevalence of the antibody curve cutoff using the *'chisq.test'* from the

package *'stats'* (version 4.3.3) [73]

As an indication of cross-reactivity in antibody response between the tested

arboviruses, pair-wise Pearson correlations were calculated on the binary results, according

to the antibody curve cutoff, of all samples using the *'corr.test'* function of R package *'psych'*

(version: 2.4.1) [78]. The cross-reactivity in antibody response was visualized using the

'heatmap.2' function of the R package *'gplots'* (version: 3.1.3) [79].

A generalized linear model (logit link function and binomial error distribution) was

constructed with the package *'stat'* (version: 4.3.1)*,* with the response variable being the

binary serostatus of each sample [73]. Age (juvenile, subadult and adult), sex and their

interaction were included as explanatory variables. The analysis of variance was performed

using a likelihood ratio test, with p-values calculated assuming a chi-squared distribution.

Pairwise comparison of the seroprevalence was performed between the six combinations of

the explanatory variables (two levels of sex and three levels of age), using the *'emmeans'*

package (version: 1.8.9) [80]. To prevent reporting statistical findings based on the reliance of

an arbitrary p-value of 0.05, we instead present significance in terms of levels of statistical

support based on p-values. P-values exceeding 0.1 are labeled as "no" support and values

around 0.05 (range 0.1 - ≥ 0.01, symbol: *) as "weak" support. "Moderate" support was

assigned to p-values clearly below 0.05 (range < 0.01 - ≥ 0.001, symbol: **), while "strong"

support is reserved for p-values lower than 0.05 (< 0.001, symbol: ***). This representation

in terms of statistical support is based on current statistical reporting practices [81].

Results

In total 1,280 DBS samples were tested of which 660 were female, consisting of 256

juveniles, 172 subadults and 232 adults, 620 samples were male with 199 juveniles, 313

subadults and 108 adults. Samples of recaptured individuals were considered as individual

samples for all analysis.

Seroprevalence

The seroprevalences according to the different cutoff methods showed at least a weak statistical support for a different seroprevalence compared to the antibody curve seroprevalence for almost all arboviruses. Histograms of the data and seroprevalence for each tested arbovirus antigen according to the different cutoff methods is shown in S1 and S2 Figs. The cutoff value according to the antibody curve based on the recaptured seroconverted individuals was used as the main cutoff value for all further calculations.

The overall arbovirus seroprevalence, defined as at least positive for one of the tested arboviruses, except ONNV, was almost 24% (95% CI: 21.89 – 26.66%; N= 1280). ONNV was excluded since the samples of the first session were not screened for antibodies against the ONNV antigen. The seroprevalence for *Flaviviridae* was 20% (95% CI: 17.99 – 22.46%; N= 1280) and for *Togaviridae*, excluding ONNV, almost 7% (95% CI: 5.48 – 8.32%; N= 1280). Overall, seroprevalences ranged from 0.62% for DENV3 (95% CI: 0.27 – 1.23%; N= 1280) and MAYV (95% CI: 0.27 – 1.23%; N= 1280) to 8.44% for DENV2 (95% CI: 6.97 – 10.10%; N= 1280), see Table 2.

Table 2. Total seroprevalence of each arbovirus and virus family in the wild-caught *M. natalensis* sample set.

	Seroprevalence (%)	95% CI (%)	Nr. positive	Cutoff
Arbovirus[a]	24.22	21.89 – 26.66	310	
Bunyaviridae				
RVFV	2.58	1.78 – 3.60	66	37.94
Flaviviridae	20.16	17.99 – 22.46	258	
YFV	2.03	1.33 – 2.96	26	3.45
ZIKV	3.44	2.51 – 4.59	44	15.23
DENV1	7.58	6.19 – 9.17	97	42.11
DENV2	8.44	6.97 – 10.10	108	101.09
DENV3	0.62	0.27 – 1.23	8	32.27
DENV4	5.62	4.43 – 7.03	72	40.66
USUV	4.61	3.53 – 5.91	59	6.47
WNV	2.42	1.65 – 3.42	31	25.60
TBEV	1.64	1.02 – 2.50	21	50.85
WSLV	3.52	2.58 – 4.68	45	4.16

Nairoviridae				
CCHFV	2.81	1.98 – 3.87	36	5513.61
Togaviridae[a]	6.80	5.48 – 8.32	87	
CHIKV	6.17	4.92 – 7.63	79	43.95
MAYV	0.62	0.27 – 1.23	8	11.91
ONNV	2.18	1.30 – 3.42	18	77.10

A 95% confidence interval (CI) is provided and the calculated cutoff value is in units. Sample size was 1280 for each tested arbovirus except for ONNV which had 826 samples. [a] Indicates that ONNV was not included for that calculation.

Pairwise arbovirus serostatus correlation

The correlations in serostatus of samples between the tested arboviruses are visualized in Fig 2. Correlation between two arboviruses is depicted in color scale with the statistical symbol, lower triangle, and the number of positive samples in the upper triangle. The matrix is accompanied by a dendrogram based on the hierarchical clustering of the correlation coefficients. The branch lengths are a proxy for relative distance between arboviruses based on the serostatus response of the samples.

Fig 2. Correlation of the serostatus response between the tested arboviruses with a dendrogram of hierarchical clustering. Symbols in lower triangle represent significance of correlation, values in upper triangle, including diagonal line, represents the number of positive individuals for the corresponding arboviruses.

304  The correlation in sample response between the tested arboviruses ranged from -

305 2.44% for TBEV and ZIKV with no statistical support (p= 0.384) to almost 59% between WSLV

306 and USUV with a strong statistical support (p< 0.001). The dendrogram based on the

307 hierarchical clustering of the correlation showed that WSLV – USUV (correlation= 58.52%; p<

308 0.001), DENV2 – DENV4 (correlation= 56.03%; p< 0.001) and DENV1 – CHIKV (correlation=

309 50.31%; p< 0.001) are relatively closer to each other than to other tested arboviruses.

Model analysis of antibody response

Sex, age, and interaction effects on serostatus

312  The generalized linear model indicated that there was a weak statistical interaction

313 between the effects of sex and age on serostatus for DENV4 (Deviance [Df.= 2; Res.Df.= 1274]= 5.44;

314 p= 0.066) and USUV (Deviance [Df.= 2; Res.Df.= 1274]= 7.72; p= 0.021). For the other tested arboviruses,

315 no support for a statistical interaction was detected, the interaction was thus removed from

316 those models. In the case of RVFV, no statistical support was found for an effect of sex, age,

317 or the interaction on the serostatus. All results from generalized linear model's analysis of

318 variance are reported in S1 Table.

319  The analysis of the sex variable showed a moderate support for males having a lower

320 seroprevalence compared to females for DENV2 (Est. males ± SE= -0.665 ± 0.242; p= 0.006) and WSLV

321 (Est. males ± SE= -1.446 ± 0.483; p= 0.003). A weak statistical effect of a lower seroprevalence in males

322 compared to females was detected in ZIKV (Est. males ± SE= -0.643 ± 0.386; p= 0.096), TBEV (Est. males ±

323 SE= -1.103 ± 0.635; p= 0.083), CCHFV (Est. males ± SE= -0.821 ± 0.460; p= 0.074) and CHIKV (Est. males ± SE= -

324 0.633 ± 0.296; p= 0.032). There was no support for a difference in seroprevalence between males

325 and females for RVFV (Est. males ± SE= 0.097 ± 0.381; p= 0.798), YFV (Est. males ± SE= 0.425 ± 0.413; p= 0.304),

326 DENV1 (Est. males ± SE= -0.377 ± 0.248; p= 0.129), DENV3 (Est. males ± SE= -17.43 ± 2021.76; p= 0.993), WNV

(Est. _males_ ± SE= -0.009 ± 0.397; p= 0.982), MAYV (Est. _males_ ± SE= -1.222 ± 1.080; p= 0.258) and ONNV (Est. _males_

± SE= -17.75 ± 1663.70; p= 0.991).

The analysis of the age variable showed a strong statistical support for a higher

seroprevalence in subadults than in juveniles for DENV1 and DENV2, a moderate support for

CHIKV and a weak support for YFV, ZIKV, TBEV, WSLV and CCHFV. There was no support for a

difference in subadult and juvenile seroprevalence in the other tested arboviruses. A

significantly higher seroprevalence in adults compared to juveniles was shown for ZIKV,

DENV1, DENV2, WSLV and CHIKV with a strong support. A moderate support for a higher

seroprevalence in adults than in juveniles was detected for YFV and CCHFV. Adults showed a

weak statistical support for a higher seroprevalence in contrast to juveniles for WNV and

TBEV. All other tested arboviruses showed no support for a statistical difference between

adults and juveniles. The comparison between subadults and adults showed a strongly

supported statistical difference for DENV1 and CHIKV with a higher seroprevalence in adults.

A moderate support for a higher seroprevalence in adults compared to subadults was

detected for ZIKV and CCHFV. Yellow Fever virus, WNV, WSLV and ONNV showed a weak

support for a statistically higher seroprevalence in adults than in subadults. The other

arboviruses showed no statistically significant difference between adults and subadults. See

Table 3 for estimates, standard errors, and p-values.

Table 3. Difference in coefficient estimate on logit scale between the age levels with standard error (SE).

	Juvenile - Subadult		Juvenile - Adult		Subadult - Adult	
	Estimate ± SE	p-value	Estimate ± SE	p-value	Estimate ± SE	p-value
RVFV	-0.227 ± 0.480	0.637	-0.649 ± 0.446	0.146	-0.422 ± 0.421	0.316
YFV	-1.368 ± 0.794	**0.085**	-2.197 ± 0.757	**0.004**	-0.829 ± 0.449	**0.065**
ZIKV	-1.349 ± 0.656	**0.040**	-2.308 ± 0.611	**< 0.001**	-0.959 ± 0.365	**0.009**
DENV1	-1.610 ± 0.456	**< 0.001**	-2.468 ± 0.434	**< 0.001**	-0.858 ± 0.243	**< 0.001**
DENV2	-1.551 ± 0.356	**< 0.001**	-1.780 ± 0.352	**< 0.001**	-0.229 ± 0.221	0.301
DENV3	-16.47 ± 2176.65	0.994	-18.14 ± 2176.65	0.993	-1.665 ± 1.073	0.121
WNV	0.007 ± 0.582	0.991	-1.20 ± 0.477	**0.012**	-1.207 ± 0.480	**0.012**
TBEV	-1.855 ± 1.083	**0.087**	-2.566 ± 1.039	**0.014**	-0.711 ± 0.499	0.155

WSLV	-0.960 ± 0.539	**0.075**	-1.656 ± 0.492	**< 0.001**	-0.697 ± 0.357	**0.051**
CCHFV	-1.999 ± 1.072	**0.062**	-3.321 ± 1.021	**0.001**	-1.321 ± 0.434	**0.002**
CHIKV	-2.267 ± 0.749	**0.002**	-3.474 ± 0.723	**< 0.001**	-1.207 ± 0.283	**< 0.001**
MAYV	0.642 ± 1.228	0.601	-0.807 ± 0.844	0.339	-1.449 ± 1.104	0.189
ONNV	-17.05 ± 1747.82	0.992	-18.15 ± 1747.82	0.992	-1.099 ± 0.641	**0.087**

Data originates from the pairwise comparison of the age class variables of the generalized linear model. P-values
marked in bold have at least a weak statistical support (p< 0.1).

In the case of DENV4, were there was a weak support for an interaction, the analysis

showed that there was a strong statistical support for a higher seroprevalence in female

adults compared to female juveniles (Est. ~female - adult~ ± SE= 1.844 ± 0.486; p< 0.001). A moderate

support was shown for a higher seroprevalence in female subadults compared to female

juveniles (Est. ~female - subadult~ ± SE= 1.379 ± 0.517; p= 0.008) and a weak support for a higher

seroprevalence in female adults compared to male adults (Est. ~female - adult~ ± SE= 1.898 ± 0.736; p=

0.010). For USUV the model analysis with a weak interaction, showed that there was weak

statistical support for a higher seroprevalence in female adults compared to female subadults

(Est. ~female - adult~ ± SE= 0.868 ± 0.345; p= 0.012) and also a weak support for a higher seroprevalence in

female adults compared to male adults (Est. ~female - adult~ ± SE= 1.515 ± 0.612; p= 0.013).

Fig 3 displays the seroprevalence for the six distinct levels (two levels of sex and three

levels of age) for all tested arboviruses, with statistical support lines based on the log odds.

Fig 3. Seroprevalence according to sex and age combinations with 95% confidence error bars for each arbovirus. Statistical support on seroprevalence difference is indicated by asterisks in the horizontal lines. Sample size: 660 females of which 256 juveniles, 172 subadults and 232 adult and 620 males with 199 juveniles, 313 subadults and 108 adults. Sample sizes for ONNV are 560 females: 150 juveniles, 150 subadults, 260 adults and 266 males: 94 juveniles, 92 subadults and 80 adults.

Discussion

In this study we optimized a high-throughput multiplex immunoassay for the simultaneous detection of IgG antibodies against 15 medically relevant arboviruses and used it to investigate the potential of *M. natalensis* to serve as a host for arboviruses. We describe the screening results of a historic set of wild *M. natalensis* DBS samples. Our results revealed an overall seroprevalence of 24% against the entire panel of tested antigens. Virus family-specific seroprevalences were approximately 2.6%, 20%, 2.8% and 7% for respectively *Bunyaviridae*, *Flaviviridae*, *Nairoviridae* and *Togaviridae*. We further found that female rodents were more likely to be classified as antibody positive for eight of the 15 tested arboviruses. Additionally, positivity increased significantly with age for almost all tested arboviruses.

The lack of realistic natural positive controls limits the possibility to determine a true cutoff, we therefore used recaptured seroconverted individuals to determine a cutoff value. The use of antibody titers at multiple time points are a standard practice to determine antibody or pathogen development and (sero)conversions in human studies [76]. However, multiple samples of an individual animal across time are often impossible or very difficult in wildlife studies. Our study is unique in that regard that we have measurements of individual recaptured *M. natalensis*. We consider that our cutoff based on seroconverted individuals is a good proxy for the natural cutoff value, since it is based on similar methods as in human studies [76]. We tried to show in our analysis that the tested mathematical methods could approximate this calculated cutoff and thus provide a method for future studies that do not have access to recaptured seroconverted wildlife samples. Unfortunately, the tested cutoff methods did not significantly approximate the seroprevalence according to the cutoff using

samples from recaptures. The negative control-based cutoff (i.e., the mean plus three times

the standard deviation of the negative control samples) gave unrealistic high seroprevalences.

This can be explained by the fact that the negative control samples originate from a breeding

colony. These animals have thus never been exposed to a natural environment and the

pathogens that occur in the environment. The statistical methods vary in their seroprevalence

with some methods approximating the estimated seroprevalence according to the recaptured

cutoff. This high degree of variation makes it difficult to decide on one method that works for

all the tested arboviruses. The cutoff value for CCHFV seems extremely high compared to the

other arboviruses, but the unit values for CCHFV are also much higher than for the other

arboviruses (see S1 Fig). The reason is that the unit values are calculated based on the positive

dilution series and the positive controls for CCHFV were not of the same magnitude as for the

other arboviruses. The value in determining a cutoff and the resulting seroprevalence is that

it allows the comparison of results with previous and future studies on arbovirus

seroprevalence in rodents or other wildlife. We are aware that the used cutoff and resulting

seroprevalences could be an over- or underestimation and might not reflect the natural

arbovirus seroprevalence. We therefore encourage future research to investigate and

compare different cutoff methods for arbovirus (or pathogen) antibody detection in wildlife

studies.

The detection of antibodies against each of 15 tested arbovirus antigens indicates that

these arboviruses, or closely related viruses, are present in *M. natalensis*. The overall

arbovirus seroprevalence of 24% suggests that this rodent species is commonly infected with

one or more arboviruses and that it could thus play a significant role in virus transmission and

persistence. Our results corroborate previous studies, which detected USUV and WNV RNA in

respectively *M. natalensis* and *M. erythroleucus*, in Senegal [24,54]. Besides in this genus,

arboviral RNA has also been found in other rodents in Africa, such as *Rattus rattus* for USUV

and WSLV and *Desmodillus auricularis* for WSLV [22,24,53]. The findings in our study thus

further corroborate that arboviruses are likely present in rodents, and specifically in the

ubiquitous *M. natalensis*. The demographic and ecological characteristics of *M. natalensis*

may have particularly important implications for arbovirus transmission. The population

densities of *M. natalensis* in Tanzania are strongly dependent on weather conditions. More

specifically, early rainfall and elevated temperatures lead to an exponential growth in the

population density, due to an increase influx of juveniles [29,30,32]. The rainfall and increased

temperatures are also beneficial for the breeding of mosquitoes and the multiplication of

arboviruses within these vectors [82]. Further, *M. natalensis* is highly abundant around

houses and in the crop fields at the fringes of the villages. These factors increase the likelihood

of arbovirus outbreaks in *M. natalensis* populations, with the possibility of spillover to

humans.

Arboviruses that show the highest seroprevalence are DENV1, DENV2, DENV4 and

CHIKV, with seroprevalences between five to nine percent. These seroprevalences could be

caused by cross-reactivity due to antibodies of other dengue virus serotypes or other

flaviviruses binding to the non-structural protein 1 (NS1 protein) of DENV1, DENV2 and

DENV4. The same effect could also be true for alphaviruses binding to the envelope protein 2

(E2 protein) of CHIKV. Whether these seroprevalences are indeed due to the presence of the

arbovirus specific antibodies or a related arbovirus remains to be investigated. Nonetheless,

it indicates that a part of the sampled *M. natalensis* population in Morogoro is exposed to

dengue virus and CHIKV or respectively to a related flavivirus and alphavirus. This hypothesis

is supported by the fact that flavi- and alphaviruses are the most prevalent arboviral genera

in humans, compared to other arbovirus genera, and potentially thus also in rodents involved

in the sylvatic cycle [83,84].

A recent health survey has shown that, in the same region as where our rodent

population was sampled, a high percentage of the human population is seropositive for CHIKV

(9.83%) [55]. Another study in the same region reported acute infection of CHIKV in 1.28% of

patients with fever and malaria-like symptoms [85]. Although these studies have not found

any indication of dengue virus in humans, a large-scale cross-sectional study in Tanzania has

found CHIKV and dengue virus antibodies in respectively 28.0% and 16.1% of the population

[56]. These studies clearly indicate that the human population in Tanzania is exposed to

arboviruses and then specifically to CHIKV and DENV.

The cross-reactivity analysis via the correlation matrix and hierarchical clustering (Fig

2) showed an antibody response correlation between WSLV – USUV (59%), DENV2 – DENV4

(56%) and DENV1 – CHIKV (50%). We expected that phylogenetically related arboviruses

would show elevated levels of correlation due to cross-reactivity [86]. A remarkable result in

this cross-reactivity analysis is that DENV1 – CHIKV cluster together with a correlation of 50%,

based on the serostatus of the tested samples. The branch DENV1/CHIKV clusters also closer

to ONNV than to the branches of WSLV/USUV and DENV2/DENV4. This is unexpected since

CHIKV belongs to the *Togaviridae* and DENV1 to the *Flaviviridae* [87]. The proteins used for

the antibody detection are also two different proteins, with the E2 protein used for the

Togaviridae and NS1 protein for the *Flaviviridae*, thus limiting the possibility of cross-

reactivity. Although we cannot exclude that there might be similar epitopes between the

different proteins, but other studies have already indicated that cross-reactivity between the

E2 protein of the *Togaviridae* and NS1 protein of the *Flaviviridae* is limited [88,89]. Given that

both *Togaviridae* and *Flaviviridae* viruses are circulating in humans in East Africa, we

hypothesize that these viral families may also both be present in rodents [5,10]. More

specifically, it is plausible that both viral families could be found in *M. natalensis,* where

pathogen co-infections are common [49]. This hypothesis is further supported by the fact that

some viruses in both families are transmitted by the same arthropod vectors, such as *Aedes*

aegypti and *Aedes albopictus* for both dengue virus and CHIKV [4,82].

For some of the tested arboviruses we found statistical support for a higher

seroprevalence in females than in males. This result is supported by previous studies where

it is shown that female mice have a stronger innate immune response than male mice [90].

Other animals (e.g., birds, fish, insects) as well as humans also display stronger immune

responses in females [91–95]. The major driving forces behind these immune differences are

genetic (i.e., X-chromosomes) and hormonal (i.e., different estrogen and testosterone levels)

[96]. In the case of *M. natalensis*, behavioral differences could also be the cause for this

divergence in seroprevalence. Previous studies have already shown that home range,

behavior and pathogen presence differ between male and female *M. natalensis* [48,97].

Besides the sex effects, we also found statistical support for a positive age effect on the

presence of antibodies in some of the tested arboviruses. This increased seroprevalence with

age corroborates previous findings for other pathogens (i.e., *Bartonella sp.*, *Anaplasma sp.*,

helminths, and arenaviruses) [48,49,98]. This age effect further supports our hypothesis that

M. natalensis is exposed to arboviruses and that individuals develop antibodies and gain

immunity via repeated exposures throughout their life. To maintain the arbovirus

transmission among the *M. natalensis* population, there needs to be a proportion of the

population that is either chronically infected or immunologically naïve. Chronic infections in

M. natalensis have already been documented for mammarenaviruses [31,99]. However, as

far as we are aware naturally occurring chronic arbovirus infections have not been reported

in humans or non-human vertebrates. Therefore, the presumable driving factor in sustained

transmission is the presence of immunologically naïve individuals. During the breeding

season, which coincides with increased rainfall and temperature, there is an influx of

immunologically naïve juveniles. This influx can reach high proportions during population

outbreak periods [30,64]. We thus expect that it is mainly juveniles who are the major factor

in sustaining the arbovirus transmission cycle. We predict that the prevalence of arboviral

genetic material will be higher in juveniles than in adults, since juveniles do not possess the

necessary antibodies to fight of the infection.

We conclude from our detected antibody responses that arboviruses, or related

viruses, are present in *M. natalensis* in Morogoro, Tanzania. The higher seroprevalence we

detect in females can be explained by genetic, hormonal, ecological and/or behavioral

differences between sexes. Individuals are exposed to these viruses throughout their life and

gain immunity as they age. We hypothesize that juvenile *M. natalensis* play an essential role

in sustaining arbovirus transmission as they are immunologically naïve and can reach high

densities in favorable climate conditions that coincide with optimal vector conditions. More

extensive screening, such as virus neutralization tests and molecular screening of these

viruses within *M. natalensis* are necessary to quantify the contribution of this rodent species

in the arbovirus transmission cycle.

Acknowledgements

503 The Ethical Committee for Animal Testing at UAntwerp approved all experiments

504 (ECD2021-79 and ECD2023-08). This study was funded by The Research Foundation – Flanders

505 (FWO) through the Senior research project G054820N (to KKA, EV and MP) and PhD fellowship

506 1171023N (to WDK).

References

[1] Boutayeb A. The Impact of Infectious Diseases on the Development of Africa. Handbook of Disease Burdens and Quality of Life Measures, 2010. https://doi.org/10.1007/978-0-387-78665-0_66.

[2] Nkengasong JN, Tessema SK. Africa Needs a New Public Health Order to Tackle Infectious Disease Threats. Cell 2020;183. https://doi.org/10.1016/j.cell.2020.09.041.

[3] Minakshi P, Brar B, Lambe UP, Ranjan K, Prasad G, Harimohan, et al. RNA viruses: Greatest global threat and one health solutions. Virusdisease 2019;30.

[4] Huang Y-JS, Higgs S, Vanlandingham DL. Emergence and re-emergence of mosquito-borne arboviruses. Curr Opin Virol 2019;34:104–9.

[5] Venter M. Assessing the zoonotic potential of arboviruses of African origin. Curr Opin Virol 2018;28:74–84.

[6] Omondi D, Masiga DK, Ajamma YU, Fielding BC, Njoroge L, Villinger J. Unraveling host-vector-arbovirus interactions by two-gene high resolution melting mosquito bloodmeal analysis in a Kenyan wildlife-livestock interface. PLoS One 2015;10:e0134375.

[7] Musa AA, Muturi MW, Musyoki AM, Ouso DO, Oundo JW, Makhulu EE, et al. Arboviruses and blood meal sources in zoophilic mosquitoes at human-wildlife interfaces in Kenya. Vector-Borne and Zoonotic Diseases 2020;20:444–53.

[8] Thompson P, van Den Bergh C, Venter E, Schade M, Swanepoel R. Co-circulation of Rift Valley fever virus and other zoonotic arboviruses at the human-livestock-wildlife interface in KwaZulu-Natal, South Africa. International Journal of Infectious Diseases 2020;101:535. https://doi.org/https://doi.org/10.1016/j.ijid.2020.09.1387.

[9]   Pierson TC, Diamond MS. The continued threat of emerging flaviviruses. Nat Microbiol

2020. https://doi.org/10.1038/s41564-020-0714-0.

[10]   Mayer S V., Tesh RB, Vasilakis N. The emergence of arthropod-borne viral diseases: A

global prospective on dengue, chikungunya and zika fevers. Acta Trop 2017.

https://doi.org/10.1016/j.actatropica.2016.11.020.

[11]   Gould E, Pettersson J, Higgs S, Charrel R, De Lamballerie X. Emerging arboviruses: why

today? One Health 2017;4:1–13.

[12]   Gould EA, Higgs S. Impact of climate change and other factors on emerging arbovirus

diseases.   Trans   R   Soc   Trop   Med   Hyg   2009;103.

https://doi.org/10.1016/j.trstmh.2008.07.025.

[13]   Shahhosseini N, Wong G, Babuadze G, Camp J V, Ergonul O, Kobinger GP, et al. Crimean-

Congo hemorrhagic fever virus in Asia, Africa and Europe. Microorganisms

2021;9:1907.

[14]   Akash S, Islam MdR, Rahman MdM. Rift Valley fever (RVF): a re-emerging zoonotic

disease, pathogenesis, epidemiology, current status, and future perspective –

correspondence. International Journal of Surgery 2023;109.

[15]   Bernstein AS, Ando AW, Loch-Temzelides T, Vale MM, Li B V, Li H, et al. The costs and

benefits of primary prevention of zoonotic pandemics. Sci Adv 2022;8:eabl4183.

[16]   Vora NM, Hannah L, Lieberman S, Vale MM, Plowright RK, Bernstein AS. Want to

prevent pandemics? Stop spillovers. Nature 2022;605:419–22.

[17]   Gubler DJ. The global emergence/resurgence of arboviral diseases as public health

problems. Arch Med Res, 2002. https://doi.org/10.1016/S0188-4409(02)00378-8.

[18]   Raulino R, Thaurignac G, Butel C, Villabona-Arenas CJ, Foe T, Loul S, et al. Multiplex

detection of antibodies to Chikungunya, O'nyong-nyong, Zika, Dengue, West Nile and

Usutu viruses in diverse non-human primate species from Cameroon and the

Democratic Republic of Congo. PLoS Negl Trop Dis 2021;15:e0009028-.

[19] Eastwood G, Sang RC, Guerbois M, Taracha ELN, Weaver SC. Enzootic circulation of

chikungunya virus in East Africa: Serological evidence in non-human Kenyan primates.

Am J Trop Med Hyg 2017;97:1399–404.

[20] Valentine MJ, Murdock CC, Kelly PJ. Sylvatic cycles of arboviruses in non-human

primates. Parasit Vectors 2019;12:463.

[21] Olive M-M, Goodman SM, Reynes J-M. The role of wild mammals in the maintenance

of Rift Valley fever virus. J Wildl Dis 2012;48:241–66.

[22] Diagne MM, Faye M, Faye O, Sow A, Balique F, Sembène M, et al. Emergence of

Wesselsbron virus among black rat and humans in Eastern Senegal in 2013. One Health

2017. https://doi.org/10.1016/j.onehlt.2017.02.001.

[23] Gora D, Yaya T, Jocelyn T, Didier F, Maoulouth D, Amadou S, et al. The potential role of

rodents in the enzootic cycle of Rift Valley fever virus in Senegal. Microbes Infect 2000.

https://doi.org/10.1016/S1286-4579(00)00334-8.

[24] Diagne MM, Ndione MHD, Di Paola N, Fall G, Bedekelabou AP, Sembène PM, et al.

Usutu virus isolated from rodents in Senegal. Viruses 2019;11:181.

[25] Dahmana H, Granjon L, Diagne C, Davoust B, Fenollar F, Mediannikov O. Rodents as

hosts of pathogens and related zoonotic disease risk. Pathogens 2020;9:202.

[26] Fiedler LA. Rodent problems in Africa. Rodent pest management, CRC Press; 2018, p.

35–65.

[27] Hánová A, Konečný A, Mikula O, Bryjová A, Šumbera R, Bryja J. Diversity, distribution,

and evolutionary history of the most studied African rodents, multimammate mice of

577  the genus Mastomys: An overview after a quarter of century of using DNA sequencing.

578  Journal of Zoological Systematics and Evolutionary Research 2021;59:2500–18.

579 [28] Chidodo S, Kilawe CJ, Mnyone LL, Broecke B Vanden, Mulungu LS. Factors affecting the

580  composition of rodent assemblages in the North Uluguru Mountains, Tanzania. J

581  Vertebr Biol 2020;69. https://doi.org/10.25225/jvb.20047.

582 [29] Leirs H, Verhagen R, Verheyen W. The basis of reproductive seasonally in Mastomys

583  rats (Rodentia: Muridae) in Tanzania. J Trop Ecol 1994;10:55–66.

584 [30] Sluydts V, Crespin L, Davis S, Lima M, Leirs H. Survival and maturation rates of the

585  African rodent, Mastomys natalensis: Density-dependence and rainfall. Integr Zool

586  2007;2:220–32.

587 [31] Mariën J, Borremans B, Verhaeren C, Kirkpatrick L, Gryseels S, de Bellocq J, et al.

588  Density dependence and persistence of Morogoro arenavirus transmission in a

589  fluctuating population of its reservoir host. Journal of Animal Ecology 2020;89:506–18.

590  https://doi.org/https://doi.org/10.1111/1365-2656.13107.

591 [32] Leirs H, Stuyck J, Verhagen R, Verheyen W. Seasonal variation in growth of Mastomys

592  natalensis (Rodentia: Muridae) in Morogoro, Tanzania. Afr J Ecol 1990;28:298–306.

593 [33] Mwanjabe PS, Sirima FB, Lusingu J. Crop losses due to outbreaks of Mastomys

594  natalensis (Smith, 1834) Muridae, Rodentia, in the Lindi region of Tanzania. Int

595  Biodeterior Biodegradation 2002;49. https://doi.org/10.1016/S0964-8305(01)00113-

596  5.

597 [34] Stenseth NC, Leirs H, Skonhoft A, Davis SA, Pech RP, Andreassen HP, et al. Mice, rats,

598  and people: The bio-economics of agricultural rodent pests. Front Ecol Environ 2003;1.

599  https://doi.org/10.1890/1540-9295(2003)001[0367:MRAPTB]2.0.CO;2.

[35]   Mulungu LS. Control of rodent pests in maize cultivation: the case of Africa, 2017.

https://doi.org/10.19103/as.2016.0002.18.

[36]   Ziwa MH, Matee MI, Hang'Ombe BM, Lyamuya EF, Kilonzo BS. Plague in Tanzania: An

overview. Tanzan J Health Res 2013;15. https://doi.org/10.4314/thrb.v15i4.7.

[37]   Holt J, Davis S, Leirs H. A model of Leptospirosis infection in an African rodent to

determine risk to humans: Seasonal fluctuations and the impact of rodent control. Acta

Trop 2006. https://doi.org/10.1016/j.actatropica.2006.08.003.

[38]   Sadlova J, Vojtkova B, Hrncirova K, Lestinova T, Spitzova T, Becvar T, et al. Host

competence of African rodents Arvicanthis neumanni, A. niloticus and Mastomys

natalensis for Leishmania major. Int J Parasitol Parasites Wildl 2019;8.

https://doi.org/10.1016/j.ijppaw.2019.01.004.

[39]   Laudisoit A, Leirs H, Makundi R, Krasnov BR. Seasonal and habitat dependence of fleas

parasitic on small mammals in Tanzania. Integr Zool 2009;4.

https://doi.org/10.1111/j.1749-4877.2009.00150.x.

[40]   Schwan TG, Lopez JE, Safronetz D, Anderson JM, Fischer RJ, Maïga O, et al. Fleas and

trypanosomes of peridomestic small mammals in sub-Saharan Mali. Parasit Vectors

2016;9. https://doi.org/10.1186/s13071-016-1818-5.

[41]   Brouat C, Duplantier JM. Host habitat patchiness and the distance decay of similarity

among gastro-intestinal nematode communities in two species of Mastomys

(southeastern Senegal). Oecologia 2007;152. https://doi.org/10.1007/s00442-007-

0680-8.

[42]   Brouat C, Kane M, Diouf M, Bâ K, Sall-Dramé R, Duplantier JM. Host ecology and

variation in helminth community structure in Mastomys rodents from Senegal.

Parasitology 2007;134. https://doi.org/10.1017/S003118200600151X.

[43]   Ribas A, López S, Makundi RH, Leirs H, De Bellocq JG. Trichuris spp. (Nematoda:

625   Trichuridae) from two rodents, Mastomys natalensis and Gerbilliscus vicinus in

626   Tanzania. Journal of Parasitology 2013;99. https://doi.org/10.1645/12-151.1.

627   [44]   Ribas A, Diagne C, Tatard C, Diallo M, Poonlaphdecha S, Brouat C. Whipworm diversity

628   in West African rodents: a molecular approach and the description of Trichuris

629   duplantieri n. sp. (Nematoda: Trichuridae). Parasitol Res 2017;116.

630   https://doi.org/10.1007/s00436-017-5404-3.

631   [45]   Ribas A, Makundi RH, Bellocq JG De. Paraconcinnum leirsi n.sp. (Trematoda:

632   Dicrocoeliidae) from rodents in Tanzania and its phylogenetic position within the

633   dicrocoeliids. Afr Zool 2012;47. https://doi.org/10.3377/004.047.0219.

634   [46]   Diouf M, Diagne CA, Quilichini Y, Dobigny G, Garba M, Marchand B. Pterygodermatites

635   (mesopectines) niameyensis n. Sp. (nematoda: Rictulariidae), a parasite of mastomys

636   natalensis (smith, 1834) (rodentia: Muridae) from Niger. Journal of Parasitology

637   2013;99. https://doi.org/10.1645/13-204.1.

638   [47]   Diagne CA, Charbonnel N, Henttonen H, Sironen T, Brouat C. Serological Survey of

639   Zoonotic Viruses in Invasive and Native Commensal Rodents in Senegal, West Africa.

640   Vector-Borne and Zoonotic Diseases 2017;17. https://doi.org/10.1089/vbz.2017.2135.

641   [48]   Vanden Broecke B, Bernaerts L, Ribas A, Sluydts V, Mnyone L, Matthysen E, et al.

642   Linking Behavior, Co-infection Patterns, and Viral Infection Risk With the Whole

643   Gastrointestinal Helminth Community Structure in Mastomys natalensis. Front Vet Sci

644   2021;8. https://doi.org/10.3389/fvets.2021.669058.

645   [49]   Vanden Broecke B, Tafompa PJJ, Mwamundela BE, Bernaerts L, Ribas A, Mnyone LL, et

646   al. Drivers behind co-occurrence patterns between pathogenic bacteria, protozoa, and

helminths in populations of the multimammate mouse, Mastomys natalensis. Acta

Trop 2023;243. https://doi.org/10.1016/j.actatropica.2023.106939.

[50] Haikukutu L, Lyaku JR, Lyimo C, Kasanga CJ, Kandusi SE, Rahelinirina S, et al. Plague in

Tanzania: first report of sylvatic plague in Morogoro region, persistence in Mbulu focus,

and ongoing quiescence in Lushoto and Iringa foci. IJID Regions 2022;4:105–10.

[51] Calvignac-Spencer S, Kouadio L, Couacy-Hymann E, Sogoba N, Rosenke K, Davison AJ,

et al. Multiple DNA viruses identified in multimammate mouse (Mastomys natalensis)

populations from across regions of sub-Saharan Africa. Arch Virol 2020;165:2291–9.

[52] Oguge N, Rarieya M, Ondiaka P. A preliminary survey of macroparasite communities of

rodents of Kahawa, Central Kenya. Belgian Journal of Zoology (Belgium) 1997;127.

[53] Weyer J, Thomas J, Leman PA, Grobbelaar AA, Kemp A, Paweska JT. Human cases of

wesselsbron disease, South Africa 2010-2011. Vector-Borne and Zoonotic Diseases

2013;13. https://doi.org/10.1089/vbz.2012.1181.

[54] Ndione MHD, Ndiaye EH, Faye M, Diagne MM, Diallo D, Diallo A, et al. Re-Introduction

of West Nile Virus Lineage 1 in Senegal from Europe and Subsequent Circulation in

Human and Mosquito Populations between 2012 and 2021. Viruses 2022;14.

https://doi.org/10.3390/v14122720.

[55] Budodo RM, Horumpende PG, Mkumbaye SI, Mmbaga BT, Mwakapuja RS, Chilongola

JO. Serological evidence of exposure to Rift Valley, Dengue and Chikungunya Viruses

among agropastoral communities in Manyara and Morogoro regions in Tanzania: A

community survey. PLoS Negl Trop Dis 2020;14:e0008061.

[56] Mwanyika GO, Sindato C, Rugarabamu S, Rumisha SF, Karimuribo ED, Misinzo G, et al.

Seroprevalence and associated risk factors of chikungunya, dengue, and Zika in eight

districts in Tanzania. International Journal of Infectious Diseases 2021;111:271–80.

[57] Chipwaza B, Mugasa JP, Selemani M, Amuri M, Mosha F. Dengue and Chikungunya

Fever among Viral Diseases in Outpatient Febrile 2014.

[58] Mboera LEG, Mweya CN, Rumisha SF, Tungu PK, Stanley G, Makange MR, et al. The Risk

of Dengue Virus Transmission in Dar es Salaam, Tanzania during an Epidemic Period of

2014. PLoS Negl Trop Dis 2016. https://doi.org/10.1371/journal.pntd.0004313.

[59] Faustine NL, Sabuni EJ, Ndaro AJ, Paul E, Chilongola JO. Chikungunya, Dengue and West

Nile virus Infections in Northern Tanzania. J Adv Med Med Res 2017:1–7.

[60] Mitchell CL, Ngasala B, Janko MM, Chacky F, Edwards JK, Pence BW, et al. Evaluating

malaria prevalence and land cover across varying transmission intensity in Tanzania

using a cross-sectional survey of school-aged children. Malar J 2022;21.

https://doi.org/10.1186/s12936-022-04107-8.

[61] Chipwaza B, Mugasa JP, Selemani M, Amuri M, Mosha F, Ngatunga SD, et al. Dengue

and Chikungunya fever among viral diseases in outpatient febrile children in Kilosa

district hospital, Tanzania. PLoS Negl Trop Dis 2014;8:e3335.

[62] Borremans B, Hughes NK, Reijniers J, Sluydts V, Katakweba AAS, Mulungu LS, et al.

Happily together forever: temporal variation in spatial patterns and complete lack of

territoriality in a promiscuous rodent. Popul Ecol 2014;56:109–18.

[63] Vanden Broecke B, Mariën J, Sabuni CA, Mnyone L, Massawe AW, Matthysen E, et al.

Relationship between population density and viral infection: A role for personality?

Ecol Evol 2019;9:10213–24.

[64] Leirs H, Kirkpatrick L, Sluydts V, Sabuni C, Borremans B, Katakweba A, et al. Twenty-

nine years of continuous monthly capture-mark-recapture data of multimammate

mice (Mastomys natalensis) in Morogoro, Tanzania. Sci Data 2023;10:798.

[65] Raulino R, Thaurignac G, Keita AK, Esteban A, Goumou S, Diallo R, et al. Seroprevalence

of IgG antibodies against multiple arboviruses in bats from Cameroon, Guinea, and the

Democratic Republic of Congo. Vector-Borne and Zoonotic Diseases 2022;22:252–62.

[66] Amorim JH, Diniz MO, Cariri FAMO, Rodrigues JF, Bizerra RSP, Gonçalves AJS, et al.

Protective immunity to DENV2 after immunization with a recombinant NS1 protein

using a genetically detoxified heat-labile toxin as an adjuvant. Vaccine 2012;30:837–

45.

[67] Li Y, Counor D, Lu P, Duong V, Yu Y, Deubel V. Protective immunity to Japanese

encephalitis virus associated with anti-NS1 antibodies in a mouse model. Virol J

2012;9:1–13.

[68] Bailey MJ, Broecker F, Duehr J, Arumemi F, Krammer F, Palese P, et al. Antibodies

elicited by an NS1-based vaccine protect mice against Zika virus. MBio 2019;10:10–

1128.

[69] Easterbrook JD, Klein SL. Immunological mechanisms mediating hantavirus persistence

in   rodent   reservoirs.   PLoS   Pathog   2008;4.

https://doi.org/10.1371/journal.ppat.1000172.

[70] Spengler JR, Haddock E, Gardner D, Hjelle B, Feldmann H, Prescott J. Experimental

Andes Virus Infection in Deer Mice: Characteristics of Infection and Clearance in a

Heterologous   Rodent   Host.   PLoS   One   2013;8.

https://doi.org/10.1371/journal.pone.0055310.

[71] Borremans B. Ammonium improves elution of fixed dried blood spots without affecting

immunofluorescence assay quality. Tropical Medicine and International Health

2014;19. https://doi.org/10.1111/tmi.12259.

[72] Ayouba A, Thaurignac G, Morquin D, Tuaillon E, Raulino R, Nkuba A, et al. Multiplex

detection and dynamics of IgG antibodies to SARS-CoV2 and the highly pathogenic

human coronaviruses SARS-CoV and MERS-CoV. Journal of Clinical Virology

2020;129:104521.

[73] R Core Team. R: A Language and Environment for Statistical Computing. R Foundation

for Statistical Computing, Vienna, Austria 2024.

[74] Killick R, Eckley IA. Changepoint: An R package for changepoint analysis. J Stat Softw

2014;58. https://doi.org/10.18637/jss.v058.i03.

[75] Donnelly Jr RA, Abdel-Raouf F. Statistics, 3E. Penguin; 2016.

[76] Piantadosi A, Kanjilal S. Diagnostic approach for arboviral infections in the united

states. J Clin Microbiol 2020;58. https://doi.org/10.1128/JCM.01926-19.

[77] Dorai-Raj S. binom: Binomial confidence intervals for several parameterizations. R

Package Version 2014;1.

[78] Revelle W. Package "psych" - Procedures for Psychological, Psychometric and

Personality Research. R Package 2015.

[79] Warnes GR, Bolker B, Bonebakker L, Gentleman R, Huber W, Liaw WHA, et al. gplots:

Various R Programming Tools for Plotting Data. R Package Version 313 2022.

[80] Lenth R V. emmeans: Estimated Marginal Means, aka Least-Squares Means. R Package

Version 189 2023.

[81] Muff S, Nilsen EB, O'Hara RB, Nater CR. Rewriting results sections in the language of

evidence. Trends Ecol Evol 2022;37. https://doi.org/10.1016/j.tree.2021.10.009.

[82] Chandra G, Mukherjee D. Effect of climate change on mosquito population and

changing pattern of some diseases transmitted by them. Advances in Animal

Experimentation and Modeling, Elsevier; 2022, p. 455–60.

[83] Mwanyika GO, Mboera LEG, Rugarabamu S, Ngingo B, Sindato C, Lutwama JJ, et al.

Dengue virus infection and associated risk factors in africa: A systematic review and

meta-analysis. Viruses 2021;13. https://doi.org/10.3390/v13040536.

[84] Madewell ZJ. Arboviruses and their vectors. South Med J 2020;113.

https://doi.org/10.14423/SMJ.0000000000001152.

[85] Mboya LB, Nkya ES, Matemba LE, Kinimi E. Evidence of Chikungunya but not Dengue

Virus Circulating among Febrile Patients during Low Transmission Period in Morogoro

Municipality, Tanzania. Int J Trop Dis Health 2020.

https://doi.org/10.9734/ijtdh/2019/v40i430236.

[86] Kasbergen LMR, Nieuwenhuijse DF, de Bruin E, Sikkema RS, Koopmans MPG. The

increasing complexity of arbovirus serology: An in-depth systematic review on cross-

reactivity. PLoS Negl Trop Dis 2023;17:e0011651.

[87] Lefkowitz EJ, Dempsey DM, Hendrickson RC, Orton RJ, Siddell SG, Smith DB. Virus

taxonomy: The database of the International Committee on Taxonomy of Viruses

(ICTV). Nucleic Acids Res 2018;46. https://doi.org/10.1093/nar/gkx932.

[88] Cho B, Jeon BY, Kim J, Noh J, Kim J, Park M, et al. Expression and evaluation of

Chikungunya virus E1 and E2 envelope proteins for serodiagnosis of chikungunya virus

infection. Yonsei Med J 2008;49. https://doi.org/10.3349/ymj.2008.49.5.828.

[89] Chang H-H, Huber RG, Bond PJ, Grad YH, Camerini D, Maurer-Stroh S, et al. Systematic

analysis of protein identity between Zika virus and other arthropod-borne viruses. Bull

World Health Organ 2017;95. https://doi.org/10.2471/blt.16.182105.

[90] Fink AL, Engle K, Ursin RL, Tang W-Y, Klein SL. Biological sex affects vaccine efficacy and

protection against influenza in mice. Proceedings of the National Academy of Sciences

2018;115:12477–82.

[91] Chrousos GP. Stress and sex versus immunity and inflammation. Sci Signal
2010;3:pe36–pe36.

[92] Kelly CD, Stoehr AM, Nunn C, Smyth KN, Prokop ZM. Sexual dimorphism in immunity
across animals: a meta-analysis. Ecol Lett 2018;21:1885–94.

[93] Vincze O, Vágási CI, Pénzes J, Szabó K, Magonyi NM, Czirják GÁ, et al. Sexual
dimorphism in immune function and oxidative physiology across birds: The role of
sexual selection. Ecol Lett 2022;25:958–70.

[94] Jacobsen H, Klein SL. Sex differences in immunity to viral infections. Front Immunol.
2021; 12: 720952 2021.

[95] Sciarra F, Campolo F, Franceschini E, Carlomagno F, Venneri MA. Gender-Specific
Impact of Sex Hormones on the Immune System. Int J Mol Sci 2023;24:6302.

[96] Wilkinson NM, Chen H-C, Lechner MG, Su MA. Sex differences in immunity. Annu Rev
Immunol 2022;40:75–94.

[97] Mlyashimbi ECM, Mlyashimbi ECM, Mariën J, Kimaro DN, Tarimo AJP, Machang'U RS,
et al. Home ranges, sex ratio and recruitment of the multimammate rat (Mastomys
natalensis) in semi-arid areas in Tanzania. Mammalia 2020;84.
https://doi.org/10.1515/mammalia-2019-0048.

[98] Borremans B, Leirs H, Gryseels S, Günther S, Makundi R, De Bellocq JG. Presence of
Mopeia virus, an African arenavirus, related to biotope and individual rodent host
characteristics: Implications for virus transmission. Vector-Borne and Zoonotic
Diseases 2011;11. https://doi.org/10.1089/vbz.2010.0010.

[99] Borremans B, Vossen R, Becker-Ziaja B, Gryseels S, Hughes N, Van Gestel M, et al.
Shedding dynamics of Morogoro virus, an African arenavirus closely related to Lassa
virus, in its natural reservoir host Mastomys natalensis. Sci Rep 2015;5:1–8.

Supporting information

**S1 Fig. Histograms of wild-caught *M. natalensis* for each tested arbovirus with on the x-axis the relative
antibody units in a logarithmic scale.** The relative antibody units are calculated according to the positive control
dilution series. The calculated cutoff values are represented by the colored vertical lines: 'CHP.m' is the
changepoint mean, 'CHP.mv' is the changepoint mean-variance, 'CHP.v' is the changepoint variance, 'NegCtrl' is
the mean plus three times the standard deviation of the negative control samples and 'Recap' is the maximum
value of an antibody development curve based on recaptured seroconverted wild-caught *M. natalensis*.

**S2 Fig. Seroprevalence, according to the calculated cutoff methods, of the wild-caught *M. natalensis* with 95%
confidence interval for each of the tested arboviruses.** The cutoff methods: 'CHP.m' is the changepoint mean,
'CHP.mv' is the changepoint mean-variance, 'CHP.v' is the changepoint variance, 'NegCtrl' is the mean plus three
times the standard deviation of the negative control samples and 'Recap' is the maximum value of an antibody
development curve based on recaptured seroconverted wild-caught *M. natalensis*. Each calculated
seroprevalence was compared to the 'Recap' seroprevalence using a Chi-square test, significant difference
is depicted in asterisk (*) symbols. P-values: * $0.1 - \geqslant 0.01$; ** $< 0.01 - \geqslant 0.001$; *** < 0.001.

**S1 Table. Analysis of variance from the generalized linear model (logit link function and binomial error
distribution) with the response variable being the binary serostatus of each sample.** Sex and age and their
interaction were included as explanatory variables. P values with at least a weak statistical support are marked
in bold ($p < 0.1$).