Sharp transitions in phytoplankton communities across estuarine to open ocean waters of

the tropical Pacific

Sarah J. Tucker [1,2,3,#] https://orcid.org/0000-0003-0541-0853

Yoshimi M. Rii[1,3] https://orcid.org/0000-0002-6486-8955

Kelle C. Freel[1] https://orcid.org/0000-0001-8350-5327

Keli‘iahonui Kotubetey[4]

A. Hi‘ilei Kawelo[4]

Kawika B. Winter[1,3] https://orcid.org/0000-0003-3762-7125

Michael S. Rappé[1] https://orcid.org/0000-0002-9829-251X

[1]Hawai‘i Institute of Marine Biology, University of Hawai‘i at Mānoa, Kāne‘ohe, Hawai‘i, USA,

96744

[2]Marine Biology Graduate Program, University of Hawai‘i at Mānoa, Honolulu, Hawai‘i, USA,

96822

[3]He‘eia National Estuarine Research Reserve, Kāne‘ohe, Hawai‘i, USA, 96744

[4]Paepae o He‘eia, Kāne‘ohe Hawai‘i, USA, 96744

[#]Josephine Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological

Laboratory, Woods Hole, MA 02543, USA

*Corresponding Author: Email: rappe@hawaii.edu

Running Title: Spatiotemporal shifts in phytoplankton

Abstract

Islands in the tropical Pacific supply elevated nutrients to nearshore waters that enhance

phytoplankton biomass and create hotspots of productivity in otherwise nutrient-poor oceans.

Despite the importance of these hotspots in supporting nearshore food webs, the fine-scale

spatial and temporal variability of phytoplankton enhancement and changes in the underlying

phytoplankton communities across nearshore to open ocean systems remain poorly understood.

In this study, a combination of flow cytometry, pigment analyses, 16S rRNA gene amplicons,

and metagenomic sequencing provide a synoptic view of phytoplankton dynamics over a four-

year, near-monthly time-series across coastal Kāneʻohe Bay, Hawaiʻi, spanning from an

estuarine Indigenous aquaculture system to the adjacent offshore environment. Through

comparisons with measurements taken at Station ALOHA located in the oligotrophic North

Pacific Subtropical Gyre, we elucidated a sharp and persistent transition between

picocyanobacterial communities, from *Synechococcus* abundant in the nearshore to

Prochlorococcus proliferating in offshore and open ocean waters. In comparison to immediately

adjacent offshore waters and the surrounding open ocean, phytoplankton biomass within

Kānʻeohe Bay was dramatically elevated. While phytoplankton community composition revealed

strong seasonal patterns, phytoplankton biomass positively correlated with wind speeds, rainfall,

and wind direction, and not water temperatures. These findings reveal sharp transitions in ocean

biogeochemistry and phytoplankton dynamics across estuarine to open ocean waters in the

tropical Pacific and provide a foundation for quantifying deviations from baseline conditions due

to ongoing climate change.

Introduction

In marine ecosystems, phytoplankton play a crucial role by forming the base of the aquatic food web, where their productivity, abundance, cell size, and community composition are greatly influenced by light and nutrient availability (Azam et al. 1983). Surrounded by oligotrophic, open ocean waters, the coastal waters of remote islands in the tropical Pacific harbor a sharp increase in nutrients through physical oceanographic, biological, geological, and anthropogenic processes that results in increased phytoplankton biomass, cell size, and productivity (i.e. the Island Mass Effect, or IME; Doty and Oguri 1956; Gove et al. 2016). The enhanced primary productivity in turn promotes secondary productivity, supporting regional fisheries (Stock et al. 2017), increased biodiversity (Messié et al. 2022), and other marine resources relied upon by island communities.

Given the importance of elevated phytoplankton biomass from near-island coastal waters to maintaining healthy and productive coastal food webs, understanding the fine-scale variability of phytoplankton communities across coastal to open ocean systems adjacent to island masses can inform both the management of local marine environments and larger ecosystem models. Currently, phytoplankton biomass and productivity in the open oceans are rapidly changing. One result of increasing sea surface temperatures due to ongoing global climate change is an increase in the intensity of water column stratification of the open ocean, which can trap nutrients at depths below where phytoplankton at the ocean's surface can access them (Li et al. 2020). This has led to an expansion of nutrient-poor "ocean deserts" in the open ocean gyres and a decline in global phytoplankton biomass and primary productivity (Kwiatkowski et al. 2018). Declines in primary productivity will likely be amplified across the trophic food web in the near future, with an expected 14% decline in zooplankton biomass as soon as 2100 (Kwiatkowski et al. 2018) and a 20% decline in global fisheries by 2300 (Moore et al. 2018).

Small phytoplankton such as the two most abundant phytoplankton globally,

Prochlorococcus and *Synechococcus*, are expected to increase in abundance under ocean

warming conditions and decreased nutrient availability at the expense of larger sized

phytoplankton groups (Flombaum et al. 2020). Small phytoplankton are often too small to be

effectively grazed by metazoans, and are consumed by intermediary microzooplankton grazers

who are then fed upon by larger zooplankton (Calbet and Landry 1999). Systems dominated by

small phytoplankton have longer food chains and potentially a reduced energy transfer efficiency

to higher trophic levels (Eddy et al. 2021). In contrast, large phytoplankton like diatoms can be

consumed directly by zooplankton grazers (Calbet and Landry 1999) such as copepods, so that

forage fish are only one trophic level apart from phytoplankton. Thus, changes in phytoplankton

size structure and community composition also have profound implications for food web

dynamics.

Unfortunately, the effects of increased open ocean stratification on food webs and

biological productivity in adjacent coastal environments is uncertain. In part, this is due to

studies predominantly focusing on coastal or oceanic systems in isolation (Xenopoulos et al.

2017), but also because the satellite-based methods that have led to an extensive understanding

of primary productivity in the global open ocean have not yet been developed for shallow coastal

waters (Carswell et al. 2017). Importantly, coastal marine food webs of islands situated in

oligotrophic waters may be particularly vulnerable to the impacts of open ocean stratification.

Defining how nutrient availability and phytoplankton community composition and

biomass vary with space and time in near-island and adjacent open ocean environments can

provide a foundation for quantifying deviations from baseline conditions and predicting food

web shifts because of climate change. To illuminate the factors influencing phytoplankton

communities across near-island to open ocean environments in the tropical Pacific, this study

examined the effect of spatial and temporal variability in biogeochemical conditions on

phytoplankton communities across multiple habitats that link the coastal environment of Oʻahu,

Hawaiʻi, with the offshore. These habitats span a tidally-influenced, estuarine environment

within an Indigenous aquaculture system, through the interior of coastal Kāneʻohe Bay, and to

the offshore ocean environment surrounding Kāneʻohe Bay. We also made comparisons to data

collected by the Hawaii Ocean Time-series (HOT), a 30+ year time-series initiative measuring

temporal trends of the adjacent ultraoligotrophic North Pacific Subtropical Gyre (NPSG; Karl

and Church 2014). Together, this extensive spatial and temporal coverage revealed dramatic

nearshore enhancement of phytoplankton biomass, pronounced seasonality in nearshore

biogeochemistry and phytoplankton biomass and composition, and distinct transitions in

phytoplankton communities spanning <6 km to >100 km across Kāneʻohe Bay to the NPSG.

Materials and Methods

Study location

The Hawaiian archipelago within the oligotrophic NPSG is the world's most remote

island chain. Kāneʻohe Bay, located on the windward side of the island of Oʻahu (21° 28′ N,

157° 48′ W), is a well-studied, coral-reef-dominated embayment (**Fig. 1a**). The bay has a total

surface area of 41.4 km^2 and is approximately 4.3 km wide, 12.8 km in length, and 10 m deep on

average (Jokiel 1991). Sharp nearshore to offshore gradients in biogeochemical parameters occur

over a short distance (<6 km) along with a diverse topography due to patch, fringing, and barrier

reefs (Jokiel 1991; Tucker et al. 2021). Localized freshwater input from streams contribute to

episodic spatial variability in environmental conditions, including salinity and inorganic nutrient

concentrations (Cox et al. 2006; Yeo et al. 2013; Tucker et al. 2021). Water residence time

within the bay varies from less than a day to over one month (Lowe et al. 2009), with the highest

residence times in the sheltered southern lobe. Oceanic water, primarily driven by wave action,

flows into the bay over a large barrier reef located in the central bay (Lowe et al. 2009). Water is

generally transported out of the bay through two nearly-parallel channels positioned in the

southern and northern portions of the bay. For most of the year, the bay is well mixed by

tradewinds. However, periods of high temperatures and low wind speeds can cause vertical

stratification in the water column (Smith 1981).

Collaboratively developed research

At the mouth of Heʻeia Stream in the southern section of Kāneʻohe Bay is an ~800-year-

old, 0.356 km^2 Indigenous aquaculture system known contemporarily as Heʻeia Fishpond, but

anciently as Pihi Loko Iʻa (Kelly 1973). Indigenous aquaculture systems in Hawaiʻi engaged in

trophic engineering to promote primary productivity that sustained the population through

abundant food fish and reef fish (Winter et al. 2020a). A 2.5 km basalt rock wall filled with coral

rubble encompasses Heʻeia Fishpond. The wall is equipped with multiple sluice gates that

increase water residence time while still allowing for exchange between coastal Kāneʻohe Bay

and Heʻeia stream waters. The sluice gate (mākāhā) system allows juvenile fish to flourish

within the high nutrient environment of the fishpond, while protecting them from predators.

While it has been understood that the fishpond provided the perfect nursery habitat for prized

food fish, little is known about the exchange of organisms from within the fishpond to the rest of

Kāneʻohe Bay, and beyond. With this interest expressed by the Indigenous stewards of Heʻeia

Fishpond, we sought to establish a current, baseline understanding of phytoplankton biomass and

community composition from within the estuarine fishpond environment out to open ocean

waters adjacent to Oʻahu. The methods employed in this study were collaboratively developed

with Heʻeia Fishpond stewards and the Heʻeia National Estuarine Research Reserve (NERR;

Winter et al. 2020b). All individuals who collaboratively developed the methods also

collaborated on interpreting the data and are listed as co-authors on this paper. Sampling

campaigns were conducted with permission from Paepae o Heʻeia, the stewards of Heʻeia

Fishpond, and the private landowner, Kamehameha Schools.

Sample collection and environmental parameters
142
Between August 2017 and June 2021, seawater was collected from a depth of 2 m at 10

sites in Kāneʻohe Bay and the adjacent offshore waters on a near-monthly basis (36 sampling

events over 46 months) as part of the Kāneʻohe Bay Time-series (KByT) using previously

described methods (**Fig. 1a, Supporting Information Table S1**; Tucker et al. 2021). Between

September 2020 and June 2021, two additional stations within Heʻeia Fishpond were also

sampled at a quarterly interval **(Fig. 1a)**: Station Wai2 (now known as Waimāʻama), located in

the northwestern corner of the Fishpond within the Heʻeia Stream mouth (known as muliwai) is a

highly turbid and brackish water environment with tidal fluctuations resulting in salinity ranges

of 10-35 ppt. Station Kahoʻokele (Kaho) is located at a sluice gate facing the ocean and receives

high exchange with coastal Kāneʻohe Bay (~30-35 ppt, Möhlenkamp et al. 2019). At all stations,

seawater samples for biogeochemical analyses and nucleic acids were collected, as were *in situ*

measurements of seawater temperature, pH, and salinity with a YSI 6600 or ProDSS multi-

parameter sonde (YSI Incorporated, Yellow Springs, OH, USA). Approximately one liter of

seawater was prefiltered with 85-μm Nitex mesh and subsequently filtered through a 25-mm

diameter, 0.1-μm pore-sized polyethersulfone (PES) filter membrane (Supor-100, Pall Gelman

Inc., Ann Arbor, MI, USA) to collect microbial cells for DNA isolation. The filters were

subsequently submerged in DNA lysis buffer (Suzuki et al. 2001; Yeo et al. 2013) and stored in

−80°C until further processing.

Fig. 1. a) Map of sampling stations located on the windward side of the island of Oʻahu Hawaiʻi.

Inset shows two stations within Heʻeia Fishpond and one immediately adjacent to the fishpond.

Contour lines mark every ten meters up until 50 m and then 50 m intervals for depths >50 m. b)

Location of Kāneʻohe Bay on Oʻahu, Hawaiʻi and the position of Station ALOHA, a sampling

station of the Hawaii Ocean Time-series (HOT) program. c) Chlorophyll *a* and d) silicate

concentrations (both plotted on a log scale) from the estuarine to open ocean environments

examined in this study. e) Ratios of dissolved inorganic nitrogen (DIN):Phosphate. Diagonal line

denotes Redfield ratio of 16:1 suggesting nitrogen limitation is characteristic of the system.

Seawater subsamples for fluorometric chlorophyll *a* concentrations (125 mL) and

photosynthetic pigments via high-performance liquid chromatography (HPLC; 2 L) were

collected on 25-mm diameter GF/F glass microfiber filters (Whatman, GE Healthcare Life

Sciences, Chicago, IL, USA) and stored in aluminum foil at −80°C until extraction. The

collection of phytoplankton pigments on the GF/F glass microfiber filters allow for comparisons

with the Hawaii Ocean Time-series data. However, because the filters have a pore size of 0.7μm,

we acknowledge that most small cyanobacteria were likely missed. Chlorophyll *a* was extracted

with 100% acetone and measured with a Turner 10-AU fluorometer (Turner Designs, Sunnyvale,

CA, USA) following standard techniques (Welschmeyer 1994). Photosynthetic pigments

measured via high performance liquid chromatography were extracted in 100% acetone and

analyzed on a Waters 2690 separations module equipped with a C18 column and full spectrum

photodiode array detector, following (Mantoura and Llewellyn 1983) and modified according to

(Bidigare et al. 1989). Chlorophyll *a* concentrations measured via the fluorometer are herein

referred to chlorophyll *a* (Chla), while chlorophyll *a* concentrations measured by high

performance liquid chromatography are specified as total chlorophyll *a* (TChla).

For cellular enumeration, seawater was preserved in 2 mL aliquots in a final

concentration of 0.95% (v:v) paraformaldehyde (Electron Microscopy Services, Hatfield, PA,

USA) at −80 °C until analyzed via flow cytometry. Cellular enumeration of cyanobacterial

picophytoplankton (*Synechococcus* and *Prochlorococcus*), eukaryotic picophytoplankton, and

non-cyanobacterial (presumably heterotrophic) bacteria and archaea (hereafter referred to as

heterotrophic bacteria) was performed on a Beckman Coulter CytoFLEX S, following the

method of (Monger and Landry 1993). Inorganic nutrients were measured using a Seal

Analytical AA3 HR Nutrient Autoanalyzer (detection limits: $NO_2^- + NO_3^-$, 0.009 µM; SiO_4,

0.09 µM; PO_4^{3-}, 0.009 µM; NH_4, 0.03 µM).

Rainfall, wind speed, and wind direction were monitored using data collected at a

meteorological station located at the Hawaiʻi Institute of Marine Biology (HIMB) on Moku o

Loʻe in Kāneʻohe Bay (http://www.pacioos.hawaii.edu/weather/obs-mokuoloe/). Either

maximum (e.g., rainfall, wind speed) or average (wind direction) values were taken across 1- to

7-day windows leading up to the sampling event, depending on data availability from the station.

One sampling event (February 5, 2021) had no data for the 7 days prior to sampling and so the

data for a 30-day window were used. Metadata from Station ALOHA (22° 45'N, 158° 00'W, ~5

m depth, August 2017 to December 2020), a sampling station of the HOT program (Karl and

Church 2014), were downloaded from https://hahana.soest.hawaii.edu/hot/hot-dogs/ (accessed on

9/12/2022).

Spatiotemporal comparisons of environmental variables, cellular abundances, and

phytoplankton pigments were conducted using the R package 'multcomp' (Hothorn et al. 2008)

with one-way ANOVAs testing for multiple comparisons of means with Holm correction and

Tukey contrasts. Summer (28 June through 28 September) and winter (27 December through 29

March) seasons were defined using harmonic regression analyses of surface seawater

temperature collected hourly between 2010–2019 at NOAA station MOKH1 in Kāneʻohe Bay

(https://www.ndbc.noaa.gov/station_page.php?station=mokh1; Tucker et al. 2021).

Spatiotemporal variation in cellular abundances and phytoplankton pigments were visualized

using 'mba.surf' from MBA (Finley et al. 2017) to interpolate data over the KByT sampling

events and stations.

The map of Kāneʻohe Bay was plotted in ArcGIS Pro v2.9. Contour lines were drawn

using bathymetry metadata (http://www.soest.hawaii.edu/hmrg/multibeam/bathymetry.php). The

map of Oʻahu was plotted in R with 'geom_sf' from ggplot2 (Wickham 2016), using shape file

from the Hawaii Statewide GIS Program (https://prod-

histategis.opendata.arcgis.com/maps/HiStateGIS::coastline).

DNA extraction, 16S rRNA gene amplicon sequencing, & metagenome sequencing

DNA extraction and 16S rRNA gene sequencing followed previously published methods

(Tucker et al. 2021). Briefly, amplicon libraries were made from polymerase chain reactions of

the 16S rRNA gene using barcoded 515F and 926R universal primers (Parada et al. 2016) and

paired-end sequenced with MiSeq v2 2x250 technology (Illumina, San Diego, CA, USA).

Genomic DNA from a subset of 32 of the 368 total samples collected between 2017-2021 were

used for metagenomic sequencing. This included samples from four sampling events between

2017 and 2019 at 6-10 stations. Libraries were constructed from approximately 100 ng of

genomic DNA using the Kappa HyperPrep Kit (Roche, Pleasanton, CA, USA) with mechanical

shearing (Covaris, Woburn, MA, USA) and paired-end sequenced on a single lane of the

NovaSeq 6000 SP 150 (Illumina, San Diego, CA, USA).

Sequence analysis

Amplicon sequence data generated from KByT sampling between July 2019 and June

2021 were analyzed in conjunction with previously published amplicon data spanning August

2017 to June 2019 (PRJNA706753; (Tucker et al. 2021). For each of the two sequencing runs,

samples were demultiplexed and quality controlled using Qiime2 v2.4 (Bolyen et al. 2019). Full

length forward reads (251 base pairs) were denoised using DADA2 (Callahan et al. 2019) to

delineate amplicon sequencing variants (ASVs). Reverse reads were not used because of

inconsistent quality. ASVs were assigned taxonomy using SILVA v138 as a reference database

(Quast et al. 2012) and the two runs were subsequently merged in Qiime2. ASVs that contained

at least 10 reads in at least two samples were retained.

ASVs classified by the SILVA v138 database as Eukaryota, unassigned at the domain

level, or classified as chloroplast at the order level were re-classified using the PR2 v 4.14.0

database (Guillou et al. 2013) in the DECIPHER R package (Wright 2016) using a 60%

confidence threshold cut off. Sequences classified as Bacteria, Archaea, or chloroplast at the

order-level were retained for further analyses, while those unclassified at the domain level or

classified as Eukaryota were excluded from further analyses. In the context of amplicon

sequence data, "phytoplankton" herein refers to ASVs classified as cyanobacteria and eukaryotic

plastid sequences, although we recognize that mixotrophic and phagotrophic lifestyles may be

included in this broad definition.

Statistical analyses were conducted using the R packages phyloseq (McMurdie and

Holmes 2013), ggplot2 (Wickham 2016), pheatmap (Kolde 2019), and microbiome (Lahti and

Shetty 2017). An Aitchison distance (Aitchison 1982), the Euclidean distance between centered

log-ratio (clr)-transformed compositions, was used on the entire quality-controlled dataset of

phytoplankton raw read counts using 'transform' in the microbiome package (Lahti and Shetty

2017). Ward D2 hierarchical clustering using 'hclust ( )' in the stats base package of R was

applied to this matrix to cluster samples with amplicon data into groups and visualized with

dendextend (Galili 2015). DESeq2 (Love et al. 2014) was used to model differential abundance

patterns of amplicon data across environmental clusters using Wald Tests and Bonferroni

correction for multiple comparisons (alpha cutoff < 0.05). Divnet (Willis and Martin 2020) was

used to estimate differences in alpha diversity and test for significance between spatiotemporal

groupings. Pearson's correlation analyses were conducted in corrplot (Wei and Simko 2021).

Lomb Scargle Periodograms (LSP) in the lomb package (Ruf 1999) were used to define

seasonality among phytoplankton genera by determining the spectrum of frequencies in a

dataset: this approach can account for unevenly sampled time-series data and has been

previously applied to microbiome time-series analyses (Auladell et al. 2021). Only genera with

annual intervals (peak frequency = 1 ± 0.25, $p<0.01$) as their most significant periodic trend were

considered as having seasonality. A starting frequency of 0.16 was used so as to not include

periodic components between two consecutive months. An inverse hyperbolic sine

transformation (asinh) was conducted on sequence data prior to LSP. Significance (q-

values<0.05) was corrected for multiple-testing using data randomization for LSP analyses using

fdrtools (Strimmer 2008).

Metagenomic read recruitment

To investigate the dominant cyanobacteria within and surrounding Kāneʻohe Bay, we

conducted metagenomic read recruitment of 32 metagenomes from KByT and 12 previously

published from the open ocean Station ALOHA (PRJNA352737; Mende et al. 2017) to 56

cyanobacterial genomes from *Prochlorococcus* (six minor clades) and the three major lineages of

the marine *Synecchococcus*/*Cyanobium* lineage [SC 5.1 (14 minor clades), SC 5.2, and SC 5.3]

(**Supporting Information Table S2**). SC 5.2 is the only clade with both *Synechococcus* and

Cyanobium members (Doré et al. 2020).

A contig database of the 56 cyanobacteria isolate genomes was constructed using anvi'o

v 8.0 (Eren et al. 2021) following previously described pipelines (Delmont and Eren 2018).

Briefly, Prodigal v2.6.3 (Hyatt et al. 2010) was used to identify open reading frames (ORFs)

from the contigs and an anvi'o database was created using 'anvi-gen-contigs-db'. Metagenomic

reads were first quality filtered using an Illumina-utils library v1.4.1 called 'iu-filterquality-

minoche' (Eren et al. 2013) that uses quality filtering parameters described previously (Minoche

et al. 2011). Quality filtered metagenomic reads were competitively mapped with Bowtie2 v2.3.5

(Langmead and Salzberg 2012) to an anvi'o contig database of cyanobacterial isolate genomes.

The 'anvi-profile' function stored coverage and detection statistics of each cyanobacterial

genomes found in the KByT and Station ALOHA metagenomic samples.

To evaluate the distribution of individual genomes, a "detection" metric, the proportion

of the nucleotides in a given sequence that are covered by at least one short read, was used to

evaluate if a population was present in a metagenomic sample. A detection value of at least 0.25

was used as a criterion to eliminate false positives, when an isolate genome was falsely found

within a sample (Utter et al. 2020). Mean coverage Q2Q3, which refers to the average depth of

coverage excluding nucleotide positions with coverages in the 1st and 4th quartiles, was mapped

for each genome. Mean coverage Q2Q3 was summed across all cyanobacterial genomes per

sample and then the genome (or all genomes in a clade) was divided by this sum to determine a

relative abundance of a genome (or a clade) in each sample. Average nucleotide identity (ANI)

was calculated using pyANI (Pritchard et al. 2015). A phylogenomic tree was estimated from the

56 cyanobacterial isolates and an outgroup using GTotree v1.4.16 (Lee 2019) with

cyanobacterial single-copy genes and visualized in FigTree v 1.4.4 (http://tree.bio.ed.ac.uk/).

Data & code availability

Sequencing data are available in the National Center for Biotechnology Information (NCBI)

Sequence Read Archive (SRA) under BioProject number PRJNA706753 as well as

PRJNA971314. Environmental data were submitted to BCO-DMO under https://www.bco-

dmo.org/project/663665. Code used in the analysis is available at

https://github.com/tucker4/Tucker_Phytoplankton_KByT_HeNERR.

Results

Biogeochemical parameters

Along the nearshore to open ocean waters of the tropical Pacific, biogeochemical

parameters sharply declined across both small spatial scales and vast stretches of ocean

(**Supporting Information Table S1**). On average, chlorophyll *a* concentrations at stations in the

coastal waters of Kāneʻohe Bay increased 18-fold (1.6±1.9 *vs.* 0.09±0.03 µg L^{-1}, mean±sd) from

the open ocean and 3-fold (1.6±1.9 *vs.* 0.6±0.4 µg L^{-1}) from the immediately adjacent offshore

waters. The estuarine waters of Heʻeia Fishpond harbored higher chlorophyll *a* concentrations

compared to coastal stations (3.9±3.8 *vs.* 1.6±1.9 µg L^{-1}, **Fig. 1c**). Mean chlorophyll *a*

concentrations increased 43-fold between the estuarine waters of Heʻeia Fishpond and the open

ocean (3.9±3.8 *vs.* 0.09±0.03 µg L^{-1}, **Supporting Information Table S1**) and 7-fold over the <6

km distance covering the interior and surrounding waters of Kāneʻohe Bay (3.9±3.8 *vs.* 0.6±0.4

µg L^{-1}; **Fig. 1c**, **Supporting Information Table S1**). Stations offshore from Kāneʻohe Bay had

elevated concentrations of chlorophyll *a* compared to the open ocean (7-fold increase; 0.6±0.4

vs. 0.09±0.03 µg L^{-1}). Elevated phytoplankton biomass was a persistent feature within Kāneʻohe

Bay, with increased chlorophyll *a* concentrations detected in at least one of the stations

positioned in the coastal environment compared to the stations offshore during all 36 sampling

events (**Supporting Information Fig. S1**).

Elevated concentrations of inorganic nutrients (silicate, nitrate+nitrite, phosphate,

ammonia) were also found in the nearshore waters of Kāne'ohe Bay compared to offshore and

open ocean stations (**Supporting Information Table S1 & S3**). Mean silicate concentrations at

Wai2 of the estuarine stations were 107.3 µM, compared to 42.2 µM in the coastal stations.

Across all KByT stations, phosphate and silicate concentrations differed significantly but were

positively correlated with increasing chlorophyll *a* concentrations. However, nitrate+nitrite

concentrations did not correlate with chlorophyll *a* concentrations (**Supporting Information**

Fig. S2). Despite the overall increase of inorganic nutrients in the estuarine and coastal stations,

all stations were below 16:1 N:P ratios using dissolved inorganic nitrogen (nitrate+nitrite plus

ammonia) and phosphate (**Fig. 1e**, (Redfield 1960).

Microbial cell counts and phytoplankton pigments

In contrast to coastal stations where *Synechococcus* cellular abundance was high,

Prochlorococcus cellular abundance was elevated in the stations positioned in the offshore

waters surrounding Kāne'ohe Bay and at Station ALOHA (**Fig. 2a, Supporting Information**

Table S1 & S3). Cellular abundances of heterotrophic bacteria and eukaryotic

picophytoplankton were also greater in the coastal stations compared to offshore (**Fig. 2a,**

Supporting Information Table S1 & S3). In coastal Kāne'ohe Bay, ratios of fucoxanthin,

peridinin, and alloxanthin to total chlorophyll *a* (Tchla) concentrations were higher than in the

offshore, indicating an increase in diatoms, dinoflagellates, and cryptophytes (respectively)

closer to shore (**Fig. 2b, Supporting Information Table S4**). In contrast, pigments relative to

Tchla for photosynthetic pigments diagnostic of prymnesiophytes (19′-hexanoyloxyfucoxanthin),

pelagophytes (19′-butanoyloxyfucoxanthin), cyanobacteria (zeaxanthin), and *Prochlorococcus*

(divinyl chlorophyll *a*) were higher in the offshore stations compared to the coastal environment

(**Fig. 2b**, **Supporting Information Table S4**).

Fig. 2. Microbial cellular abundances and pigment concentration vary through time and space

across stations from the southern sector of Kāneʻohe Bay (HP1, SB, SR8), offshore stations (SR2

and STO1), and open ocean Station ALOHA: a) Cellular abundances (cells mL^{-1}) of

heterotrophic bacteria, *Synechococcus*, photosynthetic picoueukaryotes, and *Prochlorococcus*; b)

Total chlorophyll *a* (Tchla) concentrations (µg L^{-1}) and ratios of phytoplankton pigments

indicative of specific phytoplankton groups relative to Tchla. Note, alloxanthin was below

detection levels for Station ALOHA and not presented here. Abbreviations: Fuco: Fucoxanthin,

Peri: Peridinin; Allo: Alloxanthin; Hex: 19′-hexanoyloxyfucoxanthin; But: 19′-

butanoyloxyfucoxanthin; Zea: zeaxanthin; Dvchla: divinyl chlorophyll *a*.

Phytoplankton community composition through 16S rRNA gene sequencing

We delineated 505 phytoplankton ASVs across 366 samples, including 66 from

cyanobacteria and 439 from eukaryotic plastids. Examining the distribution of phytoplankton

ASVs across samples revealed that phytoplankton communities clustered into three major

community types that coincide with spatial differences in biogeochemistry (**Fig. 3**, **Supporting

Information Table S5**). We categorized the three major community types as nearshore,

transition, and offshore because of their distinct biogeochemical characteristics and geographic

location (**Supporting Information Table S5, Fig. S3**). The nearshore cluster of 229 samples

included all samples collected from six stations found most closely located to land (Wai2,

Kahoʻokele, SB, CB, HP1, AR) and at least one sample from the six remaining stations. The

transition cluster of 85 samples consisted of samples collected from stations not immediately

next to land (NB, SR8, NR2, SR2, STO1, NTO1), while the offshore cluster encompassed 52

samples collected exclusively from the four stations located the furthest distance from land (SR2,

NR2, STO1, NTO1).

Figure 3. a) Hierarchical clustering of phytoplankton communities from 366 samples collected

at 12 sampling stations form three major groups, hereafter referred to as community types:

nearshore, transition, and offshore. b) The distribution of samples defined as nearshore,

transition, and offshore community types across stations and sampling events (36 total between

2017-2021) shows spatial and temporal persistence in the distinct community types. The size of

the circle represents the chlorophyll *a* concentration during the time of sampling and the color of

the circle represents the community type: nearshore, transition, and offshore.

ASV richness was the highest in the nearshore, while the transition community group had

the highest Shannon's diversity estimate (**Supporting Information Table S6**). Phytoplankton

relative abundance was dominated by a few highly abundant groups, including *Synechococcus*,

Prochlorococcus, Mamiellophyceae (green algae), and Bacillariophyta (diatoms) (**Fig. 4a**).

Using DESeq2 variance stabilized abundances, 20 classes (**Supporting Information Table S7**)

and 33 genera of phytoplankton (or groups unclassified at genus-level but classified at the

family-level; **Fig. 4b**, **Supporting Information Table S8**) differed significantly in abundance

across the three community types.

Fig. 4. a) Mean relative abundance of the top 10 phytoplankton groups across the nearshore (N),

transition (T), and offshore (O) environments. b) Phytoplankton genera with significantly

different distributions across the three environments. Colored boxes denote the peak in

abundance for each genus. Phytoplankton groups that were classified at the family-level but

unidentified at the genus-level are denoted with an asterisk.

Cyanobacterial population structure

Metagenomic read recruitment to 56 genomes of the cyanobacterial genera

Prochlorococcus, *Synechococcus*, and *Cyanobium* showed that *Prochlorococcus* HLI, HLII, and

LLI, and *Synechococcus* SC 5.1 II, WPC2, III, UC-A, and IX and SC 5.3 were detected in

metagenomes from the Kāneʻohe Bay Time Series and surface ocean samples from Station

ALOHA (**Fig. 5a**). Although *Cyanobium* 16S rRNA gene ASVs were detected in the amplicon

data, no *Cyanobium* representatives (SC 5.2) were detected in our metagenomic read

recruitment. Genomes from *Synechococcus* SC 5.1 II and SC 5.3 and *Prochlorococcus* HLII

were among the most abundant representatives within our samples (**Fig. 5b**). *Prochlorococcus*

HLII comprised 98.9±1.0% (mean±sd) of cyanobacterial relative abundance in the open ocean

and 83.1±17.4% of the cyanobacterial relative abundance in the offshore. *Prochlorococcus* HLII

recruited only a small proportion of reads from a handful of nearshore samples, where it made up

1.1±2.4% of the cyanobacterial relative abundance. *Synechococcus* clade II comprised

96.3±2.5% of the cyanobacterial relative abundance in the nearshore Kāneʻohe Bay community

type. *Synechococcus* SC 5.3 also recruited some metagenomic reads, but only from coastal

Kāneʻohe Bay samples and at low relative cyanobacterial abundance (2.4±1.2%) (**Fig. 5b**).

Within the *Prochlorococcus* HLII and *Synechococcus* II clades, read recruitment varied

between closely related genomes. Read recruitment was substantially higher in *Synechococcus*

clade II isolate UW86 compared to all other clade II genomes, despite sharing >95% ANI with

most other clade II genomes (**Supporting Information Fig. S4**). Within *Prochlorococcus* HLII,

isolate genomes AS9601, SB, MIT9314, and to a lesser extent MIT9301, recruited a

substantially greater proportion of reads than other members of this clade. AS9601, SB,

MIT9314, and MIT9301 shared 94% ANI, which is higher than what was shared with other HLII

isolates (<93%), with the exclusion of MIT9215 and MIT9202 who share 97% ANI with each

other (**Supporting Information Fig. S4**).

a)

b)

Fig. 5. a) Phylogenomic tree based on cyanobacterial marker genes found in 56

Prochlorococcus, *Synechococcus*, and *Cyanobium* isolate genomes and outgroup (*Gloeobacter*

violaceus- not shown). Clades detected in surface ocean metagenomic samples from the

Kāneʻohe Bay Time-series (KByT) and Station ALOHA are colored and bolded. b) The relative

abundance of cyanobacterial genomes across KByT and Station ALOHA metagenomes was

dominated by *Prochlorococcus* HLII, *Synechococcus* II (SC 5.1), and *Synechococcus* SC 5.3.

Recruitment of <0.5% not shown.

Seasonality in biogeochemistry and community composition

Surface seawater temperatures were significantly cooler in the winter than summer in

nearshore Kāneʻohe Bay and the adjacent transition and offshore waters (**Fig. 6a, Supporting**

Information Table S9). Cellular concentrations of *Synechococcus* and heterotrophic bacteria

were higher in the summer than the winter in the nearshore, transition, and offshore (**Fig.6a,**

Supporting Information Table S9). Photosynthetic picoeukaryote cellular concentrations

increased during the summer in coastal and transition environments (coastal: p=0.004, transition:

p<0.001, **Fig. 6a, Supporting Information Table S9**), but did not vary seasonally in the

offshore. *Prochlorococcus* cell concentrations only varied seasonally in the nearshore where it

increased in the winter (p<0.001, **Fig. 6a, Supporting Information Table S9**).

Chlorophyll *a* concentrations increased in the winter compared to the summer in the

nearshore (mean±sd; winter: 2.3±3.4 µg L^{-1}, summer: 0.8±0.5 µg L^{-1}; p=0.031; **Fig. 6a,**

Supporting Information Table S9), but not in the transition or offshore waters. Importantly,

given that the collection method for chlorophyll *a* concentrations likely missed most small

cyanobacteria, it is possible that seasonality has been underestimated. Nearshore chlorophyll *a*

concentrations increased with wind speed, wind direction, and rainfall, but not with seawater

temperature (**Fig. 6b**). Three sampling events with elevated chlorophyll *a* concentrations in the

nearshore occurred during anomalously high wind speeds or rainfall events (**Supporting**

Information Fig. S5). Chlorophyll *a* concentrations in the transition and offshore did not

correlate with seawater temperatures, but did positively correlate with rainfall (**Supporting**

Information Fig. S6). Phosphate concentrations increased in the summer compared to the winter

in the nearshore and transition clusters, while silicate concentrations increased in the summer

compared to the winter in the nearshore cluster (**Supporting Information Table S9**).

Ratios of phytoplankton pigments to Tchla representing cyanobacteria, diatoms, and

pelagophytes showed correlations with seawater temperature in the nearshore community type,

suggestive of seasonality in the abundance of major phytoplankton groups (**Supporting**

Information Fig. S7). Correlations between seawater temperature and pigment to Tchla ratios

were not detected in the transition and offshore cluster (**Supporting Information Fig. S7**).

Fig. 6. a) Changes in seawater temperature, cell counts of *Synechococcus*, photosynthetic

picoeukaryotes, *Prochlorococcus*, and chlorophyll *a* across seasons and environments. b)

Correlations between chlorophyll *a* concentrations and other environmental parameters in the

nearshore. c) Significant correlations between phytoplankton pigment:tchla ratios and seawater

temperature in the nearshore. An asterisk (*) denotes variables with log transformations, while a

carrot (^) denotes variables with log+1 transformations. Het.Bac: heterotrophic bacteria (cells

mL^{-1}); Syn: *Synechococcus* (cells mL^{-1}); Pro: *Prochlorococcus* (cells mL^{-1}); Picoeuk:

Photosynthetic picoeukaryotes (cells mL^{-1}); Temp: Seawater temperature (℃); Chla: Chlorophyll

a (µg L^{-1}); Salinity (ppt); Wind direction (degrees); Wind Speed (ms^{-1}); Rainfall (mm); Zea:

Zeaxanthin; Fuco: Fucoxanthin; But: 19′-butanoyloxyfucoxanthin.

Seasonal differences were found in phytoplankton alpha diversity and the relative

abundance of phytoplankton genera across the nearshore, transition, and offshore. ASV richness

and Shannon diversity increased in the winter compared to the summer in all three community

types (**Fig. 7a**, **Supporting Information Table S10**). Seasonality was observed among 23

phytoplankton genera (or groups unclassified at the genus-level, but classified at the family-

level) within each of the clusters, including 18 nearshore, 5 in the transition, and 6 offshore

(**Fig.7b, Supporting Information Table S11**). Seasonal genera accounted for 84.9±7.4,

48.6±18.9, and 23.2±13.1% of the relative abundance of the community on average in the

nearshore, transition, and offshore clusters, respectively. *Synechococcus* was the most abundant

seasonal genus, increasing in relative abundance in the summer months across all community

type clusters (**Fig.7b,c**). *Cyanobium*, *Crocosphaera*, and unidentified Pycnococcaceae also

increased in abundance in the summer months (**Fig. 7b**). Seasonal genera more often increased in

relative abundance during the winter (n=18) including those belonging to diatoms (e.g.

Chaetoceros and unclassified polar-centric Mediophyceae), green algae (e.g. unclassified

Mamiellophyceae, *Micromonas*, and *Mamiella*), prymnesiophytes (e.g. *Isochrysis*,

Chrysochromulina, and *Braarudosphaera*), and Dictyochophyceae (*Helicopedinella*,

Mesopedinella) (**Fig. 7b**). In the nearshore, the dominant phytoplankton in the winter remained

Cyanobacteria followed by Mamiellophyceae and then diatoms, although at times both

Mamiellophyceae and diatoms exceeded >30% of the total phytoplankton relative abundance

(**Fig. 7c**).

a)

Nearshore   Transition   Offshore

Shannon Diversity

p<0.001   p<0.001   p<0.001

ASV Richness

p<0.001   p=0.05   p<0.001

Summer Winter   Summer Winter   Summer Winter

b)

Mean relative abundance (%) of seasonal genera

Nearshore (n=18)   Transiton (n=5)   Offshore (n=6)

Summer Winter   Summer Winter   Summer Winter

Archaeplastida
Mamiellaceae* (N)
Micromonas (N,O)
Chloropicaceae* (N)
Pyramimonas (T)
Mamiella (N)
Prasinoderma (T)
Pycnococcaceae* (N)

Excavata
Eutreptiella (N)

Hacrobia
Teleaulax (N)
Braarudosphaera (N,T,O)
Chrysochromulina (N)
Isochrysis (N)

Rhizaria
Partenskyella (N)

Stramenopiles
Chaetoceros (N)
Polar-centric Mediophyceae* (N)
Helicopedinella (N)
Mesopedinella (N)
Pelagomonadaceae* (O)

Cyanobacteria
Crocosphaera (O)
Cyanobium (N,T)
Prochlorococcus (N)
UCYN-A (O)
Synechococcus (N,T,O)

c)

Cyanobacteria

Archaeplastida (green algae)

Stramenopiles (diatoms)

Fig. 7. a) ASV richness and Shannon's diversity across all three community types differ during

the summer and winter seasons. b) Mean relative abundances of the 29 significantly seasonal

genera across the three community types for summer and winter seasons. c) Nearshore seasonal

phytoplankton genera from Cyanobacteria, Stramenopiles, and Archaeplastida with relative

abundances >0.5% on average. A local polynomial regression fit line is shown for each genus.

Phytoplankton groups that were classified at the family-level but unidentified at the genus-level

are denoted with an asterisk.

Discussion

The activity and distribution of phytoplankton has important implications for food web

dynamics and biocultural restoration and management of near-island waters of the tropical

Pacific, especially under predicted climate change conditions that are expected to limit nutrient

availability, phytoplankton size, productivity, and biomass in open oceans. Ecosystem-level

time-series analyses spanning estuarine waters within Kāneʻohe Bay of the Hawaiian island of

Oʻahu to the North Pacific Subtropical Gyre revealed that surface ocean biogeochemistry,

phytoplankton biomass, and phytoplankton community structure varied dramatically across both

broad (e.g. nearshore Kāneʻohe Bay to the NPSG) and narrow (e.g. nearshore Kāneʻohe Bay to

adjacent offshore waters) spatial scales. Through investigations of these spatiotemporal

dynamics, we gained insight into the ecology and phenology of phytoplankton communities of

surface oceans in the tropical Pacific and identified indicators to help evaluate deviations from

current conditions.

Spatially distinct phytoplankton and biogeochemical regimes within and adjacent to Kāneʻohe

Bay

Our sampling approach provides an increased resolution upon satellite-based studies

(Gove et al. 2016; Messié et al. 2022), which are typically not reliable within ~5 km to shore due

to heavy cloud cover, along with a synoptic view of the biogeochemistry and phytoplankton

communities. Compared to offshore and open ocean waters, nearshore Kāneʻohe Bay had

elevated chlorophyll *a* concentrations- a 3-fold increase in phytoplankton biomass within 6 km

from coastal Kānʻeohe Bay and an 18-fold increase over a roughly 100 km distance. Inorganic

nutrient and chlorophyll *a* concentrations increased with decreasing salinity, suggesting that

freshwater input from streams is an important driver of nutrient delivery and subsequent

phytoplankton enhancement within Heʻeia Fishpond and Kāneʻohe Bay. Previous studies within

Kāneʻohe Bay have also revealed stream-delivered nutrient input as an important driver of

phytoplankton enhancement (Yeo et al. 2013), as well as additional processes including

submarine groundwater discharge (McKenzie et al. 2019) and human-driven pollution (Ringuet

and Mackenzie 2005). Transport from offshore, subsurface nutrient-rich waters to nearshore

waters via internal waves is likely an important process contributing to phytoplankton

enhancement of surface waters in the larger Hawaiian archipelago (Gove et al., 2016), although

its prevalence and impact specifically within Kāneʻohe Bay remains unknown.

Across Kāneʻohe Bay and the adjacent open ocean, phytoplankton communities resolved

into three distinct groups that coincide with spatial differences in biogeochemistry. Pigments

indicative of diatoms and cyanobacteria revealed that these groups were the main phytoplankton

contributing to the enhanced chlorophyll *a* concentrations in the nearshore environment of

Kāneʻohe Bay. Some phytoplankton that were more abundant closer to shore, such as *Teleaulax*

(Cryptophyta), *Pyamimonas* (Chlorophyta), *Tetraselmis* (Chlorophyta), Chlorarachniophytes

(Rhizaria), and dinoflagellates, are associated with mixotrophic lifestyles (Stoecker et al. 2016).

Mixotrophy provides a crucial trophic link in planktonic food webs by supplementing primary

production via heterotrophy, increasing carbon transfer to high trophic levels, and serving as a

source of nutrients (Stoecker et al. 2016). While establishing the relative importance of

mixotrophy across the nearshore to the adjacent offshore environment requires further

investigation, these initial insights show distinctions in food web dynamics, phytoplankton

ecologies, and contributors to primary productivity across the nearshore to open ocean waters of

the tropical Pacific.

Drivers of seasonality in nearshore chlorophyll a *concentrations*

While seasonality within transition and offshore phytoplankton communities was muted,

nearshore phytoplankton biomass was significantly elevated in winter, and members with the

highest relative abundance in the nearshore phytoplankton community showed high seasonality.

Chlorophyll *a* concentrations varied seasonally in nearshore Kāneʻohe Bay, where it increased

with wind speed, rainfall, and wind direction. In Hawaiʻi, storm events generally increase during

the winter months where storm-associated rainfall and wind may serve to elevate inorganic

nutrient concentrations through increased stream outflow, leading to short-term enhancement of

phytoplankton biomass in nearshore Kāneʻohe Bay (Ringuet and Mackenzie 2005; Cox et al.

2006; Yeo et al. 2013). Periods of intense rainfall have strong impacts on the food web dynamics

of Kāneʻohe Bay, with wet periods decreasing trophic complexity and increasing total

community biomass and the transfer of production to metazoans (Selph et al. 2018). Because

phytoplankton growth and loss processes can occur over short time scales of a day to a week, the

near-monthly sampling interval employed in KByT is unlikely to fully capture the rapid

fluctuations of chlorophyll *a* concentrations that occur in response to storm events. Thus, our

observations likely underestimate the episodic variability of phytoplankton biomass in the

nearshore (Yeo et al. 2013).

Distinct growth and nutrient uptake strategies between *Synechococcus* and diatoms, along

with changes in the environment that might favor one over the other, likely produced the

dynamic seasonal patterns observed in nearshore Kāneʻohe Bay: summer months showed a near

doubling of the cellular and relative abundance of *Synechococcus*, while winter months were

marked by sharp increases in the relative abundances of diatoms such as *Chaetoceros* and

unidentified polar-centric Mediophyceae. To understand competition outcomes under different

environmental conditions, resource competition theory characterizes trade-offs between slow-

growing nutrient specialists with high affinity for uptake but low maximum growth (e.g.

Prochlorococcus and *Synechococcus*) and fast-growing nutrient opportunists with high

maximum growth but low affinity for uptake (e.g. flagellates and diatoms) (Dutkiewicz et al.

2009). Due to their small size and high uptake capacity, *Synechococcus* are likely less limited by

the typically low nitrogen concentrations within Kāneʻohe Bay than other phytoplankton (Burson

et al. 2018). In addition, seawater temperatures have been shown to positively correlate with

Synechococcus cellular abundances, likely because seasonal increases in temperature positively

impact *Synechococcus* division rates (Hunter-Cevera et al. 2016). Thus, in the summer under

high-light, warm-water temperatures and limited nutrients, *Synechococcus* dominated Kāneʻohe

Bay. In contrast, despite higher nutrient requirements due to their larger cell sizes, diatoms have

high rates of growth allowing them to outcompete other phytoplankton under periods of elevated

nutrient availability (Laws 1975). Pulses of nutrients from the more frequent storm events during

the winter could have led to the observed fluxes in diatom relative abundances.

Phytoplankton indicators of climate change impacts

Phytoplankton biodiversity (e.g. ASV richness, Shannon's diversity) in Kāneʻohe Bay

was significantly elevated in comparison to the adjacent offshore. Near-island environments can

export diversity offshore and, importantly, biodiversity can offer functional redundancy and

ecological stability in times of environmental perturbation (Messié et al. 2022). In a recent study

that modeled changes to phytoplankton biodiversity under climate change conditions, some

tropical regions were found to face up to 30% of phytoplankton types becoming locally

extirpated (Henson et al. 2021). Importantly, phytoplankton of higher size classes, predominantly

diatoms, are expected to be lost due to increased nutrient limitation (Flombaum et al. 2020).

Diatoms serve as an important part of the diet of herbivorous fish grown in Hawaiian aquaculture

systems (Hiatt 1947), and contribute significantly to primary productivity in nearshore systems

broadly. Diatoms also contribute a large portion of the total chlorophyll *a* in the nearshore

environment of Kāneʻohe Bay. Shifts or reduction in diatom abundance within the Heʻeia

Fishpond and Kāneʻohe Bay when compared the baseline knowledge characterized here might

thus help to identify impacts of ocean warming and stratification in advance of shifts in the food

web structure.

Synechococcus and *Prochlorococcus*, the small cyanobacteria that dominated the total

phytoplankton relative abundance in our study, may also provide valuable bioindicators of

environmental change within Kāneʻohe Bay under climate change scenarios. *Prochlorococcus*

and *Synechococcus* are responsible for roughly 25% of the ocean's net primary production

(Flombaum et al. 2013) and will likely further increase in abundance with projected climate

change conditions (Flombaum et al. 2020). These marine cyanobacteria encompass fine-scale

genetic diversity that distinguishes their ecologies, metabolisms, and biogeochemical roles at the

level of major and minor clades (Berube et al. 2019). These clade identities are often difficult to

resolve with the use of single gene markers, and thus metagenomic read recruitment can increase

genetic resolution.

Across the open ocean NPSG to nearshore Kāneʻohe Bay, *Prochlorococcus* Clade HLII

and *Synechococcus* Clade II (SC 5.1) were the most abundant, consistent with previous reports

from oligotrophic oceans (Delmont and Eren 2018; Lee et al. 2019). *Prochlorococcus* Clade

HLII from the offshore waters adjacent to Kāneʻohe Bay and Station ALOHA showed high

similarity in population structure and thus may represent continuous populations with ongoing

gene flow responding to similar environmental parameters in both environments. Continued

examination of population structure, as well as cellular and relative abundances of these

cyanobacteria, could identify eventual expansions of ultraoligotrophic waters into Kāneʻohe Bay,

shifts in gene-flow, and selection for clades with unique ecological adaptations. Given the high

seasonality of *Synechococcus* cellular and relative abundance in the nearshore environment,

changes in the magnitude and timing of these metrics could also be used to identify alterations in

seasonality and associated food-web dynamics within Kāneʻohe Bay.

Implications for biocultural restoration

The enhancement of phytoplankton biomass in the estuarine Heʻeia Fishpond and coastal

Kāneʻohe Bay provides critical ecosystem services and is an important consideration for

biocultural restoration activities, community-based research efforts, and resource management.

Within Heʻeia and across the islands of Hawaiʻi, biocultural practitioners are undertaking

restoration projects to maximize primary and secondary production of the estuarine environment,

including removing invasive mangrove, managing stream use to ensure adequate flow and water

quality, and engineering water exchange through repairs and updates to fishpond walls

(Möhlenkamp et al. 2019). The baseline understanding of phytoplankton communities provided

by this study helps to inform biocultural stewards of Heʻeia and Kāneʻohe Bay of the conditions

that promote phytoplankton growth and subsequently herbivorous fish growth.

High-resolution sampling was key to identifying the magnitude of chlorophyll *a*

enhancement across the system, areas of high localized phytoplankton biomass enhancement

such as Heʻeia Fishpond, and drivers of chlorophyll *a* enhancement including freshwater input

and increased storm conditions and nutrient concentrations. Despite fine-scale biogeochemical

differences within Kāneʻohe Bay, phytoplankton communities sampled from the Heʻeia

Fishpond and across the northern, central, and southern sections of the bay grouped as one

nearshore community type. The similar phytoplankton communities found within the estuarine

fishpond and the nearshore environment of the bay emphasizes the connectedness of these

estuarine and coastal systems and highlights the need to manage them in close coordination.

The particularly wet conditions during the winter months appears to play a substantial

role in determining the variability of phytoplankton biomass and restructuring of the coastal food

web within the coastal environment studied here. Growing this baseline understanding of

phytoplankton cycling to relate to the timing and conditions documented in Hawaiian knowledge

systems like *kaulana mahina*, the Hawaiian lunar calendar (Nuʻuhiwa 2019), and in regard to life

cycles of bioculturally relevant species like ʻamaʻama (*Mugil cephalus*) that feeds on

microphytoplankton at juvenile stages (Hiatt 1947), would further advance this area of study and

utility for management within Hawaiian aquaculture systems.

Despite a poor understanding of the outcomes, near-island food webs will likely shift

with ongoing climate change impacts on phytoplankton communities and biomass. Collaborating

across diverse knowledge systems could improve the ability for local people to document and

adapt to changes in near-island marine resources (Winter et al. 2020b). Our collaboratively

developed study reveals distinct spatial and seasonal dynamics that define phytoplankton

communities and biogeochemical conditions from an estuarine aquaculture system on the coast

of Oʻahu, Hawaiʻi, to the open ocean of the North Pacific Subtropical Gyre. Understanding the

seasonal and spatial dynamics underlying phytoplankton communities and biogeochemistry of

near-island environments in the tropical Pacific provides the necessary knowledge to further co-

develop capacities to model and track changes to the marine food web, and to build resilience

now and in the future.

Acknowledgements

We thank Paepae o Heʻeia and the Heʻeia National Estuarine Research Reserve for their involvement in and perspectives on the directions on this research. We thank Jason Jones, Ciara Ratum, Hanako Mochimaru, Oscar Ramfelt, Amber Boettiger, Clarisse E.S. Sullivan, Elizabeth A. Monaghan, and Evan Freel for their assistance collecting samples, Gus Robertson for assistance with sonde calibration, Andrew McGowan for his help with the creation of maps, and Karen Selph for flow cytometry measurements. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation or the National Oceanic and Atmospheric Administration (NOAA). This research was funded by the National Oceanic and Atmospheric Administration (NOAA) Margaret A. Davidson Fellowship (No. NA20NOS4200123), National Science Foundation Graduate Research Fellowship Program under Grant No. 1842402, the Colonel Willys E. Lord, DVM and Sadina L. Lord Scholarship, and the University of Hawaiʻi at Mānoa Ecology, Evolution, and Conservation Biology's Maybelle Roth ARCS Award to SJ Tucker and NSF grants OCE-1538628 and OCE- 2149128 to MS Rappé. This is SOEST contribution XXX and HIMB contribution XXX.

Citations and References

Aitchison, J. 1982. The Statistical Analysis of Compositional Data. J Royal Statistical Soc Ser B

Methodol **44**: 139–160. doi:10.1111/j.2517-6161.1982.tb01195.x

Auladell, A., A. Barberán, R. Logares, E. Garcés, J. M. Gasol, and I. Ferrera. 2021. Seasonal

niche differentiation among closely related marine bacteria. Isme J 1–12. doi:10.1038/s41396-

021-01053-2

Azam, F., T. Fenchel, J. Field, J. Gray, L. Meyer-Reil, and F. Thingstad. 1983. The Ecological

Role of Water-Column Microbes in the Sea. Mar Ecol Prog Ser **10**: 257–263.

doi:10.3354/meps010257

Berube, P. M., A. Rasmussen, R. Braakman, R. Stepanauskas, and S. W. Chisholm. 2019.

Emergence of trait variability through the lens of nitrogen assimilation in Prochlorococcus.

Elife **8**. doi:10.7554/elife.41043

Bidigare, R., O. Schofield, and B. Prezelin. 1989. Influence of zeaxanthin on quantum yield of

photosynthesis of Synechococcus clone WH7803 (DC2). Mar Ecol Prog Ser **56**: 177–188.

doi:10.3354/meps056177

Bolyen, E., J. R. Rideout, M. R. Dillon, and others. 2019. Reproducible, interactive, scalable and

extensible microbiome data science using QIIME 2. Nat Biotechnol **37**: 852–857.

doi:10.1038/s41587-019-0209-9

Burson, A., M. Stomp, E. Greenwell, J. Grosse, and J. Huisman. 2018. Competition for nutrients

and light: testing advances in resource competition with a natural phytoplankton community.

Ecology **99**: 1108–1118. doi:10.1002/ecy.2187

Calbet, A., and M. R. Landry. 1999. Mesozooplankton influences on the microbial food web:

Direct and indirect trophic interactions in the oligotrophic open ocean. Limnol. Oceanogr. **44**:

1370–1380. doi:10.4319/lo.1999.44.6.1370

Callahan, B. J., J. Wong, C. Heiner, S. Oh, C. M. Theriot, A. S. Gulati, S. K. McGill, and M. K.

Dougherty. 2019. High-throughput amplicon sequencing of the full-length 16S rRNA gene

with single-nucleotide resolution. **4**: e2492-28. doi:10.1101/392332

Carswell, T., M. Costa, E. Young, N. Komick, J. Gower, and R. Sweeting. 2017. Evaluation of

MODIS-Aqua Atmospheric Correction and Chlorophyll Products of Western North American

Coastal Waters Based on 13 Years of Data. Remote Sensing **9**: 1063–24.

doi:10.3390/rs9101063

Cox, E. F., M. Ribes, and I. K. RA. 2006. Temporal and spatial scaling of planktonic responses

to nutrient inputs into a subtropical embayment. Mar. Ecol. Prog. Ser. **324**: 19–35.

doi:10.3354/meps324019

Delmont, T. O., and A. M. Eren. 2018. Linking pangenomes and metagenomes: the

Prochlorococcus metapangenome. PeerJ **6**: e4320. doi:10.7717/peerj.4320

Doré, H., G. K. Farrant, U. Guyet, and others. 2020. Evolutionary Mechanisms of Long-Term

Genome Diversification Associated With Niche Partitioning in Marine Picocyanobacteria.

Front Microbiol **11**: 567431. doi:10.3389/fmicb.2020.567431

Doty, M. S., and M. Oguri. 1956. The island mass effect. J. Cons. perm. int. Explor. Mer **22**: 33–

37. doi:10.1093/icesjms/22.1.33

Dutkiewicz, S., M. J. Follows, and J. G. Bragg. 2009. Modeling the coupling of ocean ecology

and biogeochemistry. Glob. Biogeochem. Cycles **23**: n/a-n/a. doi:10.1029/2008gb003405

Eddy, T. D., J. R. Bernhardt, J. L. Blanchard, and others. 2021. Energy Flow Through Marine

Ecosystems: Confronting Transfer Efficiency. Trends Ecol. Evol. **36**: 76–86.

doi:10.1016/j.tree.2020.09.006

Eren, A. M., E. Kiefl, A. Shaiber, and others. 2021. Community-led, integrated, reproducible

multi-omics with anvi'o. Nat Microbiol **6**: 3–6. doi:10.1038/s41564-020-00834-3

Eren, A. M., J. H. Vineis, H. G. Morrison, and M. L. Sogin. 2013. A Filtering Method to

Generate High Quality Short Reads Using Illumina Paired-End Technology. Plos One **8**:

e66643. doi:10.1371/journal.pone.0066643

Finley, A., B. Sudipto, and Hjelle. 2017. MBA: Multilevel B-Spline Approximation,.

Flombaum, P., J. L. Gallegos, R. A. Gordillo, and others. 2013. Present and future global

distributions of the marine Cyanobacteria Prochlorococcus and Synechococcus. Proc National

Acad Sci **110**: 9824–9829. doi:10.1073/pnas.1307701110

Flombaum, P., W.-L. Wang, F. W. Primeau, and A. C. Martiny. 2020. Global picophytoplankton

niche partitioning predicts overall positive response to ocean warming. Nat. Geosci. **13**: 116–

120. doi:10.1038/s41561-019-0524-2

Galili, T. 2015. dendextend: an R package for visualizing, adjusting and comparing trees of

hierarchical clustering. Bioinformatics **31**: 3718–3720. doi:10.1093/bioinformatics/btv428

Gove, J. M., M. A. McManus, A. B. Neuheimer, and others. 2016. Near-island biological

hotspots in barren ocean basins. Nature Communications **7**: 10581–8.

doi:10.1038/ncomms10581

Guillou, L., D. Bachar, S. Audic, and others. 2013. The Protist Ribosomal Reference database

(PR2): a catalog of unicellular eukaryote Small Sub-Unit rRNA sequences with curated

taxonomy. Nucleic Acids Res **41**: D597–D604. doi:10.1093/nar/gks1160

Henson, S. A., B. B. Cael, S. R. Allen, and S. Dutkiewicz. 2021. Future phytoplankton diversity

in a changing climate. Nat Commun **12**: 5372. doi:10.1038/s41467-021-25699-w

Hiatt, R. 1947. Food-Chains and the Food Cycle in Hawaiian Fish Ponds.–Part I. The Food and

Feeding Habits of Mullet (Mugil Cephalus), Milkfish (Chanos Chanos), and the Ten-Pounder

(Elops Machnata). Trans. Am. Fish. Soc. 250–261. doi:10.1577/1548-

8659(1944)74[250:fatfci]2.0.co;2

Hothorn, T., F. Bretz, and P. Westfall. 2008. Simultaneous Inference in General Parametric

Models. Biometrical Journal **50**: 346–363. doi:10.1002/bimj.200810425

Hunter-Cevera, K. R., M. G. Neubert, R. J. Olson, A. R. Solow, A. Shalapyonok, and H. M.

Sosik. 2016. Physiological and ecological drivers of early spring blooms of a coastal

phytoplankter. Science **354**: 326–329. doi:10.1126/science.aaf8536

Hyatt, D., G.-L. Chen, P. F. Locascio, M. L. Land, F. W. Larimer, and L. J. Hauser. 2010.

Prodigal: prokaryotic gene recognition and translation initiation site identification. BMC

Bioinformatics **11**: 119. doi:10.1186/1471-2105-11-119

Jokiel. 1991. Jokiel's illustrated scientific guide to Kāneʻohe Bay. 1–66.

doi:10.13140/2.1.3051.9360

Karl, D. M., and M. J. Church. 2014. Microbial oceanography and the Hawaii Ocean Time-series

programme. Nature Reviews Microbiology **12**: 699–713. doi:10.1038/nrmicro3333

Kelly, M. 1973. Some legendary and historical aspects of Heeia Fishpond, Koolau, Oahu, Bishop

Museum.

Kolde, R. 2019. pheatmap: Pretty Heatmaps. R package version 1.0.12,.

Kwiatkowski, L., O. Aumont, and L. Bopp. 2018. Consistent trophic amplification of marine

biomass declines under climate change. Glob Change Biol **25**: 218–229.

doi:10.1111/gcb.14468

Lahti, and S. Shetty. 2017. Tools for microbiome analysis in R.

Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat Meth **9**:

357–359. doi:10.1038/nmeth.1923

Laws, E. A. 1975. The Importance of Respiration Losses in Controlling the Size Distribution of

Marine Phytoplankton. Ecology **56**: 419–426. doi:10.2307/1934972

Lee, M. D. 2019. GToTree: a user-friendly workflow for phylogenomics. Bioinformatics **35**:

btz188. doi:10.1093/bioinformatics/btz188

Lee, M. D., N. A. Ahlgren, J. D. Kling, N. G. Walworth, G. Rocap, M. A. Saito, D. A. Hutchins,

and E. A. Webb. 2019. Marine Synechococcus isolates representing globally abundant

genomic lineages demonstrate a unique evolutionary path of genome reduction without a

decrease in GC content. Environmental Microbiology. doi:10.1111/1462-2920.14552

Li, G., L. Cheng, J. Zhu, K. E. Trenberth, M. E. Mann, and J. P. Abraham. 2020. Increasing

ocean stratification over the past half-century. Nat Clim Change **10**: 1116–1123.

doi:10.1038/s41558-020-00918-2

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol. **15**: 31–21. doi:10.1186/s13059-014-0550-8

Lowe, R. J., J. L. Falter, S. G. Monismith, and M. J. Atkinson. 2009. A numerical study of

circulation in a coastal reef-lagoon system. Journal of Geophysical Research **114**: 997.

doi:10.1029/2008jc005081

Mantoura, R. F. C., and C. A. Llewellyn. 1983. The rapid determination of algal chlorophyll and

carotenoid pigments and their breakdown products in natural waters by reverse-phase high-

performance liquid chromatography. Anal Chim Acta **151**: 297–314. doi:10.1016/s0003-

2670(00)80092-6

McKenzie, T., H. Dulai, and J. Chang. 2019. Parallels between stream and coastal water quality

associated with groundwater discharge. PLoS ONE **14**: e0224513.

doi:10.1371/journal.pone.0224513

McMurdie, P. J., and S. Holmes. 2013. phyloseq: an R package for reproducible interactive

analysis and graphics of microbiome census data. M. Watson [ed.]. PLoS ONE **8**: e61217.

doi:10.1371/journal.pone.0061217

Mende, D. R., J. A. Bryant, F. O. Aylward, J. M. Eppley, T. Nielsen, D. M. Karl, and E. F.

DeLong. 2017. Environmental drivers of a microbial genomic transition zone in the ocean's

interior. Nat Microbiol **2**: 1367–1373. doi:10.1038/s41564-017-0008-3

Messié, M., A. Petrenko, A. M. Doglioli, E. Martinez, and S. Alvain. 2022. Basin-scale

biogeochemical and ecological impacts of islands in the tropical Pacific Ocean. Nat Geosci 1–

6. doi:10.1038/s41561-022-00957-8

Minoche, A. E., J. C. Dohm, and H. Himmelbauer. 2011. Evaluation of genomic high-throughput

sequencing data generated on Illumina HiSeq and Genome Analyzer systems. Genome Biol

12: R112. doi:10.1186/gb-2011-12-11-r112

Möhlenkamp, P., C. Beebe, M. McManus, A. Kawelo, K. Kotubetey, M. Lopez-Guzman, C.

Nelson, and R. Alegado. 2019. Kū Hou Kuapā: Cultural restoration improves water budget

and water quality dynamics in Heʻeia Fishpond. Sustainability **11**: 161–25.

doi:10.3390/su11010161

Monger, B. C., and M. R. Landry. 1993. Flow cytometric analysis of marine bacteria with

Hoechst 33342. Applied and Environmental Microbiology **59**: 905–911.

doi:10.1128/aem.59.3.905-911.1993

Moore, J. K., W. Fu, F. Primeau, and others. 2018. Sustained climate warming drives declining

marine biological productivity. Science **359**: 1139–1143. doi:10.1126/science.aao6379

Nuʻuhiwa, K. 2019. Papakū makawalu: A Methodology and Pedagogy of Understanding the

Hawaiian Universe, p. 39–49. *In* N. Wilson-Hokowhitu [ed.], The Past Before Us :

Mookauhau As Methodology. University of Hawaii Press.

Parada, A. E., D. M. Needham, and J. A. Fuhrman. 2016. Every base matters: assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

global field samples. Environmental Microbiology **18**: 1403–1414. doi:10.1111/1462-

2920.13023

Pritchard, L., R. H. Glover, S. Humphris, J. G. Elphinstone, and I. K. Toth. 2015. Genomics and

taxonomy in diagnostics for food security: soft-rotting enterobacterial plant pathogens. Anal.

Methods **8**: 12–24. doi:10.1039/c5ay02550h

Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer, P. Yarza, J. Peplies, and F. O. Glöckner.

2012. The SILVA ribosomal RNA gene database project: improved data processing and web-

based tools. Nucleic Acids Res **41**: D590–D596. doi:10.1093/nar/gks1219

Redfield, A. C. 1960. The biological control of chemical factors in the environment. Sci Prog **11**:

150–70. doi:10.1086/646891

Ringuet, S., and F. T. Mackenzie. 2005. Controls on nutrient and phytoplankton dynamics during

normal flow and storm runoff conditions, southern Kaneohe Bay, Hawaii. Estuaries **28**: 327–

337. doi:10.1007/bf02693916

Ruf, T. 1999. The Lomb-Scargle Periodogram in Biological Rhythm Research: Analysis of

Incomplete and Unequally Spaced Time-Series. Biol Rhythm Res **30**: 178–201.

doi:10.1076/brhm.30.2.178.1422

Selph, K., E. Goetze, M. Jungbluth, P. Lenz, and G. Kolker. 2018. Microbial food web

connections and rates in a subtropical embayment. Mar Ecol Prog Ser **590**: 19–34.

doi:10.3354/meps12432

Smith, S. V. 1981. Responses of Kaneohe Bay, Hawaii, to Relaxation of Sewage Stress, p. 391–

410. *In* Estuaries and Nutrients. Estuaries and Nutrients. doi:10.1007/978-1-4612-5826-1_18

Stock, C. A., J. G. John, R. R. Rykaczewski, and others. 2017. Reconciling fisheries catch and

ocean productivity. Proc National Acad Sci **114**: E1441–E1449.

doi:10.1073/pnas.1610238114

Stoecker, D. K., P. J. Hansen, D. A. Caron, and A. Mitra. 2016. Mixotrophy in the Marine

Plankton. Annu Rev Mar Sci **9**: 1–25. doi:10.1146/annurev-marine-010816-060617

Strimmer, K. 2008. fdrtool: a versatile R package for estimating local and tail area-based false

discovery rates. Bioinformatics **24**: 1461–1462. doi:10.1093/bioinformatics/btn209

Suzuki, M. T., O. Béjà, L. T. Taylor, and E. F. DeLong. 2001. Phylogenetic analysis of

ribosomal RNA operons from uncultivated coastal marine bacterioplankton. Environmental

Microbiology **3**: 323–331. doi:10.1046/j.1462-2920.2001.00198.x

Tucker, S. J., K. C. Freel, E. A. Monaghan, C. E. S. Sullivan, O. Ramfelt, Y. M. Rii, and M. S.

Rappé. 2021. Spatial and temporal dynamics of SAR11 marine bacteria across a nearshore to

offshore transect in the tropical Pacific Ocean. Peerj **9**: e12274. doi:10.7717/peerj.12274

Utter, D. R., G. G. Borisy, A. M. Eren, C. M. Cavanaugh, and J. L. M. Welch. 2020.

Metapangenomics of the oral microbiome provides insights into habitat adaptation and

cultivar diversity. **1**: 808410–40. doi:10.1101/2020.05.01.072496

Wei, T., and V. Simko. 2021. corrplot: A visualization of a correlation matrix.

Welschmeyer, N. A. 1994. Fluorometric analysis of chlorophyll a in the presence of chlorophyll

b and pheopigments. Limnol. Oceanogr. **39**: 1985–1992. doi:10.4319/lo.1994.39.8.1985

Wickham, H. 2016. ggplot2: elegant graphics for data analysis. Springer-Verlag New York,.

Willis, A. D., and B. D. Martin. 2020. Estimating diversity in networked ecological

communities. Biostatistics **23**: kxaa015. doi:10.1093/biostatistics/kxaa015

Winter, K., N. Lincoln, F. Berkes, and others. 2020a. Ecomimicry in Indigenous resource

management: optimizing ecosystem services to achieve resource abundance, with examples

from Hawaiʻi. Ecol. Soc. **25**. doi:10.5751/es-11539-250226

Winter, K., Y. Rii, F. Reppun, and others. 2020b. Collaborative research to inform adaptive

comanagement: a framework for the Heʻeia National Estuarine Research Reserve. Ecol Soc

25. doi:10.5751/es-11895-250415

Wright, E. 2016. Using DECIPHER v2.0 to Analyze Big Biological Sequence Data in R. The R

Journal.

Xenopoulos, M. A., J. A. Downing, M. D. Kumar, S. Menden-Deuer, and M. Voss. 2017.

Headwaters to oceans: Ecological and biogeochemical contrasts across the aquatic continuum

M. Xenopoulos, J.A. Downing, M.D. Kumar, S. Menden-Deuer, and M. Voss [eds.]. Limnol.

Oceanogr. **62**: S3–S14. doi:10.1002/lno.10721

Yeo, S. K., M. J. Huggett, A. Eiler, and M. S. Rappé. 2013. Coastal bacterioplankton community

dynamics in response to a natural disturbance D.L. Kirchman [ed.]. PLoS ONE **8**: e56207-14.

doi:10.1371/journal.pone.0056207

Data availability statement

Data supporting the results within the manuscript are available within the main text (See

Materials & Methods). Sequencing data are available in the National Center for Biotechnology

Information (NCBI) Sequence Read Archive (SRA) under BioProject number PRJNA706753 as

well as PRJNA971314. Environmental data were submitted to BCO-DMO under

https://www.bco-dmo.org/project/663665. Code used in the analysis is available is available at

https://github.com/tucker4/Tucker_Phytoplankton_KByT_HeNERR.

Author contribution statement

900 SJT led the formal analyses and wrote the initial manuscript draft. SJT, YMR, KFC, MSR

901 collected the samples with assistance from AHK and KK. SJT, KFC, YMR, and MSR processed

902 samples. All authors were involved in the conceptualization, development of methodology,

903 interpretation of results, and providing input to the manuscript draft and revisions.