Title:

Human hypothalamic neural stem cells transplantation ameliorates age-associated

dysfunction through SNS/eNAMPT axis activation in aged mice

Yanuar Alan Sulistio[2,§] , Yuna Lee[4,5,§], Kelvin Pieknell[2,§], Sebin Hong[2-3], Jumi Kim[2-3], Min Jong

Seok[2-3,] Na-Kyung Lee[2-3], Kyu-Sang Park[6], Taeui Hong[6], Suyeon Choi[6], Ki Woo Kim[7], Dong Joo

Yang[7], Woong-Yang Park[8,9], Kyung Yeon Han[8,9], Seul Gi Yoon[10], Il Yong Kim[10,11], Je Kyung

Seong[10,11], Tae Yong Lee[12,13], Min Sung Kim[12,13], Min Soo Kim[4,5,]*, Sang-Hun Lee[1-3,]*

[1]Department of Biochemistry and Molecular Biology, College of Medicine, Hanyang University;

Seoul, South Korea

[2]Hanyang Biomedical Research Institute, Hanyang University; Seoul, South Korea

[3]Graduate School of Biomedical Science and Engineering, Hanyang University; Seoul, South

Korea

[4]Brain Science Institute, Korea Institute of Science and Technology (KIST); Seoul, South Korea

[5]Division of Bio-Medical Science and Technology, KIST School, University of Science and

Technology; Seoul, South Korea

[6]Mitohormesis Research Center, Yonsei University Wonju College of Medicine; Wonju, South

Korea

[7]Division of Physiology, Departments of Oral Biology and Applied Life Science, BK21 FOUR,

Yonsei University College of Dentistry; Seoul, Korea

[8]Geninus Inc.; Seoul, South Korea

[9]Samsung Genome Institute, Samsung Medical Center; Seoul, South Korea

[10]Korea Mouse Phenotyping Center (KMPC), Seoul National University; Seoul, South Korea

[11]Laboratory of Developmental Biology and Genomics, Research Institute for Veterinary

Science, and BK21 PLUS Program for Creative Veterinary Science Research, College of

Veterinary Medicine, Seoul National University; Seoul, South Korea

[12]Central Research Center, Corestem Inc; Seoul, South Korea

[13]College of Pharmacy, Chungbuk National University; Cheongju, South Korea

[§]These authors contributed equally.

*Correspondence: <u>Sang-Hun Lee, M.D., Ph.D.,</u> Department of Biochemistry and Molecular

Biology, College of Medicine, Hanyang University, Seoul, 04763, South Korea. Telephone:

82-2-2220-0625; Fax: 82-2-2220-2422; E-mail: leesh@hanyang.ac.kr; or <u>Min Soo Kim, Ph.D</u>.,

Brain Science Institute, Korea Institute of Science and Technology (KIST), Seoul, 02792,

South Korea. E-mail: minsoo.kim@kist.re.kr.

Abstract

The hypothalamus is the brain region that regulates systemic body metabolism and multiple brain functions. The adult hypothalamus harbors neural stem/precursor cell (NSC)-like cells. Along with age-related body changes, the hypothalamic NSC (htNSC) population declines, indicating the potential of htNSC replacement as an anti-aging strategy. Here, we developed protocol to generate htNSCs from human pluripotent stem cells (hPSCs). Implanting the hPSC-derived htNSCs into the hypothalamus of aged mice ameliorated age-related declines in metabolic fitness, physical capacity, and cognitive function. Mechanistically, these anti-aging effects were mediated by inter-tissue communication: enhanced neuronal activity in the htNSC-transplanted hypothalamus stimulated adipose tissues to produce and release the anti-aging molecule eNAMPT into systemic circulation via the sympathetic nervous system. Concurrently, the aged inflammatory environment in the hypothalamus was alleviated by peripheral anti-aging signals. Collectively, our findings support the potential of anti- or healthy aging therapies by targeting hypothalamus.

Keywords: Hypothalamus; Metabolism; NAMPT; sympathetic nervous system; Homeostasis; Hypothalamic neural stem cells; Human pluripotent stem cells; Aging; Anti-aging;

INTRODUCTION

Aging is a significant health concern, and developing methods to retard aging remains a prime interest of humanity. The hypothalamus, a brain region encompassing the third ventricle, functions as a high-order control center for body metabolism, energy balance, and homeostasis, all of which decline with age. Hypothalamic functions are mediated by regulating hormone production through the pituitary gland and overseeing the autonomic nervous system, thereby affecting peripheral organs like the liver, adipose tissue, skeletal muscle, bone, and intestine. Additionally, the hypothalamus has neurocircuitry connections with various brain regions, thus controlling brain functions, including circadian rhythm regulation and cognitive abilities, which also decline with age [1-8]. Therefore, hypothalamus is integral part of inter-organ communication network that govern metabolism that can be utilized for anti-aging interventions. [9].

Hypothalamic functions are governed by intricate neuro-circuitry networks involving a myriad of neuronal subtypes with hypothalamus-specific neuroendocrine functions. In addition, growing evidence indicates that non-neuronal glial cells play equally important roles in the hypothalamus, such orchestrating intra-hypothalamic functions and the hypothalamic response to metabolic cues from the body, by participating in synaptic transmission, synaptic plasticity, and nutrient/hormone sensing [10-14]. Adult hypothalamus, particularly within the third ventricle wall and median eminence, harbors cells with neural stem cell properties known as hypothalamic neural stem cells (htNSCs) or tanycytes, from which new neurons can form [15-18]. (Hereafter, 'tanycyte' will refer to quiescent hypothalamic NSCs in the adult hypothalamus, while 'hypothalamic NSC' will be used as a general term regardless of developmental and proliferation stages) [17-22]. Given that neurogenesis in the hypothalamus steeply drops after the postnatal period and is almost negligible in adulthood [23,24], the strategy for replenishing htNSCs might be needed to restore metabolic homeostasis during aging.

To develop a strategy for restoring metabolic balance, which is dysfunctional during aging, by targeting the hypothalamus, we differentiated hPSCs into htNSCs with neurogenic and gliogenic potential. These hPSC-derived htNSCs were then implanted into the hypothalami of aged mice. We observed improved metabolic health in the transplanted group, accompanied by several anti-

aging effects. This treatment increased neuronal activity in the dorsomedial hypothalamic

(DMH) region and subsided glial inflammation in the surrounding third ventricle region.

Furthermore, our findings suggest that the beneficial effects in these rejuvenated mice were

orchestrated by intricate inter-tissue communication between the hypothalamus and adipose

tissues, mediated by the sympathetic nervous system

RESULTS

Generation of htNSCs from hPSC-derived hypothalamus-like organoids

Despite the great clinical importance of the hypothalamus, studies on hypothalamic functions

have been made exclusively in animal models. Their translation to humans remains nascent,

mainly due to the inaccessibility of human hypothalamic tissues and cells. Several studies have

generated hypothalamic neuronal cell types from human pluripotent stem cells (hPSCs) [25-27].

Given the functional significance of glia and htNSCs in the hypothalamus, however, it is

compelling to develop research platforms that can provide diverse hypothalamic cell types,

including non-neuronal cells, of human origin. We aimed to develop a method to differentiate

hPSCs into htNSCs with neuro- and gliogenic potential.

In this work, we leveraged the principles and advantages of the organoid-based method to

prepare human htNSCs from hPSCs [28]. To this end, human embryonic stem cells (hESC, H9)

were induced to differentiate into hypothalamus-like organoids in the presence of small

molecules for inducing hypothalamic fate [25-27] (Fig. 1A and Suppl. Fig. S1A-B). The

combinations, concentrations, and treatment periods of the hypothalamic patterning molecules

were optimized based on high expression of the hypothalamus–specific markers *RAX, NKX2-1,*

OTP, ISL1, and *SF-1* and low expression of the midbrain marker Engrailed-1 (*EN1)* (Suppl. Fig.

S1A). Treatment with SHH and purmorphamine during early organoid differentiation was

indispensable for hypothalamus-specific marker expression (Fig. 1B and Suppl. Fig. S1A).

Further addition of CHIR, CNTF, IWP-2, XAV, and FGF8, which have been reported as

molecules for hypothalamic patterning [25-27], did not enhance but lower hypothalamic marker

expression (Suppl. Fig. S1A). Cells dissociated from the organoids on DIV 20 were efficiently

sub-cultured as monolayer. Generated htNSCs expressed markers specific for general NSCs

(SOX2, NESTIN), proliferating cells (KI67), and hypothalamic specific markers, indicating

efficient generation of a homogenous htNSC culture from the early stage of the organoids (Fig.

1B–C). The expressions of hypothalamus-specific markers and the percentage of Nestin+ and

RAX+ htNSCs in our method were significantly higher compared to those observed in htNSC

cultures generated using other published protocols [25,26] (Suppl. Fig. S1B-E). To further confirm

their hypothalamic identity, we performed RNA sequencing (RNA-seq) on our cultured htNSCs.

When comparing our htNSC transcriptome profile to various adult human brain transcriptome

(GTEx Database, ages of 22-68 years), the htNSC was clustered together with the human adult

hypothalamus, signifying a high hypothalamic regional identity (Fig. 1D).

Additionally, the htNSCs were expandable for extended periods (Suppl. Fig. S2A), could be

stored and re-cultured from liquid N2 without loss of viability or naïve htNSC-specific marker

expression (Suppl. Fig. S2B), establishing the htNSC culture derived from hPSC-organoids as a

stable and readily available source of human htNSCs. Proliferation, indicated by PDL, decreased

in late passages (Suppl. Fig. S2A), signifying replicative senescence.

Differentiation phenotypes of the hPSC-derived htNSCs

Upon terminal differentiation, the htNSCs could differentiate into mature neurons expressing

(TUBB3). It also expressed hypothalamus-specific neuropeptides such as POMC, NPY, SCGN,

and PMCH (Fig. 1E). Astrocyte was detected in the differentiated htNSC cultures (Fig. 1E),

which increase in proportion in the later passages (Suppl. Fig. S2C). Hypothalamic neurons and

astrocytes sense the nutritional status of the body through peptides such as leptin and ghrelin

that are generated in peripheral organs. An administration of leptin to the differentiated htNSC

cultures activated the major leptin-mediated STAT intracellular pathway, which was shown by

increased phosphorylated STAT3 (pSTAT3) levels in Western-blot analysis (Fig. 1F, left). In

addition, the phosphorylation of FOXO1, the intracellular signal transducer of ghrelin, occurred

when the differentiated htNSC cultures were treated with ghrelin (Fig. 1F, right) [29] indicating

that the differentiated hypothalamic cells functionally sensed nutritional cues. Using this

protocol, htNSCs with neurogenic and gliogenic capacities were also generated from the human

induced pluripotent stem cell (hiPSC) lines CMC-hiPSC-011 and CS-001(Suppl. Fig. S3).

Next, we performed a single-cell RNA-seq (scRNA-seq) analysis of the differentiated htNSC

culture after seventy days of terminal differentiation. After filtering doublets and clusters cells

with low-quality reads, we used 6,394 cells for the analysis. Unsupervised clustering of the

dataset revealed thirteen clusters, which were projected onto a uniform manifold approximation

(UMAP) map for visualization. On the basis of the expression levels of well-characterized

marker genes [30-32] alongside with automated identification using scType, the cluster identities

were designated as proliferating neural stem cells (NSCs), intermediate progenitor cells (Int1, 2),

interneuron progenitor cells (iNPCs), astrocyte progenitor cells (APCs), neurons (Neu1, 2),

astrocytes (Ast1, 2, 3), radial glia (RGs), proteoglycan-secreting glia (PGGs), and tanycytes

(Tan) (Fig. 1G-I and Suppl. Fig. S4A). For unbiased confirmation of the hypothalamic fate, we

mapped the gene expression signatures of the cell clusters to the mouse brain transcriptome (at

post-natal day 28) in the Allen Developing Brain Atlas using Voxhunt [33]. Significantly, our

scRNA-seq profiles showed that every identified cluster in the differentiated htNSC culture

precisely corresponded to the hypothalamic region of the atlas (Fig. 1J and Suppl. Fig. S4B).

Furthermore, the progenitor and intermediate cell clusters (Int1, Int2, iNPC, APC, NSC) were

enriched to the hypothalamic region adjacent to the ventricular area, where adult neurogenesis

occurs (Fig. 1J and Suppl. Fig. S4B). To provide additional confirmation, we cross-referenced

our single-cell transcriptome data with published datasets of human fetal hypothalamus

(gestational week (GW) 18-22) (Suppl. Fig. S4C-D) [34]. The clustering analysis indicates that our

single-cell transcriptome data align closely with the human fetal hypothalamus datasets (Suppl.

Fig. S4E-F). Expectedly, oligodendrocytes and microglia were absent in the differentiated htNSC

(Suppl. Fig. S4G). Next, we performed a gene regulatory network analysis using pySCENIC and

identified the potential upstream master regulator (regulon) of each cell cluster (Suppl. Fig.S4H).

The hypothalamus is a complex brain area that contains myriad subtypes of neurons with

neuroendocrine and neurotransmitter identities. To unravel the identity of the neuronal

populations in our differentiated htNSC cultures, we performed further clustering on the

neuronal clusters (Neu1 and Neu2). Upon re-clustering based on the signature transcripts, we

detected 8 subclusters of neurons (n1–n8) (Suppl. Fig. S5A), comprising neuronal clusters with

different neurotransmitter identities, such as glutamatergic (n6), GABAergic (n2), dopaminergic

(n8), and dopaminergic/glutamatergic (n1, n3, n4), and clusters with specific hypothalamic

neuroendocrine identities, such as those expressing *SCGN* (n5), *PMCH* (n4), *TRH* (n3), and

NR3C1 (n1) (Suppl. Fig. S5B-C). In addition, the neuronal subclusters were classified by their

unique transcriptional factors and regulons (Suppl. Fig. S5B-C). Of note, *VGF*, a gene that is

highly enriched in the adult hypothalamic neurons [35,36], was widely expressed throughout the

neuronal clusters (Suppl. Fig. S5B).

The APC cluster branched into three astrocyte trajectories (Suppl. Fig. S5D). Astrocyte cluster

Ast2 is regarded as a homeostatic astrocytic population that expresses *SLC1A3* (GLAST), a

protein that provides metabolic support for neighboring neurons (Suppl. Fig. S6F). Additionally,

Ast2 could be involved in neuroendocrine functions because of its specific expression of genes

such as *SLCO1C1* (thyroid hormone transporter), *CRYM* (Crystalin Mu, a thyroid hormone

binding protein), and *SCG2* (secretogranin II, a neuroendocrine secretory protein) (Suppl. Fig.

S5E). By contrast, Ast1 preferably expressed genes related to immune function such as *RSAD2*

and *TRIL* (Suppl. Fig. S5E). Moreover, interferon-induced genes, such as *ISG5*, *IFIT*, and

STAT1, were also specifically expressed in Ast1, suggesting an immunogenic phenotype. Aside

from the "classical" homeostatic and reactive astrocytes, we also detected a third type of

astrocyte (Ast3) that was derived from common progenitor cells (APCs) and expressed astrocyte-

related transcription factors (*SOX9*, *ID3*, *BHLHE41*) and common astrocyte markers (*APOE*,

AQP4, *SPARCL1*, *SPARC*) but was devoid of *GFAP* (Suppl. Fig. S5E). Ast3 is characterized by

gene expression specific for not only tight junction formation (*CLDN5*, *ALCAM*, *COL1A2*), but

also the astrocyte population in the adult subventricular zone (*Meis2*, *Col1a2*), which interacts

with ependymal cells or pericytes [37-39]. Of note, a cluster was designated as a tanycyte-like

population because it exclusively expressed tanycyte-specific markers such as *RAX*, *FGF10*,

COL25A1, *GPC3*, and *DIO2* (Suppl. Fig. S5E). Interestingly, *LEPR* (leptin receptor) expression

was detected in the tanycyte cluster (Suppl. Fig. S5E), supporting the involvement of tanycytes

in metabolic homeostasis via leptin signaling [19]. The presence of tanycyte-like cells post-

differentiation, further suggest hypothalamic identity of our htNSC.

In vitro **htNSC differentiation recapitulated hypothalamic brain development**

Next, we used Monocle 3 to perform pseudo-temporal ordering of the transcriptional dynamics

of the cell clusters that were differentiated from human htNSCs. Consistent with brain

development, in which neurons and astrocytes arise sequentially from common progenitors, the

UMAP clustering shows that the intermediate progenitor cell clusters (Int1, 2) were bifurcated

into neurons (Neu1, 2) and the APC cluster that gave rise to the astrocytic populations (Ast1–3)

(Suppl. Fig. S6A). Several genes showed temporal dynamic changes in gene expression that

recapitulated those in the developing brain. For example, a steep decrease in the expression of

the proliferation marker *CDK1* at an early time point was followed by a surge in the expression

of the pro-neural gene *ASCL1* at the late progenitor cell stage and increased expression of the

neuron-specific *TUBB3* and *SNAP25* at the terminal stage of the neuronal trajectory (Suppl. Fig.

S6B). The *SLIT-ROBO* signal has recently been identified as a critical regulator of

hypothalamus-specific brain development [40]. Of note, the *SLIT-ROBO* signal genes in the

differentiated htNSC cultures showed pseudo-time patterns similar to those of *in vivo*

hypothalamic brain development (Suppl. Fig. S6C), indicating our *in vitro* htNSC differentiation

recapitulates hypothalamus development *in vivo*.

Thus far, the developmental origin of tanycytes has been unidentified. Interestingly, consistent

with a developmental trajectory analysis suggesting that tanycytes originate from the astrocyte

population during hypothalamic brain development [40], our pseudo-time analysis indicates that

tanycyte-like cells (Tan) developed from the early Ast2 population (Suppl. Fig. S6D-E).

Age-associated declines in physical and cognitive activity were ameliorated in mice that

received hPSC-derived htNSCs

Physical strength, muscle endurance, motor coordination, memory, and cognitive function are all

declining with age [41], and those age-associated changes are linked to hypothalamic dysfunction

in regulating body metabolism and homeostasis [42]. A previous study has shown that age-

associated impairment of physical and cognitive activity could be delayed by implantation of

htNSCs derived from mouse pups [10]. To take a critical step toward the clinical translation of the

htNSC-based anti-aging strategy, we transplanted single dissociates of human htNSCs (passage

2-3) prepared from the hESC-derived hypothalamus-like organoids into the hypothalami of mice

at 12-months of age (Fig. 2A). The mouse htNSCs derived from newborn mice, which were used

in the previous study [10], survived poorly after implantation into the mouse hypothalamic region,

and thus genetic engineering with dominant recombinant IκB to inhibit the NF-κB inflammatory

signal was needed to overcome the survival problem [10]. By contrast, upon transplantation into the

medio-basal hypothalamus (MBH), the hPSC-derived human htNSCs survived and integrated

efficiently into the host hypothalamus, forming human-specific marker (hNCAM or human

mitochondrial-specific antigen)-positive grafts, portions of which expressed undifferentiated

htNSC (RAX), differentiated neuronal (NeuN), and astrocytic (human GFAP, hGFAP) markers

2–4 weeks post-transplantation (Fig. 2B–C).

To ensure reproducibility, we designed multiple cohorts of cell transplantation with

independently prepared human htNSCs and aged mice. In those consecutive trials of htNSC

transplantation, the mice that received transplantation displayed a significant improvement in

exploratory behavior and locomotor activity (Fig. 2D), coordination (rotarod) (Fig. 2E), muscle

endurance (grip test) (Fig. 2F), and treadmill performance (Fig. 2G), compared with the age-

matched sham-operated control mice (control, culture media-injected). Furthermore, the long-

term and spatial memory and cognitive functions of the grafted mice were significantly improved

in the Morris water-maze (MWM; Fig. 2H–I) and novel object recognition tests (Fig. 2J). The

physical activities and memory/cognitive functions in the grafted old mice were also compared

with the young mice (aged 3-6 months) (Fig. 2 and Suppl. Fig. S7).

Improvement of body energy metabolism and tissue aging by intra-hypothalamic htNSC

transplantation

The grafted mice were leaner, devoid of age-dependent body weight gain during the post-

transplantation period (12–16 months) that was manifested in the vehicle-injected control mice

(Fig. 3A). There was no difference in the food intake and physical activity changes only occurred

during the dark phase (Fig. 3B–C). Consequently, the htNSC-grafted mice displayed greater

energy expenditure in the metabolic cage compared to control aged mice (Fig. 3D). Body

composition analysis indicates that the htNSC-grafted mice had markedly lower fat mass with

increased lean mass indicating lower adiposity (Fig. 3E). Accordingly, lipid droplet size was

greatly reduced in the inguinal white adipose tissue (iWAT), epididymal white adipose tissue

(eWAT), and intrascapular brown adipose tissue (iBAT) of the grafted mice than the control

mice (Fig. 3F–H). Similarly, hepatic steatosis in the aged mice was markedly decreased in the

transplantation group (Fig. 3I).

Growth/differentiation factor 15 (GDF15) and fibroblast growth factor 21 (FGF21) are key

secretory proteins that stimulate peripheral tissues to increase energy expenditure and lipid

consumption [43-46]. Hepatic GDF15 and FGF21 proteins and their upstream regulator ATF4 were

significantly upregulated in the livers of the grafted mice (Fig. 4A). Additionally, FGF21 was

also increased in iWAT and eWAT of the grafted mice, which was accompanied by the

upregulated protein expression of MT-CO1 and SDHB, mitochondrial proteins that execute

energy consumption/thermogenesis (Fig. 4B–C) [47,48]. Consistently, rectal temperature was

significantly higher in the htNSC-grafted mice than the controls (Fig. 4D). Interestingly, this

metabolic signature closely resembles that observed in leucine-deprived diet, which is known to

promote metabolic health and prolong lifespan [49]. The hypothalamus influences the metabolism

and physiology of adipose tissues through the sympathetic nervous system (SNS) [50]. Messenger

RNA expression of β1, 2, and 3 adrenoreceptors (*Adrb1*, *Adrb2*, and *Adrb3*) and lipolytic genes

(*Dio2* and *Lipe*) was significantly upregulated in the eWAT of the grafted mice (Fig. 4E),

implying SNS activation in the WAT of the grafted mice. Along with efficient energy

metabolism, the mice that received intra-hypothalamic htNSC transplantation showed

significantly better glucose tolerance than the control mice (Fig. 4F). A bone μCT examination

showed osteoporotic changes in the tibias and femurs of the control mice at age 16 months (Fig.

4G-H). That bone loss was greatly prevented by intrahypothalamic htNSC transplantation, as

evidenced by a significant increase in bone mass density (BMD), bone volume/total volume

(BV/TV) ratio, trabecular thickness (Tb.Th.), and trabecular number (Tb.N.) in the grafted mice

(Fig. 4G-H). Notably, the increase in bone density coincided with a substantial decrease in the

osteoclast population among the transplanted mouse group (Fig. 4I), suggesting that suppression

of bone resorption underlies the observed enhancement in bone density.

293 **Increase of circulating eNAMPT through hypothalamic-adipose inter-tissue**

294 **communication underlies the systemic aging retardation in the htNSC-grafted mice**

295 To attain mechanistic hints behind the improvement in age-associated behaviors in the

296 hypothalamus-transplanted mice, we first measured biomolecules associated with body

297 metabolism/aging in circulating blood. Among those tested using a cytokine array kit, leptin

298 levels were lower in the grafted mice compared to the control mice (Suppl. Fig. S8A–B), which

299 was further validated by ELISA-based quantification (Suppl. Fig. S8C). This finding aligns with

300 the established positive correlation between plasma leptin levels and body fat mass [51,52]. The

301 reduced levels of leptin also correspond to the increased bone density observed in the grafted

302 mice [53]. Lipid blood chemistry analysis showed dramatic reduction of LDL-C but no changes in

303 the other factors examined (Suppl. Fig. S8D-G).

304

305 Additionally, we observed significantly higher levels of extracellular nicotinamide

306 phosphoribosyltransferase (eNAMPT), an anti-aging factor secreted from adipose tissues [54-56], in

307 the circulating blood of the grafted mice (Fig. 5A). The mRNA expression of NAMPT was also

308 significantly elevated in the iWAT and eWAT of the hNSC-grafted mice (Fig. 5B). Treatment

309 with a NAMPT inhibitor (FK-866, administered intraperitoneally, dosage: 10 mg/kg/day) or

310 beta-adrenergic (SNS) blocker (propranolol, administered via drinking water, dosage: 0.5 g/L)

311 two weeks post-transplantation (Fig. 5C), largely negated the anti-aging effects of the hNSC

312 transplantation, both in terms of aging-related physiological declines (Fig. 5D-J) and histologic

313 changes (Fig. 5K), indicating that the increased eNAMPT and/or symphathetic tone in peripheral

314 tissue and blood plays a crucial role in the anti-aging effects observed in the htNSC-transplanted

315 mice.

316

317 Furthermore, our blood examination revealed that circulating levels of ketone bodies were

318 significantly elevated in the transplanted mice (Fig. 5H) and reversed upon NAMPT inhibitor

319 treatment or beta-adrenergic blocker, suggesting that the increase in ketone bodies may also

320 mediates the anti-aging. Conversely, this rise in ketone bodies may act as a factor to correct

321 functional deficits and inflammation in the aged hypothalamus [57,58], indicating a potential positive

322 feedback loop between elevated ketone body levels and improved hypothalamic function and

inflammation. Ketone bodies, which increase in the blood during calorie restriction, are

recognized as anti-aging metabolites [59-61]. These findings collectively indicate that the htNSC-

grafted hypothalamus stimulates adipose tissue to synthesize and release eNAMPT into systemic

circulation via the sympathetic nervous system, thereby retarding systemic aging.

Single-cell transcriptome analysis of aged hypothalamic brains grafted with htNSCs

To identify the transcriptional profile of the grafted hypothalami, we performed a droplet-based

single-nucleus RNA-seq (snRNA-seq) analysis of cell nuclei extracted from the transplanted

hypothalamus and control groups 4 months post-transplantation. We sampled six hypothalami

(three per group) on four independent sequencing runs (two runs per group) (Fig. 6A). After

filtering and doublet removal, 12,161 nuclei were clustered and projected onto a two-

dimensional UMAP space for visualization (Fig. 6B). We identified 30 clusters: 23 neuron (10

clusters of GABAergic neurons and 13 clusters of glutamatergic neurons), two tanycyte, one

astrocyte, one oligodendrocyte, one oligodendrocyte precursor cell (OPC), one microglia, and

one pericyte cluster. For the sake of simplicity and general visualization, the 23 neuronal clusters

were merged into glutamatergic and GABAergic neuron clusters, as described above, and the

two tanycyte clusters were merged into one (Fig. 6B and Suppl. Fig. S9A). The cells were

annotated manually based on well-established cellular markers: *Snap25* for pan-neurons, *Gad1*

for GABAergic neurons, *Slc17a6* for glutamatergic neurons, *Agt* and *Slc1a3* for astrocytes, *Mbp*

and *Plp1* for oligodendrocytes, *Olig2* and *Pdgfra* for OPCs, *P2ry12* and *Cx3cr1* for microglia,

Rax and *Col23a1* for tanycytes, and *Slc4a7* and *Flt1* for pericytes (Fig. 6C and Suppl. Fig. S9B).

Many well-known hypothalamus-specific neuronal subtypes were identified in the neuron

subclusters. For example, GABA neuron subcluster 2 (GABA-2), which contained galanin-

secreting neurons, are highly enriched in the hypothalamus and could be involved in feeding

behavior [62,63], and Glut-11 is a melanin-concentrating hormone (MCH; gene name, *Pmch*)

neuron, which is a sub-population of neurons in the lateral hypothalamus (Fig. 6D–E and Suppl.

Fig. S10A). A gene ontology (GO) analysis showed that the top signature genes in each cluster

were enriched in the GOs consistent with their known biological functions. For instance, GOs

regarding 'synaptic transmission' and 'neurotransmitter' were specifically enriched in the

neuronal clusters, and GOs regarding 'immune process' were specifically enriched in the

microglial cluster (Fig. 6F). No significant difference in the number of cells in each cluster was

found between the grafted and control groups (Fig. 6G).

Next, we performed a differential gene expression (DEG) analysis on each cluster to examine the

cell-specific changes that could be mediating the anti-aging effect of the intra-hypothalamic

htNSC-transplantation. Interestingly, no statistically significant DEGs were detected in the pan-

GABA and pan-Glut clusters (Fig. 6H). By contrast, abundant DEGs were detected in non-

neuronal cell clusters. Among them, the microglia showed the most dramatic gene expression

changes following intra-hypothalamic htNSC transplantation, which was followed by tanycytes,

astrocytes, OPCs, and oligodendrocytes. These findings are consistent with the previous report

that glia-specific genes predict age with greater precision than neuron-specific genes [64-66].

The microglial DEGs that were downregulated in the grafted hypothalami were enriched in the

GOs associated with immune response, specifically 'IκB kinase/NF-κB signaling' (Fig. 6I).

Indeed, the expression of central microglial NF-κB subunits (*Nfkb1*, *Rel*, *Relb*, and *Ikbkg*) were

dramatically downregulated in the htNSC-grafted group (Fig. 7E). This is consistent with the

hypothalamic microinflammation theory of aging, which is in turn supported by a recent single-

cell transcriptomic result about the aged hypothalamic median eminence [67,68]. Specifically, it has

been shown that hypothalamic Iκκb and NF-κB are molecules responsible for aging [10,69] and that

htNSCs dampen NF-κB signaling and hypothalamic micro-inflammation in a paracrine manner

[70]. The microglial genes upregulated by htNSC transplantation were annotated to GOs related to

'metabolic process,' specifically 'lipid metabolism' (Fig. 6I). Consistently, it has been reported

that dysregulated lipid metabolism and lipid accumulation in microglia represent a dysfunctional

and pro-inflammatory state in the aging brain [71]. We further compared gene expression changes

in the microglia of grafted hypothalami (vs. vehicle-injected controls) with those of age-

associated microglial genes recorded in multiple published datasets [67,72,73]. Single-cell expression

levels of the microglial genes downregulated during aging in all the published datasets

(calculated using 'AddModuleScore' in the Seurat package) were significantly upregulated in the

microglia of grafted hypothalami, compared with the levels in the vehicle-treated control

hypothalami (p-value <0.05, statistical method Wilcoxon rank-sum test) (Fig. 6J). Concordantly,

microglial genes with age-dependent upregulation in the datasets of Chen (2022) and Ximarekis

(2019) were significantly downregulated in the microglia of the grafted hypothalami, together

indicating that age-dependent microglial signatures were corrected in aged hypothalami by

htNSC transplantation.

The tanycyte cluster had enrichment in GOs related to cell-to-cell and intracellular signaling

(Fig. 6I). That finding is consistent with the role of naïve tanycytes as the portal between the

hypothalamus and the rest of the body, the place where hormones and metabolic cues are

exchanged. The most enriched GO in the tanycytes of the grafted hypothalami was 'positive

regulation of hydrolase activity.' It has been demonstrated that after being released from

hypothalamic neurons, pro-TRH requires hydrolase-mediated activation in tanycytes [22,74].

Therefore, it is possible that htNSC transplantation also restores the tanycyte-mediated TRH

signaling process that usually decreases during aging [75].

We further analyzed the htNSC transplantation–mediated DEGs in 23 neuron subclusters

characterized by their unique gene expression signatures (Suppl. Fig. S10A). GABA-4 and

GABA-6 were the GABAergic neuron clusters whose gene expressions were most highly

modulated by htNSC transplantation, whereas Glut-5, Glut-6, Glut-7, and Glut-8 were the most

highly affected among the glutamatergic neuronal clusters (Suppl. Fig. S10B). Among those, it is

noted that GABA-4 was the *Vipr2+* neuronal population associated with circadian clock, while

GABA-6 was identified as *Lef1+* neuron whose function is unknown [76].

A recent study concluded that Ppp1r17 neuron population in the DMH regulates the aging

process in mice through activation of eNAMPT in the adipose tissues [56]. Interestingly, we

identified Glut-7 cluster as *Ppp1r17+* neuron which gene expression are highly modulated in the

transplantation group (Suppl. Fig. S10A–B). We also observed significant increase in the c-FOS

immunoreactivity in the DMH of the transplanted mice (Suppl. Fig. S10C) that is consistent with

previous finding and explained the increased plasma eNAMPT [56].

Loss of cell-to-cell communication is one of the hallmarks of aging. Using known ligand–

receptor interactions, it is possible to infer information flow from one cell type to another. A cell-

to-cell communication analysis using CellChat [77] showed that pericyte VEGF signaling was

restored by htNSC transplantation (Fig. 6K). Downregulation of the VEGF signal is a major

underlying cause of age-dependent blood brain barrier (BBB) dysfunction [78]. MCH is an orexic

neurohormone that is mainly secreted by GABAergic neurons in the hypothalamus [79]. Consistent

with the increased body energy metabolism that we saw in the grafted mice, GABAergic

neuronal output of MCH (*Pmch*) and neuronal communication with glial cells were diminished

in the grafted hypothalami (Fig. 6L). Additionally, alterations in non-canonical WNT and GDF

signal–mediated cell–cell communications were also manifested following htNSC

transplantation.

Neuro-inflammation in aged hypothalami corrected by intra-hypothalamic htNSC

implantation

Next, we validated the snRNA-seq data indicating that hypothalamic inflammation subsided

following htNSC transplantation by examining protein levels with immunofluorescence staining.

Consistent with the snRNA-seq data, the abundance of IBA1+ microglia and GFAP+ astrocytes

in the hypothalamus were not altered by htNSC transplantation (Suppl. Fig. S11A–B). No

morphologic differences in microglia or astrocytes were found between groups (Suppl. Fig.

S11A–B). However, a marked reduction of Cd68+ cells, microglial TNF-α, Casp1, and Il-1β

expression in the treatment group were confirmed both in the snRNA-seq and in

immunohistochemical analysis (Fig. 7A–D). More importantly, the number of phosphorylated

p65 (activated p65) positive cells was also greatly reduced in the transplanted group, as well as

its co-expression in microglia and astrocyte (Fig. 7E–F). Interestingly, the reduction of

hypothalamic p65 was completely reversed by NAMPT inhibitor (Suppl. Fig. S12). Chronic

hypothalamus inflammation has been identified as both cause and consequence of systemic

metabolic dysregulation [80-82]. Unexpectedly, propanolol treatment dramatically reduce p65

expression in both control and transplantation group (Suppl. Fig. S12), possibly through direct

inhibition of microglial β-AR by propranolol [83-85].

DISCUSSION

In this study, we demonstrated that htNSCs derived from hPSCs exhibit health-span-promoting

benefits when transplanted into aged hypothalamus. The correction of age-associated changes

was represented by increased energy metabolism and glucose tolerance, with upregulated

browning/thermogenic/catabolic, and sympathetic adrenergic gene and protein expression. In

addition to changes in their body/energy metabolism and physiology, the mice grafted with

human htNSCs in this study and mouse htNSCs in the study of Zhang et al. [10] showed improved

memory and cognitive functioning. Considering that the hypothalamus is the master coordinator

of a myriad of homeostatic functions that are deregulated during aging, the anti-aging effects

observed in this study are likely to reflect the central action of the grafted human htNSCs (or

their differentiated progenies) in rejuvenating host hypothalamus cells or environment to allow

systemic body metabolism and brain function to be more efficiently regulated. We identified

hypothalamus-SNS-eNAMPT axis that is required for the anti-aging effect.

Mechanistically, intra-hypothalamic htNSC transplantation increased neuronal activity in the

DMH, which activates the SNS to induce the synthesis and release of the anti-aging factor

eNAMPT from adipose tissues. The elevated levels of eNAMPT in systemic circulation are

believed to play a crucial role in enhancing body and lipid metabolism in the grafted mice [56]. In

addition, eNAMPT can be transported to the hypothalamus, where it promotes rejuvenation and

reduces inflammation through NAD+ production and SIRT1 activation [54,55]. Conversely, products

such as lipids and pro-inflammatory cytokines, increased by dysregulated peripheral energy

metabolism, cause chronic inflammation in the hypothalamus. [86,87]. Consequently, the

physiologic effects of htNSC-transplantation were abolished upon SNS or eNAMPT inhibition.

These findings support the concept of 'Inter-organ communication in aging' and highlight a feed-

forward mechanism between improved systemic metabolism and hypothalamic function in

transplanted mice [88]. In conclusion, this study has shown that implanting human htNSCs derived

from hPSC-organoids holds therapeutic promise for promoting healthy metabolic network which

may also slow aging process.

Although this study shed some insight on the potential of htNSC replacement therapy, this study

has several limitations. In this study we only utilized male mice. Sexual dimorphism of aging is a

known phenomenon that we could not cover in this study. Secondly, we also used only single

strain of mice; therefore, it is unclear although highly probable that the result observed here can

be extrapolated into another genetic background or other species.

MATERIAL AND METHODS

Reagents and resources

Detailed information of reagents and resources used in this paper can be found in Supplemental

Table S1.

Cell culture

hESCs (H9) and hiPSCs (CMC-hIPSC-011, CS-001) were cultured on Matrigel-coated plates in

feeder-free conditions in mTeSR-1 medium (Stemcell Technologies, Vancouver, Canada). As

schematized in Fig. 1A, human htNSCs were prepared by adopting the organoid-based NSC

preparation method that was recently published by our group [28]. First, to generate human

hypothalamus-like organoids , hESCs or hiPSCs were seeded at a density of 5,000 cells/well in a

low-attachment, 96-well, round-bottom plate in neural induction medium (N2:Neurobasal, 1:1,

Thermo Fisher Scientific, Waltham, Massachusetts) containing 2% B27 without vitamin A

(Thermo Fisher Scientific), 200 µM ascorbic acid (MilliporeSigma, Burlington, Massachusetts),

and 25 mg/L insulin, (Thermo Fisher Scientific). The ROCK inhibitor Y27632 (20 µM,

MilliporeSigma) was added for 24 hours after cell seeding. Hypothalamus patterning of the

differentiating hESCs/hiPSCs was induced by adding SB431542 (10 µM, Tocris, Bristol, UK)

and noggin (200 ng/ml, Peprotech, Cranbury, New Jersey) for days 0–7 and adding sonic

hedgehog (SHH, 100 ng/mL, Peprotech) and purmorphamine (2 µM, MilliporeSigma) for days

1–13. The medium was replenished every other day. Floating spheres (organoids) were

transferred into low attachment, 6-well plates (16 spheres/well) on day 13 in neural expansion

medium (N2 containing 16 µg/L insulin, 200 µM ascorbic acid, 100 U/ml penicillin G, and 100

µg/ml streptomycin) supplemented with bFGF (20 ng/mL, Peprotech), a mitogen for NSC, and

cultured on an orbital shaker (80 rpm) until day 20. The organoids enriched with htNSCs were

chopped, dissociated with Acutase (Stemcell Technologies), and plated on

poly-L-ornithine/fibronectin-coated plates in bFGF-supplemented neural expansion medium. The

htNSCs were expanded with cell passages at 5–7-day intervals. Y27632 was added for only 1

day after each cell passage. Terminal differentiation of the htNSCs was induced in N2 medium

supplemented with 200 µM ascorbic acid, 10 ng/mL BDNF (Peprotech), 10 ng/mL GDNF

(Peprotech), 1 µM db-cAMP (MilliporeSigma), and penicillin/streptomycin. As a control, NSCs

with ventral midbrain identity (ventral midbrain-type NSCs; VM-NSCs) were prepared as

described previously [28].

Animal care and study approval

Middle-aged C57BL/6J male mice (11–12 months) were purchased from the Korean Basic

Science Institute (Gwangju, South Korea) or Jackson Laboratory (Yokohama, Japan) and young-

aged C57BL/6J male mice (2–3 months) were purchased from Korean Basic Science Institute

(Gwangju, South Korea). All mice housed in a specific pathogen-free barrier facility with a

temperature of $23 \pm 1°C$, a humidity of $50 \pm 10\%$, a 12 h light/dark cycle, and free access to food

(standard chow, 5053 PicoLab Rodent Diet 20, Lab Diet, Saint Louis, Missouri) and water. All

procedures for the animal experiments were approved by the Hanyang College of Medicine

Institutional Animal Care and Use Committee (IACUC) (approval number, 2018-0217A and

2020-0142A) and the Korean Institute of Science and Technology (KIST) IACUC and

Institutional Biosafety Committee (Approval number, KIST-2019-048).

Transplantation procedures

For cell transplantation, 12-month-old mice received stereotaxic surgery under anesthesia

induced by a mixture of Zoletil (37.5 mg/kg) and Rompun (5.83 mg/kg) in isotonic saline.

Human htNSC dissociated into single-cells using Accutase and subsequently suspended in 1µl of

N2 medium (1×10^5 cells/µL) were injected into the MBH (AP: –1.5 mm, ML: ±0.4 mm, DV: –

5.6 mm) at a rate of 0.25 µL/minute for 2 minutes using 26G needle, and the injector was then

left in place for an additional 5 minutes before removal. A sham operation with an N2 medium

injection was carried out for the control group. The animals received daily injections of

cyclosporine A (10 mg/kg, i.p.) starting 1 day before the grafting and continuing for 3 weeks.

Behavioral analyses

All behavioral tests were performed in the behavioral testing room. An ANY-maze video-

tracking system (Stoelting, Wood Dale, Illinois) equipped with a digital camera connected to a

computer was used. The following behavior tests were conducted in compliance with the

ARRIVE guidelines. Detailed analyses in the Supplemental Methods.

Measurement of systemic body metabolism

Assays for body composition, metabolic cage experiment, glucose tolerance test, body

temperature, bone density measurement, and bone histomorphometry are described in the

Supplemental Methods.

Histological analysis

Mice were transcardially perfused with 30 mL of PBS (pH 7.4), followed by 30 mL of 4%

paraformaldehyde (PFA) in PBS. Their brains were dissected out and post-fixed with 4% PFA

overnight at 4°C and then transferred to 30% sucrose for 7 days for cryoprotection. The

cryoprotected brains were frozen in TissueTek OCT compound (Sakura Finetek, Torrance,

California) and sliced into 30 µm sections using a cryostat (Leica, Nussloch, Germany). For

peripheral tissues, the WAT, BAT, livers, and femurs were dissected from the mice without

fixation and divided into two halves for histochemical and Western blot analyses. The tissues for

histochemical staining were fixed overnight at 4°C in 4% PFA in PBS, paraffin embedded, and

sectioned into 4 µm slices that were stained with hematoxylin and eosin (H&E) following the

standard procedure.

Determination of protein and mRNA expressions

Protein expressions in the htNSC cultures and mouse hypothalamic tissues were assessed using

immunostaining and Western blot analyses. Serum levels of cytokines and leptin were also determined

using a Proteome Profiler Mouse XL Cytokine Array and a Mouse/Rat Leptin Quantikine ELISA

Kit (R&D Systems, Minneapolis, Minnesota), respectively. mRNA expressions were assessed

using quantitative PCR, global, and single cell transcriptomic analyses. Detailed methods are

described in the Supplemental Methods.

Statistical analysis

No statistical method was used to predetermine sample sizes, which were chosen with adequate

statistical power based on literature and experience. Data shown from representative experiments

were repeated with similar results in at least 3 independent experiments, unless otherwise

indicated by sample size. Homogeneity of variance was tested using F-test for two groups, and

Brown-Forsythe test for more than two groups. Data were analyzed using two-tailed unpaired

Student's t-testing, one-way analysis of variance (ANOVA), or Mann-Whitney U-testing.

Welch's correction was applied for datasets with unequal variance. Tukey post-hoc test was used

for one-way ordinary ANOVA test while unpaired t-testing was used for one-way Welch

ANOVA test. All multiple testing p-values were adjusted using Benjamini and Hochberg

correction. For behavioral analyses, all data points below or above mean $\pm$ 1.645 * SD

(corresponding to 5% in the normal distribution) are considered as outliers and were removed

from statistical analyses. Additional analyses using linear mixed modeling and Monte-Carlo

permutation (n = 10,000) were applied in behavior datasets. In all cases, results were considered

statistically significant when $p < 0.05$. Error bars show the mean $\pm$ standard error of the mean

(SEM) or the median, and p values are represented as: $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and

$****p < 0.0001$. Statistical analyses were performed with Prism software version 8 (GraphPad

Software) or R.

Data availability

The datasets produced in this study are available in the following databases:

hPSCs-derived htNSC bulk RNA-seq: GSE214877

hPSCs-derived htNSC organoid scRNA-seq: GSE215174

htNSC-transplanted aged mouse hypothalamus snRNA-seq: GSE215174

Values for each data point presented in this paper can be found in the Supporting Data Values

XLS file.

Acknowledgements

Fundings: This work was supported by grants 2020M3A9D8039920, KFRM 23A0104L1,

2017R1A5A2015395, 2022R1I1A1A01071091, and 2013M3A9D5072550 funded by the

Author contribution: S.H.L. and M.S.K. supervised the project. S.H.L., M.S.K., Y.A.S., Y.L.,

and K.P. designed experiments, and drafted and revised the manuscript. Y.A.S., Y.L., K.P., S.H.,

J.K., M.J.S., N-K.L., K-S.P., T.H., S.C., K.W.K., D.J.Y., W-Y.P., K.Y.H., S.G.Y., I.Y.K., and

J.K.S. performed experiments. Y.A.S., Y.L., and K.P. analyzed and interpreted data, and

performed statistical analyses. S.H.L., M.S.K., Y.A.S., J.K.S., T.Y.L., and M.S.K. acquired

funding.

Declaration of interests: S.H.L. and Y.A.S. are inventors on a patent related to the

differentiation of neural stem cells from pluripotent stem cells. The remaining authors declare no

competing interest.

References

1   Noble, E. E. *et al.* Hypothalamus-hippocampus circuitry regulates impulsivity via
melanin-concentrating hormone. *Nat Commun* **10**, 4923 (2019).
https://doi.org:10.1038/s41467-019-12895-y
2   Zhou, W. *et al.* Loss of function of NCOR1 and NCOR2 impairs memory through a
novel GABAergic hypothalamus-CA3 projection. *Nat Neurosci* **22**, 205-217 (2019).
https://doi.org:10.1038/s41593-018-0311-1
3   Chen, S. *et al.* A hypothalamic novelty signal modulates hippocampal memory. *Nature*
586, 270-274 (2020). https://doi.org:10.1038/s41586-020-2771-1
4   Li, Y. D. *et al.* Hypothalamic modulation of adult hippocampal neurogenesis in mice
confers activity-dependent regulation of memory and anxiety-like behavior. *Nat Neurosci*
25, 630-645 (2022). https://doi.org:10.1038/s41593-022-01065-x
5   Beckmann, N. D. *et al.* Multiscale causal networks identify VGF as a key regulator of
Alzheimer's disease. *Nat Commun* **11**, 3942 (2020). https://doi.org:10.1038/s41467-020-
17405-z
6   Ramirez, S. *et al.* Hypothalamic pregnenolone mediates recognition memory in the
context of metabolic disorders. *Cell Metab* **34**, 269-284 e269 (2022).
https://doi.org:10.1016/j.cmet.2021.12.023
7   Reichenbach, A. *et al.* Metabolic sensing in AgRP neurons integrates homeostatic state
with dopamine signalling in the striatum. *Elife* **11** (2022).
https://doi.org:10.7554/eLife.72668
8   Paul, A., Chaker, Z. & Doetsch, F. Hypothalamic regulation of regionally distinct adult
neural stem cells and neurogenesis. *Science* **356**, 1383-1386 (2017).
https://doi.org:10.1126/science.aal3839

629 9 Kim, K. & Choe, H. K. Role of hypothalamus in aging and its underlying cellular
630  mechanisms. *Mech Ageing Dev* **177**, 74-79 (2019).
631  https://doi.org:10.1016/j.mad.2018.04.008
632 10 Zhang, Y. *et al.* Hypothalamic stem cells control ageing speed partly through exosomal
633  miRNAs. *Nature* **548**, 52-57 (2017). https://doi.org:10.1038/nature23282
634 11 Garcia-Caceres, C. *et al.* Astrocytic Insulin Signaling Couples Brain Glucose Uptake
635  with Nutrient Availability. *Cell* **166**, 867-880 (2016).
636  https://doi.org:10.1016/j.cell.2016.07.028
637 12 Chowen, J. A. *et al.* The role of astrocytes in the hypothalamic response and adaptation to
638  metabolic signals. *Prog Neurobiol* **144**, 68-87 (2016).
639  https://doi.org:10.1016/j.pneurobio.2016.03.001
640 13 Garcia-Caceres, C. *et al.* Role of astrocytes, microglia, and tanycytes in brain control of
641  systemic metabolism. *Nat Neurosci* **22**, 7-14 (2019). https://doi.org:10.1038/s41593-018-
642  0286-y
643 14 Ameroso, D. *et al.* Astrocytic BDNF signaling within the ventromedial hypothalamus
644  regulates energy homeostasis. *Nat Metab* (2022). https://doi.org:10.1038/s42255-022-
645  00566-0
646 15 Li, J., Tang, Y. & Cai, D. IKKbeta/NF-kappaB disrupts adult hypothalamic neural stem
647  cells to mediate a neurodegenerative mechanism of dietary obesity and pre-diabetes. *Nat
648  Cell Biol* **14**, 999-1012 (2012). https://doi.org:10.1038/ncb2562
649 16 Lee, D. A. *et al.* Tanycytes of the hypothalamic median eminence form a diet-responsive
650  neurogenic niche. *Nat Neurosci* **15**, 700-702 (2012). https://doi.org:10.1038/nn.3079
651 17 Pierce, A. A. & Xu, A. W. De novo neurogenesis in adult hypothalamus as a
652  compensatory mechanism to regulate energy balance. *J Neurosci* **30**, 723-730 (2010).
653  https://doi.org:10.1523/JNEUROSCI.2479-09.2010
654 18 Kokoeva, M. V., Yin, H. & Flier, J. S. Neurogenesis in the hypothalamus of adult mice:
655  potential role in energy balance. *Science* **310**, 679-683 (2005).
656  https://doi.org:10.1126/science.1115360
657 19 Balland, E. *et al.* Hypothalamic tanycytes are an ERK-gated conduit for leptin into the
658  brain. *Cell Metab* **19**, 293-301 (2014). https://doi.org:10.1016/j.cmet.2013.12.015
659 20 Langlet, F. Tanycytes: a gateway to the metabolic hypothalamus. *J Neuroendocrinol* **26**,
660  753-760 (2014). https://doi.org:10.1111/jne.12191
661 21 Bolborea, M., Pollatzek, E., Benford, H., Sotelo-Hitschfeld, T. & Dale, N. Hypothalamic
662  tanycytes generate acute hyperphagia through activation of the arcuate neuronal network.
663  *Proc Natl Acad Sci U S A* **117**, 14473-14481 (2020).
664  https://doi.org:10.1073/pnas.1919887117
665 22 Muller-Fielitz, H. *et al.* Tanycytes control the hormonal output of the hypothalamic-
666  pituitary-thyroid axis. *Nat Commun* **8**, 484 (2017). https://doi.org:10.1038/s41467-017-
667  00604-6
668 23 Yoo, S. *et al.* Control of neurogenic competence in mammalian hypothalamic tanycytes.
669  *Sci Adv* **7** (2021). https://doi.org:10.1126/sciadv.abg3777
670 24 Yoo, S. & Blackshaw, S. Regulation and function of neurogenesis in the adult
671  mammalian hypothalamus. *Prog Neurobiol* **170**, 53-66 (2018).
672  https://doi.org:10.1016/j.pneurobio.2018.04.001

25   Wang, L. *et al.* Differentiation of hypothalamic-like neurons from human pluripotent
stem cells. *J Clin Invest* **125**, 796-808 (2015). https://doi.org:10.1172/JCI79220
26   Merkle, F. T. *et al.* Generation of neuropeptidergic hypothalamic neurons from human
pluripotent stem cells. *Development* **142**, 633-643 (2015).
https://doi.org:10.1242/dev.117978
27   Ogawa, K. *et al.* Vasopressin-secreting neurons derived from human embryonic stem
cells through specific induction of dorsal hypothalamic progenitors. *Sci Rep* **8**, 3615
(2018). https://doi.org:10.1038/s41598-018-22053-x
28   Kim, S. W. *et al.* Neural stem cells derived from human midbrain organoids as a stable
source for treating Parkinson's disease: Midbrain organoid-NSCs (Og-NSC) as a stable
source for PD treatment. *Prog Neurobiol* **204**, 102086 (2021).
https://doi.org:10.1016/j.pneurobio.2021.102086
29   Rajamani, U. *et al.* Super-Obese Patient-Derived iPSC Hypothalamic Neurons Exhibit
Obesogenic Signatures and Hormone Responses. *Cell Stem Cell* **22**, 698-712 e699
(2018). https://doi.org:10.1016/j.stem.2018.03.009
30   Artegiani, B. *et al.* A Single-Cell RNA Sequencing Study Reveals Cellular and
Molecular Dynamics of the Hippocampal Neurogenic Niche. *Cell Rep* **21**, 3271-3284
(2017). https://doi.org:10.1016/j.celrep.2017.11.050
31   Hochgerner, H., Zeisel, A., Lonnerberg, P. & Linnarsson, S. Conserved properties of
dentate gyrus neurogenesis across postnatal development revealed by single-cell RNA
sequencing. *Nat Neurosci* **21**, 290-299 (2018). https://doi.org:10.1038/s41593-017-0056-
2
32   Zeisel, A. *et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**, 999-
1014 e1022 (2018). https://doi.org:10.1016/j.cell.2018.06.021
33   Fleck, J. S. *et al.* Resolving organoid brain region identities by mapping single-cell
genomic data to reference atlases. *Cell Stem Cell* **28**, 1148-1159 e1148 (2021).
https://doi.org:10.1016/j.stem.2021.02.015
34   Herb, B. R. *et al.* Single-cell genomics reveals region-specific developmental trajectories
underlying neuronal diversity in the human hypothalamus. *bioRxiv*,
2021.2007.2020.453090 (2022). https://doi.org:10.1101/2021.07.20.453090
35   Quinn, J. P., Kandigian, S. E., Trombetta, B. A., Arnold, S. E. & Carlyle, B. C. VGF as a
biomarker and therapeutic target in neurodegenerative and psychiatric diseases. *Brain*
Commun **3**, fcab261 (2021). https://doi.org:10.1093/braincomms/fcab261
36   Lewis, J. E., Brameld, J. M. & Jethwa, P. H. Neuroendocrine Role for VGF. *Front*
Endocrinol (Lausanne) **6**, 3 (2015). https://doi.org:10.3389/fendo.2015.00003
37   Lepko, T. *et al.* Choroid plexus-derived miR-204 regulates the number of quiescent
neural stem cells in the adult brain. *EMBO J* **38**, e100481 (2019).
https://doi.org:10.15252/embj.2018100481
38   Hasel, P., Rose, I. V. L., Sadick, J. S., Kim, R. D. & Liddelow, S. A. Neuroinflammatory
astrocyte subtypes in the mouse brain. *Nat Neurosci* **24**, 1475-1487 (2021).
https://doi.org:10.1038/s41593-021-00905-6
39   Mizrak, D. *et al.* Single-Cell Analysis of Regional Differences in Adult V-SVZ Neural
Stem Cell Lineages. *Cell Rep* **26**, 394-406 e395 (2019).
https://doi.org:10.1016/j.celrep.2018.12.044

40   Romanov, R. A. *et al.* Molecular design of hypothalamus development. *Nature* **582**, 246-
252 (2020). https://doi.org:10.1038/s41586-020-2266-0
41   Suda, Y. *et al.* Normal aging induces PD-1-enriched exhausted microglia and A1-like
reactive astrocytes in the hypothalamus. *Biochem Biophys Res Commun* **541**, 22-29
(2021). https://doi.org:10.1016/j.bbrc.2020.12.086
42   Gupta, D. & Morley, J. E. Hypothalamic-pituitary-adrenal (HPA) axis and aging. *Compr*
Physiol **4**, 1495-1510 (2014). https://doi.org:10.1002/cphy.c130049
43   Cuevas-Ramos, D., Mehta, R. & Aguilar-Salinas, C. A. Fibroblast Growth Factor 21 and
Browning of White Adipose Tissue. *Front Physiol* **10**, 37 (2019).
https://doi.org:10.3389/fphys.2019.00037
44   Flippo, K. H. & Potthoff, M. J. Metabolic Messengers: FGF21. *Nat Metab* **3**, 309-317
(2021). https://doi.org:10.1038/s42255-021-00354-2
45   Mullican, S. E. & Rangwala, S. M. Uniting GDF15 and GFRAL: Therapeutic
Opportunities in Obesity and Beyond. *Trends Endocrinol Metab* **29**, 560-570 (2018).
https://doi.org:10.1016/j.tem.2018.05.002
46   Tsai, V. W. W., Husaini, Y., Sainsbury, A., Brown, D. A. & Breit, S. N. The
MIC-1/GDF15-GFRAL Pathway in Energy Homeostasis: Implications for Obesity,
Cachexia, and Other Associated Diseases. *Cell Metab* **28**, 353-368 (2018).
https://doi.org:10.1016/j.cmet.2018.07.018
47   Masand, R. *et al.* Proteome Imbalance of Mitochondrial Electron Transport Chain in
Brown Adipocytes Leads to Metabolic Benefits. *Cell Metab* **27**, 616-629 e614 (2018).
https://doi.org:10.1016/j.cmet.2018.01.018
48   Vernochet, C. *et al.* Adipose-specific deletion of TFAM increases mitochondrial
oxidation and protects mice against obesity and insulin resistance. *Cell Metab* **16**, 765-
776 (2012). https://doi.org:10.1016/j.cmet.2012.10.016
49   Solon-Biet, S. M. *et al.* The ratio of macronutrients, not caloric intake, dictates
cardiometabolic health, aging, and longevity in ad libitum-fed mice. *Cell Metab* **19**, 418-
430 (2014). https://doi.org:10.1016/j.cmet.2014.02.009
50   Tran, L. T. *et al.* Hypothalamic control of energy expenditure and thermogenesis. *Exp*
Mol Med **54**, 358-369 (2022). https://doi.org:10.1038/s12276-022-00741-z
51   Ostlund, R. E., Jr., Yang, J. W., Klein, S. & Gingerich, R. Relation between plasma leptin
concentration and body fat, gender, diet, age, and metabolic covariates. *J Clin Endocrinol*
Metab **81**, 3909-3913 (1996). https://doi.org:10.1210/jcem.81.11.8923837
52   Minocci, A. *et al.* Leptin plasma concentrations are dependent on body fat distribution in
obese patients. *Int J Obes Relat Metab Disord* **24**, 1139-1144 (2000).
https://doi.org:10.1038/sj.ijo.0801385
53   Elefteriou, F. *et al.* Serum leptin level is a regulator of bone mass. *Proc Natl Acad Sci U*
S A **101**, 3258-3263 (2004). https://doi.org:10.1073/pnas.0308744101
54   Yoon, M. J. *et al.* SIRT1-Mediated eNAMPT Secretion from Adipose Tissue Regulates
Hypothalamic NAD+ and Function in Mice. *Cell Metab* **21**, 706-717 (2015).
https://doi.org:10.1016/j.cmet.2015.04.002
55   Yoshida, M. *et al.* Extracellular Vesicle-Contained eNAMPT Delays Aging and Extends
Lifespan in Mice. *Cell Metab* **30**, 329-342.e325 (2019).
https://doi.org:10.1016/j.cmet.2019.05.015

Tokizane, K., Brace, C. S. & Imai, S. I. DMH(Ppp1r17) neurons regulate aging and lifespan in mice through hypothalamic-adipose inter-tissue communication. *Cell Metab* **36**, 377-392.e311 (2024). https://doi.org:10.1016/j.cmet.2023.12.011

Carneiro, L., Bernasconi, R., Bernini, A., Repond, C. & Pellerin, L. Elevation of hypothalamic ketone bodies induces a decrease in energy expenditures and an increase risk of metabolic disorder. *Mol Metab* **83**, 101926 (2024). https://doi.org:10.1016/j.molmet.2024.101926

Bendridi, N., Selmi, A., Balcerczyk, A. & Pirola, L. Ketone Bodies as Metabolites and Signalling Molecules at the Crossroad between Inflammation and Epigenetic Control of Cardiometabolic Disorders. *Int J Mol Sci* **23** (2022). https://doi.org:10.3390/ijms232314564

Lin, A. L., Zhang, W., Gao, X. & Watts, L. Caloric restriction increases ketone bodies metabolism and preserves blood flow in aging brain. *Neurobiol Aging* **36**, 2296-2303 (2015). https://doi.org:10.1016/j.neurobiolaging.2015.03.012

Edwards, C. *et al.* D-beta-hydroxybutyrate extends lifespan in C. elegans. *Aging (Albany NY)* **6**, 621-644 (2014). https://doi.org:10.18632/aging.100683

Wang, L., Chen, P. & Xiao, W. β-hydroxybutyrate as an Anti-Aging Metabolite. *Nutrients* **13** (2021). https://doi.org:10.3390/nu13103420

Idelevich, A. *et al.* Neuronal hypothalamic regulation of body metabolism and bone density is galanin dependent. *J Clin Invest* **128**, 2626-2641 (2018). https://doi.org:10.1172/JCI99350

Qualls-Creekmore, E. *et al.* Galanin-Expressing GABA Neurons in the Lateral Hypothalamus Modulate Food Reward and Noncompulsive Locomotion. *J Neurosci* **37**, 6053-6065 (2017). https://doi.org:10.1523/JNEUROSCI.0155-17.2017

Soreq, L. *et al.* Major Shifts in Glial Regional Identity Are a Transcriptional Hallmark of Human Brain Aging. *Cell Rep* **18**, 557-570 (2017). https://doi.org:10.1016/j.celrep.2016.12.011

Ximerakis, M. *et al.* Single-cell transcriptomic profiling of the aging mouse brain. *Nat Neurosci* **22**, 1696-1708 (2019). https://doi.org:10.1038/s41593-019-0491-3

Wang, W. *et al.* Transcriptome dynamics of hippocampal neurogenesis in macaques across the lifespan and aged humans. *Cell Res* **32**, 729-743 (2022). https://doi.org:10.1038/s41422-022-00678-y

Chen, Z. H. *et al.* Single-cell transcriptomic profiling of the hypothalamic median eminence during aging. *J Genet Genomics* **49**, 523-536 (2022). https://doi.org:10.1016/j.jgg.2022.01.001

Cai, D. & Khor, S. "Hypothalamic Microinflammation" Paradigm in Aging and Metabolic Diseases. *Cell Metab* **30**, 19-35 (2019). https://doi.org:10.1016/j.cmet.2019.05.021

Zhang, G. *et al.* Hypothalamic programming of systemic ageing involving IKK-beta, NF-kappaB and GnRH. *Nature* **497**, 211-216 (2013). https://doi.org:10.1038/nature12143

Liu, Y. *et al.* Hypothalamic Hnscr regulates glucose balance by mediating central inflammation and insulin signal. *Cell Prolif*, e13332 (2022). https://doi.org:10.1111/cpr.13332

71 Marschallinger, J. *et al.* Lipid-droplet-accumulating microglia represent a dysfunctional
and proinflammatory state in the aging brain. *Nat Neurosci* **23**, 194-208 (2020).
https://doi.org:10.1038/s41593-019-0566-1
72 Lopes, K. P. *et al.* Genetic analysis of the human microglial transcriptome across brain
regions, aging and disease pathologies. *Nat Genet* **54**, 4-17 (2022).
https://doi.org:10.1038/s41588-021-00976-y
73 Galatro, T. F. *et al.* Transcriptomic analysis of purified human cortical microglia reveals
age-associated changes. *Nat Neurosci* **20**, 1162-1171 (2017).
https://doi.org:10.1038/nn.4597
74 Rodriguez-Rodriguez, A. *et al.* Tanycytes and the Control of Thyrotropin-Releasing
Hormone Flux Into Portal Capillaries. *Front Endocrinol (Lausanne)* **10**, 401 (2019).
https://doi.org:10.3389/fendo.2019.00401
75 Daimon, C. M., Chirdon, P., Maudsley, S. & Martin, B. The role of Thyrotropin
Releasing Hormone in aging and neurodegenerative diseases. *Am J Alzheimers Dis*
(Columbia) **1** (2013). https://doi.org:10.7726/ajad.2013.1003
76 Chiba, A. & Honma, Y. Distribution of neuropeptide Y-like immunoreactivity in the
brain and hypophysis of the cloudy dogfish, Scyliorhinus torazame. *Cell Tissue Res* **268**,
453-461 (1992). https://doi.org:10.1007/BF00319152
77 Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat*
Commun **12**, 1088 (2021). https://doi.org:10.1038/s41467-021-21246-9
78 Wen, S. W. & Wong, C. H. Y. Aging- and vascular-related pathologies. *Microcirculation*
26, e12463 (2019). https://doi.org:10.1111/micc.12463
79 Jego, S. *et al.* Optogenetic identification of a rapid eye movement sleep modulatory
circuit in the hypothalamus. *Nat Neurosci* **16**, 1637-1643 (2013).
https://doi.org:10.1038/nn.3522
80 Zhang, Y., Reichel, J. M., Han, C., Zuniga-Hertz, J. P. & Cai, D. Astrocytic Process
Plasticity and IKKbeta/NF-kappaB in Central Control of Blood Glucose, Blood Pressure,
and Body Weight. *Cell Metab* **25**, 1091-1102 e1094 (2017).
https://doi.org:10.1016/j.cmet.2017.04.002
81 Dalvi, P. S. *et al.* High fat induces acute and chronic inflammation in the hypothalamus:
effect of high-fat diet, palmitate and TNF-alpha on appetite-regulating NPY neurons. *Int*
J Obes (Lond) **41**, 149-158 (2017). https://doi.org:10.1038/ijo.2016.183
82 Valdearcos, M., Xu, A. W. & Koliwad, S. K. Hypothalamic inflammation in the control
of metabolic function. *Annu Rev Physiol* **77**, 131-160 (2015).
https://doi.org:10.1146/annurev-physiol-021014-071656
83 Kota, D. J. *et al.* Propranolol and Mesenchymal Stromal Cells Combine to Treat
Traumatic Brain Injury. *Stem Cells Transl Med* **5**, 33-44 (2016).
https://doi.org:10.5966/sctm.2015-0065
84 Pilipovic, I., Stojic-Vukanic, Z., Prijic, I., Jasnic, N. & Leposavic, G. Propranolol
diminished severity of rat EAE by enhancing immunoregulatory/protective properties of
spinal cord microglia. *Neurobiol Dis* **134**, 104665 (2020).
https://doi.org:10.1016/j.nbd.2019.104665
85 Sugama, S. & Kakinuma, Y. Noradrenaline as a key neurotransmitter in modulating
microglial activation in stress response. *Neurochem Int* **143**, 104943 (2021).
https://doi.org:10.1016/j.neuint.2020.104943

86 Dutheil, S., Ota, K. T., Wohleb, E. S., Rasmussen, K. & Duman, R. S. High-fat diet
induced anxiety and anhedonia: impact on brain homeostasis and inflammation.
Neuropsychopharmacology **41**, 1874-1887 (2016).
87 Wang, X. *et al.* The effect of swimming exercise and diet on the hypothalamic
inflammation of ApoE-/-mice based on SIRT1-NF-κB-GnRH expression. *Aging (Albany*
NY) **12**, 11085 (2020).
88 Tokizane, K. & Imai, S. I. Inter-organ communication is a critical machinery to regulate
metabolism and aging. *Trends Endocrinol Metab* (2024).
https://doi.org:10.1016/j.tem.2024.11.013

Figure

Fig. 1. Derivation of hypothalamic neural stem cells (htNSCs) from hPSC-derived
hypothalamus-like organoids.

A. An overview of the protocol to yield htNSCs with proliferative and neuro-/gliogenic potential

from hPSC-derived organoid cultures.

B. mRNA expression of htNSC-specific marker genes (*NKX2-1, RAX, OTP, ISL1, SF1*) assessed

by qRT-PCR. The expression levels in the H9 hESC-derived htNSCs were compared with those

in undifferentiated hESCs and hESC-derived ventral midbrain-type neural stem cells (VM-

NSCs). n=3.[a,b]

C. Immunocytochemical detection of generic NSC (SOX2, NESTIN, KI67) and htNSC-specific

markers (NKX2-1, RAX, ISL1). Scale bars, 100 µm. n=3.[a]

D. Heatmap comparison of our H9 hESC-derived htNSC transcriptome with various human adult

brain regions in the GTEx database. HYP: hypothalamus; HIP: hippocampus; CTX: cortex; SN:

substantia nigra.

E. Terminal differentiation of htNSCs to yield hypothalamus-specific neuronal and astroglial

progeny. The cellular identities were evaluated on htNSC differentiation day 40 by the

expression of general neuronal (TUBB3, MAP2, RBFOX3, SYN1), hypothalamus neuron

subtype (α-MSH, NPY, SCGN, PMCH), and astroglial (GFAP, VIM, AQ4, A2B5) markers.

Scale bars, 100 µm.

F. Cells differentiated from htNSCs acquired the hypothalamus-specific functions of responding

to leptin and ghrelin ligands. Intracellular STAT3 and FOXO1 activation were evaluated in the

differentiated htNSC cultures before and after exogenous leptin and ghrelin treatments,

respectively. n=3.[a,b]

G. Uniform manifold approximation and projection (UMAP) plot of 6,394 single cells projected

into 13 cell clusters (indicated by color). NSCs, neural stem/progenitor cells; Int1, intermediate

progenitor cells 1, Int2, intermediate progenitor cells 2; iNPCs, interneuron progenitor cells;

APCs, astrocyte progenitor cells; Neu1, neurons 1; Neu2, neurons 2; Ast1, astrocytes 1; Ast2,

astrocytes 2; Ast3, astrocytes 3; RG, radial glia; PGG, proteoglycan-secreting glia; Tan,

tanycytes.

H–I. Violin (B) and feature (C) plots of representative cell-specific genes from the differentiated

cell clusters.

J. Hypothalamic brain identity of the cell clusters validated by voxel-based spatial brain mapping

onto the mouse brain at postnatal day 28 (from the Allen Brain Institute).

^aData shown are means ± SEM.

^b*p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001 by two-tailed Student's t testing.

Fig. 2. Age-associated decline in physical and cognitive activities is rescued by intra-
hypothalamic htNSC transplantation.

A. Experimental scheme for the animal study. hESC-derived htNSCs (P2) were dissociated into
single cells and transplanted into the medio-basal hypothalami of 12-month-old male mice.
Behavioral tests were performed 1.5 months and/or 4 months post-transplantation, and tissue
analyses were conducted 4 months post-surgery. Four rounds of transplantation experiments
were conducted using mice and donor human htNSCs that were independently prepared.

B. Overview images depict the human htNSC-derived graft localized in the mouse
hypothalamus but not in other brain regions. The htNSC-derived graft was identified using a
human mitochondria-specific antibody (huMt). CPu, caudate putamen; Hip, hippocampus; Hypo,
hypothalamus; SN, substansia nigra.

C. Identification of cell types derived from the grafted htNSCs using antibodies specific for
human-species (hNCAM, huMt), human-specific astrocytes (hGFAP), htNSCs (RAX), or
neurons (RBFOX3) at 2–4 weeks post-transplantation. Scale bars, 50 µm or 200 µm.
D–G. Physical activity, muscle coordination, strength, and endurance were estimated using the
open filed test (OFT, C), rotarod (D), grip (E), and treadmill performance (F) tests. n= 9-10
(young), 15-22 (control), 15-22 (transplanted).[a,b]
H–K. Memory and cognitive functions were evaluated using the Morris water maze (MWM,
G,H), Y-maze (I), and novel object recognition (NOR, J) tests 4 months post-surgery. n= 9
(young), 9-20 (control), 8-20 (transplanted).[a,b]
[a]Data shown are means ± SEM. The data presented in D-K represent the combined data from at
least two independent datasets.
[b]*$p < 0.05$; **$p < 0.01$; ****$p < 0.0001$ by one-way Welch analysis of variance.
918

Fig. 3. Intra-hypothalamic human htNSC transplantation improved body energy metabolism and tissue aging in aged mice.

A. The body weights of the control (vehicle-injected) and transplanted mice 1.5 months and 4 months post-transplantation (left) and representative photographs of the mice at 16 months old (right). n= 10 (young), 15 (control), 17 (transplanted).[a,b]

B. Daily food intake (normal chow diet) of the control and transplanted mice. n= 5 (control), 7 (transplanted).[a,b]

C. Measurement of x-ambulatory activity during the light and dark phases of the day at 1.5 months post-transplantation. n= 5 (young), 8 (control), 8 (transplanted).[a,b]

D. Energy expenditure measured in metabolic cages. n= 5 (young), 8 (control), 8 (transplanted).[a,b]

E. NMR-based measurement of the fat and lean masses and free fluid volumes in the vehicle-treated control (n= 5) and transplanted (n = 7) mice.[a,b]

F–H. Quantified lipid droplet areas (right) with representative H&E-stained tissue section

images (left) of the iWAT (F), eWAT (G), and iBAT (H) liver from the young (n = 10), control

(n= 10) and transplanted (n = 11) mice. Scale bars, 50 µm. [a,b]

I. Representative H&E-stained images of liver from the young (n = 10), control (n= 10) and

transplanted (n = 11) mice. Scale bars, 1 mm.

[a]Data shown are means ± SEM. The data presented in A, C-D, and F-I represent the combined

data from at least two independent datasets.

[b]*$p < 0.05$; **$p < 0.01$; ****$p < 0.0001$ by one-way analysis of variance or two-tailed Student's t

testing.

Fig. 4. Body energy and bone metabolism changes ameliorated in aged mice that received htNSC grafts.

A–C. Immunoblot-based quantification of the key regulator proteins for energy metabolism and thermogenesis (browning) in the liver (A), iWAT (B), and eWAT (C) of the control (n=5) and transplanted (n=7) mice. [a,b]

D. Rectal temperature. n= 10 (young), 16 (control), 17 (transplanted).[a,b]

E. qPCR-based quantification of gene expression for β1, 2, and 3 adrenoreceptors *(Adrb1, Adrb2,* and *Adrb3*), and lipolytic genes (Dio2 and Lipe). n= 5 (control), 7 (transplanted).[a,b]

F. Glucose tolerance test. Data represent blood glucose levels over the time points after glucose

injection (2 g/kg, i.p.) (left) and the integral of the glucose levels during the post-injection period

(right). n= 9 (young), 10 (control), 10 (transplanted).[a,b]

G. Representative μCT bone images of tibia (left) and the indices of BMD, BV/TV, Tb.Th., and

Tb.N. (right), which increased significantly in the transplanted mice. n= 5 (control), 7

(transplanted).[a,b]

H. Representative μCT bone images of femur (top) and the indices of BMD, BV/TV, Tb.Th., and

Tb.N. (below), which increased significantly in the transplanted mice. n= 5.[a,b]

I. Representative bone histomorphometry (H&E, Tartrate-resistant acid phosphatase (TRAP),

and Alkaline phosphatase (ALP) staining) images of the femurs with quantification of relative

intensity of TRAP-positive osteoclasts and ALP-positive osteoblasts. n= 5 for TRAP staining, 3

for ALP staining. Scale bar, 800 µm (H&E staining) or 100 µm (TRAP and ALP staining). [a,b]

[a]Data shown are means ± SEM. The data presented in D and F represent the combined data from

at least two independent datasets.

[b]$*p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$ by one-way analysis of variance or two-

tailed Student's t testing.

Fig. 5. NAMPT and beta-adrenergic inhibitors reversed the anti-aging effects of the hNSC transplantation.

A. ELISA-based determination of serum NAMPT levels in the transplanted and control mice. n= 4. [a,b]

B. mRNA expression of *Nampt* in iWAT and eWAT tissues assessed by qRT-PCR. n= 5 (control), 7 (transplanted). [a,b]

C. Experimental scheme for inhibitor treatment in control and transplanted mice. Treatment

began two weeks post-transplantation and continued for 2.5 months. NAMPT inhibitor (FK-866,

10 mg/kg) was administered intraperitoneally once a day. Beta-adrenergic blocker (propranolol,

0.5 g/L) was administered via drinking water. All analyses were conducted after 2.5 months

inhibitor treatment.

D–E. Bodyweight measurements (D) and NMR-based measurement of the fat and lean masses.

n= 4. [a,b]

F-G. Muscle strength and cognitive function assessment by grip test and Y-maze test,

respectively. n= 4. [a,b]

H. Fasting blood ketone levels. n= 4. [a,b]

I. Rectal temperature. n= 4. [a,b]

J. Energy expenditure measured in metabolic cages. n= 4. [a,b]

K. Quantified lipid droplet areas (right) with representative H&E-stained tissue section images

(left) of the iWAT. n= 4. [a,b]

[a]Data shown are means ± SEM.

[b]$*p < 0.05$; $**p < 0.01$; $***p < 0.001$; $****p < 0.0001$ by one-way analysis of variance or two-

tailed Student's t testing.

Fig. 6. Single nucleus RNA-seq analysis of transplanted and control hypothalami.

A. Schematic of the snRNA-seq experiment.

B–C. UMAP plot of 12,161 nuclei filtered and processed from control and transplanted

hypothalami (B) and feature plot showing representative cluster transcripts (C).

D–E. UMAP plots of the GABA-neuron subcluster (D) and Glut-neuron subcluster (E).

F. Heatmap showing the top 25 signature genes among the clusters and the two most

representative GO results.

G. Stacked bar plot showing the composition of each cell type in the dataset for each group.

H. Bar plot showing the number of upregulated (orange) and downregulated (blue) transcripts in

all cell types for the transplanted group.

I. GO (biological process) enrichment analysis of DEGs from the non-neuronal clusters.

J. Violin plot of module scores for microglial nuclei in gene sets obtained from various studies.

The statistical analysis used the Mann-Whitney test.

K. Subset of a heatmap of the cell-to-cell interaction network showing the overall (ingoing and

outgoing) signal strength of VEGF, PMCH, ncWNT, and GDF signaling.

L. Circle plots of an inferred signaling network of PMCH, ncWNT, and GDF signaling.

1011

Fig. 7. Neuro-inflammation in aged hypothalami subsided following htNSC
transplantation.

A–B. Hypothalamic inflammation, as estimated by the snRNA-seq analysis of inflammatory
gene expression (*Cd68, Tnfrsf1b, Tnfaip1, Tank, Traf6,* and *Irf5*) in the microglial clusters (C)
and the immunohistochemical analysis for the percentage of CD68+/DAPI+ and TNFA+/IBA1+
cells (D). Scale bars, 100 µm. n = 3 (control), 3 (transplanted) in (C)[c,d] and n = 4 (control), 5
(transplanted) in (D)[a,b].

C–D. SnRNA-seq- and immunohistochemistry-based determination of NLRP3-inflammasome
activation in the aged control and grafted hypothalami. Scale bars, 100 µm. n = 3 (control), 3
(transplanted) in (E)[c,d] and n = 4 (control), 5 (transplanted) in (F)[a,b].

E–F. NF-κB signal activation in the aged control and grafted hypothalami, as assessed by

snRNA-seq (A) and immunohistochemical (B) analyses. Violin plot of *Nfkb1* in microglial and

astrocyte clusters and *Rel, Relb,* and *Ikbkg* in microglia from the snRNA-seq data (A). The

percentage of p-P65+ in the total (DAPI+), microglial (IBA1+), and astrocytic (GFAP+) cell

populations (B). Scale bars, 100 µm. n = 3 (control), 3 (transplanted) in (A)[c,d] and n = 4 (control),

5 (transplanted) in (B)[a,b].

[a]Data shown are means ± SEM.

[b]*p < 0.05; ***p < 0.001; ****p < 0.0001 by two-tailed Student's t testing.

[c]Data shown are medians.

[d]*p < 0.05; **p < 0.01; ***p < 0.001; ****p < 0.0001 by Mann-Whitney U testing.

1032