Mechanism-based design of DNA-nanoparticle motor with high speed and processivity comparable to motor proteins

Takanori Harashima[1,2], Akihiro Otomo[1,2], and Ryota Iino[1,2]

[1]Institute for Molecular Science, National Institutes of Natural Sciences, Okazaki, Aichi 444-8787, Japan

[2]Graduate Institute for Advanced Studies, SOKENDAI, Hayama, Kanagawa 240-0193, Japan

Keywords: DNA, synthetic molecular motors, DNA nanotechnology, gold-nanoparticles, single-particle tracking

Abstract

DNA-nanoparticle motor is a burnt-bridge Brownian ratchet moving on RNA-modified surface

driven by Ribonuclease H (RNase H), and one of the fastest nanoscale artificial motors. However,

its speed is still much lower than those of motor proteins. Here we resolve elementary processes

of motion and reveal long pauses caused by slow RNase H binding are the bottleneck. As RNase

H concentration ([RNase H]) increases, pause lengths shorten from ~100 s to ~0.1 s, while step

sizes are constant (~20 nm). At high [RNase H], speed reaches ~100 nm s^{-1}, however, processivity,

run-length, and unidirectionality largely decrease. A geometry-based kinetic simulation reveals

switching of bottleneck from RNase H binding to DNA/RNA hybridization at high [RNase H],

and trade-off mechanism between speed and other performances. A mechanism-based newly-

designed motor with 3.8-times larger DNA/RNA hybridization rate simultaneously achieves 30

nm s^{-1} speed, 200 processivity, and 3 μm run-length comparable to motor proteins.

1 Introduction

Engineering artificial molecular motors with performances comparable to motor proteins is an

important challenge to understand their operational and design principles and to implement them

in biological and abiotic systems[1-5]. DNA is not only the fundamental molecule of living organisms

but also a versatile building block used to engineer a variety of artificial motors such as DNA

walkers and spiders[6-8]. These artificial motors have flexibility in design, but exhibit speed of only

0.1 nm s^{-1} at best[6,8-10], and are much slower than motor proteins that exhibit speed of 10–1000 nm

s^{-1} [11-14]. Furthermore, run-length is limited to the size of complementary DNA/RNA tracks,

typically tens to hundreds of nanometers.

Recent efforts have focused on overcoming these drawbacks of DNA-based artificial

motors. Choi's group used RNA-modified carbon nanotube as a track for DNAzyme-based motor,

and achieved run-length over 3 µm with the maximum speed of 0.1 nm s^{-1} [10,15]. Salaita's group

developed a rolling motor which consists of DNA-modified microparticle, RNA-modified

substrate, and Ribonuclease H (RNase H)[16,17]. In this motor, DNAs on particle and complementary

RNAs on planar substrate surface form multivalent DNA/RNA hybridizations. Then, the motion

is initiated by addition of RNase H which hydrolyzes RNA in the DNA/RNA duplex, and multiple

reaction cycles occur in parallel at the particle/substrate interface. The DNA-microparticle motor

cannot go back to the RNA cleaved sites, shows self-avoiding super-diffusion, and operates as a

burnt-bride Brownian ratchet (BBR). The motor achieved speed of 30 nm s^{-1} and run-length of

300 µm with the help of gravity on the micron-sized particle[16].

Salaita's group also developed DNA-modified gold nanoparticle (AuNP) motors[18]. The

nanoscale motors have a potential to achieve higher speed than micron-sized motors due to fast

diffusion[19,20], although its small mass can result in stochastic detachment from the substrate surface.

However, in the previous study of a 50-nm DNA-AuNP motor, the average speed was only 3 nm s^{-1} and lower than those of DNA-microparticle motors and nanoscale motor proteins, although transient 50 nm s^{-1} bursts were observed[18]. This result suggests that one or multiple elementary processes become the bottleneck of motion of the DNA-AuNP motor.

Motor proteins simultaneously realize the high speed and long run-length (high processivity). For example, ATP-driven kinesin-1 moves unidirectionally along microtubule with maximum speed of $\sim$700 nm s^{-1} and run-length longer than 1 μm (processivity higher than 125)[12,21]. In addition to the fast diffusion, high speed motion of kinesin-1 relies on fast chemical processes including ATP binding, phosphate bond cleavage, and product release with millisecond time scales [21], and long run-length (high processivity) relies on the highly-coordinated chemo-mechanical coupling where two motor domains control the timings of chemical and mechanical processes[22]. For the motor proteins, the chemo-mechanical coupling mechanisms have been elucidated by single-molecule techniques which resolve the elementary process of the motion[21,23,24].

In this study, we applied high-speed/high-precision single-particle tracking and resolved the pauses and steps in the motion of a 100-nm DNA-AuNP motor. We found that the bottleneck of motion is the long pauses caused by slow RNase H binding at nanomolar RNase H concentration ([RNase H]) used in the previous study[18]. As [RNase H] increased, the pause lengths shortened from $\sim$100 to $\sim$0.1 s, while the step sizes ($\sim$20 nm) were constant. At micromolar [RNase H], the speed reached $\sim$100 nm s^{-1}, however, processivity, run-length, and unidirectionality largely decreased.

To understand the mechanisms of motion and trade-off between speed and other performances, we also developed a geometry-based kinetic simulation which reproduces experiment quantitatively. The simulation revealed that pauses are caused by multiple DNA/RNA

hybridizations between DNA-AuNP and RNA-substrate and steps occur when the number of

hybridizations decreases by the progress of RNA hydrolysis. The simulation also revealed that as

[RNase H] increases, the bottleneck switches from RNase H binding to DNA/RNA hybridization,

resulting in low processivity, short run-length, and low unidirectionality. Furthermore, the

simulation predicted that large DNA/RNA hybridization rate improves the trade-off.

Finally, based on the prediction of the simulation, we designed new DNA-AuNP motor by

using a DNA/RNA sequence which shows 3.8-times larger hybridization rate than the original one.

As a result, the newly-designed motor simultaneously achieved speed of 30 nm s^{-1}, processivity

of 200, and run-length of 3 μm comparable to motor proteins. Our study demonstrates a

mechanism-based strategy to improve the performance of artificial molecular motors.

Results

Stepping motion of DNA-AuNP motor visualized by high-speed/high-precision tracking

The 100-nm DNA-AuNP motor was prepared by using the DNA/RNA sequences similar to the

previous study (Supplementary Table 1)[18]. The densities of the DNA on AuNP surface and the

RNA on glass substrate surface were 0.11 ± 0.02 molecules nm^{-2} and 0.022 ± 0.008 molecules

nm^{-2}, respectively (Supplementary Fig. 1 and 2).

For high-speed/high-precision single-particle tracking, we obtained image sequences at

20–1000 frames per second (fps), much higher rate than the previous study (0.2–2 fps)[18], by using

a total internal reflection dark-field microscopy (Fig. 1a)[25,26]. For long-term measurements at 36

nM RNase H, similar [RNase H] used in the previous study[18], the localization precision was 3.1

nm. At other [RNase H]s (see below), the localization precision was 0.8–1.0 nm (Supplementary

Fig. 3).

Without RNase H, the motors were stationary on the RNA substrate, and exhibited 2D

motions immediately after the RNase H addition (Fig. 1b, Supplementary Video 1). During

observations, most motors kept moving and some detached from the substrate (the ratio of

detachment depended on [RNase H], see below). Interestingly, 2D trajectories of the motor clearly

showed pauses and steps (Fig. 1c). These stepping motions were resolved for the first time.

Then, the pauses and steps were detected objectively by using a step-finding algorithm (Fig.

1d and e)[27]. Individual pause lengths and step sizes were collected for subsequent statistical

analysis. We also calculated the speed by dividing each step size with each pause length (Fig. 1f).

At 36 nM RNase H, the median pause length, step size, and speed were 70 s, 24 nm, and 0.33 nm

s^{-1}, respectively (Fig. 2a-c). Therefore, we conclude that low speed is caused by the long pauses.

Speed increases as [RNase H] increases and reaches 100 nm s^{-1}

To gain information on the chemical processes which lengthen the pauses, we next conducted

fluorescence bulk assays of DNA/RNA hybridization and RNA hydrolysis (Extended Data Fig. 1a,

b). We found that apparent hybridization rate increased linearly up to 100 nM DNA (RNA

concentration was fixed at 10 nM) (Extended Data Fig. 1c, d). Estimated rate constants for

DNA/RNA hybridization ($k_{on}^{DNA/RNA}$) and dissociation ($k_{off}^{DNA/RNA}$) were $(3.9 \pm 0.5) \times 10^5$ $M^{-1}s^{-1}$

and $(2.0 \pm 0.9) \times 10^{-3}$ s^{-1}, respectively (Supplementary Table 2). On the other hand, apparent RNA

hydrolysis rate showed a saturation above micromolar RNase H (Extended Data Fig. 1e, f). Rate

constants for RNase H binding (k_{on}^{E}) and RNA hydrolysis (k_{cat}^{E}) were estimated to be (2.2 ± 0.3)

$\times 10^6$ $M^{-1}s^{-1}$ and 1.6 ± 0.1 s^{-1}, respectively, indicating that RNase H binding is the bottleneck of

RNA hydrolysis at nanomolar [RNase H].

We then conducted single-particle tracking at different [RNase H]s from 36 to 7200 nM

(Supplementary Fig. 4, Supplementary Video 1, 2, 3). Resulting pause lengths, step sizes, and

speeds are summarized (Fig. 2a-c, Supplementary Fig. 5). We found that the pause length largely

shortened from ~100 s to ~0.1 s as [RNase H] increased (Fig. 2a), while the step size (~20 nm) did

not change largely (Fig. 2b). As results, the speed increased as [RNase H] increased (Fig. 2c).

Remarkably, at highest 7200 nM RNase H, the speed reached 100 nm s^{-1}.

Geometry-based kinetic simulation reproduces stepping motion and [RNase H] dependence

To understand the mechanism of stepping motion of the DNA-AuNP motor, we developed a

geometry-based kinetic simulation (Supplementary Note, Supplementary Fig. 6, Supplementary

Table 3). Briefly, individual DNAs on the AuNP faced with the RNA-substrate are represented by

the square grids in the accessible region (Fig. 2d, Supplementary Fig. 7), and RNAs are randomly

distributed on the same size grids on the substrate surface to reproduce density difference between

DNA and RNA. Then, reactions are stochastically progressed in parallel at individual RNA sites

in the accessible region. The reaction model is composed of three sequential processes: 1)

DNA/RNA hybridization ($k_{on}^{DNA/RNA}$), 2) RNase H binding to DNA/RNA duplex (k_{on}^{E}[RNase H]),

and 3) RNA hydrolysis (k_{cat}^{E}) (Fig. 2e). We do not consider dissociation of the intact DNA/RNA

duplex[28], because $k_{off}^{DNA/RNA}$ estimated in the bulk assay was very small (Supplementary Table 2).

The XY-coordinates of the DNA-AuNP are randomly varied within the overlapped region among

the mobile regions of all RNA sites bound with DNA (Supplementary Fig. 8).

As a result, simulated trajectories showed stepping motions similar to the experiments (Fig.

2f, Supplementary Video 4, 5, 6). Note that in the simulation, the raw trajectories showed much

smaller fluctuations than the experiments during the pauses (Fig. 2f), because the motor is

completely stationary if there is no overlap among the mobile regions of all bound sites. Thus, in

order to reproduce the experiments more precisely, we added Gaussian noise to the raw simulation

trajectories (Fig. 2g, Supplementary Note, Supplementary Fig. 9 and 10). Furthermore, based on

the values obtained by the bulk assays (Supplementary Table 2), we optimized kinetic parameters

used in the simulation by using a simulation-based fitting (Supplementary Fig. 11-16). In short,

small numbers of the simulations were conducted with different kinetic parameters, and the mean

square errors (MSE) of pause length and step size between the simulation and experiment were

calculated. The optimal set of the kinetic parameters which showed the minimum MSE was

$k_{on}^{DNA/RNA}$ = 0.3 s^{-1}, k_{on}^{E} = 1.0 × 10^6 M^{-1}s^{-1}, and k_{cat}^{E} = 4.0 s^{-1} (Supplementary Fig. 16,

Supplementary Table 3). Then, numbers of simulations were increased and each trajectory was

analyzed again. As results, the pause lengths, step sizes, and speeds obtained by the simulation

showed excellent agreement with those obtained by the experiment at all [RNase H]s (Fig. 2h-j).

[RNase H]-dependent coupling between pause-step cycle and elementary chemical processes

We investigated how the coupling between pause-step cycle and elementary chemical processes

depends on [RNase H]. In our simulation, k_{on}^{E}[RNase H] increases as [RNase H] increases, while

$k_{on}^{DNA/RNA}$ and k_{cat}^{E} are constant (Supplementary Table 3). Therefore, relative amplitude among

these rates varies depending on the [RNase H] (Fig. 3a, regions (I)–(III)). The boundaries between

(I) and (II), and (II) and (III) are 300 and 4000 nM RNase H, respectively. We then analyzed time

course of the numbers of RNAs unbound ($N_{unbound}$) and bound (N_{bound}) with DNAs and hydrolyzed

RNAs ($N_{hydrolyzed}$) in the accessible region at 36, 720, and 7200 nM RNase H, each of which

corresponds to regions (I), (II), and (III), respectively (Fig. 3b, Supplementary Video 4, 5, 6). We

focused on time points corresponding to (i) just after the step, (ii) during the pause, (iii) just before

the next step, and (iv) just after the next step (= (i)). The spatial distributions of RNA states at each

time point (i)–(iv) are also shown.

At 36 nM RNase H (Fig. 3b, left), at (i), DNAs contacted with new unbound RNAs and

started to form new hybridizations. Note that the unbound and hydrolyzed RNAs distributed in the

forward and backward half of the accessible region, respectively, because the step size (~20 nm)

is comparable to the radius of accessible region (R_{acc} = 28.0 nm, Supplementary Fig. 7). Between

(i) and (ii), N_{bound} increased faster than $N_{hydrolyzed}$, because $k_{on}^{DNA/RNA}$ is much larger than

k_{on}^{E}[RNase H] at region (I). At (ii), DNA/RNA hybridizations were completed. Between (ii) and

(iii), N_{bound} and $N_{hydrolyzed}$ decreased and increased, respectively. Then, between (iii) and (iv), the

step occurred again when N_{bound} decreased to less than 10. At (iv), DNAs became accessible to

new unbound RNAs again. Importantly, almost all RNAs under the trajectory of DNA-AuNP were

hydrolyzed during the stepping motion as the "perfect" BBR (Fig. 3c, left).

At 720 nM RNase H (Fig. 3b, center), the hybridization of unbound RNAs and hydrolysis

of bound RNAs proceeded in parallel between (i) and (iii), resulted in nearly constant N_{bound} around

10. Interestingly, there were some unbound RNAs in the forward half of the accessible region at

(iii). Then, many of them moved to the backward half after the step ((iv)), and hydrolyzed before

the next step. Thus, the BBR mechanism still works (Fig. 3c, center).

At 7200 nM RNase H (Fig. 3b, right), on the other hand, the N_{bound} was kept lower than 10

throughout (i) to (iv), because $k_{on}^{DNA/RNA}$ is much lower than k_{on}^{E}[RNase H] and k_{cat}^{E}. This resulted

in high probability of detachment and low processivity and short run-length (see below). Moreover,

there were many unbound RNAs in the entire accessible region even at (iii), and remained intact

under the trajectory of DNA-AuNP (Fig. 3c, right). This "imperfect" BBR caused frequent

backward steps which resulted in low unidirectionality (see below).

We also performed statistical analysis of the $N_{unbound}$, N_{bound}, and $N_{hydrolyzed}$ just before (iii)

and after (iv) the steps (Fig. 3d). Notably, at all [RNase H]s, at both (iii) and (iv), the N_{bound} was

always around less than 10, consistent with the notion that the steps occur when N_{bound} <10. On

the other hand, at (iii), $N_{unbound}$ increased from nearly 0 to 40, and $N_{hydrolyzed}$ decreased from 40 to

5 as [RNase H] increased, reflecting the shift from the perfect to imperfect BBR mechanism.

Trade-off between speed and processivity/run-length and mechanism of detachment

Next, we examined [RNase H] dependence of processivity and run-length (Fig. 4). For

experimental data, time courses of the fraction of motors remained attached on RNA substrate

were fitted with single exponential decay functions (Fig. 4a) to estimate time constants before

detachment (τ_{detach}) (Fig. 4b)[18,29]. Processivity and run-length estimated from τ_{detach}, mean pause

length, and mean step size (Fig. 4c) are shown in Fig. 4d and e, respectively. When [RNase H] <

1000 nM, the motor showed high processivity of 100~200 and long run-length of 3~4 μm.

However, when [RNase H] > 1000 nM, processivity and run-length largely decreased, and

importantly, similar [RNase H] dependence was observed in the simulation (Extended Data Fig.

2a, b).

Then, we investigated the mechanism of detachment in the simulation. We found that there

are two modes of detachment associated with processivity and run-length (Extended Data Fig.2c,

d). One is caused by the entrapment of DNA-AuNP into the area where surrounding RNAs are

already hydrolyzed (entrap detachment), and another is caused by the stochastic loss of all

DNA/RNA hybridizations (stochastic detachment). We also analyzed the ratio ($P_{unbound}$) of

unbound RNAs to unbound and hydrolyzed RNAs around the DNA-AuNP just before the

detachment (Extended Data Fig. 2e, f). We found that as [RNase H] increased, $P_{unbound}$ increased

in a sigmoidal fashion. When [RNase H] was higher than 1000 nM, $P_{unbound}$ was larger than 0.5,

indicating that stochastic detachment is major and results in low processivity and short run-length.

Trade-off between speed and unidirectionality

The DNA-nanoparticle and -microparticle motors show super-diffusion, where the motions are

biased forward by suppressing backward motions as 2D BBR[16,18]. In this study, to evaluate super-

diffusive motion directly, we analyzed step angles ($\Delta\theta$) between pauses (Fig. 5a, see

Supplementary Note and Supplementary Fig. 17 for the analysis of mean square displacement).

For both the experiment and simulation, when [RNase H] < 1000 nM, distribution of the $\Delta\theta$

showed a peak around 0°, indicating step directions are biased forward (Fig. 5b and d). Forward-

biased steps are more clearly shown in the polar plots (Fig. 5c and e). When [RNase H] > 1000

nM, on the other hand, the number of backward steps increased and the $\Delta\theta$ distribution became

more uniform, indicating low unidirectionality at high [RNase H] (Supplementary Fig. 18).

DNA/RNA sequence with large hybridization rate overcomes trade-off

We summarized relationships between the speed and other performances (Fig. 6, circles). Note

that to evaluate the unidirectionality quantitatively, we used the standard deviations of the $\Delta\theta$

distributions (see Methods). [RNase H] dependences of the speed, processivity, run-length, and

unidirectionality are also summarized (Supplementary Fig. 19). For both the experiment and

simulation, the trade-off between the speed and other performances was clearly observed at high

[RNase H].

Considering the mechanism shown in Fig. 3, we anticipated that large $k_{on}^{DNA/RNA}$ improves

the trade-off by expanding the perfect BBR region (Fig. 3, region (I)) to high [RNase H] region.

Actually, our simulation predicted that 4-times increase in $k_{on}^{DNA/RNA}$ results in no decreases in

processivity, run-length, and unidirectionality even at 7200 nM RNase H, although the speed

slightly decreases to 30 nm s^{-1} (Fig. 6d-f, Supplementary Fig. 19e-h, pink triangles, and

Supplementary Video 7).

5   Then, based on the prediction by the simulation, we re-designed the DNA/RNA sequence.

We introduced a repetitive sequence and increased GC content, which have been reported to

increase $k_{on}^{DNA/RNA}$ (Supplementary Table 1)[30]. Indeed, in the bulk assay, newly-designed

sequence showed 3.8-times larger $k_{on}^{DNA/RNA}$ than that of original one, while $k_{off}^{DNA/RNA}$ decreased

(Extended Data Fig. 3a, b, e, and f). We also confirmed that k_{on}^{E} and k_{cat}^{E} were not largely different

from those of the original sequence (Extended Data Fig. 3c, d, g, and h). Then, single-particle

tracking of the DNA-AuNP motor with the new sequence was performed at 1440, 3600, and 7200

nM RNase H (Supplementary Fig. 20 and 21). As a result, the trade-off between the speed and

other performances was actually improved (Fig. 6a-c, orange triangles). Notably, at 7200 nM

RNase H, the newly-designed motor simultaneously achieved speed of 30 nm s^{-1}, processivity of

200, run-length of 3 μm, and high unidirectionality (Supplementary Video 8).

Discussion

Here high-speed/high-precision single-particle tracking resolved the elementary pauses and steps

of the DNA-AuNP motor (Fig. 1). Reflecting its nanoscale, the step size (~20 nm) was comparable

to those of motor proteins such as 8 nm step for kinesin-1[12] and 36 nm step for myosin V[13,31].

Furthermore, the step size was independent on [RNase H], suggesting that it is determined by the

particle size. It has been reported that DNA-microparticle motor shows higher speed than that of

DNA-nanoparticle motor, despite its small diffusion coefficient[16,17]. Our results suggest that the large particle shows large step size and facilitates the fast motion.

In contrast to the constant step size, the pause length largely shortened as [RNase H] increased (Fig. 2). The geometry-based kinetic simulation quantitatively reproduced the experiment and revealed that RNase H binding to DNA/RNA duplex is the bottleneck at nanomolar RNase H (Fig. 2 and 3). Our result is consistent with previous studies of nano-sized BBR motors where the speed of ~1 nm s^{-1} were reported at 14–120 nM RNase H[18,29], and suggests that the speed of these motors can be increased by increasing [RNase H].

At high [RNase H], however, our study also revealed that processivity, run-length, and unidirectionality were sacrificed due to the switch of the bottleneck from RNase H binding to DNA/RNA hybridization (Fig. 4-6). Our simulation predicted that large hybridization rate improves the trade-off, and the newly-designed motor with large $k_{on}^{DNA/RNA}$ simultaneously achieved speed, processivity, and run-length comparable to motor proteins (Fig. 6)[11-14]. Note that the processivity is even higher than that of a biological BBR motor processive chitinase that moves unidirectionally with the speed of 50 nm s^{-1} and processivity of 60~80 along the single chitin chain on crystalline chitin surface[14,32].

In our simulation, we can easily change the values of $k_{on}^{DNA/RNA}$, k_{on}^{E}, and k_{cat}^{E}. In general, the larger the values of these kinetic parameters, the better performances are expected. In the experiment, large $k_{on}^{DNA/RNA}$ can be achieved by re-designing DNA/RNA sequences, as demonstrated in this study (Fig. 6). The large k_{on}^{E} and k_{cat}^{E} are more challenging, but possible by using RNase H from other species or by engineering RNase H mutant. Furthermore, our simulation is versatile for other artificial BBR motors such as a protease-modified microparticle motor moving on a peptide-modified substrate recently reported[33].

Another important issue is the fundamental limitation of processivity and run-length by the

entrap detachment (Extended Data Fig. 2), which originates from the fact that the motion of DNA-

AuNP motor is not completely unidirectional (Fig. 5). Several studies have reported that

microparticle-based BBR motors translocate in a quasi-unidirectional manner along

micropatterned fuel surfaces[16,33]. Also, DNA nanotubes and origami tiles, which allow for the

elaborate arrangement of interaction sites, can serve as a linear or more complex track for the

DNA-AuNP motors[9,34]. Furthermore, nano-sized objects with anisotropic shapes such as DNA

origami rod and gold nanorod would restrict the Brownian motion to one direction and improve

the unidirectionality[29,35]. Our next goal is development of nano-sized BBR motors capable of

directional and speed controls, which are applicable to sensing or computation based on the

changes in the motional behaviors[36].

Methods

Regents

DNAs and RNAs used in this study (Supplementary Table 1) were commercially synthesized and

purified by high-performance liquid chromatography. DNA and DNA/RNA chimera substrate

were purchased from Tsukuba Oligo Service (Ibaraki, Japan). DNA anchor (3'-alkyne-modified)

was purchased from Fasmac (Kanagawa, Japan). RNase H from *E. coli* (Product No. 2150A) was

purchased from Takara Bio Inc. (Shiga, Japan).

Preparation of 100 nm DNA-AuNPs

Twenty µL of 100 µM DNA leg T30 in the phosphate-buffered saline (PBS, pH 7.4) was added to

180 µL of a suspension of commercially purchased 100 nm diameter gold colloid (BBI Solutions)

in a glass vial. The solution was cooled and frozen by placing it in a –30 °C freezer for 20 min.

Then, the solution was allowed to thaw at room temperature for 15 min. Freeze-thaw cycles were

repeated three times. Then, the solution was centrifuged (6000 rpm for 1 min) and the pellet was

washed with 100 µL of PBS containing Triton X-100 (0.75% v/v) seven times. The obtained DNA-

AuNPs were stored at 4 °C.

Measurement of DNA density on AuNP surface

Ten µL of fluorescein (FAM) labeled DNA-AuNPs (~10 pM, Supplementary Fig. 1) was mixed

with 10 µL of 1M DTT and incubated at 50°C for 30–60 min. After the incubation, the solution

was centrifuged (6000 rpm for 1 min) and supernatant was collected. Then, 180 µL of PBS was

added and fluorescence intensity at 520 nm (excitation: 490 nm) was measured using a

fluorescence spectrometer (FP8300, JASCO). Released FAM-DNA concentration was quantified

from the fluorescence intensity using a calibration curve. The number of DNAs on AuNP was

calculated from the ratio of the concentration of released FAM-DNA to that of AuNP, and then

DNA density on AuNP surface was calculated as the number of DNAs on AuNP divided by the

surface area assuming a sphere with a 100 nm diameter.

Preparation of KOH washed cover glass used for flow cell

The cover glass (9×30 mm, #1, Matsunami Glass Industry, Japan) was washed and sonicated in

ethanol for 10 min. Then, the glass was rinsed 3 times with ultrapure water (18.2 MΩ cm^{-1}) and

sonicated in ultrapure water for 10 min. The cover glass was rinsed again with ultrapure water.

After thoroughly drying with N_2 blow, the cover glass was immersed in the KOH 10 M for 1 hour.

Then, the cover glass was well rinsed with ultrapure water. The cleaned cover glass was dried with

N_2 blow and kept on the clean rack in the closed container prior to use.

Preparation of amino-silane coated cover glass used for flow cell

The cover glass (24×32 mm, #1 SHT, Matsunami Glass Industry) was washed and sonicated in

ethanol for 10 min. Then, the glass was rinsed with ultrapure water 6 times and sonicated in

ultrapure water for 10 min. The cover glass was then rinsed again with ultrapure water 6 times and

immersed in the piranha solution for 1 hour. The cover glass was rinsed and washed well with

ultrapure water. Then, the cover glass was rinsed with ethanol and immersed in [3-aminopropyl]-

triethoxysilane in ethanol (2% v/v) for 60 min at room temperature. After that, the cover glass was

rinsed well with ethanol and cured in an oven at 110 °C for 10 min. The cover glass was kept on

the clean rack in the closed container prior to use.

Flow cell preparation for single-particle tracking experiment

The flow cell for DNA-AuNP motor observation was prepared by using the amino-silane coated glass (bottom side) and the KOH washed cover glass (top side) and a spacer (a double-sided tape, thickness of 0.62 mm) (Supplementary Fig. 2a). The length and width of each flow cell were 9.0 mm and 2.5 mm, respectively, and a grease (Dow Corning Toray) was used to isolate each flow cell. Each flow cell was then incubated with 10 mg/mL azide-peg4-NHS (Tokyo Chemical Industry) dissolved in 100 mM $NaHCO_3$ solution (pH 9.0) for 1 hour at room temperature and washed with ~50 µL ultrapure water (Supplementary Fig. 2b). Then, flow cells were washed with 1 M potassium phosphate (pH 7.4), and 30 µL of 20 µM alkyne-modified DNA anchor, 100 µM Tris(3-hydroxypropyltriazolylmethyl)amine (THPTA), 20 µM $CuSO_4$ and 2 mM sodium ascorbate, 1 M potassium phosphate (pH 7.4) was added to the flow cell and incubated for 1 hour at room temperature. After that, the flow cell was washed with PBS, and then 30 µL of 1 µM Cy3-RNA/DNA chimera in PBS was added and incubated overnight at room temperature in a box with wet filter paper attached to the inside wall. After incubation, unbound RNAs were washed out with ~50 µL PBS and stored in the box prior to use.

Measurement of RNA density on cover glass surface

The flow cell of which the bottom surface was modified with Cy3-RNA/DNA chimera was incubated with 20 µL of 100 µg mL^{-1} RNase A in PBS for 5 min at room temperature (Supplementary Fig. 2c). Then, 20 µL of the reacted solution was collected and added into a cuvette for fluorescence measurement. This procedure was repeated at least three times, to assure that all of the Cy3-RNA/DNA chimera on cover glass surface were cleaved by RNase A. The collected solution was then diluted to a final volume of 300 µL with PBS. The concentration of

Cy3-RNA was quantified from the fluorescence intensity at 560 nm (excitation: 510 nm) using a

calibration curve obtained with a fluorescence spectrometer (FP8300, JASCO). The total number

of RNA molecules was calculated as a product of concentration and volume (300 μL). RNA

surface density was calculated from the total number of RNA molecules divided by the area of the

bottom surface of the flow cell (9.0 mm × 2.5 mm).

Single-particle tracking of DNA-AuNP motor with scattering imaging

A total internal reflection dark-field microscope was built as previously described[26,37]. We used

the point illumination for observation[26,37]. Observation was conducted at $25 \pm 1°C$, and scattering

signals from the DNA-AuNPs were imaged with a high-speed CMOS camera (Fastcam AX100;

Photron). An observation buffer (50 mM Tris-HCl, 10 μM DTT, 0.75% v/v Triton X-100, 3 mM

$MgCl_2$, pH 8.3) was used. The flow cell was placed on the stage of microscope and incubated with

3 pM DNA-AuNPs solution for 15 min. Then, the flow cell was washed with the observation

buffer, and the sample stage was left untouched at least 10 min to minimize the stage drift. The

motion of DNA-AuNP was observed with designated RNase H concentration ([RNase H]).

Recording rate of 20 frames per second (fps) was used for 36, 144, and 360 nM RNase H, 250 fps

for 720 and 1440 nM RNase H, and 1000 fps for 3600 and 7200 nM RNase H. For each recording

rate, we adjusted the laser power density to avoid the saturation of the scattering signal

(Supplementary Fig. 3). At 36 nM RNase H condition, cooling fan of the CMOS camera had to be

turned on for the long-time recording for more than 300 s to avoid overheating. Then, sequential

images of individual DNA-AuNP motors were analyzed separately by home-built Python code. At

each frame, X- and Y-coordinates of the centroid of each DNA-AuNP motor were determined by

the two-dimensional Gaussian fitting. The localization precision at 36 nM RNase H condition was

3.1 nm for both X- and Y-coordinates, and those at other [RNase H] conditions were 0.8–1.0 nm

(Supplementary Fig. 3).

Analysis of step size and pause length

Trajectories of the centroid of the DNA-AuNP motors were analyzed by home-built Python code.

First, steps along X- and Y-coordinates were identified separately by using an algorithm developed

previously[27]. Then, diagonal steps were counted when the steps along X- and Y-coordinates

occurred within ± 5 frames because of the limited accuracy of the step-finding algorithm. Then,

steps along X- and Y-coordinates and diagonal steps were combined to obtain all pause lengths

and step sizes in entire trajectories.

Analysis of processivity and run-length

In the single-particle tracking experiments, especially at low [RNase H], significant fraction of the

motors did not show detachment from the RNA substrate during observation. Therefore, we

followed an approach used for other nano-sized BBR motors[18,29], in which time course of the

fraction of motors remained attached on RNA substrate was measured. Then, time courses were

fitted with single exponential decay functions to estimate time constants before detachment (τ_{detach}),

and processivity and run-length were estimated by following equations, respectively.

$$\text{Processivity} = \tau_{\text{detach}}/\text{Mean pause length}$$

$$\text{Run-length} = \text{Processivity} \times \text{Mean step size}$$

On the other hand, simulation was conducted until each motor showed the detachment. Therefore,

for simulation data, we directly estimated the processivity (number of steps before detachment)

from each trajectory, and calculated run-length by using the equation described above. For

simulation, mean values are calculated from 50 trajectories for each [RNase H] condition.

Analysis of step angle and unidirectionality

Step angles ($\Delta\theta$) were calculated from the XY coordinates of three consecutive pauses. In the case

of 2D Brownian motion, the histogram of the $\Delta\theta$ converges to a uniform distribution between

−180° and 180°, while in the case of completely unidirectional motion, the histogram of the $\Delta\theta$

has a single peak at 0°. Unidirectionality of the DNA-AuNP motor was quantified by using the

standard deviation of the $\Delta\theta$ distribution:

10
$$\text{Unidirectionality} = \frac{\sigma_{\text{uniform}} - \sigma_{\text{angle}}}{\sigma_{\text{uniform}}}$$

where σ_{angle} and σ_{uniform} are the standard deviations of the $\Delta\theta$ distribution of the data and that

assuming the uniform distribution, respectively. Theoretically, the value of the σ_{uniform} is 103.9°,

and the upper and lower limits of the unidirectionality are 1 (when σ_{angle} is 0°) and 0 (when σ_{angle}

= σ_{uniform}), respectively.

DNA/RNA hybridization and RNA hydrolysis assays in bulk solution

We applied a fluorescence-based assay in which the quantum yield of Cy3-oligonucleotide is

sensitive to the hybridization state[38,39]. A fluorescence spectrometer (FP8300, JASCO) was used

to monitor hybridization between Cy3-labeled RNA and the complementary ssDNA in a free-

solution. All measurement was performed at room temperature. The reaction buffer was the same

as that used for the DNA-AuNP motor observation. Fluorescence from 998 μL of 10 nM Cy3-

RNA in a cuvette was monitored continuously with excitation and emission wavelengths of 515

and 565 nm and slit widths of 10 and 20 nm, respectively. Two μL of concentrated DNA was

added to the 998 μL Cy3-RNA solution while stirring the solution at 500 rpm with a magnetic stirrer. The final concentrations of DNA ($[DNA]_0$) were 6–100 nM. Then, the change in fluorescence intensity was monitored for 300 s with 1 s time resolution, and apparent hybridization rate at each $[DNA]_0$ was estimated as the inverse of the relaxation time. Measurement was repeated at least 3 times for each condition. The values of $k_{on}^{DNA/RNA}$ and $k_{off}^{DNA/RNA}$ (Supplementary Table 2) were determined from the $[DNA]_0$ dependence of the apparent hybridization rate (Extended Data Fig. 1e).

For RNA hydrolysis assay, the condition of fluorescence measurement was the same with hybridization assay described above, except for the time resolution of 0.02 s. Ten nM Cy3 RNA and 20 nM DNA were mixed and incubated for more than 10 min at room temperature for hybridization. Then, 4 uL of concentrated RNase H was added to the cuvette, and final concentrations of RNase H ($[RNase\ H]_0$) were 69-6912 nM. The decay of fluorescence intensity was monitored for 130 s. Measurement was repeated at least 3 times for each condition. The decay curve was fitted by the single exponential decay to determine the apparent RNA hydrolysis rate. The values of k_{on}^{E} and k_{cat}^{E} (Supplementary Table 2) were determined from the $[RNase\ H]_0$ dependence of the apparent hybridization rate (Extended Data Fig. 1f).

Geometry-based kinetic simulation of DNA-AuNP motor

The home-built code for the geometry-based kinetic simulation of DNA-AuNP motor was implemented in Python 3.9. Details of the algorithm and the simulation parameters are described in Supplementary Information (Supplementary Note, Supplementary Fig. 6-8, Supplementary Table 3). First, a two-dimensional pixel matrix was defined to model the RNAs on the glass surface. Pixel size was normalized by the DNA density on the AuNP (0.10 molecules nm^{-2}). RNAs

were randomly distributed with a ratio of the RNA density (0.022 molecules nm^{-2}) to the DNA

density. Reaction at a single RNA site is assumed to contain three sequential elementary steps:

DNA/RNA hybridization, RNase H binding to DNA/RNA duplex, and RNA hydrolysis. The

reaction proceeds within the accessible area of DNA with the radius of 28.0 nm (Supplementary

Fig. 7). Rate constants of the three elementary steps, $k_{on}^{DNA/RNA}$, k_{on}^{E}, and k_{cat}^{E} are the simulation

parameters. The optimal set of the rate constants was obtained through the simulation-based fitting

(see below) and used as the default values. Dwell time for each elementary step at each RNA site

is determined by a random sampling from the exponential distribution: $P(\tau) = k\exp(-kt)$, and the

site with the shortest dwell time is changed to the next state. Simulation steps proceeded in units

of reaction events. Constraint of the motor position was considered by introducing mobile region

of each DNA/RNA hybrid (25.2 nm) (Supplementary Fig. 8). The motor position was determined

randomly within the region where all mobile regions overlap. Unless all mobile regions overlapped,

the motor position was fixed. To reproduce the experimental trajectory accurately, Gaussian noise

with standard deviation of 2 or 6 nm was added to the raw simulated trajectories (Supplementary

Fig. 9 and 10).

Simulation-based fitting to optimize kinetic parameters

Simulation-based fitting was employed to obtain the optimal set of the rate constants ($k_{on}^{DNA/RNA}$,

k_{on}^{E}, k_{cat}^{E}) which reproduce the experimental results more precisely (Supplementary Fig. 11-16),

because the rate constants on the surface can change from those estimated in the bulk assay

(Extended Data Fig. 1, Supplementary Table 2). The geometry-based kinetic simulation of the

DNA-AuNP motor was performed with all combinations of $k_{on}^{DNA/RNA} = \{0.01, 0.05, 0.1, 0.2, 0.3,$

$0.4, 0.5, 1.0, 5.0\}$ s^{-1}, $k_{on}^{E} = \{0.5, 1.0, 1.5, 2.0, 2.5\} \times 10^{6}$ $M^{-1}s^{-1}$, and $k_{cat}^{E} = \{0.5, 1.0, 2.0, 3.0,$

4.0, 5.0, 6.0} s^{-1}. The [RNase H] of 36, 144, 360, 720, 1440, 3600, and 7200 nM were tried for

each parameter set, and 20 simulations were carried out for each combination. Pause length and

step size were calculated from the simulated trajectories by a step-finding algorithm. As the

evaluation function, the sum of the mean squared errors (MSE) of the pause length and step size

against the experimental data was calculated by using following equation:

6
$$\text{MSE} = \frac{1}{n}\sum_{i}^{n}\left\{\log \tilde{\tau}_{i,\text{sim}} - \log \tilde{\tau}_{i,\text{exp}}\right\}^2 + \left\{\log \tilde{\Delta}_{i,\text{sim}} - \log \tilde{\Delta}_{i,\text{exp}}\right\}^2$$

7   where $\tilde{\tau}_{i,\text{sim}}$ and $\tilde{\tau}_{i,\text{exp}}$ are the median pause length in simulation and experiment, $\tilde{\Delta}_{i,\text{sim}}$ and $\tilde{\Delta}_{i,\text{exp}}$

8   are the median step sizes in simulation and experiment, and n is the number of the conditions of

9   [RNase H]. The parameter set ($k_{\text{on}}^{\text{DNA/RNA}}$, k_{on}^{E}, $k_{\text{cat}}^{\text{E}}$) which showed the minimal MSE was used

10   for the subsequent 50 simulations (Supplementary Figure 16, Supplementary Table 3).

Figure 1. Observation of elementary processes of motion of DNA-AuNP motor

(a) Schematic illustration of the experimental system for high-speed/high-precision single particle tracking of DNA-AuNP motor. DNAs on the AuNP surface are hybridized with RNAs on the glass surface. RNAs that hybridized with DNAs are selectively hydrolyzed by RNase H. The motion of DNA-AuNP motor is visualized by using total-internal reflection dark-field microscopy. **(b)** An image sequence of motion of DNA-AuNP motor on RNA surface at 36 nM RNase H. Recording rate: 20 fps. Red lines indicate trajectories of the particle centroid. Dark yellow squares show the regions of interest to calculate the centroid for each frame. Scale bar: 200 nm. **(c)** Whole trajectory of the centroid of DNA-AuNP motor shown in (b). **(d)** Detected pauses (green dots) and steps (magenta lines) superimposed to raw trajectory shown in (c). **(e)** Time-course of X- and Y-coordinates. Green solid lines are fittings to the raw trajectories, and magenta dashed lines indicate the positions of detected steps. **(f)** Schematic illustration and definitions of pause length, step size, and speed.

Figure 2. [RNase H] dependence of pause length, step size, and speed measured by single-particle tracking experiments and geometry-based kinetic simulations

(a-c) [RNase H] dependence of pause length (a), step size (b), and speed (c) in the experiments. Individual data are shown by cyan circles. Box plot shows interquartile range (IQR). Error bars show 5th/95th percentile for each [RNase H] condition. Recording rates were 20, 20, 250, 250, 250, 1000, 1000 fps for 36, 144, 360, 720, 1440, 3600, and 7200 nM RNase H, respectively. **(d)** Schematic illustration of geometry-based kinetic simulation. Hybridization between RNA and DNA occurs stochastically within an area with the accessible radius (R_{acc}) of 28.0 nm, where single DNA-AuNP motor is accessible to the RNA surface (dark yellow circle). Multivalent DNA/RNA hybridizations immobilize the motor and result in the pauses, and the steps occur after the successive degradations of RNAs by RNase H. **(e)** Kinetic model at each RNA-modified site on the surface. $k_{on}^{DNA/RNA}$, k_{on}^{E}, and k_{cat}^{E} correspond to the rate constants of DNA/RNA hybridization, RNase H binding to DNA/RNA duplex, and RNA hydrolysis, respectively. Empty sites are inert in the simulation. The values of $k_{on}^{DNA/RNA}$, k_{on}^{E}, and k_{cat}^{E} were set to 0.3 s^{-1}, 1.0×10^6 M^{-1}s^{-1}, and 4 s^{-1}, respectively. **(f)** Representative trajectory of the DNA-AuNP motor translocation superimposed on spatial distribution of RNA states at 36 nM RNase H. **(g)** Whole trajectory of the centroid after the addition of the Gaussian noise ($\sigma_{noise} = 6$ nm). **(h-j)** [RNase H] dependence of pause length, step size, and speed in the simulation. Individual data are shown by green circles.

1
Figure 3. Stepping mechanisms depending on [RNase H]

(a) Relationships among three reaction rates considered in this study ($k_{on}^{DNA/RNA}$, k_{on}^E[RNase H],
and k_{cat}^E) depending on [RNase H]. Blue, yellow, and red shaded regions correspond to different
order of magnitude of these rates (I)–(III). **(b)** Time courses of the numbers of unbound RNAs,
RNAs bound with DNAs, and hydrolyzed RNAs at 36 (left), 720 (middle), and 7200 (right) nM
RNase H. X- and Y-coordinates of the motion are also shown. Black dashed lines labeled by (i)–
(iv) show the representative frames at (i, iv) just after step, (ii) during pause, (iii) just before step.
Spatial distributions of RNA around DNA-AuNP motor corresponding to the frames of (i)–(iv)
are shown in the right of the time courses. Black circles represent accessible region of the DNA-

AuNP motor. Gray dashed circles in (iv) show the accessible region in (iii). Pink arrows indicate the step directions. **(c)** Schematic illustrations of the stepping mechanisms at different [RNase H]s at regions (I)–(III). **(d)** Violin plots of the number of RNA sites within the accessible region just before and after steps. Black solid line shows the median value.

Figure 4. [RNase H] dependence of processivity and run-length of the experimental data

(a) Typical plots of fraction of motors remained attached to the RNA surface as a function of time. Each plot was fitted by single exponential decay function, $\exp(-t/\tau_{detach})$, that is superimposed as the black dashed line. (b) [RNase H] dependence of τ_{detach}. Black bars and error bars correspond to mean ± s.d. The mean and s.d. values were calculated from at least three independent experiments for each [RNase H] condition (cyan circles). (c) Schematic illustration of whole kinetics from attachment to detachment from RNA surface. Processivity was defined as the number of steps before detachment and calculated by dividing the τ_{detach} by mean pause length. Run-length was estimated by multiplying the processivity with mean step size. (d, e) [RNase H] dependence of processivity (d) and run-length (e).

Figure 5. [RNase H] dependence of step angle

(a) Schematic illustration describing the calculation of step angle ($\Delta\theta$). Leftward and rightward steps are defined as positive and negative, respectively. **(b, c)** Histograms (b) and polar plots (c) of step angles at each [RNase H] in the experiments. **(d, e)** Histograms (d) and polar plots (e) of step angles at each [RNase H] in the simulations.

Figure 6. Trade-off between motor speed and other performances

(a-c) Relationship between the speed and processivity (a), or run-length (b), or unidirectionality (c) in the experiments. Blue circles indicate the data using original DNA/RNA sequence at 36, 144, 360, 720, 1440, 3600, and 7200 nM RNase H. Orange triangles indicate the data using newly-designed DNA/RNA sequence at 1440, 3600, and 7200 nM RNase H. The upper right and lower left regions of each panel correspond to high and low performances, respectively. Diagonal red dashed lines in (a) and (b) show the trade-off between speed and processivity/run-length due to the stochastic detachment, and horizontal brown dashed lines show the upper limit of processivity/run-length due to the entrap detachment. In (c), high and low unidirectionalities correspond to the perfect and imperfect BBR, respectively. **(d-f)** Relationship between the speed and processivity (d) or run-length (e) or unidirectionality (f) in the simulations. Green circles indicate the data using $k_{\text{on}}^{\text{DNA/RNA}} = 0.3$ s^{-1}, $k_{\text{on}}^{\text{E}} = 1.0 \times 10^6$ M^{-1}s^{-1}, and $k_{\text{cat}}^{\text{E}} = 4$ s^{-1} at 36, 144, 360, 720, 1440, 3600, and 7200 nM RNase H. Pink triangles indicates the data using $k_{\text{on}}^{\text{DNA/RNA}} = 1.2$ s^{-1}, $k_{\text{on}}^{\text{E}} = 1.0 \times 10^6$ M^{-1}s^{-1}, and $k_{\text{cat}}^{\text{E}} = 4$ s^{-1} at 1440, 3600, and 7200 nM RNase H.

1
Extended Data Fig. 1. Determination of kinetic rate constants in bulk solution. (a, b)
Schematic illustration and kinetic model of (a) DNA/RNA hybridization and (b) RNA hydrolysis
assays. Cy3-labeled RNA was used as a probe of DNA/RNA hybridization. The same sequence to
that of the DNA/RNA duplex part of the original sequence and the same solution composition for
single-particle tracking experiments were used. **(c)** Normalized time courses of fluorescence
intensity of the DNA/RNA hybridization assay. Initial RNA concentration was 10 nM, and
measurements were carried out at least in triplicate for each initial DNA concentration ($[DNA]_0$).
Error bars represent s.d. Time resolution was 1 s, and excitation and emission wavelengths were
set at 510 nm and 560 nm, respectively. **(d)** Plots of apparent hybridization rate as a function of
$[DNA]_0$. The values of $k_{on}^{DNA/RNA}$ and $k_{off}^{DNA/RNA}$ were determined as the slope and intercept of
the linear fitting (red line), respectively. **(e)** Normalized time courses of fluorescence intensity of
the RNA hydrolysis assay. Initial DNA and RNA concentrations were 10 nM, and measurements
were carried out at least in triplicate for each RNase H concentration ($[RNase\ H]_0$). Time resolution
was 0.02 s, and same excitation and emission wavelengths were used as (c). **(f)** Plots of apparent
hydrolysis rates as a function of $[RNase\ H]_0$. Error bars represent s.d. The value of k_{on}^E was
calculated by dividing the rates with $[RNase\ H]_0$ at 69 and 173 nM RNase H. The value of k_{cat}^E
was estimated from the saturating hydrolysis rate at 3456 and 6912 nM RNase H.

Extended Data Fig. 2. [RNase H] dependence of processivity and run-length, and two
different mechanisms of detachment revealed by the simulation. (a, b) Plots for processivity
(a) and run-length (b) at different [RNase H] conditions in the simulations. Mean values are
calculated from 50 trajectories for each condition (black bars). **(c, d)** Two representative
trajectories which show entrap (c) and stochastic (d) detachment at 720 nM RNase H. Black lines
correspond to the trajectories of the centroid of the motors. Large black dots indicate the start and
end of each trajectory. Regions around the end points are indicated by cyan squares and enlarged
in lower panels. Dark yellow solid and dashed circles indicate the accessible region at the final
frame and the region with the twice radius of the accessible region ($2 \times R_{acc}$), respectively.
Trajectories are overlaid on the distribution of RNA states on the surface. White, red, and gray
dots correspond to the empty, hydrolyzed, and unbound RNA sites, respectively. **(e)** An example
of 3D view corresponding to the last frame before entrap detachment. The $P_{unbound}$ is defined as
the ratio of unbound RNA sites to unbound and hydrolyzed RNA sites in the doughnut region with
the radius between R_{acc} to $2 \times R_{acc}$. Small or large value of the $P_{unbound}$ corresponds to entrap or
stochastic detachment is major, respectively. **(f)** [RNase H] dependence of the $P_{unbound}$. Median
values are calculated for each condition and shown by black horizontal bars.

Extended Data Fig. 3. Determination of kinetic rate constants in bulk solution with newly-designed DNA/RNA sequence. (a) Normalized time courses of fluorescence intensity of the DNA/RNA hybridization assay. Initial RNA concentration was 10 nM, and measurements were carried out at least in triplicate for each initial DNA concentration ($[DNA]_0$). ⎯Error bars represent s.d. Time resolution was 1 s, and excitation and emission wavelengths were set at 510 nm and 560 nm, respectively. **(b)** Plots of apparent hybridization rate as a function of $[DNA]_0$. The values of $k_{on}^{DNA/RNA}$ and $k_{off}^{DNA/RNA}$ were determined as the slope and intercept of the linear fitting (red line), respectively. **(c)** Normalized time courses of fluorescence intensity of the RNA hydrolysis assay. Initial DNA and RNA concentrations were 10 nM, and measurements were carried out at least in triplicate for each RNase H concentration ($[RNase\ H]_0$). Time resolution was 0.02 s, and same excitation and emission wavelengths were used as (a). **(d)** Plots of hydrolysis rates as a function of $[RNase\ H]_0$. Error bars represent s.d. **(e-h)** Bar graphs of rate constants for comparison between original (cyan) and designed (orange) DNA/RNA sequences. Error bars represent s.d. Black dashed line in (f) indicates the minimum pause-step cycle rate in single-particle tracking experiments calculated by 1/median pause length at 36 nM RNase H.

ASSOCIATED CONTENT

Supporting Information.

The Supporting Information is available free of charge.

AUTHOR INFORMATION

Corresponding Authors

Takanori Harashima − Institute for Molecular Science, National Institutes of Natural Sciences, Okazaki, Aichi 444-8787, Japan; Molecular Science Program, Graduate Institute for Advanced Studies, The Graduate University for Advanced Studies, SOKENDAI, Hayama, Kanagawa 240-0193, Japan

orcid.org/0000-0002-1322-5618; Email: harashima@ims.ac.jp

Ryota Iino − Institute for Molecular Science, National Institutes of Natural Sciences, Okazaki, Aichi 444-8787, Japan; Molecular Science Program, Graduate Institute for Advanced Studies, The Graduate University for Advanced Studies, SOKENDAI, Hayama, Kanagawa 240- 0193, Japan

orcid.org/0000-0003-0110-5704; Email: iino@ims.ac.jp

Authors

Akihiro Otomo − Institute for Molecular Science, National Institutes of Natural Sciences, Okazaki, Aichi 444-8787, Japan; Molecular Science Program, Graduate Institute for Advanced Studies, The Graduate University for Advanced Studies, SOKENDAI, Hayama, Kanagawa 240-0193, Japan

orcid.org/0000-0001-6709-4206; Email: otomoa@ims.ac.jp

Author Contributions

R.I. conceived the research. T.H., A.O., and R.I. designed the experiments. T.H. performed sample preparations, biochemical and fluorescence imaging-based characterizations, and single-particle tracking. T.H. and A.O. performed the data analysis. T.H. developed geometry-based kinetic simulation. R.I. supervised the project and preparing the manuscript with T.H. All authors discussed the results.

Funding Sources

MEXT | Japan Society for the Promotion of Science (JSPS) - 23K13645, 24H01732 [Harashima]

Tsugawa Foundation. [Harashima]

MEXT | Japan Society for the Promotion of Science (JSPS) - 18H05424, 23H04434 [Iino]

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

The authors thank Yasuko Okuni, Yayoi Kon, and Mayuko Yamamoto for their technical
assistance in sample preparations, Kosuke Matsumoto and Rosie Graham for helpful discussions.

REFERENCES

1.   Erbas-Cakmak, S., Leigh, D. A., McTernan, C. T. & Nussbaumer, A. L. Artificial

Molecular Machines. *Chem. Rev.* **115**, 10081-10206, (2015).

2.   Kassem, S. *et al.* Artificial molecular motors. *Chem. Soc. Rev.* **46**, 2592-2621, (2017).

3.   Li, S. *et al.* A DNA nanorobot functions as a cancer therapeutic in response to a molecular

trigger in vivo. *Nat. Biotechnol.* **36**, 258-264, (2018).

4.   Chen, Y. J., Groves, B., Muscat, R. A. & Seelig, G. DNA nanotechnology from the test

tube to the cell. *Nat. Nanotechnol.* **10**, 748-760, (2015).

5.   Iino, R., Kinbara, K. & Bryant, Z. Introduction: Molecular Motors. *Chem. Rev.* **120**, 1-4,

(2020).

6.   Lund, K. *et al.* Molecular robots guided by prescriptive landscapes. *Nature* **465**, 206-210,
(2010).

7.   Gu, H., Chao, J., Xiao, S. J. & Seeman, N. C. A proximity-based programmable DNA
nanoscale assembly line. *Nature* **465**, 202-205, (2010).

8.   Yin, P., Yan, H., Daniell, X. G., Turberfield, A. J. & Reif, J. H. A unidirectional DNA
walker that moves autonomously along a track. *Angew. Chem. Int. Ed.* **43**, 4906-4911,
(2004).

9.   Wickham, S. F. *et al.* A DNA-based molecular motor that can navigate a network of tracks.
Nat. Nanotechnol. **7**, 169-173, (2012).

10.   Pan, J. *et al.* Visible/near-infrared subdiffraction imaging reveals the stochastic nature of
DNA walkers. *Sci. Adv.* **3**, e1601600, (2017).

11.   Imai, H. *et al.* Direct observation shows superposition and large scale flexibility within
cytoplasmic dynein motors moving along microtubules. *Nat. Commun.* **6**, 8179, (2015).

12.   Schnitzer, M. J. & Block, S. M. Kinesin hydrolyses one ATP per 8-nm step. *Nature* **388**,
386-390, (1997).

13.   Ali, M. Y. *et al.* Myosin V is a left-handed spiral motor on the right-handed actin helix.
Nat. Struct. Mol. Biol. **9**, 464-467, (2002).

14.   Nakamura, A., Okazaki, K. I., Furuta, T., Sakurai, M. & Iino, R. Processive chitinase is
Brownian monorail operated by fast catalysis after peeling rail from crystalline chitin. *Nat.*
Commun. **9**, 3814, (2018).

15.   Cha, T. G. *et al.* A synthetic DNA motor that transports nanoparticles along carbon
nanotubes. *Nat. Nanotechnol.* **9**, 39-43, (2014).

16. Yehl, K. *et al.* High-speed DNA-based rolling motors powered by RNase H. *Nat. Nanotechnol.* **11**, 184-190, (2016).

17. Blanchard, A. T. *et al.* Highly Polyvalent DNA Motors Generate 100+ pN of Force via Autochemophoresis. *Nano Lett.* **19**, 6977-6986, (2019).

18. Bazrafshan, A. *et al.* DNA Gold Nanoparticle Motors Demonstrate Processive Motion with Bursts of Speed Up to 50 nm Per Second. *ACS Nano* **15**, 8427-8438, (2021).

19. Ron, M. & Rob, P. *Cell Biology by the Numbers* (Garland Science, 2015).

20. Howard, J. *Mechanics of motor proteins and the cytoskeleton* (Sinauer Associates, 2001).

21. Isojima, H., Iino, R., Niitani, Y., Noji, H. & Tomishige, M. Direct observation of intermediate states during the stepping motion of kinesin-1. *Nat. Chem. Biol.* **12**, 290-297, (2016).

22. Nishiyama, M., Higuchi, H. & Yanagida, T. Chemomechanical coupling of the forward and backward steps of single kinesin molecules. *Nat. Cell. Biol.* **4**, 790-797, (2002).

23. Nishizaka, T. *et al.* Chemomechanical coupling in F1-ATPase revealed by simultaneous observation of nucleotide kinetics and rotation. *Nat. Struct. Mol. Biol.* **11**, 142-148, (2004).

24. Wolff, J. O. *et al.* MINFLUX dissects the unimpeded walking of kinesin-1. *Science* **379**, 1004-1010, (2023).

25. Ueno, H. *et al.* Simple dark-field microscopy with nanometer spatial precision and microsecond temporal resolution. *Biophys. J.* **98**, 2014-2023, (2010).

26. Ando, J. *et al.* Single-Nanoparticle Tracking with Angstrom Localization Precision and Microsecond Time Resolution. *Biophys. J.* **115**, 2413-2427, (2018).

27. Kerssemakers, J. W. *et al.* Assembly dynamics of microtubules at molecular resolution. *Nature* **442**, 709-712, (2006).

28. Zhang, L., Piranej, S., Namazi, A., Narum, S. & Salaita, K. "Turbo-Charged" DNA Motors with Optimized Sequence Enable Single-Molecule Nucleic Acid Sensing. *Angew. Chem. Int. Ed.* **63**, e202316851, (2024).

29. Bazrafshan, A. *et al.* Tunable DNA Origami Motors Translocate Ballistically Over μm Distances at nm/s Speeds. *Angew. Chem. Int. Ed.* **59**, 9514-9521, (2020).

30. Hertel, S. *et al.* The stability and number of nucleating interactions determine DNA hybridization rates in the absence of secondary structure. *Nucleic Acids Res.* **50**, 7829-7841, (2022).

31. Mehta, A. D. *et al.* Myosin-V is a processive actin-based motor. *Nature* **400**, 590-593, (1999).

32. Nakamura, A. *et al.* Rate constants, processivity, and productive binding ratio of chitinase A revealed by single-molecule analysis. *Phys. Chem. Chem. Phys.* **20**, 3010-3018, (2018).

33. Korosec, C. S. *et al.* Motility of an autonomous protein-based artificial motor that operates via a burnt-bridge principle. *Nat. Commun.* **15**, 1511, (2024).

34. Ibusuki, R. *et al.* Programmable molecular transport achieved by engineering protein motors to move on DNA nanotubes. *Science* **375**, 1159-1164, (2022).

35. Zhou, C., Duan, X. Y. & Liu, N. A plasmonic nanorod that walks on DNA origami. *Nat. Commun.* **6**, (2015).

36. Piranej, S., Bazrafshan, A. & Salaita, K. Chemical-to-mechanical molecular computation using DNA-based motors with onboard logic. *Nat. Nanotechnol.* **17**, 514-523, (2022).

37. Ando, J. *et al.* Small stepping motion of processive dynein revealed by load-free high-speed single-particle tracking. *Sci. Rep.* **10**, 1080, (2020).

38. Sanborn, M. E., Connolly, B. K., Gurunathan, K. & Levitus, M. Fluorescence properties

and photophysics of the sulfoindocyanine Cy3 linked covalently to DNA. *J. Phys. Chem.*

B **111**, 11064-11074, (2007).

39. Peterson, E. M., Manhart, M. W. & Harris, J. M. Single-Molecule Fluorescence Imaging

of Interfacial DNA Hybridization Kinetics at Selective Capture Surfaces. *Anal. Chem.* **88**,

1345-1354, (2016).

a DNA-AuNP (100 nm diameter)
RNase H
RNA monolayer
Glass substrate
Laser (532 nm)
Scattering light

b t = 100 s   600 s   1100 s   1600 s   2100 s

c Start   End   Time (s)

d Data / Pause / Step   **100 nm**

e

f Pause length : τ_i
Step size : Δ_i
Speed : Δ_i / τ_i

Experiment

a
Pause length (s)

b
Step size (nm)

c
Speed (nm s⁻¹)

[RNase H] (nM)

d
DNA-AuNP

Accessible radius : R_{acc}

RNA monolayer

e
$k_{on}^{DNA/RNA}$ — Unbound
k_{on}^{E}[RNase H] — Bound
k_{cat}^{E} — Hydrolyzed
Empty

f
[RNase H] = 36 nM
Start
End
Empty
Hydrolyzed
Unbound
100 nm

g
[RNase H] = 36 nM
Start
End
Time (s)
2500
0
100 nm

Simulation

h
Pause length (s)

i
Step size (nm)

j
Speed (nm s⁻¹)

[RNase H] (nM)

a [RNase H]
- 36 nM
- 144 nM
- 360 nM
- 720 nM
- 1440 nM
- 3600 nM
- 7200 nM

c Processivity = τ_{detach} / <Pause length>
Run-length = Processivity × <Step size>

Attach → Pause → Step → Pause → ⋯ → Pause → Step → Pause → Detach

τ_{detach}

a

Fluorescence / Time

Cy3 + DNA $\xrightleftharpoons[k_{off}^{DNA/RNA}]{k_{on}^{DNA/RNA}}$

RNA — DNA

DNA

b

RNase H

$\xrightarrow{k_{on}^{E}[\text{RNase H}]} \xrightarrow{k_{cat}^{E}}$

Original
DNA: 3′-GACACTAACTAATGA-5′
RNA: 5′-C$_5$-CUGUGAUUGAUUACU-3′-Cy3

c

Normalized Fluorescence (a.u.)

[DNA]$_0$ = 6 nM
[DNA]$_0$ = 12 nM
[DNA]$_0$ = 25 nM
[DNA]$_0$ = 50 nM
[DNA]$_0$ = 67 nM
[DNA]$_0$ = 100 nM

Time (s)

d

Hybridization rate (s^{-1})

R^2 = 0.97

[DNA]$_0$ (nM)

e

Normalized Fluorescence (a.u.)

[RNase H]$_0$ = 69 nM
[RNase H]$_0$ = 173 nM
[RNase H]$_0$ = 346 nM
[RNase H]$_0$ = 864 nM
[RNase H]$_0$ = 3456 nM
[RNase H]$_0$ = 6912 nM

Time (s)

f

RNA hydrolysis rate (s^{-1})

[RNase H]$_0$ (nM)

a

Processivity

[RNase H] (nM)

b

Run length (nm)

[RNase H] (nM)

c Entrap detachment

[RNase H] = 720 nM

- Empty
- Hydrolyzed
- Unbound

Start

End

100 nm

50 nm

d Stochastic detachment

[RNase H] = 720 nM

End

Start

100 nm

50 nm

e

Unbound

Hydrolyzed

Empty

R_{acc} R_{acc}

End

$$P_{unbound} = \frac{N_{unbound}\,(R_{acc} < r < 2R_{acc})}{N_{unbound}\,(R_{acc} < r < 2R_{acc}) + N_{hydrolyzed}\,(R_{acc} < r < 2R_{acc})}$$

f

$P_{unbound}$

[RNase H] (nM)

Stochastic detachment

Entrap detachment

Newly-designed

DNA: 3′-CACACACACACACAC -5′
RNA: 5′-C$_5$-GUGUGUGUGUGUGUG -3′-Cy3

a

[DNA]$_0$ = 6 nM

[DNA]$_0$ = 12 nM

[DNA]$_0$ = 25 nM

[DNA]$_0$ = 50 nM

[DNA]$_0$ = 67 nM

[DNA]$_0$ = 100 nM

c

[RNase H]$_0$ = 69 nM

[RNase H]$_0$ = 173 nM

[RNase H]$_0$ = 346 nM

[RNase H]$_0$ = 864 nM

[RNase H]$_0$ = 3456 nM

[RNase H]$_0$ = 6912 nM

b
R^2 = 0.95

Original
Newly-designed

d

Original
Newly-designed

e $k_{on}^{DNA/RNA}$ (10^6 M^{-1} s^{-1})

f $k_{off}^{DNA/RNA}$ (10^{-3} s^{-1})

g k_{on}^E (10^6 M^{-1} s^{-1})

h k_{cat}^E (s^{-1})