1   Early-Life Gut Inflammation Drives Sex-Dependent Shifts in the
2   Microbiome-Endocrine-Brain Axis
3

Olivia Sullivan[1,2, †], Claire Sie[3,†], Katharine M. Ng[3], Sophie Cotton[3], Cal Rosete[2], Jordan E.
Hamden[2,4], Ajay Paul Singh[5], Kristen Lee[2,4], Jatin Choudhary[2], Jennifer Kim[1,2], Huaxu Yu[6],
Charlotte A. Clayton[3], Natalia A. Carranza Garcia[3], Kateryna Voznyuk[2], Brian D. Deng[3], Nadine
Plett[2], Sana Arora[2], Hans Ghezzi[7], Tao Huan[6], Kiran K. Soma[1,2,8], John-Paul J. Yu[5], Carolina
Tropini[3,9,10]*, Annie Vogel Ciernia[1,2,4]*

[1]Graduate Program in Neuroscience, University of British Columbia, Vancouver, Canada
[2]Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver,
Canada
[3]Department of Microbiology & Immunology, University of British Columbia, Vancouver,
Canada
[4]Department of Biochemistry and Molecular Biology, University of British Columbia,
Vancouver, Canada
[5]Department of Radiology, University of Wisconsin School of Medicine and Public Health,
Madison, Wisconsin, USA.
[6]Department of Chemistry, University of British Columbia, Vancouver, Canada
[7]Department of Bioinformatics, University of British Columbia, Vancouver, Canada
[8]Department of Psychology, University of British Columbia, Vancouver Canada
[9]School of Biomedical Engineering, University of British Columbia, Vancouver, Canada
[10]Humans and the Microbiome Program, Canadian Institute for Advanced Research (CIFAR),
Toronto, Canada
24

†Co-first authors

*Corresponding Authors:

Carolina Tropini

Life Sciences Centre, Rm 3557
2350 Health Sciences Mall
Vancouver, British Columbia V6T 1Z3

carolina.tropini@ubc.ca

Annie Vogel Ciernia
Centre for Brain Health
2215 Wesbrook Mall
Vancouver, BC V6T2A1
annie.ciernia@ubc.ca

CRediT authorship contribution statement:

OS: Methodology, Formal Analysis, Investigation, Data Curation, Writing - Original Draft, Writing - Review & Editing, Visualization
CS: Methodology, Formal Analysis, Investigation, Writing - Original Draft, Writing - Review & Editing, Visualization
KN: Methodology, Formal Analysis, Investigation, Data Curation
SC: Investigation, Data Curation, Writing - Review & Editing
CR: Methodology, Formal Analysis, Investigation, Data Curation, Writing - Original Draft, Writing - Review & Editing
JEH: Investigation, Formal Analysis, Writing - Original Draft, Writing - Review & Editing
APS: Investigation, Formal Analysis
KL: Methodology, Investigation
JC: Investigation, Formal Analysis
JK: Investigation, Writing – Review & Editing
HY: Investigation, Formal Analysis
CAC: Investigation, Data Curation
NACG: Methodology
BDD: Investigation
NP: Formal Analysis
KV: Investigation
SA: Investigation
HG: Investigation, Writing – Review & Editing
TH: Formal Analysis
KS: Methodology, Writing - Review & Editing
JP: Resources, Writing – original draft, Writing– review & editing, Supervision, Project administration, Funding acquisition
CT: Conceptualization, Resources, Writing – original draft, Writing– review & editing, Supervision, Project administration, Funding acquisition.
AVC: Conceptualization, Resources, Visualization, Writing – original draft, Formal Analysis, Writing– review & editing, Supervision, Project administration, Funding acquisition.

Declarations of interest: none

Word count: 13,787

Keywords: gut-brain axis, inflammatory bowel disease, gut inflammation, microglia, androgens, sexual behaviours, microbiome, development

Highlights

Early-life gut inflammation produces sex-specific effects on i) microbiome, ii) sex hormones and iii) behaviour.

Both sexes show disrupted gut bacterial members that regulate sex hormone levels.

Male mice demonstrate deficits in mate seeking, which may be mediated by reduced androgen levels.

80
81   Both male and female mice demonstrate shifts in hippocampal microglial morphology.
82
83

Abstract

Despite recent advances in understanding the connection between the gut microbiota and the
adult brain, there remains a wide knowledge gap in how gut inflammation impacts brain
development. We hypothesized that intestinal inflammation in early life would negatively affect
neurodevelopment through dysregulation of microbiota communication to the brain. We
therefore developed a novel pediatric chemical model of inflammatory bowel disease (IBD), an
incurable condition affecting millions of people worldwide. IBD is characterized by chronic
intestinal inflammation, and has comorbid symptoms of anxiety, depression and cognitive
impairment. Significantly, 25% of patients with IBD are diagnosed during childhood, and the
effect of chronic inflammation during this critical period of development is largely unknown.
This study investigated the effects of early-life gut inflammation induced by DSS (dextran
sulfate sodium) on a range of microbiota, endocrine, and behavioral outcomes, focusing on sex-
specific impacts. DSS-treated mice exhibited increased intestinal inflammation, altered
microbiota membership, and changes in microbiota-mediated circulating metabolites. The
majority of behavioral measures were unaffected, with the exception of impaired mate-seeking
behaviors in DSS-treated males. DSS-treated males also showed significantly smaller
seminal vesicles, lower circulating androgens, and decreased intestinal hormone-activating
enzyme activity. In the brain, microglia morphology was chronically altered with DSS treatment
in a sex-specific manner. The results suggest that early-life gut inflammation causes changes in
gut microbiota composition, affecting short-chain fatty acid (SCFA) producers and
glucuronidase (GUS) activity, correlating with altered SCFA and androgen levels. The findings
emphasize the developmental sensitivity to inflammation-induced changes in endocrine
signalling and underscore long-lasting physiological and microbiome changes associated with
juvenile IBD.

108
Introduction

Inflammatory bowel disease (IBD) is a group of chronic gastrointestinal (GI) tract conditions
that affects over 6.8 million people globally and remains incurable (Alatab et al., 2020). The risk
of IBD is correlated with industrialization and, as such, is globally rising in prevalence (Kaplan
and Windsor, 2021). Genetic factors account for less than one-third of IBD risk, underscoring the
importance of environmental factors such as gut microbiota composition, dietary habits,
smoking, and stress levels on disease development (Ananthakrishnan, 2013; Ye et al., 2015).
Individuals with IBD frequently undergo cycles of intense inflammatory flare-ups and remission
periods (Roda et al., 2020). Over 25% of patients with IBD are diagnosed during childhood or
adolescence with symptoms that persist through adulthood (Däbritz et al., 2017). Importantly,
little is known about the specific impacts of pediatric-onset IBD on the developing brain.
Children and adolescents with IBD have an increased risk of developing anxiety and depression,
experience a lower quality of life, and face educational and social challenges along with mild
cognitive impairments (Attree et al., 2003; Mackner et al., 2013; Thavamani et al., 2019).
Furthermore, neuroimaging studies in adult female and male patients with IBD have revealed

altered brain structure and function (Bao et al., 2015; Lv et al., 2017) some of which correlate
with neuropsychiatric impairments (Hou et al., 2020), highlighting the need for similar research
in pediatric populations.

Research into inflammatory phenotypes through various animal models – including genetic,
chemically induced, and adoptive transfer T cell models – has significantly advanced our
understanding of IBD (Wirtz and Neurath, 2007). The widely-used dextran sodium sulfate
(DSS)-induced colitis model damages the gut epithelial lining and mimics the gut inflammation
and microbiota changes seen in human ulcerative colitis (UC), one of the two main types of IBD
(Chassaing et al., 2014). Adult mice treated with DSS have increased anxiety (Nyuyki et al.,
2018) and reduced sucrose preference, indicative of a depression endophenotype (Zhou et al.,
2023). Additionally, active gut inflammation in adult mice led to anxiety-like behavior and
recognition memory impairments (Emge et al., 2016) and social impairments (Brown et al.,
2024). Importantly, anxiety and memory deficits were reversed upon resolution of acute
inflammation (Emge et al., 2016), which is encouraging for therapeutic resolution of
neurophysiological symptons in humans with IBD. While DSS-colitis studies predominantly
focus on adult rodents and capture many of the extraintestinal manifestations of IBD seen in
adult human patients, research in juvenile rodent models remains limited. In contrast to adult
mice, a recent DSS-colitis model in juvenile mice revealed persistent cognitive deficits and
anxiety-like behaviors even upon GI-symptom resolution (Salvo et al., 2020). These observations
suggest that GI inflammation during developmental periods may have unique and long-lasting
effects on brain development, leading to distinct behavioral impacts in adulthood.

How GI inflammation triggers brain and behavior dysfunction is not well understood, but is
theorized to involve altered signaling along the gut-brain-axis. The gut microbiota can signal to
the brain by immune cell trafficking, peripheral nerve activity, and release of metabolites that
circulate to the brain (Banfi et al., 2021; Günther et al., 2021). One of the key brain cells
responsive to gut microbe derived metabolites are microglia, the brain's resident innate immune
cells. Microglia sculpt synaptic physiology during brain development (Sullivan and Ciernia,
2022), are acutely sensitive to changes in microbiota in early life (Erny et al., 2015; Thion et al.,
2017) and are chronically functionally altered from early life inflammation (Vogel Ciernia et al.,
2018). Microbiota-derived short chain fatty acids (SCFAs) can enter the blood stream and cross
the blood brain barrier to impact microglia development and function (Cryan and Dinan, 2012;
Erny et al., 2015; Sullivan and Ciernia, 2022). In both adult and juvenile IBD mouse models,
microglia adopt a pro-inflammatory phenotype characterized by altered cellular morphology and
increased pro-inflammatory cytokine secretion (Caetano-Silva et al., 2024; Gampierakis et al.,
2021; Salvo et al., 2020; Sroor et al., 2019; Talley et al., 2021). Together, this suggests that early
life changes to microbiota-derived signalling to brain microglia may negatively impact brain
development in IBD.

Beyond impacting brain function and behavior, pediatric IBD poses several unique challenges in
other aspects of development, including delayed onset of puberty in both males and females
(Ballinger et al., 2003). While undernutrition was inititally thought to be the main driver of
delayed puberty, restoring nutritional status fails to normalize the timing of puberty onset,
suggesting the involvement of additional factors (Azooz et al., 2001; Brain and Savage, 1994;
DeBoer et al., 2010; Deboer and Li, 2011). Animal models of IBD have linked intestinal
inflammation to disruption in testosterone levels, estrogen receptor function, and reproductive

organ size, with effects that extend beyond what is seen with undernutrition alone (Azooz et al.,
2001; Ballinger et al., 2003; DeBoer et al., 2010; Deboer and Li, 2011). Similarly, findings in
human studies show that some patients with IBD (Klaus et al., 2008; Tigas and Tsatsoulis, 2012)
and irritable bowel syndrome (Rastelli et al., 2022) have disrupted sex hormone levels and higher
rates of sexual dysfunction (Chen et al., 2024; Gaidos et al., 2020). One connection between
inflammation and altered hormone levels is thought to be mediated by the gut microbiota, which
acts as a regulator of estrogen and androgen synthesis and metabolism (Colldén et al., 2019; He
et al., 2021; Maffei et al., 2022; Neuman et al., 2015). In mouse models of IBD, the loss of
bacterial species such as *Faecalibacterium prausnitzii* (Sokol et al., 2008) and *Roseburia*
hominis, known for producing anti-inflammatory SCFAs, is associated with changes in sex
hormone levels (Sui et al., 2021; Torres et al., 2018). The depletion of specific bacteria such as
F. prausnitzii in IBD, which are crucial for converting estrogen to its active form, can directly
impact hormone availability, potentially affecting the onset and progression of sexual
development (Ervin et al., 2019). Additionally, the recent discovery of testosterone-degrading
bacteria such as *Mycobacterium* contributing to inflammation in IBD underscores the
microbiota's profound impact on sexual development, suggesting avenues for targeted microbial
interventions to mitigate impairments in IBD (Li et al., 2022; Naser et al., 2014). Altogether,
these studies highlight the complex relationships between the gut microbiota and sexual
development (Lavelle and Sokol, 2020; Machiels et al., 2014).

There is a critical gap in understanding the interactions between early life intestinal
inflammation, brain function, and reproductive development. To bridge this gap, here we
introduce a mouse model of pediatric-onset IBD that captures recurrent episodes of gut
inflammation throughout childhood and adolescence, modeling the human condition. In both
sexes, we assessed the acute and chronic effects of early-life gut inflammation on gut microbiota
composition, short-chain fatty acid production, a wide variety of behaviors, sex steroid levels,
brain white matter tract integrity and microglial morphology, offering a comprehensive view of
IBD's impact on juvenile development. Our findings reveal novel sex-specific changes in
reproductive behavior, alterations in microbiota metabolites and hormone signaling, and
microglial morphology adaptations. These results highlight the unique long-term consequences
for human patients with early-onset IBD, emphasizing a need for early diagnosis and
microbiome-targeted intervention strategies.

Methods

Animals

Experimentally naïve 3-week-old C57BL/6J wildtype mice (strain #000664, Jackson
Laboratories) were housed in groups of 3-5 with sex and treatment-matched cage mates. Mice
were housed in ventilated cages under a 12:12 light-dark cycle with lights on at 0700h and lights
off at 1900h in a temperature (22ºC) and humidity (40-60%) controlled environment. Purina
LabDiet 5k67 diet and reverse osmosis chlorinated water were given *ad libitum* throughout the
experiment unless stated otherwise for specific tasks. All housing conditions and testing
procedures were in accordance with the guidelines of the Canadian Council on Animal Care, and
all protocols were approved by the Animal Care Committee of the University of British
Columbia *(protocols A19-0078, A23-0115, and A23-0086).*

Pediatric IBD mouse model

Mice randomly assigned to the treatment groups were administered three rounds of 5-day
treatments of 3-3.5% colitis-grade DSS (36-50 kDa, MP Biomedicals #160110) while the vehicle
control group (VEH) received water only. Due to the reported batch-dependent effects of DSS,
the optimal concentration of DSS to cause colitis was determined in preceding pilot experiments
(Eichele and Kharbanda, 2017). The treatment periods occurred during postnatal days (P) P21-
26, P42-47, and P63-68 (Figure 1A). During treatment and for three days following each
treatment, mice were weighed daily to monitor for weight loss (Supplemental Table 2). Regular
monitoring of weight was continued during the recovery period intervals. One cohort of mice
was euthanized three days after the final DSS treatment on P71 (n=4-5/treatment/sex). The
remaining two cohorts of mice underwent behavioral testing as young adults between P85-P130
(n=11-13/treatment/sex).

Serum and organ collection

Animals were deeply anesthetized with 5% isoflurane, and blood was collected by cardiac
puncture. Whole blood was collected into BD Microtainer™ Serum Separator Tubes (BD) and
allowed to coagulate for at least 30 minutes at room temperature according to manufacturer
instructions. To isolate serum for later analyses, blood samples were spun down at 1200 x g for
20 minutes at 4ºC, and the serum was removed and placed in a new tube for storage at -80ºC.
Animals were then transcardially perfused with 1X phosphate buffered saline (PBS). Intestinal
measures were conducted as follows: first, intestines were dissected out from the gastroduodenal
junction to the rectum. Colon length was then measured from the ileocecal junction to the
rectum. The contents of the cecum were gently scraped and collected into a sterile tube, then
immediately flash-frozen on dry ice. The colon was manipulated into rolls in histological
cassettes and immediately fixed in freshly prepared Methacarn solution (60% dry methanol, 30%
chloroform, 10% glacial acetic acid) (Ng and Tropini, 2021). Brains were collected and drop
fixed in 4% paraformaldehyde (PFA) for 24 hours at 4°C and then transferred to 30% sucrose
solution and kept at 4°C until sectioned. For brain connectivity analysis, mice were perfused
with PBS followed by 4% PFA and brains were dissected out and then stored in 4% PFA until
scanning.

Gut staining and histological scoring

Within 2 weeks of fixation in Methacarn solution, intestinal tissues were processed and
infiltrated with paraffin wax as previously described (Ng and Tropini, 2021). Briefly, tissues
were washed twice with absolute methanol for 30 minutes each, followed by two washes with
absolute ethanol for 20 minutes each. Next, tissues were washed twice with xylene for 15
minutes before being placed in melted paraffin wax and incubated for 2 hours at 60ºC. Samples
were then sent to the BC Children's Hospital Histology Core for embedding in paraffin wax and
sectioning into 4 – 5 μm sections. Histological staining was performed with hematoxylin and
eosin as previously described (Stahl et al., 2017). Finally, slides were imaged using a
PANNORAMIC® MIDI II (3DHISTECH).

Mucus and bacteria visualization

Visualization of gut microbes was performed as previously described (Ng and Tropini, 2021).
Briefly, to deparaffinize tissues, slides were heated at 60°C in Coplin jars for 10 minutes,
followed by incubation in xylene that was pre-warmed to 60°C. A second xylene wash was
performed at room temperature for 10 minutes, then slides were placed in 99.5% ethanol for 5
minutes. After allowing slides to air-dry, tissues were encircled using a liquid blocker PAP pen.
FISH probe hybridization solution (20 mM Tris-HCl, pH 7.4; 0.9 M NaCl; 0.1% sodium
dodecylsulfate in nuclease-free water) was pre-warmed to 50°C before addition of Eub-388 (5'-
AF488-GCTGCCTCCCGTAGGAGT-3') to a final concentration of 10 ng/µL (Amann et al.,
1990). Slides were then incubated for 3 hours at 50°C in a humid chamber. Following incubation,
tissues were washed once with FISH washing buffer (20 mM Tris-HCl, pH 7.4; 0.9 M NaCl, pre-
warmed to 50°C) for 10 minutes at 50°C, then additionally with 1x PBS. DNA and intestinal
mucus were counter-stained using 10 µg/mL DAPI and 40 µg/mL UEA-1-Rhodamine Red
respectively in PBS for 45 minutes at 4°C. Finally, slides were washed thrice with PBS before
mounting with ProLong Gold AntiFade mounting medium (Invitrogen). Fluorescence images
were captured using a Zeiss LSM900 confocal microscope using a 100x objective lens.

Microbiome sequencing and analysis

Microbiome composition was determined via sequencing of the 16S rRNA hypervariable region.
Total DNA was extracted from 30-60 mg of cecal contents using the DNeasy PowerSoil Pro kit
(Qiagen) in accordance with manufacturer protocols. Quantification of extracted DNA
concentration prior to library preparation was performed using the Quant-iT™ dsDNA HS Assay
(Thermo Fisher). 16S library preparation was conducted at either the Gut4Health facility at BC
Children's Hospital or the Biofactorial High-Throughput Biology (Bio!) Facility at the
University of British Columbia. For samples collected 3 days after cessation of DSS, library
preparation targeting the V4 hypervariable region of the 16S rRNA gene was performed at the
Gut4Health facility using established protocols (De Wolfe and Wright, 2023). Amplification of
the V4 region was performed with 515F/926R primers (515F, 5'-
GTGYCAGCMGCCGCGGTAA-3'; 926R, 5'-CCGYCAATTYMTTTRAGTTT-3'). Pooled
libraries were then submitted to the Bio! facility, where sequencing was performed on the
Illumina MiSeq™ platform with v2 2 x 250 bp paired-end read chemistry. For all other samples,
library preparation targeting the V4-V5 16S rRNA genes was performed at the Bio! Facility as
previously described (Ng et al., 2023), and sequencing was run on the Illumina MiSeq™
platform with v3 2 x 300 bp paired-end read chemistry. FASTQC (Andrews, 2010) was run on
the generated FASTQ files to assess read quality. Reads were then imported into QIIME2-2023.2
for subsequent analyses (Bolyen et al., 2019). Denoising and quality filtering was performed
using DADA2 (via q2-dada2), and reads were trimmed to remove primer sequences while
maintaining mean Phred quality scores >Q30 (Callahan et al., 2016). Multiple sequence
alignment and phylogenetic tree generation was performed using MAFFT (via q2-alignment) and
FastTree2 respectively (via q2-phylogeny) (Katoh et al., 2002; Price et al., 2010). Using the
QIIME classification plugin (q2-feature-classifier), amplicon sequence variants (ASVs) were
classified via a naïve Bayes machine-learning taxonomic classifier against the SILVA 138 99%
identity reference sequence database (Quast et al., 2013).

Further analyses were conducted using R v4.2.2 (R Core team, 2021). Analyses were visualized
through the tidyverse (Wickham et al., 2019) and ggplot2 (Hadley, 2016) packages. For sample
rarefaction, calculation, and visualization of alpha and beta diversity metrics, the phyloseq

(McMurdie and Holmes, 2013), ggpubr (Kassambara, 2023), and vegan (Dixon, 2003) packages
were employed. Differential abundance analysis (DAA) was conducted using the
metagenomeseq package (Paulson et al., 2013) and Analysis of Composition of Microbiomes
(ANCOM) (Mandal et al., 2015) packages through the microbiomeMarker package (Cao et al.,
2022). The relative abundance of those taxa identified as significantly differentially abundant
were then plotted with FDR-corrected p-values obtained with metagenomeseq. Differentially
abundant taxa were plotted using the ComplexHeatmap (Gu et al., 2016) package. The relative
abundance of select differentially abundant taxa was then plotted against serum hormone and
SCFA concentrations, and Spearman's correlation was calculated using smplot2 (Min and Zhou,
2021).

Behavior testing

All behavior testing was performed under white light (600 lux) between the hours of 0900-1800
except for the sucrose preference test, which is a 24-hour task. Animals were habituated to the
researchers after the anxiety tasks but prior to all other tests through 3 days of handling (1-2
min/mouse). For each behavior task, animals were first habituated to room conditions for 30
minutes before testing. For all tasks that did not involve bedding, the apparatus was cleaned with
diluted saber solution (Vert-2-Go saber diluted 1:16 in water) between mice and between stages.
ANY-maze tracking software (Stoelting) was used to record videos and track animal movement
during the task. Mice began testing 17 days following the final round of DSS with the exception
of a subset cohort, which underwent only the urine preference test and mating behavior test
beginning 28 days following the final round of DSS treatment.

Urine Preference Test (UPT)
Mice were placed in the centre of an open-top box (40 cm x 40 cm x 40 cm, L x W x H)
containing two 10 cm diameter inverted pencil cups in opposite corners. A petri dish with
blotting paper (1 cm^2) was placed under each pencil cup. Mice were given a 10-minute
habituation period where 20 µL of deionized water was dropped onto both blotting papers.
Following habituation, mice were given another 10-minutes to explore the two cups, this time
with 20 µL of pooled male or female C57BL/6 mouse urine dropped onto one of the blotting
papers and 20 µL of deionized water dropped onto the other. In the final stage, mice were given
another 10-minutes to explore the two cups, with the urine cup location and urine sex switching
from the previous stage (Malkesman et al., 2010). Presentation order of same or opposite sex
urine as well as location of urine in the first and second stages was counterbalanced across mice
and treatment conditions. ANY-maze tracked the total investigation time when the test mouse's
nose was within 1 cm of the inverted pencil cup. Exclusion criteria for data collected during the
urine preference test include side preference during habituation (defined as a discrimination
index (DI) over 20 or under -20, where DI is the difference in time spent investigating the
stimulus cups divided by the total time spent investigating) and side preference exceeding three
standard deviations from the mean.

Mounting Behavior
Sexually inexperienced VEH- and DSS-treated male mice were paired with a novel sexually
experienced C57BL/6 female mouse in an empty 35 cm x 20 cm x 15 cm (L x W x H) clear
plastic box. ANY-maze (Stoelting) was used to record a 15-minute free interaction period.
Videos were hand-scored for the frequency and duration of male mounting by a researcher blind

to treatment conditions. Mounting behavior was defined as male contact with two paws on the
front or back of the female as previously described (Bayless et al., 2023).

Open Field Test
The open field was performed in an open-topped grey Plexiglass arena (65 cm x 65 cm x 40 cm;
L x W x H). Mice were placed in the centre of the arena and given 20 minutes to freely explore.
ANY-maze tracked the total distance travelled and the amount of time spent in the inner and
outer zones. The outer zone was defined at 8 cm from the wall of the apparatus.

Elevated Plus Maze
The plus-shaped apparatus was elevated 50 cm above the floor and had two open arms and two
closed arms, both 30 cm x 8 cm (L x W). The closed arms were surrounded by 15 cm high walls.
Mice were placed in the centre of the apparatus and allowed to freely explore for 5 minutes.
ANY-maze tracked the total distance travelled and the amount of time spent in the open arms
and closed arms.

Three Chamber Test
The test apparatus was an open-topped rectangular box (60 cm x 40 cm x 22 cm ; L x W x H)
divided lengthwise into three identically sized compartments (Yang et al., 2011). The task
consisted of three 10-minute phases: habituation, sociability, and social novelty. The sociability
and social novelty phases require sex and age matched stimulus mice. The novel mouse was not
a cage mate of the familiar mouse in the social novelty phase. Prior to the task, stimulus mice
spent 20 minutes/day for 3 days habituating to sitting under inverted pencil cups.

ANY-maze was used to track distance travelled, time in each chamber, and time spent
investigating the cup in each phase of the task. When the mouse's nose was within 1 cm of the
pencil cup and pointing directly at it, investigation was scored. Looking above or lateral to the
cup and climbing on top of the object were not counted as investigation.

Olfaction Habituation-Dishabituation Test
The olfaction test was performed as previously described (Yang and Crawley, 2009). Briefly,
mice were habituated to a clean cage without bedding for 30 minutes prior to being repeatedly
presented a cotton swab with non-social (vanilla or banana extract diluted 1:1000 in water) and
social odours (swab of cage bottom of novel mouse). The order of odour presentation was: water,
vanilla extract, banana extract, same sex mice, opposite sex mice. Videos were recorded and
hand-scored by researchers blinded to experimental conditions for the time spent investigating
each odoured cotton swab, defined as the nose within 1 cm of the swab. Biting and otherwise
playing with the cotton swab was not counted as olfaction investigation.

Object Location Memory (OLM) Test
The OLM test was performed in an open-topped Plexiglass square box (40 cm x 40 cm x 40 cm ;
L x W x H) lined with bedding ~1 cm in depth. Bedding was kept in the same box each day. One
wall of the box was lined with duct tape as a visual cue. The OLM took 8 days in total with 6
days of habituation, 1 day of training and 1 day of testing (Vogel-Ciernia and Wood, 2014).
Identical cylindrical spice tins filled with cement were used as training objects and were cleaned
with diluted saber solution between trials and allowed to fully dry before re-use. The 5-minute

testing phase took place 24 hours after training where one object was moved to a novel location
and the other remained in the original training location.

ANY-maze tracked the total distance travelled on all 8 days. ANY-maze tracked investigation
time and the number of investigations of each object on training and testing days. Investigation
time was scored when the mouse's nose was within 1 cm of the object and pointing directly at
the object. Looking over or past the object and climbing on top of the object were not counted as
investigation. For the training and testing days, a discrimination index (DI) was calculated where
DI is the difference in time spent investigating the moved object and the unmoved object divided
by the total time spent investigating both objects x100.

Mice who had a DI over 20 or under -20 during the training phase were determined as having an
innate preference and excluded from final analysis. Mice who explored the objects more or less
than 3 standard deviations from the mean were also excluded from analysis as we have done
previously (Vogel-Ciernia and Wood, 2014).

Self-Grooming and Forced Grooming
Mice were individually placed in a new cage without bedding. After a 5 minute habituation
period, the cumulative time spent naturally self-grooming was recorded for 5 minutes. The mice
were then lightly sprayed with water from a mist bottle and forced self-grooming was scored in
the following 5 minutes. All hand scoring of grooming behavior was performed by a researcher
blinded to experimental conditions.

Forced Swim Test
Mice were placed individually into 1 L glass beakers filled with 900 mL of water at a
temperature of 23-25°C. Mice were recorded for 6 minutes and then removed from the water as
previously described(Can et al., 2011). The last 4 minutes of footage was hand-scored for time
spent immobile, defined as the absence of escape-related movements. The hand-scorer was
blinded to experimental conditions.

Sex steroid quantification

Serum (10 µl) was pipetted into 2 mL polypropylene bead ruptor tubes with 5 ceramic zirconium
oxide beads (Hamden et al., 2021, 2019). Calibration curves (0.05 – 1000 pg), blanks, and
quality controls were extracted alongside samples via liquid-liquid extraction and analyzed by
specific and ultrasensitive LC-MS/MS as before (Hamden et al., 2021). Progesterone,
testosterone (T), androstenedione (AE), 5α-dihydrotestosterone (DHT), and 17β-estradiol (E_2)
were measured using multiple reaction monitoring with 2 mass transitions for each analyte.
Deuterated internal standards (progesterone-d9, testosterone-d5, and 17β-estradiol-d4) were
included in all standards, blanks, controls, and samples to correct for analyte loss and matrix
effects. Steroid concentrations were acquired by a Sciex 6500 Qtrap triple quadrupole tandem
mass spectrometer. A steroid was considered non-detectable if either the quantifier or qualifier
transition was not present. If 50% or more of the values in a group were detectable, then missing
values were replaced with the lower limit of quantification∕√2(Handelsman and Ly, 2019)
(Handelsman and Ly 2019). DHT and E_2 were non-detectable in all samples, and AE was non-
detectable in all female samples.

Brain microglia immunofluorescence staining

After PFA fixation and sucrose cryoprotection, brain hemispheres were embedded in Optimal
Cutting Temperature (OCT) compound (Scigen, 23-730-625), cryosectioned at 30 μm, and
stored in 1X PBS at 4°C. For staining, free-floating sections were permeabilized with 1X PBS +
0.5% Triton X-100 for 5 minutes and then washed 3 x 5 minutes with 1X PBS. Sections were
placed in a blocking solution (5% normal donkey serum, and 1% bovine serum albumin
dissolved in 1X PBS + 0.03% Triton) for 1 hour at room temperature with gentle shaking and
then incubated with anti-IBA1 primary antibody (chicken; Synaptic Systems (234009); 1:1000)
for 48 hours. Sections were washed 3 x 5 minutes in 1X PBS + 0.03% Triton and then incubated
in a donkey anti-chicken secondary antibody (Alexa Fluor® 488 Jackson ImmunoResearch
(703545185); 1:500) and DAPI (Biolegend (422801); 1:1000) for two hours at room temperature
on a gentle tilt. Sections were washed 3 x 5 minutes in 1X PBS + 0.03% Triton and 1x 5 minutes
in 1X PBS and then mounted onto glass slides and coverslipped using Invitrogen ProLong Glass
Antifade Mountant (Invitrogen, P36984).

Reverse transcription quantitative polymerase chain reaction
At time of euthanasia, the hippocampus tissue was dissected, snap frozen on dry ice and stored at
-80ºC until RNA extraction. Total RNA was isolated and water blanks using a Monarch® Total
RNA Miniprep Kit (New England Biolabs, Whitby, ON) and RNA quantity and quality were
confirmed by Nanodrop spectrophotometer. Complementary DNA (cDNA) was reverse
transcribed using a LunaScript RT Supermix Kit (New England Biolabs, Ipswich, MA). The
amount of RNA used for reverse transcription was normalized (227 – 400ng) and experimenters
were blinded to treatment group and sex during RNA isolation and cDNA synthesis.
Transcripts (*Tlr4, Nod1, Nod2, Il1β, Tnfα, Il6*) were quantified via SYBR Green assays
(Integrated DNA Technologies, Inc., Coralville, IA) (Table 1). Relative expression levels of the
reference gene and genes of interest were assessed using Luna® Universal qPCR Master Mix
(New England Biolabs, Whitby, ON). Samples were run in duplicate. A reference gene (*Hprt*)
was measured in each sample. Negative controls (water blanks, no reverse transcriptase controls)
confirmed the specificity of assays. Melt curve analyses verified that the expected targets were
amplified. One sample was excluded from the Male DSS *Nod1* analysis due to lack of specificity
in the melt curve.

Gene	Forward	Reverse
Tlr4	GAAGCTTGAATCCCTGCATAG	AGCTCAGATCTATGTTCTTGGTTG
Il1β	TGGCAACTGTTCCTGAACTCA	GGGTCCGTCAACTTCAAAGAAC
Tnfα	GGGTGATCGGTCCCCAAA	TGAGGGTCTGGGCCATAGAA
Il6	CGATGATGCACTTGCAGAAA	ACTCCAGAAGACCAGAGGAA
Nod1	TTCCTCACATAGCACCTTCAC	TCAATGACTATGGCGTGCAG
Nod2	CGAAGCCAAACCTCCAGAA	TCGCAGAAGGACTCAAGAGA
Hprt	CAGTACAGCCCCAAAATGGTTA	AGTCTGGCCTGTATCCAACA

Table 1. RT-qPCR primers.
The cycle threshold (Ct) of the reference gene *Hprt* for each treatment group and sex were
analyzed by two-tailed t-test to determine whether expression of reference gene differed between
VEH and LPS groups. There was no significant difference between VEH and DSS groups for the
females nor males (p = .76 and .31, respectively). The mean of the reference gene was used to

452 calculate the ΔCq and ΔCq values were normalized to the VEH group within each sex. Statistics
453 were run and data were graphed using the fold change for each transcript within each sex. Data
454 were analyzed for an effect of treatment by unpaired t-tests.

Microglial morphology analysis

Slides were imaged using the Zeiss Axioscan7 slide scanner microscope at 20X magnification.
Maximum intensity extended depth of focus (EDF) images were acquired from Z-stacks
containing 16 slices at a step size of 1 μm. EDF images were created using maximum projection
settings and converted to tiffs using ZEN (Zeiss). Maximum projection EDF images compile the
pixels of highest intensity at any given z-stack position and construct a new 2D image retaining
the 3D information. The hippocampus and hypothalamus anatomical regions were traced using
the FASTMAP FIJI plugin with reference to the Allen Brain Atlas (Terstege et al., 2022). Within
these segmented regions, microglial morphology was analyzed using the pipeline described in
Kim et al., 2024 using MicrogliaMorphology (ImageJ tool) and MicrogliaMorphologyR (R package)
(Kim et al., 2024a) which uses Image-J plugins Skeletonize (2D/3D) and AnalyzeSkeleton (2D/3D)
(Arganda-Carreras et al., 2010) to collect 27 metrics of morphology per microglial cell
(Supplemental Figure 4A). We conducted dimensionality reduction using principal component analysis
followed by k-means clustering on the first three principal components (Supplemental Figure 4C) to
define four morphological classes within our dataset (hypertrophic, rod-like, ramified, ameboid). We
identified the optimal clustering parameter using exploratory data analysis methods including the within
sum of squares and silhouette methods (Supplemental Figure 4B). We assigned cluster identities by
examining the relationship between each cluster and the 27 morphological features (Supplemental Figure
4D) and by using the ColorByCluster feature in MicrogliaMorphology to visually examine the different
cluster morphologies (Figure 7A).

We then calculated the percentage of cells in each morphology cluster for each treatment and sex using
the 'clusterpercentage' function within MicrogliaMorphologyR. To assess how cluster membership
changes with treatment for each sex, we fit a generalized linear mixed model using a beta distribution to
model the percentage of cluster membership as a factor of Cluster identity and DSS treatment with
MouseID as a repeated measure ("percentage ~ Cluster*treatment + (1|MouseID)") using the
`stats_cluster.animal` function from MicrogliaMorphologyR, which is wrapped around the glmmTMB R
package (Brooks et al., 2017). We ran a 2-way Analysis of Deviance (Type II Wald chisquare test) test on
the model to assess the contribution of Cluster and treatment interactions. Tests between treatment
conditions were corrected for multiple comparisons across Clusters (~treatment|Cluster) using the Sidak
method and q-values < 0.05 were considered statistically significant.

Short and medium chain fatty acid quantification

Serum samples were stored in -80 freezer prior to lipid extraction. A total of 50 μL of serum was
mixed with 450 μL ice-cold methanol, vortexed for 10 seconds, and incubated at -20 ºC for 4
hours for protein precipitation. Samples were then centrifuged at 14,000 rpm at 4 ºC for 15
minutes, and the supernatant were collected and dried down via SpeedVac at 4 ºC. The dried
extract was reconstituted in 100 μL of water and acetonitrile solution (1:1, v:v) for LC-MS
analysis. Quality control (QC) samples were prepared by pooling 20 μL aliquot from each
sample to monitor the instrument performance and optimize the injection volume. A method
blank sample was prepared using the identical analytical workflow but without serum. An UHR-

QqTOF (Ultra-High Resolution Qq-Time-Of-Flight) Impact II (Bruker Daltonics, Bremen, Germany) mass spectrometry interfaced with an Agilent 1290 Infinity II Ultrahigh-Performance Liquid Chromatography (UHPLC) system (Agilent Technologies, Santa Clara, CA, USA) was used for analysis. LC separation was achieved using a Waters reversed phase (RP) UPLC Acquity BEH C18 Column (1.7 μm, 1.0 mm ×100 mm, 130 Å) (Milford, MA, USA) maintained at 25 °C. The mobile phase A was 5% ACN in H_2O with 10 mM ammonium acetate (pH = 9.8, adjusted ammonium hydroxide) and the mobile phase B was 95% ACN in H_2O. The LC gradient was set as follow: 0 min, 5% B; 8 min, 40% B; 14 min, 70% B; 20 min, 95% B; 23 min, 95% B; 24 min, 5% B; 33 min, 5% B. The flow rate was 0.1 mL/min. Negative ion mode was applied with the following MS settings: dry gas temperature, 220 °C; dry gas flow, 7 L/min; nebulizer gas pressure, 1.6 bar; capillary voltage, 3000 V. External calibration was applied using sodium formate to ensure the *m/z* accuracy before sample analysis.

SCFA and MCFA measurements from cecal contents were obtained as follows. In brief, 20-40 mg of cecal contents were mixed with 0.4 – 0.8 mL of 25% phosphoric acid in water (LabChem). Sample mixtures were thoroughly homogenized via vortexing, then centrifuged at 15,000 x g for 10 minutes at 4ºC. Following centrifugation, the sample supernatant was transferred to a new tube, then centrifuged again at 15,000 x g for 10 minutes at 4ºC. The resulting supernatant was then frozen at -20ºC before being sent to the University of Alberta Agricultural, Food & Nutritional Science (AFNS) Chromatography Facility for volatile fatty acid (VFA) analysis using gas chromatography/mass spectrometry (GC/MS) as previously described (Ng et al., 2023). There, the supernatant was filtered through 0.45 μm filters and mixed in a 4:1 ratio with internal standard solution (24.5 mmol/L isocaproic acid). The samples were then injected into a Stabilwax-DA column (length: 30 m, inner diameter: 0.53 mm, film thickness: 0.5 μm, Restek Corporation) on a Bruker Scion 456 GC with a model 8400 autosampler (Bruker Ltd) using a helium carrier gas. The samples were run using a 10 mL/min constant flow with a 5:1 split ratio with a column temperature gradient as follows: 80ºC initial temperature, immediately increased to 180ºC at 20ºC/min, then held for 3 min at 180ºC. The injector and detector temperature were at 250ºC. The peaks were analyzed using CompassCDS software. The concentration of SCFAs/MCFAs per sample was then normalized to mass of the cecal contents for statistical testing.

Bacterial culturing and preparation for SCFA quantification

Selected bacterial strains (Table 2) from public culture collections and lab strain libraries were cultured anaerobically in a vinyl anaerobic chamber maintained with an atmosphere of 5% CO_2, 5% H_2, and 90% N_2 for 1-5 days (Coy Lab Products, Grass Lake, MI, USA) on pre-reduced Anaerobic Akkermansia Media plates (AAM:18.5 g/L Brain Heart Infusion, 5.0 g/L Yeast Extract, 8.5 g/L Tryptone, 1.5 g/L Soytone, 2.5 g/L NaCl, 3.75 g/L K_2HPO_4, 1.75 g/L *D*-glucose, 0.4 g/L Na_2CO_3, 1.0 mg/L Hemin, 0.5 g/L L-Cysteine HCl, 0.5 mg/L Vitamin K1, and 3% v/v heat-inactivated fetal calf serum supplemented with 15 g/L Agar) until visible colonies were observed. Subsequently, strains were inoculated in AAM for 3 days (or 1 day for fast-growing strains) before being centrifuged at 6000 xg for 5 minutes to remove bacteria. 1 mL of bacterial supernatants was then combined with 200 μl of 25% v/v phosphoric acid before being frozen and quantified at the AFNS Chromatography Facility as described above.

Strain	Culture Collection	Reference number

Eubacterium eligens	DSMZ	3376
Eubacterium ventriosum	DSMZ	3988
Eubacterium hallii	DSMZ	3353
Eubacterium biforme	DSMZ	3989
Eubacterium rectale	ATCC	33656
Roseburia inulinivorans	DSMZ	16841
Ruminococcus lactaris	ATCC	29176
Ruminococcus torques	ATCC	27756
Ruminococcus gnavus	ATCC	29149
Ruminococcus bromii	ATCC	27255
Lactobacillus salivarius HA-118	Probiotic	
Lactobacillus acidophilus A118	Probiotic	
Lactobacillus reuteri NM12	CIAMIB	(Wong et al., 2022)
Lactobacillus murinus NM26	CIAMIB	(Wong et al., 2022)
Lactobacillus murinus NM28	CIAMIB	(Wong et al., 2022)
Lactobacillus johnsonii NM60	CIAMIB	(Wong et al., 2022)
Lactobacillus intestinalis NM61	CIAMIB	(Wong et al., 2022)
Akkermansia muciniphila	ATCC	BAA-835
Escherichia coli NM84	CIAMIB	(Wong et al., 2022)
Bacteroides thetaiotaomicron 8702		(Han et al., 2021)
Bacteroides thetaiotaomicron 8713		(Han et al., 2021)
Bacteroides thetaiotaomicron 3731		(Ng et al., 2023)
Bacteroides thetaiotaomicron VPI 5482		(Ng et al., 2023)

Table 2. Cultured bacterial strains used for *in vitro* SCFA measurements.

Bacterial metagenomics shotgun sequencing and analysis

DNA from cecal contents collected from mice 3 days after cessation of the third round of DSS
was extracted and quantified as described above in "*Microbiome sequencing and analysis*".
Shotgun metagenomics library preparation was done using the Illumina DNA Prep library
preparation kit in accordance with manufacturer instructions, and sequencing was then
performed on the Illumina NextSeq^{TM} platform with 2x150 bp paired-end read chemistry.
Library preparation and sequencing were performed at the UBC Sequencing + Bioinformatics
Consortium.

For shotgun sequencing analysis, the quality of raw reads was evaluated using FASTQC v0 12.1
(Andrews, 2010) and MultiQC v 1.19 (Ewels et al., 2016). Data was then run through
SqueezeMeta v 1.6.5 pipeline in coassembly mode (Tamames and Puente-Sánchez, 2019). The
coassembly method pools reads from all samples to create a single shared metagenome
assembly. Briefly, sequences were trimmed and quality-filtered using Trimmomatic (Bolger et
al., 2014); metagenome assembly was performed using Megahit (Li et al., 2015); ORF and rRNA
prediction was done using Prodigal (Hyatt et al., 2010) and barrnap (Seemann, 2024),
respectively before classification was performed using RDP classifier (Wang et al., 2007);
Diamond software (Buchfink et al., 2015) was used to assign taxonomy against the Genbank nr
database, COGs/NOGs against the eggNOG database (Huerta-Cepas et al., 2016) and KEGG ID
against the KEGG database (Kanehisa and Goto, 2000); gene coverage and abundance

estimation were performed by mapping individual reads to the reference coassembly contig using
Bowtie2 (Langmead and Salzberg, 2012). Finally, SqueezeMeta outputs were analyzed using the
SQMTools package (Puente-Sánchez et al., 2020). Normalization of gene abundance was
performed using trimmed mean of M-values (TMM) with the edgeR package (Robinson et al.,
2009) and plotted using ComplexHeatmap (Gu et al., 2016). The TMM method of normalization
can robustly account for differences in sequencing depth between samples by adjusting raw
counts using a unique scaling factor for each sample. The scaling factor is derived by comparing
gene abundances against a reference sample (which is automatically determined to be the sample
that has the 75th percentile of log-counts closest to the median across all samples), then using a
weighted trimmed mean over the log-transformed fold-change differences between the sample
and the reference.

β-glucuronidase assay

Beta-glucuronidase activity was measured using the QuantiChrom^{TM} β-Glucuronidase Assay Kit
(BioAssay Systems, San Francisco, CA, USA). Cecal contents collected from mice acutely
following DSS (P71) and after ~40 days of recovery (P110) were processed as follows: 10-50
mg of cecal contents were aliquoted and suspended in 600 µl of sterile dH$_2$O. Samples were then
homogenized at 30 Hz for 2 minutes using the TissueLyser II (QIAGEN) before being
centrifuged at 1000 xg for 1 minute at 4ºC to remove cecal debris without pelleting bacteria. The
supernatant was then centrifuged at 13000 xg for 5 minutes at 4ºC. The pelleted bacteria were
then resuspended in 600 µl of chilled sterile dH$_2$O and vortexed for 10 seconds. Bacterial lysis
was achieved through 3 rounds of sonication at 50 watts for 20 seconds each using an XL2020
sonicator equipped with a Microtip^{TM} (Misonix, Farmingdale, NY, USA). Samples were diluted
1:2-1:10 as needed before performing the β-glucuronidase activity assay according to
manufacturer instructions. Enzyme activity was normalized to total protein concentration in each
sample, measured using the Pierce BCA Protein Assay Kit (ThermoFisher).

MRI acquisition and data processing

Ex vivo imaging and analysis, including standard data preprocessing and region of interest (ROI)
analyses, were performed as previously described (Yi et al., 2019). Briefly, brains were extracted
from the cranial vault between P95 and P130 and post fixed in 4% PFA (n = 54;13 vehicle per
sex, 14 DSS per sex). Brains were placed in a custom-built holder immersed in Fluorinert (FC-
3283, 3M, St. Paul, MN, USA) and imaged with a 4.7-T Agilent MRI system with a 3.5 cm
diameter quadrature volume RF coil. Multi-slice, diffusion-weighted, spin echo images were
used to acquire 10 nondiffusion weighted images (b=0 s•mm^{-2}) and 75 diffusion-weighted
images (25: b=800 s•mm^{-2}, 50: b=2,000 s•mm^{-2}), using non-colinear diffusion-weighting
directions. Diffusion imaging was performed with a TE/TR=24.17/2000-ms, FOV=30×30 mm^2,
and a matrix=192×192 reconstructed to 256×256 for an isotropic voxel size of 0.25-mm over two
signal averages. Raw data files were converted to NIfTI format and FSL was used to correct for
eddy current artifacts (Smith et al., 2004). A DTI-based mouse brain atlas (Jiang and Johnson,
2011) was used as a study space template and to define regions of interest (ROI) including the
left and right hippocampus, frontal association cortex, striatum, amygdala, thalamus, and
hypothalamus. Multi-shell diffusion data were fit with the Microstructure Diffusion Toolbox
(MDT) (Harms et al., 2017) to the neurite orientation dispersion and density imaging (NODDI)
model (Zhang et al., 2012). An additional compartment of isotropic restriction was included to

account for potential fixative effects as recommended (Alexander et al., 2010). The mean values
for neurite density index (NDI) and orientation dispersion index (ODI), which represent indices
of intracellular and extracellular diffusion respectively, were calculated in each ROI using
Pyradiomics (van Griethuysen et al., 2017) and compared among experimental groups.

Statistical analyses

Statistics and graphs were created with GraphPad Prism 10.1.1 (GraphPad Software), Python 3.8
and RStudio software (RStudio Team, 2022). A repeated measures ANOVA was used to assess
body weight changes during treatment. All organ comparisons, SCFA measurements, gut
microbial β-glucuronidase (GUS) activity measurements, and almost all behavioral tests were
assessed using a two-way ANOVA to assess main effects of treatment and sex and Šidák
corrected post-hoc comparisons were performed to determine statistical significance within sex if
main effects or interactions were observed. The urine preference test was assessed by comparing
each group to chance (50%) using a single sample t-test. To directly compare preferences
between treatments, an unpaired two-tailed t-test was conducted. Steroid levels were analyzed by
Mann-Whitney U. Statistical significance of alpha diversity metrics (Observed features) was
assessed using a two-way ANOVA to assess main effects of treatment and sex. Beta diversity
metrics were evaluated using PERMANOVA analysis. FDR-corrected p-values obtained from
DAA with metagenomeseq were reported.

For the MRI analysis, a 2×2 analysis of variance (ANOVA) was used to determine the effect of
treatment, biological sex, and their interactions on the ROI values. The effects of ROI laterality
and imaging age effects were regressed from the data before the ANOVA was performed as
these effects were not pertinent to the tested hypotheses. Benjamini-Hochberg false discovery
rate (FDR) procedure was used to control for multiple hypothesis testing across ROIs within
each imaging signal. Statistical significance of an ANOVA is defined as an FDR adjusted p-
value <.05. Post-hoc pairwise difference testing was performed using Fisher's LSD test Partial η^2
explains how much of the unexplained variance in the model is described by the addition of that
variable into the model. The plotted imaging metric values represent the metric values after the
laterality and batch effects have been removed.

Results

An early-life DSS model of pediatric IBD

To establish a model of juvenile-onset IBD that mimics the remitting and recurrent cycles of gut
inflammation often observed in juvenile patients with IBD, we administered DSS in drinking
water repeatedly across early postnatal development starting on postnatal day 21 (P21) (Figure
1A). Colon length is indicative of inflammatory state, with colon shortening being induced by
effective DSS treatment (Chassaing et al., 2014). Three days after the final treatment, colon
length was significantly decreased by DSS in both sexes (main effect of treatment (F (1, 21) =
39.30, *p*.0001) (Figure 1B). Both female (*p*=.009) and male (*p*=.0002) mice treated with DSS
had colon lengths that were significantly shorter than untreated mice, indicating successful
induction of intestinal inflammation in our model. Furthermore, body weights were recorded

throughout the treatment period (Chassaing et al., 2014) (Figures 1C, D). Female mice showed
mild weight loss in response to DSS, with only the second round producing significant decreases
from VEH littermates (treatment x day interaction $F(50,714) = 6.693$, $p<.0001$) (Figure 1C).
Conversely, male DSS-treated mice showed more dramatic weight loss across the three treatment
periods (treatment x day interaction $F(50,686) = 12.372$, $p<.0001$) (Figure 1D). To further
investigate inflammation caused by repeated administration of DSS, we examined the
histological architecture of colonic tissues using hematoxylin and eosin staining. Following
chronic DSS treatment, the gut epithelial layer was disturbed, showing increased crypt length
(crypt abscess), depletion of mucus-producing goblet cells, increased incidence of inflammatory
cell infiltration, and ruffling of the normally smooth epithelium (Figures 1E, F). Furthermore,
UEA-1 lectin staining targeting the intestinal mucus layer revealed that DSS-treated juvenile
mice experienced profound alterations in mucus structure, with loss of the normal striated bilayer
organization (Ng and Tropini, 2021) (Figure 1F). Finally, to examine changes to gut microbiota
composition caused by repeated cycles of early life inflammation, we performed 16S rRNA
sequencing targeting the bacterial V4-V5 variable regions. Consistent with previous studies in
adult mice, DSS treatment resulted in significant losses to microbiota alpha diversity, with the
number of observed features, a proxy for the number of bacterial species present (Kleine
Bardenhorst et al., 2022), being significantly decreased in DSS-treated animals compared to
vehicle (significant effect of treatment $F(1,12) = 5.3$, $p = .0387$) (Figure 1G) (Munyaka et al.,
2016). Similarly, between-group microbiota compositional differences were revealed to be
significantly different when compared using calculated Bray-Curtis distance matrices (treatment
x sex interaction $F(1,12) = 3.2158$, $p < .046$) (Figure 1H). Altogether, these results suggest that
our DSS colitis model of juvenile IBD induces profound physiological and microbiological
changes to the gut.

Figure 1. Juvenile mouse DSS treatment models early-life intestinal inflammation. *A*.
Experimental schematic. Control mice were given water throughout (VEH), while treatment

mice were given 3 rounds of 5-day long DSS treatments beginning at P21, P42, and P63 (DSS).
B. Post-treatment colon lengths measured at P71, three days after the final round of DSS, were
shorter in DSS-treated mice. Each graph bar and error bar represent the mean ± SEM, n=5-
7/group. *C, D.* Female and male percent weight change from baseline (P21) starting weight. Each
point and error bar represents the mean ± SEM, n=11-13/group. *E.* Representative images of
hematoxylin and eosin-stained colonic tissues from mice collected at P71 confirm that DSS
treatment induced intestinal inflammation. Scale bar, 100 μm. *F.* Representative fluorescence *in*
situ hybridization (FISH) images from VEH and DSS treated mice shows gut disruption in DSS
treated mice. Scale bar, 100 μm. Colors: host tissue (DAPI, blue), mucus (UEA-1, green), and
bacteria (Eub-338, red). *G.* Microbiota alpha diversity as measured by number of observed
unique features (a proxy for number of species) per sample is significantly lower in DSS-treated
mice, n=3-5/group. *H.* Principal Component Analysis of Bray-Curtis diversity index shows
separation by DSS treatment, n=3-5/group. *$p<.05. ***$p<.0001.

Early-life DSS affects self-grooming behavior in male mice

Previous work in early life gut-inflammatory models, including those for IBD, identified altered
mouse behavior (Salvo et al., 2020). To fully profile behavioral phenotypes in our pediatric IBD
model, we performed a battery of behavioral testing beginning 17 days after DSS treatment to
examine anxiety-like (elevated plus maze, light-dark box, open field test), memory (spatial
object location), sociability and social memory (three chamber task), repetitive behavior (self-
grooming), and depression-like behaviors (forced swim). Unlike previous reports, we did not
observe any significant effects of treatment for either sex in most of these tests (Figure 2 and
Supplemental Table 1), with the single exception of decreased self-grooming (main effect of
treatment ($F (1, 24) = 6.514$, $p=.018$)) in the DSS-treated males ($p=.031$). There were no
significant main effects of treatment or interactions with treatment on distance travelled or total
exploration on any of the tasks (Supplemental Figure 1 and Supplemental Table 1). This suggests
that that the majority of behaviors examined are not impacted by previous repeated bouts of
postnatal gut inflammation in early-life, potentially due to the normal development of most
neuronal circuits prior to treatment beginning at P21.

Figure 2. Early-life DSS treatment has minimal impacts on adult anxiety, social, memory, and depression-like behaviors. *A*. Experimental design. After a 16-day recovery period, a behavioral battery compared VEH and DSS treated mice on *B*. the elevated plus maze (n=10-13/group) *C*. the light dark box (n=11-13/group) *D*. the open field test (n=11-13/group) *E*. the object location memory test (24hr) (n=6-13/group) *F*. the three-chamber social task (n=11-13/group) *G*. the three-chamber novel social task (n=10-13/group) *H*. self-grooming (n=6-8/group) *I*. forced grooming (n=7-8/group) and *J*. the forced swim test (n=11-13/group). All graph bars and error bars represent the mean ± SEM. **p<.05*

Early-life DSS disrupts male preference for female odour

To test the impact of repeated bouts of inflammation on sexual behaviors in our model, ~1 month following DSS, mice were given a preference test for same- or opposite-sex urine (Figure 3A). Female VEH- and DSS-treated mice both failed to show a significant preference for either same-sex urine (VEH one sample t-test vs 50% $t(6) = 0.383$, $p=.714$ or DSS one sample t-test vs 50% $t(6) = 0.406$, $p=.70$) or opposite-sex urine (VEH one sample t-test vs 50% $t(6) = 0.350$, $p=.74$ or one sample t-test vs 50% DSS $t(6) = 0.256$, $p=.81$) compared to water (Figure 3B). In contrast, male VEH-treated mice showed a significant preference for same-sex urine compared to water (one sample t-test vs 50% $t(7) = 2.468$, $p=.04$) and a trend for a preference for opposite-sex urine $t(7) = 1.972$, $p=.086$. However, male DSS-treated mice failed to show a preference for same-sex urine ($t(7) = 1.153$, $p=.29$) as well as opposite-sex urine over water ($t(7) = 0.99$, $p=.36$) (Figure 3C). When comparing preferences directly, there was no significant difference in VEH-treated and DSS-treated males in their preference for male urine ($t(14) = 1.45$, $p=.17$). However, VEH-treated males showed a significantly higher preference for female urine than DSS-treated males ($t(14) = 2.16$, $p=.048$) (Figure 3C). Together, these data demonstrate that DSS treatment produced a significant deficit in opposite-sex scent-seeking behaviors, a key aspect of mate-seeking behavior in male rodents (Malkesman et al., 2010).

Figure 3. Early-life DSS treatment induces male sex-specific impairments in mate-seeking behavior. *A*. Experimental design. After a 30-day recovery period, mate-seeking behavior was

assessed by the Urine Preference Test. *B*. Female mice showed no preference for female or male
urine over water and VEH and DSS female mice performed similarly. n=7/group. Each graph bar
and error bar represent the mean ± SEM. Dotted line is 50% preference. *C*. Male VEH mice
significantly preferred both male and female urine over water (above 50% dotted line). DSS
treated males showed no preference for urine over water. n=8/group. *D, E*. Olfactory habituation-
dishabituation test for females (*D*) and males (*E*) revealed no differences between DSS- and
VEH- treated mice and normal exploration, habituation and then dishabituation patterns for each
odour n=6-7/group. *F*. DSS and VEH male mice showed similar number of mounts and latency
to first mount a sexually experienced female. n=4-5/group. *p<0.05.

To verify that deficits in the urine preference test in DSS-treated males were not due to
impairments in olfaction, we performed the olfaction habituation-dishabituation task (Yang and
Crawley, 2009). We tested two extract odours (vanilla and banana) and two social odours (one
same sex and one opposite sex novel mouse odours). Both treatment groups and sexes showed
the expected patterns of active exploration of each odour followed by habituation across repeated
trials and then active exploration of the next novel odour (dishabituation) (Figure 3D and E and
supplemental Table 1), indicating that DSS treatment did not negatively impact odour processing
or recognition. Together, these findings demonstrate a long-term shift in male-specific
preferences for female odours following early-life gut inflammation.

To further explore impacts on consummatory mating behaviors, we compared sexually naïve
mounting behavior of the VEH- and DSS-treated male mice when paired with a sexually
experienced untreated female wildtype control mouse. Over a 15-minute mating session, DSS-
treated male mice showed no significant differences in copulatory behaviors compared to VEH
males including frequency of mounting ($t(8) = 1.06$, $p=.32$) and latency to first mount ($t(6) =$
1.30, $p=.42$) (Figure 3F). Together, this suggests that early-life DSS treatment has specific
deficits on mate-seeking through olfactory cues and not on consummatory sexual behavior.

Early-life intestinal inflammation disrupts androgens in males

Given the observed differences in urine odor preferences, we investigated reproductive organ
development in DSS- and VEH- treated mice. Male and female reproductive organs were
collected following behavior testing and weighed (P110). In males, there was a significant
decrease in the weight of the androgen-sensitive seminal vesicles in the DSS-treated animals
compared to VEH ($t(23) = 2.44$, $p=.03$) (Figure 4A). However, there was no significant
difference in testes weight between treatments ($t(23) = 1.05$, $p=.30$) (Figure 4B). In females, we
observed no difference in uterus weight ($t(23) = 0.42$, $p=.68$) (Figure 4C). To further investigate
the impairment in male seminal vesicle development, we examined changes in circulating steroid
hormones in VEH- and DSS-treated male mice both acutely (P71, 3 days after final DSS
treatment) (Figure 4D, E) and chronically (after behavioral testing, P110) (Figure 4F, G). At the
acute timepoint (P71), the DSS-treated males showed a significant decrease in serum
testosterone ($U = 0$, $p =.03$) and androstenedione ($U = 0$, $p =.03$) levels. This decrease was
maintained through behavioral testing for androstenedione ($U = 27$, $p =.01$) but not for
testosterone ($U = 46$, $p =.22$). There were no significant differences in serum progesterone in
males at either timepoint and no impacts in females for androstenedione, testosterone, or

progesterone (Supplemental Figure 2). All together, these data indicate that early-life DSS
treatment reduces male reproductive organs, potentially through reduced levels of circulating
androgens.

Figure 4. Early-life DSS treatment induces male sex-specific impairments in reproductive
physiology. *A*. Seminal vesicles were significantly lower in weight in DSS-treated male mice
than VEH-treated mice at P110. n=12-13/group. *B*. There were no significant differences in
testes weight between DSS- and VEH-treated males at P110. n=12-13/group. *C*. No significant
differences in uterus and ovaries weights between DSS- and VEH-treated females were observed
at P110. n=12-13/group. *D-E*. 3 days after the last DSS treatment (P71), male serum testosterone
(*D*) and androstenedione (*E*) levels were significantly lower in DSS- than in VEH-treated mice.
n=4/group. *F-G*. Following the last behavioral experiment (P110), male serum testosterone (*D)*
levels were not significantly different between VEH and DSS mice, but androstenedione (*E*)
levels were significantly lower in DSS- than in VEH-treated mice. n=12-13/group. Each graph
bar and error bar represent the mean ± SEM. *$p<0.05$.

Early-life intestinal inflammation alters SCFA/MCFA production and gut microbiome
composition to favor decreased abundance of SCFA/MCFA-producing taxa

Since we observed changes in microbiome composition immediately following DSS treatment,
we then examined circulating short and medium-chain fatty acids (SCFAs and MCFAs
respectively) (Figure 5A), which are produced by the gut microbiota. Importantly, SCFAs are
known to be disrupted in patients with IBD and can alter sex steroids (Acharya et al., 2024) and
brain function (Cryan and Dinan, 2012; Erny et al., 2015; Sullivan and Ciernia, 2022). Acutely

following the last round of DSS (P71), DSS treatment significantly reduced female (p = .014)
and male (p = .037) serum levels of valeric acid (see Supplemental Table 1 for full statistics). In
females, DSS also decreased serum levels of capric acid (p = .022) (Figure 5A). Interestingly, we
found no significant differences in levels of acetic, propionic, and butyric acid, which are the
most abundant SCFAs and play important roles in maintaining intestinal homeostasis (Venegas
et al., 2019). We also measured SCFA/MCFA levels in cecal contents, where butyric acid was
the only SCFA that was significantly different between VEH- and DSS-treated males (p = .0498)
(Figure 5B, see Supplemental Table 1 for full statistics). Critically, butyric acid is an important
anti-inflammatory agent that is known to be reduced in patients with IBD (Hodgkinson et al.,
2023).

We then investigated whether SCFA-producing bacteria were differentially present in VEH- and
DSS-treated mice acutely following the last round of DSS at P71 (Figure 5C, Supplemental
Figure 3A). Differential abundance analysis revealed that DSS-treated female mice possessed
decreased relative abundance of many bacteria genera belonging to the phylum Bacillota
(Supplemental Figure 3B, See Supplemental Statistics Table for full list), which contains many
SCFA-producing bacterial species (Fusco et al., 2023; Mandal et al., 2015). Indeed, the relative
abundance of Bacillota was moderately correlated with cecal acetic (R = 0.54, p = .005) and
butyric acid (R = 0.76, p < .001) levels (Supplemental Figure 3C). Specifically, DSS-treated
mice possessed decreased relative abundance of the genus *Roseburia* ($p_{females}$ = .0039, p_{males} =
.0201) along with several *Eubacterium* groups ([*Eubacterium*] ventriosum group, $p_{females}$ < .0001,
p_{males} < .0001; [*Eubacterium*] xylanophilum group, $p_{females}$ < .0001, p_{males} = .0062; [*Eubacterium*]
coprostanoligenes group, $p_{females}$ = .0059, p_{males} < .0001) – bacterial taxa that are known to
produce SCFAs (Figure 5C, D, Supplemental Figure 3B) (Akhtar et al., 2021; Fusco et al., 2023;
Mirzaei et al., 2021; Ziętek et al., 2021). Differential abundance analysis also revealed that the
species *Bacteroides thetaiotaomicron* ($p_{females}$ = .0001, p_{males} < .0001) and *Akkermansia*
muciniphila ($p_{females}$ < .0001, p_{males} = ns) were enriched in DSS-treated mice, bacteria not known
to produce butyric acid (Effendi et al., 2022; Kim et al., 2021). To investigate these correlations,
we cultured a variety of strains belonging the above identified taxa, then measured production of
SCFAs *in vitro* (Figure 5D). This analysis revealed that *Eubacterium* sp. and *Roseburia sp.* are
both capable of producing butyric acid and are depleted following DSS treatment, potentially
contributing to the decreased concentration of butyric acid in cecal contents. Interestingly, strains
that are both more (*Bacteroides* sp. and *Akkermansia* sp.) and less (*Ruminoccocus* sp. and
Lactobacillus sp.) abundant after DSS treatment were capable of producing acetic acid, which
may contribute to the lack of significant difference in cecal acetic acid concentration following
DSS.

Due to the differences observed in mating behaviour and circulating sex hormone levels
following DSS treatment, we also examined the potential role of the DSS-altered microbiome in
driving these phenotypes. Of the differentially abundant taxa observed, we noted that many have
been shown to express β-glucuronidases (GUS) and β-glucosidases, enzymes that convert
conjugated hormones to their active form (Ervin et al., 2019; Patel et al., 2023). Strikingly, the
relative abundance of several GUS-producing taxa was correlated with serum testosterone and
androstenedioine concentrations (Figure 5E). For instance, the relative abundance of
[*Eubacterium*] *coprostanoligenes* group was positively correlated with androgen levels
(androstenedione, R = 0.91, p = .002; testosterone, R = 0.91, p = .002). Similarly, the relative
abundance of *Lactobacillus* was positively correlated with serum androstenedione (R = 0.81, p =

.022) and testosterone ($R = 0.90$, $p = .005$). In contrast, the relative abundance of *Bacteroides*
thetaiotaomicron, which is known to express a sulfotransferase that can deactivate hormones
(Cotton et al., 2023), was negatively correlated with testosterone concentrations ($R = -0.74$, $p =$
.046) (Supplemental Figure 3D).

Following this observation, we performed shotgun metagenomic sequencing analysis to examine
the genomic potential of the microbiota to express hormone-modifying and degrading enzymes
(Figure 5F). Supporting the genetic potential of the microbiota in our mice to modify active
levels of hormones, we found the presence of genes mapping to various hydroxysteroid
dehydrogenases (K00038 and K22418) as well as β-glucuronidase (K01195). Importantly,
genomic potential is not a predictor of expression levels or enzyme activity due to various pre-
and post-translational regulatory mechanisms. Therefore, we aimed to measure GUS activity
directly from cecal contents using a fluorometric molecule conjugated to a glucuronide.
Interestingly, GUS activity was significantly decreased in males ($p = .0006$) following DSS
treatment, with activity in females trending lower as well ($p = .0856$) (Figure 5G). Taken
together, these findings indicate that early life inflammation alters microbiome composition,
leading to differences in SCFA producers and GUS activity which correlates with SCFA and
androgen levels.

Figure 5. Early-life DSS treatment causes decreases in short- and medium-chain fatty acid
levels and relative abundance of associated microbiota. *A*. Serum levels of valeric and caproic
acid were significantly lower in DSS-treated females compared to VEH at P71. Valeric acid was
also significantly lower in DSS-treated male serum. n=4-5/group. See Supplement for full set of
serum statistics. *B*. Cecal levels SCFAs at P71, represented in nmol/mg of contents.
Concentration of butyric acid was significantly different in DSS-treated males compared to VEH
at P71. n=4-5/group. See Supplement for full set of cecal SCFA statistics. *C*. Relative abundance
of select SCFA-producing bacteria. n=4-5/group. *D. In vitro* SCFA production from bacterial
genera represented in the experimental microbiota. *E*. Correlation of select bacteria with serum
testosterone and androstenedione concentrations in males. n=4/group. *F*. Metagenomic
abundance of steroid-modifying and steroid-degrading enzymes normalized by TMM, trimmed
mean of M-values. Numbers represent mean TMM-normalized counts per treatment for each
enzyme. n=4-6/treatment. *G*. GUS activity in cecal contents was significantly lower in DSS-
treated males. Activity was measured from cecal contents and normalized to total protein. n=3-
5/group. Error bars in bar plots represent SEM. *$p<0.05$, ***$p<.001$, ****$p<.0001$.

Early life intestinal inflammation causes long-lasting physiological differences and changes to
gut microbiota composition

Having observed changes in microbially-produced and modified metabolites and bacterial
abundance acutely post-inflammation, we sought to measure whether they persisted into
adulthood (P110). We first examined microbiome diversity as an indicator of overall lasting
effect of DSS. At P110, the number of observed microbiota features remained significantly
decreased in female DSS-treated mice compared to VEH ($p = .0207$) (Figure 6A). Additionally,
although microbiota compositions (as calculated using Bray-Curtis distances) became more
similar between DSS-treated and VEH mice, differences in sex and treatment remained
significant (PERMANOVA, $p_{sex} = .001$, $p_{treatment} = .004$) (Figure 6B). Consistent with these long-
lasting microbiota changes, analysis of colon histology revealed that DSS-treated colons at P110
retained several indicators of inflammation at sparsely found regions throughout the gut,
specifically intestinal crypt hyperplasia and increased immune cell infiltration (representative
images shown, Figure 6C). Nevertheless, the majority of the epithelium along the length of the

colon appeared to have recovered after ~40 days post cessation of DSS

Figure 6. Early-life DSS treatment causes long-lasting physiological differences and
changes to microbiota composition. *A.* Diversity of microbiota composition ~30 days after
cessation of DSS at P110, measured in number of unique features per sample. n=5/group. *B.*
Principal Component Analysis of Bray-Curtis diversity index at P110. n=5/group. *C.*

Representative histological images of hematoxylin and eosin-stained colons at P110. *D.* Cecal
concentrations of SCFAs at P110. n=5/group. See Supplement for full set of cecal
measurements. *E.* Relative abundance bar plots of select SCFA-producing bacteria. n=5/group.
F. GUS activity in cecal contents at P110. Activity measured from cecal contents and normalized
to total protein/sample. n=5/group. $*p<.05$, $****p<.0001$. Error bars in bar plots represent SEM.

Following these observations, we next sought to examine whether gut bacteria and microbial
signaling molecules showed long-lasting changes after recovery. First, we found that there were
no significant differences in cecal SCFA levels between VEH- and DSS-treated mice at P110
(Figure 6D). Consistent with these results, we found that many of the differentially abundant
SCFA-producing taxa identified to be differentially present at P71 such as *Eubacterium*
coprostanoligenes group and *Roseburia* were no longer significantly different between
treatments. In addition, there were fewer significantly different taxa identified at P110 (Figure
6E) compared to the P71 (Supplementary Statistics Table, Supplemental Figure 3A) acute
timepoint, indicating that the microbiome composition shifted towards recovery. Of the taxa that
remained differentially abundant, *Eubacterium* ventriosum group ($p_{females} < .0001$) is a known
SCFA-producing taxa (Figure 6E). Second, we measured GUS activity from cecal contents at
this timepoint and found that, consistent with a shift towards recovery, GUS activity was no
longer depleted in DSS-treated animals (Figure 6F). Ultimately, these findings support the notion
that early life inflammation causes acute changes to the microbiome and microbial signaling
molecules, some of which may equilibrate in absence of further inflammation in adulthood.

Early-life intestinal inflammation disrupts microglial morphology and brain connectivity in both sexes

SCFAs have been shown to alter microglial morphology and function (Erny et al., 2015). We
therefore examined changes in microglial activation state in several brain regions. Using
MicrogliaMorphology and MicrogliaMorphologyR (Kim et al., 2024a), we measured 27
morphological features from 12,318 microglia in the dorsal hippocampus (average of 513 per
mouse) (Figure 7), 5,971 microglia in the hypothalamus (average of 426 per mouse) and 8,482
microglia in the cortex (average of 446 per mouse) (Supplemental Figure 4).We identified four
clusters of different morphologies (Figure 7A, Supplemental Figure 4D) including microglia
showing ameboid, ramified, rod-like and hypertrophic characteristics. Within the dorsal
hippocampus there was a significant treatment by cluster interaction within each sex
(Supplemental Table 1). Sidak corrected posthoc tests identified a significant decrease in
ramified microglia in DSS-treated females (Figure 7B). In males there was a significant increase
in hypertrophic microglia in the DSS-treated males. There were no significant differences in
hypothalamic or cortical microglia between treatments within either sex (Supplemental Figure
4G, H). We further examined expression of cytokine genes and several immune receptors in
brain tissue of DSS and VEH treated mice (Supplemental Figure 5). There were no significant
differences between treatments within either sex, indicating that the sex-specific impacts of early
life DSS treatment on microglia were not solely driven by neuroinflammation.

Figure 7. Long-Term Microglial Morphology and Extraneurite Microstructure Changes Following Early-Life DSS. *A*. Clusters were assigned to morphology descriptions based on relationship to individual metrics (see supplemental Figure 4). *B*. Example images from dorsal hippocampus of female (P110) and male (P72 and P110) vehicle and DSS treated mice reveals sex-specific changes in microglial morphology. Top panel is IBA1 staining. Middle panel is the overlay with individual cell morphology clusters and the bottom panel is the individual cell labels for each cluster. All scale bars are 50 μm. *B*. Quantification of the percentage of microglia within each cluster between treatment conditions for females and males. DSS significantly decreased the proportion of ramified microglia in females and increased hypertrophic microglia in males. n=4-8/group *C*. Orientation dispersion index (ODI) was calculated for the bilateral hippocampus across all samples. Data points are mean ODI values from each sample. Both males and females had significant increases in ODI in the DSS treatment when compared to treatment with the vehicle alone. n= 25-28/group * = *p* < .05, *** = *p* < .001.

Clinical-translational quantitative neuroimaging reveals extraneuritic hippocampal
microstructural changes following early life inflammation in both sexes

We performed ex-vivo multicompartment (MC) diffusion MRI and fit it to the neurite orientation
and dispersion index (NODDI) model to assess and compare microstructural changes across
biologically relevant regions (hippocampus, amygdala, striatum, thalamus, hypothalamus, frontal
association cortex) between the VEH- and DSS-treated mice. MC diffusion MRI enables non-
invasive characterization of neuronal cytoarchitecture at the mesoscale (1- 100μm). Specifically,
NODDI diffusion MRI permits non-invasive quantification of neuronal microstructure by
measuring neurite density and cellular and morphometric changes occurring in the extra-
neuronal space. In our imaging experiments, the mean orientation dispersion index (ODI, extra-
neurite) and mean neurite density index (NDI, intra-neurite) values were tested using a 2x2
ANOVA considering the effects of treatment and sex. For ODI, a significant main effect of
treatment was identified in the hippocampus (FDR-adjusted p = .0002, partial η^2 = .1560)
(Figure 7D). Post-hoc testing identified that both males and females had a significant increase in
their ODI value when treated with DSS compared to VEH (males: Fisher's LSD p = .0001;
females: Fisher's LSD p = .0269). Additionally, a significant main effect of treatment was found
in the thalamus (FDR-adjusted p = .0095, partial η^2 = .0822) (Supplemental Figure 6). Post-hoc
testing indicated that females had a significant increase in their ODI value when treated with
DSS compared to VEH, but that the increase in males did not reach significance (males: Fisher's
LSD p = .2353; females: Fisher's LSD p = .0023). No significant treatment by sex interactions
were identified in either the thalamus or hippocampus. No significant treatment or treatment by
sex interaction effects were found in the frontal association cortex, amygdala, striatum, and
hypothalamus. Furthermore, for NDI, no significant effects were identified in any of the regions
of interest. Supplemental Table 1 contains the full findings for all regions tested.

Early-life intestinal inflammation specifically during adolescence decreases seminal vesicle
weight

To examine whether the developmental timing of one DSS treatment is sufficient to induce
impairments in male sex organ development, we treated male mice with DSS for 5 days either at
weaning (1xDSS Weaning) or during adolescence (1xDSS Adolescence) and collected seminal
vesicles at P71 (Figure 8). We found that DSS treatment produced significant decreases in body
weight (treatment x day interaction F(46, 318) = 15.111, p<.0001) following DSS treatment in
both groups (Figure 8 and Supplemental Table S1 and S2). DSS treatment during adolescence
significantly decreased seminal vesicle weights, highlighting this developmental period as
specifically sensitive to inflammatory effects on endocrine function (F(2, 12) =7.514, p=.009,
p=.0075). Conversely, DSS treatment during weaning did not alter seminal vesicle weight at P71
(p=.7565). Furthermore, there was no difference in testes weight between treatments F(2, 11)
=.575, p =.5788.

Figure 8. DSS Treatment During Adolescence Decreases Seminal Vesicle Weight. *A*. Male mice
were treated with a single puse of DSS at either weaning (P21-26) or adolescence (P43-48) and
were euthanized on P71. *B*. Body weight decreases following DSS treatment for the weaning
group at P28-33 and following DSS treatment for the adolescent group beginning on P50 through
the remainder of the experiment. *p<0.05 Sidak corrected posthoc tests. *C*. Seminal vesicle
weights were significantly lower in the adolescent DSS treatment group compared to vehicle
controls. *D*. Testes weights were not significantly different across treatments. n=4/group. Each
graph bar and error bar represent the mean ± SEM. *p<.05*.

Discussion

The incidence of IBD is high and increasing, especially in early life (Benchimol et al., 2017,
2014; Ng et al., 2017)., and the disease is particularly challenging when it develops in children.
Delay in the onset and progression through puberty is common in this population, and can result
in reduced adult height and sexual immaturity (Gupta et al., 2020). In this study, we assessed the
microbiome, intestinal, endocrine, brain, and behavioral effects of repeated DSS-induced gut
inflammation in early life by developing a novel model of juvenile IBD. While early life gut
inflammation did not impair the majority of the adult rodent behavioral repertoire, we identified
a unique deficit in preference for female urine in DSS-treated male mice. These deficits in mate-
seeking behavior corresponded to a decrease in seminal vesicle size, blunted circulating levels of
androgen hormones and SCFAs, an altered gut microbiota, and altered hormone-modifying
enzyme activity. Microglia were altered in the hippocampus, indicating hippocampal dysfunction
may contribute to reproductive behavioral deficits in males (Kight and McCarthy, 2020).

In our model, DSS significantly decreased seminal vesicle weight (Figure 4A), circulating sex
hormone levels (Figure 4D-E), and impaired male-specific preference for female urine (Figure
3C) – a key aspect of mate-seeking behavior in rodents (Keller et al., 2008; Malkesman et al.,
2010). These differences were not driven by impaired olfactory discrimination, as both sexes and
treatments performed normally on the olfactory habituation-dishabituation paradigm (Figure 3D-
E). In both adolescent males and females with IBD, puberty onset is often delayed and
progression through puberty is often prolonged, especially in patients with disease relapses
(Hildebrand et al., 1994). Furthermore, adults with IBD report increased rates of sexual
dysfunction (Bel et al., 2015), and infertility in both men (Hammami and Mahadevan, 2020) and
women (Zhang et al., 2021; Zhao et al., 2019). This is consistent with findings from our study,
where we found that DSS-treated male mice had decreased androgen hormones (Figure 4D-E)
that may have directly contributed to the decrease in seminal vesicle weight (Yamane et al.,
1986). Additionally, we observed that seminal vesicles in DSS-treated male mice were often not
descended fully within the abdominal cavity (data not shown), and that a single DSS treatment
during adolescence was sufficient to cause a decrease in seminal vesicle size (Figure 8C). While
we did not observe direct impacts on male mounting behavior, future work will be required to
fully examine the impacts of early life DSS on adult male reproduction and fertility.

Research suggests that hormone supplementation could potentially alleviate symptoms of IBD
(Nasser et al., 2015; Rosen et al., 2015). However, the use of hormone supplementation in
adolescents with IBD is contentious and can pose substantial risks due to potential side effects

such as weight gain, mood changes, sleep disruption, and increased susceptibility to infections
(Bishop et al., 2014). This necessitates new, non-hormone based therapies for treating adolescent
IBD. The microbiota represents a potential lever for management of IBD-mediated alterations in
hormones and development. Germ-free mice are known to have changes in sex hormones and
pubertal onset (Markle et al., 2013; Weger et al., 2019), suggesting that the gut microbiome
regulates sex hormone levels of the host. Furthermore, sex hormones can in turn regulate the gut
microbiome and gastrointestinal transit time (Rastelli et al., 2022). This bidirectional relationship
develops during puberty (Korpela et al., 2021). In both sexes, puberty is marked by increasing
abundance of estrogen-metabolizing Clostridia and decreasing abundance of Bacteroidia
(Korpela et al., 2021). Specific gut microbiota members can also secrete β-glucuronidase and β-
glucosidase enzymes that deconjugate inactivated estrogens and androgens back to their active
form (Ervin et al., 2019; Patel et al., 2023). These re-activated hormones are then secreted back
into the circulating blood, where they can act throughout the body. In humans, estrogen levels
are strongly and significantly associated with fecal Clostridia taxa, including three genera in the
Ruminococcaceae family (Flores et al., 2012). This suggests that gut microbial families such as
Ruminococcaceae may directly influence levels of host sex-hormones such as estrogen.
Disruption of β-glucuronidase-producing microbiota results in a reduction in circulating
estrogens (Baker et al., 2017), impairing the feedback loop between sex hormone levels and gut
microbiota composition. In our novel model of pediatric IBD, we report differences in the key β-
glucuronidase-expressing genus *Roseburia* acutely following the final round of DSS treatment,
along with several *Eubacterium* groups that are known to deconjugate estrogens (Figure 5C)
(Ervin et al., 2019). The relative abundance of these taxa were positively correlated with serum
testosterone and androstenedione concentrations (Figure 5E), highlighting the potential
involvement of these gut microbes in regulating systemic hormone levels. In addition, we
observed significant decreases in cecal β-glucuronidase activity in DSS-treated males, with
activity also trending lower in DSS-treated females (Figure 5G). Subsequently, we found that
after resolution of inflammation, the microbiome shifts towards recovery – with most of the
alterations in microbiome composition and microbial signalling molecules resolving (Figure 6).
Together, these findings suggest that future microbiota-targeted therapeutics specifically
impacting hormone-regulating microbes could potentially offset the negative effects of IBD on
sexual health.

How changes in gut inflammation drive endocrine, brain and behavior changes is not well
understood, but metabolite signalling molecules have been proposed as a potential link between
all three systems (Silva et al., 2020). A number of metabolites have been implicated in IBD
pathogenesis, including SCFAs, secondary bile acids (SBA) and tryptophan (Lavelle and Sokol,
2020; Sinha et al., 2020). These microbiota-derived metabolic signals can cross the blood-brain
barrier (Mertens et al., 2017; Mitchell et al., 2011; Wenzel et al., 2020) and are critical for
normal microglial development, gene regulation and metabolism (Erny et al., 2021, 2015; Thion
et al., 2017). In our study, we found more differences in SCFA/MCFA abundances at the acute
timepoint in the serum of DSS mice compared to VEH than in cecal contents (Figure 5A-B).
This difference could be due to SCFAs being absorbed by host cells for use as energy sources. In
particular, intestinal epithelial cells utilize SCFAs (particularly butyric acid) as a main source of
energy, causing the concentration of SCFAs in the blood to decrease compared to what is
measured in the cecum (den Besten et al., 2013). It has also been shown that butyrate uptake by
colonocytes is impaired in inflamed IBD patient tissues (Thibault et al., 2010). Thus, DSS-
induced inflammation could cause decoupling of cecal and serum SCFA concentrations. The

reduced levels of some SCFAs observed in our pediatric IBD model may be responsible for the
abnormal microglial morphology we observed (Figure 7B-C), with potential negative impacts on
microglial regulation of neural circuit formation during formative years in childhood and
adolescence (Sullivan and Ciernia, 2022). While previous work has shown that DSS can increase
microglial inflammatory gene expression acutely in adult animals, with males showing a stronger
response than females (Caetano-Silva et al., 2024), we did not observe changes in cytokine
expression in our experiments (Supplemental Figure 5). However, prior work on the impacts of
SCFAs on neuroinflammation is mixed. For example, increasing dietary fiber can offset the
impacts of aging on microglial gene expression (Vailati-Riboni et al., 2022); however, a high
fiber diet enhanced microglial inflammatory gene expression in an adult DSS model (Caetano-
Silva et al., 2024) and acetate treatment exacerbated pathology in an Alzheimer's disease mouse
model (Erny et al., 2021), suggesting more work is needed to understand how SCFAs modulate
microglia. We found decreases in valeric acid in both males and females treated with DSS.
Valerate, a ester of valeric acid, suppressed inflammation induced microglial phagocytosis in an
immortalized human microglia cell line (Wenzel et al., 2020). Consistent with these findings, we
observed decreased proportions of ramified microglia in DSS-treated females (Figure 7B,
indicating a decrease in homeostatic microglia. We also observed an increase in hypertrophic
microglia in DSS-treated males (Figure 7B), a phenotype observed previously during
neuroinflammation(Kim et al., 2024b). Together, these findings suggest that early life gut
inflammation can alter metabolite signaling to the brain, influencing microglial morphology in a
sex-specific manner. Future work will be needed to connect the alterations in SCFA directly to
microglial morphological state shifts and functions related to behaviors during development.

1097

In order to test the therapeutic efficacy of any future intervention in humans, diagnostic
biomarkers and biomarkers of therapeutic response are critically needed. Herein, we
demonstrated that early-life DSS treatment altered increased extra-neurite signal in NODDI
within the hippocampus and thalamus, but not other brain regions. The NODDI model is has
been used previously to calculate the diffusion signal specific to intra-neurite and extra-neurite
compartments of the brain and is sentisitve to various neuropathological states (Barnett et al.,
2019; Singh et al., 2023; Stowe et al., 2024; Yi et al., 2022). Consistent with our prior work (Yi
et al., 2019), changes in the ODI signal alone are consistent with its sensitivity to microglial
morphology and density (Figure 7C). Importantly, this imaging technique is being developed for
use in humans (Garcia-Hernandez et al., 2022), paving the way for future studies to examine
microglial changes in humans with IBD.

Finally, our main behavioral battery tested mouse anxiolytic, repetitive, cognitive, social and
depressive-like behavior and found no significant differences between control and DSS-treated
mice (Figure 2). This is in contrast to previous findings (Salvo et al., 2020), where 5 days of DSS
at weaning (P21) resulted in adult (P56+) behavioral deficits on the novel object recognition task
and light-dark task. This finding is rather unique in that the majority of adult DSS models show
altered anxiety and memory only during active inflammation (Emge et al., 2016) and would be
consistent with our findings that after recovery behavior is largely normal. One of the major
differences between our work in Salvo et al., 2020 is that we used three repeated DSS treatments.
The chronic nature of the current study may have induced resiliency and a faster recovery than in
a single DSS treatment. Previous work in adult mice comparing one versus repeated rounds of

DSS treatment found that acute-treated mice showed impairments in memory, repetitive,
depression and anxiety like behaviors while chronically treated mice only showed deficits in
repetitive behaviors (Matisz et al., 2020). The authors suggest that the lack of behavior deficits in
the repeated DSS treatment group may have been due to an adaptive or tolerizing effect of
repeated DSS cycles (Matisz et al., 2020). Similar findings in adult mice treated with repeated
rounds of DSS found no changes in locomotion, anxiety, depression or compulsive like
behaviors. However, they did find lasting alterations in social interactions (Brown et al., 2024),
which we did not observe. Salvo et al. did not examine sex-specific behaviors in their model; in
contrast, we have observed impacts specifically on male development. When we specifically
treated adolescent mice with DSS we found impacts on seminal vesicle development (Figure 8C)
similar to those observed after three rounds of DSS (Figure 4A), suggesting that DSS impacts on
male sexual development are driven by impacts during adolescence. Future work is required to
replicate and extend our findings to identify how timing of DSS treatment impacts reproductive
health including the onset of puberty and life-long fertility.

The incidence of pediatic onset IBD disease has doubled in the last 10 years (Ashton and Beattie,
2024), and young children with IBD often have more severe disease and more frequent health
visits than adults (Benchimol et al., 2011). Their illness also imposes a large emotional and
financial toll on families and caregivers (Herzer et al., 2011). Developing effective treatments for
adolescents with IBD holds great promise, as reducing inflammation early in the disease course
may prevent long-term complications, need for surgery, and hospitalization. Our novel mouse
model of pediatric onset IBD, which captures alterations in the microbiome, microbial signaling
molecules and brain-behavior disruptsion, represents a unique avenue for advancing our
mechanistic understanding of IBD in this key subpopulation. Our findings open the door for new
treatment avenues for therapeutics targeted to adolescents with IBD, with minimal side effects
and maximal benefits for growth, development, and quality of life.

1144

Funding

Research Corporation for Science Advancement, Scialog SA-MND-2021-001 (to CT, JPY) and
Scialog SA-MND-2023-028a and SA-MND-2023-035c (to AVC). Canadian Institutes of Health
Research (CIHR) Project Grant (to AVC, CT), Michael Smith Foundation Health Research
Scholar Award (to CT, AVC), Canada Tier 2 Research Chair, Quantitative Microbiota Biology
for Health Applications (to CT), Canada Tier 2 Research Chair, Understanding Gene Expression
in the Brain (to AVC), Paul Allen Distinguished Investigator Award (to CT), Johnson & Johnson
WiSTEM2D Award (to CT), Brain Canada Future Leaders Award (to AVC), Research
Corporation for Science Advancement Award (to AVC), UBC DMCBH pilot award (to AVC),
National Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant
(RGPIN-2020-04895) (to TH), PEO International Peace Scholarship and World Universities
Ramsay Postgraduate Scholarship (to SC). None of the funding sources influenced any aspect of
the study design, data collection, analysis, interpretation or writing of the manuscript.

1158
1159

1160 **Acknowledgement**s

The authors acknowledge that the land we performed this research on is the traditional, ancestral, and unceded territory of the xwməθkwəy̓əm (Musqueam) Nation. The land it is situated on has always been a place of learning for the Musqueam people, who for millennia have passed on in their culture, history, and traditions from one generation to the next on this site. We encourage others to learn more about the native lands in which they live and work at https://native-land.ca/.

This work would not be possible without the help of numerous core facilities. In particular, the authors would like to acknowledge the staff at the Centre for Disease Modeling (CDM) at UBC, Ingrid Barta at UBC Animal Care Services (ACS) Histology, and the UBC Biofactorial High-Throughput Biology Facility, supported by the University of British Columbia Global Research Excellence Biological Resilience Initiative. Thank you to the Djavad Mowafaghian Centre for Brain Health NeuroImaging and NeuroComputation Core (NINC) for data storage and computational resources. This research was supported in part through computational resources and services provided by the Advanced Research Computing at the University of British Columbia. Additionally, we would like to thank the Kopp Lab at UBC for allowing use of their tissue processor, embedder, microtome, and slide scanner. Finally, this work received support from the BC Children's Hospital Histology Core, Dr. Andy Sham, Dr. Catherine Chan, and the team from the Gut4Health Microbiome Core Facility at the BC Children's Hospital Research Institute, Lisa Nikolai from the Chromatography Facility at the Faculty of Agricultural, Life & Environmental Sciences at the University of Alberta, and Dr. Ian Nation from Animal Pathology Services Ltd. (Canmore, Alberta). Thank you to Dr. Shelly McErlane for consultation on sex organ collection in mice. Thank you to the Centre for Disease Modeling staff and animal technicians for assistance with mouse experiments and animal care.

References

Acharya, A., Shetty, S.S., Kumari N, S., 2024. Role of gut microbiota derived short chain fatty acid metabolites in modulating female reproductive health. Human Nutrition & Metabolism 36, 200256. https://doi.org/10.1016/j.hnm.2024.200256

Akhtar, M., Chen, Y., Ma, Z., Zhang, X., Shi, D., Khan, J.A., Liu, H., 2021. Gut microbiota-derived short chain fatty acids are potential mediators in gut inflammation. Anim Nutr 8, 350–360. https://doi.org/10.1016/j.aninu.2021.11.005

Alatab, S., Sepanlou, S.G., Ikuta, K., Vahedi, H., Bisignano, C., Safiri, S., Sadeghi, A., Nixon, M.R., Abdoli, A., Abolhassani, H., Alipour, V., Almadi, M.A.H., Almasi-Hashiani, A., Anushiravani, A., Arabloo, J., Atique, S., Awasthi, A., Badawi, A., Baig, A.A.A., Bhala, N., Bijani, A., Biondi, A., Borzì, A.M., Burke, K.E., Carvalho, F., Daryani, A., Dubey, M., Eftekhari, A., Fernandes, E., Fernandes, J.C., Fischer, F., Haj-Mirzaian, Arvin, Haj-Mirzaian, Arya, Hasanzadeh, A., Hashemian, M., Hay, S.I., Hoang, C.L., Househ, M., Ilesanmi, O.S., Balalami, N.J., James, S.L., Kengne, A.P., Malekzadeh, M.M., Merat, S., Meretoja, T.J., Mestrovic, T., Mirrakhimov, E.M., Mirzaei, H., Mohammad, K.A., Mokdad, A.H., Monasta, L., Negoi, I., Nguyen, T.H., Nguyen, C.T., Pourshams, A., Poustchi, H., Rabiee, M., Rabiee, N., Ramezanzadeh, K., Rawaf, D.L., Rawaf, S., Rezaei, N., Robinson, S.R., Ronfani, L., Saxena, S., Sepehrimanesh, M., Shaikh, M.A., Sharafi, Z., Sharif, M., Siabani, S., Sima, A.R., Singh, J.A., Soheili, A., Sotoudehmanesh, R., Suleria, H.A.R., Tesfay, B.E., Tran, B., Tsoi, D., Vacante, M., Wondmieneh, A.B., Zarghi, A., Zhang, Z.-J.,

Dirac, M., Malekzadeh, R., Naghavi, M., 2020. The global, regional, and national burden
of inflammatory bowel disease in 195 countries and territories, 1990–2017: a systematic
analysis for the Global Burden of Disease Study 2017. The Lancet Gastroenterology &
Hepatology 5, 17–30. https://doi.org/10.1016/S2468-1253(19)30333-4
Alexander, D.C., Hubbard, P.L., Hall, M.G., Moore, E.A., Ptito, M., Parker, G.J.M., Dyrby, T.B.,
2010. Orientationally invariant indices of axon diameter and density from diffusion MRI.
Neuroimage 52, 1374–1389. https://doi.org/10.1016/j.neuroimage.2010.05.043
Amann, R.I., Binder, B.J., Olson, R.J., Chisholm, S.W., Devereux, R., Stahl, D.A., 1990.
Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for
analyzing mixed microbial populations. Appl Environ Microbiol 56, 1919–1925.
https://doi.org/10.1128/aem.56.6.1919-1925.1990
Ananthakrishnan, A.N., 2013. Environmental Risk Factors for Inflammatory Bowel Disease.
Gastroenterol Hepatol (N Y) 9, 367–374.
Andrews, S., 2010. FastQC: A Quality Control tool for High Throughput Sequence Data [WWW
Document]. FASTQC: A Quality Control tool for High Throughput Sequence Data. URL
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/ (accessed 5.15.24).
Arganda-Carreras, I., Fernández-González, R., Muñoz-Barrutia, A., Ortiz-De-Solorzano, C., 2010.
3D reconstruction of histological sections: Application to mammary gland tissue.
Microscopy Research and Technique 73, 1019–1029.
https://doi.org/10.1002/jemt.20829
Ashton, J.J., Beattie, R.M., 2024. Inflammatory bowel disease: recent developments. Arch Dis
Child 109, 370–376. https://doi.org/10.1136/archdischild-2023-325668
Attree, E.A., Dancey, C.P., Keeling, D., Wilson, C., 2003. Cognitive Function in People With
Chronic Illness: Inflammatory Bowel Disease and Irritable Bowel Syndrome. Applied
Neuropsychology 10, 96–104. https://doi.org/10.1207/S15324826AN1002_05
Azooz, O.G., Farthing, M.J.G., Savage, M.O., Ballinger, A.B., 2001. Delayed puberty and response
to testosterone in a rat model of colitis. American Journal of Physiology-Regulatory,
Integrative and Comparative Physiology 281, R1483–R1491.
https://doi.org/10.1152/ajpregu.2001.281.5.R1483
Baker, J.M., Al-Nakkash, L., Herbst-Kralovetz, M.M., 2017. Estrogen–gut microbiome axis:
Physiological and clinical implications. Maturitas 103, 45–53.
https://doi.org/10.1016/j.maturitas.2017.06.025
Ballinger, A.B., Savage, M.O., Sanderson, I.R., 2003. Delayed Puberty Associated with
Inflammatory Bowel Disease. Pediatr Res 53, 205–210.
https://doi.org/10.1203/01.PDR.0000047510.65483.C9
Banfi, D., Moro, E., Bosi, A., Bistoletti, M., Cerantola, S., Crema, F., Maggi, F., Giron, M.C.,
Giaroni, C., Baj, A., 2021. Impact of Microbial Metabolites on Microbiota–Gut–Brain Axis
in Inflammatory Bowel Disease. Int J Mol Sci 22, 1623.
https://doi.org/10.3390/ijms22041623
Bao, C.H., Liu, P., Liu, H.R., Wu, L.Y., Shi, Y., Chen, W.F., Qin, W., Lu, Y., Zhang, J.Y., Jin, X.M.,
Wang, X.M., Zhao, J.M., Liu, X.M., Tian, J., Wu, H.G., 2015. Alterations in brain grey
matter structures in patients with Crohn's disease and their correlation with
psychological distress. Journal of Crohn's and Colitis. https://doi.org/10.1093/ecco-
jcc/jjv057

Bayless, D.W., Davis, C.-H.O., Yang, R., Wei, Y., de Andrade Carvalho, V.M., Knoedler, J.R., Yang,
T., Livingston, O., Lomvardas, A., Martins, G.J., Vicente, A.M., Ding, J.B., Luo, L., Shah,
N.M., 2023. A neural circuit for male sexual behavior and reward. Cell 186, 3862-
3881.e28. https://doi.org/10.1016/j.cell.2023.07.021
Bel, L.G.J., Vollebregt, A.M., Van der Meulen-de Jong, A.E., Fidder, H.H., Ten Hove, W.R., Vliet-
Vlieland, C.W., Ter Kuile, M.M., de Groot, H.E., Both, S., 2015. Sexual Dysfunctions in
Men and Women with Inflammatory Bowel Disease: The Influence of IBD-Related
Clinical Factors and Depression on Sexual Function. J Sex Med 12, 1557–1567.
https://doi.org/10.1111/jsm.12913
Benchimol, E.I., Bernstein, C.N., Bitton, A., Carroll, M.W., Singh, H., Otley, A.R., Vutcovici, M., El-
Matary, W., Nguyen, G.C., Griffiths, A.M., Mack, D.R., Jacobson, K., Mojaverian, N.,
Tanyingoh, D., Cui, Y., Nugent, Z.J., Coulombe, J., Targownik, L.E., Jones, J.L., Leddin, D.,
Murthy, S.K., Kaplan, G.G., 2017. Trends in epidemiology of pediatric inflammatory
bowel disease in Canada: Distributed network analysis of multiple population-based
provincial health administrative databases. American Journal of Gastroenterology 112,
1120–1134. https://doi.org/10.1038/ajg.2017.97
Benchimol, E.I., Manuel, D.G., Guttmann, A., Nguyen, G.C., Mojaverian, N., Quach, P., Mack,
D.R., 2014. Changing age demographics of inflammatory bowel disease in Ontario,
Canada: A population-based cohort study of epidemiology trends. Inflammatory Bowel
Diseases. https://doi.org/10.1097/MIB.0000000000000103
Benchimol, E.I., To, T., Griffiths, A.M., Rabeneck, L., Guttmann, A., 2011. Outcomes of pediatric
inflammatory bowel disease: Socioeconomic status disparity in a universal-access
healthcare system. Journal of Pediatrics 158.
https://doi.org/10.1016/j.jpeds.2010.11.039
Bishop, J., Lemberg, D., Day, A., 2014. Managing inflammatory bowel disease in adolescent
patients. Adolesc Health Med Ther 5, 1–13. https://doi.org/10.2147/AHMT.S37956
Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30, 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170
Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H.,
Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A.,
Brislawn, C.J., Brown, C.T., Callahan, B.J., Caraballo-Rodríguez, A.M., Chase, J., Cope,
E.K., Da Silva, R., Diener, C., Dorrestein, P.C., Douglas, G.M., Durall, D.M., Duvallet, C.,
Edwardson, C.F., Ernst, M., Estaki, M., Fouquier, J., Gauglitz, J.M., Gibbons, S.M., Gibson,
D.L., Gonzalez, A., Gorlick, K., Guo, J., Hillmann, B., Holmes, S., Holste, H., Huttenhower,
C., Huttley, G.A., Janssen, S., Jarmusch, A.K., Jiang, L., Kaehler, B.D., Kang, K.B., Keefe,
C.R., Keim, P., Kelley, S.T., Knights, D., Koester, I., Kosciolek, T., Kreps, J., Langille, M.G.I.,
Lee, J., Ley, R., Liu, Y.-X., Loftfield, E., Lozupone, C., Maher, M., Marotz, C., Martin, B.D.,
McDonald, D., McIver, L.J., Melnik, A.V., Metcalf, J.L., Morgan, S.C., Morton, J.T.,
Naimey, A.T., Navas-Molina, J.A., Nothias, L.F., Orchanian, S.B., Pearson, T., Peoples, S.L.,
Petras, D., Preuss, M.L., Pruesse, E., Rasmussen, L.B., Rivers, A., Robeson, M.S.,
Rosenthal, P., Segata, N., Shaffer, M., Shiffer, A., Sinha, R., Song, S.J., Spear, J.R.,
Swafford, A.D., Thompson, L.R., Torres, P.J., Trinh, P., Tripathi, A., Turnbaugh, P.J., Ul-
Hasan, S., van der Hooft, J.J.J., Vargas, F., Vázquez-Baeza, Y., Vogtmann, E., von Hippel,

M., Walters, W., Wan, Y., Wang, M., Warren, J., Weber, K.C., Williamson, C.H.D., Willis,
A.D., Xu, Z.Z., Zaneveld, J.R., Zhang, Y., Zhu, Q., Knight, R., Caporaso, J.G., 2019.
Reproducible, interactive, scalable and extensible microbiome data science using QIIME
2. Nat Biotechnol 37, 852–857. https://doi.org/10.1038/s41587-019-0209-9
Brain, C.E., Savage, M.O., 1994. Growth and puberty in chronic inflammatory bowel disease.
Baillieres Clin Gastroenterol 8, 83–100. https://doi.org/10.1016/s0950-3528(06)80020-5
Brooks, M.E., Kristensen, K., Benthem, K.J. van, Magnusson, A., Berg, C.W., Nielsen, A., Skaug,
H.J., Mächler, M., Bolker, B.M., 2017. glmmTMB Balances Speed and Flexibility Among
Packages for Zero-inflated Generalized Linear Mixed Modeling. The R Journal 9, 378–
400.
Brown, D.G., Murphy, M., Cadeddu, R., Bell, R., Weis, A., Chiaro, T., Klag, K., Morgan, J., Coon,
H., Stephens, W.Z., Bortolato, M., Round, J.L., 2024. Colitis reduces active social
engagement in mice and is ameliorated by supplementation with human microbiota
members. Nat Commun 15, 2769. https://doi.org/10.1038/s41467-024-46733-7
Buchfink, B., Xie, C., Huson, D.H., 2015. Fast and sensitive protein alignment using DIAMOND.
Nat Methods 12, 59–60. https://doi.org/10.1038/nmeth.3176
Caetano-Silva, M.E., Rund, L., Vailati-Riboni, M., Matt, S., Soto-Diaz, K., Beever, J., Allen, J.M.,
Woods, J.A., Steelman, A.J., Johnson, R.W., 2024. The emergence of inflammatory
microglia during gut inflammation is not affected by FFAR2 expression in intestinal
epithelial cells or peripheral myeloid cells. Brain Behav Immun 118, 423–436.
https://doi.org/10.1016/j.bbi.2024.03.016
Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., Holmes, S.P., 2016.
DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods 13,
581–583. https://doi.org/10.1038/nmeth.3869
Can, A., Dao, D.T., Arad, M., Terrillion, C.E., Piantadosi, S.C., Gould, T.D., 2011. The Mouse
Forced Swim Test. JoVE 3638. https://doi.org/10.3791/3638
Cao, Y., Dong, Q., Wang, D., Zhang, P., Liu, Y., Niu, C., 2022. microbiomeMarker: an
R/Bioconductor package for microbiome marker identification and visualization.
Bioinformatics 38, 4027–4029. https://doi.org/10.1093/bioinformatics/btac438
Chassaing, B., Aitken, J.D., Malleshappa, M., Vijay-Kumar, M., 2014. Dextran Sulfate Sodium
(DSS)-Induced Colitis in Mice. Curr Protoc Immunol 104, Unit-15.25.
https://doi.org/10.1002/0471142735.im1525s104
Chen, D., Zhou, C., Luo, Q., Chen, C., Liu, G., 2024. A Mendelian randomization study on causal
effects of inflammatory bowel disease on the risk of erectile dysfunction. Sci Rep 14,
2137. https://doi.org/10.1038/s41598-024-52712-1
Colldén, H., Landin, A., Wallenius, V., Elebring, E., Fändriks, L., Nilsson, M.E., Ryberg, H.,
Poutanen, M., Sjögren, K., Vandenput, L., Ohlsson, C., 2019. The gut microbiota is a
major regulator of androgen metabolism in intestinal contents. American Journal of
Physiology-Endocrinology and Metabolism 317, E1182–E1192.
https://doi.org/10.1152/ajpendo.00338.2019
Cotton, S., Clayton, C.A., Tropini, C., 2023. Microbial endocrinology: the mechanisms by which
the microbiota influences host sex steroids. Trends in Microbiology 31, 1131–1142.
https://doi.org/10.1016/j.tim.2023.03.010

Cryan, J.F., Dinan, T.G., 2012. Mind-altering microorganisms: The impact of the gut microbiota
1336   on brain and behaviour. Nature Reviews Neuroscience 13, 701–712.
1337   https://doi.org/10.1038/nrn3346
1338 Däbritz, J., Gerner, P., Enninger, A., Claßen, M., Radke, M., 2017. Inflammatory Bowel Disease in
1339   Childhood and Adolescence. Dtsch Arztebl Int 114, 331–338.
1340   https://doi.org/10.3238/arztebl.2017.0331
1341 De Wolfe, T.J., Wright, E.S., 2023. Multi-factorial examination of amplicon sequencing
1342   workflows from sample preparation to bioinformatic analysis. BMC Microbiol 23, 107.
1343   https://doi.org/10.1186/s12866-023-02851-8
1344 Deboer, M.D., Li, Y., 2011. Puberty Is Delayed in Male Mice With Dextran Sodium Sulfate Colitis
1345   Out of Proportion to Changes in Food Intake, Body Weight, and Serum Levels of Leptin.
1346   Pediatr Res 69, 34–39. https://doi.org/10.1203/PDR.0b013e3181ffee6c
1347 DeBoer, M.D., Li, Y., Cohn, S., 2010. Colitis causes delay in puberty in female mice out of
1348   proportion to changes in leptin and corticosterone. J Gastroenterol 45, 277–284.
1349   https://doi.org/10.1007/s00535-009-0192-x
1350 den Besten, G., van Eunen, K., Groen, A.K., Venema, K., Reijngoud, D.-J., Bakker, B.M., 2013. The
1351   role of short-chain fatty acids in the interplay between diet, gut microbiota, and host
1352   energy metabolism. Journal of Lipid Research 54, 2325–2340.
1353   https://doi.org/10.1194/jlr.R036012
1354 Dixon, P., 2003. VEGAN, a package of R functions for community ecology. Journal of Vegetation
1355   Science 14, 927–930. https://doi.org/10.1111/j.1654-1103.2003.tb02228.x
1356 Effendi, R.M.R.A., Anshory, M., Kalim, H., Dwiyana, R.F., Suwarsa, O., Pardo, L.M., Nijsten,
1357   T.E.C., Thio, H.B., 2022. Akkermansia muciniphila and Faecalibacterium prausnitzii in
1358   Immune-Related Diseases. Microorganisms 10, 2382.
1359   https://doi.org/10.3390/microorganisms10122382
1360 Eichele, D.D., Kharbanda, K.K., 2017. Dextran sodium sulfate colitis murine model: An
1361   indispensable tool for advancing our understanding of inflammatory bowel diseases
1362   pathogenesis. World J Gastroenterol 23, 6016–6029.
1363   https://doi.org/10.3748/wjg.v23.i33.6016
1364 Emge, J.R., Huynh, K., Miller, E.N., Kaur, M., Reardon, C., Barrett, K.E., Gareau, M.G., 2016.
1365   Modulation of the microbiota-gut-brain axis by probiotics in a murine model of
1366   inflammatory bowel disease. American Journal of Physiology-Gastrointestinal and Liver
1367   Physiology 310, G989–G998. https://doi.org/10.1152/ajpgi.00086.2016
1368 Erny, D., De Angelis, A.L.H., Jaitin, D., Wieghofer, P., Staszewski, O., David, E., Keren-Shaul, H.,
1369   Mahlakoiv, T., Jakobshagen, K., Buch, T., Schwierzeck, V., Utermöhlen, O., Chun, E.,
1370   Garrett, W.S., Mccoy, K.D., Diefenbach, A., Staeheli, P., Stecher, B., Amit, I., Prinz, M.,
1371   2015. Host microbiota constantly control maturation and function of microglia in the
1372   CNS. Nature Neuroscience 18, 965–977. https://doi.org/10.1038/nn.4030
1373 Erny, D., Dokalis, N., Mezö, C., Castoldi, A., Mossad, O., Staszewski, O., Frosch, M., Villa, M.,
1374   Fuchs, V., Mayer, A., Neuber, J., Sosat, J., Tholen, S., Schilling, O., Vlachos, A., Blank, T.,
1375   Agüero, M.G. de, Macpherson, A.J., Pearce, E.J., Prinz, M., 2021. Microbiota-derived
1376   acetate enables the metabolic fitness of the brain innate immune system during health
1377   and disease. Cell Metabolism 33, 2260-2276.e7.
1378   https://doi.org/10.1016/j.cmet.2021.10.010

Ervin, S.M., Li, H., Lim, L., Roberts, L.R., Liang, X., Mani, S., Redinbo, M.R., 2019. Gut microbial β-
glucuronidases reactivate estrogens as components of the estrobolome that reactivate
estrogens. Journal of Biological Chemistry 294, 18586–18599.
https://doi.org/10.1074/jbc.RA119.010950
Ewels, P., Magnusson, M., Lundin, S., Käller, M., 2016. MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics 32, 3047–3048.
https://doi.org/10.1093/bioinformatics/btw354
Flores, R., Shi, J., Fuhrman, B., Xu, X., Veenstra, T.D., Gail, M.H., Gajer, P., Ravel, J., Goedert, J.J.,
2012. Fecal microbial determinants of fecal and systemic estrogens and estrogen
metabolites: a cross-sectional study. Journal of Translational Medicine 10, 253.
https://doi.org/10.1186/1479-5876-10-253
Fusco, W., Lorenzo, M.B., Cintoni, M., Porcari, S., Rinninella, E., Kaitsas, F., Lener, E., Mele, M.C.,
Gasbarrini, A., Collado, M.C., Cammarota, G., Ianiro, G., 2023. Short-Chain Fatty-Acid-
Producing Bacteria: Key Components of the Human Gut Microbiota. Nutrients 15, 2211.
https://doi.org/10.3390/nu15092211
Gaidos, J.K.J., Naik, K., Dave, J., Yao, M., Hou, J.K., Cipher, D.J., Smith, A.D., Feagins, L.A., 2020.
High Prevalence of Male Sexual Dysfunction in a Prospective Multicenter VA
Inflammatory Bowel Disease Population. Crohns Colitis 360 2, otaa004.
https://doi.org/10.1093/crocol/otaa004
Gampierakis, I.-A., Koutmani, Y., Semitekolou, M., Morianos, I., Polissidis, A., Katsouda, A.,
Charalampopoulos, I., Xanthou, G., Gravanis, A., Karalis, K.P., 2021. Hippocampal neural
stem cells and microglia response to experimental inflammatory bowel disease (IBD).
Mol Psychiatry 26, 1248–1263. https://doi.org/10.1038/s41380-020-0651-6
Garcia-Hernandez, R., Cerdán Cerdá, A., Trouve Carpena, A., Drakesmith, M., Koller, K., Jones,
D.K., Canals, S., De Santis, S., 2022. Mapping microglia and astrocyte activation in vivo
using diffusion MRI. Science Advances 8, eabq2923.
https://doi.org/10.1126/sciadv.abq2923
Gu, Z., Eils, R., Schlesner, M., 2016. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics 32, 2847–2849.
https://doi.org/10.1093/bioinformatics/btw313
Günther, C., Rothhammer, V., Karow, M., Neurath, M., Winner, B., 2021. The Gut-Brain Axis in
Inflammatory Bowel Disease—Current and Future Perspectives. Int J Mol Sci 22, 8870.
https://doi.org/10.3390/ijms22168870
Gupta, N., Lustig, R.H., Andrews, H., Sylvester, F., Keljo, D., Goyal, A., Gokhale, R., Patel, A.S.,
Guthery, S., Leu, C.-S., 2020. Introduction to and Screening Visit Results of the
Multicenter Pediatric Crohn's Disease Growth Study. Inflammatory Bowel Diseases 26,
1945–1950. https://doi.org/10.1093/ibd/izaa023
Hadley, W., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
Hamden, J.E., Gray, K.M., Salehzadeh, M., Kachkovski, G.V., Forys, B.J., Ma, C., Austin, S.H.,
Soma, K.K., 2021. Steroid profiling of glucocorticoids in microdissected mouse brain
across development. Developmental Neurobiology 81, 189–206.
https://doi.org/10.1002/dneu.22808
Hamden, J.E., Salehzadeh, M., Jalabert, C., O'Leary, T.P., Snyder, J.S., Gomez-Sanchez, C.E.,
Soma, K.K., 2019. Measurement of 11-dehydrocorticosterone in mice, rats and

songbirds: Effects of age, sex and stress. General and Comparative Endocrinology 281,
173–182. https://doi.org/10.1016/j.ygcen.2019.05.018
Hammami, M.B., Mahadevan, U., 2020. Men With Inflammatory Bowel Disease: Sexual
Function, Fertility, Medication Safety, and Prostate Cancer. Am J Gastroenterol 115,
526–534. https://doi.org/10.14309/ajg.0000000000000515
Han, S., Treuren, W.V., Fischer, C.R., Merrill, B.D., DeFelice, B.C., Sanchez, J.M., Higginbottom,
S.K., Guthrie, L., Fall, L.A., Dodd, D., Fischbach, M.A., Sonnenburg, J.L., 2021. A
metabolomics pipeline for mechanistic interrogation of the gut microbiome. Nature 595,
415. https://doi.org/10.1038/s41586-021-03707-9
Handelsman, D.J., Ly, L.P., 2019. An Accurate Substitution Method To Minimize Left Censoring
Bias in Serum Steroid Measurements. Endocrinology 160, 2395–2400.
https://doi.org/10.1210/en.2019-00340
Harms, R.L., Fritz, F.J., Tobisch, A., Goebel, R., Roebroeck, A., 2017. Robust and fast nonlinear
optimization of diffusion MRI microstructure models. Neuroimage 155, 82–96.
https://doi.org/10.1016/j.neuroimage.2017.04.064
He, S., Li, H., Yu, Z., Zhang, F., Liang, S., Liu, H., Chen, H., Lü, M., 2021. The Gut Microbiome and
Sex Hormone-Related Diseases. Frontiers in Microbiology 12.
Herzer, M., Denson, L.A., Baldassano, R.N., Hommel, K.A., 2011. Patient and parent
psychosocial factors associated with health-related quality of life in pediatric
inflammatory bowel disease. Journal of Pediatric Gastroenterology and Nutrition 52,
295–299. https://doi.org/10.1097/MPG.0b013e3181f5714e
Hildebrand, H., Karlberg, J., Kristiansson, B., 1994. Longitudinal growth in children and
adolescents with inflammatory bowel disease. J Pediatr Gastroenterol Nutr 18, 165–173.
https://doi.org/10.1097/00005176-199402000-00008
Hodgkinson, K., El Abbar, F., Dobranowski, P., Manoogian, J., Butcher, J., Figeys, D., Mack, D.,
Stintzi, A., 2023. Butyrate's role in human health and the current progress towards its
clinical application to treat gastrointestinal disease. Clinical Nutrition 42, 61–75.
https://doi.org/10.1016/j.clnu.2022.10.024
Hou, J., Dodd, K., Nair, V.A., Rajan, S., Beniwal-Patel, P., Saha, S., Prabhakaran, V., 2020.
Alterations in brain white matter microstructural properties in patients with Crohn's
disease in remission. Scientific Reports. https://doi.org/10.1038/s41598-020-59098-w
Huerta-Cepas, J., Szklarczyk, D., Forslund, K., Cook, H., Heller, D., Walter, M.C., Rattei, T.,
Mende, D.R., Sunagawa, S., Kuhn, M., Jensen, L.J., von Mering, C., Bork, P., 2016.
eggNOG 4.5: a hierarchical orthology framework with improved functional annotations
for eukaryotic, prokaryotic and viral sequences. Nucleic Acids Research 44, D286–D293.
https://doi.org/10.1093/nar/gkv1248
Hyatt, D., Chen, G.-L., LoCascio, P.F., Land, M.L., Larimer, F.W., Hauser, L.J., 2010. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC
Bioinformatics 11, 119. https://doi.org/10.1186/1471-2105-11-119
Jiang, Y., Johnson, G.A., 2011. Microscopic diffusion tensor atlas of the mouse brain.
NeuroImage 56, 1235–1243. https://doi.org/10.1016/j.neuroimage.2011.03.031
Kanehisa, M., Goto, S., 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids
Research 28, 27–30. https://doi.org/10.1093/nar/28.1.27

Kaplan, G.G., Windsor, J.W., 2021. The four epidemiological stages in the global evolution of
inflammatory bowel disease. Nat Rev Gastroenterol Hepatol 18, 56–66.
https://doi.org/10.1038/s41575-020-00360-x
Kassambara, A., 2023. ggpubr: "ggplot2" Based Publication Ready Plots [WWW Document]. URL
https://rpkgs.datanovia.com/ggpubr/
Katoh, K., Misawa, K., Kuma, K., Miyata, T., 2002. MAFFT: a novel method for rapid multiple
sequence alignment based on fast Fourier transform. Nucleic Acids Research 30, 3059–
3066. https://doi.org/10.1093/nar/gkf436
Keller, M., Baum, M.J., Bakker, J., 2008. Olfactory Control of Sex-Recognition and Sexual
Behavior in Mice, in: Hurst, J.L., Beynon, R.J., Roberts, S.C., Wyatt, T.D. (Eds.), Chemical
Signals in Vertebrates 11. Springer, New York, NY, pp. 241–250.
https://doi.org/10.1007/978-0-387-73945-8_23
Kight, K.E., McCarthy, M.M., 2020. Androgens and the developing hippocampus. Biol Sex Differ
11, 30. https://doi.org/10.1186/s13293-020-00307-6
Kim, J., Pavlidis, P., Ciernia, A.V., 2024a. Development of a high-throughput pipeline to
characterize microglia morphological states at a single-cell resolution. eNeuro.
https://doi.org/10.1523/ENEURO.0014-24.2024
Kim, J., Sullivan, O., Lee, K., Jao, J., Tamayo, J., Madany, A.M., Wong, B., Ashwood, P., Ciernia,
A.V., 2024b. Repeated LPS induces training and tolerance of microglial responses across
brain regions. https://doi.org/10.1101/2024.04.08.588502
Kim, K., Choe, D., Song, Y., Kang, M., Lee, S.-G., Lee, D.-H., Cho, B.-K., 2021. Engineering
Bacteroides thetaiotaomicron to produce non-native butyrate based on a genome-scale
metabolic model-guided design. Metabolic Engineering 68, 174–186.
https://doi.org/10.1016/j.ymben.2021.10.005
Klaus, J., Reinshagen, M., Adler, G., Boehm, B., von Tirpitz, C., 2008. Bones and Crohn's:
Estradiol deficiency in men with Crohn's disease is not associated with reduced bone
mineral density. BMC Gastroenterol 8, 48. https://doi.org/10.1186/1471-230X-8-48
Kleine Bardenhorst, S., Vital, M., Karch, A., Rübsamen, N., 2022. Richness estimation in
microbiome data obtained from denoising pipelines. Computational and Structural
Biotechnology Journal 20, 508–520. https://doi.org/10.1016/j.csbj.2021.12.036
Korpela, K., Kallio, S., Salonen, A., Hero, M., Kukkonen, A.K., Miettinen, P.J., Savilahti, E., Kohva,
E., Kariola, L., Suutela, M., Tarkkanen, A., de Vos, W.M., Raivio, T., Kuitunen, M., 2021.
Gut microbiota develop towards an adult profile in a sex-specific manner during
puberty. Sci Rep 11, 23297. https://doi.org/10.1038/s41598-021-02375-z
Langmead, B., Salzberg, S.L., 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9,
357–359. https://doi.org/10.1038/nmeth.1923
Lavelle, A., Sokol, H., 2020. Gut microbiota-derived metabolites as key actors in inflammatory
bowel disease. Nature Reviews Gastroenterology and Hepatology 17, 223–237.
https://doi.org/10.1038/s41575-019-0258-z
Li, D., Liu, C.-M., Luo, R., Sadakane, K., Lam, T.-W., 2015. MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics 31, 1674–1676. https://doi.org/10.1093/bioinformatics/btv033
Li, D., Liu, R., Wang, M., Peng, R., Fu, S., Fu, A., Le, J., Yao, Q., Yuan, T., Chi, H., Mu, X., Sun, T.,
Liu, H., Yan, P., Wang, S., Cheng, S., Deng, Z., Liu, Z., Wang, G., Li, Y., Liu, T., 2022. 3β-

Hydroxysteroid dehydrogenase expressed by gut microbes degrades testosterone and is
linked to depression in males. Cell Host & Microbe 30, 329-339.e5.
https://doi.org/10.1016/j.chom.2022.01.001
Lv, K., Fan, Y.H., Xu, L., Xu, M.S., 2017. Brain changes detected by functional magnetic
resonance imaging and spectroscopy in patients with Crohn's disease. World Journal of
Gastroenterology. https://doi.org/10.3748/wjg.v23.i20.3607
Machiels, K., Joossens, M., Sabino, J., De Preter, V., Arijs, I., Eeckhaut, V., Ballet, V., Claes, K.,
Van Immerseel, F., Verbeke, K., Ferrante, M., Verhaegen, J., Rutgeerts, P., Vermeire, S.,
2014. A decrease of the butyrate-producing species Roseburia hominis and
Faecalibacterium prausnitzii defines dysbiosis in patients with ulcerative colitis. Gut 63,
1275–1283. https://doi.org/10.1136/gutjnl-2013-304833
Mackner, L.M., Greenley, R.N., Szigethy, E., Herzer, M., Deer, K., Hommel, K.A., 2013.
Psychosocial Issues in Pediatric Inflammatory Bowel Disease: A Clinical Report of the
North American Society for Pediatric Gastroenterology, Hepatology and Nutrition. J
Pediatr Gastroenterol Nutr 56, 449–458.
https://doi.org/10.1097/MPG.0b013e3182841263
Maffei, S., Forini, F., Canale, P., Nicolini, G., Guiducci, L., 2022. Gut Microbiota and Sex
Hormones: Crosstalking Players in Cardiometabolic and Cardiovascular Disease.
International Journal of Molecular Sciences 23, 7154.
https://doi.org/10.3390/ijms23137154
Malkesman, O., Scattoni, M.L., Paredes, D., Tragon, T., Pearson, B., Shaltiel, G., Chen, G.,
Crawley, J.N., Manji, H.K., 2010. The Female Urine Sniffing Test: A Novel Approach for
Assessing Reward-Seeking Behavior in Rodents. Biol Psychiatry 67, 864–871.
https://doi.org/10.1016/j.biopsych.2009.10.018
Mandal, S., Van Treuren, W., White, R.A., Eggesbø, M., Knight, R., Peddada, S.D., 2015. Analysis
of composition of microbiomes: a novel method for studying microbial composition.
Microb Ecol Health Dis 26, 27663. https://doi.org/10.3402/mehd.v26.27663
Markle, J.G.M., Frank, D.N., Mortin-Toth, S., Robertson, C.E., Feazel, L.M., Rolle-Kampczyk, U.,
von Bergen, M., McCoy, K.D., Macpherson, A.J., Danska, J.S., 2013. Sex Differences in
the Gut Microbiome Drive Hormone-Dependent Regulation of Autoimmunity. Science
339, 1084–1088. https://doi.org/10.1126/science.1233521
Matisz, C.E., Vicentini, F.A., Hirota, S.A., Sharkey, K.A., Gruber, A.J., 2020. Behavioral
adaptations in a relapsing mouse model of colitis. Physiol Behav 216, 112802.
https://doi.org/10.1016/j.physbeh.2020.112802
McMurdie, P.J., Holmes, S., 2013. phyloseq: An R Package for Reproducible Interactive Analysis
and Graphics of Microbiome Census Data. PLOS ONE 8, e61217.
https://doi.org/10.1371/journal.pone.0061217
Mertens, K.L., Kalsbeek, A., Soeters, M.R., Eggink, H.M., 2017. Bile Acid Signaling Pathways from
the Enterohepatic Circulation to the Central Nervous System. Frontiers in Neuroscience
11.
Min, S.H., Zhou, J., 2021. smplot: An R Package for Easy and Elegant Data Visualization. Front.
Genet. 12. https://doi.org/10.3389/fgene.2021.802894
Mirzaei, R., Bouzari, B., Hosseini-Fard, S.R., Mazaheri, M., Ahmadyousefi, Y., Abdi, M., Jalalifar,
S., Karimitabar, Z., Teimoori, A., Keyvani, H., Zamani, F., Yousefimashouf, R., Karampoor,

S., 2021. Role of microbiota-derived short-chain fatty acids in nervous system disorders.
Biomed Pharmacother 139, 111661. https://doi.org/10.1016/j.biopha.2021.111661
Mitchell, R.W., On, N.H., Del Bigio, M.R., Miller, D.W., Hatch, G.M., 2011. Fatty acid transport
protein expression in human brain and potential role in fatty acid transport across
human brain microvessel endothelial cells. J Neurochem 117, 735–746.
https://doi.org/10.1111/j.1471-4159.2011.07245.x
Munyaka, P.M., Rabbi, M.F., Khafipour, E., Ghia, J., 2016. Acute dextran sulfate sodium (DSS)-
induced colitis promotes gut microbial dysbiosis in mice. J. Basic Microbiol. 56, 986–998.
https://doi.org/10.1002/jobm.201500726
Naser, S.A., Sagramsingh, S.R., Naser, A.S., Thanigachalam, S., 2014. Mycobacterium avium
subspecies paratuberculosis causes Crohn's disease in some inflammatory bowel
disease patients. World J Gastroenterol 20, 7403–7415.
https://doi.org/10.3748/wjg.v20.i23.7403
Nasser, M., Haider, A., Saad, F., Kurtz, W., Doros, G., Fijak, M., Vignozzi, L., Gooren, L., 2015.
Testosterone therapy in men with Crohn's disease improves the clinical course of the
disease: data from long-term observational registry study. Horm Mol Biol Clin Investig
22, 111–117. https://doi.org/10.1515/hmbci-2015-0014
Neuman, H., Debelius, J.W., Knight, R., Koren, O., 2015. Microbial endocrinology: the interplay
between the microbiota and the endocrine system. FEMS Microbiology Reviews 39,
509–521. https://doi.org/10.1093/femsre/fuu010
Ng, K.M., Pannu, S., Liu, S., Burckhardt, J.C., Hughes, T., Van Treuren, W., Nguyen, J., Naqvi, K.,
Nguyen, B., Clayton, C.A., Pepin, D.M., Collins, S.R., Tropini, C., 2023. Single-strain
behavior predicts responses to environmental pH and osmolality in the gut microbiota.
mBio 14, e00753-23. https://doi.org/10.1128/mbio.00753-23
Ng, K.M., Tropini, C., 2021. Visualization of Gut Microbiota-host Interactions via Fluorescence In
Situ Hybridization, Lectin Staining, and Imaging. JoVE (Journal of Visualized Experiments)
e62646. https://doi.org/10.3791/62646
Ng, S.C., Shi, H.Y., Hamidi, N., Underwood, F.E., Tang, W., Benchimol, E.I., Panaccione, R.,
Ghosh, S., Wu, J.C.Y., Chan, F.K.L., Sung, J.J.Y., Kaplan, G.G., 2017. Worldwide incidence
and prevalence of inflammatory bowel disease in the 21st century: a systematic review
of population-based studies. The Lancet. https://doi.org/10.1016/S0140-
6736(17)32448-0
Nyuyki, K.D., Cluny, N.L., Swain, M.G., Sharkey, K.A., Pittman, Q.J., 2018. Altered brain
excitability and increased anxiety in mice with experimental colitis: Consideration of
hyperalgesia and sex differences. Frontiers in Behavioral Neuroscience 12, 1–11.
https://doi.org/10.3389/fnbeh.2018.00058
Patel, J., Chaudhary, H., Rajput, K., Parekh, B., Joshi, R., 2023. Assessment of gut microbial β-
glucuronidase and β-glucosidase activity in women with polycystic ovary syndrome. Sci
Rep 13, 11967. https://doi.org/10.1038/s41598-023-39168-5
Paulson, J.N., Stine, O.C., Bravo, H.C., Pop, M., 2013. Differential abundance analysis for
microbial marker-gene surveys. Nat Methods 10, 1200–1202.
https://doi.org/10.1038/nmeth.2658

Price, M.N., Dehal, P.S., Arkin, A.P., 2010. FastTree 2 – Approximately Maximum-Likelihood
Trees for Large Alignments. PLOS ONE 5, e9490.
https://doi.org/10.1371/journal.pone.0009490
Puente-Sánchez, F., García-García, N., Tamames, J., 2020. SQMtools: automated processing and
visual analysis of 'omics data with R and anvi'o. BMC Bioinformatics 21, 358.
https://doi.org/10.1186/s12859-020-03703-2
Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., Glöckner, F.O.,
2013. The SILVA ribosomal RNA gene database project: improved data processing and
web-based tools. Nucleic Acids Res 41, D590–D596.
https://doi.org/10.1093/nar/gks1219
R Core team, 2021. R: A language and environment for statistical computing [WWW
Document]. URL https://www.r-project.org/ (accessed 4.1.23).
Rastelli, D., Robinson, A., Lagomarsino, V.N., Matthews, L.T., Hassan, R., Perez, K., Dan, W., Yim,
P.D., Mixer, M., Prochera, A., Shepherd, A., Sun, L., Hall, K., Ballou, S., Lembo, A., Nee, J.,
Rao, M., 2022. Diminished androgen levels are linked to irritable bowel syndrome and
cause bowel dysfunction in mice. J Clin Invest 132, e150789.
https://doi.org/10.1172/JCI150789
Robinson, M.D., McCarthy, D.J., Smyth, G.K., 2009. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26, 139.
https://doi.org/10.1093/bioinformatics/btp616
Roda, G., Chien Ng, S., Kotze, P.G., Argollo, M., Panaccione, R., Spinelli, A., Kaser, A., Peyrin-
Biroulet, L., Danese, S., 2020. Crohn's disease. Nat Rev Dis Primers 6, 1–19.
https://doi.org/10.1038/s41572-020-0156-2
Rosen, M.J., Dhawan, A., Saeed, S.A., 2015. Inflammatory Bowel Disease in Children and
Adolescents. JAMA Pediatr 169, 1053–1060.
https://doi.org/10.1001/jamapediatrics.2015.1982
Salvo, E., Stokes, P., Keogh, C.E., Brust-Mascher, I., Hennessey, C., Knotts, T.A., Sladek, J.A.,
Rude, K.M., Swedek, M., Rabasa, G., Gareau, M.G., 2020. A murine model of pediatric
inflammatory bowel disease causes microbiota-gut-brain axis deficits in adulthood. Am J
Physiol Gastrointest Liver Physiol 319, G361–G374.
https://doi.org/10.1152/ajpgi.00177.2020
Seemann, T., 2024. tseemann/barrnap.
Silva, Y.P., Bernardi, A., Frozza, R.L., 2020. The Role of Short-Chain Fatty Acids From Gut
Microbiota in Gut-Brain Communication. Frontiers in Endocrinology 11.
https://doi.org/10.3389/fendo.2020.00025
Sinha, S.R., Haileselassie, Y., Nguyen, L.P., Tropini, C., Wang, M., Becker, L.S., Sim, D., Jarr, K.,
Spear, E.T., Singh, G., Namkoong, H., Bittinger, K., Fischbach, M.A., Sonnenburg, J.L.,
Habtezion, A., 2020. Dysbiosis-Induced Secondary Bile Acid Deficiency Promotes
Intestinal Inflammation. Cell Host Microbe 27, 659-670.e5.
https://doi.org/10.1016/j.chom.2020.01.021
Smith, S.M., Jenkinson, M., Woolrich, M.W., Beckmann, C.F., Behrens, T.E.J., Johansen-Berg, H.,
Bannister, P.R., De Luca, M., Drobnjak, I., Flitney, D.E., Niazy, R.K., Saunders, J., Vickers,
J., Zhang, Y., De Stefano, N., Brady, J.M., Matthews, P.M., 2004. Advances in functional

and structural MR image analysis and implementation as FSL. Neuroimage 23 Suppl 1,
S208-219. https://doi.org/10.1016/j.neuroimage.2004.07.051
Sokol, H., Pigneur, B., Watterlot, L., Lakhdari, O., Bermúdez-Humarán, L.G., Gratadoux, J.J.,
Blugeon, S., Bridonneau, C., Furet, J.P., Corthier, G., Grangette, C., Vasquez, N., Pochart,
P., Trugnan, G., Thomas, G., Blottière, H.M., Doré, J., Marteau, P., Seksik, P., Langella, P.,
2008. Faecalibacterium prausnitzii is an anti-inflammatory commensal bacterium
identified by gut microbiota analysis of Crohn disease patients. Proceedings of the
National Academy of Sciences of the United States of America 105, 16731–16736.
https://doi.org/10.1073/pnas.0804812105
Sroor, H.M., Hassan, A.M., Zenz, G., Valadez-Cosmes, P., Farzi, A., Holzer, P., El-Sharif, A.,
Gomaa, F.A.-Z.M., Kargl, J., Reichmann, F., 2019. Experimental colitis reduces microglial
cell activation in the mouse brain without affecting microglial cell numbers. Sci Rep 9,
20217. https://doi.org/10.1038/s41598-019-56859-0
Stahl, M., Graef, F.A., Vallance, B.A., 2017. Mouse Models for Campylobacter jejuni Colonization
and Infection, in: Butcher, J., Stintzi, A. (Eds.), Campylobacter Jejuni: Methods and
Protocols. Springer, New York, NY, pp. 171–188. https://doi.org/10.1007/978-1-4939-
6536-6_15
Sui, Y., Wu, J., Chen, J., 2021. The Role of Gut Microbial β-Glucuronidase in Estrogen
Reactivation and Breast Cancer. Front Cell Dev Biol 9, 631552.
https://doi.org/10.3389/fcell.2021.631552
Sullivan, O., Ciernia, A.V., 2022. Work hard, play hard: how sexually differentiated microglia
work to shape social play and reproductive behavior. Frontiers in Behavioral
Neuroscience 16.
Talley, S., Valiauga, R., Anderson, L., Cannon, A.R., Choudhry, M.A., Campbell, E.M., 2021. DSS-
induced inflammation in the colon drives a proinflammatory signature in the brain that
is ameliorated by prophylactic treatment with the S100A9 inhibitor paquinimod. Journal
of Neuroinflammation 18, 263. https://doi.org/10.1186/s12974-021-02317-6
Tamames, J., Puente-Sánchez, F., 2019. SqueezeMeta, A Highly Portable, Fully Automatic
Metagenomic Analysis Pipeline. Front. Microbiol. 9.
https://doi.org/10.3389/fmicb.2018.03349
Terstege, D.J., Oboh, D.O., Epp, J.R., 2022. FASTMAP: Open-Source Flexible Atlas Segmentation
Tool for Multi-Area Processing of Biological Images. eNeuro 9, ENEURO.0325-21.2022.
https://doi.org/10.1523/ENEURO.0325-21.2022
Thavamani, A., Umapathi, K.K., Khatana, J., Gulati, R., 2019. Burden of Psychiatric Disorders
among Pediatric and Young Adults with Inflammatory Bowel Disease: A Population-
Based Analysis. Pediatr Gastroenterol Hepatol Nutr 22, 527–535.
https://doi.org/10.5223/pghn.2019.22.6.527
Thibault, R., Blachier, F., Darcy-Vrillon, B., de Coppet, P., Bourreille, A., Segain, J.-P., 2010.
Butyrate utilization by the colonic mucosa in inflammatory bowel diseases: a transport
deficiency. Inflamm Bowel Dis 16, 684–695. https://doi.org/10.1002/ibd.21108
Thion, M.S., Low, D., Silvin, A., Chen, J., Grisel, P., Schulte-Schrepping, J., Blecher, R., Ulas, T.,
Squarzoni, P., Hoeffel, G., Coulpier, F., Siopi, E., David, F.S., Scholz, C., Shihui, F., Lum, J.,
Amoyo, A.A., Larbi, A., Poidinger, M., Buttgereit, A., Lledo, P.-M., Greter, M., Chan,
J.K.Y., Amit, I., Beyer, M., Schultze, J.L., Schlitzer, A., Pettersson, S., Ginhoux, F., Garel, S.,

2017. Microbiome Influences Prenatal and Adult Microglia in a Sex-Specific Manner. Cell
1–17. https://doi.org/10.1016/j.cell.2017.11.042

Tigas, S., Tsatsoulis, A., 2012. Endocrine and metabolic manifestations in inflammatory bowel
disease. Ann Gastroenterol 25, 37–44.

Torres, P.J., Siakowska, M., Banaszewska, B., Pawelczyk, L., Duleba, A.J., Kelley, S.T., Thackray,
V.G., 2018. Gut Microbial Diversity in Women With Polycystic Ovary Syndrome
Correlates With Hyperandrogenism. J Clin Endocrinol Metab 103, 1502–1511.
https://doi.org/10.1210/jc.2017-02153

Vailati-Riboni, M., Rund, L., Caetano-Silva, M.E., Hutchinson, N.T., Wang, S.S., Soto-Díaz, K.,
Woods, J.A., Steelman, A.J., Johnson, R.W., 2022. Dietary Fiber as a Counterbalance to
Age-Related Microglial Cell Dysfunction. Front Nutr 9, 835824.
https://doi.org/10.3389/fnut.2022.835824

van Griethuysen, J.J.M., Fedorov, A., Parmar, C., Hosny, A., Aucoin, N., Narayan, V., Beets-Tan,
R.G.H., Fillion-Robin, J.-C., Pieper, S., Aerts, H.J.W.L., 2017. Computational Radiomics
System to Decode the Radiographic Phenotype. Cancer Res 77, e104–e107.
https://doi.org/10.1158/0008-5472.CAN-17-0339

Venegas, D.P., Fuente, M.K.D. la, Landskron, G., González, M.J., Quera, R., Dijkstra, G., Harmsen,
H.J.M., Faber, K.N., Hermoso, M.A., 2019. Short Chain Fatty Acids (SCFAs)-Mediated Gut
Epithelial and Immune Regulation and Its Relevance for Inflammatory Bowel Diseases.
Frontiers in Immunology 10, 277. https://doi.org/10.3389/fimmu.2019.00277

Vogel Ciernia, A., Careaga, M., LaSalle, J.M., Ashwood, P., 2018. Microglia from offspring of
dams with allergic asthma exhibit epigenomic alterations in genes dysregulated in
autism. Glia 66, 505–521. https://doi.org/10.1002/glia.23261

Vogel-Ciernia, A., Wood, M.A., 2014. Examining object location and object recognition memory
in mice. Current Protocols in Neuroscience 2014, 8.31.1-8.31.17.
https://doi.org/10.1002/0471142301.ns0831s69

Wang, Q., Garrity, G.M., Tiedje, J.M., Cole, J.R., 2007. Naïve Bayesian Classifier for Rapid
Assignment of rRNA Sequences into the New Bacterial Taxonomy. Applied and
Environmental Microbiology 73, 5261–5267. https://doi.org/10.1128/AEM.00062-07

Weger, B.D., Gobet, C., Yeung, J., Martin, E., Jimenez, S., Betrisey, B., Foata, F., Berger, B.,
Balvay, A., Foussier, A., Charpagne, A., Boizet-Bonhoure, B., Chou, C.J., Naef, F., Gachon,
F., 2019. The Mouse Microbiome Is Required for Sex-Specific Diurnal Rhythms of Gene
Expression and Metabolism. Cell Metab 29, 362-382.e8.
https://doi.org/10.1016/j.cmet.2018.09.023

Wenzel, T.J., Gates, E.J., Ranger, A.L., Klegeris, A., 2020. Short-chain fatty acids (SCFAs) alone or
in combination regulate select immune functions of microglia-like cells. Mol Cell
Neurosci 105, 103493. https://doi.org/10.1016/j.mcn.2020.103493

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund, G.,
Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M., Müller,
K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C.,
Woo, K., Yutani, H., 2019. Welcome to the Tidyverse. Journal of Open Source Software
4, 1686. https://doi.org/10.21105/joss.01686

Wirtz, S., Neurath, M.F., 2007. Mouse models of inflammatory bowel disease. Advanced Drug
Delivery Reviews, Prediction of Therapeutic and Drug Delivery Outcomes Using Animal
Models 59, 1073–1083. https://doi.org/10.1016/j.addr.2007.07.003
Wong, E.O.-Y., Brownlie, E.J.E., Ng, K.M., Kathirgamanathan, S., Yu, F.B., Merrill, B.D., Huang,
K.C., Martin, A., Tropini, C., Navarre, W.W., 2022. The CIAMIB: a Large and Metabolically
Diverse Collection of Inflammation-Associated Bacteria from the Murine Gut. mBio 13,
e02949-21. https://doi.org/10.1128/mbio.02949-21
Yamane, T., Kitamura, Y., Terada, N., Matsumoto, K., 1986. Proliferative response of seminal
vesicle cells to androgen and estrogen in neonatally castrated mice. Journal of Steroid
Biochemistry 24, 703–708. https://doi.org/10.1016/0022-4731(86)90846-0
Yang, M., Crawley, J.N., 2009. Simple behavioral assessment of mouse olfaction. Curr Protoc
Neurosci Chapter 8, Unit 8.24. https://doi.org/10.1002/0471142301.ns0824s48
Ye, Y., Pang, Z., Chen, W., Ju, S., Zhou, C., 2015. The epidemiology and risk factors of
inflammatory bowel disease. Int J Clin Exp Med 8, 22529–22542.
Yi, S.Y., Barnett, B.R., Torres-Velázquez, M., Zhang, Y., Hurley, S.A., Rowley, P.A., Hernando, D.,
Yu, J.-P.J., 2019. Detecting Microglial Density With Quantitative Multi-Compartment
Diffusion MRI. Front Neurosci 13, 81. https://doi.org/10.3389/fnins.2019.00081
Zhang, H., Schneider, T., Wheeler-Kingshott, C.A., Alexander, D.C., 2012. NODDI: practical in
vivo neurite orientation dispersion and density imaging of the human brain. Neuroimage
61, 1000–1016. https://doi.org/10.1016/j.neuroimage.2012.03.072
Zhang, J., Wei, S., Zeng, Q., Wu, X., Gan, H., 2021. Prevalence and risk factors of sexual
dysfunction in patients with inflammatory bowel disease: systematic review and meta-
analysis. Int J Colorectal Dis 36, 2027–2038. https://doi.org/10.1007/s00384-021-03958-
y
Zhao, S., Wang, J., Liu, Y., Luo, L., Zhu, Z., Li, E., Luo, J., Zhao, Z., 2019. Inflammatory Bowel
Diseases Were Associated With Risk of Sexual Dysfunction in Both Sexes: A Meta-
analysis. Inflamm Bowel Dis 25, 699–707. https://doi.org/10.1093/ibd/izy345
Zhou, Y., Ji, G., Yang, X., Chen, Z., Zhou, L., 2023. Behavioral abnormalities in C57BL/6 mice with
chronic ulcerative colitis induced by DSS. BMC Gastroenterology 23, 84.
https://doi.org/10.1186/s12876-023-02718-2
Ziętek, M., Celewicz, Z., Szczuko, M., 2021. Short-Chain Fatty Acids, Maternal Microbiota and
Metabolism in Pregnancy. Nutrients 13, 1244. https://doi.org/10.3390/nu13041244
1757
1758

Supplemental Figure 1. DSS treatment did not alter distance travelled or exploration behavior. There are no significant main effects of treatment or interactions with treatment and sex. *A*. Total distance traveled in meters (m) in the A. elevated plus maze n=10-13/group, *B*. light-dark box task n=11-13/group and *C*. open field n=11-13/group. Total investigation time in seconds for object location memory *D*. testing and n=6-13/group. *E*. training n=6-13/group. *F*. Discrimination Index for OLM training n=6-13/group. *G*. Total investigation time in the 3 chamber social task in seconds. n=11-13/group. *H*. Total distance travelled in meters for the three chamber social task. n=11-13/group. *I*. Total investigation time in the 3 chamber novel task in seconds. n=10-13/group. *H*. Total distance travelled in meters for the three chamber novel task. n=10-13/group. Each bar and error bar represent the mean ± SEM.

Supplemental Figure 2. Effects of DSS treatment on male and female steroid levels. *A*. Three days after the last DSS treatment (P71), male serum progesterone levels were not significantly different between VEH and DSS treated males. n=4/group. *B*. Following the last behavioural test (P110), male serum progesterone levels were not significantly different between VEH and DSS treated males. n=12/group. *C-D*. Following the last behavioural test (P110), female serum progesterone (C) and testosterone (D) levels were not significantly different between VEH and DSS treated females. n=10-12/group. Androstenedione levels were non-detectable in VEH and DSS treated females. Each bar and error bar represent the mean ± SEM.

Supplemental Figure 3. Relative abundance of select microbial taxa correlate with cecal SCFA and serum testosterone concentrations acutely following DSS. *A.* Heatmap of differentially abundant gut bacterial taxa identified by differential abundance testing at P71. n=4-5/group. *B.* Relative abundance of Bacillota, *Lactobacillus*, *Eubacterium* xylanophilum group, and *Akkermansia muciniphila*. n=4-5/group. *C.* Correlation of relative abundance of Bacillota, *Roseburia,* and *Lactobacillus* vs. acetic and butyric acid concentrations in cecal contents at P71. *R* values from Spearman's correlation. n=5-8/group. *D.* Correlation of relative abundance of the species *Bacteroides thetaiotaomicron* vs. serum testosterone levels at P71. n=4/group. Error bars in bar plots represent SEM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Supplemental Figure 4. Microglial Morphology Analysis. A. Correlations between individual microglial morphology metrics. B. Percentage of variance explained by each principal component (PC) after clustering the 27 features. The first 3 PCs capture ~85% of the total variance and were

used for k-means clustering. C. The correlation between each PC and individual metrics. D. The silhouette plot reveals an optimal number of 4 clusters. E. K means clustering of 4 morphologyc clusters. F. Clusters were assigned morphology names based on their relationship to each individual metric. G. Percentage of hypothalamic microglia in each cluster between treatment groups for females and males. No significant differences were identified between treatments for either sex. n=2-5/group H. Percentage of cerebral cortical microglia in each cluster between treatment groups for females and males. No significant differences were identified between treatments for either sex. n=3-5/group.

Supplemental Figure 5. Gene Expression is Not Altered in Hippocampal Tissue from DSS and VEH treated male and female mice. Expression by RT-qPCR for *Tlr4* in A. females and B. males. Expression by RT-qPCR for *Nod1* in C. females and D. males. Expression by RT-qPCR for *Nod2* in E. females and F. males. Expression by RT-qPCR for *Tnfa* in G. females and H. males. Expression by RT-qPCR for *Il1b* in I. females and J. males. Expression by RT-qPCR for *Il6* in K. females and L. males. Fold change calculated to VEH controls within each sex for each gene. n=6-8/group. Each bar and error bar represent the mean ± SEM.

Supplemental Figure 6. Orientation dispersion index (ODI) was calculated for the thalamus, frontal association cortex, and hypothalamus. Data points are mean ODI values from each sample. A significant effect of treatment was identified in the thalamus, and post-hoc tests found a significant increase for DSS females compared to VEH females. n= 25-28/group ** $p < 0.01$, Fisher's LSD test.