1 **Title**: Plant hormone manipulation impacts salt spray tolerance, which preempts herbivory as a
2 driver of local adaptation in the yellow monkeyflower, *Mimulus guttatus*
3

4 **Authors**: Katherine Toll[1,2*†], Megan Blanchard[3,4†], Anna Scharnagl[5], Liza M. Holeski[3], and
5 David B. Lowry[1,6]
6
Affiliations:
[1] Department of Plant Biology, Michigan State University, East Lansing, MI, 48824
[2] Department of Biological Sciences, University of South Carolina, Columbia, SC, 29208
[3] Department of Biological Sciences and Center for Adaptable Western Landscapes, Northern
Arizona University, Flagstaff, AZ, 86011
[4] Museum of Natural History, University of Colorado Boulder, Boulder, CO, 80309
[5] Department of Integrative Biology, University of California, Berkeley, Berkeley, CA, 94720
[6] Plant Resilience Institute, Michigan State University, East Lansing, MI, 48824
[†] These authors contributed equally to this work.
[*] Corresponding author email: tollkath@gmail.com; mailing address: Department of Biological
Sciences, University of South Carolina, 715 Sumter Street, Columbia, SC 29208

18
Author contributions: Katherine Toll: Methodology, Investigation, Formal Analysis, Data
Curation, Writing - Original Draft, Writing - Review & Editing, Visualization. Megan
Blanchard: Methodology, Investigation, Formal Analysis, Data Curation, Writing - Original
Draft, Writing - Review & Editing, Visualization. Anna Scharnagl: Investigation. Liza Holeski:
Conceptualization, Methodology, Writing - Review & Editing, Resources, Supervision, Funding
acquisition. David B. Lowry: Conceptualization, Methodology, Writing - Review & Editing,
Resources, Supervision, Funding acquisition.
26

Acknowledgements: We would like to thank Ben Blackman and Jack Collichio for germinating
seeds at UC Berkeley. This work would not have been possible without the support of staff at the
Bodega Marine Lab and Reserve, particularly Jackie Sones, Luis Morales, and Al Carranza, who
facilitated field work and use of the greenhouse and lab. Michael Gillogly and Michelle Halbur
of Pepperwood Preserve were instrumental to this research through their support of our work at
the Preserve, including performing emergency plot-monitoring when the preserve temporarily
closed to visitors due to the COVID-19 pandemic. Isabella Johnson and Billie Fraser assisted
with chemical sample processing. This work was funded by National Science Foundation
Division of Integrative Organismal Systems Grants to DBL and LMH (IOS-1855927) and DBL
(IOS-2153100).

37
38 **Conflict of Interest Statement:** The authors declare no conflict of interest.

Abstract

A major challenge in evolutionary biology is identifying the selective agents and phenotypes

underlying local adaptation. Local adaptation along environmental gradients may be driven by

trade-offs in allocation to reproduction, growth, and herbivore resistance. To identify

environmental agents of selection and their phenotypic targets, we performed a manipulative

field reciprocal transplant experiment with coastal perennial and inland annual ecotypes of the

common yellow monkeyflower (*Mimulus guttatus*). We manipulated herbivory with exclosures

built in the field and exogenously manipulated gibberellin and jasmonic acid to shift allocation of

plant resources among growth, reproduction, and herbivore resistance. Our hormone treatments

influenced the timing of allocation to reproduction and allocation to phytochemical defense, but

this shift was small relative to ecotype differences in allocation. Herbivore exclosures reduced

herbivory and increased fitness of plants at the coastal site. However, this reduction in herbivory

did not decrease the homesite advantage of coastal perennials. Unexpectedly, we found that the

application of exogenous gibberellin increased mortality due to salt spray at the coastal site for

both ecotypes. Our results suggest that divergence in salt spray tolerance, potentially mediated

by ecotype differences in gibberellin synthesis or bioactivity, is a strong driver of local

adaptation and preempts any impacts of herbivory in coastal habitats that experience salt spray.

Key words: local adaptation, herbivory, salt spray, gibberellin, monkeyflower

Introduction

Organisms experience dramatically different environmental conditions throughout their geographic ranges. Spatial gradients in abiotic factors, such as temperature, salinity, and water availability, as well as biotic factors, such as the presence of competitors, predators, and mutualists, can generate divergent natural selection (Kawecki & Ebert, 2004; Maron *et al.*, 2014; López-Goldar & Agrawal, 2021). This divergent selection can in turn lead to evolutionary responses in traits that increase fitness in local environments, and result in the evolution of local adaptation (Clausen *et al.*, 1940; Kawecki & Ebert, 2004; Leimu & Fischer, 2008; Hereford, 2009; Wadgymar *et al.*, 2022). Identifying the causal environmental factors contributing to adaptation is a major challenge because environmental conditions often co-vary and thus, experimental manipulations are necessary to identify the environmental agents of selection (Briscoe Runquist *et al.*, 2020; Hargreaves *et al.*, 2020). Likewise, the phenotypic targets of selection are challenging to identify because traits are often highly correlated, so approaches that minimize trait correlations or manipulate trait variation independently of other traits are necessary to identify adaptive traits (Wadgymar *et al.*, 2017, 2022). Despite their importance, experiments that simultaneously manipulate putative environmental selective agents and their phenotypic targets are uncommon (Wadgymar *et al.*, 2017, 2022). These field experiments are critical because the interactions of experimental manipulations and environmental factors are complex and can have unexpected outcomes. In this study, we isolate the effect of a putative selective agent, herbivory, at sites that vary in two abiotic factors, salt spray and soil moisture, and manipulate trait variation using hormone applications to identify the environmental and biotic drivers of local adaptation.

Traits that increase fitness on one end of an environmental gradient can reduce fitness on the

opposite end of that gradient, resulting in fitness trade-offs (Kawecki & Ebert, 2004; Wadgymar

et al., 2022). Trade-offs are often directly caused by evolutionary changes in the allocation of

limited resources to critical biological functions, including growth, reproduction, and defense

(Bazzaz *et al.*, 1987; Herms & Mattson, 1992), but may be indirectly caused by interactions with

other species (Strauss *et al.*, 2002). Theory predicts that resource allocation to herbivore defense

should depend on the risk and consequences of herbivory on fitness, and models of the evolution

of plant defense assume a cost to the production of herbivore defenses (Rhoades, 1979; Stamp,

2003). Within species, herbivore resistance frequently trades-off with reproduction (Agren &

Schemske, 1993; Heil & Baldwin, 2002; Strauss *et al.*, 2002; Stowe & Marquis, 2011; Cipollini

et al., 2017), and increased herbivore resistance is associated with longer growing seasons

(Kooyers *et al.*, 2017).

Recent studies have shown that shifts in the allocation of resources from rapid growth to

herbivore resistance are made through a set of interacting gene networks that are involved in the

biosynthesis and downstream signaling of plant hormones (Kazan & Manners, 2012; Huot *et al.*,

2014; Campos *et al.*, 2016; Havko *et al.*, 2016; Aerts *et al.*, 2021; Monson *et al.*, 2022).

Jasmonic acid (JA) is a key regulatory hormone involved in the response of plants to herbivore

attack (Zhang & Turner, 2008; Havko *et al.*, 2016). While JA production increases herbivore

defense, it also can inhibit rapid plant growth through interactions with other gene networks

(Yan *et al.*, 2007; Zhang & Turner, 2008; Kazan & Manners, 2012; Yang *et al.*, 2012). Further,

the antagonistic interactions of genes in the gibberellin and jasmonic acid signalling pathways

are thought to contribute to the trade-off in resource allocation in plants between rapid

growth/reproduction (promoted by gibberellin) and defense (promoted by jasmonic acid) (Yang

et al., 2012; Hou *et al.*, 2013; Havko *et al.*, 2016). However, empirical evidence that

evolutionary changes in the GA pathway lead to changes in the relative allocation of resources to

reproduction, long-term growth, and herbivore resistance is lacking.

We used locally-adapted ecotypes of the yellow monkeyflower, *Mimulus guttatus* (syn.

Erythranthe guttata; (Barker *et al.*, 2012)) to investigate the mechanisms responsible for the

evolution of locally adaptive trade-offs in growth, reproduction, and resistance. In prior

reciprocal transplant experiments in this system, the primary environmental factor contributing to

local adaptation at inland sites was the onset of summer drought (Hall & Willis, 2006; Lowry *et*

al., 2008). In contrast, above-ground factors, potentially salt spray and/or herbivory, contribute to

adaptation in coastal habitats (Lowry *et al.*, 2009; Popovic & Lowry, 2020). Oceanic salt spray

limits plant growth, injures plant tissues, and contributes to coastal plant community zonation

through the differential tolerance of species to salt spray (Boyce, 1954). Inland populations of *M.*

guttatus are typically small annuals that flower in early spring to avoid the onset of summer

drought. Coastal populations, which occur in habitats with year-round soil moisture, are large

perennials that allocate resources primarily to long-term vegetative growth (Hall & Willis, 2006;

Lowry *et al.*, 2008; Hall *et al.*, 2010; Baker & Diggle, 2011; Baker *et al.*, 2012). Further, coastal

populations have higher levels of phytochemical defenses (phenylpropanoid glycosides, PPGs)

and are thought to experience higher levels of herbivory than inland annual populations (Holeski

et al., 2010, 2013; Lowry *et al.*, 2019). Although these patterns are suggestive, we lack direct

evidence that coastal populations are more resistant to herbivory than annuals when grown in the

field. In the greenhouse, coastal populations are more responsive to exogenous applications of

gibberellin (GA_3) than annuals and respond by recapitulating the elongated growth habit of

inland annual populations (Lowry *et al.*, 2019). As a result, we hypothesize that natural variation

in the timing of reproduction, and allocation to vegetative growth and resistance is the result of

molecular changes that alter the interactions of the gibberellin (GA) and jasmonic acid (JA)

pathways.

In this study, we performed a manipulative reciprocal transplant experiment with coastal

perennial and inland annual ecotypes of *M. guttatus* to test whether resource allocation trade-offs

contribute to local adaptation at opposite ends of an environmental gradient. We predicted that

GA treatment would cause earlier reproduction, that in turn would increase perennial fitness at

the inland site, where earlier flowering would rescue fitness for individuals that typically perish

before the onset of summer drought. We also predicted that JA and paclobutrazol (a GA

inhibitor) treatment would cause increased allocation to vegetative growth (via delayed

flowering) and herbivore resistance, that in turn would increase annual fitness at the coast, where

we expected herbivore pressure to be higher. Finally, we predicted that reduction of herbivory

via exclosures would rescue inland annual fitness on the coast, and thus decrease perennial home

site advantage. While our study was designed to focus on the role of hormone manipulation on

herbivore resistance, we instead discovered that our hormone manipulations had a much larger

role in causing susceptibility to stress imposed by oceanic salt spray. To assess the generality of

our observations in the manipulative reciprocal transplant experiment, we performed an

additional experiment focusing on the role GA plays in regulating plant height and salt spray

sensitivity across multiple populations spanning the coast to inland gradient.

Materials and Methods

Study locations and plants used in the reciprocal transplant experiment

In 2020, we performed a reciprocal transplant experiment at a coastal seep at the Bodega Marine Reserve in Bodega Bay, CA (Latitude: 38.3157, longitude: −123.0686), and an inland seep at the Pepperwood Preserve near Santa Rosa, CA (latitude: 38.5755, longitude: −122.7009). We used outbred maternal families from a coastal perennial population that experiences salt spray from Bodega Bay, CA ($n = 5$ families, BHW population coordinates: 38.303783, -123.064483) and an inland annual population north of Sonoma ($n = 4$ families, CAV population coordinates: 38.342817, -122.4854). Since we were primarily interested in ecotype differences, we generated outbred maternal families by intercrossing a small number of field-collected maternal families ($n = 6$ and $n = 4$, for BHW and CAV respectively) in the greenhouses at Michigan State University. Seeds from these outbred families were planted and placed in a 4°C cold room on January 27, 2020 at UC Berkeley. We staggered perennial and annual germination to synchronize their development, following (Popovic & Lowry, 2020). One and two weeks after initiating stratification, perennial and annual seeds were moved into a 16-hr day length growth chamber, respectively. Seedlings were transported from UC Berkeley to the Bodega Marine Reserve (BMR) greenhouse on February 20, 2020, where they experienced ambient light and temperature conditions. In the greenhouse, seedlings were transplanted into 288 cell plug trays filled with Sunshine Mix #4 Professional Growing Mix (Sun Gro Horticulture, Agawam, MA, USA) over a week (February 21- February 28, 2020).

Hormone treatments applied in the reciprocal transplant experiment

We altered allocation phenotypes of each ecotype by manipulating hormone levels of plants with

exogenous applications of gibberellin (GA_3), paclobutrazol (a gibberellin inhibitor), and methyl

jasmonate (MeJA, the bioactive form of Jasmonic Acid) to test the role of those hormone

pathways in adaptive trade-offs. Following a week of transplanting in the BMR greenhouse, on

February 28, 2020 we randomly assigned trays to one of three hormone treatments or a no-

hormone control. We sprayed plants with a 100 µM solution of GA_3 ($n = 252$ seedlings per

ecotype, Consolidated Chemical Solvents LLC, following Lowry *et al.*, 2019), 10 mM methyl

jasmonate ($n = 250$ seedlings per ecotype; TCI America, Portland, Oregon, USA), or 14.3 mg/L

solution of paclobutrazol ($n = 250$ seedlings per ecotype, General Hydroponics, Santa Rosa,

California, USA). Concentrations of methyl jasmonate and paclobutrazol were chosen after

conducting dose response experiments at the MSU greenhouses in winter 2019, based on the

minimum concentration needed to elicit a phenotypic change relative to no-hormone controls

without detrimental effects (e.g., leaf damage, stunting, death). Using a spray bottle, we sprayed

individual plants 5 times, corresponding to 3.5 mL of solution. The no-hormone control

consisted of spraying plants ($n = 248$ seedlings per ecotype) with 3.5mL of a 0.25% ethanol

solution since a dilute ethanol solution was needed to dissolve methyl jasmonate and all

hormones were dissolved in a 0.25% ethanol solution. Hormones were applied once on a single

day in the greenhouse prior to field transplanting. All trays were covered with clear plastic

domes for 24 hours (February 28 - February 29, 2020) and moved to different greenhouse

benches to prevent cross contamination.

Field planting of the reciprocal transplant experiment

Prior to transplanting, we removed vegetation from ten (1.08 x 0.84 m) plots at each site. We dug

trenches along the edge of each plot to bury the bottom of our open control and exclosure

structures. At the Pepperwood Preserve, we transplanted 800 seedlings on March 9, 2020, and

200 seedlings on March 12, 2020. At the Bodega Marine Reserve, we transplanted 800 seedlings

on March 10, 2020, and 200 seedlings on March 13, 2020. All seedlings had two expanded leaf

pairs at transplanting. Plants were fully randomized within each block (n = 100 seedlings/block).

Herbivore exclosures used in the reciprocal transplant experiment

To lessen the effect of herbivory, we deployed herbivory exclosures on four out of ten plots at

each site (Figure 1). A previous reciprocal transplant experiment at our study sites used

agrofabric exclosures that blocked all above-ground factors (Popovic & Lowry, 2020), and could

not separate the effects of salt spray and herbivory on plant fitness. Thus, we designed exclosures

that excluded many herbivores but allowed salt spray to pass through. The exclosures were 1.08

m long x 0.84 m deep x 0.87 m tall and constructed of 2.67 cm diameter PVC pipe covered with

fiberglass window-screen (1.4 x 1.6 mm mesh) that was affixed with fishing line and marine

epoxy. Each exclosure had screen doors along both long sides that were attached with velcro to

allow access to the plots. The screen extended 10 cm down into the soil around the plots. To

control for shading or moisture-collection due to the screen, the remaining six plots (open

structures) at each site were covered with window screen only on the top of each structure and 30

cm down each side.

Field Censuses of the reciprocal transplant experiment

After transplanting seedlings at the two-leaf stage, we performed regular censuses of our

transplant sites, recording survival, the presence of herbivore damage, the presence of salt-spray

damage, and the presence and number of reproductive structures (buds, flowers, and fruits). In

our census, we distinguished damage and death caused by salt spray from herbivory: salt-

damaged leaves appeared necrotic and brown and exhibited no sign of herbivore damage (i.e., no

missing tissue). When salt damage spread to the entire plant and no green tissue remained, we

considered plants to be killed by salt spray. When possible, we distinguished between sources of

herbivory. Most notably, we observed damage to plants consistent with mammalian damage at

the coastal site (removal of flowering stalks leaving plant stumps) that we could visually

distinguish from insect chewing damage and leaf miners. We were prevented from accessing our

transplant sites for two weeks at Bodega Marine Reserve and seven weeks at Pepperwood

Preserve after transplanting due to the 2020 COVID-19 pandemic lockdowns. Due to site

differences in growing season length, and restricted access due to the 2020 COVID-19

pandemic, we censused each site at different intervals and for different lengths of time

(Pepperwood Preserve (inland site): 12 censuses over 139 days, April 3 through July 29, 2020;

Bodega Marine Reserve (coastal site): 22 censuses over 194 days, March 24 through September

22, 2020). Since we were unable to access our sites for weeks because of the pandemic, we

missed observing the first flower opening for many annual plants at the inland transplant site. For

these plants, we estimated the onset of flowering as the date we first observed any reproductive

structures. Our censuses occurred on a roughly weekly basis after we were able to re-access our

sites.

Plant chemistry measured in the reciprocal transplant experiment

We sampled leaves for chemical analysis 55 to 57 (May 7-9, 2020) and 59 to 64 (May 11-16,

2020) days after transplanting at the coastal and inland site, respectively. To minimize the

potential effect of diurnal fluctuation in PPGs (phenylpropanoid glycosides), we sampled from

9am until 1pm, and to minimize the effect of leaf position, we sampled two leaves from the 3rd

node when possible, using leaves from the 4th and 5th nodes if leaves at the 3rd node were

damaged. After sampling, leaves were flash-frozen with liquid nitrogen and then freeze-dried.

For samples that did not meet the minimum dry mass (3mg), we either pooled them with other

low-mass samples (by grouping within all fixed and random factors as discussed below) or

excluded them. Our final sample size was 216 from Bodega (perennials only due to high annual

mortality) and 599 from Pepperwood. To determine the PPG concentrations in sampled leaves,

we ground, extracted, and prepped extract aliquots as described in Holeski *et al.* (2013). We then

used high-performance liquid chromatography (HPLC) to quantify PPGs. The HPLC method is

described in (Kooyers *et al.*, 2017) and was run on an Agilent 1260 HPLC with a diode array

detector and Poroshell 120 EC-C18 analytical column [4.6□×□250□mm, 2.7 μm particle size];

Agilent Technologies). We calculated concentrations of individual PPGs as verbascoside

equivalents, using a standard verbascoside solution (Santa Cruz Biotechnology, Dallas, Texas),

as described in (Holeski *et al.*, 2013, 2014).

GA$_3$-salt spray exposure experiment (2023)

Since we observed a large effect of GA$_3$ treatment on plant height (although we did not measure

it), necrosis and mortality at our coastal transplant site in 2020, we wanted to confirm the

combined impact of GA$_3$ and salt spray and assess how common this effect was across multiple

populations. In 2023, we grew one outbred accession from four coastal perennial (HEC, OPB,

SWB, BMR) and four inland annual (LMC, PPW, CAV, MOR), one inland perennial (SAL), one

near-coastal annual (AJEN, 1.5km from the coast), and two near-coastal perennial populations

(JEN and GUL, 3 and 4.7 km from the coast respectively) in a greenhouse in the Bodega Marine

Reserve (Table S17). Near-coastal populations experience the same maritime climate as coastal

populations but are not directly exposed to salt spray. Seeds for perennials were planted a week

earlier than annuals to synchronize germination (March 2 vs March 9, 2023). Seedlings were

transplanted into individual 8.9 cm square pots (T.O. Plastics, Clearwater, MN, USA) as they

germinated and randomly assigned to a GA_3 treatment or no-hormone control. We used the exact

same protocol as in 2020 for the GA_3 treatment and no-hormone control, except that we did not

add ethanol to the spray bottle which was used as a solvent for methyl jasmonate. A total of 22

and 23 seedlings per population were sprayed with GA_3 and water, respectively, on March 24,

2023. GA_3-treated and no-hormone control plants were sprayed on separate greenhouse benches

and covered with clear plastic domes for 24 hours to prevent cross contamination. Two weeks

after the GA_3 or water treatments, we randomly assigned seedlings into blocks (six 1020 trays

[T.O. Plastics, Clearwater, MN, USA] per block). We measured height from the base of each

plant to the apical meristem using a ruler in the greenhouse on April 19, 2023 and then

immediately placed plants ($n = 540$) outdoors on a coastal bluff for one week (latitude: 38.31841,

longitude: -123.07329; Figure S1). On the bluff, we either protected the plants from salt spray

using two agrofabric covered exclosures ($n = 218$ plants) or exposed them to salt spray using

three open structures ($n = 324$ plants) using the same structure dimensions and agrofabric as

(Popovic & Lowry, 2020). The two agrofabric exclosures contained ten GA_3-treated and eight

water-sprayed plants per population, and the three open structures contained 12 GA_3-treated and

15 water-sprayed plants per population. After one week of exposure, we transported plants back

to the greenhouse and assessed levels of leaf damage on each leaf pair by eye. Standard methods

for estimating whole plant damage typically break up whole plants into units (Herbivory

Variability Network*† *et al.*, 2023). Leaf pair was chosen as the unit to visually estimate damage

because monkeyflowers produce alternate leaves at each node. Whole plant damage was

estimated by averaging visual estimates of leaf damage for each leaf pair within each plant.

Because plants were maintained in pots during the entirety of the 2023 experiment, we could be

assured that damage was caused by oceanic salt spray applied to experimental plants by wind

coming off of the Pacific Ocean. We did not observe any herbivore damage to plants exposed or

protected from the ocean in 2023.

Statistical analyses

The overarching goal of our statistical analyses was to understand how ecotype, hormone

treatment, field site, and herbivore exclosure treatment contribute to within plant resource

allocation and fitness. To establish how plants with different ecotype and treatment combinations

allocate resources to reproduction, we compared flowering time among plants. To compare

allocation to herbivore resistance, we compared the presence or absence of herbivore damage to

each plant and the concentration and composition of the defensive compounds in leaves. To

compare fitness, we quantified survival across the season, the presence of flowers, and seasonal

fruit production. For annuals, these measures indicate lifetime fitness, whereas perennials that

survived the season have the potential to reproduce in subsequent years. We performed all

statistical analysis in R version 4.3.1 (R core team 2023).

Univariate analysis

To test whether ecotypes differed in resource allocation and whether hormone treatments altered

these differences at each transplant site, we fit mixed models for each analysis that included

ecotype, hormone treatment, and exclosure type as interactive fixed factors. The response

variables were flowering time, the presence or absence of herbivory, survival, total PPG

concentration (summed concentration for all PPG compounds), whether an individual produced a

reproductive structure (buds, flowers, or fruit), or the number of fruits produced by plants that

flowered at the end of the season. When possible we categorized herbivore damage by the

putative source, based on patterns of plant tissue damage. Methods and results of analyses by

herbivore type are detailed in the supplement. All models also included two random effects for

maternal family and experimental plot. We fit all mixed models except for the survival model

with the R package *glmmTMB* (Brooks *et al.*, 2017), and modeled survival using the R package

coxme (Therneau, 2022). We identified the best fitting error distributions by evaluating model

diagnostics with the R package *DHARMa* (Hartig, 2022). We fit mixed models for flowering

time with Gaussian error distributions, mixed models for herbivory and flowering probability

with binomial error distributions, and mixed models for log-transformed total PPGs with gamma

distributions. We modeled survival using a mixed-effects Cox Proportional Hazards model, and

modeled fruit number with a zero-inflated negative binomial error distribution at the coastal site

and a negative binomial error distribution at the inland site.

To prevent model overfitting, we used an analysis of deviance (Wald χ^2 test) to assess the

significance of model terms and sequentially removed unsupported model terms (R package *car*,

(Fox & Weisberg, 2018). We compared fits of complex versus reduced models using likelihood

ratio tests (LRT) to find the minimum adequate model for each response variable in each site

(Tables S1 and S2). We compared treatment groups using post-hoc tests on the minimum

adequate model with the R package *emmeans* (Lenth *et al.*, 2020). For one of the binary response

variables (flowering probability), there was no variance in one of the treatment levels (quasi-

complete separation) and a binomial model was not able to accurately estimate the parameter for

that treatment group. For this reason, we used a randomization test to estimate the probability

that this result was due to chance, by reassigning treatments to all outcomes 10,000 times and

calculating the proportion of these reshuffles that resulted in a difference as extreme as the

observed (Gotelli & Ellison, 2004). We predicted the mean and 95% confidence intervals for

each response variable from our models using the R package *ggeffects* (Lüdecke, 2018). For all

non-binary response variables, we predicted confidence intervals via bootstrapping (n=500

iterations). We plotted raw data and predictions in *ggplot2* (Wickham, 2016) and combined plots

with *patchwork* in R (Pedersen, 2019).

Multivariate PPG analysis

Within each transplant site, we modeled the concentration of all nine different PPGs (the PPG

arsenal) using mixed models for each analysis that included ecotype (at Pepperwood only),

hormone treatment, and exclosure type as interactive fixed factors and block as a random factor.

We fit all models with PERMANOVA with Bray-Curtis distance using the adonis2 function

from the R package *vegan* (Oksanen, 2016). We dropped all non-significant factors for the

minimum adequate model (Table S3). To test for homogeneity of variance among treatment

groups, which can influence inference, we used the betadisper function from the *vegan* package.

The only factor that had heterogeneity of variance among levels was ecotype. Due to the strength

of the signal for ecotype, and confirmation from other studies that annuals and perennials have

different PPG arsenals (Holeski *et al.*, 2013), we are confident that differences due to ecotype are

not attributable only to heterogeneity of variance. We compared treatment groups using post-hoc

tests on the minimum adequate model with the function pairwise.adonis2 from the R package

pairwiseAdonis (Arbizu, 2019). To visualize how multivariate PPG arsenals are influenced by

our factors, we used non-metric multidimensional scaling (NMDS) (*MetaMDS* function in *vegan*

package with Bray-Curtis distance to determine dissimilarity) and added standard-error ellipses

at 95% confidence around the centroid of each cluster (function *ordiellipse* from package vegan).

Analysis of GA_3-salt spray exposure experiment (2023)

To analyze the effect of GA_3 treatment on plant height in the greenhouse, we fit a linear mixed

model with height as the response variable and population ($n = 12$), GA_3 treatment ($n = 2$), and

their interaction as the predictor variables and flat ($n = 30$) as a random effect. Since only 10%

(21/216) of seedlings in the exclosures showed any evidence of necrosis after one week of being

outdoors, compared to 78% (254/324) of seedlings in the open structures, we only analyzed the

effect of hormone treatment and population on plants exposed to salt spray in the three open

structures. We first logit transformed proportional leaf necrosis (Warton & Hui, 2011), and then

fit a linear mixed model with logit-transformed leaf necrosis as the response variable, population,

hormone treatment, and their interaction as fixed effects, and flat ($n = 18$) nested within block (n

$= 3$) as random effects. Within each population, we performed pairwise comparisons of no-

hormone control vs GA_3-treated plants (function *pairs* from package emmeans). We used a

multivariate *t*-adjustment for multiple comparisons ($n = 12$ comparisons). We plotted predicted

leaf necrosis by transforming logit-transformed proportions back to proportions.

Results

Annuals flowered earlier, investing less time in vegetative growth than perennials, and hormone treatments slightly delayed flowering time

At the coastal site, flowering time was significantly associated with ecotype, hormone treatment, and an ecotype by hormone treatment interaction (Table S2: all $p \leq 0.03$). At the inland site, flowering time was significantly associated with ecotype, hormone treatment, an ecotype by hormone treatment interaction, and a three-way ecotype by hormone treatment by exclosure treatment interaction (Table S2: all $p \leq 0.01$). Annuals flowered 37-62 and 47-63 days earlier than perennials at the inland and coastal sites, respectively (within each hormone and exclosure treatment), respectively (Tables S4 & S5: all annual vs perennial comparison p-values < 0.001, Figure 2 A&B). Annuals treated with GA_3 flowered 9 and 16 days later than annuals not treated with hormones at the inland site in open structures and at the coastal site, respectively (Tables S4 and S5, coastal site: GA_3 vs control annuals $p = 0.043$; inland site: GA_3 vs no-hormone control annuals in open structures $p < 0.009$, in exclosures $p = 1$). Annuals treated with paclobutrazol at the coastal site flowered 6 days later than annuals not treated with hormones (Tables S4 and S5, coastal site: paclobutrazol vs no-hormone control annuals $p = 0.027$). Mesh exclosure treatment did not influence flowering time (Table S1: $p = 0.66$; Table S5: all comparisons between exclosure treatments within ecotype and hormone treatment at the inland site $p \geq 0.463$). Detailed results can be found in the supplemental materials.

Perennials were more likely to be attacked by herbivores at both sites and GA$_3$ reduced the probability of herbivory at the coastal site

At the coastal site, the probability of herbivory was significantly associated with ecotype, mesh

exclosure treatment, an ecotype by exclosure treatment interaction, and a hormone treatment by

exclosure treatment interaction (all $p < 0.001$, Table S2). At the inland site, the probability of

herbivory was significantly associated with an ecotype by hormone treatment interaction, and an

ecotype by exclosure treatment interaction (all $p \leq 0.03$, Table S2). Contrary to expectations,

perennials were 21-287 and 6-10 times more likely to experience herbivory than annuals at the

coastal site in open structures and exclosures, respectively (all annual vs perennial comparisons

within hormone and exclosure treatment $p < 0.001$, Table S6, Figure 2 C). Perennials were 6

times more likely to experience herbivory than annuals in open structures when treated with

MeJA or no hormone at the inland site (no-hormone and MeJA-treated annual vs perennial

comparisons in open structures $p < 0.001$, all other within hormone and exclosure treatment

contrasts $p \geq 0.28$; Table S7, Figure 2D). However, this result is somewhat misleading at the

coastal site, since annuals perished quickly and had less time to encounter herbivores and accrue

herbivory. GA_3 was the only hormone to influence the probability of herbivory, and only at the

coastal site, where, again contrary to expectations, it reduced the probability of herbivory for

both ecotypes. Annuals and perennials were 77-97% and 59-73% less likely to experience

herbivory when treated with GA_3 relative to no-hormone controls (all no-hormone control vs

GA_3-treatment contrasts within each ecotype and exclosure treatment $p < 0.001$, Table S6, Figure

2C). This is potentially due to an interaction with salt-spray resistance rather than allocation to

herbivore resistance.

The mesh-size of screen used in our exclosures, while necessary to allow salt spray to enter, did

allow some small insects, including leaf miners and weevils, to enter the exclosures (or they

were present when the exclosures were erected) and damage plants mildly. As a result, our

herbivore exclosures did not significantly reduce the probability of herbivory for perennials at

either site (Table S6 & S7: all open vs exclosure structure contrasts within each hormone

treatment for perennials $p \geq 0.06$, Figure 2C). However, we observed the removal of flowering

stalks from perennial transplants consistent with mammalian damage (likely deer, based on the

nature of the cuts, hoof prints, and observations of the resident deer population, per obs.) only in

open structures at the coastal site. Exclosures significantly reduced deer herbivory by 100%, but

did not significantly reduce other sources of herbivory for coastal perennials at the coastal site

(detailed in the supplemental materials, Figure S2). At the coastal site, GA_3-treated annuals were

29% more likely to experience herbivory in exclosures compared to open structures, although

this is likely due to an interaction with salt-spray (Table S6: $p = 0.02$, all other open vs exclosure

structure contrasts within hormone treatments for annuals $p = 1$, Figure 2C). At the inland site,

hormone-treated annuals and perennials did not significantly differ from their respective no-

hormone controls and exclosure treatment did not influence the probability of herbivory for

either ecotype (Table S7: all open vs exclosure structure contrasts within ecotype and hormone

treatment $p \geq 0.96$, Figure 3D).

Perennials produced more PPGs, and methyl jasmonate and GA_3

increased PPGs at the inland site while GA_3 decreased PPGs at the

coastal site

At the coastal site, total PPG concentrations were significantly associated with hormone

treatment, and a hormone treatment by mesh exclosure treatment interaction (Table S2: all $p \leq$

0.008). At the inland site, total PPG concentrations were significantly associated with ecotype,

hormone treatment, and an ecotype by hormone treatment interaction (Table S2: all $p \leq 0.01$).

Perennials produced 75-87 mg/g more PPGs than annuals at the inland site (Table S9: all annual

vs perennial comparisons within hormone treatments $p < 0.001$, Figure 3B), while annual

mortality at the coast prevented this comparison (Figure 3A). Exclosure treatment did not

significantly influence total PPG concentrations at either transplant site (Table S8: all open vs

exclosure structure comparisons within hormone treatment at the coastal transplant site $p \geq 0.76$,

Table S2: $p = 0.67$). At the coastal site in the open structures, perennials treated with GA$_3$

produced 43 mg/g fewer PPGs than perennials not treated with hormones (Table S8: no-hormone

control vs GA$_3$ in open structures: $p < 0.001$, in exclosures: $p = 0.76$). At the inland site, annuals

and perennials treated with GA$_3$ produced 14 mg/g and 5 mg/g more PPGs than annuals and

perennials not treated with hormones, respectively (Table S9: no-hormone control vs GA$_3$-

treatment comparisons for each ecotype: $p < 0.001$). Annuals treated with methyl jasmonate

produced 17 mg/g more PPGs than annuals not treated with hormones at the inland field site

(Table S9: no-hormone control vs MeJA-treatment contrast $p = 0.001$ for annuals, $p = 0.99$ for

perennials).

At the coastal site, multivariate PPG arsenals were significantly associated with hormone

treatment; GA$_3$-treated perennials differed from all other perennials (Table S3: all $p \leq 0.02$,

Figure 3C). At the inland site, multivariate PPG arsenals were significantly associated with

ecotype, hormone treatment, mesh exclosure treatment, and an ecotype by hormone interaction

(Table S3: all $p < 0.003$, Figure 3D). Detailed results can be found in the supplemental materials.

Perennials survived longer at both sites, and GA₃ reduced survival at the coastal site

At the coastal transplant site, survival was significantly associated with ecotype, hormone treatment, mesh exclosure treatment, an ecotype by hormone treatment interaction, and a three-way ecotype by hormone treatment by exclosure treatment interaction (Table S2: all $p \leq 0.04$). At the inland transplant site, survival was only significantly associated with ecotype (Table S2: $p < 0.001$). At both transplant sites, perennials survived significantly longer than annuals (Tables S12 & S13: all annual vs perennial comparisons [within hormone and exclosure treatment at the coastal site] $p < 0.001$, Figure 3A&B). At the inland site, the median death date for annuals (97 days from transplanting) was 21 days earlier than perennials (118 days). At the coastal site, the median death date for annuals was 46 days, but since fewer than 50% of perennials before the end of our experiment, we could not estimate their median death date. At the coastal site, we observed tissue necrosis prior to the death of all annuals and 93% (109/117) of the perennials that died. For the remaining 8 perennial plants, we did not observe tissue necrosis but did observe herbivory prior to death. The majority of plants at the inland site died as the site began to dry out in summer.

At the coastal site, GA₃ treatment significantly increased the risk of death for both ecotypes relative to their respective no-hormone controls (GA₃:no-hormone cumulative hazard ratios - annuals: 2.65 [open structure], 3.34 [exclosure], perennials: 28.6 [open structure], 9.87 [exclosure]; Table S12: no-hormone control vs GA₃-treatment comparisons for both ecotypes, $p < 0.001$). GA₃-treated plants senesced and died faster in the open structures at the coastal site; the median death date in the open structures was 53 days compared to 96 days in the exclosures

(Figure 4A). We observed tissue necrosis on GA_3-treated plants prior to death at the coastal site.

GA_3-treated perennials were more upright compared to prostrate no-hormone controls, and

elongated their stems early in development like annuals which may have increased exposure to

salt spray (Figure 5). Mesh herbivore exclosures decreased the risk of death in GA_3-treated

perennials at the coastal site (open structure:exclosure cumulative hazard ratio: 2.53, Table S12:

open vs exclosure structure comparison $p < 0.001$, Figure 4A). Coastal fog sometimes condensed

on the screens that we used to exclude herbivores, which may have slightly decreased the

transmission of salt spray into exclosures.

Annuals were more likely to flower at the inland site, and GA_3 reduced flowering probability at the coastal site

At the coastal site, the probability of flowering was significantly associated with hormone

treatment, mesh exclosure treatment, and an ecotype by exclosure treatment interaction (Table

S2: all $p \le 0.007$). Despite high early season annual mortality at the coastal site, annuals and

perennials did not significantly differ in the probability of flowering (Table S14: all annual vs

perennial comparisons within exclosure treatment $p \ge 0.18$, Figure 6A). Due to their rapid

phenology, 40% (198/500) of annual transplants flowered prior to dying, although death

occurred quickly after flowering so very few annuals produced fruit (detailed below). Perennials

were 3-11 more likely to flower in herbivore exclosures than open structures (Table S14: $p <$

0.014 for open vs exclosure structure contrasts, Figure 6A), which may be due to the reduction of

large mammalian herbivory and/or minor buffering of salt spray from condensation collecting on

mesh screens. In addition, transplants that were not treated with hormones were more likely to

flower than transplants treated with GA_3 (8x in open structures, 3-4x in exclosures), potentially

due to the effect of GA_3 on sensitivity to salt spray (Table S14: all no-hormone control vs GA_3-

treatment contrasts $p < 0.001$, Figure 6A).

At the inland transplant site, the probability of flowering was significantly associated with

ecotype and an ecotype by hormone treatment interaction (Table S2: both $p \leq 0.01$). At the

inland site, annuals were 65-305% and 34-90% more likely to flower than perennials in open

structures and exclosures, respectively (Table S15: all annual vs perennial comparisons within

each hormone treatment in open structures $p < 0.001$, in exclosures $p = 1$, Figure 6B). Since all

annuals flowered in the exclosures, the binomial model could not accurately estimate flowering

probabilities for annuals in exclosures. Therefore, we used a randomization test to estimate the

probability that the difference between annuals and perennials in exclosures was due to chance.

Randomization tests showed that 0 (all hormone treatments) or 15 (the no-hormone controls) of

10,000 treatment reshuffles resulted in the observed difference between annual and perennial

flowering probabilities in the exclosures, making it unlikely that this pattern arose by chance (no-

hormone controls: $p = 0.0015$, GA-, methyl jasmonate- and paclobutrazol- treated plants $p =$

0.0). Perennials treated with GA_3 and MeJA were 30-51% and 36-59% less likely to flower than

perennials not treated with hormones, but hormone treatment did not affect the probability of

flowering in annuals (Table S15: $p \leq 0.002$ for no-hormone control vs GA_3 and no-hormone

control vs MeJA-treatment contrasts in perennials within each exclosure treatment, Figure 6B).

GA_3 reduced fruit production at the inland site, and herbivore exclosures

drastically increased fruit production at the coastal site

At the inland site, fruit number among plants that flowered was significantly associated with

hormone treatment and marginally associated with mesh exclosure treatment (Table S2: $p <$

0.001 and $p = 0.06$). Annuals and perennials that flowered did not significantly differ in fruit

production in either open structures or exclosures (Table S16: all annual vs perennial

comparisons $p = 1$, Figure 6D). Each ecotype produced 2-4 fewer fruits when treated with GA_3

relative to no-hormone controls (Table S16: all no-hormone control vs GA_3-treatment

comparisons $p < 0.001$, Figure 6).

At the coastal site, only plants protected by mesh exclosures successfully produced fruit by the

end of the season (Figure 6C) because of complete inflorescence herbivory by mule deer

(*Odocoileus hemionus*). Since no plants produced fruit outside of the exclosures, and few

annuals produced fruit inside the exclosures, we analyzed only the effect of hormone treatments

on perennial fruit production inside the exclosures at the coastal site (Figure 6C). Fruit

production in perennial plants that flowered in exclosures at the coastal site was not significantly

associated with hormone treatment (Table S1: $p = 0.48$).

GA_3 increased plant height and salt spray susceptibility

In the subsequent experiment in 2023, greenhouse plant height was significantly associated with

population (Analysis of deviance: Wald Type III $\chi^2 = 628.78$, $df = 11$, $p < 0.001$), GA_3 treatment

($\chi^2 = 1318.39$, $df = 1$, $p < 0.001$), and a population by GA_3 treatment interaction ($\chi^2 = 362.10$, df

$= 11$, $p < 0.001$). In pairwise comparisons, GA_3 treated plants from 12 out of 13 populations were

significantly taller than their respective no-hormone controls (Figure 7; Table S18). Three inland

annual populations (LMC, CAV, and MOR), the inland perennial population (SAL), and one

near-coastal perennial population (JEN) were the most responsive to GA_3 treatment (effect on

height > 19 mm vs < 11 mm in the remaining populations; Figure 7).

Agrofabric exclosures effectively protected seedlings from salt spray damage: 10% (21/216) of

seedlings had tissue damage after a week in the exclosures compared to 78% (254/324) of

seedlings in the open structures. In ocean-exposed open structures, leaf damage was significantly

associated with population (Analysis of deviance: Wald Type III $\chi^2 = 37.40$, $df = 11$, $p < 0.001$)

and a population by GA_3 treatment interaction ($\chi^2 = 28.41$, $df = 11$, $p = 0.003$), but was not

significantly associated with hormone treatment ($\chi^2 = 2.34$, $df = 1$, $p = 0.13$). GA_3 treatment

significantly increased leaf damage by 224-952% only in four of the five accessions that

increased most in height in response to GA_3 (Table S19: $p \leq 0.03$, Figure 7, Figure S3).

Discussion

Across environmental gradients, shifts in allocation between reproduction, growth, and defense

follow predictable patterns, suggesting that these shifts underlie local adaptation (Bazzaz *et al.*,

1987; Hahn & Maron, 2016; Züst & Agrawal, 2017). However, multiple abiotic and biotic

factors co-vary across environmental gradients and multiple traits often differ between locally

adapted populations, making the identification of selective agents and their phenotypic targets a

major challenge (Wadgymar *et al.*, 2017, 2022; López-Goldar & Agrawal, 2021). In this study,

we used a manipulative reciprocal transplant experiment to test the hypothesis that herbivory and

divergence in reproductive timing, vegetative growth, and defense against herbivores contributes

to local adaptation across a coastal to inland environmental gradient.

At our coastal site, herbivore exclosures dramatically increased fecundity of local coastal

perennials. However, contrary to our predictions, herbivory did not contribute to local

adaptation, since excluding herbivores did not increase inland annual fitness at the coastal site.

This is likely due to the abiotic effect of salt-spray pre-empting the impacts of herbivory on

annuals. Our hormone treatments slightly shifted allocation between vegetative growth,

reproduction and defense in each ecotype, but did not recapitulate the full effect size of

differences previously observed in controlled greenhouse conditions (Lowry *et al.*, 2019).

Nevertheless, we observed dramatic effects of our hormone treatments on survival and fecundity

across our transplant sites. Despite delaying flowering, the GA_3 application caused obvious

earlier bolting and taller heights in the perennial transplants. A follow-up experiment confirmed

that GA_3 application both increased plant height and susceptibility to oceanic salt spray. Overall,

our results suggest that oceanic salt spray is the dominant environmental factor driving local

adaptation near the ocean and that coastal perennials significantly lose much of their locally

adaptive salt spray tolerance advantage over inland annuals in response to GA_3 application.

Role of biotic interactions in local adaptation

The organisms a plant interacts with vary across the landscape, causing different selective

pressures (Thompson, 2005; Urban, 2011; Friberg *et al.*, 2019). Given the differences in the

abiotic environment at our two sites–cool and foggy on the coast, hot and dry inland–the

communities of organisms which our plants interact with differ substantially. The moist coastal

environment has far more molluscan herbivores (snails, slugs), and a rare leaf-mining fly

(Eiseman *et al.*, 2023), which we did not see at the much drier inland site. Voles and deer also

only contributed to herbivory at the coastal site. Given the differences in communities,

differences in defense-levels, and prior research suggesting differences in intensity of herbivory

on the coast and inland (Holeski, 2007), we predicted herbivory would be a driving factor in

local adaptation. Surprisingly, we found no effects of herbivory on local adaptation at these two

specific sites. At both sites, coastal perennials were more likely to be damaged by herbivores.

Although exclosures reduced deer herbivory at the coastal site, this reduction only benefited the

local perennial ecotype since almost all annual transplants died of salt spray. While we did not

find an effect of herbivory on local adaptation, we stress that insect-plant interactions regularly

occur in a complex mosaic across the landscape and greatly vary temporally across years (Rotter

et al., 2022).

High rate of herbivory at the coastal site did not contribute to local adaptation

At the coastal site, we predicted that high rates of herbivore attack would result in herbivory

playing a strong role in local adaptation. A previous reciprocal transplant experiment found

dramatic fitness differences between inland annuals grown in agrofabric exclosures that blocked

all above ground factors, including herbivores and oceanic salt spray, and open structures at a

coastal site (Popovic & Lowry, 2020). While we observed high rates of herbivory at the coastal

field site, reducing deer herbivory with exclosures did not increase the success of inland annual

plants because salt spray could get through the exclosures and had such a deleterious effect.

Annuals transplanted at the coastal site quickly exhibited necrosis from salt spray before dying;

the window that they could have received herbivory was short, and they were likely poor quality

host plants during that time. The perennials, in comparison, were larger, healthier, and had many

more days in which to encounter an herbivore and receive damage. All reproductive herbivory at

the coastal site (which had a very strong effect on fecundity) came after the median death date of

our annual plants and thus was only experienced by perennials. These observations highlight the

importance of the timing of selective events, particularly for local adaptation of ecotypes with

differing life-history strategies. The importance of fecundity versus survival are likely to differ

between ecotypes (DeMarche *et al.*, 2016), and early-season factors (like coastal salt spray) that

impact survival might disproportionately contribute to fitness differences between populations

relative to late season factors that affect fecundity (such as herbivory; (Crone, 2001; Wadgymar

et al., 2022).

While this study suggests that herbivory is preempted from playing a role in keeping annuals out

of coastal environments, it does not mean it is unimportant. In the open structures on the coast,

perennials completely failed to reproduce due to deer herbivory of inflorescences. By virtue of

allocating growth to clonal expansion and non-reproductive tissue, perennials are likely

increasing both tolerance (Stevens *et al.*, 2008) and reducing the impacts of high-herbivory years

on their lifetime fitness. Some populations are completely sterilized (i.e., all inflorescences are

completely consumed by herbivores) in certain years (Toll, pers. obs.) and thus, herbivory may

be an extremely strong selective pressure in the morphology, allocation to clonal growth, and

reproductive timing of these coastal perennials. The results of this study were also clearly

influenced by the close proximity of our coastal field site to the open ocean (within 50 meters of

the shoreline). While it is common for coastal perennials to grow in close proximity to the ocean,

where they are impacted by high levels of salt spray, it is also common for them to grow slightly

further inland, where salt spray is greatly reduced (Boyce, 1954; Barbour, 1978; Du & Hesp,

2020). Coastal perennial populations of *M. guttatus* growing in close proximity to the ocean are

generally shorter than coastal perennial populations that are further from the ocean, and

presumably more protected from oceanic salt spray (Zambiasi & Lowry, 2024).

Life-history contributed to differences in herbivore attack at the inland site

Our finding that herbivory did not influence local adaptation inland is somewhat less surprising,

as there is evidence that herbivore damage is generally less extensive there (Holeski, 2007). It

was unexpected, however, that perennials were also more likely to be attacked by herbivores

than annuals at the inland site, as we predicted that perennials would be more resistant to

herbivory due to ecotype differences in phytochemical defenses (PPGs). Although early

flowering can be a highly effective herbivory avoidance strategy (Krimmel & Pearse, 2016),

perennials were more likely to be attacked even during periods of time when both ecotypes were

alive in the same site suggesting the difference in attack rates was not due to a shorter lifespan

for annuals. Higher attack rates for perennials could be due to differences in apparency caused by

differences in plant size (Feeny, 1976), or herbivore preference due to nutritional differences or

water content. In addition, while PPGs are feeding deterrents to generalists, some can be feeding

stimulants for specialist herbivores (Holeski *et al.*, 2013; Rotter *et al.*, 2018), and thus perennials

may be more likely to get attacked by specialists. Our presence-absence measure of herbivory

also may have missed differences in degree of herbivory among plants that were attacked, which

may have greater impacts on fitness.

Oceanic salt spray sensitivity increased with gibberellin treatment

The most surprising result of our experiment was how dramatically GA_3 application decreased

survival of coastal perennial genotypes at the coastal field site in the 2020 reciprocal transplant

experiment. Based on the patterns of plant tissue death, we attributed this mortality primarily to

oceanic salt spray. Our 2023 experiment confirmed that GA_3 addition made perennial and annual

genotypes more susceptible to natural oceanic salt spray.

There are two non-mutually exclusive ways that GA_3 could have decreased fitness in the coastal

environment with regard to salt spray. First, the addition of GA_3 increased plant height, as

evident by increased internode elongation of plants here as well as in a prior study (Lowry *et al.*,

2019). Increased plant height could put the aboveground portions of these plants more directly in

the path of prevailing wind delivering the salt spray (Zambiasi & Lowry, 2024). In 2023, all

seedlings treated with GA_3 responded by increasing height, but populations that grew tallest in

response to GA_3 had the most tissue damage after a week of ocean exposure, consistent with

increased height directly increasing plant exposure to salt spray.

A second hypothesis is that the GA_3 treatment may directly increase the susceptibility of tissues

to salt spray, independent of changes in plant height. The second hypothesis is particularly

intriguing, as it is the opposite of what would be expected based on the soil salinity literature.

For example, previous experiments in rice (Rodríguez *et al.*, 2006), wheat (Iqbal & Ashraf,

2013), apple (Wang *et al.*, 2019), cucumber (Wang *et al.*, 2020), and sorghum (Liu *et al.*, 2023)

have all found that the application of GA_3 increases yields under saline conditions. The

conflicting results of those studies and our experiment make it clear that findings from the soil

salinity literature cannot be directly extrapolated to what is experienced by plants growing in

coastal environments, where salt spray is a major source of stress on plant aboveground tissues

(Boyce, 1954; Du & Hesp, 2020; Itoh *et al.*, 2024). The exact mechanisms by which GA_3

increases salt spray susceptibility are still not clear, but are an active focus of our current

research.

While our study made the surprising discovery that GA_3 addition makes plants more susceptible

to salt spray, it is unclear the extent to which the adaptive changes in hormone pathways

contributed to the evolution of the more salt spray-tolerant coastal perennial ecotype. Other

recent studies have found evidence that evolutionary changes in both the gibberellin and

jasmonic acid pathways contribute to the divergence between coastal perennial and inland annual

ecotypes of *M. guttatus* (Olsen *et al.*, 2024; Kollar *et al.*, 2025). However, these evolutionary

genetic changes in hormone pathways have yet to be directly linked to salt spray tolerance in this

system or in other coastal ecotypes where hormone shifts have been implicated in their evolution

(Wilkinson *et al.*, 2021; James *et al.*, 2023; Broad *et al.*, 2024).

688   Limitations and future directions

Although our study provided clear evidence that salt spray is a major abiotic factor contributing

to local adaptation to coastal environments, our experiments had several limitations. First, we

could not monitor transplants for the first few weeks of the COVID-19 lockdown. Future studies

should monitor transplant fitness more frequently to capture early mortality. Second, we applied

only one dose of each hormone and observed minimal trait responses. Applying multiple

hormone doses in the field would allow researchers to test whether hormone effects on traits and

fitness are dosage dependent. Third, mesh exclosures were chosen to reduce herbivory while also

not excluding salt spray. The use of insecticides or molluscicides might further reduce different

sources of herbivory, though those compounds can have unintended impacts on plants, such as

through alterations of soil microbial communities. Our understanding of herbivory in this study

was also limited by measuring damage as presence or absence. Attempts to quantify damage

levels could be used to more precisely explore the role of herbivory in this system. Finally, we

were unable to examine the role of herbivory in the absence of salt spray. Future studies should

test the effect of herbivory on local adaptation in sites not directly exposed to salt spray.

703   Conclusions

The evolution of local adaptation is inherently complex, involving multiple environmental

factors that differ in importance among habitats and the evolution of multiple molecular and

biochemical pathways. The results of our study illustrate the complexity of local adaptation and

how studies to identify its underlying environmental, physiological, and genetic basis can lead to

surprising yet important findings. The initial goal of our manipulative reciprocal transplant

experiment was to investigate how hormone manipulation and herbivore exclosures affect the

fitness of locally adaptive ecotypes in both local and foreign habitats. To our surprise, hormone

manipulations ultimately had a minor impact on defensive compound production and herbivory.

Instead, we accidentally discovered that the manipulation of gibberellin concentrations, through

the addition of GA_3, had a profound impact on the fitness of coastal plants in their home habitat.

While it is well-established that interactions among hormone pathways mediate differences in

allocation to growth, reproduction, and resistance in plants (Murphy, 2015; VanWallendael *et*

al., 2019; Gasperini & Howe, 2024), few studies have investigated how evolutionary changes in

hormone pathways contribute to the formation of plant ecotypes and local adaptation (Wilkinson

et al., 2021; James *et al.*, 2023; Broad *et al.*, 2024), especially in the field. Future studies of local

adaptation should aim to more explicitly understand how shifts in hormone pathways contribute

to local adaptation and how those changes may have a broad and unexpected range of impacts on

adaptations to various environmental factors. Critically, functional molecular studies should be

conducted to test the hypothesis that adaptive genetic changes in hormone pathways underlie the

evolution of locally adaptive shifts in resource allocation. While transgenic plants produced by

those studies can readily be evaluated in the laboratory, government regulations discourage

evaluation of those plants in natural habitats. The application of synthetic hormones and their

antagonists, as conducted here, offer a much more acceptable way to test hypotheses about the

roles of hormones in local adaptation in the field.

Literature Cited

Aerts, N., Pereira Mendes, M. & Van Wees, S.C.M. 2021. Multiple levels of crosstalk in hormone networks regulating plant defense. *Plant J.* **105**: 489–504.

Agren, J. & Schemske, D.W. 1993. The cost of defense against herbivores: an experimental study of trichome production in Brassica rapa. *Am. Nat.* **141**: 338–350.

Arbizu, P.M. 2019. pairwiseAdonis: pairwise multilevel comparison using adonis. 2017. *R package version 00* **1**.

Baker, R.L. & Diggle, P.K. 2011. Node-specific branching and heterochronic changes underlie population-level differences in Mimulus guttatus (Phrymaceae) shoot architecture. *Am. J. Bot.* **98**: 1924–1934. Wiley.

Baker, R.L., Hileman, L.C. & Diggle, P.K. 2012. Patterns of shoot architecture in locally adapted populations are linked to intraspecific differences in gene regulation. *New Phytol.* **196**: 271–281.

Barbour, M.G. 1978. Salt spray as a microenvironmental factor in the distribution of beach plants at point reyes, California. *Oecologia* **32**: 213–224.

Barker, W., Nesom, G., Beardsley, P. & Fraga, N. 2012. A taxonomic conspectus of Phrymaceae: A narrowed circumscription for Mimulus, new and resurrected genera, and new names and combinations. 1–60.

Bazzaz, F.A., Chiariello, N.R., Coley, P.D. & Pitelka, L.F. 1987. Allocating Resources to Reproduction and Defense. *Bioscience* **37**: 58–67. [American Institute of Biological Sciences, Oxford University Press].

Boyce, S.G. 1954. The Salt Spray Community. *Ecol. Monogr.* **24**: 29–67. Ecological Society of America.

Briscoe Runquist, R.D., Gorton, A.J., Yoder, J.B., Deacon, N.J., Grossman, J.J., Kothari, S., *et al.* 2020. Context Dependence of Local Adaptation to Abiotic and Biotic Environments: A Quantitative and Qualitative Synthesis. *Am. Nat.* **195**: 412–431. journals.uchicago.edu.

Broad, Z., Lefreve, J., Wilkinson, M.J., Barton, S., Barbier, F., Jung, H., *et al.* 2024. Gravitropic gene expression divergence associated with adaptation to contrasting environments in an Australian wildflower. *Mol. Ecol.* e17543. Wiley.

Brooks, M.E., Kristensen, K. & Van Benthem, K.J. 2017. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R.* research-collection.ethz.ch.

Campos, M.L., Yoshida, Y., Major, I.T., de Oliveira Ferreira, D., Weraduwage, S.M., Froehlich, J.E., *et al.* 2016. Rewiring of jasmonate and phytochrome B signalling uncouples plant growth-defense tradeoffs. *Nat. Commun.* **7**: 12570. nature.com.

Cipollini, D., Walters, D. & Voelckel, C. 2017. Costs of resistance in plants: From theory to
evidence. John Wiley & Sons, Ltd, Chichester, UK.

Clausen, J., Keck, D.D. & Hiesey, W.M. 1940. *Experimental studies on the nature of species. I.*
Effect of varied environments on western North American plants. Carnegie Institution of
Washington Publication No. 520., Washington D.C.

Crone, E.E. 2001. Is survivorship a better fitness surrogate than fecundity? *Evolution* **55**: 2611–
2614.

DeMarche, M.L., Kay, K.M. & Angert, A.L. 2016. The scale of local adaptation in Mimulus
guttatus: comparing life history races, ecotypes, and populations. *New Phytol.* **211**: 345–
356.

Du, J. & Hesp, P.A. 2020. Salt Spray Distribution and Its Impact on Vegetation Zonation on
Coastal Dunes: a Review. *Estuaries Coasts* **43**: 1885–1907. Springer.

Eiseman, C.S., Namayandeh, A., Linden, J.V.A.N.D.E.R. & Palmer, M.W. 2023. Metriocnemus
erythranthei sp. nov. and Limnophyes viribus sp. nov. (Diptera: Chironomidae:
Orthocladiinae): leafminers of monkeyflowers, speedwells, and other herbaceous plants,
with new observations on the ecology and habitats of other leaf-mining Chironomidae.
Zootaxa **5249**: 41–68.

Feeny, P. 1976. Plant Apparency and Chemical Defense. In: *Biochemical Interaction Between*
Plants and Insects (J. W. Wallace & R. L. Mansell, eds), pp. 1–40. Springer US, Boston,
MA.

Fox, J. & Weisberg, S. 2018. *An R Companion to Applied Regression*. SAGE Publications.

Friberg, M., Schwind, C., Guimarães, P.R., Jr, Raguso, R.A. & Thompson, J.N. 2019. Extreme
diversification of floral volatiles within and among species of Lithophragma
(Saxifragaceae). *Proc. Natl. Acad. Sci. U. S. A.* **116**: 4406–4415.

Gasperini, D. & Howe, G.A. 2024. Phytohormones in a universe of regulatory metabolites:
lessons from jasmonate. *Plant Physiol.* **195**: 135–154. Oxford University Press (OUP).

Gotelli, N.J. & Ellison, A.M. 2004. *A primer of ecological statistics*. Sinauer Associates,
Sunderland, MA.

Hahn, P.G. & Maron, J.L. 2016. A Framework for Predicting Intraspecific Variation in Plant
Defense. *Trends Ecol. Evol.* **31**: 646–656. cell.com.

Hall, M.C., Lowry, D.B. & Willis, J.H. 2010. Is local adaptation in Mimulus guttatus caused by
trade-offs at individual loci? *Mol. Ecol.* **19**: 2739–2753. Wiley.

Hall, M.C. & Willis, J.H. 2006. Divergent selection on flowering time contributes to local
adaptation in Mimulus guttatus populations. *Evolution* **60**: 2466–2477. Wiley Online
Library.

Hargreaves, A.L., Germain, R.M., Bontrager, M., Persi, J. & Angert, A.L. 2020. Local
Adaptation to Biotic Interactions: A Meta-analysis across Latitudes. *Am. Nat.* **195**: 395–
411. journals.uchicago.edu.

Hartig, F. 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)
Regression Models (R Package Version 0.4 5). 2022.

Havko, N.E., Major, I.T., Jewell, J.B., Attaran, E., Browse, J. & Howe, G.A. 2016. Control of
Carbon Assimilation and Partitioning by Jasmonate: An Accounting of Growth–Defense
Tradeoffs. *Plants* **5**: 7. Multidisciplinary Digital Publishing Institute.

Heil, M. & Baldwin, I.T. 2002. Fitness costs of induced resistance: emerging experimental
support for a slippery concept. *Trends Plant Sci.* **7**: 61–67.

Herbivory Variability Network*†, Robinson, M.L., Hahn, P.G., Inouye, B.D., Underwood, N.,
Whitehead, S.R., *et al.* 2023. Plant size, latitude, and phylogeny explain within-population
variability in herbivory. *Science* **382**: 679–683.

Hereford, J. 2009. A quantitative survey of local adaptation and fitness trade-offs. *Am. Nat.* **173**:
579–588. journals.uchicago.edu.

Herms, D.A. & Mattson, W.J. 1992. The Dilemma of Plants: To Grow or Defend. *Q. Rev. Biol.*
67: 283–335. The University of Chicago Press.

Holeski, L.M. 2007. Quantitative trait evolution in Mimulus guttatus (yellow monkeyflower).
search.proquest.com.

Holeski, L.M., Chase-Alone, R. & Kelly, J.K. 2010. The genetics of phenotypic plasticity in
plant defense: trichome production in Mimulus guttatus. *Am. Nat.* **175**: 391–400.

Holeski, L.M., Keefover-Ring, K., Bowers, M.D., Harnenz, Z.T. & Lindroth, R.L. 2013. Patterns
of phytochemical variation in Mimulus guttatus (yellow monkeyflower). *J. Chem. Ecol.* **39**:
525–536.

Holeski, L.M., Monnahan, P., Koseva, B., McCool, N., Lindroth, R.L. & Kelly, J.K. 2014. A
high-resolution genetic map of yellow monkeyflower identifies chemical defense QTLs and
recombination rate variation. *G3* **4**: 813–821.

Hou, X., Ding, L. & Yu, H. 2013. Crosstalk between GA and JA signaling mediates plant growth
and defense. *Plant Cell Rep.* **32**: 1067–1074. Springer.

Huot, B., Yao, J., Montgomery, B.L. & He, S.Y. 2014. Growth-defense tradeoffs in plants: a
balancing act to optimize fitness. *Mol. Plant* **7**: 1267–1287. Elsevier BV.

Iqbal, M. & Ashraf, M. 2013. Gibberellic acid mediated induction of salt tolerance in wheat
plants: Growth, ionic partitioning, photosynthesis, yield and hormonal homeostasis.
Environ. Exp. Bot. **86**: 76–85. Elsevier.

Itoh, M., Fukunaga, K. & Osako, T. 2024. Local adaptation in parapatric and sympatric mosaic
coastal habitats through trait divergence of Setaria viridis. *J. Ecol.* **112**: 784–799. Wiley.

James, M.E., Allsopp, R.N., Groh, J.S., Kaur, A., Wilkinson, M.J. & Ortiz-Barrientos, D. 2023.
Uncovering the genetic architecture of parallel evolution. *Mol. Ecol.* **32**: 5575–5589.

Kawecki, T.J. & Ebert, D. 2004. Conceptual issues in local adaptation. *Ecol. Lett.* **7**: 1225–1241.
Wiley.

Kazan, K. & Manners, J.M. 2012. JAZ repressors and the orchestration of phytohormone
crosstalk. *Trends Plant Sci.* **17**: 22–31.

Kollar, L.M., Stanley, L.E., Kenchanmane Raju, S.K., Lowry, D.B. & Niederhuth, C.E. 2025.
The evolution of locally adaptive chromosome inversions in Mimulus guttatus. *Mol. Ecol.*
e17708. Wiley.

Kooyers, N.J., Blackman, B.K. & Holeski, L.M. 2017. Optimal defense theory explains
deviations from latitudinal herbivory defense hypothesis. *Ecology* **98**: 1036–1048.

Krimmel, B. & Pearse, I.S. 2016. Tolerance and phenological avoidance of herbivory in tarweed
species. *Ecology* **97**: 1357–1363. Wiley.

Leimu, R. & Fischer, M. 2008. A meta-analysis of local adaptation in plants. *PLoS One* **3**:
e4010. journals.plos.org.

Lenth, R., Buerkner, P., Herve, M., Love, J., Riebl, H. & Singmann, H. 2020. emmeans:
Estimated marginal means, aka least-squares means (1.5. 2-1)[Computer software].

Liu, J., Wu, Y., Dong, G., Zhu, G. & Zhou, G. 2023. Progress of Research on the Physiology and
Molecular Regulation of Sorghum Growth under Salt Stress by Gibberellin. *Int. J. Mol. Sci.*
24.

López-Goldar, X. & Agrawal, A.A. 2021. Ecological interactions, environmental gradients, and
gene flow in local adaptation. *Trends Plant Sci.* **26**: 796–809. Elsevier BV.

Lowry, D.B., Hall, M.C., Salt, D.E. & Willis, J.H. 2009. Genetic and physiological basis of
adaptive salt tolerance divergence between coastal and inland Mimulus guttatus. *New*
Phytol. **183**: 776–788. Wiley Online Library.

Lowry, D.B., Popovic, D., Brennan, D.J. & Holeski, L.M. 2019. Mechanisms of a locally
adaptive shift in allocation among growth, reproduction, and herbivore resistance in
Mimulus guttatus. *Evolution* **73**: 1168–1181.

Lowry, D.B., Rockwood, R.C. & Willis, J.H. 2008. Ecological reproductive isolation of coast
and inland races of Mimulus guttatus. *Evolution* **62**: 2196–2214. academic.oup.com.

Lüdecke, D. 2018. Ggeffects: Tidy data frames of marginal effects from regression models. *J.*
Open Source Softw. **3**: 772. The Open Journal.

Maron, J.L., Baer, K.C. & Angert, A.L. 2014. Disentangling the drivers of context☐dependent
plant–animal interactions. *J. Ecol.* **102**: 1485–1496. Wiley.

Monson, R.K., Trowbridge, A.M., Lindroth, R.L. & Lerdau, M.T. 2022. Coordinated resource
allocation to plant growth-defense tradeoffs. *New Phytol.* **233**: 1051–1066. Wiley.

Murphy, A. 2015. Hormone crosstalk in plants. *J. Exp. Bot.* **66**: 4853–4854. academic.oup.com.

Oksanen, J. 2016. Vegan: ecological diversity. *R project*. mirror.linux.duke.edu.

Olsen, J.D., Gould, B.A., Chen, Y. & Lowry, D.B. 2024. Gene expression divergence between
locally adapted inland annual and coastal perennial ecotypes of Mimulus guttatus across
developmental stages.

Pedersen, T.L. 2019. Package "patchwork." *R package http://CRAN. R-project. org/package=*
patchwork. Cran.

Popovic, D. & Lowry, D.B. 2020. Contrasting environmental factors drive local adaptation at
opposite ends of an environmental gradient in the yellow monkeyflower (Mimulus
guttatus). *Am. J. Bot.* **107**: 298–307. Wiley Online Library.

Rhoades, D.F. 1979. Evolution of plant chemical defenses against herbivores. *Herbivores-Their*
interaction with secondary plant metabolites 3–48. Academic Press.

Rodríguez, A.A., Stella, A.M., Storni, M.M., Zulpa, G. & Zaccaro, M.C. 2006. Effects of
cyanobacterial extracellular products and gibberellic acid on salinity tolerance in Oryza
sativaL. *Saline Systems* **2**: 7.

Rotter, M.C., Christie, K. & Holeski, L.M. 2022. Climate and the biotic community structure
plant resistance across biogeographic groups of yellow monkeyflower. *Ecol. Evol.* **12**:
e9520.

Rotter, M.C., Couture, J.J. & Rothwell, E.M. 2018. Evolutionary ecology of plant resistance
traits across the herbivore diet spectrum: a test in the model plant Mimulus guttatus.
Evolutionary. evolutionary-ecology.com.

Stamp, N. 2003. Out of the quagmire of plant defense hypotheses. *Q. Rev. Biol.* **78**: 23–55.

Stevens, M.T., Kruger, E.L. & Lindroth, R.L. 2008. Variation in Tolerance to Herbivory Is
Mediated by Differences in Biomass Allocation in Aspen. *Funct. Ecol.* **22**: 40–47. [British
Ecological Society, Wiley].

Stowe, K.A. & Marquis, R.J. 2011. Costs of defense: correlated responses to divergent selection
for foliar glucosinolate content in Brassica rapa. *Evol. Ecol.* **25**: 763–775. Springer.

Strauss, S.Y., Rudgers, J.A., Lau, J.A. & Irwin, R.E. 2002. Direct and ecological costs of
resistance to herbivory. *Trends Ecol. Evol.* **17**: 278–285. Elsevier.

Therneau, T.M. 2022. Mixed Effects Cox Models [R package coxme version 2.2-18.1].
Comprehensive R Archive Network (CRAN).

Thompson, J.N. 2005. *The Geographic Mosaic of Coevolution*. University of Chicago Press.

Urban, M.C. 2011. The evolution of species interactions across natural landscapes. *Ecol. Lett.*
14: 723–732.

VanWallendael, A., Soltani, A., Emery, N.C., Peixoto, M.M., Olsen, J. & Lowry, D.B. 2019. A
molecular view of plant local adaptation: Incorporating stress-response networks. *Annu.*
Rev. Plant Biol. **70**: 559–583. Annual Reviews.

Wadgymar, S.M., DeMarche, M.L., Josephs, E.B., Sheth, S.N. & Anderson, J.T. 2022. Local
adaptation: Causal agents of selection and adaptive trait divergence. *Annu. Rev. Ecol. Evol.*
Syst. **53**: 87–111.

Wadgymar, S.M., Lowry, D.B. & Gould, B.A. 2017. Identifying targets and agents of selection:
innovative methods to evaluate the processes that contribute to local adaptation. *Methods*
Ecol. Evol. Wiley Online Library.

Wang, X., Chen, X., Wang, Q., Chen, M., Liu, X., Gao, D., *et al.* 2019. MdBZR1 and MdBZR1-
2like Transcription Factors Improves Salt Tolerance by Regulating Gibberellin Biosynthesis
in Apple. *Front. Plant Sci.* **10**: 1473.

Wang, Y., Gong, X., Liu, W., Kong, L., Si, X., Guo, S., *et al.* 2020. Gibberellin mediates
spermidine-induced salt tolerance and the expression of GT-3b in cucumber. *Plant Physiol.*
Biochem. **152**: 147–156. Elsevier.

Warton, D.I. & Hui, F.K.C. 2011. The arcsine is asinine: the analysis of proportions in ecology.
Ecology **92**: 3–10. Wiley.

Wickham, H. 2016. Data Analysis. In: *ggplot2: Elegant Graphics for Data Analysis* (H.
Wickham, ed), pp. 189–201. Springer International Publishing, Cham.

Wilkinson, M.J., Roda, F., Walter, G.M., James, M.E., Nipper, R., Walsh, J., *et al.* 2021.
Adaptive divergence in shoot gravitropism creates hybrid sterility in an Australian
wildflower. *Proc. Natl. Acad. Sci. U. S. A.* **118**.

Yang, D.-L., Yao, J., Mei, C.-S., Tong, X.-H., Zeng, L.-J., Li, Q., *et al.* 2012. Plant hormone
jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade.
Proc. Natl. Acad. Sci. U. S. A. **109**: E1192–200. National Acad Sciences.

Yan, Y., Stolz, S., Chételat, A., Reymond, P., Pagni, M., Dubugnon, L., *et al.* 2007. A
downstream mediator in the growth repression limb of the jasmonate pathway. *Plant Cell*
19: 2470–2483. academic.oup.com.

Zambiasi, T. & Lowry, D.B. 2024. Ocean exposure and latitude drive multiple clines within the
coastal perennial ecotype of the yellow monkeyflower, Mimulus guttatus. *Am. J. Bot.* **111**:

e16402. Wiley.

Zhang, Y. & Turner, J.G. 2008. Wound-induced endogenous jasmonates stunt plant growth by
inhibiting mitosis. *PLoS One* **3**: e3699.

Züst, T. & Agrawal, A.A. 2017. Trade-Offs Between Plant Growth and Defense Against Insect
Herbivory: An Emerging Mechanistic Synthesis. *Annu. Rev. Plant Biol.* **68**: 513–534.
annualreviews.org.

Figure Legends

Figure 1. Photographs depicting transplant sites and structures used in the reciprocal transplant experiment. For our reciprocal transplant experiment, each site had ten plots, each with 100 plants. Each site had six control plots (with mesh tops and partially-mesh sides) and four exclosure plots (fully enclosed with mesh on all sides). A control and exclosure plot are shown side by side in the foreground of the inland site.

Figure 2. Flowering time and probability of herbivory of annuals (circles) and perennials (triangles) treated with gibberellic acid (GA, yellow), paclobutrazol (blue), and methyl jasmonate (MeJA, green), and the no-hormone controls (black) in open structures and herbivore exclosures at the coastal site, Bodega Marine Reserve (A & C), and the inland site, Pepperwood Preserve (B & D). Larger symbols in the foreground are the mean predictions and 95% confidence intervals from the minimum adequate mixed models, smaller and lighter symbols in the background are the raw data. Since no annuals treated with paclobutrazol in exclosures at the inland site were damaged by herbivores (d), the binomial model did not accurately estimate that parameter and thus we did not plot those confidence intervals (0-100%). Results of Tukey post-hoc contrasts within each site are indicated above each prediction; shared letters indicate that groups do not significantly differ, while non-overlapping letters indicate that groups significantly differ within each site. Exclosure type was not plotted for flowering time on the coast (A) because the minimum adequate model did not include exclosure type as a fixed effect.

Figure 3. Allocation to chemical defense: total concentration of all PPGs and differences in multivariate PPG arsenals of annuals (circles) and perennials (triangles) treated with gibberellic

acid (GA, yellow), paclobutrazol (blue), and methyl jasmonate (MeJA, green), and the no-

hormone controls (black) in open structures and herbivore exclosures at the coastal site, Bodega

Marine Reserve (A & C), and the inland site, Pepperwood Preserve (B & D). In the Total PPG

figures (A & B), larger symbols in the foreground are the mean predictions and 95% confidence

intervals from the minimum adequate mixed models, smaller and lighter symbols in the

background are the raw data. Exclosure type was not plotted for Total PPG at the inland site (B)

because the minimum adequate model did not include exclosure type as a fixed effect. PPG

arsenal figures (C & D) use non-metric multidimensional scaling (NMDS) with Bray-curtis

distance (and 95% confidence interval ellipses) to visualize multivariate differences among

plants. Exclosure type was either not a significant factor in the multivariate model (C) or did not

interact with other fixed effects (D) and was therefore not included in these plots.

Figure 4. GA_3 application decreased survival at the coastal site. Survival probabilities for annual

(solid line) and perennial (dashed line) transplants at the coastal site, Bodega Marine Reserve

(A), and the inland site, Pepperwood Preserve (B). Survival probabilities and 95% confidence

intervals for control (black), GA (yellow), methyl jasmonate (green) and paclobutrazol (blue)

treatments were predicted from Cox Proportional Hazards models. At the inland transplant site

(B), survival probabilities and 95% confidence intervals were only plotted for ecotypes (grey)

because the minimum adequate model did not include exclosure or hormone treatment as a fixed

effect. Results of Tukey post-hoc contrasts within each site are indicated above each final

predicted survival; shared letters indicate that groups do not significantly differ, while non-

overlapping letters indicate that groups significantly differ within each site.

Figure 5. GA$_3$ application on perennials resulted in stem-elongation relative to controls. The

plants pictured (on day 16 after transplantation) are from the same family and were grown in the

same plot at the coastal site.

Figure 6. Herbivore exclosures tended to increase, while GA$_3$ applications tended to decrease

components of fecundity: probability of flowering and fruit number of annuals (circles) and

perennials (triangles) that flowered treated with gibberellic acid (GA, yellow), paclobutrazol

(blue), and methyl jasmonate (MeJA, green), and the no-hormone controls (black circles) in open

structures and herbivore exclosures at the coastal site, Bodega Marine Reserve (A & C), and the

inland site, Pepperwood Preserve (B & D). Larger symbols in the foreground are the mean

predictions and 95% confidence intervals from mixed models, smaller and lighter symbols in the

background are the raw data. Since all annuals flowered in exclosures at the inland site (b), the

binomial model did not accurately estimate that parameter and thus we did not plot those

confidence intervals (0-100%). Results of Tukey post-hoc contrasts within each site are indicated

above each prediction; shared letters indicate that groups do not significantly differ, while non-

overlapping letters indicate that groups significantly differ within each site. Predictions are not

plotted for fruit production at the coastal site because no fixed factors were in the minimum

adequate model. To improve visualization, one outlier that produced 164 fruit was not plotted at

the coastal site.

1006

Figure 7. GA$_3$ treatment increased plant height in a greenhouse, and increased leaf damage of

plants exposed to salt spray in 2023. (a) Mean predictions and 95% confidence intervals from a

linear mixed model, where the response variable was height in the greenhouse, the fixed

predictor variables were population, GA$_3$ treatment and their interaction, and flat was a random

effect. (b) Mean predictions and 95% confidence intervals of a linear mixed model, where the

response variable was logit transformed leaf damage in blocks exposed to the ocean, the fixed

predictor variables were population, GA$_3$ treatment, and their interaction, and flat nested within

block was a random effect. Horizontal lines and asterisks (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$)

depict significant pairwise comparisons between GA$_3$-treated plants and controls.

1016
1017 **Figure 1.**
1018

1019
1020 **Figure 2.**
1021

1022
1023　**Figure 3.**

1024
1025　**Figure 4.**
1026

1027
1028 **Figure 5.**

1029
1030 **Figure 6.**

1031
1032 **Figure 7.**