Virulence of *Fusarium oxysporum* strains causing corneal or plant disease is associated with their

distinct accessory chromosomes

Dilay Hazal Ayhan[1,2,3], Serena Abbondante[4], Domingo Martínez-Soto[5], Siyuan Wu[1,2], Ricardo

Rodriguez-Vargas[5], Shira Milo[1], Katherine Rickelton[1,2], Vista Sohrab[1‡], Shunsuke Kotera[6†], Tsutomu

Arie[6], Michaela Ellen Marshall[4], Marina Campos Rocha[7], Sajeet Haridas[8], Igor V. Grigoriev[8], Neta

Shlezinger[7], Eric Pearlman[4], Li-Jun Ma [1,2 #]

[1] Biochemistry and Molecular Biology, [2] Molecular and Cellular Biology, University of Massachusetts

Amherst, Amherst, MA, USA

[3]Graduate School of Natural and Applied Sciences, Acibadem University, Istanbul, Türkiye

[4]Physiology and Biophysics and Ophthalmology, University of California, Irvine, USA

University of Massachusetts Amherst, Amherst, MA, USA

[5] Department of Microbiology, Centro de Investigación Científica y Educación Superior de Ensenada

(CICESE), Ensenada, Baja California, Mexico.

[6] Laboratory of Plant Pathology, Graduate School of Agriculture, Tokyo University of Agriculture and

Technology (TUAT), Fuchu, Tokyo, Japan

[7] The Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of

Jerusalem, Jerusalem, Israel

[8] US Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley,

CA 94720, USA

†Present address: Mitsui chemicals crop & life solutions, Inc., Mobara-shi, Chiba, 297-0017, Japan

‡ Undergraduate student.

Running head (53 out of 54 total): Accessory chromosomes in plant versus human pathogens

[#] **Address correspondence to Li-Jun Ma** (lijun@biochem.umass.edu).

Keywords

Fusarium oxysporum, cross-kingdom fungal pathogen, corneal infection, tomato vascular wilt,

accessory chromosomes, comparative genomics

ABSTRACT (231 out of 250 words)

Fusarium oxysporum is a cross-kingdom pathogen. While some strains cause disseminated

fusariosis and blinding corneal infections in humans, others are responsible for devastating

vascular wilt diseases in plants. To better understand the distinct adaptations of *F. oxysporum* to

animal or plant hosts, we conducted a comparative phenotypic and genetic analysis of two

strains: MRL8996 (isolated from a keratitis patient) and Fol4287 (isolated from a wilted tomato

[*Solanum lycopersicum*]). Infection of mouse corneas and tomato plants revealed that, while both

strains cause symptoms in both hosts, MRL8996 caused more severe corneal disease in mice,

whereas Fol4287 induced more pronounced wilting symptoms in tomato plants. *In vitro* assays

using abiotic stress treatments revealed that the human pathogen MRL8996 was better adapted to

elevated temperatures, whereas the plant pathogen Fol4287 was more tolerant to osmotic and cell

wall stresses. Both strains displayed broad resistance to antifungal treatment, with MRL8996

exhibiting the paradoxical effect of increased tolerance to higher concentrations of the antifungal

caspofungin. We identified a set of accessory chromosomes (ACs) that encode genes with

different functions and have distinct transposon profiles between MRL8996 and Fol4287.

Interestingly, ACs from both genomes also encode proteins with shared functions, such as

chromatin remodeling and post-translational protein modifications. Our phenotypic assays and

comparative genomics analyses lay the foundation for future studies correlating genotype with

phenotype and for developing targeted antifungals for agricultural and clinical uses.

Importance (146 out of 150 words)

Fusarium oxysporum is a cross-kingdom fungal pathogen that infects both plants and animals. In

addition to causing many devastating wilt diseases, this group of organisms was recently

recognized by the World Health Organization as a high-priority threat to human health. Climate

change has increased the risk of *Fusarium* infections, as *Fusarium* strains are highly adaptable to

changing environments. Deciphering fungal adaptation mechanisms is crucial to developing

appropriate control strategies. We performed a comparative analysis of *Fusarium* strains using

an animal (mouse) and plant (tomato) host and *in vitro* conditions that mimic abiotic stress. We

also performed comparative genomics analyses to highlight the genetic differences between

human and plant pathogens and correlate their phenotypic and genotypic variations. We

uncovered important functional hubs shared by plant and human pathogens, such as chromatin

modification, transcriptional regulation, and signal transduction, which could be used to identify

novel antifungal targets.

Main text (4,971 out of 5,000 words)

INTRODUCTION

Fusarium oxysporum, a cross-kingdom pathogen, is included in the list of health-

threatening fungi by the World Health Organization (WHO) (1) and is considered to be among

the top five most important plant pathogens (2). Corneal infections (keratitis) caused by

Fusarium pathogens are an important cause of blindness worldwide, resulting in over one million

new cases of blindness annually (3, 4). Indeed, *Fusarium* spp. are the most common cause of

fungal keratitis in India (5, 6), China (7, 8), South Africa (9), and Brazil (10). *F. oxysporum* was

also the cause of the 2005–2006 keratitis outbreak among contact lens wearers in the US and

other temperate regions of the world (11, 12). As a plant pathogen, *F. oxysporum* causes

devastating vascular wilt diseases in many economically important crops, including tomato

(*Solanum lycopersicum*) (13), cotton (*Gossypium hirsutum*) (14, 15), and banana (*Musa* sp.) (16),

and is responsible for billions of dollars in annual yield losses. Disease severity caused by this

cross-kingdom pathogen is compounded by the lack of effective drugs, as only a limited number

of antifungal agents are available to control eukaryotic fungal pathogens, and most *Fusarium*

isolates are notorious for their broad resistance to many of these drugs (17–19).

Host-specific virulence is well recognized among plant pathogenic *F. oxysporum* isolates,

as a single *F. oxysporum* species complex (FOSC) member typically infects only one or two

plant species and is recognized as a *forma specialis* among plant pathogens. Within the FOSC,

over 100 recognized *F. oxysporum formae speciales* infect more than 100 diverse plant hosts

(20). We previously identified horizontally inherited accessory chromosomes (ACs) that lack

genes associated with housekeeping functions but are enriched for genes related to fungal

virulence as determinants of host-specific pathogenicity within the FOSC (21–24). Therefore,

ACs (also referred to as pathogenicity chromosomes) encode several functionally validated

virulence factors toward plant hosts, such as Secreted in Xylem (*SIX*) effectors (25), transcription

factors (26), and kinases (27, 28). The importance of *F. oxysporum* ACs was demonstrated by

the finding that a non-pathogenic *F. oxysporum* strain became virulent upon receiving ACs from

a pathogenic strain (21, 29, 30).

No specific *forma specialis* has been reported for clinical isolates of *F. oxysporum*. The

conventional wisdom was that disseminated fusariosis is the result of opportunistic infection of

immunocompromised patients by environmental isolates, and corneal infections occur as a result

of ocular trauma or contaminated contact lenses. The finding that a plant pathogenic isolate was

able to infect immunosuppressed mice and cause systemic disease supports this notion (31).

However, we reported that distinct sets of ACs were reported in the genomes of two clinical

FOSC strains (32) and found to lack the genomic signatures that define plant pathogenic ACs,

including *SIX* effector genes and some repeat elements commonly present upstream of these

genes (33). By contrast, these two clinical isolates share AC regions that are enriched in genes

encoding metal ion transporters and cation transporters (32).

Using the well-established murine model of fungal keratitis (4) and the tomato vascular

wilt models (25), this study compared the human pathogenic strain MRL8996 isolated from a

keratitis patient and the plant pathogenic strain Fol4287 isolated from a diseased tomato to test

the hypothesis that distinct ACs contribute to the unique adaptation of fungal strains to animal or

plant hosts. Both strains exhibited greater virulence in their respective hosts. Although we

observed cross-kingdom virulence, these interactions resulted in milder disease in plants and in

animals. These observations were supported by *in vitro* studies showing that MRL8996 was

better adapted to higher temperatures, while Fol4287 tolerated conditions imposing osmotic or

cell wall stress. These phenotypic assays revealed the unique adaptations of human and plant

pathogens, providing a platform to dissect their different interactions with diverse hosts.

Comparative genomics highlighted distinct genetic elements unique to each strain, offering

potential testable hypotheses to correlate phenotypic to genotypic variations between a human

and a plant pathogen. We also uncovered important functional hubs in ACs used by both human

and plant pathogens, including chromatin modification, transcriptional regulation, and signal

transduction, potentially identifying novel antifungal targets.

RESULTS

COMPARATIVE PHENOMICS REVEALS HOST-SPECIFIC ADAPTATION

We selected two *F. oxysporum* strains, one isolated from a keratitis patient and one from

a tomato plant showing wilting symptoms as representative human and plant pathogens.

MRL8996 was originally isolated from an infected cornea of a patient in the 2005–2006 contact

lens–associated keratitis outbreak cohort (34). In addition, the genome of this strain has been

published (35) and a clinically relevant mouse model of keratitis has been established (36).

Phylogenetically, this keratitis strain is grouped with other human pathogenic isolates that cause

systemic disease (37, 38). The tomato pathogenic strain Fol4287 was originally isolated from a

diseased tomato plant in Spain (39, 40) has been adopted by the international Fusarium research

community as a reference strain (21).

MRL8996 is more virulent than Fol4287 in infected mouse corneas

To identify potential differences in virulence between Fol4287 and MRL8996 during

corneal infection, we infected corneas of C57BL/6 mice with 5,000 *Fusarium* conidia from each

strain (inoculum concentrations from 5,000 to 20,000 conidia are shown in Supplementary Fig.

S1). After 24 h, we observed significantly higher corneal opacification MRL8996-inoculated eyes (**Fig. 1A, B**), compared to Fol428-inoculated corneas (**Fig. 1A, B**). We also examined fungal viability following eye infection with either fungal strain and detected significantly more colony-forming units (CFUs) in corneas infected with MRL8996 compared to Fol4287 (**Fig. 1C, Fig. S1**), indicating that the host was more efficient at containing the infection and possibly killing the Fol4287 strain compared to MRL8996.

The major cause of corneal disease following fungal infection is the recruitment of CD45+ myeloid cells, including neutrophils and monocytes to the corneal stroma (4). To quantify these cells following infection with each *F. oxysporum* strain, we dissected corneas that were infected with MRL8996 or Fol4287, digested them with collagenase, and incubated the infiltrating cells with antibodies against CD45 and CD11b (which recognize total myeloid cells) and Ly6G and Ly6C (which identify neutrophils and monocytes, respectively) (41). Consistent with disease severity, corneas infected with MRL8996 had a higher percentage of CD45+ myeloid cells than those infected with Fol4287 (86.6% and 75.1%, respectively) (**Fig. 1D-i**). However, ~90% of CD45+ myeloid cells were Ly6G+ neutrophils in corneas infected with each fungal strain (**Fig. 1D-ii**). Due to the higher percentage of CD45+ myeloid cells, we observed significantly more Ly6G+/CD11b+ neutrophils in corneas infected with the keratitis strain MRL8996 compared to the plant pathogenic strain Fol4287 (**Fig. 1E**). There was no significant difference in the number of Ly6C+/CD11b+ monocytes (**Fig. 1D-iii, E**).

Collectively, these findings indicate that the clinical isolate MRL8996 causes more severe disease and survives better than the plant pathogen Fol4287 in mouse corneas.

Fol4287 is more virulent than MRL8996 in infected tomato plants

To examine the relative disease severity caused by these two *F. oxysporum* strains in plants, we inoculated tomato seedlings with MRL8996 or Fol4287 using a well-established wilting disease assay (42–44). Briefly, we soaked cleaned roots of tomato seedlings in a 10^6 spores/mL suspension for 45 min before gently transplanting the inoculated seedlings in sterilized soil. As a mock control, roots were incubated in sterile distilled water. We used a disease index scale from 1 to 5 to measure disease severity (45), with 1 for healthy plants; 2 for plants showing wilted leaves with chlorotic areas; 3 for plants with necrotic spots; 4 for plants with wilted leaves and whole plants showing chlorosis, areas of necrosis, and defoliation; and 5 for dead plants.

At 10 days post-inoculation (dpi), the mock-inoculated seedlings were completely healthy, with a disease index of 1. By contrast, all tomato seedlings infected with the plant pathogen Fol4287 developed severe wilt symptoms, with an average disease index of 4, with wilting leaves showing areas of chlorosis, necrosis, and plant defoliation. Although the keratitis strain MRL8996 also caused chlorosis, the disease symptoms were significantly less pronounced, with an average disease index of 2 (**Fig. 2A**, **B**). The distinction between these strains was even more pronounced when tracking fungal colonization by confocal microscopy after wheat germ agglutinin (WGA)-Alexa Fluor 488 (green) staining of fungal cell walls and propidium iodide staining (red) of plant cell walls (46). As reported for wilt pathogens (44), the plant pathogenic strain Fol4287 invaded the xylem tissues in both primary and lateral roots at 4 dpi. By contrast, the keratitis strain MRL8996 was restricted to the epidermis and part of the cortex cell layers in both primary or lateral roots (**Fig. S2**). Furthermore, we documented that only Fol4287 colonized

the vascular vessels of primary roots, lateral roots, and stems (**Fig. 2E**) and could be re-isolated

from inoculated plant stems (**Fig. 2 C**, **D**).

Overall, these findings demonstrate that the plant pathogen caused significantly greater

disease severity in tomato seedlings than the clinical pathogen.

MRL8996 and Fol4287 exhibit resistance to different abiotic stress conditions

To explore cross-kingdom adaptation to different environmental conditions, we exposed

the two *F. oxysporum* strains to the abiotic stresses of high salinity (0.6 M NaCl), oxidative

stress (1 mM H_2O_2), and cell wall stress (1 mg/mL Congo Red), or to different temperatures

(28°C or 34°C) and pH (5.0 or 7.4). The higher temperature and pH were chosen since they

reflect the conditions of the human cornea (47). We calculated the growth rates of each strain at

3 dpi under each condition as the slope of the growth curve (diameter of the colony/dpi).

Notably, the keratitis strain MRL8996 formed larger colonies than the plant pathogen Fol4287 in

rich and minimal media at both temperatures tested whereas Fol4287 had a slower growth rate

(**Fig. 3A, Table S1**, **Fig. S3A**).

To examine the tolerance of these strains to different stresses, we calculated the ratio of

their growth rates under two different conditions. For instance, to assess tolerance to different

temperatures, we calculated the ratio of the fungal growth rates between 34°C and 28°C

($GR_{34°C}/GR_{28°C}$). While both strains grew more slowly at the higher temperature, the growth rate

of the plant pathogen Fol4287 was significantly slower at this temperature compared to the

keratitis strain MRL8996 under all conditions tested (**Fig. 3B, C**). At 34°C, lowering the pH

from the human physiological pH (7.4) to an acidic pH (5) environment decreased the growth of

the keratitis strain by 40%. Our observation that the human pathogenic strain MRL8996 has

better tolerance to an elevated temperature of 34°C at pH 7.4 may reflect the adaptation of

MRL8996 to the mammalian environment.

Conversely, the plant pathogenic strain Fol4287 exhibited significantly higher tolerance

to osmotic stress (0.6 M NaCl) than MRL8996 (**Fig. 3C, Fig. S2B**). Compared to growth in

complete medium alone at 28°C, the growth of MRL8996 in the presence of 0.6 M NaCl

decreased by 16.7% relative to the control condition, whereas the growth of strain Fol4287

increased by 22.7%, revealing a significant difference in their response to osmotic stress ($p < 10^{-5}$

). At 34°C, both strains grew better under higher salinity conditions compared to the complete

medium alone. While growth of MRL8996 increased by 99.4% under these conditions, growth of

Fol4287 increased by 187.6% ($p < 10^{-9}$). Therefore, the plant pathogenic strain Fol4287 is better

adapted to higher salinity conditions than MRL8996 at both temperatures (**Fig. 3C, Fig. S2B**).

Surprisingly, 1 mM H_2O_2 treatment (to induce oxidative stress) did not affect the growth

rates of either strain at 28°C. However, the human pathogenic strain MRL8996 exhibited

resistance to oxidative stress, with a 28.4% increase in growth at 34°C compared to samples

grown in complete medium alone. Further, cell wall stress imposed by the addition of 1 mg/mL

Congo Red to the medium inhibited the growth of both strains. At 28°C, compared to growth in

YPD medium alone, the growth rates of MRL8996 and Fol4287 were significantly lower (by

58.8% and 52.9%, respectively). At 34°C, the growth rate was reduced by 33.3% for MRL8996

and 43.8% for Fol4287 compared to the control.

We repeated *in vitro* phenotyping of the two reported strains and added one additional

plant pathogenic isolate Fo5176 (infecting *Arabidopsis thaliana*), and one additional clinical

isolate NRRL 32931 (isolated from a leukemia patient) (**Fig. S3, Table S1**) and observed

consistent results. While growth rates were reduced for all four strains at 34ºC compared to 28ºC,

the reduction of the two plant strains was significantly greater than the two clinical isolates.

MRL8996 exhibited the highest tolerance to heat compared with Fo5176. The clinical isolate

NRRL 32931 grew significantly better at physiological pH (7.4) compared to the acidic pH (5)

for both Minimal Media and Complete Media (**Fig. S3B**). Also consistent with the previous

findings, the plant isolates show higher resistance to high salinity in both temperatures compared

to the two human clinical isolates. Fo5176 surprisingly shows high resistance to salinity and cell

wall stress under both temperatures, with the difference more pronounced at 34ºC (**Fig. S3C**).

Together, these findings indicate that the clinical isolate MRL8996 is better adapted to

the physical conditions of the animal host, such as elevated temperature, whereas the plant

pathogenic strain Fol4287 is more tolerant to increased salinity. These findings reflect a

complex, interconnected regulatory relationship between the physiological adaptation of the

fungus and fungal–host interactions.

MRL8998, but not Fol4287, exhibits tolerance to high caspofungin concentrations

Caspofungin acetate (CFA) is an antifungal agent which targets 1,3-β-glucan synthase that

mediates biosynthesis of one of the main components of the fungal wall (48, 49). However,

different fungi show diverse responses to CFA, as clinical isolates of *Candida albicans* and

Aspergillus fumigatus are sensitive to CFA (50), whereas some less common fungal human

pathogens, including *Fusarium* spp., are resistant to clinically relevant levels of CFA (51).

In the current study, the keratitis strain MRL8998 exhibited enhanced tolerance to high

caspofungin concentrations compared with Fol4287 (**Fig. 4A, Fig. S4**). The growth rate of the

plant pathogen Fol4287 decreased from 80% to 70% with an increase in caspofungin

concentration from 0.2 µg/mL to 8 µg/mL compared to growth in the absence of caspofungin,

showing a strong tolerance to caspofungin without a clear dose-dependent response. In contrast,

the growth rate of the keratitis strain MRL8996 dropped by 50% under 2 µg/mL caspofungin,

with a further reduction to a 35% decrease at 8 µg/mL, relative to the control (**Fig. 4B**). This

paradoxical caspofungin effect was also observed when measuring the time required to reach

50% conidial germination (**Fig. 4C**). While there was no significant difference in the time

needed for the plant pathogen Fol4287 to reach a 50% germination rate in the presence of

caspofungin, the human keratitis strain required a significantly longer time to reach 50%

gemination under 0.5 µg/mL caspofungin treatment compared to the no caspofungin control or 8

µg/mL caspofungin treatment. Together, these findings indicate the different responses toward

the antifungal caspofungin.

COMPARATIVE GENOMICS REVEALS STRAIN-SPECIFIC ACCESSORY

CHROMOSOMES WITH DISTINCT TRANSPOSON PROFILES

We used a contour-clamped homogeneous electric field (CHEF) gel to observe the

diversity of karyotypes among *F. oxysporum* genomes and confirmed the presence of ACs in

Fol4287 (21, 52), MRL8996 and NRRL 32931 (32) (**Fig. 5A**). Comparative genomics confirmed

the conservation of core chromosomes (CCs) (**Fig. 5B**, **Table S2**) and revealed three small CCs

(chromosomes 11, 12, and 13) that were less conserved, with an average of 94.0% sequence

identity over 35.1% coverage. These findings support the three-speed evolutionary hypothesis

(53). The conservation of CCs enabled us to identify ACs and sequences unique to the plant and

human strains. As highlighted in darker blue and darker green boxes in **Fig. 5B**, the genomes of

MRL8996 and Fol4287 contained 6.4 Mb and 9.8 Mb of accessory sequences corresponding to

12.8% and 18.7% of the total genome, respectively.

Our comparative analysis also revealed overall conservation of the mitochondrial genome

(99.0%) and highlighted a divergent sequence around open reading frame 2285 (ORF2285) (**Fig.**

S3), a known variable mitochondrial region in the genus *Fusarium* (54).

The repeat contents in the Fol4287 and MRL8996 genomes were similar, with 6.6% and

5.4% values, respectively (**File S1, S2**). Transposable elements (TEs), a signature of accessory

sequences (55), were enriched in the ACs of both genomes (27.2% and 23.8%, respectively)

(**Table S3**), but with distinct transposon profiles. In the Fol4287 genome, approximately one-

third of identified TEs were type I retrotransposons, and the another third was type II DNA

transposons. In the MRL8896 genome, more than half were identified as type I and less than

20% as type II TEs (**Fig. 5C**).

Some class I retrotransposons are conserved within the genus of *Fusarium* (21).

Similarly, the long terminal repeat (LTR) transposon *Copia*, was present in both the Fol4287 and

MRL8996 genomes (**Fig. 5D, Table S4**). However, some LTR elements with the highest copy

numbers in MRL8996, such as several *Gypsy* elements, were not present in Fol4287. Similarly,

the most abundant long interspersed nuclear element (LINE) transposon in MRL8996, the *Cnl1*

CRE-type non-LTR retrotransposon, was not present in Fol4287. Of the three short interspersed

nuclear element (SINE) transposon families, the *Foxy2* family was the most abundant in

Fol4287, while the *Foxy5* family was the most abundant in MRL8996. Conversely, Fol4287

contained unique class II DNA transposons (**Fig. 5D, Table S4**).

The most abundant DNA TE in Fol4287, *Hormin* (a miniature *Hornet* element), was

absent from the MRL8996 genome. Miniature impala elements (MIMPs) (33) were not present

in MRL8996. Similarly, Fol4287 contained significantly more Helitrons (56) compared to

MRL8996. There were also many unclassified repeat elements that were not shared between the

Fol4287 and MRL8996 genomes (**Fig. S4**). The distinction between different TE profiles is

important, as class I and class II TEs play different roles in the genome dynamics of MRL8996

and Fol4287.

STRAIN-SPECIFIC ACCESSORY CHROMOSOMES CONTRIBUTE TO DISTINCT

FUNCTIONAL ADAPTATIONS

We annotated the genome of MRL8996 using the Joint Genome Institute annotation

pipeline (57) and assigned Gene Ontology (GO) terms to genes in both genomes using the

Mycocosm GO annotation pipeline (**Table 1**). Among the 16,631 and 20,925 predicted protein-

coding genes in the MRL8996 and Fol4287 genomes, respectively, we identified 2,017 (12.1%)

and 3,890 (18.6%) genes located on their respective ACs. GO functional enrichments (*p*-value <

10^{-3}) revealed some functional categories shared by both pathogens (**Fig. 6** category III), in

addition to strain-specific functions (**Fig. 6** categories I and II, **Table S5, S6**).

Functional groups enriched in both genomes due to different sets of AC genes

1) *Chromatin assembly or disassembly*. The most significantly enriched functional term

in both genomes was chromatin assembly or disassembly (GO:0006333), with $p = 6 \times 10^{-12}$ for

MRL8996 AC genes and $p = 7 \times 10^{-15}$ for Fol4287 AC genes. Most genes under this annotation

encode CHROMO domain-containing proteins that bind to the H3K9 di/trimethyl modification

on histone H3, a hallmark of heterochromatic regions (58, 59) that are typically involved in the

assembly or remodeling of chromatin and associated proteins or RNAs (60). Phylogenetic

analysis grouped these genes into three clades (**Fig. S5**). The first clade shares homologs of

CHROMO domain-containing protein 2 (Chp2) from fission yeast (*Schizosaccharomyces*

pombe). The second clade encodes homologs of putative Heterochromatin protein 1 (HP1)

(A0A0C4BKY0) based on AlphaFold prediction (61). The third clade includes genes encoding

Zinc finger C2H2-type domain-containing proteins. Notably, Chp2 and HP1 both interact with

chromatin in DNA–protein complexes (62) and functioning as a key regulator coordinating

eukaryotic chromatin compaction, HP1 can bind to heterochromatin marks and recruits other

factors to promote heterochromatin formation (63). *F. oxysporum* ACs are primarily composed

of facultative heterochromatin (64). The equal expansion in this protein family suggests that

establishing the ability to effectively open or close AC regions, either to exploit transcriptional

networks or to allow genome maintenance activities, likely plays a crucial role in the functional

regulation of *F. oxysporum* ACs.

2) *Post-translational modifications.* The other significantly enriched term ($p = 1.95 \times 10^{-8}$

for MRL8996 AC genes and $p = 3 \times 10^{-13}$ for Fol4287 AC genes) was protein phosphorylation

(GO:0006468), an important cellular regulatory mechanism through kinases. This result was

expected, as we previously identified a positive correlation between the total number of protein

kinases encoded in a genome and the size of the proteome of an organism (65). However, each

genome has its own distinct set of encoded proteins. For instance, the major AC-enriched kinases

in MRL8996 are Halotolerance protein 4-like (HAL4) serine/threonine kinases (which regulate

ion transporters), Calmodulin Dependent Protein Kinase Kinase 2 (which is involved in energy

balance and perhaps mobile DNA repair activity), and Cell Division Cycle (CDC)-like kinases

(which function in chromatin remodeling and DNA metabolism/repair). The kinases encoded by

the AC of Fol4287 include the second copy of a TOR kinase paralog (a top regulator that dictates

cellular stress responses) (65); choline kinase (a conserved APH superfamily member involved

in antibiotic resistance); a TOMM system kinase/cyclase fusion protein; and the serine/threonine

kinase protein kinase B (involved in stress responses and signal transduction).

In addition to phosphorylation, the other shared GO term involved in the post-

translational modification of proteins is protein ADP-ribosylation (GO:0006471) ($p = 2.99 \times$

10^{-8} for MRL8996 AC genes and $p = 5 \times 10^{-3}$ for Fol4287 AC genes). Several transporter genes,

including a chromate transporter gene (GO:0015703), genes enriched in the terms potassium ion

transport and iron ion transport, and a hydroxymethylglutaryl-CoA reductase gene

(GO:0015936), the rate-limiting enzyme for ergosterol biosynthesis, were also enriched in both

genomes (**Fig. 5C**, **Table S5**, **S6**).

Fol4287 AC genes are uniquely enriched for defense responses and signaling: Among the

best characterized Fol4287 AC genes are *SIX* effector genes, encoding *bona fide* fungal virulence

factors (66–68) that are directly involved in fungal–plant interactions. Many genes identified in

this study were reported previously. In addition, this study revealed a significant enrichment of

Fol4287 AC genes in the GO term defense response (GO:0006952) compared to the core genes

and the AC genes from MRL8896 (**Figure 6** category I, **Table S6**). The AC genes in this group

include two chitinase II genes and one patatin-like phospholipase gene involved in the

manipulation of plant resistance proteins (69).

MRL8996 AC genes are uniquely enriched for metal ion transport and catabolism:

Whereas MRL8996 ACs lack *SIX* effector genes and genes encoding plant cell wall–degrading

enzymes, unlike plant pathogenic *F. oxysporum* strains (32), other effectors are also detected.

Also, MRL8996 ACs are uniquely enriched in genes involved in the transport and catabolism of

metal ions such as copper, zinc, magnesium (GO:0006825 and GO:0030001), and mercury

(GO:0046413) **(Figure 6** category II**, Table S5)**. Nine genes are specifically involved in copper

ion transport, with seven Ctr-type copper transporter paralogs and two copper-exporting P-type

ATPase (CopA) paralogs encoding proteins containing multiple copies of the copper chaperone

(CopZ) domain. Other genes included in these categories encode a CorA-like magnesium/zinc

transporter, with multiple copies of ankyrin repeats, and a Zrt-like, Irt1-like protein (ZIP)-type

zinc ion transporter. Most of these genes are present in the genome of another clinical isolate,

NRRL32931 (25 out of 33 genes). One of these shared genes encodes a mammalian

ceruloplasmin homolog with two multicopper oxidase domains. Homologous sequences were

previously detected among a few other opportunistic fungal pathogens and bacteria isolated from

extreme environments, but not in plant pathogenic *Fusarium* species (32), suggesting that a host-

specific gene network evolved in the human pathogens. Another GO term that was exclusively

enriched in the MRL8996 ACs is regulation of ligase activity (GO:0051340).

Taken together, our results provide strong evidence that different ACs acquired by a plant

or a human pathogen perform shared functions that are essential for chromatin modifications,

transcriptional regulation, and post-translational modification of proteins. These essential

regulatory mechanisms, which are involved in environmental sensing and cellular signal

transduction, may hold key to understanding the crosstalk between the core and AC genomic

regions. However, the strain-specific AC gene repertoires are uniquely enriched for defense

responses and signaling in the plant pathogen and metal ion transport and catabolism in the

human pathogen. We propose that these unique adaptations are important for fungal survival in

different environments and hosts.

DISCUSSION

The FOSC, a group of cross-kingdom fungal pathogens, includes both plant and animal pathogens. Our *in vivo* assays using animal and plant hosts confirmed the host specificity of the keratitis strain MRL8996 and the tomato pathogen Fol4287, as corneal infection with the keratitis strain resulted in significantly more corneal ulceration, while the plant pathogen caused significantly more wilt symptoms among all inoculated tomato seedlings. Consistent with host-specific adaptations, the keratitis strain MRL8996 grew significantly better at elevated temperature, whereas the tomato pathogen Fol4287 exhibited more tolerance to osmotic and cell wall stress. As a line of fungal adaptation, a human pathogen must overcome the host defenses associated with mammalian endothermy and homeothermy, while a plant pathogen must handle different stresses (70, 71). Our findings reveal interconnected fungal responses toward biotic stress from plant and animal hosts and abiotic stress in distinct environments. The distinct phenotypes observed in our *in vivo* and *in vitro* assays, together with the identification of unique ACs in each genome, lay the foundation for dissecting cross-kingdom fungal pathogenicity using this comparative model.

The widespread occurrence of the caspofungin paradoxical effect underscores the complexity of antifungal drug responses in filamentous fungi and provides an early warning about the potential complexity in treating infectious diseases caused by *Fusarium* species (72). In the current study, we detected the caspofungin paradoxical effect, but only for the keratitis strain MRL8996. Mechanisms underlying this phenomenon likely involve specific drug–target interactions for each strain(73), fungal stress responses (74), and cellular signaling pathways (72, 75–79). Future studies will identify underlying mechanisms that may lead to strategies to overcome the paradoxical effect and for improving the efficacy of antifungal therapies.

In addition, we observed severe disease phenotypes from *Fusarium* keratitis compared to

other fungal infections. For instance, corneal infection with *Aspergillus fumigatus* requires

40,000 conidia as the inoculum, which still causes severe corneal disease (80). When we used the

same titer of *F. oxysporum* conidia, almost all animals developed corneal ulceration and

perforation within 24 h. Therefore, we lowered the inoculum to 5,000 conidia. This

hypervirulence may contribute to *Fusarium* species being the leading cause of blindness among

fungal keratitis patients (3, 5).

A unique advantage of our comparative system is the compartmentalized genomic

structures for both pathogens, allowing us to characterize two distinct sets of ACs to be which

revealed distinct transposon profiles: while DNA transposons dominate Fol4287 ACs,

retrotransposons, especially LINE transposons, are highly abundant among MRL8996 ACs.

Intriguingly, ACs from both genomes also encode proteins with shared functions, including

chromatin assembly/disassembly, protein phosphorylation, and transcriptional regulation, which

support our previous findings (65, 26) Our observation that there is shared enrichment of genes

involved in chromatin assembly and disassembly points to the importance of chromatin

remodeling regardless of host-specific functions. Identifying master regulators of the crosstalk

between ACs and core components, especially those regulating the activities of both plant and

animal pathogens, should uncover new targets for control of this cross-kingdom pathogen.

In summary, this is the first study documenting similarities and differences in the

genotypes and phenotypes of two closely related pathogens: one infecting humans and one

infecting plants. Our *in vivo* and *in vitro* assays allowed us to examine strain adaptation under

differenet environmental stress conditions feasible. In addition to host-specific virulence, we also

observed cross-kingdom virulence, as the human pathogen also colonized plant roots and the

plant pathogenic strain infected corneas, although both resulted in milder disease symptoms.

More in-depth research is needed on the molecular mechanisms underlying species divergence

and adaptation among this group of pathogens.

MATERIALS AND METHODS

Fungal strains

The genome of *F. oxysporum* Fol4287 was first sequenced in 2010 (21), and an improved

genome assembly was subsequently produced in 2018 (52). The strain is deposited in multiple

public strain repositories, including the Fungal Genetics Stock Center (FGSC 9935),

NCAUR/USDA (NRRL 34936), and CBS-KNAW (CBS 123668). The keratitis strain *F.*

oxysporum MRL8996 was originally isolated in 2006 from the cornea of a patient with the

contact lens–associated multistate outbreak fungal keratitis at Cleveland Clinic Foundation in

Ohio, USA (34). The strain is available at NCAUR/USDA (NRRL 47514). MRL8996 is grouped

with other human pathogenic isolates belonging to clade FOSC 3-a. The genome assembly for

MRL8996 was produced using the same sequencing technologies and computational strategies

used to assemble the Fol4287 genome to facilitate effective comparative analysis (32).

Fungal growth conditions

The conidia of the fungal strains were stored at −80°C in an ultra-freezer in 25–50% (v/v)

glycerol for long-term storage and propagated from stocks in liquid potato dextrose broth (PDB,

BD Difco™, USA) or potato sucrose broth (PSB: 25% [w/v] boiled potato and 0.5% [w/v]

sucrose) for at least 4 days in a shaking incubator at 28°C. The conidia were collected by

filtering the liquid fungal cultures through two layers of Miracloth (EMD Millipore). The filtrate

was centrifuged at 3700 g for 5 minutes and the spores were resuspended in sterile water to the

desired concentration. The spore titers were determined using a hemocytometer.

Mouse corneal infection

Fungal conidia were harvested from Sabouraud Dextrose Agar (SDA) medium and resuspended

in phosphate buffered saline (PBS, pH 7.0). Mice were anesthetized by intraperitoneal injection

with ketamine/xylazine. A 30-gauge needle was used to make a pocket in the corneal stroma,

after which a 33-gauge Hamilton syringe was inserted, and 5×10^3 conidia in 2 µl PBS was

injected into the corneal stroma. Corneal opacity was photographed using a high-resolution

stereo fluorescence MZFLIII microscope (Leica Micro-systems) and a Spot RT Slider KE

camera (Diagnostics Instruments). All images were captured using SpotCam software (RT Slider

KE; Diagnostics Instruments). Corneal opacity was quantified using ImageJ software. The

experiments used 5 mice per group and were repeated 4 times.

Colony-forming units (CFUs) from *Fusarium*-infected corneas

462   Infected whole eyes were homogenized in 1 mL of sterile PBS using a Retsch Mixer Mill

463   MM300 (Qiagen, Valencia, CA) at 33 Hz for 4 min. Ten-fold dilutions were made, and the

464   samples were plated on SDA plates and incubated at 34°C for 48 h. CFUs were determined by

465   direct counting.

Flow cytometry

Infected corneas were dissected, the vascularized iris was removed, and the corneas were

incubated in 500 µL collagenase type I (Sigma-Aldrich) at a titer of 82 U per cornea for 1–2 h at

37°C. Cells were resuspended in 200 µL FACS buffer containing 4 µg/cornea Fc blocking Ab

(anti-mouse CD16/32; eBioscience) on ice for 10 min, followed by incubation with anti-mouse

antibodies against CD45, Ly6G CD11b, or Ly6C, all from BioLegend. Total cells were

quantified using an ACEA Novocyte™ flow cytometer.

Phenotyping of isolates

Yeast extract peptone dextrose (YPD) plates were used as complete medium, and modified

Czapek-Dox plates were used as minimal medium. The pH levels of both media were adjusted to

pH 7.4 using 6.5% (w/v) 0.1 M citric acid and 43.5% (w/v) 0.2 M Na_2HPO_4. YPD plates

containing 0.6 M NaCl, 1 mM H_2O_2, and 1 mg/mL Congo Red were prepared for osmotic,

oxidative, and cell wall stress treatments, respectively. Each isolate (Fol4387, MRL8996, NRRL

32931, and Fo5176) was grown in PSB for 4 days at 28°C, and the conidia were filtered as

described above. Plates containing the same medium were inoculated with 2 μL of spore

suspensions at concentrations of 5×10^6, 5×10^5, or 5×10^4 conidia/mL, each in three replicates.

The plates were incubated at 28°C or 34°C for 3 days and photographed once a day. The

experiments were repeated twice.

Statistical analysis of growth rates

The growth rates of each replicate and dilution were calculated as the slope of the 3-day growth

curve. Five-way ANOVA with a linear model was performed for the following groups: strain,

temperature, medium, dilution, and replicate, followed by a multi-comparison with strains,

temperatures, and media groups to identify significant differences using MATLAB. To quantify

adaptation to different temperatures, the growth rates of cultures with an initial inoculum of $1 \times$

10^3 conidia were measured in triplicate. The values of the three replicates incubated at 34°C were

normalized to the values of the three replicates incubated at 28°C (total of nine values). A two-

way Student's *t*-test was performed between Fol4287 and MRL8996 values. Similarly, to

quantify the effects of stress treatment, the values of the three replicates of samples in medium

containing different stress components were normalized to the values of the three replicates in

control YPD medium (total of nine values), followed by a two-way Student's *t*-test between

Fol4287 and MRL8996 in MATLAB.

Caspofungin sensitivity assay

Both strains were maintained in potato dextrose agar (PDA) medium. For solid medium, 2%

(w/v) agar was added. The growth rate was determined by spotting 1×10^4 conidia in the center

of a 90-mm Petri plate containing 20 mL of solid PDA medium and incubating the plates at

28°C. The diameter was scored at 24-h intervals until 5 days (96 h) of incubation with three

biological replicates. *To assess the strains' germination kinetics*, 1×10^4 conidia of each strain

were inoculated onto glass coverslips containing 200 µL liquid PDA medium, and the samples

were incubated at 28°C. A conidiospore was counted as germinated if it possessed a germ tube,

which was readily detectable as a small protuberance on the spherical spore surface. Two

hundred conidia were counted in each experiment. Times when 50% of conidia reach the stage of

germination are recorded.

Plant infection assay

Seeds of tomato (*Solanum lycopersicum*) cv. M82 were maintained in the dark at 4°C for 3 days.

The seeds were surface sterilized in 70% (v/v) ethanol for 5 min and washed with 2.7% (w/v)

sodium hypochlorite (NaClO) for 5 min. After removing the NaClO solution, the seeds were

rinsed with sterile distilled water (SDW) three times. The seeds were then sown in pots filled

with autoclaved soil (Promix BX) and watered with SDW. The soil was gently removed from the

roots of 10-day-old tomato seedlings by rinsing them with abundant distilled water and SDW

while avoiding root tissue damage. The roots of the seedlings were inoculated by dipping them

into the respective human or plant pathogenic *Fusarium* spore suspension (10^6 spores/ml) or in

SDW as a mock infection control for 45 min before replanting the infected seedlings in soil.

After infection, all seedlings were maintained in a growth chamber at 28°C under a 14-h-

light/10-h-dark photoperiod.

Observe *F*. *oxysporum* cells within plant tissue

The staining process using wheat germ agglutinin (WGA) coupled to the fluorophore Alexa

Fluor 488 (WGA-Alexa Fluor 488) and Propidium Iodide (PI) is employed for observing fungal

colonization in plant leaves and roots following the staining method described previously (46,

81). The infected roots were observed under an Olympus FluoView FV1000 confocal

microscope (Tokyo, JP) and photographed with a Hamamatsu camera (Hamamatsu, JP). The

experiments were repeated twice.

Detect *F*. *oxysporum* from above-ground tissues

To detect Fol4287 and MRL8996 from above-ground tissues, tomato stems were harvested after

16 dpi and washed with abundant SDW. Under the laminar flow hood, stems were dipped in

10% NaClO for 1 min, washed several times with SDW, and blotted dry with Kimwipe paper

towels. Small stem pieces were placed on a PDA medium and incubated at 28°C. After 3 days,

the observed fungal growth was transferred to new Petri dishes with PDA to obtain axenic

cultures. The Qiagen plant DNA extraction kit (Hilden, Germany) was used to extract the DNA

of three different fungal colonies isolated from the tomato stems. Fol4287 strain-specific genes

FOXG_22560 and FOXG_18682 were amplificated by PCR using the primers:

Fol4287_22560_For: ATGCGCTTC   AATGTTCTCGC   and   Fol4287_22560_Rev:

ACAACAGACAGTACCAGCGG,   as   well   as   Fol4287_18682_For:

GCTGCTACGGCGATACTGTC and Fol4287_18682_Rev: GACTCGTCTGGGCTGTACTC

with 63 and 64 °C of alignment temperature respectively, 35 cycles and following the

manufacturer instructions of Phusion High–Fidelity DNA Polymerase (Thermo Scientific).

Finally, the PCR products were visualized in 1.2% of agarose gels.

Contour-clamped homogeneous electric field (CHEF) electrophoresis

The harvested conidia sample was washed with sterile water, followed by 1.2 M KCl,

resuspended in protoplasting solution (25 mg/mL driselase, 5 mg/mL lysing enzyme from

Sigma-Aldrich, and 1.2 M KCl), and incubated overnight at 30°C under shaking at 80 rpm. The

protoplasts were centrifuged at 4°C for 15 min at 1,000 g, washed by slowly resuspending in 10

mL STC buffer (1 M sorbitol, 10 mM Tris-HCl pH 7.4, and 10 mM $CaCl_2$), and centrifuged

again as above. The supernatant was carefully poured out, and the protoplasts were resuspended

in STE buffer (2% [w/v] SDS, 0.5 M Tris-HCl pH 8.1, and 10 mM EDTA) to a titer of 2×10^8

conidia/mL. The protoplasts were incubated at 50°C for 10 min and mixed at a 1:1 (v/v) ratio

with 1.2% (w/v) Bio-Rad Certified Low Melt Agarose. The mixture was transferred to CHEF

Disposable Plug Molds and stored at 4°C for long-term storage. CHEF gel electrophoresis was

performed as previously described (82). Briefly, the CHEF Mapper System (Bio-Rad) with 1%

(w/v) SeaKem® Gold Agarose (Lonza) in 0.5× Tris borate EDTA (TBE) buffer was used to

separate chromosomes at 4–7°C for 260 h. The switching time was 1,200 to 4,800 s at

1.5 V/cm, and the angle was 120°. The running buffer was changed every 2–3 days. The gels

were stained with 3× GelGreen (Biotium).

Genome alignment

The alignments of the Fol4287 and MRL8996 nuclear and mitochondrial genomes with repeats

masked were generated using MUMmer version 3.23 (83) with the option 'nucmer --maxmatch'.

Alignments less than 1 kb were removed using 'delta-filter -g -l 1000'. The extent of sequence

identity was calculated for each chromosome by multiplying the percentage sequence identity for

each alignment by the length of the alignment averaged over the total length of alignments for

the chromosome. The circular plot was generated using Circos (84). Only alignments longer than

5 kb and contigs longer than 5 kb were considered. Contigs from the same chromosome were

concatenated. A sequence alignment plot for the mitochondrial genome was generated via a

PlotMUMmerAlignments.m script (available at https://github.com/d-ayhan/tools). The

annotation of the Fol4287 genome was obtained from the 2010 assembly, while the MRL8996

genome was annotated *de novo* at JGI (85, 86). The genes in ACs were identified and their

enriched Gene Ontology (GO) terms were analyzed. For Fol4287 genes, locus tag IDs starting

with 'FOXG_' were used, while for MRL8996, the 'protein_id' numbers assigned by JGI were

used.

Repeat analysis

Repeats were *de novo* identified and classified using RepeatModeler v1.0.11 (87). A previously

curated library of 69 transposable elements (TEs) was included in the repeat sequence database.

Usearch with the option '-id 0.75' was used to cluster the sequences around centroid repeat

sequences, and the annotations were fixed manually (88). RepeatMasker version 4.0.7 was used

to mask and annotate the genome assemblies (89). TECNEstimator, a bioinformatics pipeline

used to quantify the copy numbers of repeat sequences using short reads from whole-genome

sequencing data (SRA accessions: SRP140697 and SRP214968), was used to estimate read

counts (available at https://github.com/d-ayhan/TECNEstimator). The counts were normalized to

the median read coverages of the samples. For both genomes, only one representative sequence

in a cluster with the highest copy number was selected for downstream analysis.

GO term enrichment analysis

The genes harbored by ACs were identified as those located in AC contigs of MRL8996 and AC

and unmapped contigs of Fol4287. The orthologous genes were identified using OrthoFinder

version 2.3.3 with default options (90). GO terms for the Fol4287 and MRL8996 genomes were

downloaded from JGI Mycocosm (JGI-specific genome identifiers: Fusox2 and FoxMRL8996,

respectively) (85). The GO_term_enrichment_analysis.m script (available at

https://github.com/d-ayhan/tools), which utilizes the hypergeometric cumulative distribution

function, was used to analyze the enriched GO terms in ACs. The proteins encoded by the genes

included in each term were subjected to conserved domain analysis using an InterPro-based

annotated file provided by JGI and a NCBI conserved domain database CD-Search

(https://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi).

Phylogenetic analysis

The protein sequence alignments were generated by MEGA11 (91) using the ClustalW algorithm.

The phylogenetic tree was reconstructed using FastTree (92) with default options and visualized

in iTOL (93).

Acknowledgements

The authors thank the University of Massachusetts High-Performance Computing Center for

providing high-performance computing capacities and assisting in data analysis; we also thank

Chris Joyner and David O'Neil of the College of Natural Sciences Greenhouse at the University

of Massachusetts, Amherst, for continued support with plant propagation. This project was

supported by the National Eye Institute of the National Institutes of Health (R01EY030150) and

the Natural Science Foundation (IOS-165241). L.-J.M. is also supported by an Investigator

Award in Infectious Diseases and Pathogenesis by the Burroughs Wellcome Fund BWF-

1014893; SM is also supported by a Vaadia-BARD Postdoctoral Fellowship. The work

(proposal: 10.46936/10.25585/60001360) conducted by the U.S. Department of Energy Joint

Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is

supported by the Office of Science of the U.S. Department of Energy operated under Contract

No. DE-AC02-05CH11231. The funding bodies played no role in the design of the study, the

collection, analysis, and interpretation of the data, or the writing of the manuscript.

Author contributions

L.-J.M., D.H.A., and E.P. designed this study. D.H.A. and L.-J.M. wrote the manuscript with

input from all authors. D.H.A., S.M., V.S. S.H, and S.W. performed data analysis. D.H.A., S.A.,

D.M.S, K.R., S.K., M.E.M, M.C.R, S.Y, and R.R.V. performed experiments. T.A, I.V.G, N.S,

E.P and L.-J.M. provided funding support for all experiments and analyses.

623
Figure 1. *In vivo* pathogenicity of clinical (MRL8996) and agricultural (Fol4287) *F.*

***oxysporum* strains in infected corneas.**

A. Representative mouse corneas (n = 4) 24 h after intrastromal injection of 5,000 swollen

conidia. **B.** Disease severity, as measured by the corneal opacification (94) (*** $p < 0.001$). **C.**

Viable conidia from infected corneas at t = 0 (inoculum; 5,000 conidia) and after 24 h, as

indicated by the number of colony-forming units (CFUs) (** $p < 0.01$). **D.** Infiltrating

neutrophils and monocytes identified by flow cytometry. Total cells were isolated from

collagenase-digested corneas at 24 h post-infection (hpi) with MRL8996 (top panels) or Fol4287

(bottom panels). Cells were identified using (i) CD45 for total myeloid cells (ii); CD45+ Ly6G+

and CD11b+ to identify neutrophils; and (iii) CD11b+ Ly6C+ monocytes. **E.** Quantification of

neutrophils (top) and monocytes (bottom) in multiple infected corneas at 24 hpi. Significance

was determined by a paired Student's *t*-test, where * $p < 0.05$ was considered significant.

Figure 2. *In vivo* pathogenicity assay of clinical (MRL8996) and agricultural (Fol4287) *F.*

***oxysporum* strains using the tomato vascular disease model. A.** Representative photographs of

tomato plants inoculated with Fol4287, MRL8996, or water (mock treatment). Ten-day-old

tomato seedlings were infected and imaged 10 days later. **B.** A disease index of 1–5 was used to

measure disease severity as described previously (45): 1 for healthy plants; 2 for plants showing

wilted leaves with chlorotic areas; 3 for plants with necrotic spots; 4 for plants with wilted leaves

and whole plants showing chlorosis, areas of necrosis, and defoliation; and 5 for dead plants. **C.**

Re-isolation of fungal inoculum from inoculated plant stems. The red bracket indicates the area

of the plant stem used for fungal isolation. Small slides of the stem were inoculated in Petri

dishes with PDA medium. A total of 6 plants from each treatment were used for this assay. We

isolated fungal colonies only from the stems of tomato plants inoculated with Fol4287. **D**. PCR

amplification of 100 bp performed for two specific genes (FOXG_22560 and FOXG_18682) of

Fol4287 in three different isolated colonies. **E**. Tracking the colonization of Fol4287, MRL8996,

and mock infection using confocal microscopy. Fungal hyphae (in green) were stained with

WGA-Alexa Fluor 488 and detected by excitation at 488 nm and emission at 500 to 540 nm.

Plant tissues (in red) were stained with propidium iodide and detected with excitation at 561 nm

and emission at 580–660 nm. Xylem is indicated by the dotted lines. The primary roots, the

lateral roots, and the stems are visualized top to bottom.

Figure 3. *In vitro* growth of clinical (MRL8996) and agricultural (Fol4287) *F. oxysporum*

strains subjected to abiotic stress.

A. Colony morphology of MRL8996 and Fol4287 on minimal medium (modified Czapek-Dox

agar) pH ~5, minimal medium pH 7.4, complete medium (CM, yeast extract peptone dextrose

[YPD] agar), pH ~5, complete medium pH 7.4, osmotic stress medium (potato dextrose agar

[PDA] containing 0.6 M NaCl), oxidative stress medium (PDA with 1 mM hydrogen peroxide

[H_2O_2]), and cell wall stress medium (YPD with 1 mg/mL Congo Red). The plates were

incubated at 28°C or 34°C. The images are representative of three replicates and were taken at 2

days post-inoculation **B.** Ratio of the growth rates (GRs) of colonies under the same conditions

as in **A**. at 34°C and 28°C ($GR_{34°C}/GR_{28°C}$) as a representation of temperature adaptation. **C.**

Ratio of mean rates of growth under stress conditions (GR_{stress}) and in CM (GR_{normal}) as a

representation of stress tolerance at 28°C and 34°C. Data points indicate each growth rate value

at 34°C normalized to each growth rate value at 28°C, with each bar showing the mean value. *

$p < 0.05$, ** $p < 10^{-5}$; *** $p < 10^{-9}$, calculated by a two-sample *t*-test.

Figure 4. *F. oxysporum* **keratitis strain MRL8896 exhibits a paradoxical effect to**

caspofungin.

A. Representative images of radial fungal growth in the presence of caspofungin. **B.** Ratio of

growth rates (GRs) for colonies in the presence of caspofungin (treatment) and in the absence of

caspofungin (control). **C.** Effect of caspofungin concentration on germination time. For each

strain, 1×10^4 conidia were inoculated onto glass coverslips containing 200 µL of liquid PDA

medium and incubated at 28°C for 24 h. Two hundred conidia were counted, and the percent

germination was calculated at 0, 2, 4, 6, 8, and 12 h. The time when 50% of the spores

germinated was calculated based on the closest datapoint in a log-logistic regression.

Figure 5. Compartmentalized genomes of the clinical (MRL8996) and agricultural

(Fol4287) *F. oxysporum* **strains.**

A. Contour-clamped homogeneous electric field (CHEF) gel of small accessory chromosomes

(ACs) of the agricultural isolate Fol4287, the control clinical isolate NRRL32931, and the

clinical isolate MRL8996. *Schizosaccharomyces pombe* (left) and *Hansenula wingei* (right) were

used as markers. The band sizes are shown in Mb. ACs are typically below 2 Mb in size, except

for two known ACs, chr3 and chr6 in Fol4287, which are much larger due to recent segmental

duplications (21). **B.** Whole-genome comparison between MRL8996 and Fol4287 revealed 11

homologous core chromosomes (light green and light blue, respectively) and accessory

sequences (dark green and dark blue, respectively) (i). The accessory sequences typically

displayed low gene density (ii) and high repetitive sequence composition (iii). The syntenic

alignment between MRL8996 and Fol4287 using nucmer (iv) indicates the accessory regions

lacking synteny between MRL8996 and Fol4287. **C.** Distribution of all identified transposable

elements in the Fol4287 (upper panel) and MRL8996 (lower panel) genomes, consisting of class

I retrotransposons, class II DNA transposons, and unknown transposable elements. **D.**

Transposon abundance in the Fol4287 and MRL8996 genomes. Class I retrotransposons include

LTRs (long terminal repeats), LINEs (long interspersed nuclear elements), and SINEs (short

interspersed nuclear elements). Class II DNA transposons are classified into TIRs (terminal

inverted repeats) and Helitrons.

Figure 6. Enriched Gene Ontology terms among accessory genes of the clinical (MRL8996)

and agricultural (Fol4287) *F. oxysporum* **isolates.**

Enriched Gene Ontology (GO) terms harbored by the ACs from Fol4287 (green) and MRL8996

(blue). Category I represents GO terms enriched in Fol4287 ACs. Category II represents GO

terms enriched in the ACs of both genomes. Category III represents GO terms enriched in

MRL8996 ACs. For graphical representation, a p-value $= 0$ is shown as $-\log10(p\text{-value}) = 30$.

703   **Table 1. Comparative annotation of the Fol4287 and MRL8996 genomes.**

	Fol4287	MRL8996
Total genes	20,925	16,631
Genes with GO annotation	9,783	7,958
Total AC genes	3,890	2,017
AC genes with GO annotation	1,788	542

REFERENCES

1.   World Health Organization. 2022. WHO fungal priority pathogens list to guide research,

development and public health action. https://www.who.int/publications-detail-

redirect/9789240060241.

2.   Dean R, Van Kan JA, Pretorius ZA, Hammond-Kosack KE, Di Pietro A, Spanu PD, Rudd JJ,

Dickman M, Kahmann R, Ellis J, Foster GD. 2012. The Top 10 fungal pathogens in molecular plant

pathology. Mol Plant Pathol, 2012/04/05 ed. 13:414–430.

3.   Kredics L, Narendran V, Shobana CS, Vágvölgyi C, Manikandan P, Indo-Hungarian Fungal Keratitis

Working Group. 2015. Filamentous fungal infections of the cornea: a global overview of

epidemiology and drug sensitivity. Mycoses 58:243–260.

4.   Abbondante S, Leal SM, Clark HL, Ratitong B, Sun Y, Ma L-J, Pearlman E. 2023. Immunity to

pathogenic fungi in the eye. Seminars in Immunology 67:101753.

5.   Hassan AS, Al-Hatmi AMS, Shobana CS, van Diepeningen AD, Kredics L, Vágvölgyi C, Homa M,

Meis JF, de Hoog GS, Narendran V, Manikandan P, IHFK Working Group. 2016. Antifungal

Susceptibility and Phylogeny of Opportunistic Members of the Genus Fusarium Causing Human

Keratomycosis in South India. Med Mycol 54:287–294.

6.   Lalitha P, Prajna NV, Manoharan G, Srinivasan M, Mascarenhas J, Das M, D'Silva SS, Porco TC,

Keenan JD. 2015. Trends in bacterial and fungal keratitis in South India, 2002-2012. Br J

Ophthalmol 99:192–194.

7.   He D, Hao J, Zhang B, Yang Y, Song W, Zhang Y, Yokoyama K, Wang L. 2011. Pathogenic

spectrum of fungal keratitis and specific identification of Fusarium solani. Invest Ophthalmol Vis Sci

52:2804–2808.

8.   Wang L, Sun S, Jing Y, Han L, Zhang H, Yue J. 2009. Spectrum of fungal keratitis in central China.

Clin Exp Ophthalmol 37:763–771.

9.   O'Sullivan J, Gilbert C, Foster A. 1997. The causes of childhood blindness in South Africa. S Afr

Med J 87:1691–1695.

10. Ibrahim MM, Vanini R, Ibrahim FM, Fioriti LS, Furlan EMR, Provinzano LMA, De Castro RS, Sousa

SJDFE, Rocha EM. 2009. Epidemiologic aspects and clinical outcome of fungal keratitis in

southeastern Brazil. Eur J Ophthalmol 19:355–361.

11. O'Donnell K, Sarver BA, Brandt M, Chang DC, Noble-Wang J, Park BJ, Sutton DA, Benjamin L,

Lindsley M, Padhye A, Geiser DM, Ward TJ. 2007. Phylogenetic diversity and microsphere array-

based genotyping of human pathogenic Fusaria, including isolates from the multistate contact lens-

associated U.S. keratitis outbreaks of 2005 and 2006. J Clin Microbiol, 2007/05/18 ed. 45:2235–

2248.

12. Gower EW, Keay LJ, Oechsler RA, Iovieno A, Alfonso EC, Jones DB, Colby K, Tuli SS, Patel SR,

Lee SM, Irvine J, Stulting RD, Mauger TF, Schein OD. 2010. Trends in fungal keratitis in the United

States, 2001 to 2007. Ophthalmology 117:2263–2267.

13. Michielse CB, Rep M. 2009. Pathogen profile update: Fusarium oxysporum. Mol Plant Pathol,

2009/04/30 ed. 10:311–324.

14. Berger S, Chazli YE, Babu AF, Coste AT. 2017. Azole resistance in Aspergillus fumigatus: A

consequence of antifungal use in agriculture? Frontiers in Microbiology.

15. Pfaller MA. 2012. Antifungal drug resistance: mechanisms, epidemiology, and consequences for

treatment. Am J Med 125:S3-13.

16. Altus, Viljoen, Li-Jun, Ma, Augusta, Molina. 2020. Fusarium Wilt (Panama Disease) and

Monoculture in Banana Production: Resurgence of a Century-Old Disease, p. 159–184. *In*

Emerging plant diseases and global food security. A. Records & J. Ristaino, APS Press.

17. Scheel CM, Hurst SF, Barreiros G, Akiti T, Nucci M, Balajee SA. 2013. Molecular analyses of

Fusarium isolates recovered from a cluster of invasive mold infections in a Brazilian hospital. BMC

Infect Dis 13:49.

18.   Nucci M, Marr KA, Vehreschild MJGT, De Souza CA, Velasco E, Cappellano P, Carlesse F,

755   Queiroz-Telles F, Sheppard DC, Kindo A, Cesaro S, Hamerschlak N, Solza C, Heinz WJ, Schaller

756   M, Atalla A, Arikan-Akdagli S, Bertz H, Galvão Castro C, Herbrecht R, Hoenigl M, Härter G,

757   Hermansen NEU, Josting A, Pagano L, Salles MJC, Mossad SB, Ogunc D, Pasqualotto AC, Araujo

758   V, Troke PF, Lortholary O, Comely OA, Anaissie E. 2014. Improvement in the outcome of invasive

759   fusariosis in the last decade. Clinical Microbiology and Infection 20:580–585.

760   19.   Prajna NV, Krishnan T, Rajaraman R, Patel S, Srinivasan M, Das M, Ray KJ, O'Brien KS,

761   Oldenburg CE, McLeod SD, Zegans ME, Porco TC, Acharya NR, Lietman TM, Rose-Nussbaumer J,

762   for the Mycotic Ulcer Treatment Trial II Group. 2016. Effect of Oral Voriconazole on Fungal Keratitis

763   in the Mycotic Ulcer Treatment Trial II (MUTT II): A Randomized Clinical Trial. JAMA Ophthalmol

764   134:1365.

765   20.   Kistler H. 2001. Evolution of host specificity in Fusarium oxysporum, p. 70–82. *In* Nelson, P,

766   Summerell, B (eds.), Fusarium: Paul E. Nelson Memorial Symposium. APS Press.

767   21.   Ma LJ, van der Does HC, Borkovich KA, Coleman JJ, Daboussi MJ, Di Pietro A, Dufresne M,

768   Freitag M, Grabherr M, Henrissat B, Houterman PM, Kang S, Shim WB, Woloshuk C, Xie X, Xu JR,

769   Antoniw J, Baker SE, Bluhm BH, Breakspear A, Brown DW, Butchko RA, Chapman S, Coulson R,

770   Coutinho PM, Danchin EG, Diener A, Gale LR, Gardiner DM, Goff S, Hammond-Kosack KE, Hilburn

771   K, Hua-Van A, Jonkers W, Kazan K, Kodira CD, Koehrsen M, Kumar L, Lee YH, Li L, Manners JM,

772   Miranda-Saavedra D, Mukherjee M, Park G, Park J, Park SY, Proctor RH, Regev A, Ruiz-Roldan

773   MC, Sain D, Sakthikumar S, Sykes S, Schwartz DC, Turgeon BG, Wapinski I, Yoder O, Young S,

774   Zeng Q, Zhou S, Galagan J, Cuomo CA, Kistler HC, Rep M. 2010. Comparative genomics reveals

775   mobile pathogenicity chromosomes in Fusarium. Nature, 2010/03/20 ed. 464:367–373.

776   22.   Ma L-J, Geiser DM, Proctor RH, Rooney AP, O'Donnell K, Trail F, Gardiner DM, Manners JM,

777   Kazan K. 2013. Fusarium Pathogenomics. Annu Rev Microbiol 67:399–416.

778   23.   Van Dam P, Fokkens L, Schmidt SM, Linmans JHJ, Kistler HC, Ma L-J, Rep M. 2016. Effector

779   profiles distinguish *formae speciales* of *Fusarium oxysporum*: Effector profiles distinguish *Formae*

780   *speciales* of Fo. Environ Microbiol 18:4087–4102.

781   24.   Yu H, Ayhan DH, Martínez-Soto D, Cochavi SM, Ma L-J. 2023. Accessory Chromosomes of the

782   Fusarium oxysporum Species Complex and Their Contribution to Host Niche Adaptation, p. 371–

783   388. *In* Scott, B, Mesarich, C (eds.), Plant Relationships. Springer International Publishing, Cham.

784   25.   Rep M. 2005. Small proteins of plant-pathogenic fungi secreted during host colonization. FEMS

785   Microbiology Letters 253:19–27.

786   26.   Yu H, Yang H, Haridas S, Hayes RD, Lynch H, Andersen S, Newman M, Li G, Martínez-Soto D,

787   Milo-Cochavi S, Hazal Ayhan D, Zhang Y, Grigoriev IV, Ma L-J. 2023. Conservation and Expansion

788   of Transcriptional Factor Repertoire in the Fusarium oxysporum Species Complex. JoF 9:359.

789   27.   Turrà D, Segorbe D, Di Pietro A. 2014. Protein Kinases in Plant-Pathogenic Fungi: Conserved

790   Regulators of Infection. Annu Rev Phytopathol 52:267–288.

791   28.   López-Berges MS, Rispail N, Prados-Rosales RC, Di Pietro A. 2010. A Nitrogen Response Pathway

792   Regulates Virulence Functions in *Fusarium oxysporum* via the Protein Kinase TOR and the bZIP

793   Protein MeaB. The Plant Cell 22:2459–2475.

794   29.   Kistler HC, Rep M, Ma L-J. 2013. Structural dynamics of Fusarium genomes In: Fusarium:

795   genomics, molecular and cellular biology. Eds. Brown, D.W. and Proctor, R.H., Horizon Scientific

796   Press, Norwich, United Kingdom.

797   30.   Van Dam P, Fokkens L, Ayukawa Y, Van Der Gragt M, Ter Horst A, Brankovics B, Houterman PM,

798   Arie T, Rep M. 2017. A mobile pathogenicity chromosome in Fusarium oxysporum for infection of

799   multiple cucurbit species. Scientific Reports 7.

31.   Schäfer K, Di Pietro A, Gow NAR, MacCallum D. 2014. Murine Model for Fusarium oxysporum

Invasive Fusariosis Reveals Organ-Specific Structures for Dissemination and Long-Term

Persistence. PLoS ONE 9:e89920.

32.   Zhang Y, Yang H, Turra D, Zhou S, Ayhan DH, DeIulio GA, Guo L, Broz K, Wiederhold N, Coleman

JJ, Donnell KO, Youngster I, McAdam AJ, Savinov S, Shea T, Young S, Zeng Q, Rep M, Pearlman

E, Schwartz DC, Di Pietro A, Kistler HC, Ma L-J. 2020. The genome of opportunistic fungal

pathogen Fusarium oxysporum carries a unique set of lineage-specific chromosomes. 1. Commun

Biol 3:1–12.

33.   Schmidt SM, Houterman PM, Schreiver I, Ma L, Amyotte S, Chellappan B, Boeren S, Takken FLW,

Rep M. 2013. MITEs in the promoters of effector genes allow prediction of novel virulence genes in

Fusarium oxysporum. BMC Genomics 14:119.

34.   Chang DC, Grant GB, O'Donnell K, Wannemuehler KA, Noble-Wang J, Rao CY, Jacobson LM,

Crowell CS, Sneed RS, Lewis FMT, Schaffzin JK, Kainer MA, Genese CA, Alfonso EC, Jones DB,

Srinivasan A, Fridkin SK, Park BJ, Fusarium Keratitis Investigation Team. 2006. Multistate outbreak

of Fusarium keratitis associated with use of a contact lens solution. JAMA 296:953–963.

35.   Zhang Y, Yang H, Turra D, Zhou S, Ayhan DH, DeIulio GA, Guo L, Broz K, Wiederhold N, Coleman

JJ, Donnell KO, Youngster I, McAdam AJ, Savinov S, Shea T, Young S, Zeng Q, Rep M, Pearlman

E, Schwartz DC, Di Pietro A, Kistler HC, Ma L-J. 2020. The genome of opportunistic fungal

pathogen Fusarium oxysporum carries a unique set of lineage-specific chromosomes. Commun Biol

3:50.

36.   Leal SM, Vareechon C, Cowden S, Cobb BA, Latgé J-P, Momany M, Pearlman E. 2012. Fungal

antioxidant pathways promote survival against neutrophils during infection. J Clin Invest 122:2482–

2498.

37.   O'Donnell K, Sutton DA, Rinaldi MG, Magnon KC, Cox PA, Revankar SG, Sanche S, Geiser DM,

Juba JH, Van Burik J-AH, Padhye A, Anaissie EJ, Francesconi A, Walsh TJ, Robinson JS. 2004.

Genetic Diversity of Human Pathogenic Members of the *Fusarium oxysporum* Complex Inferred from Multilocus DNA Sequence Data and Amplified Fragment Length Polymorphism Analyses: Evidence for the Recent Dispersion of a Geographically Widespread Clonal Lineage and Nosocomial Origin. J Clin Microbiol 42:5109–5120.

38. O'Donnell K, Sarver BAJ, Brandt M, Chang DC, Noble-Wang J, Park BJ, Sutton DA, Benjamin L, Lindsley M, Padhye A, Geiser DM, Ward TJ. 2007. Phylogenetic Diversity and Microsphere Array-Based Genotyping of Human Pathogenic Fusaria, Including Isolates from the Multistate Contact Lens-Associated U.S. Keratitis Outbreaks of 2005 and 2006. J Clin Microbiol 45:2235–2248.

39. Di Pietro A, Roncero MI. 1996. Purification and characterization of an exo-polygalacturonase from the tomato vascular wilt pathogen Fusarium oxysporum f.sp. lycopersici. FEMS Microbiol Lett 145:295–299.

40. O'Donnell K, Sutton DA, Rinaldi MG, Magnon KC, Cox PA, Revankar SG, Sanche S, Geiser DM, Juba JH, van Burik J-AH, Padhye A, Anaissie EJ, Francesconi A, Walsh TJ, Robinson JS. 2004. Genetic Diversity of Human Pathogenic Members of the Fusarium oxysporum Complex Inferred from Multilocus DNA Sequence Data and Amplified Fragment Length Polymorphism Analyses: Evidence for the Recent Dispersion of a Geographically Widespread Clonal Lineage and Nosocomial Origin. J Clin Microbiol 42:5109–5120.

41. Christensen JE, Andreasen SØ, Christensen JP, Thomsen AR. 2001. CD11b expression as a marker to distinguish between recently activated effector CD8+ T cells and memory cells. International Immunology 13:593–600.

42. Di Pietro A, Roncero MI. 1998. Cloning, expression, and role in pathogenicity of pg1 encoding the major extracellular endopolygalacturonase of the vascular wilt pathogen Fusarium oxysporum. Mol Plant Microbe Interact, 1998/02/05 ed. 11:91–98.

848   43.   Guo L, Yu H, Wang B, Vescio K, DeIulio GA, Yang H, Berg A, Zhang L, Edel-Hermann V, Steinberg

849   C, Kistler HC, Ma L-J. 2021. Metatranscriptomic comparison of endophytic and pathogenic

850   *Fusarium* –Arabidopsis interactions reveals plant transcriptional plasticity. preprint. Plant Biology.

851   44.   Martínez-Soto D, Yu H, Allen KS, Ma L-J. 2023. Differential Colonization of the Plant Vasculature

852   Between Endophytic Versus Pathogenic *Fusarium oxysporum* Strains. MPMI MPMI-08-22-0166-SC.

853   45.   García-Maceira FI, Di Pietro A, Roncero MIG. 2000. Cloning and Disruption of *pgx4* Encoding an In

854   Planta Expressed Exopolygalacturonase from *Fusarium oxysporum*. MPMI 13:359–365.

855   46.   Ghareeb H, Becker A, Iven T, Feussner I, Schirawski J. 2011. Sporisorium reilianum Infection

856   Changes Inflorescence and Branching Architectures of Maize. Plant Physiology 156:2037–2052.

857   47.   Kessel L, Johnson L, Arvidsson H, Larsen M. 2010. The Relationship between Body and Ambient

858   Temperature and Corneal Temperature. Investigative Opthalmology & Visual Science 51:6593.

859   48.   Pfaller MA, Diekema DJ, Messer SA, Hollis RJ, Jones RN. 2003. In Vitro Activities of Caspofungin

860   Compared with Those of Fluconazole and Itraconazole against 3,959 Clinical Isolates of *Candida*

861   spp., Including 157 Fluconazole-Resistant Isolates. Antimicrob Agents Chemother 47:1068–1071.

862   49.   Bowman JC, Hicks PS, Kurtz MB, Rosen H, Schmatz DM, Liberator PA, Douglas CM. 2002. The

863   Antifungal Echinocandin Caspofungin Acetate Kills Growing Cells of *Aspergillus fumigatus* In Vitro.

864   Antimicrob Agents Chemother 46:3001–3012.

865   50.   Ha Y, Covert SF, Momany M. 2006. FsFKS1, the 1,3-β-Glucan Synthase from the Caspofungin-

866   Resistant Fungus *Fusarium solani*. Eukaryot Cell 5:1036–1042.

867   51.   Pfaller MA, Marco F, Messer SA, Jones RN. 1998. In Vitro Activity of Two Echinocandin Derivatives,

868   LY303366 and MK-0991 (L-743,792), Against Clinical Isolates of Aspergillus, Fusarium, Rhizopus,

869   and Other Filamentous Fungi. Diagnostic Microbiology and Infectious Disease 30:251–255.

52. Ayhan DH, López-Díaz C, Di Pietro A, Ma L-J. 2018. Improved Assembly of Reference Genome

Fusarium oxysporum f. sp. lycopersici Strain Fol4287. Microbiology Resource Announcements

https://doi.org/10.1128/mra.00910-18.

53. Fokkens L, Shahi S, Connolly LR, Stam R, Schmidt SM, Smith KM, Freitag M, Rep M. 2018. The

multi-speed genome of *Fusarium oxysporum* reveals association of histone modifications with

sequence divergence and footprints of past horizontal chromosome transfer events. preprint.

Genomics.

54. Brankovics B, Van Dam P, Rep M, De Hoog GS, J. Van Der Lee TA, Waalwijk C, Van Diepeningen

AD. 2017. Mitochondrial genomes reveal recombination in the presumed asexual Fusarium

oxysporum species complex. BMC Genomics 18:735.

55. Yang H, Yu H, Ma L-J. 2020. Accessory Chromosomes in *Fusarium oxysporum*. Phytopathology®

110:1488–1496.

56. Chellapan BV, Van Dam P, Rep M, Cornelissen BJC, Fokkens L. 2016. Non-canonical Helitrons in

Fusarium oxysporum. Mobile DNA 7:27.

57. Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A, Zhao X,

Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2014. MycoCosm portal: gearing

up for 1000 fungal genomes. Nucl Acids Res 42:D699–D704.

58. Lachner M, O'Carroll D, Rea S, Mechtler K, Jenuwein T. 2001. Methylation of histone H3 lysine 9

creates a binding site for HP1 proteins. Nature 410:116–120.

59. Nakayama J, Rice JC, Strahl BD, Allis CD, Grewal SIS. 2001. Role of Histone H3 Lysine 9

Methylation in Epigenetic Control of Heterochromatin Assembly. Science 292:110–113.

60. Mondal T, Rasmussen M, Pandey GK, Isaksson A, Kanduri C. 2010. Characterization of the RNA

content of chromatin. Genome Res 20:899–907.

61.   Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, Yuan D, Stroe O, Wood G,

894   Laydon A, Žídek A, Green T, Tunyasuvunakool K, Petersen S, Jumper J, Clancy E, Green R, Vora

895   A, Lutfi M, Figurnov M, Cowie A, Hobbs N, Kohli P, Kleywegt G, Birney E, Hassabis D, Velankar S.

896   2022. AlphaFold Protein Structure Database: massively expanding the structural coverage of

897   protein-sequence space with high-accuracy models. Nucleic Acids Research 50:D439–D444.

898   62.   Leopold K, Stirpe A, Schalch T. 2019. Transcriptional gene silencing requires dedicated interaction

899   between HP1 protein Chp2 and chromatin remodeler Mit1. Genes Dev 33:565–577.

900   63.   Schoelz JM, Riddle NC. 2022. Functions of HP1 proteins in transcriptional regulation. Epigenetics &

901   Chromatin 15:14.

902   64.   Fokkens L, Shahi S, Connolly LR, Stam R, Schmidt SM, Smith KM, Freitag M, Rep M. 2018. The

903   multi-speed genome of *Fusarium oxysporum* reveals association of histone modifications with

904   sequence divergence and footprints of past horizontal chromosome transfer events. preprint.

905   Genomics.

906   65.   DeIulio G, Guo L, Zhang Y, Goldberg J, Kistler H, Ma L-J. 2018. Kinome expansion in the Fusarium

907   oxysporum species complex driven by accessary chromosomes. mSphere mSphere 3:e00231-18.

908   66.   van der Does HC, Duyvesteijn RG, Goltstein PM, van Schie CC, Manders EM, Cornelissen BJ, Rep

909   M. 2008. Expression of effector gene SIX1 of Fusarium oxysporum requires living plant cells.

910   Fungal Genet Biol, 2008/07/09 ed. 45:1257–1264.

911   67.   Lievens B, Houterman PM, Rep M. 2009. Effector gene screening allows unambiguous identification

912   of Fusarium oxysporum f. sp. lycopersici races and discrimination from other formae speciales.

913   FEMS Microbiol Lett, 2009/10/06 ed. 300:201–215.

914   68.   van Dam P, Fokkens L, Schmidt SM, Linmans JH, Kistler HC, Ma LJ, Rep M. 2016. Effector profiles

915   distinguish formae speciales of Fusarium oxysporum. Environ Microbiol

916   https://doi.org/10.1111/1462-2920.13445.

69.   Houterman PM, Ma L, van Ooijen G, de Vroomen MJ, Cornelissen BJ, Takken FL, Rep M. 2009.

The effector protein Avr2 of the xylem colonizing fungus Fusarium oxysporum activates the tomato

resistance protein I-2 intracellularly. Plant J, 2009/02/21 ed. 58:970–978.

70.   Lamb C, Dixon RA. 1997. THE OXIDATIVE BURST IN PLANT DISEASE RESISTANCE. Annu Rev

Plant Physiol Plant Mol Biol 48:251–275.

71.   Segal LM, Wilson RA. 2018. Reactive oxygen species metabolism and plant-fungal interactions.

Fungal Genet Biol 110:1–9.

72.   Valero C, Colabardini AC, De Castro PA, Amich J, Bromley MJ, Goldman GH. 2022. The

Caspofungin Paradoxical Effect is a Tolerant "Eagle Effect" in the Filamentous Fungal Pathogen

Aspergillus fumigatus. mBio 13:e00447-22.

73.   Sumiyoshi M, Miyazaki T, Makau JN, Mizuta S, Tanaka Y, Ishikawa T, Makimura K, Hirayama T,

Takazono T, Saijo T, Yamaguchi H, Shimamura S, Yamamoto K, Imamura Y, Sakamoto N, Obase

Y, Izumikawa K, Yanagihara K, Kohno S, Mukae H. 2020. Novel and potent antimicrobial effects of

caspofungin on drug-resistant Candida and bacteria. Sci Rep 10:17745.

74.   Day AM, Quinn J. 2019. Stress-Activated Protein Kinases in Human Fungal Pathogens. Front Cell

Infect Microbiol 9:261.

75.   García R, Bravo E, Diez-Muñiz S, Nombela C, Rodríguez-Peña JM, Arroyo J. 2017. A novel

connection between the Cell Wall Integrity and the PKA pathways regulates cell wall stress

response in yeast. Sci Rep 7:5703.

76.   Iyer KR, Robbins N, Cowen LE. 2022. The role of Candida albicans stress response pathways in

antifungal tolerance and resistance. iScience 25:103953.

77.   Pianalto KM, Billmyre RB, Telzrow CL, Alspaugh JA. 2019. Roles for Stress Response and Cell

Wall Biosynthesis Pathways in Caspofungin Tolerance in *Cryptococcus neoformans*. Genetics

213:213–227.

78.   Ries LNA, Rocha MC, De Castro PA, Silva-Rocha R, Silva RN, Freitas FZ, De Assis LJ, Bertolini

MC, Malavazi I, Goldman GH. 2017. The Aspergillus fumigatus CrzA Transcription Factor Activates

Chitin Synthase Gene Expression during the Caspofungin Paradoxical Effect. mBio 8:e00705-17.

79.   De Castro PA, Colabardini AC, Manfiolli AO, Chiaratto J, Silva LP, Mattos EC, Palmisano G,

Almeida F, Persinoti GF, Ries LNA, Mellado L, Rocha MC, Bromley M, Silva RN, De Souza GS,

Loures FV, Malavazi I, Brown NA, Goldman GH. 2019. Aspergillus fumigatus calcium-responsive

transcription factors regulate cell wall architecture promoting stress tolerance, virulence and

caspofungin resistance. PLoS Genet 15:e1008551.

80.   Leal SM, Roy S, Vareechon C, Carrion S deJesus, Clark H, Lopez-Berges MS, diPietro A, Schrettl

M, Beckmann N, Redl B, Haas H, Pearlman E. 2013. Targeting Iron Acquisition Blocks Infection

with the Fungal Pathogens Aspergillus fumigatus and Fusarium oxysporum. PLoS Pathog

9:e1003436.

81.   Martínez-Soto D, Yu H, Allen KS, Ma L-J. 2023. Differential Colonization of the Plant Vasculature

Between Endophytic Versus Pathogenic *Fusarium oxysporum* Strains. MPMI 36:4–13.

82.   Ayukawa Y, Asai S, Gan P, Tsushima A, Ichihashi Y, Shibata A, Komatsu K, Houterman PM, Rep M,

Shirasu K, Arie T. 2021. A pair of effectors encoded on a conditionally dispensable chromosome of

Fusarium oxysporum suppress host-specific immunity. Commun Biol 4:1–12.

83.   Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile

and open software for comparing large genomes. Genome biology 5:R12.

84.   Krzywinski M, Schein J, Birol İ, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA. 2009.

Circos: An information aesthetic for comparative genomics. Genome Res 19:1639–1645.

85.   Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, Riley R, Salamov A, Zhao X,

Korzeniewski F, Smirnova T, Nordberg H, Dubchak I, Shabalov I. 2014. MycoCosm portal: gearing

up for 1000 fungal genomes. Nucleic Acids Research 42:D699–D704.

86.   Grigoriev IV, Cullen D, Goodwin SB, Hibbett D, Jeffries TW, Kubicek CP, Kuske C, Magnuson JK,

966   Martin F, Spatafora JW, Tsang A, Baker SE. 2011. Fueling the future with fungal genomics. null

967   2:192–209.

968   87.   Smit A, Hubley R. 2015. RepeatModeler Open-1.0<http://www.repeatmasker.org>.

969   88.   Edgar R. 2010. Usearch. Lawrence Berkeley National Lab. (LBNL), Berkeley, CA (United States).

970   89.   Smit A, Hubley R, Green P. 2015. RepeatMasker Open-4.0<http://www.repeatmasker.org>.

971   http://www.repeatmasker.org.

972   90.   Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics.

973   Genome Biology 20:238.

974   91.   Tamura K, Stecher G, Kumar S. 2021. MEGA11: Molecular Evolutionary Genetics Analysis Version

975   11. Molecular Biology and Evolution 38:3022–3027.

976   92.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2 – Approximately Maximum-Likelihood Trees for

977   Large Alignments. PLoS ONE 5:e9490.

978   93.   Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

979   display and annotation. Nucleic Acids Research 49:W293–W296.

980   94.   Beisel KW, Hazlett LD, Berk RS. 1983. Dominant Susceptibility Effect on the Murine Corneal

981   Response to Pseudomonas aeruginosa. Experimental Biology and Medicine 172:488–491.

A

Fol4287　　MRL8996　　Mock

B

Disease index

Mock　MRL8996　Fol4287

****

C

Isolated colonies

Mock

Fol4287

MRL8996

Number of colonies

Mock　MRL8996　Fol4287

****

D

FOXG_22560

100bp　C1　C2　C3

FOXG_18682

100bp　C1　C2　C3

E

Fol4287　　MRL8996　　Mock

Primary root

Lateral root

Stem

A

[Caspofungin] (μg/ml)

	Control	0.2	0.5	1	2	4	8

MRL8996

Fol4287

B

MRL8996 Fol4287

$GR_{treatment}/GR_{control}$

[Caspofungin] (μg/ml)

C

50% germination time (h)

[Caspofungin] (μg/ml)