Structure-activity relationships of coumarin and its analogs and mechanistic insights into

germination inhibition by coumarin

Kazuma Fukuda[1], Sota Hyakutake[1], Taiga Oishi[1], Michinari Yoshida[1], Mizuho Koga[1], Chisato

Egami[1], Hyuga Matsuura[1], Ryusei Ito[1], Kosei Tsukahara[1], Mitsuki Noda[1], Kosei Tomonaga[1], Kohei

Chifu[1], Rinsei Ichimaru[1], Eita Fujito[1], Kanae Mori[2], Yuka Tokuyama[2], Takako Yoshida[2], Noriko

Ryuda[2,*], Yukio Nagano[2] & Kazuhide Matsutaka[1,*]

[1]*Saga Prefectural Chienkan High School, Saga, Japan*

[2]*Analytical Research Center for Experimental Sciences, Saga University, Saga, Japan.*

*Corresponding authors:

E-mail: si3264@cc.saga-u.ac.jp

E-mail: matsutaka-kazuhide-a@education.saga.jp

Abstract

Coumarin is produced by various land plants, such as cherry trees, and is known to exhibit a wide range of physiological activities. Notably, one of its primary functions is the inhibition of seed germination. However, the structural basis underlying coumarin's germination-inhibitory effect remains poorly understood. In this study, we analyzed the structure-activity relationships of coumarin analogs in two plant species, white clover and Arabidopsis, and demonstrated that the benzene ring and ester bond within the coumarin skeleton constitute the minimal structural requirements for germination inhibition. The inhibitory effects of dihydrocoumarin, which lacks the double bond in the coumarin skeleton, varied among plant species, while those of various other modified coumarin analogs also differed across species. In both plant species, coumarin consistently proved to be one of the most potent inhibitors. Furthermore, the observation that dihydrocoumarin exhibited a strong germination inhibitory effect only in white clover suggests that certain coumarin analogs may act in a species-specific manner. Additionally, the mechanism of coumarin-mediated germination inhibition remains largely unclear. Using RNA-Seq analysis on Arabidopsis seeds, we uncovered a novel and critically important mechanism whereby coumarin suppresses the expression of genes encoding expansins, proteins that are responsible for loosening plant cell walls. This study is significant for understanding how coumarin contributes to plant competition in nature.

32   **Keywords:** Natural inhibitors, Coumarin structure, Cheminformatics, RNA-Seq, Cell wall loosening,

33   Expansin

Introduction

Coumarin, classified as a benzopyrone, is a heterocyclic compound composed of a benzene ring fused with a 2-pyrone ring that is produced by land plants[1,2]. In Japan, traditional products such as "sakurayu" (cherry blossom tea) and "sakuramochi"—a sweet made with mochi (Japanese rice cake) and preserved cherry leaves—derive their characteristic aroma primarily from coumarin[3]. Thus, coumarin is deeply intertwined with Japanese cultural heritage.

Coumarin is a versatile compound with a wide range of functions. In plants, it acts as an allelochemical by inhibiting the germination and root elongation of neighboring vegetation[4]. In this study, we focus on one of its primary roles: the inhibition of seed germination. The minimal structural elements required for coumarin-mediated germination inhibition remain unclear—that is, it is not known which parts of the coumarin skeleton are critical for this effect.

In addition, studies on the structure–activity relationships of coumarin-modified compounds are still limited. Numerous investigations have explored how modifications to coumarin affect seed germination. For example, using compounds such as 6-nitrocoumarin and 3-chlorocoumarin, Mayer et al. conducted a comparative analysis and found that structural modifications do not completely abolish coumarin's germination inhibitory effect but only partially attenuate it[5]. This observation suggests that the ability to inhibit germination is conserved among coumarin analogs. Moreover,

various coumarin analogs—including 4-hydroxycoumarin, 7-hydroxycoumarin (umbelliferone)[6],

4-methylumbelliferone[7], scopoletin (6-methoxy-7-hydroxycoumarin), Ayapin (presumed to be

6-acetoxy-7-hydroxycoumarin), and psoralen (a type of furanocoumarin)[8]—have been reported to

exhibit germination inhibitory effects in different plant species. However, the relationship between

molecular descriptors (such as molecular volume, molecular surface area, the lipophilicity index

[LogP], and topological polar surface area [TPSA]) and the germination inhibitory effects of these

compounds has not been thoroughly explored, highlighting the need for further research.

A critical aspect of understanding how coumarin influences seed germination is its effect on cellular

processes. Several studies have investigated this mechanism, with the work by Chen et al. being

particularly notable[9,10]. They demonstrated that coumarin negatively regulates the expression of

abscisic acid (ABA) metabolic enzyme genes (OsABA8'ox2/3) and reduces levels of gibberellin A4

(GA4). Furthermore, coumarin suppresses the accumulation of reactive oxygen species (ROS) and

the expression of key metabolic enzymes such as superoxide dismutase and catalase, thereby

providing physiological evidence for the delayed germination observed upon coumarin treatment.

Omics analyses have significantly contributed to elucidating the mechanisms underlying germination

inhibition[11,12]. For instance, Araniti et al. performed metabolomic analyses, revealing that coumarin

extensively alters the seed's metabolic profile by impacting amino acid metabolism, the citric acid

(TCA) cycle, and aminoacyl-tRNA biosynthesis[11]. A similar study by Zhang et al. yielded consistent

findings. Additionally, Zhang et al. conducted transcriptomic analyses 72 hours after coumarin

treatment, demonstrating that coumarin affects the expression of genes involved in amino acid

metabolism, the TCA cycle, and protein translation[12]. Although transcriptomic analysis has been

used to investigate the biosynthetic pathway of coumarin[13,14,15,16], this study remains the only one to

examine its germination inhibitory effect, rendering it invaluable. However, because these analyses

were conducted 72 hours after treatment—by which time multiple downstream responses have likely

been initiated (given that germination typically occurs one to two days after water

absorption)—many previous studies may not have clearly identified the primary targets of

coumarin-induced germination inhibition or fully accounted for the associated cellular structural

changes.

In this study, we addressed these unresolved questions. First, we examined which structural elements

of coumarin are minimally required for germination inhibition. Next, using coumarin-modified

compounds, we investigated the correlation between germination inhibitory effects and molecular

descriptors predicted by chemoinformatics, such as molecular volume and polar surface area. Finally,

to evaluate the early effects of coumarin on germination inhibition, we conducted a transcriptomic

analysis 24 hours after the addition of coumarin.

Results

Structural elements of coumarin minimally required for germination inhibition

Our experimental investigation into the minimal structural elements of coumarin required for

germination inhibition revealed that the ester bond and benzene ring within its substructure are

important for this activity, while the role of the double bond is species-dependent. In this study, we

evaluated the germination effects on both white clover and Arabidopsis (Supplementary Tables 1 and

2) seeds using compounds with substructures analogous to that of coumarin (Fig. 1a): α pyrone (Fig.

1b), which lacks the benzene ring present in coumarin; β tetralone (Fig. 1c), which is missing both

the double bond and the ester bond from coumarin's lactone ring; and 3,4 dihydrocoumarin (Fig.

1d), which lacks only the double bond from the lactone ring.

For clover seeds, the germination rate for the negative control (water only) was 94% compared to

29% for coumarin-treated seeds. Seeds treated with α pyrone and β tetralone showed germination

rates of 82% and 86%, respectively—values significantly higher than those of the coumarin-treated

group and only slightly lower than the negative control (Fig. 1e). In contrast, the germination rate for

seeds treated with 3,4 dihydrocoumarin was 12%, markedly lower than both the coumarin-treated

seeds and the negative control.

For Arabidopsis seeds, the negative control yielded a germination rate of 66%, while no germination

occurred in the coumarin-treated group. Seeds treated with α‑pyrone and β‑tetralone exhibited

germination rates of 44% and 44%, respectively—again, significantly higher than those for

coumarin-treated seeds and only marginally lower than the control (Fig. 1f). Notably, the

germination rate for seeds treated with 3,4‑dihydrocoumarin was 41%; although this rate was

significantly lower than the negative control, it was considerably higher than that observed for

coumarin, indicating a limited inhibitory effect.

Collectively, these studies using two different plant species demonstrate that the ester bond and

benzene ring are important for coumarin-mediated germination inhibition, while the contribution of

the double bond varies between species.

Effect of coumarin-modified compounds on germination inhibition

After determining which substructures of the coumarin skeleton are critical for germination

inhibition, we next examined how modifications to coumarin affect its inhibitory activity.

Specifically, we focused on compounds that retain the important ester bond and benzene ring while

preserving the double bond, given that the importance of the double bond for germination inhibition

varies among plant species. Using these coumarin-modified compounds, we evaluated their

structure–activity relationships.

Our results demonstrate that the inhibitory effects of modified coumarin analogs differ among plant

species; however, coumarin consistently proves to be one of the most potent. To evaluate the factors

influencing this activity, we investigated various molecular descriptors for the coumarin-modified

compounds. TPSA is defined as the sum of the surface areas of all polar atoms—primarily oxygen

and nitrogen, along with their attached hydrogens—in a molecule, reflecting its potential to interact

with aqueous environments and traverse cell membranes. In addition, we calculated the molecular

volume, which represents the three-dimensional space occupied by the molecule; the molecular

surface area, defined as the total surface area of its three-dimensional structure; and the lipophilicity

index LogP, defined as the logarithm of the octanol–water partition coefficient, serving as an

indicator of hydrophobicity. These descriptors were determined for coumarin and the compounds

shown in Fig. 2a–d (Supplementary Tables 3), and we examined the correlations between these

values and germination inhibition.

Based on a correlation coefficient threshold of $r = 0.5$, experiments using white clover seeds showed

that a correlation was observed between germination inhibition and logP ($r = -0.60$) as well as TPSA

($r = 0.77$), but no correlation was found with molecular volume ($r = 0.36$) or molecular surface area

($r = 0.45$) (Fig. 2e–h). Furthermore, in experiments using Arabidopsis seeds, correlations were

observed between germination inhibition and molecular volume ($r = 0.73$), molecular surface area (r

$= 0.77$), and TPSA ($r = 0.58$), but no correlation was found with logP ($r = 0.23$) (Fig. 2i–l). In all

experiments, the p-values calculated from the test of no correlation exceeded the commonly used

threshold value of 0.05.

Thus, in both plant species, TPSA was the only molecular descriptor that showed a correlation with

germination. Although this suggests that TPSA might be important, it should be noted that none of

the p-values were sufficiently low. The key finding is that, although the effects of modified coumarin

analogs vary among plant species, the compound that exhibited the strongest germination inhibitory

activity is coumarin itself.

Coumarin suppresses the expression of genes encoding expansin proteins

Utilizing Arabidopsis—a model organism with a well-established genetic profile—we examined the

impact of coumarin on seeds using RNA-Seq analysis. Our results revealed that coumarin notably

alters the expression of genes encoding expansins, which are responsible for loosening plant cell

walls. The RNA-Seq analysis compared gene expression in seeds treated with coumarin for 24 hours

to that in control seeds treated with water only. Three biological replicates were obtained for each

group, yielding a total of six samples with an average of 3.9 Gbp of nucleotide sequences per sample.

Principal component analysis (Supplementary Fig 1A) demonstrated that the samples clustered into

two groups (coumarin-treated and water-treated), with the coumarin-treated group exhibiting greater

intra-group variability compared to the water-treated group. Using an adjusted p-value threshold of

<0.05, we identified 3,416 differentially expressed genes (DEGs) out of 21,133 detected genes

(Supplementary Spreadsheets 1). Differences in gene expression were shown in an MA plot

(Supplementary Fig 1B).

Analysis of the DEGs revealed that many members of the expansin family (*EXPA2/3/9/15*,

ATEXPA1/10) exhibited a significant reduction in expression (Fig. 3a, Supplementary Tables 4).

Among these expansin genes, we further investigated the temporal expression of *EXPA2*, *ATEXPA1*,

and *ATEXPA10*—which showed the most pronounced changes in RNA-Seq analysis—using

real-time RT-PCR (Fig. 3b–d). In the water-treated control group, the expression of these genes

increased until 36 hours after water absorption and then declined. In contrast, in the coumarin-treated

group, the expression levels were markedly suppressed.

Based on the association with expansin genes, we performed a Gene Ontology (GO) analysis

focusing on categories such as "cell wall," "plant-type cell wall," and "unidimensional cell growth"

(Supplementary Tables 5). This analysis revealed the enrichment of numerous expansin family genes

within these categories. Additionally, other genes involved in xyloglucan metabolism, such as

α-Xylosidase1 (*XYL1*) and members of the Xyloglucan endotransglucosylase/hydrolase (*XTH*)

family, were also identified, with their expression data presented in Fig. 3e (Supplementary Tables 6

and 7).

Coumarin influences the expression of various genes

We utilized gene set enrichment analysis (GSEA) (Table 1) to investigate the impact of coumarin on

gene expression. In coumarin-treated seeds, pathways related to ribosome function—specifically

"Ribosome" and "Ribosome, eukaryotes"—were downregulated. Notably, genes encoding

components of the large ribosomal subunit (*RPL*; 111 genes) and the small ribosomal subunit (*RPS*;

80 genes) were markedly suppressed (Fig. 4a and 4b, Supplementary Spreadsheets 2 and 3).

Conversely, we observed upregulation in three pathways: "glutathione-mediated detoxification II,"

"β-Oxidation," and "Jasmonic acid biosynthesis" (Supplementary Table 8).

Further insights were obtained through Gene Ontology (GO) analysis (Supplementary Spreadsheets

4). The GO Process analysis revealed activation of genes involved in "sulfate assimilation" and

"response to toxic substances," whereas genes associated with "ribosome biogenesis," "translation,"

and "unidimensional cell growth" were inhibited (Fig. 4c). Additionally, the GO Component analysis

indicated that while genes encoding vacuolar components were upregulated, those forming part of

the ribosomes and cell wall were downregulated (Fig. 4d). Under the GO Function category, genes

related to "RNA binding" and "glutathione gamma-glutamylcysteinyltransferase activity" were

upregulated (Fig. 4e).

Moreover, the expression of gibberellin 20-oxidase 1 (*GA20ox1*), a key enzyme in gibberellin

synthesis, was significantly reduced (Fig. 4f, Supplementary Tables 9). Within the "unidimensional

cell growth" category, *PIN1*—a member of the auxin transporter PIN family—and other PIN family

members (*PIN1/2/3/7*) were also downregulated (Fig. 4g, Supplementary Table 10).

Discussion

One of the most significant findings of this study is that coumarin suppresses the expression of genes

encoding expansins, which play a crucial role in loosening plant cell walls[17,18,19]. It is well

established that the induction of expansin expression immediately following water uptake is critical

for germination. For example, studies on tomato seeds have shown that *LeEXP4* expression

increases after water absorption, peaks at 24 hours, and then declines[20]. This pattern is consistent

with our observation that *EXPA2*, *ATEXPA1*, and *ATEXPA10* expression peaks at 36 hours after water

uptake and subsequently decreases. Moreover, in the tomato study, puncture force analysis revealed

that the weakening of the endosperm cap gradually progresses after water uptake, reaching a

minimum at 36 hours in tandem with the expression pattern of *LeEXP4*, thereby supporting its role

in germination[20]. Additionally, research on Arabidopsis *EXPA2* (*EXP2*) has demonstrated that its

expression peaks at 24 hours after water uptake before declining[21]. Although there is a slight time

lag—likely due to experimental conditions—the overall pattern of an initial increase followed by a

decrease aligns with our observations. In that study, the delayed germination observed in an *expa2*

mutant further underscored the essential role of *EXPA2* in germination. In summary, based on these

previous studies and our findings, it is evident that one of the final targets of coumarin's germination

inhibitory effect is the loosening of cell walls mediated by expansins.

In the transcriptome analysis of *Eleusine indica* conducted by Zhang et al.[12], no suppression of

expansin gene expression by coumarin was observed. In contrast, our analysis—performed 24 hours

after water uptake—revealed significant suppression. This discrepancy is likely due to the timing of

the analyses, as expansin gene expression is induced between 24 and 36 hours after water uptake and

then declines. By analyzing gene expression at this earlier stage, we were able to detect the

suppression of expansin gene expression by coumarin.

Our study indicates that the primary mechanism underlying coumarin-induced germination

inhibition is the suppression of the expression of expansin genes, which are critical for loosening

plant cell walls. Moreover, coumarin not only suppresses expansin gene expression but also

downregulates genes involved in cell wall synthesis. In particular, we observed that the expression of

genes encoding endoxyloglucan transferase A4 (*XTH5*) and α-xylosidase 1 (*XYL1*)—enzymes that

interact with xyloglucan, a hemicellulose abundantly present in cell walls and essential for cell

elongation[22,23]—is dramatically reduced by coumarin treatment. Collectively, these results suggest

that coumarin exerts a multifaceted effect on cell walls to inhibit germination; further research is

warranted to fully elucidate the underlying mechanisms.

Our research demonstrated that coumarin downregulates genes involved in the gibberellin synthesis

pathway, linking this suppression to a decrease in cell wall loosening—a process critical for seed

germination. Gibberellins (GA) regulate various germination-associated biological processes[24]. In

addition, numerous studies have shown that GA induces expansin expression[25,26,27,28,29], with one

particularly significant report confirming that GA induces expansin expression during seed

germination[29] Notably, Chen's discovery that coumarin reduces gibberellin synthesis further

supports our transcriptomic findings[10].

It has been reported that coumarin interferes with polar auxin transport in the Arabidopsis root apical

meristem[30]. Although the phenomenon we observed differs, we discovered that coumarin suppresses

the expression of genes encoding members of the auxin transporter PIN family. In addition to its

effect on the gibberellin signaling pathway, coumarin likely also influences the auxin signaling

pathway.

A secondary discovery in this transcriptomic research is the significant suppression of genes

encoding ribosomal proteins by coumarin. In line with our findings, Zhang et al. reported through

transcriptomic analysis that coumarin also inhibits the expression of these genes in *Eleusine indica*[12].

Furthermore, their metabolomic analysis indicated that the synthesis of aminoacyl tRNAs is

inhibited. While we examined gene expression 24 hours after water uptake and Zhang et al.

conducted their analysis after 72 hours, both studies consistently show that coumarin suppresses the

expression of genes encoding ribosomal components. Therefore, the translation system in seeds is

likely another major target of coumarin. Given that ribosome biogenesis requires a substantial

energy investment from the cell[31], it is conceivable that seeds exposed to chemical stressors like

coumarin would suppress ribosome production. Moreover, since these suppressions were observed

as early as 24 hours, it suggests that the inhibition of ribosome assembly occurs at an early stage,

prior to the onset of additional cellular changes.

Furthermore, our study revealed that within the coumarin skeleton, the benzene ring and ester bond

are important for its germination inhibitory activity. In contrast, the role of the double bond appears

to be species-specific: it is non-essential in clover but plays an important role in Arabidopsis. Our

Arabidopsis results are consistent with Mayer et al.'s findings, which indicated that the absence of

the double bond weakens the germination inhibitory effect[5]. These observations suggest that the

impact of dihydrocoumarin on germination is species-specific, reflecting differences in the

underlying mechanisms of action.

Our research using two plant species suggests that the germination inhibitory effect of

coumarin-modified compounds is correlated with their topological polar surface area (TPSA). In line

with this observation, Mayer et al. demonstrated that the germination inhibition of modified

coumarin compounds decreases in the following order: NO_2 (TPSA: 49.36) > OH (TPSA: 20.23) >

Cl (TPSA: 0)[5]. These findings imply that higher TPSA values may enhance the compounds' ability

to interact with cellular targets, possibly by increasing cell membrane permeability or facilitating

receptor binding[32]. However, our experiments with Arabidopsis revealed a weak correlation between

TPSA and germination inhibition, suggesting that the influence of TPSA on activity may be

modulated by species-specific physiological or biochemical factors.

Moreover, we did not observe significant correlations common to both plant species between

germination inhibition and other molecular descriptors—namely, molecular volume, molecular

surface area, and the lipophilicity index (LogP). Although these parameters are widely used in

animal pharmacology and agrochemical research (e.g., in pesticide development), our results

indicate that their impact on the inhibitory effect is limited.

An important observation is that while the inhibitory effects of various modified coumarin analogs

differed among species, coumarin itself consistently proved to be one of the most potent germination

inhibitors in both plant species examined. This consistency might be related to the ecological role of

coumarin as a naturally produced compound in many plants, where its structure has been

evolutionarily optimized for competitive interactions.

Despite the challenges in linking molecular descriptors directly to germination inhibition, the finding

that dihydrocoumarin exhibited a stronger inhibitory effect than coumarin in white clover is

particularly intriguing. This result suggests that specific modifications to the coumarin structure can

enhance activity in a species-specific manner, opening up potential avenues for the development of

targeted herbicides.

In summary, our study underscores the complexity of the structure–activity relationships in coumarin

and its analogs. Future research should aim to identify additional molecular descriptors and

investigate plant-specific factors that influence the bioactivity of these compounds. Such efforts will

not only deepen our understanding of coumarin's role in plant competition but also aid in the design

of more effective, targeted herbicidal agents.

Methods

Evaluation of the importance of coumarin substructures in germination inhibition

Coumarin (Tokyo Chemical Industry, Tokyo, Japan) was employed as a positive control. For the

assessment of the benzene ring, α-pyrone (Sigma-Aldrich Japan, Tokyo, Japan) was utilized, while

3,4-dihydrocoumarin (Tokyo Chemical Industry, Tokyo, Japan) was used for the evaluation of the

double bond, and β-tetralone (Sigma-Aldrich Japan, Tokyo, Japan) was used for the assessment of

the ester bond. These compounds were diluted with water and prepared at a concentration of 0.10

mmol/L. Seeds of white clover (*Trifolium repens*, purchased from Takii, Kyoto, Japan) were sown

onto cotton-lined petri dishes, after which the solutions were added. Ion-exchanged water, without

the addition of compounds, served as the negative control. Subsequently, the petri dishes were

incubated in darkness at 20°C for 48 hours in an incubator (A4201, IKUTA Sangyo Co, Ltd, Nagano,

Japan). For the germination test using *Arabidopsis thaliana* seeds (ecotype Col-0, purchased from

Inplanta innovations, Kanagawa, Japan), in order to achieve a sufficient germination rate in the control

group, the seeds were pre-soaked in 1 mL of solution at 20°C for 24 hours. After this pre-treatment, the

germination test was conducted in the same manner as that for the white clover seeds. The number of

germinations per petri dish was counted, and the percentage of germinated seeds out of the total

number sown was calculated as the germination rate.

Correlation analysis between cheminformatics-estimated indices and germination inhibition

effects

Among the coumarin-modified compounds, five compounds were prepared: coumarin (KANTO

CHEMICAL CO., INC., Tokyo, Japan), 7-methoxy coumarin (Tokyo Chemical Industry, Tokyo,

Japan), aesculetin (Apollo Scientific Ltd, Manchester, UK), psoralen (Tokyo Chemical Industry,

Tokyo, Japan), and umbelliferone (FUJIFILM Wako Pure Chemical Co., Osaka, Japan). The

germination rates were calculated in the same manner as described above.   Various indices

estimated by cheminformatics for each compound were calculated using the following methods. The

Topological Polar Surface Area (TPSA) and the octanol/water partition coefficient (LogP) were

computed using SwissADME (http://www.swissadme.ch/)[33]. The molecular surface area and van der

Waals volume were determined using winmostar (V11.5.8, https://winmostar.com/en/). The

correlations between these indices and the germination rates were investigated. This analysis was

performed on the germination rates of both *Arabidopsis thaliana* seeds and white clover seeds.

Transcriptome analysis by RNA-Seq

Arabidopsis seeds, known for their well-characterized genetic information, were used. Initially,

20 mg of seeds were treated with either a 10 mmol/L coumarin solution or water and incubated for

24 hours. Compared to the structure–activity relationship experiments—which used 0.10 mmol/L

due to solubility limitations—a higher concentration of 10 mmol/L coumarin was employed in the

transcriptome analysis. This was because solubility was not an issue in this experiment, and using a

higher concentration allowed us to clearly demonstrate its impact on the transcriptome. Three

biological replicates were used for each condition. The seeds were then manually crushed with a

pestle, and RNA extraction was performed using ISOSPIN Plant RNA (NIPPON GENE, Tokyo,

Japan). Subsequently, RNA concentration was measured using the Qubit RNA BR Assay Kit

(Thermo Fisher Scientific, MA, USA). RNA samples were sent to Novogene (Singapore) for quality

testing using a Bioanalyzer (Agilent, California, USA), library construction using the NEBNext

Ultra II RNA Library Prep Kit for Illumina, and sequencing on the NovaSeq 6000 platform (Illumina,

California, USA). The resulting FASTQ data were analyzed using RaNA-seq (https://ranaseq.eu/)[34],

where differential gene expression analysis was conducted based on the DESeq2 method[35]. Using

the DESeq2 method with a pValue cutoff of 0.05 and a parametric Wald test, differentially expressed

genes were detected. Genes with an adjusted p-value (padj) below 0.05 were considered significantly

differentially expressed. PCA plots and MA plots were also generated using RaNA-seq. In addition,

GSEA[36,37] was performed on these DEGs. The GSEA, including pathway and gene ontology

analyses, determined the extent of impact by the normalized enrichment score (NES). The pathway

analysis referenced the KEGG PATHWAY Database, and the gene ontology analysis was carried out

with reference to the National Center for Biotechnology Information (NIH) database, analyzing

biological process, cellular component, and molecular function domains.

Real-time RT-PCR Analysis

The time-course analysis of expansin gene expression after coumarin treatment was performed using

real-time RT-PCR. Arabidopsis seeds were treated with either a 10 mmol/L coumarin solution or

water and incubated for 6, 12, 24, 36, and 48 hours, after which RNA extraction and quantification

were performed as described above. cDNA synthesis was performed using the ReverTra Ace qPCR

RT Master Mix with gDNA Remover (TOYOBO, Osaka, Japan) according to the manufacturer's

protocol. Real-time RT-PCR was carried out with the QuantStudio 3 Real-Time PCR system

(Thermo Fisher Scientific, MA, USA) using the PowerUp SYBR Green Master Mix (Thermo Fisher

Scientific, MA, USA) in triplicate. The primers used for real-time RT-PCR were as follows:

ATEXPA1-F: 5'-TGCACGAGGCCTTGATGATG-3', *ATEXPA1*-R:

5'-AACTCTCCCTCTCGCTTCGA-3', *EXPA2*-F: 5'-ATGCTACCTTCTATGGTGGAGCTG-3',

EXPA2-R: 5'-AAAGCAGGCCCCACATTTCTG-3', *ATEXPA10*-F:

5'-CCGTTAAAAAGGGCAAGTTGGTTAATCTCT-3', *ATEXPA10*-R:

5'-GATTAGACGTCATCAAGGCCTTCTGCAAAA-3', *PP2A*-F:

5'-ATCGCTACTCAGTTTAACCACA-3', *PP2A*-R: 5'-TAGCAATAGTTTGGGGCACTAA-3'. The

PCR program was as follows: 50°C for 2 min, 95°C for 2 min, followed by 40 cycles of 95°C for 1 s

and 60°C for 30 s. The results were normalized to the expression level of *PP2A*, and the fold change

relative to the 0 h time point was calculated.

Utilization of generative AI

ChatGPT (https://chat.openai.com/) was used to rewrite the manuscript into more appropriate

English and to proofread the text. The correctness of the revisions was subsequently verified by the

authors.

Legal statement

Permissions or licenses were not required for the laboratory use of these seeds. Additionally, there

are no specific guidelines or legislation required for the use of the seeds in our study.

References

1. Jain, P. K., & Joshi, H. Coumarin: chemical and pharmacological profile. J. Appl. Pharm. Sci. 2,

236–240 (2012).

2. Bourgaud, F. et al. Biosynthesis of coumarins in plants: a major pathway still to be unravelled for

cytochrome P450 enzymes. Phytochem. Rev. 5, 293–308 (2006).

3. Takahashi, K., Tsutsumi, Y., Ohtani, H., & Katsuki, T. Variation of fragrance constituents in the

leaves of *Prunus*. Biochem. Syst. Ecol. 34, 127–135 (2006).

4. Audus, L. J., & Quastel, J. H. Coumarin as a selective phytocidal agent. Nature 159, 320–324

(1947).

5. Mayer, A. M., & Evenari, M. The relation between the structure of coumarin and its derivatives,

and their activity as germination inhibitors. J. Exp. Bot. 3, 246–252 (1952).

6. Mino, Y., Kawabe, M., & Idonuma, T. Effect of coumarins on germination and growth of some

forage grasses and legumes. Japanese J. of Grassland Sci. 21, 91–95 (1975).

7. Chattha, F. A., Munawar, M. A., & Kousar, S. Plant growth regulating activity of coumarin

derivatives. Horticult. Int. J. 2, 225–30 (2018).

8. Rial, C., Varela, R. M., Molinillo, J. M. G., Peralta, S., & Macías, F. A. Sunflower metabolites

involved in resistance mechanisms against broomrape. Agronomy 11, 501;

10.3390/agronomy11030501 (2021).

9.   Chen, B. X. et al. Coumarin-induced delay of rice seed germination is mediated by suppression of

abscisic acid catabolism and reactive oxygen species production. Front. Plant Sci. 10, 828;

10.3389/fpls.2019.00828 (2019).

10. Chen, B. X., Peng, Y. X., Yang, X. Q. & Liu, J. Delayed germination of *Brassica parachinensis*

seeds by coumarin involves decreased GA4 production and a consequent reduction of ROS

accumulation. Seed Sci. Res. 10, 224–235 (2021).

11. Araniti, F., Prinsi, B. & Espen, L. The delay of *Raphanus raphanistrum* subsp. sativus (L.)

Domin seed germination induced by coumarin is mediated by a lower ability to sustain the

energetic metabolism. Plants (Basel) 11, 843; 10.3390/plants11070843 (2022).

12. Zhang, T. J. et al. Metabolic pathways modulated by coumarin to inhibit seed germination and

early seedling growth in *Eleusine indica*. Plant. Physiol. Biochem. 203, 108035;

10.1016/j.plaphy.2023.108035 (2023).

13. Shi, Y. et al. De novo transcriptome analysis of *Cnidium monnieri* (L.) Cuss and detection of

genes related to coumarin biosynthesis. PeerJ 8, 10157; 10.7717/peerj.10157 (2020).

14. Zhang, X., Li, C., Hao, Z., & Liu, Y. Transcriptome analysis provides insights into coumarin

biosynthesis in the medicinal plant *Angelica dahurica* cv. Yubaizhi. Gene 888, 147757;

10.1016/j.gene.2023.147757 (2023).

15. Luo, K., Wu, F., Zhang, D. et al. Transcriptomic profiling of *Melilotus albus* near-isogenic lines

contrasting for coumarin content. Sci. Rep. 7, 4577; 10.1038/s41598-017-04111-y (2017).

16. Yang, B. et al. Integration of transcriptomics and metabolomics to identify key coumarin

biosynthetic genes in *Bupleurum chinense*. Biotechnol. Biotechnol. Equip. 3, 1858–1868 (2021).

17. McQueen-Mason, S., Durachko, D. M., & Cosgrove, D. J. Two endogenous proteins that induce

cell wall extension in plants. Plant Cell 4, 1425–1433 (1992).

18. Cosgrove, D. J. et al. The growing world of expansins. Plant. Cell Physiol. 43, 1436–1444

(2002).

19. Cosgrove, D. J. Plant expansins: diversity and interactions with plant cell walls. Curr. Opin. Plant

Biol. 25, 162–172 (2015).

20. Chen, F. & Bradford, K. J. Expression of an expansin is associated with endosperm weakening

during tomato seed germination. Plant Physiol. 124, 1265–1274 (2000).

21. Yan, A., Wu, M., Yan, L., Hu, R., Ali, I., & Gan, Y. *AtEXP2* is involved in seed germination and

abiotic stress response in *Arabidopsis*. PLoS ONE 9, e85208; 10.1371/journal.pone.0085208

(2014).

22. Rose, J. K., Braam, J., Fry, S. C. & Nishitani, K. The XTH family of enzymes involved in

xyloglucan endotransglucosylation and endohydrolysis: current perspectives and a new unifying

nomenclature. Plant. Cell Physiol. 43, 1421–1435 (2002).

23. Günl, M. & Pauly, M. *AXY3* encodes a α-xylosidase that impacts the structure and accessibility of

the hemicellulose xyloglucan in Arabidopsis plant cell walls. Planta 233, 707–719 (2011).

24. Urbanova, T. & Leubner-Metzger, G. Gibberellins and seed germination. Annu. Plant Rev. 49,

253–284 (2016).

25. Jones, R. L. & Kaufman, P. B. The role of gibberellins in plant cell elongation. Crit. Rev. Plant.

Sci. 1, 23–47 (1983).

26. Butcher, D. N. & Street, H. E. The effects of gibberellins on the growth of excised tomato roots. J.

Exp. Bot. 11, 206–216 (1960).

27. Xu, Q., Krishnan, S., Merewitz, E., Xu, J. & Huang, B. Gibberellin-regulation and genetic

variations in leaf elongation for tall fescue in association with differential gene expression

controlling cell expansion. Sci. Rep. 6, 30258; 10.1038/srep30258 (2016).

28. Xu, Q., Burgess, P., Xu, J., Meyer, W. & Huang, B. Osmotic stress- and salt stress-inhibition and

gibberellin-mitigation of leaf elongation associated with up-regulation of genes controlling cell

expansion. Environ. Exp. Bot. 131, 101–109 (2016).

29. Ogawa, M. et al. Gibberellin biosynthesis and response during Arabidopsis seed germination.

Plant Cell 15, 591–604 (2003).

30. Bruno, L. et al. Coumarin interferes with polar auxin transport altering microtubule cortical array

organization in *Arabidopsis thaliana* (L.) Heynh. root apical meristem. Int. J. Mol. Sci. 22, 7305;

10.3390/ijms22147305 (2021).

31. Shore, David et al. Ribosome biogenesis and the cellular energy economy. Curr. Biol. 32,

611–617 (2022).

32. Silva, A. M. et al. In vitro assessment of pesticides toxicity and data correlation with pesticides

physicochemical properties for prediction of toxicity in gastrointestinal and skin contact

exposure. Toxics 10, 378; 10.3390/toxics10070378 (2022).

33. Daina, A., Michielin, O. & Zoete, V. SwissADME: a free web tool to evaluate pharmacokinetics,

drug-likeness and medicinal chemistry friendliness of small molecules. Sci. Rep. 7, 42717;

10.1038/srep42717 (2017).

34. Prieto, C. & Barrios, D. RaNA-Seq: Interactive RNA-Seq analysis from FASTQ files to

functional analysis. Bioinformatics 36, 1955–1956 (2020).

35. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 15, 550; 10.1186/s13059-014-0550-8 (2014).

36. Subramanian, A. et al. Gene set enrichment analysis: a knowledge-based approach for

interpreting genome-wide expression profiles. Proc. Natl. Acad. Sci. USA 102, 15545–15550

(2005).

37. Mootha, V. K. et al. PGC-1α-responsive genes involved in oxidative phosphorylation are

coordinately downregulated in human diabetes. Nat. Genet. 34, 267–273 (2003).

Acknowledgments

The authors would like to express gratitude for the technical support provided by Nana Nakahara,

Ayane Hirose, Itsuki Chiwata, Saori Nakagawa, Syunsuke Saeki, Yuki Takubo, Haruka Mine,

Tatsuhiro Tanaka, Kotaro Yamamoto, Minoru Nishio, Masato Hara, Sota Fukushima, Shion Tanaka,

and Hikari Kaneshige. RNA extraction and real-time RT-PCR experiments were facilitated at the

Analytical Research Center for Experimental Sciences, Saga University. This work was supported by

grants from the Nabeshima-Hokokai and the Japan Science and Technology Agency under the Super

Science High School project of the Ministry of Education, Culture, Sports, Science and Technology.

Author Contributions

K.F., T.O., S.H., M.Y., and K.M. conceptualized the research plan. S.H., T.O., M.Y., H.M., M.K.,

C.E., K.T., K.T., K.C., R.Ich., and E.F. collected the germination test data. S.H. and T.O. conducted

the cheminformatics analysis. K.F., S.H., T.O., Y.T., K.M., T.Y., and N.R. executed RNA extraction.

K.F., S.H., T.O., Y.T., K.M., and T.Y. conducted real-time RT-PCR. K.F., S.H., T.O., N.R., Y.N., and

K.M. conducted the bioinformatics analysis. K.F. and R.Ito completed the statistical analysis. As

Ph.D. holders, N.R. and Y.N. provided advisory input to the research. The manuscript was drafted by

K.F., S.H., and T.O. and edited by Y.N. The manuscript was finalized by K.F., S.H., T.O., M.Y., N.R.,

Y.N., and K.M. The manuscript was reviewed by all authors.

Data Availability Statement

The raw sequencing data for each sample have been deposited in the DDBJ Sequence Read Archive

(https://www.ddbj.nig.ac.jp/dra/index-e.html; accession numbers DRR546110–DRR546115).

Competing Interests

The authors declare no competing interests. Yukio Nagano is conducting collaborative research with

Senoo Suisan, Inc. and Sante Laboratories, Inc.; however, the research conducted in collaboration

with these for-profit companies is unrelated to the present study.

 Figure Legends

Figure 1. Effects of structural changes in the coumarin skeleton on seed germination. (**a–d**)

Chemical structures of the coumarin-like substances used in this study are shown: coumarin (**a**),

α-pyrone (**b**), β-tetralone (**c**), and 3,4-dihydrocoumarin (**d**). The solutions used in the experiments

were prepared at a concentration of 0.10 mmol/L for both plant species. (**e–f**) Germination rates after

48 hours of incubation with the addition of coumarin-like substances are presented: (**e**) white clover

(*Trifolium repens*), $n = 300$; (**f**) *Arabidopsis thaliana*, $n = 300$ for the control, 3,4-dihydrocoumarin,

and coumarin, and $n = 50$ for α-pyrone and β-tetralone. Below the graph, the presence (+) or absence

(–) of the benzene ring, ester bond, and double bond within the lactone ring for each substance is

indicated. Group comparisons of germination rates were performed using chi-squared tests, with an

asterisk (*) indicating statistical significance ($p < 0.05$). Error bars represent the 95% confidence

intervals of the proportions.

Figure 2. Effects of coumarin-modified compounds on seed germination. (**a–d**) Chemical structures

of additional compounds used in this study, alongside coumarin as shown in Fig. 1, are displayed:

7-methoxy-coumarin (**a**), aesculetin (**b**), psoralen (**c**), and umbelliferone (**d**). The solutions for the

experiments were prepared at a concentration of 0.10 mmol/L for both plant species. (**e–l**)

Relationships between the germination rates of seeds treated with the coumarin-modified

compounds and four calculated structural indicators are shown (white clover (**e–h**) and Arabidopsis

thaliana (**i–l**)). The y-axis represents the germination rates of seeds treated with each

coumarin-modified compound ($n = 300$), and the x-axis represents the molecular descriptors

calculated from their structures: Molecular Volume (**e** and **i**), Molecular Surface Area (**f** and **j**), LogP

(**g** and **k**), and TPSA (**h** and **l**). Pearson's correlation coefficient (r) and the p-value from the test of

no correlation are provided. For each pair with an absolute correlation coefficient greater than 0.5, a

scatter plot with a least-squares linear regression model is shown.

Figure 3. Down-regulation of expansin and cell wall-related genes by coumarin. (**a**) The bar plot

shows the log2-transformed expression ratios of expansin family genes between coumarin-treated

and water-treated control groups, as determined by RNA-Seq. Experiments were performed in

triplicate for each group, with each column representing the expression profile of an individual

sample. (**b–d**) Real-time RT-PCR analysis of expansin family transcript levels: *ATEXPA1* (**b**), *EXPA2*

(**c**), and *ATEXPA10* (**d**). The data were normalized to the expression of the control gene (the *PP2A*

gene encoding Protein Phosphatase 2A) and are presented as the fold change relative to the

expression level at 0 hours. (**e**) Bar plots display the log2 expression ratios between the

coumarin-treated and water-treated control groups for the XYL1 gene and members of the XTH

family. An asterisk (*) indicates statistical significance based on the adjusted p-value (padj < 0.05).

Figure 4. Coumarin Addition Impacts Multiple Pathways. (**a–b**) The heatmaps display the

expression ratios of RPL family genes (**a**) and RPS family genes (**b**) between coumarin-treated and

water-treated control groups as obtained by RNA-Seq. (**c–e**) Bar plots show the normalized

enrichment scores (NES) from GSEA using the NIH Gene Ontology database. The top five activated

and repressed pathways are presented for each gene ontology category: biological process (**c**),

cellular component (**d**), and molecular function (**e**). (**f–g**) Bar plots depict the log2 expression ratios

between the coumarin-treated and water-treated control groups for gibberellin 20-oxidase 1 and 3 (**f**)

and for members of the PIN family (**g**). The asterisk (*) indicates statistical significance based on the

adjusted p-value (padj < 0.05).

Table 1. GSEA Pathway Analysis Results

Pathway_ID	Pathway	NES	padj
ath03010	Ribosome	-2.409	0.011
ath_M00177	Ribosome, eukaryotes	-2.290	0.011
ath_M00049	Adenine ribonucleotide biosynthesis, IMP => ADP,ATP	-1.812	0.011
ath_M00087	beta-Oxidation	1.969	0.011
ath00480	Glutathione metabolism	2.004	0.013
ath_M00113	Jasmonic acid biosynthesis	2.053	0.018
ath00945	Stilbenoid, diarylheptanoid and gingerol biosynthesis	1.902	0.024
ath00230	Purine metabolism	-1.674	0.049