Structure-activity relationships and crucial mechanisms of the coumarin family as germination

inhibitors

Kazuma Fukuda[1], Sota Hyakutake[1], Taiga Oishi[1], Michinari Yoshida[1], Mizuho Koga[1], Chisato Egami[1],

Hyuga Matsura[1], Ryusei Ito[1], Kosei Tsukahara[1], Mitsuki Noda[1], Takako Yoshida[2], Noriko Ryuda[2,*],

Yukio Nagano[2] & Kazuhide Matsutaka[1,*]

[1]*Saga Prefectural Chienkan High School, Saga, Japan*

[2]*Analytical Research Center for Experimental Sciences, Saga University, Saga, Japan.*

*Corresponding authors:

✉E-mail: si3264@edu.cc.saga-u.ac.jp

✉E-mail: matsutaka-kazuhide-a@education.saga.jp

Abstract

Members of the coumarin family, including coumarin and its derivatives, are molecules produced by

various higher plants, such as cherry trees, and are noted for their diverse physiological activities.

Notably, these compounds serve as germination inhibitors, which is considered their primary role.

However, the structural basis underpinning the germination-inhibitory effects of the coumarin family

is not well understood. Our research investigated the structure-activity relationships using coumarin

analogs and revealed that the benzene ring and ester bond within the coumarin skeleton play a crucial

role in germination inhibition. We also identified a correlation between the topological polar surface

area (TPSA) and the inhibitory effects on germination. Furthermore, while the physiological effects

of coumarin on seeds prior to germination have been reported variably, the mechanisms of germination

inhibition remain largely unknown. Using RNA-Seq analysis, we have discovered a novel mechanism

that is pivotal for germination inhibition, where coumarin suppresses the expression of genes

associated with cellulose fiber reassembly. This study holds significant importance in understanding

how the coumarin family contributes to plant competition in nature and promises future contributions

to the agricultural field through the development of novel growth regulation methods.

Keywords: Natural inhibitors, Coumarin structure, Cheminformatics, RNA-Seq, Cell wall

reconfiguration, Ribosome biogenesis

Introduction

Coumarin belongs to the benzopyrone family and is produced by various higher plants, including the cherry tree[1,2]. In Japan, there is a tradition of enjoying the aroma of "sakurayu," literally "cherry blossom tea," and "sakuramochi," a sweet made with mochi (Japanese rice cake) and preserved cherry leaves, which primarily derives its scent from coumarin[3]. Thus, coumarin has a deep association with Japanese traditional culture. Furthermore, not only is coumarin itself produced by some plants, but a variety of coumarin derivatives are also known to be naturally synthesized[2], and others are artificially created. For simplicity, both coumarin and its derivatives will be collectively referred to as the "coumarin family." Members of the coumarin family are known for their diverse physiological activities and are particularly noteworthy in medicine, where, for example, warfarin, a representative member of the coumarin family, is used as an anticoagulant to prevent thromboembolic disorders[4]. Additionally, artificially synthesized coumarin derivatives are gaining attention as potential candidates for new anticancer agents[5]. Thus, members of the coumarin family are gaining interest as compounds with potential applications in pharmaceuticals and other fields.

However, the primary role of the coumarin family in plants is to inhibit the growth of surrounding vegetation through their production. The main mechanism by which members of the coumarin family

inhibit growth is through a strong inhibitory effect on seed germination[6]. Regarding the germination

inhibition by members of the coumarin family, it is unclear which structures are important, as well as

how and where these compounds impact the seeds.

One of the unresolved questions is which structural elements of the coumarin family members are

critical for germination inhibition, namely, the correlation between the molecular structures of various

coumarin family members and their germination inhibitory effects. Mayer et al. conducted a

comparison and reported that structural modifications to coumarin do not destroy the germination

inhibitory effect but only partially attenuate it[7]. This suggests that the germination inhibition effect is

conserved among many members of the coumarin family. On the other hand, coumarin derivatives

with more potent phytotoxic effects than coumarin itself have been discovered[8], indicating a need for

further research into the impact of such modifications.

Another unresolved issue is where and how coumarin family members influence seeds to achieve

germination inhibition. A critical factor in determining where coumarin family members affect seeds

is their impact on the seed coat. As the rupture of the seed coat is crucial for germination, there is a

possibility that the members of the coumarin family act on the seed coat they first come into contact

with. Recent studies have shown that a thick and low water-permeable seed coat plays a role in

negatively regulating the initiation of seed germination[9]. Interestingly, Aliotta et al. demonstrated that

the removal of the seed coat significantly diminishes the effect of coumarin[10]. Moreover, the

interaction of coumarin family members with the seed coat may alter its properties, potentially

changing the water absorption rate of the seed. Thus, while there is a likelihood that coumarin family

members act primarily on the seed coat as described above, it is also necessary to verify whether they

penetrate the seed coat and act inside it.

A critical aspect of understanding how the coumarin family members influence seed germination is

their effect on cellular processes. Several studies have explored this mechanism, with the studies by

Chen et al. particularly notable[11,12]. They demonstrated that coumarin negatively regulates the

expression of the abscisic acid (ABA) metabolic enzyme genes (OsABA8' ox2/3) and reduces the

levels of gibberellin A4 (GA_4). Furthermore, coumarin also suppresses the accumulation of reactive

oxygen species (ROS) and the expression of metabolic enzymes such as superoxide dismutase and

catalase. These outcomes physiologically substantiate the delay in seed germination induced by

coumarin. Omics analyses have greatly contributed to the elucidation of the mechanisms behind

germination inhibition[13,14]. Araniti et al. conducted metabolomic analyses, revealing that coumarin

extensively affects the seed's metabolic profile, impacting amino acid metabolism, the citric acid

(TCA) cycle, and aminoacyl-tRNA biosynthesis[13]. A similar analysis by Zhang et al. is consistent with

Araniti et al.'s findings[14]. Additionally, Zhang et al. conducted transcriptomic analyses 72 hours after

coumarin addition, showing that coumarin influences the expression of genes involved in amino acid

metabolism, the TCA cycle, and protein translation. The outcomes of these transcriptomic analyses

are invaluable. However, given that they were conducted 72 hours after the addition of coumarin, they

may reflect gene expression after a range of changes had already occurred. Many previous studies may

not have clearly identified the final targets of coumarin-induced germination inhibition, implying they

did not account for the cellular structure changes expected during inhibition of germination.

In this study, we addressed the unresolved questions we had raised. To identify the structures critical

for germination inhibition, we explored the structure-activity relationships and pinpointed the

responsible substructures for germination inhibition effects using various coumarin analogs. We also

compared the germination inhibition effects with various molecular indices, such as molecular volume

and polar surface area, as predicted by cheminformatics. To investigate where coumarin affects seeds,

we measured the changes they induce in the seed coat structure and the seed's water uptake. In addition,

we examined the localization of coumarin by studying their absorption spectrum. Finally, to assess the

early effects of coumarin, we performed a transcriptomic analysis 24 hours after the addition of

coumarin.

Results

The impact of coumarin substructures on germination inhibition

Our experimental investigation into the structure-activity relationship of coumarin (Fig. 1A) has

revealed that the ester bond and benzene ring within its substructure are critical for germination

inhibition, while the presence of a double bond is not essential. In this study, we tested the germination

effects on clover seeds using compounds that have similar substructure to coumarin: α-pyrone, which

lacks coumarin's benzene ring (Fig. 1B), β-tetralone, missing both the double bond and ester bond

from coumarin's lactone ring (Fig. 1C), and 3,4-dihydrocoumarin, lacking only the double bond from

the lactone ring (Fig. 1D). The germination rate for seeds with just water added (negative control) was

94%, and for those with coumarin added, it was 29%. The germination rates for seeds treated with α-

pyrone and β-tetralone were 82% and 86%, respectively. These rates were significantly higher than

those observed for seeds treated with coumarin and were only slightly lower than the negative control,

albeit significantly so (Fig. 1E). These results demonstrate the importance of the ester bond and

benzene ring in the substructure of coumarin for germination inhibition. On the other hand, the

germination rate for seeds treated with dihydrocoumarin was 12%, which was significantly lower than

the coumarin-treated seeds and the negative control. This indicates that the absence of the double bond

in coumarin's substructure does not play a significant role in germination inhibition.

Prediction of germination inhibition effects by the coumarin family through TPSA (topological

polar surface area)

Our experimental investigations into the germination inhibition effects of various members of the

coumarin family revealed that the topological polar surface area (TPSA) is an effective predictor of

germination inhibition. In addition to coumarin and 3,4-dihydrocoumarin, we used the compounds

shown in Fig. 2A–D for our analysis. We calculated the molecular volume, molecular surface area,

lipophilicity index LogP, and another lipophilicity index TPSA (Topological Polar Surface Area), for

these substances (Supplementary Table 1) and examined the correlation between these values and the

effects on germination inhibition. Our analysis detected no significant correlation between

germination inhibition and molecular volume (p=0.31 > 0.05), surface area (p=0.25 > 0.05), or LogP

(p=0.26 > 0.05) (Fig. 2E–G). However, we identified a significant correlation between germination

inhibition and TPSA (p=0.019 < 0.05), with a high correlation coefficient (r=0.8349) (Fig. 2H).

Observation of changes in seed coats treated with coumarin

We investigated the possibility of coumarin acting on the outermost layer of the seed, the seed coat, to

inhibit germination and found no changes in the structure or permeability of the seed coat. Examination

of pea seed coats with a scanning electron microscope revealed a three-layered structure consisting of

palisade parenchyma, hourglass cells, and compressed parenchyma, regardless of whether coumarin

was administered (Fig. 3A). Furthermore, measurements of the overall thickness of the seed coat

indicated that both the coumarin-treated and control seeds had seed coats approximately 90 µm thick,

with no detectable difference. Similarly, measurements of the thickness of each tissue layer also did

not reveal any significant differences due to coumarin treatment (Fig. 3B). Additionally, when

measuring water absorption of the seeds, both the coumarin-treated and control seeds began to absorb

water immediately upon immersion. For both, the water absorption rate reached a near plateau after

about 12 hours, and thereafter, very little additional water was absorbed (Fig. 3C).

Penetration of coumarin beyond the seed coat into the seed interior

Based on the above results, we inferred that coumarin penetrates the seed coat and enters the interior

of the seed, which we then demonstrated. We measured the absorption spectrum of the liquid obtained

from crushed pea seeds from which the seed coat had been removed. The results showed a peak around

the wavelength of 270 nm, which corresponds to the absorption maximum of coumarin, only in the

extract from the coumarin-treated seeds (Fig. 3D).

Transcriptomic impact of coumarin on seed germination

Utilizing Arabidopsis, which has a well-established genetic profile, we examined the impact of

coumarin on seeds using RNA-Seq analysis. We found that coumarin notably alters the expression of

gene groups involved in protein translation and cellular elongation growth, which is mediated by the

reassembly of cellulose fibers. The RNA-Seq analysis compared gene expression in seeds treated with

coumarin for 24 hours with control seeds treated with water only. Three biological replicates for each

group resulted in a total of six samples for analysis, with an average of 3.9 Gbp of nucleotide sequences

obtained per sample. Out of 21,133 detected genes, 3,416 were identified as differentially expressed

genes (DEGs) (Supplementary Spreadsheets 1). Notably, gene groups coding for proteins that

constitute the large subunit of ribosomes, RPL (111 genes), and the small subunit, RPS (80 genes),

had their expression significantly suppressed by coumarin treatment (Fig. 4A and 4B, Supplementary

Spreadsheet 2 and 3). Many genes of the expansin family (EXPA2/3/4/8/9/15, ATEXPA1/10) also

exhibited a significant reduction in expression (Fig. 4C, Supplementary Table 2)

The results of the pathway analysis of the gene set enrichment analysis (GSEA) are presented in Table

1. In seeds treated with coumarin, a downregulation of gene expression was observed in "Ribosome"

and "Ribosome, eukaryotes," while an upregulation was noted in three pathways including

"glutathione-mediated detoxification II," "β-Oxidation," and "Jasmonic acid biosynthesis" (Fig. 5A).

Subsequent results of the GSEA Gene Ontology (GO) analysis are detailed (Fig. 5B–D, Supplementary

Spreadsheet 4). The analysis of GO Process analysis indicated that genes related to "sulfate

assimilation" and "response to toxic substances" were activated, while those involved in "ribosome

biogenesis," "translation," and "unidimensional cell growth" were inhibited (Fig. 5B). Furthermore,

the analysis of GO Component revealed that genes forming parts of "vacuoles" were activated, while

the expression of genes constituting "ribosomes" and the "cell wall" was suppressed (Fig. 5C).

Additionally, under GO Function, genes associated with "RNA binding" and "glutathione gamma-

glutamylcysteinyltransferase activity" showed increased expression (Fig. 5D).

From the association with expansins, we focused on the enrichment of "cell wall," "plant-type cell

wall," and "unidimensional cell growth" in the GO analysis (Supplementary Table 3). Analysis of these

enriched groups included many expansin family genes. Other genes involved in the metabolism of

xyloglucan, such as α-Xylosidase1 (XYL1) and the Xyloglucan endotransglucosylase/hydrolase

(XTH) family, were also present, with expression data shown in Fig. 5E (Supplementary Table 4 and

5). Additionally, the SKU similar (sks) family, coding for proteins of unknown function, and

gibberellin 20-oxidase 3 (GA20ox3) involved in gibberellin synthesis were included, and their

expressions were significantly suppressed as well (Fig. 5F and 5G, Supplementary Table 6 and 7).

Moreover, within the "unidimensional cell growth" category was PIN1 from the auxin transporter PIN

family, and the expression of many from the PIN family (PIN1/2/3/7) was also reduced (Fig. 5H,

Supplementary Table 8).

Discussion

The most significant finding of this study is that coumarin suppresses the expression of genes coding

for proteins involved in the reconfiguration of the cell wall, such as expansins, suggesting that one of

the final targets of coumarin's germination inhibitory effect is the cell wall. Expansins are a type of

plant cell wall protein that plays a crucial role in plant growth and development[15,16]. They have a

distinct structure comprising two domains: an N-terminal domain structurally similar to glycoside

hydrolase family 45 (GH45) and a C-terminal domain distantly related to group 2 grass pollen

allergens. While no enzymatic activity has been reported for expansins, they can induce pH-dependent

wall extension and stress relaxation, appearing to weaken non-covalent bonds between cell wall

polysaccharides. Expansins are known to play important physiological roles in various plant growth

and developmental processes, including cell enlargement, fruit softening, pollen tube growth, root hair

emergence, and abscission.

The suppression of expansin gene expression by coumarin is significant. Furthermore, coumarin also

downregulated the expression of genes coding for endoxyloglucan transferase A4 (XTH5) and α-

xylosidase1 (XYL1), which are known to interact with xyloglucans, a hemicellulose abundantly

present in cell walls and critical for cellular elongation growth[17,18]. The SKU similar (sks) family

proteins, while functionally uncharacterized, may be implicated in the biosynthesis of primary cell

walls in coordination with SKU[19]. Thus, similar to expansins, XTH5, and XYL1, coumarin may target

these proteins to inhibit cellular elongation growth.

Cell wall reorganization, directed by genes such as expansins, enables cellular elongation growth by

orienting cellulose fibers in a specific direction[20]. However, seeds treated with coumarin, exhibiting

reduced expression of such genes, would likely be hindered in this directional elongation growth. The

inhibition of expansin expression induction could potentially interfere with the final stages of

germination, where the radicle breaks through the seed coat. Notably, this suppression of expression

was evident at the 24-hour mark, indicating that the inhibition of cell wall reconfiguration occurs early

before various other changes can occur.

Our research also showed that genes involved in the gibberellin synthesis pathway were

downregulated by coumarin, linking cell wall reconfiguration to this suppression. Gibberellins (GA)

control various biological events associated with germination, including cell elongation, which

involves cell wall reconstruction[21,22]. There are also reports of GA inducing expansin

expression[23,24,25,26]. For instance, in tall fescue (*Festuca arundinacea*) leaves, GA was shown to induce

expansin expression differently between fast-growing and dwarf cultivars[23]. Similarly, GA likely

induces expansin expression during seed germination.

Studies on the relationship between plant hormones, including GA, and coumarin provide essential

insights into coumarin's mechanism of germination inhibition. Particularly notable is Chen's discovery

that coumarin reduces the synthesis of gibberellin, a hormone crucial for germination and elongation

growth[12]. These findings are consistent with our transcriptomic analysis. Additionally, Bruno et al.

demonstrated that the expression of the auxin transporter PIN family in Arabidopsis roots is suppressed

by the addition of coumarin[27]. Our results clarify that this effect is also present in pre-germinative

seeds. Therefore, it is plausible that coumarin's impact on GA and auxin plays a vital role in the

inhibition of germination.

Another important discovery in this research is the significant suppression by coumarin of genes

coding for ribosomal proteins. In line with our findings, Zhang et al., through transcriptomic analysis,

have also reported the inhibition of expression of these genes by coumarin in *Eleusine indica*.

Furthermore, Zhang et al. have reported that the synthesis of aminoacyl tRNAs is inhibited as a result

of their metabolomic analysis[14]. We examined gene expression at 24 hours post-coumarin treatment,

while Zhang et al. did so after 72 hours. Despite the difference in timing of the analyses, coumarin

consistently inhibits the expression of genes encoding components of ribosomes. Therefore, the

translation system in seeds is likely another major target of coumarin. However, the pathway through

which coumarin affects the synthesis of ribosomal components remains unclear. Ribosome biogenesis

is a process that demands a substantial energy investment from the cell, so it is conceivable that seeds

exposed to chemical stressors like coumarin may suppress ribosome biogenesis. Indeed, in eukaryotic

cells such as yeast and animal cells, ribosome biogenesis is known to be intricately regulated[28,29]. Yet,

it is currently unclear whether plants possess such a system. Moreover, since these suppressions of

expression were observed as early as 24 hours, it implies that inhibition of ribosome assembly occurs

at an early stage before various changes take place.

Additionally, our study has discovered that benzene rings and ester bonds are the structures responsible

for coumarin's effects. Knowledge regarding the relationship between coumarin's structure and activity

is limited and largely dates back to decades-old research by Mayer et al., which indicated the

importance of double bonds in dihydrocoumarins for germination inhibition[7]. Our germination tests

using three types of coumarin analogs suggest that benzene rings and ester bonds play key roles in its

effects, while double bonds are not crucial. Comparing the structures of coumarin with a double bond

and dihydrocoumarin without it, the change due to the absence of the double bond is minimal, limited

to a slight protrusion of the third carbon atom, contributing to the perceived non-essentiality of the

double bond. However, our results conflict with those of Mayer, who reported that the loss of the

double bond could attenuate germination inhibition[7]. While this discrepancy could be due to different

experimental conditions, such as the plant species used, the details remain unclear.

Our research also shows that the effects of the coumarin family depend on one of the lipophilicity

indices, TPSA. Mayer et al. have conducted extensive experiments on the effects of substituent

introduction to coumarin, showing that substituents in the order of NO_2, OH, and Cl may decrease

germination inhibition[7]. This order aligns with the TPSA values (NO_2: 49.36, OH: 20.23, Cl: 0). Given

that two independent studies have arrived at similar results, TPSA is likely a significant factor affecting

the germination inhibitory effect of the coumarin family, regardless of the experimental conditions.

Additionally, TPSA is known to be more closely associated with cell membrane permeability

compared to LogP, another lipophilicity index[30]. Therefore, the introduction of substituents to the

coumarin skeleton might alter germination inhibition by changing cell membrane permeability.

However, it is important to note that discussions regarding TPSA and cell membrane permeability

have primarily focused on mammalian cells.

Furthermore, the ability to quantitatively assess germination inhibition using TPSA paves the way for

more efficient searches for more potent coumarin derivatives. Such derivatives hold promise for

applications in weed control.

Additionally, this study suggests that the seed coat is unlikely to be the primary site of coumarin action.

The seed coat serves as the first line of defense against the external environment and plays a critical

role in relaying environmental conditions to the interior[9]. Furthermore, Aliotta et al. have shown that

the removal of the seed coat significantly diminishes the germination inhibitory effect of coumarin[10].

On the other hand, our analyses of the seed coat's cross-sectional structure and water permeability,

which have a substantial impact on germination, did not reveal any significant effects of coumarin.

While there remains room for further analysis, it seems that the seed coat may not be the main site of

action for coumarin. Additionally, the absorption spectrum of the seed extract suggests that coumarin

penetrates the seed within 24 hours. It is highly probable that coumarin exerts a more direct effect

inside the seed.

Considering the overall results obtained, it is possible that coumarin negatively regulates the

expression of cell wall-associated gene groups, including expansins, and inhibits germination by

obstructing the reorganization of cellulose fibers in the radicle.

Methods

Evaluation of the importance of coumarin substructures in germination inhibition

Coumarin (Tokyo Chemical Industry, Tokyo, Japan) was employed as a positive control. For the

assessment of the benzene ring, α-pyrone (Sigma-Aldrich Japan, Tokyo, Japan) was utilized, while

3,4-dihydrocoumarin (Tokyo Chemical Industry, Tokyo, Japan) was used for the evaluation of the

double bond, and β-tetralone (Sigma-Aldrich Japan, Tokyo, Japan) for the ester bond assessment.

These compounds were diluted with water and prepared at a concentration of 0.10 mmol/L. Thirty

seeds of white clover (*Trifolium repens*) were sown onto each cotton-lined petri dish, and these

solutions were then added. Ion-exchanged water, without the addition of compounds, served as the

negative control. Subsequently, the petri dishes were incubated in darkness at 20°C for 48 hours in an

incubator (A4201, IKUTA Sangyo Co, Ltd, Japan). The number of germinations per petri dish was

counted, and the percentage of germinated seeds out of the total number sown was calculated as the

germination rate. The experiment to investigate the effect of one type of compound was repeated ten

times, using a total of 300 seeds.

Correlation analysis between cheminformatics-estimated indices and germination inhibition

effects

As members of the coumarin family, six compounds were prepared: coumarin (KANTO CHEMICAL

CO., INC., Tokyo, Japan), 3,4-dihydrocoumarin, 7-methoxy coumarin (Tokyo Chemical Industry,

Tokyo, Japan), aesculetin (Apollo Scientific Ltd, Manchester, UK), psoralen (Tokyo Chemical

Industry, Tokyo, Japan), and umbelliferone (FUJIFILM Wako Pure Chemical Co., Osaka, Japan).

These compounds were diluted with water and prepared at a concentration of 0.10 mmol/L, and the

germination rates were calculated in a manner similar to the above. Various indices estimated by

cheminformatics for each compound were calculated using the following methods. The Topological

Polar Surface Area (TPSA), and the octanol/water partition coefficient (LogP) were computed using

SwissADME (http://www.swissadme.ch/)[31]. The molecular surface area and van der Waals volume

were determined using winmostar (V11.5.8, https://winmostar.com/en/). The correlations between

these indices and the germination rates were investigated.

Observation of seed cross-sections with scanning electron microscopy

Pea (*Pisum sativum*) seeds were selected for their ease of observation. Three seeds were immersed for

24 hours in either water or a coumarin solution adjusted to 10 mmol/L within an incubator set to 20°C

under dark conditions. Subsequently, the seeds were spread on cotton-lined Petri dishes, and an

additional 10 mL of the same solution was added. After another 24 hours, the seeds were allowed to

air-dry naturally for 24 hours. The fractured cross-sections of these seeds were then prepared using a

razor blade (FEATHER Safety Razor Co., Ltd) and sputter-coated with Pt-Pd. The coated seeds were

observed under a scanning electron microscope (SEM; S-3400N, Hitachi, Ltd, Osaka, Japan) at an

accelerating voltage of 15 kV and a magnification of 250x. Images of the seeds were captured from

five locations on the cross-sections of the three cleaved seeds. The thickness from the palisade tissue

to the spongy parenchyma of the seeds was measured using ImageJ Fiji (ver. 1.54f,

https://imagej.nih.gov)[32]. Additionally, the thickness of each tissue was measured in the same manner.

Measurement of seed water absorption

Ten pea seeds were permeated in either water or a coumarin solution prepared at 10 mmol/L, and their

surrounding moisture was removed with a paper towel before the fresh weight was measured over

time. Fresh weight measurements were taken hourly from 1 to 24 hours. The seeds were then dried for

over 24 hours in a constant temperature drying oven set to 90°C, and the dry weight was measured.

The water absorption of the seeds from 1 to 24 hours was calculated based on the difference between

each time point's fresh weight and the dry weight.

Detection of coumarin within seed interiors

352   Five pea seeds were immersed in a coumarin solution prepared at 10 mmol/L and were incubated for

353   24 hours in an incubator set at 20°C. Subsequently, the seed coats were delicately scored with a razor

354   blade and removed, after which the inner part of the seed coat was pulverized using a mortar and pestle.

355   The absorption spectrum of 1 mL of this pulverized solution, diluted 100-fold, was measured using a

356   spectrophotometer (PD-3000 Uve, APEL Co., Ltd., Saitama, Japan).

Transcriptome analysis by RNA-Seq

Arabidopsis seeds, known for their well-characterized genetic information, were used. Initially, 20 mg

of seeds were treated with either a 10 mmol/L coumarin solution or water and allowed to incubate for

24 hours. Three biological replicates were used for each condition. The seeds were then manually

crushed with a pestle, and RNA extraction was performed using ISOSPIN Plant RNA (NIPPON GENE,

Japan). Subsequently, RNA concentration was measured using the Qubit RNA BR Assay Kit (Thermo

Fisher Scientific, CA, USA). RNA samples were sent to Novogene (Singapore) for RNA sample

quality testing using a Bioanalyzer (Agilent, California, USA), library construction using the

NEBNext Ultra II RNA Library Prep Kit for Illumina, and sequencing on the NovaSeq 6000 platform

(Illumina, California, USA). The resulting FASTQ data were analyzed using RaNA-seq

(https://ranaseq.eu/)[33], where differential gene expression analysis was conducted based on the

DESeq2 method[34]. In addition, GSEA[35,36] was performed on these DEGs. The GSEA, including

pathway and gene ontology analyses, determined the extent of impact by the normalized enrichment

score (NES). The pathway analysis referenced the KEGG PATHWAY Database, and the gene ontology

analysis was carried out with reference to the database of the National Center for Biotechnology

Information (NIH), analyzing biological process, cellular component, and molecular function domains.

Utilization of generative AI

ChatGPT (https://chat.openai.com/) was employed to rewrite the manuscript into more appropriate

English and to proofread the text. The correctness of the revisions was subsequently verified by the

authors.

References

1. Jain, P. K. & Joshi, H. Coumarin: chemical and pharmacological profile. *J. Appl. Pharm. Sci.* **2**, 236−240 (2012).

2. Bourgaud, F. *et al.* Biosynthesis of coumarins in plants: a major pathway still to be unravelled for cytochrome P450 enzymes. *Phytochem. Rev.* **5**, 293−308 (2006).

3. Takahashi, K., Tsutsumi, Y., Ohtani, H. & Katsuki, T. Variation of fragrance constituents in the leaves of *Prunus*. *Biochem. Syst. Ecol.* **34**, 127−135 (2006).

4. Witt, D. M., Clark, N. P., Kaatz, S., Schnurr, T. & Ansell, J. E. Guidance for the practical management of warfarin therapy in the treatment of venous thromboembolism. *J. Thromb. Thrombolysis* **41**, 187−205 (2016).

5. Bhattarai, N., Kumbhar, A. A., Pokharel, Y. R. & Yadav, P. N. Anticancer potential of coumarin and its derivatives. *Mini-Rev. Med. Chem.* **21**, 2996−3029 (2021).

6. Audus, L. J. & Quastel, J. H. Coumarin as a selective phytocidal agent. *Nature* **159**, 320−324 (1947).

7. Mayer, A. M. & Evenari, M. The relation between the structure of coumarin and its derivatives, and their activity as germination inhibitors. *J. Exp. Bot.* **3**, 246−252 (1952).

8. Araniti, F. *et al*. Phytotoxic potential and biological activity of three synthetic coumarin

derivatives as new natural-like herbicides. *Molecules* **20**, 17883−17902 (2015).

9.   Wu, Y. & Shen, Y. B. Seed coat structural and permeability properties of *Tilia miqueliana* seeds.

J. Plant Growth Regul. **40**, 1198−1209 (2021).

10. Aliotta, G., Fuggi, A. & Strumia, S. Coat-imposed dormancy by coumarin in radish seeds: the

influence of light. *Plant Biosyst.* **126**, 631−637 (1992).

11.   Chen, B. X. *et al.* Coumarin-induced delay of rice seed germination is mediated by suppression

of abscisic acid catabolism and reactive oxygen species production. *Front. Plant Sci.* **10**, 828;

10.3389/fpls.2019.00828 (2019).

12. Chen, B. X., Peng, Y. X., Yang, X. Q. & Liu, J. Delayed germination of *Brassica parachinensis*

seeds by coumarin involves decreased GA_4 production and a consequent reduction of ROS

accumulation. *Seed Sci. Res.* **10**, 224−235 (2021).

13. Araniti, F., Prinsi, B. & Espen, L. The delay of *Raphanus raphanistrum* subsp. *sativus* (L.) Domin

seed germination induced by coumarin is mediated by a lower ability to sustain the energetic

metabolism. *Plants (Basel)* **11**, 843; 10.3390/plants11070843 (2022).

14. Zhang, T. J. *et al.* Metabolic pathways modulated by coumarin to inhibit seed germination and

early seedling growth in *Eleusine indica*. *Plant. Physiol. Biochem.* **203**, 108035;

10.1016/j.plaphy.2023.108035 (2023).

15. Cosgrove, D. J. *et al.* The growing world of expansins. *Plant. Cell Physiol.* **43**, 1436−1444 (2002).

16. Cosgrove, D. J. Plant expansins: diversity and interactions with plant cell walls. *Curr Opin. Plant*

Biol. **25**, 162−172 (2015).

17. Rose, J. K., Braam, J., Fry, S. C. & Nishitani, K. The XTH family of enzymes involved in

xyloglucan endotransglucosylation and endohydrolysis: current perspectives and a new unifying

nomenclature. *Plant. Cell Physiol.* **43**, 1421−1435 (2002).

18. Günl, M. & Pauly, M. AXY3 encodes a α-xylosidase that impacts the structure and accessibility

of the hemicellulose xyloglucan in Arabidopsis plant cell walls. *Planta* **233**, 707−719 (2011).

19. Mutwil, M. *et al.* Transcriptional wiring of cell wall-related genes in *Arabidopsis*. *Mol. Plant* **2**,

1015−1024 (2009).

20. Brett, C. T. Cellulose microfibrils in plants: biosynthesis, deposition, and integration into the. cell

wall. *Int. Rev. Cytol.* **199**, 161−199 (2000).

21. Jones, R. L. & Kaufman, P. B. The role of gibberellins in plant cell elongation. *Crit. Rev. Plant.*

Sci. **1**, 23−47 (1983).

22. Butcher, D. N. & Street, H. E. The effects of gibberellins on the growth of excised tomato roots.

J. Exp. Bot. **11**, 206−216 (1960).

23. Xu, Q., Krishnan, S., Merewitz, E., Xu, J. & Huang, B. Gibberellin-regulation and genetic

variations in leaf elongation for tall fescue in association with differential gene expression

controlling cell expansion. *Sci. Rep.* **6**, 30258; 10.1038/srep30258 (2016).

24. Xu, Q., Burgess, P., Xu, J., Meyer, W. & Huang, B. Osmotic stress- and salt stress-inhibition and

gibberellin-mitigation of leaf elongation associated with up-regulation of genes controlling cell

expansion. *Environ. Exp. Bot.* **131**, 101−109 (2016).

25. Azeez, A., Sane, A. P., Tripathi, S. K., Bhatnagar, D. & Nath, P. The gladiolus GgEXPA1 is a GA-

responsive alpha-expansin gene. expressed ubiquitously during expansion of all floral tissues and

leaves but repressed during organ senescence. *Postharvest Biol. Tec.* **58**, 48−56 (2010).

26. Lee, Y. & Kende, H. Expression of β-expansins is correlated with internodal elongation in

deepwater rice. *Plant Physiol.* **127**, 645−654 (2001).

27. Bruno, L. *et al*. Coumarin interferes with polar auxin transport altering microtubule cortical array

organization in *Arabidopsis thaliana* (L.) Heynh. root apical meristem. *Int. J. Mol. Sci.* **22**, 7305;

10.3390/ijms22147305 (2021).

28. Turi, Z., Lacey, M., Mistrik, M. & Moudry, P. Impaired ribosome biogenesis: mechanisms and

relevance to cancer and aging. *Aging (Albany NY)* **11**, 2512–2540 (2019).

29. Lempiäinen, H. & Shore, D. Growth control and ribosome biogenesis. *Curr. Opin. Cell Biol.* **21**,

855–863 (2009).

30. Silva, A. M. *et al.* In vitro assessment of pesticides toxicity and data correlation with pesticides

physicochemical properties for prediction of toxicity in gastrointestinal and skin contact exposure.

450   *Toxics* **10**, 378; 10.3390/toxics10070378 (2022).

451 31. Daina, A., Michielin, O. & Zoete, V. SwissADME: a free web tool to evaluate pharmacokinetics,

452   drug-likeness and medicinal chemistry friendliness of small molecules. *Sci. Rep.* **7**, 42717;

453   10.1038/srep42717 (2017).

454 32. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

455   analysis. *Nat. Methods* **9**, 671–675 (2012).

456 33. Prieto, C. & Barrios, D. RaNA-Seq: Interactive RNA-Seq analysis from FASTQ files to functional

457   analysis. *Bioinformatics* **36**, 1955–1956 (2020).

458 34. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-

459   seq data with DESeq2. *Genome Biol.* **15**, 550; 10.1186/s13059-014-0550-8 (2014).

460 35. Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for interpreting

461   genome-wide expression profiles. *Proc. Natl. Acad. Sci. USA* **102**, 15545–15550 (2005).

462 36. Mootha, V. K. *et al.* PGC-1α-responsive genes involved in oxidative phosphorylation are

463   coordinately downregulated in human diabetes. *Nat. Genet.* **34**, 267–273 (2003).

Acknowledgments

The authors would like to express gratitude for the technical support provided by Nana Nakahara,

Ayane Hirose, Kenta Araki, Rinsei Ichimaru, Kohei Chifu, Eita Fujito, Kosei Tomonaga, Itsuki

Chiwata, Saori Nakagawa, Syunsuke Saiki, Yuki Takubo, Haruka Mine, Tatsuhiro Tanaka, Kotaro

Yamamoto, Minoru Nishio, Masato Hara, Sota Fukushima, Shion Tanaka and Hikari Kaneshige. RNA

extraction and SEM observation were facilitated at the Analytical Research Center for Experimental

Sciences, Saga University. This work was supported by grants from the Nabeshima-Hokokai and the

Japan Science and Technology Agency under the Super Science High School project of the Ministry

of Education, Culture, Sports, Science and Technology.

Author Contributions

K.F., T.O., S.H., M.Y., and K.M. conceptualized the research plan. S.H., T.O., M.Y., H.M., M.K., and

C.E. collected the germination test data. S.H. and T.O. conducted the cheminformatics analysis. M.Y.,

M.K., K.F., M.N., and N.R. carried out the SEM observations. K.F. performed the measurements of

water absorption. R.I. and K.T. were engaged in the experiments to investigate the spectral properties

of seed extractions. K.F., S.H., T.O., T.Y., and N.R. executed RNA extraction. K.F., S.H., T.O., N.R.,

Y.N., and K.M. conducted the bioinformatics analysis. K.F. and R.I. completed the statistical analysis.

As Ph.D. holders, N.R. and Y.N. provided advisory input to the research. The manuscript was drafted

by K.F., S.H., and T.O., edited by Y.N. The manuscript was finalized by K.F., S.H., T.O., M.Y., N.R.,

Y.N., and K.M. The manuscript was reviewed by all authors.

Data Availability Statement

The raw sequencing data for each sample have been deposited in the DDBJ Sequence Read Archive

(https://www.ddbj.nig.ac.jp/dra/index-e.html; accession numbers DRR546110–DRR546115).

Competing Interests

The authors declare no competing interests.

Figure Legends

Figure 1. Loss of the benzene ring and ester linkage from coumarin attenuates its germination inhibitory effect. (**A–D**) Chemical structures of coumarin-like substances used in this study are shown (coumarin (**A**),α-pyrone (**B**), β-tetralone (**C**), and 3,4-dihydrocoumarin (**D**)). (**E**) Germination rates of white clover *(Trifolium repens)* seeds after 48 hours of incubation with the addition of coumarin-like substances (n = 300). The solutions used in the experiments were prepared at a concentration of 0.10 mmol/L. Below the graph, the presence or absence of the benzene ring, ester bond, and double bond within the lactone ring for each substance is indicated (+ means the structure is present, - means it is absent). Group comparisons of germination rates were made using Chi-squared tests, where the asterisk (*) indicates statistical significance ($p < 0.05$). Error bars represent the 95% confidence interval of the proportion.

Figure 2. A strong correlation is observed between TPSA and the germination inhibition effect. (**A–D**) Structures of additional compounds used in this study, in addition to coumarin and dihydrocoumarin, as shown in Fig. 1 (7-methoxy-coumarin (**A**), aesculetin (**B**), psoralen (**C**), and umbelliferone (**D**)). (**E–H**)Relationships between germination rates of seeds treated with coumarin derivatives and four calculated structural indicators are depicted. The y-axis shows the germination

rates of seeds treated with each member of the coumarin family (n = 300), and the x-axis shows the

calculated indicators from the structure (Molecular Volume (**E**), Molecular Surface Area (**F**), LogP

(**G**), and TPSA (**H**)). Pearson's correlation coefficient r and the p-value from the test of no correlation

are provided. For TPSA, where a significant correlation was observed, a scatter plot with a least-

squares linear regression model is shown.

Figure 3. No observable changes in seed coat structure with coumarin treatment. (**A**) SEM image of

a pea (*Pisum sativum*) seed coat cross-section after 72 hours of treatment with coumarin. Seeds were

sputter-coated with Pt-Pd, observed under an acceleration voltage of 1.5 kV and magnification of 250x.

PAL indicates palisade parenchyma, HOU represents hourglass cells, and COM signifies compressed

parenchyma. Scale bar = 100 µm. (**B**) Thickness measurements of pea seed coat tissues treated with

coumarin. Measurements were obtained from images captured using the same method as Fig. 3A and

analyzed with ImageJ Fiji software. "All" indicates the total thickness from palisade parenchyma to

compressed parenchyma. The thickness of each tissue was compared between coumarin-treated and

untreated seeds using Student's t-test. n.s. denotes no statistical significance (p > 0.05), and error bars

represent the 95% confidence interval for the mean. (**C**) Water absorption of pea seeds from 1 to 24

hours post-immersion. Absorption was calculated based on the difference between wet weight

measured post-absorption and dry weight after drying. No significant differences were observed at any

time point (p > 0.05). (**D**) Absorption spectra of homogenates from ten pea seeds immersed in a

coumarin solution for 24 hours. Experiments were conducted after removing seed coats using a 10

mol/L coumarin solution, replicated three times. Coumarin's maximum absorption wavelength at 270

nm is indicated with a dashed line. Absorbance at 270 nm was compared between treatments with and

without coumarin using Student's t-test, with the asterisk (*) indicating statistical significance (p <

0.05).

Figure 4. Down-regulation in genes related to ribosomes and expansins by coumarin. (**A–B**) The

heatmaps display the expression ratio of RPL family (**A**) and RPS family genes (**B**) between coumarin-

treated and control groups obtained through RNA-Seq. Experiments were replicated thrice for each

group, with columns corresponding to the expression profiles of individual samples. (**C**) The bar plot

represents the log2-transformed expression ratios of expansin family genes between the coumarin-

treated and control groups. The asterisk (*) denotes statistical significance based on the adjusted p-

value (padj<0.05).

Figure 5. Coumarin Addition Impacts Multiple Pathways. (**A**) Rug plots and enrichment score walk

544 from GSEA based on the KEGG Pathways database, showing the maximum distance from zero along

545 the walk, calculated as the enrichment score. This analysis reveals the impact of coumarin on eight

546 pathways in Arabidopsis, with particularly interesting data for four pathways shown. (**B–D**) Bar plots

547 display the normalized enrichment score (NES) from GSEA using the NIH Gene Ontology database.

548 The top five activated and repressed pathways are shown for each gene ontology category (biological

549 process (**B**), cellular component (**C**), and molecular function (**D**)). (**E–H**)Bar plots show the log2

550 expression ratio between the coumarin-treated and control groups for genes of the XYL1 and XTH

551 families (**E**), the sks family (**F**), GA20ox1/3 genes (**G**), and the PIN family genes (**H**). The asterisk

552 (*) indicates statistical significance based on the adjusted p-value (padj < 0.05).

554   **Table 1.** GSEA Pathway Analysis Results

Pathway_ID	Pathway	NES	padj
ath03010	Ribosome	-2.409	0.011
ath_M00177	Ribosome, eukaryotes	-2.290	0.011
ath_M00049	Adenine ribonucleotide biosynthesis, IMP => ADP,ATP	-1.812	0.011
ath_M00087	beta-Oxidation	1.969	0.011
ath00480	Glutathione metabolism	2.004	0.013
ath_M00113	Jasmonic acid biosynthesis	2.053	0.018
ath00945	Stilbenoid, diarylheptanoid and gingerol biosynthesis	1.902	0.024
ath00230	Purine metabolism	-1.674	0.049