Multimodal identification of the mouse brain using simultaneous Ca^{2+} imaging and fMRI

Francesca Mandino[1]*, Corey Horien[2,3]*, Xilin Shen[1], Gabriel Desrosiers-Gregoire[4,5], Wendy Luo[6], Marija Markicevic[1], R.
Todd Constable[1,2,3,6,7], Xenophon Papademetris[1,6,8], Mallar M. Chakravarty[4], Richard F. Betzel[9], Evelyn M.R. Lake[1,6,10]

[1]Department of Radiology and Biomedical Imaging, Yale University School of Medicine, New Haven, CT, USA
[2]Interdepartmental Neuroscience Program, Yale University School of Medicine, New Haven, CT, USA
[3]MD/PhD program, Yale University School of Medicine, New Haven, CT, USA
[4]Computational Brain Anatomy Laboratory, Douglas Mental Health University Institute, Montreal, Canada
[5]Integrated Program in Neuroscience, McGill University, Montreal, QC, Canada
[6]Department of Biomedical Engineering, Yale University School of Medicine, New Haven, CT, USA
[7]Department of Neurosurgery, Yale University School of Medicine, New Haven, CT, USA
[8]Deparment of Biomedical Informatics and Data Science, Yale University, New Haven, CT, USA
[9]Department of Psychological and Brain Sciences, Indiana University
[10]Wu Tsai Institute, Yale University, New Haven, CT, USA

Corresponding authors: francesca.mandino@yale.edu, evelyn.lake@yale.edu

* Co-first authorship

Keywords: functional connectivity, human, connectome, fingerprinting, GCaMP, resting-state, wide-field fluorescence
imaging.
18

ABSTRACT

Individual differences in neuroimaging are of interest to clinical and cognitive neuroscientists based on their potential for guiding the personalized treatment of various heterogeneous neurological conditions and diseases. Despite many advantages, the workhorse in this arena, BOLD (blood-oxygen-level-dependent) functional magnetic resonance imaging (fMRI) suffers from low spatiotemporal resolution and specificity as well as a propensity for noise and spurious signal corruption. To better understand individual differences in BOLD-fMRI data, we can use animal models where fMRI, alongside complementary but more invasive contrasts, can be accessed. Here, we apply simultaneous wide-field fluorescence calcium imaging and BOLD-fMRI in mice to interrogate individual differences using a connectome-based identification framework adopted from the human fMRI literature. This approach yields high spatiotemporal resolution cell-type specific signals (here, from glia, excitatory, as well as inhibitory interneurons) from the whole cortex. We found mouse multimodal connectome-based identification to be successful and explored various features of these data.

1. INTRODUCTION

Individual differences are of growing interest in clinical and cognitive neuroscience[1,2], joining more

established examples of precision medicine, including personalized treatments in oncology[3] and cardiovascular

disease[4]. For nearly a decade, we have known that functional connectivity (FC) by means of non-invasive BOLD

(blood-oxygen-level-dependent) functional magnetic resonance imaging (fMRI) exhibits individual-specific

features that make a person's brain distinguishable from others[5]. This framework identifies subjects from a pool

using their connectome (a summary of inter-regional BOLD signal (a)synchronies or connectivity strengths). High

rates of correct identification (ID) are typical in human populations (e.g., 88% in **Fig. 1**)[5-17].

Fig. 1 | Connectome ID framework using random nodes and functional networks in humans. *Connectomes were*
computed for human subjects from the Human Connectome Project[18] using the Shen-268 atlas following established
methods[5,19](see 'Human BOLD-fMRI data' in Methods). ***A.*** *Connectome-to-connectome correlation (Pearson) across*
imaging sessions. The clear diagonal indicates that connectomes from the same individual (collected during different
sessions) are more similar (have a higher correlation value) than data from unmatched individuals (in-line with previous
work). ***B.*** *The Shen-268[5] atlas has been sub-divided into 8 functional networks: (1) frontoparietal, (2) medial frontal, (3)*
visual, (4) salience, (5) subcortical, (6) cerebellum, (7) motor, and (8) default mode. These networks contain 20-50 nodes
each. We repeated a 'winner-take-all' ID procedure using each functional network, and random selections of 'x' nodes (N =
100 iterations). 'x' was varied between 5 & 65 in increments of 5. As the number of randomly selected nodes increased, the
ID rate also increased (reaching a plateau by approximately 40 nodes, or only 15% of all nodes). Relative to collections of
randomly selected nodes, functional networks showed variable ID rates. Some performed worse (e.g., the subcortical and
cerebellum networks), while others performed better (e.g., medial frontal and frontoparietal networks) than expected based
on their size. Thus, as has been shown previously[5,13], we affirmed that ID rates using BOLD-fMRI data from humans required
only a small fraction of the whole-brain and showed clear functional network dependence.
80

81   This discovery garnered widespread attention by spawning the notion that clinically accessible measures of

82   brain connectivity could be used to guide the personalized care of various neurological conditions and diseases[20].

83   Yet, to-date, there are no applications within the clinic. In part, this lack of translation is due to our limited

understanding of the biological basis for BOLD-fMRI measures of brain connectivity, which stem from the BOLD signal being a relatively low-spatiotemporal resolution summary metric of brain activity accessed via hemodynamic coupling to neural activity.

To address knowledge gaps, BOLD-fMRI applied in animal models allows for invasive techniques (not applicable in humans[21]) to offer complementary measures of brain activity, and the opportunity for manipulations that can help to determine causal relationships[22,23]. Despite this potential (and some promising leads[24,25]), animal studies investigating individual differences are rare. Instead, animal studies have typically focused on group averages, which leverages the tight control animal work has over environment and genetic variability but fails to capture the clear heterogeneity that is evident in the human conditions being modeled[26-28]. Encouragingly, recent pioneering work by Bergmann et al. in 2020 established that laboratory mice, despite being homogenized by design, are identifiable using the same BOLD-fMRI connectome-based ID framework as has been used on human BOLD-fMRI data[5,29]. To the best of our knowledge, this finding has not been replicated or investigated further, until the present work.

We use a unique simultaneous wide-field calcium (WF-Ca^{2+}) and BOLD-fMRI acquisition system[30] to investigate connectome-based ID in mice using multimodal neuroimaging data. To gain insight into relationships between the BOLD signal and the underlying cellular activity, the use of optical modes, which report on brain activity using genetically encoded or virally transfected fluorescent indicators (e.g., GCaMP), yields a complementary high specificity and spatiotemporal resolution signal source[31,32]. Here, we implement simultaneous WF-Ca^{2+} and BOLD-fMRI in healthy adult transgenic mice (N = 45) expressing GCaMP in one of five cell types (herein called 'groups'): (1) excitatory neurons ('SLC'), (2) glia, as well as inhibitory interneurons expressing (3) parvalbumin ('PV'), (4) somatostatin ('SOM'), or (5) vasoactive intestinal (poly)peptide ('VIP'). Mice were imaged three times (~7-days apart) under low-dose isoflurane anesthesia.

This is the first study to apply a connectome based-ID framework using simultaneously obtained multimodal data. That we found significant rates of correct IDs in both modalities affirms that connectome-based ID is likely driven by cellular activity, rather than noise. We also found preliminary evidence for differing rates of ID across canonical networks, echoing observations made in human BOLD-fMRI studies (**Fig. 1**)[5,8,12]. The group structure of our dataset (based on cell-type expressing GCaMP) allowed us to investigate group membership in addition to individual mouse ID rates. Data from both modalities resulted in significant group IDs – with an expected

discrepancy in performance (WF-Ca^{2+} outperforming BOLD-fMRI). Differing group ID rates, in WF-Ca^{2+} imaging data, as well as clear structure in overall connectome (dis)similarity across groups (and networks), highlights the power of optical imaging measures – specifically at the 'mesoscale' – to uncover cell-type specific network structure. We also investigated inter-modal connectome-based ID and found that it did not result in above chance-level ID rates. These observations illustrate the complementarity of WF-Ca^{2+} and BOLD-fMRI and, to an extent, a limitation of coarse static measures like connectome similarity.

Altogether, our work affirms using connectome based-ID as a framework for studying individual differences in mouse models. It highlights the power of multimodal methods, that can only be applied in animals, for informing on clinically accessible imaging modes within a framework adopted from the human fMRI literature. Through studying individual differences, across species, neuroimaging can move closer to becoming a means through which to conduct personalized medicine.

2. RESULTS

Simultaneous WF-Ca^{2+} and BOLD-fMRI data were acquired from free-breathing lightly anesthetized (0.5-0.75% isoflurane) mice (N = 45, **Table S1**) as we have described previously[30,33] (**Methods**). To acquire WF-Ca^{2+} imaging data, mice underwent a minimally invasive surgical procedure (7-days prior to the first imaging session, **Methods**) where the skull was exposed and thinned to translucency (but left intact) and a custom glass and plastic head-plate was affixed to the bone for permanent optical access to the cortical surface and immobilization during imaging. All (N = 45) mice underwent three multimodal imaging sessions, with a minimum of 7-days between each session (**Fig. 2A**, left), where structural (for image registration) and functional (BOLD) MRI data were acquired (**Methods**). WF-Ca^{2+} imaging data were acquired simultaneously with the BOLD-fMRI data while mice were in the 'resting-state' (i.e., no exogenous stimuli were presented). Forty-minutes of functional multimodal data were acquired at each imaging session across four 10-minute runs. Data were processed using BioImageSuite (BIS) and RABIES (Rodent Automated Bold Improvement of EPI Sequences)[34] software and registered to an in-house template that has been registered to the Allen Institute Common Coordinate Framework Reference Atlas (CCFv3)[35] (**Fig. 2B**, **Methods**), as we have described previously[30,33,36].

A. Experimental timeline and data acquired included in all analyses

GLIA   Aldh1l1-Cre/Ai162, glia, N=6 mice

PV   PV-Cre/Ai162, interneurons, N=9 mice

SLC   Slc17a7/Cam/TeTG6f, excitatory neurons, N=9 mice

SOM   Som-Cre/i162, inhibitory neurons, N=9 mice

VIP   Vip-Cre/Ai162, interneurons, N=12 mice

B. Framework for multimodal data registration and registration to common-space

Fig. 2 | Dataset overview and multimodal data registration framework. A. *Experiment timeline (left top), group breakdown (left bottom), and number of 10-minute runs included across groups in our analyses (right).* **B.** *Multimodal registration framework. Structural and functional MR-images (peach and orange), from each mouse at each acquisition (imaging session), were registered to an in-house reference space (****Methods****). This reference space (created from N = 162 datasets collected on our scanner, see [36] has been registered to the Allen Institute CCvf3 and Atlas [35] (pink). We use an MRI time-of-flight angiogram (yellow) to capture blood vessels on the cortical surface that also appear in the static WF-Ca2+ imaging data (green). These serve as anatomical landmarks to link across imaging modalities, as described previously[30] (****Methods****). Together, these transformations (black arrows) enable us to move atlases, regions of interest, or data from individual to common space and between 2 and 3 dimensions (using forward or backward projection, see [30]. All transformations were concatenated before being applied. The tools to create all transformations, do projections, the atlases, and our in-house reference space are all freely available (https://bioimagesuiteweb.github.io/webapp/ or [34]).*

Stringent data inclusion criteria were applied (**Methods**) such that a minimum of 20-minutes of functional

data (or two runs) were retained per mouse per session (**Fig. 2A**, right & **Table S1**). Data were also matched

across modalities such that all the WF-Ca^{2+} and BOLD-fMRI data in our analyses were temporally matched. For

WF-Ca^{2+} imaging data, the majority of the power tended to be concentrated in the infra-slow range and aligned

with previous work[37] (**Fig. S1; Table S2 and S3**). Hereon, we explored a slow and a fast WF-Ca^{2+} imaging

frequency band: (1) 0.008-0.2Hz, WF-Ca$^{2+}_{Slow}$ and (2) 0.4-4Hz, WF-Ca$^{2+}_{Fast}$, as we have done previously[37]. The

WF-Ca$^{2+}_{Slow}$ band was chosen to match the BOLD-fMRI data.

Connectomes were computed (using frequency filtered data), using the product-moment correlation, for

the whole-brain (BOLD-fMRI only) and cortex (BOLD-fMRI and WF-Ca^{2+} imaging data) as the WF-Ca^{2+} imaging

has a limited field-of-view (FOV). Thus, we consider data in four 'conditions': (1) WF-Ca$^{2+}_{Slow}$, (2) WF-Ca$^{2+}_{Fast}$,

(3) BOLD-fMRI cortex-only, and (4) BOLD-fMRI whole-brain. For the BOLD-fMRI whole-brain condition, we used

172 regions from the Allen Atlas (**Fig. S2**). Forty-four of these were 'visible' from above (in all N = 45 mice at all

sessions) and comprise the regions within our cortical connectomes or remaining three 'conditions'. The

multimodal data registration pipeline and images of both the whole-brain and cortex-only atlases are shown in

Fig. 2B. In all subsequent figures we indicate the imaging modality, FOV (whole-brain or cortex), and 'condition'

using black-and-white icons.

2.1 Average connectomes showed group and condition-specific features.

Connectomes were computed at the level of each run (i.e., using 10-minutes of WF-Ca^{2+} or BOLD-fMRI

data) with data residing within our in-house reference space. After data inclusion criteria were applied (**Methods**),

we retained N = 437 connectomes per condition. All following analyses were performed on these N = 437

connectomes as well as the within-session, within mouse, averaged connectomes (i.e., averaging across 2-4

run-level connectomes per mouse per session which yielded N = 135 connectomes). None of our findings were

affected by using run-level or averaged connectomes. As an overview of these data, 'grand' average

connectomes for each group and condition were computed (**Fig. 3A**). Data were averaged across all mice, runs,

and sessions (within group and condition). See corresponding standard deviations (SD) in **Fig. S3**. As expected,

WF-Ca^{2+} imaging data showed universally high correlation strengths.

A. 'Grand' average connectomes (all mice, all sessions, averaged within group and condition)

B. Difference between 'grand' average connectomes and the dataset average (all mice, all sessions, & all groups)

All groups and conditions showed clear bilateral symmetry which indicated greater connectivity between homotopic regions. To highlight inter-group differences, within condition, the difference between each 'grand' average connectome (**Fig. 3A**) and the dataset average (all mice, runs, sessions and groups) was computed (**Fig. 3B**). For this calculation, the number of runs included from each group was held constant (N = 56 per group). As anticipated, GLIA WF-Ca^{2+}$_{Slow}$ and WF-Ca^{2+}$_{Fast}$ were the most different from the dataset average. This group exhibited globally higher connectivity strengths than excitatory (SLC) ($p=9.2e^{-48}$ and $p=8.9e^{-107}$ for WF-Ca^{2+}$_{Slow}$ and WF-Ca^{2+}$_{Fast}$, respectively) or inhibitory interneuron (PV, SOM, and VIP) groups ($p=7.0e^{-76}$ and $p=4.1e^{-184}$ for WF-Ca^{2+}$_{Slow}$ and WF-Ca^{2+}$_{Fast}$, respectively) (MATLAB, *ttest2*). The BOLD-fMRI data showed more inter-group features than expected. For example, high correlation strengths between cortical areas within SLC (relative to the dataset average), and opposing patterns between PV and SOM.

2.2 Inter-connectome, intra- and inter-condition correlations showed group and condition-specific features.

Inter-connectome correlations were computed within (e.g., WF-Ca^{2+}$_{Fast}$ to WF-Ca^{2+}$_{Fast}$) and between (e.g., WF-Ca^{2+}$_{Slow}$ to BOLD-fMRI) conditions across sessions (e.g., session 1 → 2) or within session, respectively. Within condition comparisons were computed across time, while between condition comparisons were computed using simultaneously acquired multimodal data (**Fig. 4**). WF-Ca^{2+} connectomes were compared within matched frequency bands (slow and fast in **Fig. 4A** & **B**, respectively) and to BOLD-fMRI cortex-only data. Data were organized by group (top-to-bottom: GLIA, PV, SLC, SOM, & VIP) to highlight between group features. For example, GLIA (topmost) and VIP (bottommost) showed lower within and between condition inter-connectome correlation strengths relative to other groups for the WF-Ca^{2+}$_{Fast}$ condition (**Fig. 4B**). Specifically, for session 1 → 2, within condition (WF-Ca^{2+}$_{Fast}$), GLIA vs. GLIA relative to all other within group pairs $p=6.7e^{-58}$ and GLIA vs. any other group relative to all other between group pairs p~0; whilst, within session 1, between condition (WF-Ca^{2+}$_{Fast}$ vs. BOLD-fMRI), GLIA vs. GLIA relative to all other within group pairs $p=2.4e^{-94}$ and GLIA vs. any other group relative to all other between group pairs $p=5.4e^{-204}$ (MATLAB, *ttest2*). Likewise, for session 1 → 2, within

condition, VIP vs. VIP relative to all other within group pairs p=4.8e^{-181} and VIP vs. any other group relative to all

other between group pairs p=1.7e^{-79}; whilst, within session 1, between condition, VIP vs. VIP relative to all other

within group pairs p=2.8e^{-22} and VIP vs. any other group relative to all other between group pairs p=5.3e^{-5}

(MATLAB, *ttest2*). Further, correlation magnitude was on average stronger within WF-Ca^{2+} signals than BOLD-

fMRI (as an example, for session 1 → 2, within Ca$^{2+}_{Fast}$ vs. within BOLD-fMRI, p~0, MATLAB *ttest2*).

A. Within and cross-condition connectome-to-connectome correlation (WF-Ca$^{2+}_{Slow}$ and cortex-only BOLD-fMRI)

B. Within and cross-condition connectome-to-connectome correlation (WF-Ca$^{2+}_{Fast}$ and cortex-only BOLD-fMRI)

Fig. 4 | Within and between condition inter-connectome correlation. *Inter-connectome correlation strengths within (upper left and lower right quadrants) and between (upper right and lower left quadrants) conditions using data from within or between sessions, respectively, such that within condition comparisons were across time (i.e., from different sessions), and between condition comparisons used simultaneously collected multimodal data. Each row/column in the matrices is a single run.* ***A.*** *WF-Ca$^{2+}_{Slow}$ data and BOLD-fMRI cortex-only data.* ***B.*** *WF-Ca$^{2+}_{Fast}$ data and BOLD-fMRI cortex-only data. Inter-connectome comparisons were organized by group (top-to-bottom: GLIA, PV, SLC, SOM, & VIP) as indicated on the far right.*

2.3 All intra-condition ID rates of mouse and group were above chance-level with WF-Ca^{2+} outperforming BOLD-fMRI and WF-Ca$^{2+}_{Fast}$ outperforming WF-Ca$^{2+}_{Slow}$.

For intra-condition ID analyses, of mouse (**Fig. 5A** & **C**) or group (**Fig. 5B** & **D**), all six possible inter-session pairings (i.e., session 1→2, as well as 2→1) were considered. A correct ID was a binary decision based on whether the connectomes that were most similar (highest inter-connectome correlation strength) between sessions belonged to the same mouse or group. For group ID, the mouse being queried was removed from the target session to avoid bleed-through of mouse ID into group ID. As in our analysis of human BOLD-fMRI data (**Fig. 1**), subsets of randomly selected nodes were used to perform mouse or group ID (N = 50 iterations, colored dots in **Fig. 5A** & **B**). These ranged from 5 to all nodes in our cortex-only (N = 44) or whole-brain (N = 172) atlases. Statistically significant ID rates were computed relative to a null distribution generated by shuffling mouse or group ID labels in the target session (N = 50 iterations, greyscale dots in **Fig. 5A** & **B**). All intra-condition mouse and group ID rates quickly rose above chance-levels (**Fig. 5A** & **B**). Only five nodes were needed for WF-Ca$^{2+}_{Fast}$ (p=1.9e^{-186} and p=1.7e^{-222}, for mouse and group ID respectively), WF-Ca$^{2+}_{Slow}$ (p=1.8e^{-146} and p=1.6e^{-165}), BOLD-fMRI cortex-only (p=9.5e^{-31} and p=6.0e^{-42}), and BOLD-fMRI whole-brain (p=7.4e^{-27} and p=2.2e^{-26}) (MATLAB, *ttest2*). Similarly, in all cases, correct mouse and group ID rates appear to reach a plateau when approximately half of all available nodes were being considered (based on a visual inspection of the results, **Fig. 5A** & **B**). A subtle difference in WF-Ca^{2+} imaging-based ID rates was noted across session pairings. Specifically, longer inter-session time (sessions 1 & 3, yellow) outperformed sessions that were closer in time (sessions 1 & 2 and 2 & 3, green and pink). For example, for 40/44 cortical nodes, sessions 1 & 3 relative to the other two pairings performed better: p=5.6e^{-52} and p=1.1e^{-25} for mouse and group ID, respectively, for WF-Ca$^{2+}_{Slow}$, p=7.9e^{-4}, and WF-Ca$^{2+}_{Fast}$, p=0.02 (MATLAB, *ttest2*). For both mouse and group ID, WF-Ca$^{2+}_{Fast}$ performed better than WF-Ca$^{2+}_{Slow}$ (for 40/44 cortical nodes, p=1.3e^{-35} and p=8.4e^{-176} for mouse and group ID, respectively, MATLAB *ttest2*). Overall, higher mouse and group ID rates were observed when WF-Ca^{2+} results were compared to BOLD-fMRI results (for 40/44 cortical nodes, relative to WF-Ca$^{2+}_{Slow}$, p=2.5e^{-79} and p=5.7e^{-279} for mouse and group ID, respectively, while relative to WF-Ca$^{2+}_{Fast}$, p=1.1e^{-163} and p~0, MATLAB *ttest2*). Notably, this discrepancy was more modest than anticipated for mouse ID rates; especially given the inherent group differences that are backed-into the WF-Ca^{2+} imaging data. Accordingly, group ID rates based on WF-Ca^{2+}

249 imaging data were substantially greater than those based on BOLD-fMRI data (above). However, the still clearly

250 significant group ID rates achieved with BOLD-fMRI data were noteworthy.

Fig. 5 | Intra-condition mouse and group ID. A. *Mouse ID rates for all conditions (left-to-right: WF-Ca$^{2+}_{Slow}$ WF-Ca$^{2+}_{Fast}$, BOLD-fMRI cortex-only, and BOLD-fMRI whole-brain).* **B.** *Group ID rates for all conditions (ordered as in* **A.***).* *In* **A.** *and* **B.**, *colored dots (green, yellow, & pink) – from intact data – were compared to a null distribution (greyscale dots) where the mouse or group ID labels in the target session were shuffled (N = 50 iterations). For group ID (***B.***), the mouse being queried was removed from the target session (to avoid mouse ID bleed-through). As in* **Fig. 1***, mouse or group ID was repeated*

*using randomly selected nodes ranging from 5 to all-nodes within the atlas (N = 50 iterations). In all cases, intact data outperformed shuffled data. As in **Fig. 1**, mouse **C.**, and group **D.**, IDs from random selections of nodes were compared to results obtained with a priori functional networks. See **Fig. S2** for 2D and 3D network node definitions. Thus, colored dots in **A.** and **B.** (of appropriate size) are re-plotted in greyscale in **C.** and **D.** alongside the results from canonical networks (in color). For reference, column one, in all subplots in **C.** and **D.**, were results from all nodes (color) and shuffled data (greyscale) – replicas of the rightmost columns in all subplots within **A.** and **B.** – to bound expected high and low rates of ID for each condition. Statistically significant results in **C.** and **D.** (two-tailed, MATLAB t-test, ttest2, Bonferroni corrected) were indicated by an *.*

Finally, differences in mouse and group ID rates between cortex-only and whole-brain BOLD-fMRI data were

small (despite still being significant for 60 or more nodes, $p<3.5e^{-5}$, MATLAB *ttest2*) which indicated that

subcortical areas played a limited role in ID success rate (as anticipated based on work in human subjects, **Fig.**

1).

For both mouse and group, we computed whether correct IDs were shared across conditions (**Fig. S4**). For

example, if mouse-1 was correctly matched in the WF-Ca^{2+}$_{Slow}$ calculation (leftmost panel, **Fig. 5A**), was mouse-

1 also correctly matched in the BOLD-fMRI whole-brain calculation (rightmost panel, **Fig. 5A**)? Both intra- and

inter-modality comparisons were compared to a null distribution generated by shuffling the correct hit labels from

one of the two calculations (N = 50 iterations). Overall, correct IDs of mouse and group were not shared when

results from different modalities were compared (**Fig. S4A** & **B**). Conversely, correct IDs of mouse and group

showed consistent above chance-level subject overlap within modality (**Fig. S4C** & **D**).

2.4 Intra-condition ID rates of mouse and group showed *a priori* functional network dependence.

Human BOLD-fMRI data-based connectome ID shows a strong dependence on *a priori* functional networks

(**Fig. 1**)[5,7]. To search for parallels in mouse data, we investigated 12 whole-brain (BOLD-fMRI) *a priori* canonical

networks (somatosensory, auditory, visual, olfactory, sensory, cortex, limbic, lateral cortical, DMN, salience,

subcortical, and brainstem). Six of these (somatosensory, visual, sensory, limbic, lateral cortical, and DMN) were

'visible' from above (**Fig. S2**) and used in our WF-Ca^{2+} imaging and BOLD-fMRI cortex-only analyses.

As in **Fig. 1**, we determined whether canonical networks played a disproportionate role in mouse or group

ID by comparing ID rates obtained when a network was used to random size-matched selections of nodes (N =

50 iterations) (**Fig. 5C** & **D**). For WF-Ca^{2+} imaging data, none of the canonical networks performed better than

this benchmark for mouse or group ID. Somewhat surprisingly, several canonical networks under-performed for

group ID (**Fig. 5D**). Conversely, the BOLD-fMRI data showed some parallels to results obtained using human

BOLD-fMRI data. The whole-brain lateral cortical network (a proxy for the human frontal parietal network) and

somatosensory network produced greater mouse ID rates while the subcortical and brainstem networks under-

performed (**Fig. 5C**). Notably, the nodes in the lateral cortical and somatosensory networks that were 'visible'

from above and were thus included in our BOLD-fMRI cortex-only analysis, did not produce the same results as

in our whole-brain analyses. These observations suggest that connections to regions outside of the WF-Ca^{2+}

imaging FOV may play a disproportionately important role in increasing mouse ID rates and could help explain

why we did not see similar results in the WF-Ca^{2+} imaging data.

2.5 Intra-condition ID rates depended on *a priori* canonical network and group (e.g., from WF-Ca$^{2+}_{Fast}$).

The preceding subsection considered all groups together. Here, we searched for differences in mouse

ID rates across groups (e.g., are SLC more identifiable than GLIA?) and structure within the incorrect IDs. For

example, are GLIA that were incorrectly identified most similar to others in the GLIA group or are they frequently

misidentified as members of another group (e.g., PV)? For this aim, we interrogated a range of thresholds for

"top-hits" (i.e., most similar inter-session intra-contrast connectomes) in-place of selecting only one top-hit (as

done in the preceding subsections). We identified the top 5, 15, or 25 most correlated inter-session connectome

pairs (when considering runs, N = 146/144/147 for sessions 1/2/3), and the top 2, 4, or 6 most correlated inter-

session connectome pairs (when considering averaged runs, N = 45 per session). An example is shown in **Fig.**

6A for WF-Ca$^{2+}_{Fast}$ (sessions 1 & 2), all nodes. We quantified the binarized "top-hit" matrices (**Fig. 6A**, right) as

the fraction of hits within each area of interest where the areas of interest were each within- or between-group

pairing (e.g., GLIA-to-GLIA, indicated in **Fig. 6**). The fraction of top-hits within each area of interest was

expressed as a matrix (**Fig. 6B**) for all session pairs, thresholds, conditions, and canonical networks. As above,

a null distribution was generated by shuffling the target session labels (N = 50 iterations).

The example matrices in **Fig. 6B** were organized into three clusters to highlight three patterns. First, all

nodes, visual and sensory networks (top row) showed a strong diagonal which indicated high within-group ID

rates (although, VIP performed poorly relative to the rest) and low between group incorrect IDs. The second

pattern (bottom row, left), evident in the somatosensory and lateral cortical networks, showed low GLIA ID rates

(GLIA being incorrectly identified as SOM), reasonable PV, SLC, and SOM ID rates – albeit with frequent errors

between PV and SLC as well as between SOM and VIP – and, as in the first pattern, low correct VIP ID rates.

In the final pattern, limbic and DMN, showed low (but still significant) within-group ID rates with more diffuse

errors (than seen in the second pattern). Results were summarized further, across session pairs and canonical

networks, using boxplots (**Fig. 6C**).

The example results in **Fig. 6C** (all runs, threshold 25, WF-Ca$^{2+}_{Fast}$, GLIA and VIP) showed that where 'top-

hits' were found, had group-pairing and network dependence. For instance, limbic and default mode networks,

showed poor correct GLIA ID rates (same as chance) (**Fig. 6C**, top). Conversely, VIP (**Fig. 6C**, bottom) showed

more modest correct ID rates without clear differences between networks. In terms of where incorrect IDs

resided, GLIA were unlikely to be misidentified as PV or SLC (below chance) but were mistaken for SOM or VIP

at chance-level (mostly regardless of network). Similarly, VIP were unlikely to be misidentified as SLC, or PV –

albeit to a lesser extent – and were mistaken for GLIA at chance-level. VIP were, however, misidentified as SOM

at greater than chance-level (with some network dependence). These observations, and others for remaining

groups, held across 'top-hit' thresholds, and the averaging of runs.

2.6 Groups showed some consistent cross-condition network and inter-group ID patterns.

The analyses outlined in **subsection 2.5** were performed on all groups using data from all conditions.

Representative results from all four conditions (for GLIA, SLC, and VIP) are shown in **Fig. 7**. Results from PV

and SOM are shown in (**Fig. S5**). WF-Ca$^{2+}_{Slow}$ and WF-Ca$^{2+}_{Fast}$ conditions showed nearly identical results across

groups and canonical networks. SLC, followed closely by PV, showed the highest fraction of correct IDs with

GLIA, SOM, and finally VIP showing progressively lower (but still above chance) fractions of correct IDs. Overall,

limbic and default mode networks performed poorly, while all nodes and the sensory network performed best.

SOM and PV were more likely to be misidentified for one another than either were to be misidentified as SLC.

Similarly, PV and SLC showed some propensity for being misidentified for one another; specifically, when the

somatosensory and lateral cortical networks were being considered. Finally, it was very unlikely for GLIA to be

misidentified as PV or SLC (and vice versa).

Fig 7 | Network influence on intra-condition ID separated by group (for GLIA, SLC, and VIP). *Boxplots are constructed as described in* ***Fig. 6C*** *(topmost left and right panels were replicated from* ***Fig. 6C***). *Results for PV and SOM were shown in* ***Fig. S5***. ***A.*** *Results (top-to-bottom) for WF-Ca$^{2+}_{Slow}$, WF-Ca$^{2+}_{Fast}$, and BOLD-fMRI cortex-only. Six canonical networks, in addition to all ROIs.* ***B.*** *Results for BOLD-fMRI whole-brain. Twelve canonical networks, in addition to all ROIs. For* ***A.*** *and* ***B.*** *group pairings where a statistically significant (two-tailed, MATLAB t-test, ttest2, Bonferroni corrected) surplus, or paucity, of 'top-hits' were found were indicated by an* *.*

Cortex-only BOLD-fMRI results mirrored some features that were evident in the WF-Ca^{2+} imaging data. As in the WF-Ca^{2+} imaging conditions, SLC showed the highest fraction of correct IDs. Although PV were still above chance, they performed more poorly, and not as well as GLIA. SOM were still above chance-level, but VIP were not. These findings roughly paralleled what was seen in the WF-Ca^{2+} imaging conditions. Network effects in the

cortex-only BOLD-fMRI condition, were only evident in SLC, and showed that visual and limbic networks performed poorly. Similar patterns, with more clear network dependence, were evident in the whole-brain BOLD-fMRI condition where we considered 12 canonical networks (and all nodes). Here, auditory, visual, and olfactory performed poorly, as did limbic, default mode, salience, subcortical, and brainstem, relative to somatosensory, sensory, cortex, the lateral cortical network, and all nodes. Findings were consistent across thresholds and the averaging of runs.

2.7 Cross-modality instances of high connectome-to-connectome correlation strengths showed group and network dependence.

Overall, our findings were consistent with our expectations given the appearance of the matrices in **Fig. 4**. Inter-condition (top left, and bottom right quadrants) comparisons showed the classic diagonal structure which indicated that the highest connectome-to-connectome correlations were (on average) between the same animals. The block diagonal structure also indicated that within-group errors were likely to be more frequent than between-group errors. This was abundantly clear when a threshold was applied (**Fig. 6A**). Notably, the same pattern was not evident when connectomes were compared across imaging modalities (top right, and bottom left quadrants in **Fig. 4**), which led to the unsurprising result that when we performed cross-modality ID (BOLD-fMRI cortex-only to WF-Ca^{2+}$_{Fast}$ or WF-Ca^{2+}$_{Slow}$), using the whole dataset in a 'winner-take-all' framework we found no significant results (data not shown). Nevertheless, we pursued a more subtle interrogation of cross-modality similarity quantification using a framework that was similar to the one applied in subsections **2.5** & **2.6**.

We took the average of the "forward" and "backward" (e.g., session 1→2, and 2→1) connectome-to-connectome correlation matrices for each pair of sessions (i.e., top right and bottom left quadrants in **Fig. 4**). Rather than threshold for "top-hits", we applied a range of correlation strength thresholds (0.45, 0.40, and 0.55) to assess where the greatest agreement was between imaging modalities in terms of connectivity patterns. Null distributions were generated as before by shuffling the target session labels (N = 50 iterations). As in the previous subsections, we computed the fraction of "hits" within each area of interest for all group pairings and canonical networks.

377
378 **Fig 8 | Inter-modality high connectome-to-connectome correlation strengths by network and group pairing. A.** *Inter-*
379 *modality instances of high correlation values for BOLD-fMRI cortex-only → WF-Ca^{2+}$_{Fast}$.* **B.** *Same as* **A.**, *for BOLD-fMRI*
380 *cortex-only → WF-Ca^{2+}$_{Slow}$. Subplots where matched group pairings (e.g., GLIA-to-GLIA) showed a significant result are*
381 *outlined with a solid black line. A statistically significant (two-tailed, MATLAB t-test, ttest2, Bonferroni corrected) surplus, or*
382 *paucity, of high correlation strengths were indicated by an *.*
383

Results, boxplots in **Fig. 8** for BOLD-fMRI cortex-only → WF-Ca$^{2+}_{Fast}$ or WF-Ca$^{2+}_{Slow}$, were organized

differently than boxplots in **Fig. 6**, **7**, and **S7** due to clear differences in correlation values between modalities

across canonical networks (see y-axis ranges in **Fig. 8**). For example, limbic and default mode networks showed

a much higher fraction of high inter-modal correlations than other canonical networks. Thus, boxplots were

organized with canonical networks in rows, and group pairings in columns.

For BOLD-fMRI cortex-only → WF-Ca$^{2+}_{Fast}$ (**Fig. 8A**), GLIA, SOM, and VIP showed no instances where

matched groups (e.g., GLIA-to-GLIA) showed higher fractions of high inter-modal correlation values than the null

distribution. Conversely, PV and SLC showed select instances where fractions of high inter-modal correlation

values were above chance. Curiously, these were apparent within different canonical networks (**Fig. 8**). For

BOLD-fMRI cortex-only → WF-Ca$^{2+}_{Slow}$ (**Fig. 8B**), GLIA, PV, and SLC showed no instances where matched

groups showed a higher fraction of high inter-modal correlation values than chance whereas SOM and VIP

showed some select instances that were. However, the specificity of these occurrences was low (i.e., they

appeared coincident with high fractions of high inter-modal correlation values in unmatched group pairings, e.g.,

Fig. 8B, somatosensory, SOM-to-PV, SOM-to-SOM, as well as PV-to-SLC, PV-to-PV, and PV-to-SOM, were all

above chance).

3. DISCUSSION

We investigated functional connectome-based ID in mice using simultaneously acquired WF-Ca^{2+} and

BOLD-fMRI data. In-line with previous work in mice[29] and humans[5-16], animals were identifiable using BOLD-

fMRI data. Building on this foundation and affirming the likely cellular-basis of these findings, animals were also

identifiable using WF-Ca^{2+} imaging data. To the best of our knowledge, this is the first study to investigate

connectome-based ID using this modality. Further, the unique structure of our dataset allowed us to explore

group ID rates (where groups bore GCaMP fluorescent labels in different cell populations), as well as inter-modal

connectome-based ID for the first time.

Translational work across species can be aided by using common tools (e.g., imaging modalities) and

analytical frameworks[38,39] (e.g., connectome-based ID). Work of this nature contributes by establishing whether

(and the degree to which) specific measurements are possible in multiple species[40] and by extension how useful

animal models might be within a specific clinical context. Given a foothold, i.e., when overlap across species can

be established, work in animals can offer insights that are not attainable from human subjects[21].

Connectome-based ID, using human BOLD-fMRI data, produces high rates of correct IDs (typically about

60-99%, depending on sample size[5,8,13]; here, 88% in the human data, **Fig. 1**). Previous work on mouse BOLD-

fMRI data, by Bergmann et al., established that ID rates in mice are lower (60-70%)[29] but still well above chance.

Here, we observe even lower, but again still significant, ID rates (20-40%) using BOLD-fMRI data. As expected,

ID rates based on WF-Ca^{2+} imaging data are higher (30-50%), but still below what Bergmann et al. observed.

This gap was not addressed by repeating our analyses using a framework designed to match the one

implemented in their paper. Specifically, creating a query and target set by splitting the dataset within each

session, i.e. using runs from different timepoints such that each timepoint was represented in each averaged

connectome (or query and target set) (**Fig. S6**). The remaining notable differences between the study by

Bergmann et al. and our own is that our mice were lightly anesthetized (0.5-0.75% isoflurane) whilst theirs were

awake, and dataset size (their N = 16 vs. our N = 45; which has previously been shown to affect ID rates[13]).

Similarly, it is very likely that our use of anesthesia reduced our observed ID rates given recent work in

anesthetized human subjects[41]. This is notable given that the majority of fMRI studies in rodents use anesthesia,

and that differences in resting-state fMRI signals between awake and anesthetized mice have not been

extensively characterized (see [42] for a recent systematic review). The inherent challenges (e.g., motion and

stress), and significant time required, to perform fMRI experiments on awake mice necessitates clear and specific

scientific reasons for doing so. It is also exceptionally difficult, and resource intensive, to generate large datasets

(typical of ID-based or similar analyses) from awake mice[43]. Future work is needed to confirm whether the use

of anesthesia in mice truly reduces connectome-based ID rates and the role this may play in studies focused on

inter- and intra- subject differences. In combination with follow-up studies, the findings in this work may provide

some key evidence for using awake animals within the context of exploring individual differences.

Human BOLD-fMRI data-based connectome ID shows clear canonical network dependence (**Fig. 1**)[1,23,25].

This feature was not evident in the study by Bergmann et al. where ID rates were comparable between all cortical

nodes, association, and sensory systems. Here, we investigated a wider array of networks, as well as subcortical

areas, and found that networks did influence ID rate. Specifically, paralleling findings from human BOLD-fMRI

data, we found somatosensory, and lateral cortical networks performed well (for their size), whilst subcortical

and brainstem networks performed poorly (**Fig. 5**). Notably, data from the whole-brain was necessary for this

observation as these findings were not evident in cortex-only BOLD-fMRI or WF-Ca^{2+} imaging data. That these

effect sizes were small was unsurprising given the overall low ID rates in our dataset, and that mice lack what

are believed to be the main drivers of individual differences in humans (higher order association networks)[44,45].

Our unique multimodal dataset allowed us to explore new aspects of the connectome-based ID framework.

With our WF-Ca^{2+} imaging data, we showed that the framework extends beyond applications in BOLD-fMRI data,

which helps affirm that cellular activity, and not noise or spurious signals, are the likely drivers of correct IDs.

Although we did not observe cross-modal agreement in terms of which mice are correctly identified across all

comparisons, we did see significant agreement between WF-Ca$^{2+}_{Slow}$ and cortex-only BOLD-fMRI (**Fig. S4A**) –

the most closely matched cross-modal conditions. These data also allowed us to explore group ID, which was

well above chance using data from either modality. The high rates from the WF-Ca^{2+} imaging data indicated that

there are differences in connectivity structure that are apparent in data from different cellular sources at our

'mesoscale' spatiotemporal resolution – even at low frequencies. This is encouraging for future work aimed at

examining cell-type specific activity or network-based measures at the 'mesoscale' using WF-Ca^{2+} imaging data.

This finding runs counter to the assumption that if cells of different types are closely linked at the microscale,

their apparent meso- or macro-scale organizational structures will be indistinguishable.

We also used these data to characterize inter-group connectome similarity and dissimilarity (**Fig. 6**, **7**, & **S5**).

Clear patterns in the WF-Ca^{2+} imaging results emerged between groups that aligned with how closely related

different cell-types are at the micro-circuit level. For example, GLIA were more easily dissociated from PV and

SLC, whilst the later were more similar to one another. Similarly, SOM and VIP were similar to one another and

unlike SLC. The broad similarities between inhibitory interneurons (and differences with respect to excitatory

neurons and glial cells) generally aligns with previous WF-Ca^{2+} work[37] and is consistent with the hierarchical

nature of cell type classes[46]. Curiously, some of these patterns were recapitulated in the fMRI data where the

signal source (BOLD) should be independent from group membership except for the role played by closer

lineage. Specifically, we expected that mice might be more similar within group, but not that trends across groups

would emerge (e.g., for SLC to be the most identifiable, and for VIP to be at chance-level). This could indicate

that there are differences across groups (e.g., sensitivity to anesthesia) that contributed to some of the observed

inter-group differences in both modalities. Future work in awake mice, and animals bearing more than one

fluorescent cell-label are needed to investigate this finding further. Yet, the difference in effect size, as well as

the clear dissimilarities evident in the WF-Ca^{2+} imaging data - but not the BOLD-fMRI data – indicate that the

cell-type specific findings in the WF-Ca^{2+} imaging data are still likely present.

Finally, we found that a strict winner-take-all ID framework failed when applied across imaging modalities

(i.e., WF-Ca^{2+} → BOLD-fMRI and vice versa). To us, this outcome emphasized the complementary information

available from each modality and encourages future experiments that investigate how these contrasts, when

applied together, can further inform on brain function[47]. To this end, we investigated whether different groups, or

networks, showed greater connectome similarity across modes than would be expected by chance (**Fig. 8**). We

found some preliminary, but intriguing evidence that SLC and PV groups, showed greater inter-modal similarity

(albeit unexpectedly using WF-Ca$^{2+}_{Fast}$ whilst some, more tenuous patterns, were found for SOM and VIP (using

WF-Ca$^{2+}_{Slow}$). More in-depth investigations into these data are warranted and necessitates a framework that goes

beyond measures of connectome similarity. Future work could also use data obtained via other imaging

modalities[48], as well as examining the extent to which identifiability is affected by disorders commonly modelled

in mice (e.g.[49]).

In conclusion, using a unique simultaneously acquired WF-Ca^{2+} and BOLD-fMRI dataset, we showed that

mice are identifiable using a connectome-based ID framework adopted from the human BOLD-fMRI literature.

This work extends this framework beyond BOLD-fMRI and explores features of connectome identifiability,

similarity, and dissimilarity across subjects, groups, and imaging modalities. Although a strength of animal

research is the ability to conduct experiments within a tightly controlled environment and on nearly identical

subjects, as neuroimaging methods in mice become more sophisticated, studies of inter- and intra- subject

features stand to contribute in new ways. Areas where this may be most relevant include studies of mental

illness, or within any arena where heterogeneity – both in terms of brain function and behavior – of unknown

origin and significance are the norm[27,50].

4. METHODS

All procedures were performed following the Yale Institutional Animal Care and Use Committee (IACUC)

and in agreement with the National Institute of Health Guide for the Care and Use of Laboratory Animals.

4.1 Group and dataset overview

Mice were housed on a 12-hour light/dark cycle. Food and water were available *ad libitum*. Animals were 6-8 weeks old, 25-30g, at the time of the first imaging session. Five mouse lines were used: (1) GLIA, (2) PV, (3) SLC, (4) SOM, and (5) VIP (**Fig. 2A** & **Table S1**). All groups were of mixed sex.

Four groups (all but GLIA) shared a C57BL/6J background (**Table S1**). Male CRE mice were selected from the offspring of parents with different genotypes; this avoided leaking of CRE expression. See **Table S1** for the full breeding scheme. Briefly, the Ai162 genotype resulted from tTA and TITL-GCaMP6s (TIGRE1.0). Differences between TIGRE1.0 and 2.0 have been reported[51] and TIGRE1.0 mice have been previously described[30]. The number of inbred generations may have caused some genetic variation between groups. Specifically, VIP mice were bred in-house for a prolonged period, which may have led to slight variations in their background. Given the careful monitoring of sublines by Jackson's lab, this factor was likely negligible for other groups. The fifth group (GLIA) was derived from the Aldh1l1-CreER/Ai162 line, where Aldh1l1-CreER is on a C57BL/6;FVB/N genetic background. The GLIA group received tamoxifen injections for three consecutive days after weaning to elicit GCaMP expression. No visible physiological or phenotypic effects due to tamoxifen toxicity have been reported in these animals.

4.2 Surgical procedure for permanent optical access to the cortical surface and immobilization

All mice underwent a minimally invasive surgical procedure were an in-house built head-plate (acrylonitrile butadiene styrene plastic, TAZ-5 printer, 0.35mm nozzle, Lulzbot) was attached to the thinned, but intact, skull for permanent optical access to the cortical surface, necessary for WF-Ca^{2+} imaging, and immobilization. The surgical procedure has been described by us previously[30,33,37]. Briefly, mice were initially anesthetized with isoflurane (5%, 70/30 medical air/O$_2$) and head-fixed in a stereotaxic frame (KOPF, USA). Isoflurane was then reduced to 2%. An eye ointment was applied; meloxicam (2mg/(kg)body weight) administered subcutaneously and bupivacaine (0.1%) injected locally, at the incision site. After shaving the hair, the scalp was washed with betadine followed by ethanol 70%, (×3). The scalp was surgically removed, and the skull cleaned and dried. Antibiotic powder (Neo-Predef) was applied to the incision site, and isoflurane reduced (1.5%). Skull-thinning of the frontal and parietal skull plates was performed with a hand-held drill (FST), diameter

of the tips 1.4mm and 0.7mm. Superglue (Locite) was applied to the exposed skull, followed by transparent

dental cement C&B Metabond (Parkell). The head-plate was attached to the dental cement before solidification.

The head-plate had a double-dovetail plastic frame with a microscope slide hand-cut to match the size and shape

of the mouse skull[19].

4.3 Imaging protocol

Mice were allowed 1-week to recover post-surgery before undergoing three simultaneous WF-Ca^{2+} and BOLD-

fMRI sessions on an 11.7T MRI scanner (Bruker, Billerica, MA) using custom MRI-compatible WF-Ca^{2+} imaging

equipment[30]. The interval between each imaging sessions was at least 1-week.

4.3.1 BOLD-fMRI data acquisition

Data were acquired as described previously[30,33] using ParaVision 6.0.1. Mice were scanned whilst lightly

anesthetized with isoflurane and free breathing (0.5-0.75%, 70/30 medical air/O$_2$). Body temperature was

continuously monitored (Spike2, Cambridge Electronic Design Limited), and maintained with a circulating water

bath. Imaging data and physiological recordings were synchronized (Master-8 A.M.P.I., Spike2 Cambridge

Electronic Design Limited). A gradient-echo, echo-planar-imaging (GE-EPI) sequence with a 1.0s repetition time

(TR) and 9.1ms echo time (TE) was used with an isotropic resolution of 0.4x0.4x0.4mm^3. Twenty-eight slices

yielded near whole-brain coverage (**Fig. 2B**). Each functional run was comprised of 600 volumes or 10mins of

data. On average, 7 functional runs were acquired per mouse per session. Three included a unilateral LED-

stimulation whilst 4 were acquired in the resting-state (no external stimuli were presented). Here, we consider

the resting-state data only.

In addition to the functional data, 4 structural images were acquired during each imaging session for data

registration purposes[30,33]. (1) A multi-spin-multi-echo (MSME) image sharing the same FOV as the functional

data with a TR/TE of 2500/20ms, 28 slices, two averages, and resolution of 0.1x0.1x0.4mm^3 in 10mins and 40s.

(2) A whole-brain isotropic MSME image with a TR/TE of 5500/20ms, 78 slices, two averages, and resolution of

0.2x0.2x0.2mm^3 in 11mins and 44s. (3) A fast-low-angle-shot (FLASH) time-of-flight (TOF) angiogram, with a

TR/TE of 130/4ms, resolution of 0.05x0.05x0.05mm^3 and FOV of 2.0x1.0x2.5cm^3 in 18mins. (4) A FLASH image

of the angiogram FOV, with a TR/TE of 61/7ms, four averages, and resolution of 0.1×0.1×0.1mm^3 in 11mins and

24s. Each average of images (1) and (2) were interleaved with functional scans such that the functional data

were acquired at approximately evenly spaced intervals throughout each imaging session.

4.3.2 WF-Ca^{2+} imaging data acquisition

WF-Ca^{2+} imaging data were acquired simultaneously with all BOLD-fMRI data as we have described

previously[30,33] using CamWare software version 3.17. During each acquisition, GCaMP-sensitive (470/24nm)

and GCaMP-insensitive (395/25nm) illumination (LLE 7Ch Controller from Lumencor) were interleaved at 20Hz.

Exposure time for each wavelength was 40ms to avoid artifacts caused by the rolling shutter refreshing. Thus,

the sequence was: 10ms blank, 40ms GCaMP-sensitive wavelength on, 10ms blank, 40ms GCaMP-insensitive

wavelength on. Following data processing, where the GCaMP-insensitive channel is regressed from the GCaMP-

sensitive channel, the data were at an effective 10Hz. The raw WF-Ca^{2+} imaging data had a raw spatial resolution

of 25×25µm^2 and a FOV of 1.4×1.4cm^2 (as in [30]).

4.4 Data processing

4.4.1 MRI data registration and preprocessing

BOLD-fMRI data processing was performed using RABIES (Rodent automated BOLD improvement of EPI

sequences[36]) v0.4.8 (https://rabies.readthedocs.io/en/stable/). All resting-state functional and isotropic MSME

structural scans (from all imaging sessions) were input and processed together. Within native subject space, the

BOLD-fMRI timeseries were slice time corrected, and head motion was estimated. Structural images were

corrected for inhomogeneities (N3 nonparametric nonuniform intensity normalization). The within-sample

anatomical template, created by averaging all structural images following non-linear registration, was registered

to our in-house template (using non-linear registration), which has been previously registered to the Allen Atlas

reference space CCfv3[33,35,36].

For each BOLD-fMRI scan, a representative mean image (averaged across time) was derived. These data

were corrected for intensity inhomogeneities and non-linearly registered to the corresponding isotropic structural

MSME image of each mouse acquired during each imaging session. This minimized distortions caused by

susceptibility artifacts. Then, the BOLD-fMRI data were moved to common space using four concatenated

transformations: (1) framewise rigid head motion correction, (2) representative mean BOLD-fMRI to individual

mouse/session isotropic MSME image, (3) individual mouse/session isotropic MSME image to within-dataset

template, and (4) within-dataset template to out-of-sample in-house template. As part of this transformation, the

data were resampled to the template resolution ($0.2 \times 0.2 \times 0.2 mm^3$). After this step, the Allen Atlas and BOLD-

fMRI data resided in the same space. Registration performance was visually inspected using the RABIES quality

control report. Following timeseries normalization to common space, the six-parameter motion estimates were

regressed from the timeseries and used to compute framewise-displacement. Frame scrubbing for motion using

a conservative 0.075mm threshold was applied. Data were filtered 0.008-0.2Hz and 30 timepoints discarded

from the beginning and end of each run to avoid filter-related edge artifacts. Cerebrospinal fluid and white matter

signals were regressed, and the data were smoothed with 0.4mm sigma. Global signal regression was not

applied as there is no equivalent for WF-Ca^{2+} imaging data and we aim to harmonize data processing as much

as possible across modalities.

4.4.2 WF-Ca^{2+} imaging data registration and processing

WF-Ca^{2+} imaging data (GCaMP-sensitive and -insensitive channels) were smoothed with a 0.1mm

sigma. Data were down-sampled by a factor of two in each spatial dimension and the GCaMP-insensitive channel

regressed from the GCaMP-sensitive channel. $\Delta F/F_0$ (F = fluorescence) was computed for each run. The data

were filtered, within both a slow (0.008-0.2Hz) and fast (0.4-4Hz) band. 300 timepoints were discarded from the

beginning and end of each run to match the BOLD-fMRI timeseries and eliminate photobleaching artifacts.

Timepoints censored (based on framewise displacement) from the BOLD-fMRI timeseries were also removed

from the WF-Ca^{2+} imaging timeseries. Data were registered to our in-house template using three transformations,

two linear (1) WF-$Ca^{2+} \rightarrow$ Angiogram, and (2) Angiogram $\rightarrow$ MSME and one nonlinear (3) MSME to in-house

template space, as described previously[30].

4.4.3 Connectomes

To compute connectomes, BOLD-fMRI or WF-Ca^{2+} imaging signals within each node (**Fig. S2**) was

averaged, and the inter-regional Pearson's correlation computed, and Fisher transformed. Connectomes were

computed for each run using data residing in our in-house template space.

4.5 Inclusion criteria

All data were temporally matched across imaging modalities. Thus, if a run (or timepoint) was excluded

based on criteria applied to the BOLD-fMRI data, the corresponding simultaneously acquired WF-Ca^{2+} imaging

data were also excluded (and vice versa). Data from both modalities were excluded if framewise displacement,

estimated using the BOLD-fMRI timeseries, exceeded 0.075mm. If 2/3 timepoints within a run exceeded this

cutoff, the run was excluded (none excluded). If registration to the in-house template space failed visual

inspection, the corresponding run was excluded (6 runs excluded across the dataset, and one subjects excluded

fully across all sessions). If data from either modality were corrupted, the data were excluded (two instances in

the WF-Ca^{2+} imaging dataset, corresponding BOLD-fMRI also excluded). If the WF-Ca^{2+} imaging signal was

poor, or the MR-angiogram scan was not acquired, the data were excluded (8 runs excluded). If the software

showed acquisition issues (4 runs excluded). Finally, all data from a mouse were excluded if less than two runs

in all sessions passed all preceding criteria (none excluded). Thus, a minimum of 20mins of functional data per

mouse per session were retained. Due to FOV variability, and common artifacts at the periphery of the WF-Ca^{2+}

imaging data, some regions at the edge of the cortical surface were excluded from WF-Ca^{2+} and cortex-only

BOLD-fMRI conditions (**Fig. S2**).

4.6 Analyses

4.6.1 Human BOLD-fMRI data

LR resting state BOLD-fMRI data from the human connectome project S900 release were downloaded

and processed following standard procedures, as described previously[18,19]. A 268-node functional atlas[5] was

used to compute connectomes using Person's correlation and a winner-take-all ID procedure applied, as

described previously[5,8]. Nodes were grouped into eight functional networks[1].

4.6.2 Connectome-based ID

The winner-take-all procedure has been described in detail previously[1]. Briefly, a query (e.g., session 1

connectomes) and target (e.g., session 2 connectomes) are selected. In an iterative process, each connectome

in the query and target set are compared using Pearson's correlation. If the highest inter-connectome correlation

coefficient belonged to the same subject (i.e. within-subject correlation > all other between-subject correlations),

this was denoted as a correct ID. For group ID, the mouse being queried was removed from the target session

(to avoid mouse ID bleed-through). Network-based ID was performed using only nodes within a given network

(**Fig. S2**). For random networks, we iteratively (N = 50) selected nodes at random and performed the ID

procedure using the resulting data.

4.6.3 Node and network definitions

Nodes belonging to the default mode network were selected based on the description by Whitesell and

colleagues[55]. Nodes belonging to the salience and lateral cortical networks were selected based on the triple-

network model[36,43,52-54]. Nodes belonging to other networks mentioned (e.g. somatosensory) were selected

based on the recent comparison across species[36,52,53].

4.7 Statistics

Statistical significance tests were computed in MATLAB (R2023b & R2021bV5).

ACKNOWLEDGEMENTS

Drs. Joel Greenwood, Omer Mano, and Paul Shamble from the Neurotechnology Core of the Yale Kavli Institute for Neuroscience for their technological expertise. Gabriel Desrosiers-Gregoire for RABIES software support. Dr. Xinxin Ge for help in acquiring the multimodal imaging data. Dr. Peter Herman for performing the head-plate implant surgeries. All members of the Yale Multiscale Imaging and Spontaneous Activity in Cortex (MISAC) group for help in conceptualizing and building the in-scanner WF-Ca^{2+} imaging apparatus.

FUNDING

CH is supported by a Medical Scientist Training Program training grant (NIH/NIGMS T32GM007205). This work was supported in-part by funding from the NIH: R01 MH111424 - RTC as well as U01 N2094358 - RTC. M.M. is funded by SNSF Postdoc.Mobility grant (214392).

AUTHOR CONTRIBUTIONS

FM and CH wrote the manuscript, processed the multimodal data, and conducted analyses. XS and GDG helped with analyses. WL downloaded and processed the human BOLD-fMRI data. MM processed the WF-Ca^{2+} imaging data. RTC supervised WL and the multimodal data collection. XP wrote code for multimodal data processing and registration (BioImage Suite software). RFB help conceptualize and frame the project and edited the manuscript. MC edited the manuscript and supervised GDG. EMRL collected the data, conducted analyses, supervised the study, and edited the manuscript.

CONFLICTS

XP is a consultant for the Brain Electrophysiology Laboratory Company. XP also consults and has an ownership stake in Veridat.

DATA & CODE AVAILABILITY

Data are not available in a public repository due to ongoing work by the authors on this dataset. Access can be obtained by contacting the corresponding authors. Data from the Allen Reference Atlas and CCFv3 are available on their website: https://portal.brain-map.org/. Software packages for data processing are available: multimodal data registration and processing of human data (BioImage Suite, https://bioimagesuiteweb.github.io/webapp/), and fMRI preprocessing (RABIES, https://github.com/CoBrALab/RABIES). ID code can be found here: https://www.nitrc.org/frs/?group_id=51

REFERENCES

Dubois, J. & Adolphs, R. Building a Science of Individual Differences from fMRI. *Trends Cogn Sci* **20**, 425-443 (2016). https://doi.org:10.1016/j.tics.2016.03.014

Bergmann E, K. I. in *Advances in Resting-State Functional MRI: Methods, Interpretation, and Applications* (ed Catie Chang Jean Chen) Ch. Chapter 13, 297-318. (Academic Press, 2023).

Schilsky, R. L. Personalized medicine in oncology: the future is now. *Nat Rev Drug Discov* **9**, 363-366 (2010). https://doi.org:10.1038/nrd3181

Leopold, J. A. & Loscalzo, J. Emerging Role of Precision Medicine in Cardiovascular Disease. *Circ Res* **122**, 1302-1315 (2018). https://doi.org:10.1161/CIRCRESAHA.117.310782

Finn, E. S. *et al.* Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nat Neurosci* **18**, 1664-1671 (2015). https://doi.org:10.1038/nn.4135

Miranda-Dominguez, O. *et al.* Heritability of the human connectome: A connectotyping study. *Netw Neurosci* **2**, 175-199 (2018). https://doi.org:10.1162/netn_a_00029

Miranda-Dominguez, O. *et al.* Connectotyping: Model Based Fingerprinting of the Functional Connectome. *Plos One* **9** (2014). https://doi.org:ARTN e111048
10.1371/journal.pone.0111048

Horien, C., Shen, X. L., Scheinost, D. & Constable, R. T. The individual functional connectome is unique and stable over months to years. *Neuroimage* **189**, 676-687 (2019). https://doi.org:10.1016/j.neuroimage.2019.02.002

Kaufmann, T. *et al.* Delayed stabilization and individualization in connectome development are related to psychiatric disorders. *Nat Neurosci* **20**, 513-+ (2017). https://doi.org:10.1038/nn.4511

Amico, E. & Goni, J. The quest for identifiability in human functional connectomes. *Sci Rep* **8**, 8254 (2018). https://doi.org:10.1038/s41598-018-25089-1

Horien, C. *et al.* Considering factors affecting the connectome-based identification process: Comment on Waller et al. *Neuroimage* **169**, 172-175 (2018). https://doi.org:10.1016/j.neuroimage.2017.12.045

Jalbrzikowski, M. *et al.* Functional connectome fingerprinting accuracy in youths and adults is similar when examined on the same day and 1.5-years apart. *Hum Brain Mapp* **41**, 4187-4199 (2020). https://doi.org:10.1002/hbm.25118

Waller, L. *et al.* Evaluating the replicability, specificity, and generalizability of connectome fingerprints. *Neuroimage* **158**, 371-377 (2017). https://doi.org:10.1016/j.neuroimage.2017.07.016

Demeter, D. V. *et al.* Functional Connectivity Fingerprints at Rest Are Similar across Youths and Adults and Vary with Genetic Similarity. *Iscience* **23** (2020). https://doi.org:ARTN 100801
10.1016/j.isci.2019.100801

Graff, K. *et al.* Benchmarking common preprocessing strategies in early childhood functional connectivity and intersubject correlation fMRI. *Dev Cogn Neurosci* **54**, 101087 (2022). https://doi.org:10.1016/j.dcn.2022.101087

Graff, K. *et al.* Functional connectomes become more longitudinally self-stable, but not more distinct from others, across early childhood. *Neuroimage* **258**, 119367 (2022). https://doi.org:10.1016/j.neuroimage.2022.119367

Jo, Y., Faskowitz, J., Esfahlani, F. Z., Sporns, O. & Betzel, R. F. Subject identification using edge-centric functional connectivity. *Neuroimage* **238**, 118204 (2021). https://doi.org:10.1016/j.neuroimage.2021.118204

Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: An overview. *Neuroimage* **80**, 62-79 (2013). https://doi.org:10.1016/j.neuroimage.2013.05.041

Luo, W. & Constable, R. T. Inside information: Systematic within-node functional connectivity changes observed across tasks or groups. *Neuroimage* **247**, 118792 (2022). https://doi.org:10.1016/j.neuroimage.2021.118792

Michon, K. J., Khammash, D., Simmonite, M., Hamlin, A. M. & Polk, T. A. Person-specific and precision neuroimaging: Current methods and future directions. *Neuroimage* **263**, 119589 (2022). https://doi.org:10.1016/j.neuroimage.2022.119589

Zerbi, V. On the use of preclinical imaging to explore the principles of brain function in rodent models and their relevance for illnesses of the human mind. *Transl Psychiatry* **12**, 155 (2022). https://doi.org:10.1038/s41398-022-01924-y

22 Ryali, S. *et al.* Combining optogenetic stimulation and fMRI to validate a multivariate dynamical
systems model for estimating causal brain interactions. *Neuroimage* **132**, 398-405 (2016).
https://doi.org:10.1016/j.neuroimage.2016.02.067
23 Gozzi, A. & Zerbi, V. Modeling Brain Dysconnectivity in Rodents. *Biol Psychiatry* **93**, 419-429 (2023).
https://doi.org:10.1016/j.biopsych.2022.09.008
24 Murdy, T. J. *et al.* Leveraging genetic diversity in mice to inform individual differences in brain
microstructure and memory. *Front Behav Neurosci* **16**, 1033975 (2022).
https://doi.org:10.3389/fnbeh.2022.1033975
25 Melozzi, F. *et al.* Individual structural features constrain the mouse functional connectome. *Proc Natl*
Acad Sci U S A **116**, 26961-26969 (2019). https://doi.org:10.1073/pnas.1906694116
26 Allsopp, K., Read, J., Corcoran, R. & Kinderman, P. Heterogeneity in psychiatric diagnostic
classification. *Psychiatry Res* **279**, 15-22 (2019). https://doi.org:10.1016/j.psychres.2019.07.005
27 Feczko, E. *et al.* The Heterogeneity Problem: Approaches to Identify Psychiatric Subtypes. *Trends*
Cogn Sci **23**, 584-601 (2019). https://doi.org:10.1016/j.tics.2019.03.009
28 Feczko, E. & Fair, D. A. Methods and Challenges for Assessing Heterogeneity. *Biol Psychiatry* **88**, 9-17
(2020). https://doi.org:10.1016/j.biopsych.2020.02.015
29 Bergmann, E., Gofman, X., Kavushansky, A. & Kahn, I. Individual variability in functional connectivity
architecture of the mouse brain. *Commun Biol* **3**, 738 (2020). https://doi.org:10.1038/s42003-020-
01472-5
30 Lake, E. M. R. *et al.* Simultaneous cortex-wide fluorescence Ca(2+) imaging and whole-brain fMRI. *Nat*
Methods **17**, 1262-1271 (2020). https://doi.org:10.1038/s41592-020-00984-6
31 Lin, M. Z. & Schnitzer, M. J. Genetically encoded indicators of neuronal activity. *Nat Neurosci* **19**, 1142-
1153 (2016). https://doi.org:10.1038/nn.4359
32 Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature* **499**, 295-300
(2013). https://doi.org:10.1038/nature12354
33 Vafaii, H. *et al.* Multimodal measures of spontaneous brain activity reveal both common and divergent
patterns of cortical functional organization. *Nat Commun* **15**, 229 (2024).
https://doi.org:10.1038/s41467-023-44363-z
34 Desrosiers-Gregoire, G., Devenyi, G. A., Grandjean, J, Chakravarty, MM. Rodent Automated Bold
Improvement of EPI Sequences (RABIES): A standardized image processing and data quality platform
for rodent fMRI. *bioRxiv* (2023). https://doi.org:https://doi.org/10.1101/2022.08.20.504597
35 Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas. *Cell*
181, 936-953 e920 (2020). https://doi.org:10.1016/j.cell.2020.04.007
36 Mandino F, S. X., Desrosiers-Gregoire G, O'Connor D, Mukherjee B, Owens A, Qu A,Onofrey
J,Papademetris X, Chakravarty MM, Strittmatter, SM Lake EMR. Aging-Dependent Loss of Connectivity
in Alzheimer's Model Mice with Rescue by mGluR5 Modulator. *bioRxiv* (2023).
https://doi.org:https://doi.org/10.1101/2023.12.15.571715
37 O'Connor, D. *et al.* Functional network properties derived from wide-field calcium imaging differ with
wakefulness and across cell type. *Neuroimage* **264**, 119735 (2022).
https://doi.org:10.1016/j.neuroimage.2022.119735
38 Gutierrez-Barragan D, R. J., Panzeri S, Xu T, Gozzi A. Evolutionarily conserved fMRI network
dynamics in the mouse, macaque, and human brain. *bioRxiv* (2024).
https://doi.org:https://doi.org/10.1101/2023.07.19.549681
39 Greene, A. S., Horien, C., Barson, D., Scheinost, D. & Constable, R. T. Why is everyone talking about
brain state? *Trends Neurosci* **46**, 508-524 (2023). https://doi.org:10.1016/j.tins.2023.04.001
40 Pagani, M., Gutierrez-Barragan, D., de Guzman, A. E., Xu, T. & Gozzi, A. Mapping and comparing fMRI
connectivity networks across species. *Commun Biol* **6**, 1238 (2023). https://doi.org:10.1038/s42003-
023-05629-w
41 Luppi AI, G. D., Ranft A, Ilg R, Jordan D, Bzdok D, Owen AM, Naci L, Stamatakis EA, Amico E, Misic B.
General anaesthesia reduces the uniqueness of brain connectivity across individuals and across
species. *bioRxiv* (2023). https://doi.org:10.1101/2023.11.08.566332
42 Mandino, F., Vujic, S., Grandjean, J. & Lake, E. M. R. Where do we stand on fMRI in awake mice?
Cereb Cortex **34** (2024). https://doi.org:10.1093/cercor/bhad478

43 Grandjean, J. *et al.* Common functional networks in the mouse brain revealed by multi-centre resting-
state fMRI analysis. *Neuroimage* **205**, 116278 (2020).
https://doi.org:10.1016/j.neuroimage.2019.116278
44 Mantwill, M., Gell, M., Krohn, S. & Finke, C. Brain connectivity fingerprinting and behavioural prediction
rest on distinct functional systems of the human connectome. *Commun Biol* **5**, 261 (2022).
https://doi.org:10.1038/s42003-022-03185-3
45 Dufford, A. J., Noble, S., Gao, S. & Scheinost, D. The instability of functional connectomes across the
first year of life. *Dev Cogn Neurosci* **51**, 101007 (2021). https://doi.org:10.1016/j.dcn.2021.101007
46 Zeng, H. & Sanes, J. R. Neuronal cell-type classification: challenges, opportunities and the path
forward. *Nat Rev Neurosci* **18**, 530-546 (2017). https://doi.org:10.1038/nrn.2017.85
47 Meyer-Baese, L., Watters, H. & Keilholz, S. Spatiotemporal patterns of spontaneous brain activity: a
mini-review. *Neurophotonics* **9**, 032209 (2022). https://doi.org:10.1117/1.NPh.9.3.032209
48 Markicevic, M., Savvateev, I., Grimm, C. & Zerbi, V. Emerging imaging methods to study whole-brain
function in rodent models. *Transl Psychiatry* **11**, 457 (2021). https://doi.org:10.1038/s41398-021-01575-
5
49 Zerbi, V. *et al.* Brain mapping across 16 autism mouse models reveals a spectrum of functional
connectivity subtypes. *Mol Psychiatry* **26**, 7610-7620 (2021). https://doi.org:10.1038/s41380-021-
01245-4
50 Voelkl, B. *et al.* Reproducibility of animal research in light of biological variation. *Nat Rev Neurosci* **21**,
384-393 (2020). https://doi.org:10.1038/s41583-020-0313-3
51 Daigle, T. L. *et al.* A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced Brain-Cell-
Type Targeting and Functionality. *Cell* **174**, 465-480 e422 (2018).
https://doi.org:10.1016/j.cell.2018.06.035
52 Zerbi, V., Grandjean, J., Rudin, M. & Wenderoth, N. Mapping the mouse brain with rs-fMRI: An
optimized pipeline for functional network identification. *Neuroimage* **123**, 11-21 (2015).
https://doi.org:10.1016/j.neuroimage.2015.07.090
53 Xu, N. *et al.* Functional Connectivity of the Brain Across Rodents and Humans. *Front Neurosci* **16**,
816331 (2022). https://doi.org:10.3389/fnins.2022.816331
54 Menon, V. Large-scale brain networks and psychopathology: a unifying triple network model. *Trends*
Cogn Sci **15**, 483-506 (2011). https://doi.org:10.1016/j.tics.2011.08.003