Perception of task duration affects metabolic cost during split-belt adaptation

S.N. Jeffcoat[1], A. Aragon[2], A. Kuch[1], S. Farrokhi[1], N. Sanchez[1,3*]

[1.] Department of Physical Therapy, Crean College of Health and Behavioral Sciences, Chapman University

[2.] Department of Applied Human Physiology, Crean College of Health and Behavioral Sciences, Chapman University

[3.] Department of Electrical Engineering and Computer Science, Fowler School of Engineering, Chapman University

* Corresponding Author
Natalia Sanchez
sanchezaldana@chapman.edu
9401 Jeronimo Road
Irvine CA 92618

Task duration influences metabolic cost of adaptation

Abstract

Humans continuously adapt locomotor patterns. Whether metabolic cost reduction is the primary objective or a by-product of the observed biomechanical changes during adaptation is not known. The main goal of our study is to determine if perception of task duration affects the adaptation of locomotor patterns to reduce energetic cost during split-belt walking. We tested the hypothesis that individuals who believe they will sustain a locomotor adaptation task for a prolonged time will reduce metabolic cost by adapting toward a walking pattern associated with lower mechanical work. N=14 participants walked on a split-belt treadmill for 10 minutes with knowledge of task duration (group K), while N=15 participants performed the task under the assumption that they would walk for 30 minutes (group U). Both groups walked for 10 minutes with the belts moving at 1.5 and 0.5 m/s, followed by 6 minutes of walking with both belts at 1.0 m/s. We observed a significant main effect of Time ($p<0.001$, observed power 1.0) and the interaction of Time×Group ($p=0.004$, observed power 0.84) on metabolic cost. Participants in the U group had a metabolic cost that was 12% lower during adaptation compared to the K group, which did not reduce metabolic cost during adaptation. The metabolic cost reduction observed in group U was not associated with biomechanical changes during adaptation. Our results indicate that metabolic cost reduction has a primary role in tasks that need to be sustained for a prolonged time, and this reduction is not only related to biomechanical factors.

New and Noteworthy

Individuals can markedly reduce the metabolic cost of a walking adaptation task if they believe they must sustain the task for a prolonged time. The variables typically used to track adaptation do not track the reductions in metabolic cost observed in our study. Metabolic cost reduction can occur without clear changes in biomechanical factors.

Keywords

Adaptation, locomotion, split-belt, metabolic cost

Introduction

Humans continuously adapt locomotor patterns in response to environmental conditions, such as changes in walking terrain, footwear, and load carriage (Bruijn and van Dieën 2018; Kowalsky et al. 2021; Marigold and Patla 2008). In laboratory settings, split-belt treadmills have been successfully used to study locomotor adaptation, where individuals walk on a dual-belt treadmill in which each belt moves at different speeds (Dietz et al. 1994; Reisman et al. 2005). Split-belt walking leads to immediate changes in intralimb coordination and gradual changes in interlimb coordination (Reisman et al. 2005), which are attributed to distinct neural control mechanisms (Gonzalez-Rubio et al. 2019; Reisman et al. 2005). A vast amount of literature in the field has shown that the neuromotor system can adapt to the split-belt's asymmetric environment and retain some of the adaptive changes in locomotion in the short and long term (Malone and Bastian 2010, 2016; Reisman et al. 2005; Torres-Oviedo et al. 2011; Torres-Oviedo and Bastian 2010). While the kinematic, kinetic, and physiological changes that occur during locomotor adaptation have been thoroughly characterized (Betschart et al. 2017; Finley et al. 2013; Hinton et al. 2019; Jacobsen and Ferris 2023; Kambic et al. 2023; Mawase et al. 2013; Roemmich et al. 2012, 2016; Sánchez et al. 2019, 2021; Selgrade et al. 2017a, 2017b; Stenum and Choi 2020), the factors driving split-belt walking adaptation are still under debate. Identifying the factors that shape adaptive locomotor processes allows manipulating them to accelerate adaptation and long-term retention of the newly learned locomotor patterns, and this information is vital in the retraining of walking behaviors as part of rehabilitation interventions (Charlton et al. 2021; Reisman et al. 2010).

Empirical studies of split-belt adaptation (Brinkerhoff et al. 2023; Park and Finley 2022), as well as simulation studies of split-belt treadmill walking (Price et al. 2023; Seethapathi et al. 2021) have shown that the early stages of locomotor adaptation are associated with an increase in balance control, whereas the later stages of locomotor adaptation are associated with reductions in energetic cost. An area of interest regarding locomotor adaptation is whether metabolic cost reduction is the driving cost function or a by-product of the observed biomechanical changes (Sparrow and Newell 1994, 1998). Studies in cycling have provided evidence for energetic cost shaping motor behaviors: cyclists generate power proportionally to the distance to the endpoint (Joseph et al. 2008; Swart et al. 2009). These findings indicate that the central nervous system uses efferent feedback and modulates central commands to couple the metabolic and biomechanical limits of the body to the demands of physical tasks in a

feedforward process that allows individuals to complete a task while maintaining a reserve capacity (Hampson et al. 2001). Thus, individuals can modify effortful motor tasks to maintain the task for prolonged time. What remains to be determined is whether maintaining an energetic reserve is also prioritized when performing a novel task, such as adapting to a split-belt treadmill, and how maintaining an energetic reserve might shape locomotor adaptation.

Split-belt walking can be considered an effortful behavior, where an individual's mechanical work and energetic cost markedly increase during initial exposure to split-belt walking (Brinkerhoff et al. 2023; Finley et al. 2013; Sánchez et al. 2021; Selgrade et al. 2017a, 2017b). As individuals adapt their walking pattern to the split-belt treadmill, they reduce positive power and positive work generated by their legs (Brinkerhoff et al. 2023; Sánchez et al. 2021; Selgrade et al. 2017a, 2017b), which is accompanied by a reduction in metabolic cost (Finley et al. 2013; Sánchez et al. 2021). Recent studies have shown that energetic optimization is prioritized during a locomotor adaptation task, even when a secondary cognitive task is superimposed (McAllister et al. 2021). Also, a recent study showed that fitness level influences adaptation to split-belt walking, with fitter individuals adapting the kinematics related to energetic cost reduction at a slower rate than sedentary individuals (Brinkerhoff et al. 2023). Given that walking patterns can change continuously, even for small energetic savings (Sánchez et al. 2019; Selinger et al. 2015), our working hypothesis is that task duration will influence walking behavior during locomotor adaptation as individuals aim to maintain an energetic reserve, further supporting the role of energetics in driving adaptation of locomotor behaviors.

The main goal of our study is to determine if perception of task duration affects the adaptation of locomotor patterns to reduce energetic expenditure during split-belt walking. We tested the hypothesis that individuals who believe they will sustain a locomotor adaptation task for a prolonged time, will have a lower energy cost. Previous research has shown that more positive step length asymmetries are associated with lower work by the fast leg and lower energetic cost (Sánchez et al. 2019, 2021). Thus, we hypothesize that individuals who believe they will sustain the locomotor task for a longer time will adapt faster towards a more positive asymmetry and reduce positive work by the fast leg. By confirming our hypotheses, our results will support the role of energetic optimization in being a primary goal during the adaptation of locomotor behaviors as opposed to simply being a by-product of the adaptation of walking behaviors.

Materials and Methods

We recruited 29 participants between the ages of 18-35. Participants were excluded if they had a history of lower extremity surgery at any point in their lives or lower extremity injury in the last two years, performed more than 10 hours of weekly exercise, reported current cough, cold, congestion, or any respiratory ailment that affected breathing, had previous experience walking on a split-belt treadmill, or had a history of neurological disorders or severe head trauma. The Chapman University Institutional Review Board approved all experimental procedures, with approval number IRB-23-189, and participants provided written, informed consent before testing. All study aspects conformed to the principles described In the Declaration of Helsinki.

Data were collected at the Gait Behavior Lab at Chapman University's Department of Physical Therapy using a Gait Real-time Analysis Interactive Lab (GRAIL) system (Motek Medical Base.V., Houten, Netherlands). The GRAIL is equipped with 10 motion capture cameras (Vicon Motion Systems Ltd, Oxford, UK), and an instrumented dual-belt treadmill (Motek Medical Base.V.). We used a metabolic cart to measure energetic expenditure (TrueOne 2400, Parvomedics, Salt Lake City, UT, USA). Participants were instructed not to eat or have caffeine two hours before the experiment. Before each experiment, we calibrated the motion capture system and the metabolic cart per manufacturer specifications. During all walking trials, participants wore a harness to prevent falls, without providing body weight support.

Experimental Protocol

After informed consent, participants were fitted with a mask covering their nose and mouth to measure their breathing (Fig. 1A). First, participants' resting caloric expenditure was measured while standing for 6 minutes. Then, participants underwent three walking trials with breaks between each to ensure resting conditions before continuing to the next trial. The first trial, Baseline (Base), had participants walking for 6 minutes with the belts under each leg moving at 1 m/s. The second trial had participants walking for 10 minutes with the left/fast belt moving at 1.5 m/s and the right/slow belt at 0.5 m/s (Fig. 1Base, C). The third trial had participants walking again during a post-adaptation trial for 6 minutes with the belts under each leg moving at 1 m/s. Participants were assigned to either the knowledge group (K) or the uncertainty group (U) via a coin flip to ensure randomization. Group K participants performed the split-belt adaptation trial with knowledge of how long the trial would last, and they were given an update on the time

remaining every minute. Group U participants were given the instruction: "You will be walking until we obtain the data needed, which can take about 30 minutes". Participants in the U group were given no updates on time elapsed or time remaining, and they also walked for 10 minutes. At the end of the 10th minute, we stopped the treadmill and told participants that we had collected all the necessary data.

Participants were asked to rate their level of exertion via the Borg Rating of Perceived Exertion (RPE) (Borg, 1962) using their fingers to indicate, on a scale of 1-10 which level of the RPE table they perceived. During the baseline walking trial (Base), participants reported RPE at the 5-minute mark. During the split-belt trial, they were asked to rate their level of exertion at the 1 (early adaptation – EA), 5 (mid-adaptation – MA), and 9-minute (late adaptation – LA) marks. During the tied post-adaptation trial, participants rated their RPE at the 1 (early post-adaptation – EP) and 5-minute marks (late post-adaptation – LP) (Fig. 1C).

Data Acquisition

We recorded marker data to calculate spatiotemporal variables and force data to calculate work by the legs. The positions of reflective markers located bilaterally on the greater trochanter, lateral epicondyle of the femur, and lateral malleolus were recorded at 100 Hz (Fig.1A). Force plates embedded into the split-belt treadmill recorded the ground reaction forces generated by each leg with a sampling frequency of 1000 Hz. Metabolic cost was recorded by determining the rates of oxygen consumption (VO_2) and carbon dioxide production (VCO_2). We also collected heart rate (HR) as a measure of cardiovascular intensity using a Polar H10 chest strap monitor (Polar Electro Inc., Bethpage, NY, USA), synchronized to the metabolic cart.

Data Processing and Analysis

We used custom-written code in MATLAB R2023b (Mathworks, Natick, MA, USA) for all data processing and analyses. We estimated the energy consumed using the standard Brockway equation to obtain metabolic rate (Brockway, 1987). We subtracted each participant's standing metabolic rate from each walking trial. Thus, all metabolic rate values presented here are net metabolic rates. Biomechanical data processing was done as in (Sánchez et al. 2019, 2021). A fourth-order low-pass digital Butterworth filter smoothed marker data and ground reaction force data using cut-off frequencies of 10 Hz and 20 Hz, respectively. We calculated step lengths using marker data, defined as the distance between the lateral malleoli markers at the instance

of peak anterior position of the respective marker. Step length asymmetry was then defined as follows:

$$SLA = \frac{SL_{fast} - SL_{slow}}{SL_{fast} + SL_{slow}} \text{ (Eq. 1)}$$

With SL_{fast} representing the step length at heel strike on the fast belt and SL_{slow} representing the step length at heel strike on the slow belt. We calculated leading and trailing limb placement for each limb as the fore-aft distance of the ankle markers relative to the midpoint between markers placed bilaterally on the greater trochanters.

We estimated the mechanical work performed by the legs using an extension of the individual limbs method (Donelan et al. 2002a, 2002b; Sánchez et al. 2019, 2021; Selgrade et al. 2017a, 2017b). This method approximates the legs as massless pistons and the entire body as a point mass. We measured individual leg ground reaction forces from the instrumented treadmill, and segmented forces into strides using a vertical ground reaction force threshold of 32 N (Selgrade et al. 2017a) to identify the beginning and end of each stride. We calculated the mediolateral, fore-aft, and vertical center of mass velocities by integrating the center of mass accelerations. We then calculated mechanical power as the dot product between the center of mass velocity and the ground reaction force from each leg. To determine the total positive and negative work performed by a leg or a belt, we calculated the time integral of the positive or negative portion of the mechanical power over the stride cycle. Work was normalized by leg length and mass to derive unitless mechanical work.

Statistical Analyses

Statistical analyses were run in SPSS version 29 (IBM, Armonk, NY, USA) and MATLAB R2023b. We assessed data for normality using the Shapiro-Wilk test. We compared age, height, and weight between the two experimental groups using independent samples t-tests or Wilcoxon Signed Rank test for not normally distributed data. All summary statistics are reported as mean ± standard deviation unless otherwise specified.

The metabolic power for the Base condition was computed by averaging the power for the last 2 minutes. The metabolic power for EA, MA, and LA was averaged between the second and third minutes, the fifth and sixth minutes, and the ninth and tenth minutes, respectively, of split-belt walking to match the RPE reporting periods (Figure 2). We used the second instead of the first

minute for metabolic data as the average duration to reach peak metabolic power is around 1 minute (Bogard et al. 2023; Finley et al. 2013). Finally, the metabolic power for post-adaptation was averaged between the second and third minutes and the fifth and sixth minutes of this condition to obtain average values of metabolic cost for EP and LP, respectively.

We obtained average values for step length asymmetry, step lengths, leading and trailing limb placement, double support time, swing time, step width, and positive and negative work for the left/fast and right/slow legs for the last five strides of the Base trial. For EA, MA, and LA we averaged the first, middle, and last five strides of the split-belt trial (Park and Finley 2022). For the post-adaptation trial, we averaged the five first and last strides (EP, LP, respectively). These are close to the timepoints when we obtained RPE and average measures of metabolic cost.

We compared differences between both groups for all average variables via 2×6 Mixed ANOVAs, with group (U or K) being the between-subjects effect and timepoint (Base, EA, MA, LA, EP, LP) being the within-subjects effect. Data were tested for sphericity using Mauchly's test. If the sphericity assumption was not fulfilled, we used the Huyn-Feldt corrected p-values to assess for significant effects or interaction effects. We obtained the post-hoc estimated statistical power for each significant effect in the mixed ANOVA and effect size estimates using partial eta-squared, with values of .01, .06, and .14 to indicate small, medium, or large effect sizes, respectively (Cohen 1988; Richardson 2011). We ran multiple comparisons using LSD-adjusted confidence intervals for significant effects in the ANOVA.

To determine whether the observed differences in metabolic cost during LA were due to group-specific differences in gait biomechanics during EA, MA, and LA, we ran correlation analyses between biomechanical variables and metabolic cost for each group and between biomechanical variables. We also used linear regression models of the form: $MetCost \sim W^+_{Fast} + W^+_{Slow} + Group$ during LA and $\Delta MetCost \sim \Delta W^+_{Fast} + \Delta W^+_{Slow} + Group$. We used positive work in our model based on previous work showing that the positive work reductions track metabolic cost reductions (Sánchez et al. 2019, 2021; Selgrade et al. 2017a).

Results

N = 29 participants volunteered for this study. No significant differences were observed in participant demographics between groups (Table 1, Supplementary Figures 1, 2). Metabolic

cost and kinematic data were collected for all participants. Force data was collected in all participants except one participant in the U group for whom force data were corrupted, which hindered performing mechanical calculations. Metabolic and kinematic data for this participant were included in all other analyses.

Perception of task duration influenced metabolic cost during adaptation

We observed a significant main effect of Time ($p<0.001$, observed power 1.0, partial eta-squared 0.714) and the interaction of Time×Group ($p=0.004$, observed power 0.84, partial eta-squared 0.141) on metabolic power (Fig. 2A, 2B). We did not observe a main effect of Group ($p=0580$). Across groups, the metabolic cost during EA ($p<0.001$), MA ($p<0.001$), LA ($p<0.001$), and EP ($p<0.001$) was significantly different from Base. Participants in the K group did not significantly decrease their metabolic cost during adaptation (post-hoc $p=0.233$ between EA and MA, and $p=0.092$ between EA and LA). In contrast, participants in the U group significantly decreased their metabolic cost during adaptation (post-hoc $p<0.001$ between EA and MA and $p=0.005$ between EA and LA). The metabolic cost during EA was 3.806 ± 0.632 Watts/kg for the K group and 3.646 ± 0.646 for the U group. The metabolic cost during MA was 3.657 ± 0.702 Watts/kg for the K group and 3.204 ± 0.564 for the U group. The metabolic cost during LA was 3.533 ± 0.698 Watts/kg for the K group and 3.182 ± 0.583 for the U group.

For heart rate, we observed a significant main effect of Time ($p<0.001$, observed power 1.0, partial eta-squared 0.703) and the interaction of Time×Group ($p=0.024$, observed power 0.69, partial eta-squared 0.109) but no main effect of Group ($p=0.206$) (Fig. 2C). Across groups, heart rate during EA ($p<0.001$), MA ($p=0.003$), EP ($p<0.001$), and LP ($p<0.001$) was significantly different from Base. We did not observe significant post-hoc differences between the K and U groups during the different timepoints. Heart rate during LA for the K group was 117 ± 16 beats per minute and 107 ± 16 for the U group.

Despite differences in metabolic cost and heart rate in the U group compared to the K group, participants' perception of exertion was not significantly different between groups. We observed a significant main effect of Time ($p<0.001$, observed power 1.0, partial eta-squared 0.572). We did not observe a significant main effect of Group or for the interaction of Time×Group ($p=0.369$ and $p=0.165$ respectively) (Fig. 2D). Across groups, RPE during EA ($p<0.001$), MA ($p<0.001$), and LA ($p<0.001$) was significantly different from Base.

Task duration influences metabolic cost of adaptation

We did not observe any differences between groups during post-adaptation in metabolic cost, heart rate, or RPE (p>0.050).

Perception of task duration affected right/slow step lengths during adaptation

We observed a significant main effect of Time (p<0.001, observed power 1.0, partial eta-squared 0.896) but no significant effect of Group (p=0.191) or the interaction of Time×Group (p=0.170) on step length asymmetry (Fig. 3A-B). Across groups, step length asymmetry during EA (p<0.001), MA (p=0.003), EP (p<0.001), and LP (p<0.001) was significantly different from the Base condition. Step length asymmetry during LA was -0.035 ± 0.089 for the K group and 0.004 ± 0.041 for the U group. Step length asymmetry during EP was 0.313 ± 0.080 for the K group and 0.413 ± 0.136 for the U group.

Step length asymmetry is a combination of right/slow and left/fast step lengths. No significant differences between groups were observed for the left/fast step length (main effect of Time p<0.001, observed power 1.0, no effect of the interaction of Time×Group, p=0.976 or Group, p=0.403) (Fig. 3C). Across groups, step lengths during EA (p<0.001), MA (p<0.001), LA (p<0.001), EP (p<0.001) and LP (p<0.001) were significantly different from the Base condition (Fig. 3C).

We observed significant differences in the right/slow leg step lengths (Fig. 3D). We observed a significant main effect of Time (p<0.001, observed power 1.0, partial eta-squared 0.845) and the interaction of Time×Group (p=0.031, observed power 0.73, partial eta-squared 0.098) and no significant effect of Group (p=0.591) on step lengths of the right/slow leg. The step lengths with the right/slow leg were shorter in the U group during LA compared to Base (p=0.002) but not in the K group compared to Base (p=0.437). Step lengths for the right/slow leg during LA were 491 ± 59 mm for group K and 463 ± 46 mm for group U (p=0.166). During post-adaptation, step lengths for the right/slow leg during EP were 296 ± 75 mm for group K and 247 ± 56 mm for group U (p=0.095).

Perception of task duration influenced work by the right/slow leg during adaptation

We did not observe any significant differences between groups in work by the left/fast leg during adaptation or post-adaptation (Fig. 4A, 5C): we observed a significant main effect of Time (p<0.001, observed power 1.0) but not for the interaction of Time×Group (p=0.652) or the main effect of Group (p=0.278). Across groups, work during EA (p<0.001), MA (p<0.001), LA (p<0.001), EP (p=0.007), and LP (p=0.017) was significantly different from the Base condition.

For negative work by the left/fast leg (Fig. 4E), we observed a significant main effect of Time (p<0.001, observed power 1.0) but not for the interaction of Time×Group (p=0.573) or the main effect of Group (p=0.802). Across groups, work during EA (p<0.001), MA (p<0.001), LA (p<0.001), EP (p=0.007), and LP (p=0.017) was significantly more negative than the Base condition.

We observed a significant main effect of Time (p<0.001, observed power 1.0, partial eta-squared 0.690) and the interaction of Time×Group (p=0.031, observed power 0.69, partial eta-squared 0.111) and no significant effect of Group (p=0.597) on the positive work by the right/slow leg (Fig. 4Base, 4D). Participants in the U group generated marginally less positive work with the right/slow leg during EA compared to Base (p=0.052) and significantly less work during MA and LA compared to Base (p<0.005). In contrast, participants in the K group generated work that was significantly lower than Base only during MA (p=0.017) (Fig. 4D).

For negative work by the right/slow leg (Fig. 4F), we observed a significant main effect of Time (p<0.001, observed power 1.0) but not for the interaction of Time×Group (p=0.462) or the main effect of Group (p=0.419).

For net work, which is the sum of positive and negative work by both legs, we observed a significant main effect of Time (p<0.001, observed power 1.0) but not for the interaction of Time×Group (p=0.320) or the main effect of Group (p=0.185).

Perception of task duration did not systematically influence other spatiotemporal variables during adaptation and post-adaptation

Task duration influences metabolic cost of adaptation

We observed a significant main effect of Time ($p<0.001$), but not of Group ($p>0.500$) or the interaction of Time×Group ($p>0.100$) on double support times, swing times, leading and trailing limb placement, and step widths for the left/fast or right/slow leg. We did not observe significant differences in cadence between the groups during split-belt walking ($p=0.660$), which could explain the differences in metabolic cost.

Lack of association between metabolic cost and gait biomechanics during adaptation

We assessed correlations between the change in metabolic cost from EA to LA and the change in biomechanical variables (Fig. 5). The change in metabolic cost during adaptation was not associated with a change in step length asymmetry in the K group ($r=-0.257$, $p=0.370$) or in the U group ($r=-0.175$, $p=0.534$) (Fig. 5A). The change in metabolic cost was not correlated with the change in positive work by the left/fast leg in the K group ($r=0.449$, $p=0.107$) or in the U group ($r=0.328$, $p=0.252$) (Fig. 5B). The change in metabolic cost was marginally correlated with the change in positive work by the right/slow leg in the U group ($r=0.552$, $p=0.054$) but not in the K group ($r=0.029$, $p=0.920$) (Fig. 5C). The change in net work by the legs was not correlated with the change in metabolic cost ($p=0.113$ for the K group and $p=0.947$ for the U group) (Fig. 5D).

Regarding the association between gait biomechanics, we found a significant association between change in step length asymmetry and change in positive work by the left/fast leg ($r=-0.580$, $p=0.030$ for the K group, and $r=-0.905$, $p<0.001$ for the U group) (Fig. 5E), but no association between change in step length asymmetry and change in positive work by the right/slow leg (Fig. 5F).

The model using work to predict metabolic cost during LA was not significant ($p=0.751$). None of the biomechanical variables measured were significantly associated with metabolic cost ($p>0.100$) during LA. The model using change in work from EA to LA to predict change in metabolic cost from EA to LA was also not significant ($p=0.235$).

Comparison to previous studies of split-belt adaptation

To determine whether we observed a reduction in energy cost in the U group or an increase in energy cost of the K group that led to the observed metabolic cost differences, we compared the energy cost at 10 minutes of adaptation (LA) to previous studies using the same 3:1 split-belt ratio at similar timepoints during adaptation. We report results as mean ± standard errors in this

section to match previous studies (Finley et al. 2013). The metabolic cost during LA for the K group in our study was 3.53 ± 0.19 W/kg and for the U group was 3.18 ± 0.14 W/kg. Finley *et al.*, 2013 measured a metabolic cost of 3.05 ± 0.24 W/kg at 12 minutes of adaptation. Sánchez *et al.*, 2021 measured a metabolic cost of 2.37 ± 0.17 W/kg at 10 minutes of a 45-minute adaptation trial.

Using previous data by the authors (Sánchez et al. 2021), we ran a one-way ANOVA with group as the independent variable and metabolic cost at minute 10 as the dependent variable (Sánchez 2021 data and U group N=15, K group N=14) and found that the cost reported in Sánchez 2021 was significantly lower than the K and U groups: one way ANOVA $p < 0.001$, post-hoc $p < 0.001$ Sanchez 2021 compared to K, $p = 0.025$ Sanchez 2021 compared to U and $p = 0.013$ for K compared to the U group. The comparison of all these studies indicates both an increased cost in participants who have detailed knowledge of task duration (K group) relative to Finley 2013, and a reduction in cost in individuals who plan to maintain the task for a prolonged time (U group and Sanchez 2021).

We also compared step length asymmetry across the three groups. We did not observe significant differences in step length asymmetry between Sanchez 2021 and participants in the U or K groups (one-way ANOVA $p = 0.208$).

Discussion

While studies of motor adaptation have consistently observed reductions in metabolic cost that parallel the kinematic and kinetic changes during adaptation (Finley et al. 2013; Huang et al. 2012; Sánchez et al. 2021), whether metabolic cost drives these biomechanical changes is still under debate. To test whether metabolic cost indeed shapes the adaptation of locomotor behaviors or is a byproduct of adaptation, we manipulated the perception of task duration during split-belt walking adaptation in neurotypical young adults. Manipulating perception of task duration can lead to differences in performance between individuals in order to maintain an energetic reserve (Hampson *et al.*, 2001). Our results support the idea that when participants believe they must sustain a task for a prolonged time, they prioritize energy reduction and markedly reduce energy cost. When participants know they must sustain a task for a short time,

they do not reduce energy cost. However, these differences in metabolic cost are not associated with clear differences in walking biomechanics. We observed small biomechanical differences between groups in step lengths and positive work by the right/slow leg, which might explain the marked metabolic differences. We also observed more consistent behaviors across individuals who believed they would sustain the task for a longer time as shown by the small variances in the U group, pointing at the idea that these individuals avoided any movement strategies associated with potential energetic penalties. Thus, we conclude that metabolic cost reduction is prioritized in tasks that need to be sustained for a prolonged time, yet the mechanisms underlying metabolic cost reductions are idiosyncratic and vary between participants, and do not depend on biomechanical strategies to reduce energy cost.

During adaptation, we observed different metabolic costs between groups such that by the end of adaptation, the metabolic cost in the U group was 0.351 W/kg or ~12% lower than in the K group. Note that this difference is about half the overall reduction in metabolic cost from EA to LA reported in (Finley et al. 2013), and twice that observed from 15 to 45 minutes of adaptation reported in (Sánchez et al. 2021). This difference in metabolic cost is well above the 5% energetic savings that have been reported to lead to walking adjustments in order to reduce energetic cost (Selinger et al. 2015). The lack of reduction in metabolic cost for the K group might indicate that for a relatively short task of 10 minutes, our population of healthy young adults might not prioritize metabolic cost reduction, consistent with recent studies showing prioritization of other cost functions over metabolic cost (McDonald et al. 2022). Note that the split-belt adaptation task duration is comparable to that used in (Finley et al. 2013) of 12 minutes, and (Buurke et al. 2018) of 9 minutes, where they observed significant reductions in metabolic cost during adaptation. The only difference in our study is the added updates about remaining time, which potentially provided contextual cues that led to an increased sympathetic response in the K group (Gordan et al. 2015) as indicated by the higher heart rate and metabolic cost. In contrast, the 12% reduction in cost in the U group indicates that for a perceived task duration three times longer than the actual task duration, metabolic cost reduction to maintain an energetic reserve is a priority, whereas for shorter tasks, it is not.

Our results show that metabolic cost can be reduced through biomechanical and non-biomechanical factors. Studies of motor adaptation during reaching showed continued metabolic power reductions even when kinematics and muscle activation patterns had already reached stable behaviors (Huang et al. 2012; Huang and Ahmed 2014). These findings during reaching

adaptation, in combination with our results, point at additional mechanisms that might explain metabolic cost, beyond the cost of reaching or walking, such as autonomic response associated with the different contexts (walking for 10 min or the instruction to walk "around 30 minutes") (Robinson *et al.*, 1966; Lay *et al.*, 2002), differences in neural activity during the task between groups as they received regular updates vs. no information (Hasenstaub et al. 2010), value placed on task duration (Summerside et al. 2018), postural demands and comfort during the task between groups (Rosenbaum et al. 1995) among others. These non-biomechanical factors are evidenced by differences in heart rate that accompany the metabolic cost differences, thus pointing at differences in sympathetic response (Stauss 2003) that might influence metabolic cost. Overall, our findings show that metabolic cost during adaptation can be significantly reduced without being able to attribute these changes to specific biomechanical changes.

Participants in our study did not perceive a reduced effort that aligned with the differences in metabolic cost between groups. Our previous work has also indicated that people's perception of effort does not necessarily reflect their actual metabolic expenditure, even when metabolic cost between conditions varies by 30% (Sánchez et al. 2017). In a prior study, participants used visual feedback to adjust their foot placements, leading to varying levels of step length asymmetry and metabolic costs (Sánchez et al. 2017). These findings suggest that the perception of effort is influenced by factors beyond just energy expenditure and may be related to the perceived difficulty of the task (Sparrow and Newell 1998), which might relate to both physiological and psychological factors associated with an unknown task duration (Morgan 1994) and have a greater influence on metabolic cost than what we observed in previous studies.

The lack of association between biomechanical variables and metabolic cost contrasted previous work. Contrary to previous studies (Bogard et al. 2024; Finley et al. 2013; Sánchez et al. 2021), the change in metabolic cost during adaptation was not associated with a change in step length asymmetry. Contrary to previous studies, the reduction in positive work by the left/fast leg here was not associated with the reduction in metabolic cost (Sánchez et al. 2021). Also, surprisingly, here, the reduction in work seemed to be associated with the reduction in work by the slow leg, but only in participants in the U group, also contrary to previous work (Sánchez et al. 2019, 2021). Based on our previous studies, it might be the case that any change in work by the left/fast leg might incur a large energetic penalty, such that individuals in the U group modify the work in the right/slow leg to avoid energetic penalties when modifying

the larger work generated by the fast leg (Sánchez et al. 2021). What is also interesting is the narrower variability in the U group, as evidenced by the smaller standard deviations of the biomechanical variables. This also supports the idea of the U group avoiding metabolic penalties by keeping a more consistent behavior (Abram et al. 2022). Thus, our findings indicate that the metabolic cost reductions cannot be attributed to a change in a single biomechanical variable, but small coordinated changes across multiple biomechanical variables.

While step length asymmetry has been consistently used as a metric to track adaptation (Leech et al. 2018; Reisman et al. 2005; Roemmich et al. 2016; Roemmich and Bastian 2015; Sánchez et al. 2019), our study shows that it cannot serve as a proxy to track adaptation when the goal prioritized is metabolic cost. Despite no differences in step length asymmetry, we observed marked differences in metabolic cost between groups. We also observe a lack of association between step length asymmetry and positive work by the right/slow leg, the latter being marginally associated with metabolic cost reduction. While step length asymmetry provides a simple measure that is easily observable and easily quantified, it is also a global measure that involves both limbs and might not be sensitive enough to detect smaller adjustments during adaptation. This is also supported by our results showing that while step length asymmetry did not show differences between groups, the right/slow step lengths did differ between groups. Thus, research should quantify step length asymmetry and the underlying factors that produce this asymmetry to understand locomotor adaptation further.

Our study has multiple limitations. First, we did not systematically assess fitness levels across participants. While our criteria excluded people who exercised more than 10 hours/week, we did not exclude sedentary participants and, as such, cannot determine if either group had different levels of fitness, which has been shown to affect energy expenditure (Morgan et al. 1995) and adaptation (Brinkerhoff et al. 2023). As previously mentioned, the different contexts created by the instructions provided between groups may have led to differences in autonomic response. While we measured heart rate, we were not able to obtain measures of heart rate variability to determine sympathetic activation during the task in each group that could explain differences in heart rate and metabolic cost (Stauss 2003). It might also be the case that the metabolic cost differences observed between groups are due to differences in muscle coactivation during adaptation, detected using electromyography of both lower extremity and postural muscles, which cannot be detected using the biomechanical variables we measured. Finally, our results indicate that metabolic cost reduction is not attributed to a single biomechanical variable but

necessitates significant coordination of gait biomechanics. While we are well-powered to measure differences in metabolic cost between groups (post-hoc power 0.84), the changes in gait biomechanics that occur during our adaptation protocol that might explain metabolic cost reduction are small in magnitude and might involve a combination of kinematic, kinetic, and temporal changes. There are multiple aspects of gait that individuals can adapt to reduce metabolic cost, including limb power, positive and negative work by each leg, and the timing of power and work generation during the gait cycle (Selgrade et al. 2017a). As such, a lack of consistency in strategies to reduce energy costs among participants might have hindered us from detecting a specific strategy in our sample. It could be that we can predict metabolic cost using a multidimensional model with interactions between multiple biomechanical terms, which would require a larger sample size than the one used here.

Conclusion

We observed significant differences in the metabolic cost of split-belt walking in individuals who were uncertain of task duration compared to individuals who had definitive information about task duration without clear differences in biomechanical variables that explained metabolic cost reduction. Our findings indicate that metabolic cost reduction is not a by-product of the walking pattern; metabolic cost reduction is prioritized during movements that need to be sustained over a prolonged time to maintain an energetic reserve, and this metabolic cost reduction can occur using a combination of biomechanical, physiological, and psychological factors. The simplest implication of our findings is that the metabolic cost associated with a task depends on perceived task duration, and as such, task duration information should be controlled and informed during experimental paradigms. Our findings also imply that information about task duration can be manipulated to elicit different cardiopulmonary and metabolic responses, which can have implications for movement interventions that require achieving different levels of cardiovascular intensity or aim to prevent fatigue in patient populations. Future work should assess how the perception of task duration might affect locomotor learning, metabolic cost, and cardiovascular responses in pathological populations, to determine whether these reductions can occur even with impaired biomechanics.

Data Availability

Data used for analyses in this manuscript are available at:
[https://github.com/GaitBehaviorLab/TimeUncertaintyData/tree/d0437e6242c050fdbc35f4aafad1f f7b3a0f8d1c](https://github.com/GaitBehaviorLab/TimeUncertaintyData/tree/d0437e6242c050fdbc35f4aafad1ff7b3a0f8d1c)

Acknowledgments

We would like to thank Lorcan Graham, Makenna Howard, Julia Malloy, Khloe McCarthy, Marissa Vasquez, Laura Corona, Jessica Cota, Emi Heisterkamp, Brandon Nakano, Rebecca Wingen, and Joseph Yeung for helping with data collection for this study. We would also like to thank Alejandro Aguirre-Ramirez and Hailey Hashiguchi for helping with data collection and labeling. Finally, we would like to thank all participants who volunteered for this study.
This work was supported by grants NIH NCMRR R03HD107630 and NIH NCATS R03TR004248 to N. Sánchez.

REFERENCES

Abram SJ, **Poggensee KL**, **Sánchez N**, **Simha SN**, **Finley JM**, **Collins SH**, **Donelan JM**. General variability leads to specific adaptation toward optimal movement policies. *Current Biology* 32: 2222-2232.e5, 2022.

Bakkum A, **Marigold DS**. Learning from the Physical Consequences of Our Actions Improves Motor Memory. *eNeuro* 9: ENEURO.0459-21.2022, 2022.

Betschart M, **Lauzière S**, **Miéville C**, **McFadyen BJ**, **Nadeau S**. Changes in lower limb muscle activity after walking on a split-belt treadmill in individuals post-stroke. *Journal of Electromyography and Kinesiology* 32: 93–100, 2017.

Bogard AT, **Hembree TG**, **Pollet AK**, **Smith AC**, **Ryder SC**, **Marzloff GE**, **Tan AQ**. Intermittent hypoxia-induced enhancements in corticospinal excitability predict gains in motor learning and metabolic efficiency. 2024.

Bogard AT, **Hemmerle MR**, **Smith AC**, **Tan AQ**. Enhanced motor learning and motor savings after acute intermittent hypoxia are associated with a reduction in metabolic cost. *The Journal of Physiology* JP285425, 2023.

Brinkerhoff SA, **Sánchez N**, **Roper JA**. Habitual exercise evokes fast and persistent adaptation during split-belt walking. *PLoS ONE* 18: e0286649, 2023.

Bruijn SM, **van Dieën JH**. Control of human gait stability through foot placement. *J R Soc Interface* 15: 20170816, 2018.

Bruijn SM, **Van Impe A**, **Duysens J**, **Swinnen SP**. Split-belt walking: adaptation differences between young and older adults. *Journal of Neurophysiology* 108: 1149–1157, 2012.

Buurke TJW, **Lamoth CJC**, **Vervoort D**, **Van Der Woude LHV**, **Den Otter R**. Adaptive control of dynamic balance in human gait on a split-belt treadmill. *Journal of Experimental Biology* jeb.174896, 2018.

Charlton JM, **Eng JJ**, **Li LC**, **Hunt MA**. Learning Gait Modifications for Musculoskeletal Rehabilitation: Applying Motor Learning Principles to Improve Research and Clinical Implementation. *Physical Therapy* 101: pzaa207, 2021.

Cohen J. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed. Academic Press, New York, 1988.

Dietz V, **Zijlstra W**, **Duysens J**. Human neuronal interlimb coordination during split-belt locomotion. *Exp Brain Res* 101: 513–520, 1994.

Donelan JM, **Kram R**, **Kuo AD**. Mechanical work for step-to-step transitions is a major determinant of the metabolic cost of human walking. *Journal of Experimental Biology* 205: 3717–3727, 2002a.

Donelan JM, **Kram R**, **Kuo AD**. Simultaneous positive and negative external mechanical work in human walking. *Journal of Biomechanics* 35: 117–124, 2002b.

Ellmers TJ, **Durkin M**, **Sriranganathan K**, **Harris DJ**, **Bronstein AM**. The influence of postural threat-induced anxiety on locomotor learning and updating. *Journal of Neurophysiology* 131: 562–575, 2024.

Finley JM, **Bastian AJ**, **Gottschall JS**. Learning to be economical: the energy cost of walking tracks motor adaptation: Split-belt adaptation reduces metabolic power. *The Journal of Physiology* 591: 1081–1095, 2013.

Gonzalez-Rubio M, **Velasquez NF**, **Torres-Oviedo G**. Explicit Control of Step Timing During Split-Belt Walking Reveals Interdependent Recalibration of Movements in Space and Time. *Front Hum Neurosci* 13: 207, 2019.

Gordan R, **Gwathmey JK**, **Xie L-H**. Autonomic and endocrine control of cardiovascular function. *WJC* 7: 204, 2015.

Hampson DBase, Gibson AS, **Lambert MI**, **Noakes TD**. The Influence of Sensory Cues on the Perception of Exertion During Exercise and Central Regulation of Exercise Performance: *Sports Medicine* 31: 935–952, 2001.

Hasenstaub A, **Otte S**, **Callaway E**, **Sejnowski TJ**. Metabolic cost as a unifying principle governing neuronal biophysics. *Proc Natl Acad Sci USA* 107: 12329–12334, 2010.

Hinton DC, **Thiel A**, **Soucy J-P**, **Bouyer L**, **Paquette C**. Adjusting gait step-by-step: Brain activation during split-belt treadmill walking. *NeuroImage* 202: 116095, 2019.

Huang HJ, **Ahmed AA**. Reductions in muscle coactivation and metabolic cost during visuomotor adaptation. *Journal of Neurophysiology* 112: 2264–2274, 2014.

Huang HJ, **Kram R**, **Ahmed AA**. Reduction of Metabolic Cost during Motor Learning of Arm Reaching Dynamics. *Journal of Neuroscience* 32: 2182–2190, 2012.

Huang VS, **Shadmehr R**. Persistence of Motor Memories Reflects Statistics of the Learning Event. *Journal of Neurophysiology* 102: 931–940, 2009.

Jacobsen NA, **Ferris DP**. Electrocortical activity correlated with locomotor adaptation during split-belt treadmill walking. *The Journal of Physiology* 601: 3921–3944, 2023.

Joseph T, **Johnson Base, Battista RA**, **Wright G**, **Dodge C**, **Porcari JP**, **De Koning JJ**, **Foster C**. Perception of Fatigue during Simulated Competition. *Medicine & Science in Sports & Exercise* 40: 381–386, 2008.

Kambic RE, **Roemmich RT**, **Bastian AJ**. Joint-level coordination patterns for split-belt walking across different speed ratios. *Journal of Neurophysiology* 129: 969–983, 2023.

Kowalsky DBase, Rebula JR, **Ojeda LV**, **Adamczyk PG**, **Kuo AD**. Human walking in the real world: Interactions between terrain type, gait parameters, and energy expenditure. *PLoS ONE* 16: e0228682, 2021.

Lay BS, **Sparrow WA**, **Hughes KM**, **O'Dwyer NJ**. Practice effects on coordination and control, metabolic energy expenditure, and muscle activation. *Human Movement Science* 21: 807–830, 2002.

Leech KA, **Day KA**, **Roemmich RT**, **Bastian AJ**. Movement and perception recalibrate differently across multiple days of locomotor learning. *Journal of Neurophysiology* 120: 2130–2137, 2018.

Malone LA, **Bastian AJ**. Thinking About Walking: Effects of Conscious Correction Versus Distraction on Locomotor Adaptation. *Journal of Neurophysiology* 103: 1954–1962, 2010.

Malone LA, **Bastian AJ**. Age-related forgetting in locomotor adaptation. *Neurobiology of Learning and Memory* 128: 1–6, 2016.

Marigold DS, **Patla AE**. Age-related changes in gait for multi-surface terrain. *Gait & Posture* 27: 689–696, 2008.

Mawase F, **Haizler T**, **Bar-Haim S**, **Karniel A**. Kinetic adaptation during locomotion on a split-belt treadmill. *Journal of Neurophysiology* 109: 2216–2227, 2013.

McAllister MJ, **Blair RL**, **Donelan JM**, **Selinger JC**. Energy optimization during walking involves implicit processing. *Journal of Experimental Biology* 224: jeb242655, 2021.

McDonald KA, **Cusumano JP**, **Hieronymi A**, **Rubenson J**. Humans trade off whole-body energy cost to avoid overburdening muscles while walking. *Proceedings of the Royal Society B: Biological Sciences* 289, 2022.

Morgan DW, **Bransford DR**, **Costill DL**, **Daniels JT**, **Howley ET**, **Krahenbuhl GS**. Variation in the aerobic demand of running among trained and untrained subjects. *Med Sci Sports Exerc* 27: 404–409, 1995.

Morgan WP. Psychological components of effort sense. *Med Sci Sports Exerc* 26: 1071–1077, 1994.

Park S, **Finley JM**. Manual stabilization reveals a transient role for balance control during locomotor adaptation. *Journal of Neurophysiology* 128: 808–818, 2022.

Price M, **Huber ME**, **Hoogkamer W**. Minimum effort simulations of split-belt treadmill walking exploit asymmetry to reduce metabolic energy expenditure. *Journal of Neurophysiology* 129: 900–913, 2023.

Reisman DS, **Bastian AJ**, **Morton SM**. Neurophysiologic and Rehabilitation Insights From the Split-Belt and Other Locomotor Adaptation Paradigms. *Physical Therapy* 90: 187–195, 2010.

Reisman DS, **Block HJ**, **Bastian AJ**. Interlimb Coordination During Locomotion: What Can be Adapted and Stored? *Journal of Neurophysiology* 94: 2403–2415, 2005.

Richardson JTE. Eta squared and partial eta squared as measures of effect size in educational research. *Educational Research Review* 6: 135–147, 2011.

Robinson BF, **Epstein SE**, **Beiser GD**, **Braunwald E**. Control of Heart Rate by the Autonomic Nervous System: Studies in Man on the Interrelation Between Baroreceptor Mechanisms and Exercise. *Circulation Research* 19: 400–411, 1966.

Roemmich RT, **Bastian AJ**. Two ways to save a newly learned motor pattern. *Journal of Neurophysiology* 113: 3519–3530, 2015.

Roemmich RT, **Long AW**, **Bastian AJ**. Seeing the Errors You Feel Enhances Locomotor Performance but Not Learning. *Current Biology* 26: 2707–2716, 2016.

Roemmich RT, **Stegemöller EL**, **Hass CJ**. Lower extremity sagittal joint moment production during split-belt treadmill walking. *Journal of Biomechanics* 45: 2817–2821, 2012.

Rosenbaum DA, **Loukopoulos LD**, **Meulenbroek RGJ**, **Vaughan J**, **Engelbrecht SE**. Planning reaches by evaluating stored postures. *Psychological Review* 102: 28–67, 1995.

Sánchez N, **Park S**, **Finley JM**. Evidence of Energetic Optimization during Adaptation Differs for Metabolic, Mechanical, and Perceptual Estimates of Energetic Cost. *Sci Rep* 7: 7682, 2017.

Sánchez N, **Simha SN**, **Donelan JM**, **Finley JM**. Taking advantage of external mechanical work to reduce metabolic cost: the mechanics and energetics of split-belt treadmill walking. *J Physiol* 597: 4053–4068, 2019.

Sánchez N, **Simha SN**, **Donelan JM**, **Finley JM**. Using asymmetry to your advantage: learning to acquire and accept external assistance during prolonged split-belt walking. *Journal of Neurophysiology* 125: 344–357, 2021.

Seethapathi N, **Clark Base, Srinivasan M**. Exploration-based learning of a step to step controller predicts locomotor adaptation. Neuroscience.

Selgrade BP, **Thajchayapong M**, **Lee GE**, **Toney ME**, **Chang Y-H**. Changes in mechanical work during neural adaptation to asymmetric locomotion. *Journal of Experimental Biology* jeb.149450, 2017a.

Selgrade BP, **Toney ME**, **Chang Y-H**. Two biomechanical strategies for locomotor adaptation to split-belt treadmill walking in subjects with and without transtibial amputation. *Journal of Biomechanics* 53: 136–143, 2017b.

Selinger JC, **O'Connor SM**, **Wong JD**, **Donelan JM**. Humans Can Continuously Optimize Energetic Cost during Walking. *Current Biology* 25: 2452–2456, 2015.

Smith MA, **Ghazizadeh A**, **Shadmehr R**. Interacting Adaptive Processes with Different Timescales Underlie Short-Term Motor Learning. *PLoS Biol* 4: e179, 2006.

Sparrow WA, **Newell KM**. Energy expenditure and motor performance relationships in humans learning a motor task. *Psychophysiology* 31: 338–346, 1994.

Sparrow WA, **Newell KM**. Metabolic energy expenditure and the regulation of movement economy. *Psychonomic Bulletin & Review* 5: 173–196, 1998.

Stauss HM. Heart rate variability. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 285: R927–R931, 2003.

Stenum J, **Choi JT**. Step time asymmetry but not step length asymmetry is adapted to optimize energy cost of split-belt treadmill walking. *The Journal of Physiology* 598: 4063–4078, 2020.

Summerside EM, **Kram R**, **Ahmed AA**. Contributions of metabolic and temporal costs to human gait selection. *J R Soc Interface* 15: 20180197, 2018.

Swart J, **Lamberts RP**, **Lambert MI**, **St Clair Gibson A**, **Lambert EV**, **Skowno J**, **Noakes TD**. Exercising with reserve: evidence that the central nervous system regulates prolonged exercise performance. *British Journal of Sports Medicine* 43: 782–788, 2009.

Torres-Oviedo G, **Bastian AJ**. Seeing Is Believing: Effects of Visual Contextual Cues on Learning and Transfer of Locomotor Adaptation. *Journal of Neuroscience* 30: 17015–17022, 2010.

Torres-Oviedo G, **Vasudevan E**, **Malone L**, **Bastian AJ**. Locomotor adaptation. In: *Progress in Brain Research*. Elsevier, p. 65–74.

Task duration influences metabolic cost of adaptation

Table 1

Group	N	Age (yrs)	Height (cm)	Weight (kg)	Sex
K	14	25.71	165.82	67.67	9F/5M
U	15	26.00	167.91	66.59	9F/6M
		p=0.851	p=0.532	p=0.824	p=0.999

Participant demographics for both experimental groups. P-values for independent samples t-tests and for Chi-Square tests show no differences in group demographics.

Task duration influences metabolic cost of adaptation

Figure 1.

Figure 1. Experimental setup and protocol. A) Experimental setup showing experimental groups, metabolic mask, and marker location to calculate spatiotemporal variables. B) Split-belt configuration. For all participants, the left belt was set at 1.5 m/s and the right belt at 0.5 m/s. 27 participants reported the right leg as the leg they would use to kick a ball. C) Experimental protocol. Bars indicate each belt, and numbers indicate belt speed. Bar width indicates the duration of each experimental condition. Participants reported their rate of perceived exertion during the timepoints identified by the arrows. S: Standing. B: Baseline. EA: Early adaptation. MA: Mid-adaptation. LA: Late adaptation. EP: Early post-adaptation. LP: Late post-adaptation. GTRO: Greater Trochanter. LEPI: Lateral Epicondyle. LMAL: Lateral Malleolus.

Figure 2.

Figure 2. Metabolic cost, heart rate, and perception of effort. A) Net metabolic power timeseries during the experiment between the Knowledge (Red) and Uncertainty (Blue) groups. Time windows indicating Base (B), EA, MA, LA, EP, and LP are indicated in the figure. B) Average metabolic cost between groups during 30 seconds corresponding to Base, EA, MA, LA, EP, and LP. In the U group, metabolic cost during EA was higher than MA (p<0.001) and LA (p<0.001). C) Average heart rate between groups during 30 seconds corresponding to Base, EA, MA, LA, EP and LP. D) Rating of perceived exertion as indicated by participants during Base, EA, MA, LA, EP, and LP. +: Significant differences between EA and Base, EP and LP (p<0.001). *: Significant differences between EA and all other timepoints (p<0.005). K: Knowledge. U: Uncertainty. HR: Heart Rate. RPE: Rating of Perceived Exertion. Other abbreviations as in Figure 1.

Figure 3.

Figure 3. Step length asymmetry and step lengths. A) Step length asymmetry during the experiment between the Knowledge (Red) and Uncertainty (Blue) groups. Time windows indicating Base (B), EA, MA, LA, EP, and LP are indicated in the figure. B) Distribution of average step length asymmetry between groups during the 5 strides corresponding to Base, EA, MA, LA, EP, and LP. C) Distribution of average step lengths for the left/fast leg between groups during the 5 strides corresponding to Base, EA, MA, LA, EP, and LP. D) Distribution of average step lengths for the right/slow leg between groups during the 5 strides corresponding to Base, EA, MA, LA, EP, and LP. *p<0.050 significant differences from the Base condition. Abbreviations as in Figures 1 and 2.

Figure 4.

Figure 4. Work by the fast and slow legs. A) Positive work by the fast leg during the experiment between the Knowledge (Red) and Uncertainty (Blue) groups. Time windows indicating Base, EA, MA, LA, EP and LP are indicated in the figure. B) Positive work by the slow

leg during the experiment between the Knowledge (Red) and Uncertainty (Blue) groups. C) Distribution of average positive work by the left/fast leg between groups during the 5 strides corresponding to Base, EA, MA, LA, EP, and LP. D) Distribution of average positive work by the right/slow leg between groups during the 5 strides corresponding to Base, EA, MA, LA, EP, and LP. E) Distribution of average negative work by the fast leg between groups during the 5 strides corresponding to Base, EA, MA, LA, EP, and LP. F) Distribution of average negative work by the slow leg between groups during the 5 strides corresponding to Base, EA, MA, LA, EP, and LP. Abbreviations as in Figures 1 and 2.

Figure 5

Figure 5. Associations between biomechanical and metabolic measures. A) Association between change in step length asymmetry and change in metabolic cost from EA to LA. B) Association between change in positive work by the left/fast leg and change in metabolic cost from EA to LA. C) Association between change in positive work by the right/slow leg and change in metabolic cost from EA to LA. D) Association between change in net work and change in metabolic cost from EA to LA. E) Association between change in positive work by the left/fast leg and change in step length asymmetry from EA to LA. F) Association between change in positive work by the right/slow leg and change in step length asymmetry from EA to LA.

Task duration influences metabolic cost of adaptation

Supplementary Figure 1

Supplementary Figure 1. Individual step length asymmetry data during adaptation. N=14 participants in the Knowledge group and N=15 participants in the uncertainty group.

Task duration influences metabolic cost of adaptation

Supplementary Figure 2

Supplementary Figure 2. Individual step length asymmetry data during post-adaptation.

N=14 participants in the Knowledge group and N=15 participants in the uncertainty group.