The mammary glands of cows abundantly display receptors for circulating avian H5 viruses

María Ríos Carrasco[1], Andrea Gröne[2], Judith M.A. van den Brand[2], Robert P. de Vries[1*]

[1] Department of Chemical Biology & Drug Discovery, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Universiteitsweg 99, 3584CG, Utrecht, The Netherlands

[2] Division of Pathology, Department of Biomolecular Health Sciences, Faculty of Veterinary Medicine of Utrecht University, Utrecht University, Yalelaan 1, 3584 CL Utrecht, The Netherlands

Abstract

Influenza A viruses (IAV) from the H5N1 2.3.4.4b clade are circulating in dairy farms in the United States of America, and goat infections have also been reported. These ruminants were presumed not to be hosts for IAVs. Previously, IAV-positive mammalian species were hunters and scavengers, possibly getting infected while feeding on infected birds. It is now presumed that H5N1 viruses that circulate in US dairy cattle transmit through a mammary gland route, in contrast to transmission by aerosols via the respiratory tract, although the latter cannot be excluded. The receptor display in the mammary and respiratory tract in cows for IAVs is unknown. Here, we used recombinant HA proteins representing current circulating and classical H5 viruses to determine the distribution of IAV receptors in the respiratory and mammary tract tissues of cows and goats. Most of the sialome of the cow and goat respiratory tract is lined with sialic acid modifications such as N-glycolyl and O-acetyl, which are not bound by IAV. Interestingly, the H5 protein representing the cow isolates bound significantly in the mammary gland, whereas classical H5 proteins failed to do so. Furthermore, whereas the 9-O-acetyl modification is prominent in all tissues tested, the 5-N-glycolyl modification is not, resulting in the display of receptors for avian IAV hemagglutinins. This could explain the high levels of virus found in these tissues and milk, adding supporting data to this possible virus transmission route.

Introduction

Since late 2020, we have been experiencing an unprecedented global outbreak of high pathogenic H5Nx influenza A viruses (IAV) (1, 2). These viruses circulate year-round from the Northern to Southern hemisphere and cause incredibly high mortality in avian species with significant transmission to mammals (3-5). The transmission route to mammals is under debate as most infected mammals are hunters and scavengers, possibly getting infected while

consuming bird remains (6-9). Seal infections, on the other hand, have been known to occur

for decades due to their proximity to wild waterfowl (10); however, the spread into pinnipeds

of the southern hemisphere was unprecedented. The appearance of the viruses in the

Antarctic and the current zoonotic event to ruminants add even more novelties to this

remarkable outbreak (11). The latest isolation of IAVs in cows and goats was an unexpected

transmission event as these species were believed not to be hosts of IAVs, while they are

hosts for influenza D viruses (IDV) (12, 13). However, it quickly became apparent that

mammary tract tissues are probably essential in the transmission route, as milk samples most

frequently contain high viral titers (14).

The molecular determinant of the current widespread H5Nx viruses is currently unknown.

While the E627K mutation in the polymerase is commonly found in viruses isolated from

mammalian species, the moment of occurrence is whether viruses with this mutation circulate

in birds or is this mutation is immediately selected in mammals. Only the human isolate

associated with the current cow outbreak contains this mutation (15). The HA gene has been

remarkably stable with no mutations in the receptor binding site (RBS) in the 2.3.4.4b virus for

years. Genotypes such as 2.3.4.4a,c, and e harbor some RBS mutations; however, these

clades are minority species. Thus, it is likely that 2.3.4.4b virus HA proteins already have an

optimal receptor binding specificity, allowing their panzootic nature.

Most IAVs use sialic acids (SIA) to enter a host cell. The glycome that presents these SIAs in

the respiratory tract of farm animals such as goats and cows is poorly defined (16), in contrast

to several other bovine proteins such as submaxillary gland mucins and fetuin (17-20). It is,

however, known that ruminants display a set of modified SIAs that other IAV host species do

not. These include the N-glycolyl modification at the C5 position (5-N-glycolyl), which is

created by the cytidine monophospho-N-acetylneuraminic acid hydroxylase (CMAH) (21).

CMAH is a mammalian-specific enzyme that is non-functional in various influenza hosts,

including humans, ferrets, dogs, and seals (22). Cows and goats express a functional enzyme

and thus abundantly display Neu5Gc (23) which is not a receptor for most influenza A viruses

(24). The other abundant SIA modifications in the cow respiratory tract are O-acetyls, which

can be present on the 4, 7, 8, and 9 positions. The latter is the essential receptor for influenza

C, D (25), and a variety of coronaviruses (26, 27), but a non-ligand for influenza A virus (28-

30). The current dogma is that these SIA modifications, not present in the avian reservoir, are

decoy or blocking moieties for IAVs.

To determine if 2.3.4.4b H5 influenza A viruses can bind to available receptors in the

respiratory and mammary tract of cows, we used formalin-fixed and paraffin-embedded

(FFPE) nasal, tracheal, lung, and mammary gland tissues of cows and goats (31). The animals

were submitted to the Veterinary Pathology Diagnostic Centre (VPDC) for postmortem

evaluation. No animals were euthanized for this study.

Results

2.3.4.4b viruses isolated from goats and dairy cows have a conserved receptor

binding site.

The receptor binding site, the HA of IAVs, is composed of the 190 helices (AA180-200), the

130-(AA130-140), and 220-(AA220-230) loops. Since the introduction of the 2.3.4.4b clade a

decade ago, the receptor binding domain has remained relatively conserved (Fig 1).

Significant changes in 2.3.4.4 viruses, compared to the classical A/Vietnam/1203/2004

(H5VN) and A/Indonesia/05/2005 (H5IN), include the 130 loop, the loss of a glycosylation site

at position 158, and significant perturbations in the 190 helix and the 222 and 227 positions in

the 220-loop that are directly involved in receptor binding (32, 33). Taking

A/duck/France/161108h/2016 (H5FR) as a reference, we only observe minimal amino acid

changes in the recent North American mammalian-derived viruses. The mutations observed

are outside the canonical RBS, namely L122Q, T144A, T199I, and V214A (H3 numbering). Of

note, some cow sequences isolated later now contain A160T and thus restore the

glycosylation site at position 158. Conclusively, although the 2.3.4.4b H5 viruses have been

circulating for several years around the globe in different hosts, the receptor binding domain

remained conserved.

Figure 1. **HA receptor binding site amino acid alignment of the H5 hemagglutinins used in this study.** Alignment of the receptor binding site residues with amino acid positions (H3 numbering) indicated above the alignment, non-conserved residues highlighted in black, dots indicating identical amino acids. Several mammalian sequences isolate in North America are shown to demonstrate the close relationship.

Contemporary 2.3.4.4b HA proteins bind efficiently to α2,3 linked Neu5Ac containing sialosides in mammary tract tissues of cows.

A central observation of IAV-infected dairy cows is mammary gland infections (14). To determine if this gland has receptors for H5 IAVs to support active replication, we used both classical and contemporary H5 proteins for tissue binding studies. We used the classical H5 derived from A/Indonesia/05/2005 (H5IN), A/Vietnam/1203/2004 (H5VN), and the Y161A (H5VN Y161A) mutant that confers binding to Neu5Gc (34). To detect 9-O-acetylated structures, we employed an enzymatically inactive influenza D virus hemagglutinin esterase fusion protein (D/OK) (12). We also used commonly employed plant lectins SNA and MAL-II (biotinylated, Vector laboratories) to determine the distribution of α2,3 and α2,6 linked SIAs. To detect α2,6 linked SIAs, we employed a human H1 HA derived from A/Puerto Rico/8/34 (H1PR8) (35).

We used FFPE tissues of the mammary glands of three cows and applied our library of plant-IAV- and IDV-derived glycan-binding proteins. A protein that binds a glycan is defined as a lectin, and an overview of the specificities of all lectins used is provided in the supplementary section (Fig S1). No signal was observed when we applied our antibody mix as a negative control. The classical H5 proteins, H5IN & H5VN, showed variable binding in the mammary gland, with H5IN binding significantly to cows 1 and 2 but not 3, exemplifying biological variability (Fig 2). The H5VN Y161A mutant did not bind the alveoli but bound in the mammary gland to connective tissues and blood vessels. The staining of this lectin was not as intense as expected, as cows are known for displaying 5-N-glycolyl-modified SIAs in their respiratory tract (23). The IDV HEF, on the other hand, is bound throughout the mammary gland with high intensity, confirming the abundance of 9-O-acetylated SIA.

We then applied a 2.3.4.4a (H5KU) and a 2.3.4.4b HA (H5FR) and observed that the H5KU did not engage SIAs in the mammary gland, whereas the H5FR did. Intense staining is observed within the mammary glands of all three cows. A similar observation is made for both plant lectins, MAL-II and SNA. Notably, the H1 protein of a human H1N1 virus hardly bound in the mammary gland, indicating that SNA might bind α2,6 linked 5-N-glycolyl SIA (24). We confirmed the biological activity of all the lectins used using a traditional hemagglutination assay in which the proteins that bind erythrocytes form a mesh. All proteins can hemagglutinate chicken erythrocytes (Fig S2), except for the H5VN Y161A mutant, as there is no 5-N-glycolyl in avian species. H5KU also binds less avidly than the other H5 proteins but is biologically active.

Conclusively, the mammary gland of cows displays receptors for 2.3.4.4b H5 proteins and further confirms the possibility of virus binding as the first step of active replication in and transmission from this organ.

Figure 2. **Immunohistochemical analyzes of three cow mammary glands, stained with small library of viral and plant lectins.** The binding to mammary glands of three different cows was investigated for different IAV H5 proteins and a human H1 protein, as well as an IDV HEF and plant lectins MAL-II and SNA. AEC staining was used to visualize tissue binding.

The respiratory tract of cows and goats hardly display receptors for IAVs

In its wild waterfowl reservoir, IAVs are transmitted by the oral-fecal route, whereas in mammalian hosts, they are transmitted by the respiratory tract. Although all preliminary data points to a non-traditional transmission by milking practices, we did not want to exclude the respiratory tract of cows and goats as a possible transmission route.

We used FFPE upper (nasal or trachea) and lower (lung) respiratory tract tissues from cows and goats (N=2 and 1, respectively) and applied our library of lectins. The only H5 protein significantly binding to the upper respiratory tract of both cow and goat was the H5VN Y161A mutant, which binds to 5-N-glycolyl but that is not a receptor for circulating 2.3.4.4b H5N1 viruses. The lung tissues were bound by almost all HA proteins, including the human H1PR8,

Figure 3. Immunohistochemical analyzes of cow and goat upper and lower respiratory tract tissues. The binding to upper (nasal or tracheal) and lower (lung) respiratory tract tissues of a cow and goat was investigated for different IAV H5 proteins and a human H1 protein, as well as an IDV HEF and plant lectins MAL-II and SNA. AEC staining was used to visualize tissue binding.

with various intensities. Furthermore, the upper respiratory tract of both cow and goat also
abundantly displays 9-O-acetylated SIAs, as indicated by the high signal intensity of D/OK.
Both plant lectins bound to all four tissues tested, contrasting the avian and human HAs in the
tracheal and nasal tissues. This illustrates that plant-derived lectins, such as SNA and MAL-
II, are poor predictors of IAV receptor distribution (36, 37).

The inability of our H5 proteins to engage in the upper respiratory tract of cows and goats and
the high abundance of IAV non-receptors such as 5-N-glycolyl and 9-O-acetyl most likely
exclude this transmission route. Thus, receptors are available in the lungs, and lower
respiratory tract infections are often not efficiently transmitted and cause severe disease (38).

Discussion

Here, we demonstrate that the mammary gland of cows abundantly displays avian-type
receptors for circulating H5 viruses. We also show that this organ lacks human-type receptors,
which contradicts a previous study that only relied on plant lectins (16). Thus, we deem the
adaptation of 2.3.4.4b H5N1 viruses to human-type receptor specificity during replication in
the cow mammary gland unlikely. Using direct immunohistochemical binding of H5 proteins,
we demonstrate that the upper respiratory tract of goats and cows is devoid of receptors for
IAV. Thus, although the zoonotic transmission of 2.3.4.4b H5N1 viruses to ruminants is
unprecedented, we would suggest it does not pose an immediate pandemic risk, as there is
no need to adapt to human-type receptors. However, the continuous and widespread
circulation of these zoonotic viruses in primary livestock is a concern. Importantly, in human
infections, the virus is almost only isolated from eye swabs (39). Although conjunctivitis in
humans caused by AIV has been observed before and can be modeled in the ferret model
(40, 41), after such ocular inoculation, the virus could further spread to the upper respiratory
tract, in which H5N1 viruses might adapt to human-type receptors.

We focused our attention on the display of sialylated receptors for IAV HA proteins. We have
previously shown that the receptor specificity of HA proteins is very similar to that of whole
viruses, although with a lower degree of multivalency, leading to lower signals (24). However,
we can easily observe binding in the cow's mammary gland. Using multiple cows was vital as
the signals in cow 3 are significantly lower, which probably presents natural variation. We
would like to emphasize our use of an IDV HEF protein that stains almost all tested tissues
with high intensity. Cows are the reservoir for IDVs, and the abundant display of 9-O-
acetylated SIAs could be a block binding of IAV. Finally, we would like to point out that sole
use of plant lectins to study the differential display of avian- and human-type receptors (36,
37), leads to an over-interpretation as their binding profile is more promiscuous.

We expect that our studies into the display of SIA modifications in ruminants will aid in our
understanding of how different IAV viruses with distinct glycan specificities infect and transmit.
Indeed, a recent unofficial report paints a picture of severe infections in cows, which might be
related to lower respiratory tract infections, as ample receptors are available there (42). The
glycome of horses and pigs are not expected to differ greatly from cows, as both abundantly
display 5-N-glycolyl and 9-O-acetyls. While horses are hardly ever infected with avian IAVs,
pigs are, and experimental infection of pigs with 2.3.4.4b viruses have shown that they are
susceptible (43, 44), so a question is why we do not observe natural 2.3.4.4b H5N1 infections
in pigs.

Finally, MS-based analyses are essential to further map the sialome, and controlled infection
studies are desperately needed to understand this zoonotic event.

Acknowledgments

This research was made possible by funding from ICRAD, an ERA-NET co-funded under the
European Union's Horizon 2020 research and innovation programme
(https://ec.europa.eu/programmes/horizon2020/en), under Grant Agreement n°862605 (Flu-
Switch) to, and a Mizutani Foundation for Glycoscience Research Grant 2023 to Robert P.
de Vries.

We thank the staff of the Veterinary Pathology Diagnostic Centre of Utrecht University for
their assistance in collecting the tissues.

Materials and methods

Expression and purification of trimeric influenza A hemagglutinins

Recombinant trimeric IAV hemagglutinin ectodomain proteins (HA) were cloned into the pCD5
expression vector (an example is addgene plasmid #182546) in frame with a GCN4
trimerization motif (KQIEDKIEEIESKQKKIENEIARIKK), a super folder GFP or mOrange2 (45)
and the Twin-Strep-tag (WSHPQFEKGGGSGGGSWSHPQFEK); IBA, Germany). The
trimeric HAs were expressed in HEK293S GnTI(-) cells with polyethyleneimine I (PEI) in a 1:8
ratio (µg DNA:µg PEI) for the HAs as previously described. The transfection mix was replaced
after 6 hours by 293 SFM II suspension medium (Invitrogen, 11686029), supplemented with
sodium bicarbonate (3.7 g/L), primatone RL-UF (3.0 g/L, Kerry, NY, USA), glucose (2.0 g/L),
glutaMAX (1%, Gibco), valproic acid (0.4 g/L) and DMSO (1.5%). According to the
manufacturer's instructions, culture supernatants were harvested five days post-transfection
and purified with sepharose strep-tactin beads (IBA Life Sciences, Germany).

Protein histochemical tissue staining

Sections of formalin-fixed, paraffin-embedded cow (*Bos taurus*) and goat (*Capra aegagrus hircus*) were obtained from the Division of Pathology, Department of Biomolecular Health Sciences, Faculty of Veterinary Medicine of Utrecht University, the Netherlands. In the figures, representative images of at least two individual experiments are shown. Protein histochemistry was performed as previously described (46, 47). In short, tissue sections of 5µm were deparaffinized and rehydrated, after which antigens were retrieved by heating the slides in 10 mM sodium citrate (pH 6.0) for 10 min. Endogenous peroxidase was inactivated using 1% hydrogen peroxide in MeOH for 30 min at RT. Tissues were blocked at 4°C using 3% BSA (w/v) in PBS for at least 90 minutes. Subsequently, slides were stained for 90 minutes with 10µg/mL solution of precomplexed proteins of interest. HAs were precomplexed with human α-strep-tag primary antibody and goat-α-mouse-HRP secondary antibody at 4:2:1 molar ratio as previously described (12). For biotinylated plant lectins, we used streptavidin-HRP at 4:1 molar ratio. We used 3-amino-9-ethylcarbazole (AEC) (Sigma-Aldrich, Steinheim, Germany) to visualize protein binding. Tissue sections were then counterstained with hematoxylin and mounted with coverslips using AquaTex (Merck).

References

1. Lewis NS, Banyard AC, Whittard E, Karibayev T, Al Kafagi T, Chvala I, Byrne A, Meruyert Akberovna S, King J, Harder T, Grund C, Essen S, Reid SM, Brouwer A, Zinyakov NG, Tegzhanov A, Irza V, Pohlmann A, Beer M, Fouchier RAM, Akhmetzhan Akievich S, Brown IH. 2021. Emergence and spread of novel H5N8, H5N5 and H5N1 clade 2.3.4.4 highly pathogenic avian influenza in 2020. Emerg Microbes Infect 10:148-151.
2. Xie R, Edwards KM, Wille M, Wei X, Wong SS, Zanin M, El-Shesheny R, Ducatez M, Poon LLM, Kayali G, Webby RJ, Dhanasekaran V. 2023. The episodic resurgence of highly pathogenic avian influenza H5 virus. Nature 622:810-817.
3. Floyd T, Banyard AC, Lean FZX, Byrne AMP, Fullick E, Whittard E, Mollett BC, Bexton S, Swinson V, Macrelli M, Lewis NS, Reid SM, Núñez A, Duff JP, Hansen R, Brown IH. 2021. Encephalitis and death in wild mammals at a rehabilitation center after infection with highly pathogenic avian influenza A virus (H5N8), United Kingdom. Emerg Infect Dis 27:2856-2863.
4. Hiono T, Kobayashi D, Kobayashi A, Suzuki T, Satake Y, Harada R, Matsuno K, Sashika M, Ban H, Kobayashi M, Takaya F, Fujita H, Isoda N, Kimura T, Sakoda Y. 2023. Virological, pathological, and glycovirological investigations of an Ezo red fox and a tanuki naturally infected with H5N1 high pathogenicity avian influenza viruses in Hokkaido, Japan. Virology 578:35-44.
5. Agüero M, Monne I, Sánchez A, Zecchin B, Fusaro A, Ruano MJ, del Valle Arrojo M, Fernández-Antonio R, Souto AM, Tordable P, Cañás J, Bonfante F, Giussani E, Terregino C, Orejas JJ. 2023. Highly pathogenic avian influenza A(H5N1) virus infection in farmed minks, Spain, October 2022. Eurosurveillance 28:2300001.
6. Bessière P, Gaide N, Croville G, Crispo M, Fusade-Boyer M, Abou Monsef Y, Dirat M, Beltrame M, Dendauw P, Lemberger K, Guérin JL, Le Loc'h G. 2024. High pathogenicity avian influenza A (H5N1) clade 2.3.4.4b virus infection in a captive Tibetan black bear (Ursus thibetanus): investigations based on paraffin-embedded tissues, France, 2022. Microbiol Spectr 12:e0373623.
7. Chestakova IV, van der Linden A, Bellido Martin B, Caliendo V, Vuong O, Thewessen S, Hartung T, Bestebroer T, Dekker J, Jonge Poerink B, Gröne A, Koopmans M, Fouchier R, van den Brand JMA, Sikkema RS. 2023. High number of HPAI H5 virus infections and antibodies in wild carnivores in the Netherlands, 2020-2022. Emerg Microbes Infect 12:2270068.
8. Cruz CD, Icochea ME, Espejo V, Troncos G, Castro-Sanguinetti GR, Schilling MA, Tinoco Y. 2023. Highly Pathogenic Avian Influenza A(H5N1) from Wild Birds, Poultry, and Mammals, Peru. Emerg Infect Dis 29:2572-2576.
9. Elsmo EJ, Wünschmann A, Beckmen KB, Broughton-Neiswanger LE, Buckles EL, Ellis J, Fitzgerald SD, Gerlach R, Hawkins S, Ip HS, Lankton JS, Lemley EM, Lenoch JB, Killian ML, Lantz K, Long L, Maes R, Mainenti M, Melotti J, Moriarty ME, Nakagun S, Ruden RM, Shearn-Bochsler V, Thompson D, Torchetti

MK, Van Wettere AJ, Wise AG, Lim AL. 2023. Highly Pathogenic Avian Influenza A(H5N1) Virus Clade
2.3.4.4b Infections in Wild Terrestrial Mammals, United States, 2022. Emerg Infect Dis 29:2451-2460.
10.   Herfst S, Zhang J, Richard M, McBride R, Lexmond P, Bestebroer TM, Spronken MIJ, de Meulder D,
van den Brand JM, Rosu ME, Martin SR, Gamblin SJ, Xiong X, Peng W, Bodewes R, van der Vries E,
Osterhaus A, Paulson JC, Skehel JJ, Fouchier RAM. 2020. Hemagglutinin Traits Determine
Transmission of Avian A/H10N7 Influenza Virus between Mammals. Cell Host Microbe 28:602-613.e7.
11.   Fusaro A, Gonzales JL, Kuiken T, Mirinavičiūtė G, Niqueux É, Ståhl K, Staubach C, Svartström O,
Terregino C, Willgert K, Baldinelli F, Delacourt R, Georganas A, Kohnle L. 2024. Avian influenza
overview December 2023-March 2024. Efsa j 22:e8754.
12.   Nemanichvili N, Berends AJ, Wubbolts RW, Gröne A, Rijks JM, de Vries RP, Verheije MH. 2021. Tissue
Microarrays to Visualize Influenza D Attachment to Host Receptors in the Respiratory Tract of Farm
Animals. Viruses 13.
13.   Song H, Qi J, Khedri Z, Diaz S, Yu H, Chen X, Varki A, Shi Y, Gao GF. 2016. An Open Receptor-
Binding Cavity of Hemagglutinin-Esterase-Fusion Glycoprotein from Newly-Identified Influenza D Virus:
Basis for Its Broad Cell Tropism. PLOS Pathogens 12:e1005411.
14.   Burrough E, Magstadt D, Petersen B, Timmermans S, Gauger P, Zhang J, Siepker C, Mainenti M, Li G,
Thompson A, Gorden P, Plummer P, Main R. 2024. Highly Pathogenic Avian Influenza A(H5N1) Clade
2.3.4.4b Virus Infection in Domestic Dairy Cattle and Cats, United States, 2024. Emerging Infectious
Disease journal 30.
15.   Uyeki TM, Milton S, Hamid CA, Webb CR, Presley SM, Shetty V, Rollo SN, Martinez DL, Rai S,
Gonzales ER, Kniss KL, Jang Y, Frederick JC, Cruz JADL, Liddell J, Di H, Kirby MK, Barnes JR, Davis
CT. Highly Pathogenic Avian Influenza A(H5N1) Virus Infection in a Dairy Farm Worker. New England
Journal of Medicine 0.
16.   Kristensen C, Jensen HE, Trebbien R, Webby RJ, Larsen LE. 2024. The avian and human influenza A
virus receptors sialic acid (SA)-α2,3 and SA-α2,6 are widely expressed in the bovine mammary gland.
bioRxiv doi:10.1101/2024.05.03.592326:2024.05.03.592326.
17.   Song X, Ju H, Lasanajak Y, Kudelka MR, Smith DF, Cummings RD. 2016. Oxidative release of natural
glycans for functional glycomics. Nat Methods 13:528-34.
18.   Vos GM, Weber J, Sweet IR, Hooijschuur KC, Sastre Toraño J, Boons GJ. 2023. Oxidative Release of
O-Glycans under Neutral Conditions for Analysis of Glycoconjugates Having Base-Sensitive
Substituents. Anal Chem 95:8825-8833.
19.   Safferthal M, Bechtella L, Zappe A, Vos GM, Pagel K. 2024. Labeling of Mucin-Type O-Glycans for
Quantification Using Liquid Chromatography and Fluorescence Detection. ACS Measurement Science
Au doi:10.1021/acsmeasuresciau.3c00071.
20.   Zhang T, Madunić K, Holst S, Zhang J, Jin C, ten Dijke P, Karlsson NG, Stavenhagen K, Wuhrer M.
2020. Development of a 96-well plate sample preparation method for integrated N- and O-glycomics
using porous graphitized carbon liquid chromatography-mass spectrometry. Molecular Omics 16:355-
363.
21.   Irie A, Koyamat S, Kozutsumi Y, Kawasaki T, Suzuki A. 1998. The molecular basis for the absence of N-
glycolylneuraminic acid in humans. Journal of Biological Chemistry 273:15866-15871.
22.   Peri S, Kulkarni A, Feyertag F, Berninsone PM, Alvarez-Ponce D. 2018. Phylogenetic distribution of
CMP-Neu5Ac hydroxylase (CMAH), the enzyme synthetizing the proinflammatory human xenoantigen
Neu5Gc. Genome Biology and Evolution 10:207-219.
23.   Jahan M, Thomson PC, Wynn PC, Wang B. 2021. The non-human glycan, N-glycolylneuraminic acid
(Neu5Gc), is not expressed in all organs and skeletal muscles of nine animal species. Food Chem
343:128439.
24.   Broszeit F, Tzarum N, Zhu X, Nemanichvili N, Eggink D, Leenders T, Li Z, Liu L, Wolfert MA,
Papanikolaou A, Martínez-Romero C, Gagarinov IA, Yu W, García-Sastre A, Wennekes T, Okamatsu M,
Verheije MH, Wilson IA, Boons GJ, de Vries RP. 2019. N-Glycolylneuraminic Acid as a Receptor for
Influenza A Viruses. Cell Rep 27:3284-3294.e6.
25.   Vlasak R, Luytjes W, Spaan W, Palese P. 1988. Human and bovine coronaviruses recognize sialic acid-
containing receptors similar to those of influenza C viruses. Proc Natl Acad Sci U S A 85:4526-9.
26.   Li Z, Liu L, Unione L, Lang Y, de Groot RJ, Boons GJ. 2022. Synthetic O-Acetyl-N-glycolylneuraminic
Acid Oligosaccharides Reveal Host-Associated Binding Patterns of Coronaviral Glycoproteins. ACS
Infect Dis 8:1041-1050.
27.   Tomris I, Unione L, Nguyen L, Zaree P, Bouwman KM, Liu L, Li Z, Fok JA, Ríos Carrasco M, van der
Woude R, Kimpel ALM, Linthorst MW, Kilavuzoglu SE, Verpalen E, Caniels TG, Sanders RW, Heesters
BA, Pieters RJ, Jiménez-Barbero J, Klassen JS, Boons GJ, de Vries RP. 2023. SARS-CoV-2 Spike N-
Terminal Domain Engages 9-O-Acetylated α2-8-Linked Sialic Acids. ACS Chem Biol 18:1180-1191.
28.   Barnard KN, Alford-Lawrence BK, Buchholz DW, Wasik BR, LaClair JR, Yu H, Honce R, Ruhl S, Pajic
P, Daugherity EK, Chen X, Schultz-Cherry SL, Aguilar HC, Varki A, Parrish CR. 2020. Modified Sialic
Acids on Mucus and Erythrocytes Inhibit Influenza A Virus Hemagglutinin and Neuraminidase Functions.
J Virol 94.
29.   Rogers GN, Herrler G, Paulson JC, Klenk HD. 1986. Influenza C virus uses 9-O-acetyl-N-
acetylneuraminic acid as a high affinity receptor determinant for attachment to cells. J Biol Chem
261:5947-51.

30. Vos GM, Hooijschuur KC, Li Z, Fjeldsted J, Klein C, de Vries RP, Toraño JS, Boons G-J. 2023. Sialic
acid O-acetylation patterns and glycosidic linkage type determination by ion mobility-mass spectrometry.
Nature Communications 14:6795.
31. Nemanichvili N, Spruit CM, Berends AJ, Gröne A, Rijks JM, Verheije MH, de Vries RP. 2022. Wild and
domestic animals variably display Neu5Ac and Neu5Gc sialic acids. Glycobiology 32:791-802.
32. Hiono T, Okamatsu M, Igarashi M, McBride R, de Vries RP, Peng W, Paulson JC, Sakoda Y, Kida H.
2016. Amino acid residues at positions 222 and 227 of the hemagglutinin together with the
neuraminidase determine binding of H5 avian influenza viruses to sialyl Lewis X. Arch Virol 161:307-16.
33. Guo H, de Vries E, McBride R, Dekkers J, Peng W, Bouwman KM, Nycholat C, Verheije MH, Paulson
JC, van Kuppeveld FJ, de Haan CA. 2017. Highly Pathogenic Influenza A(H5Nx) Viruses with Altered
H5 Receptor-Binding Specificity. Emerg Infect Dis 23:220-231.
34. Wang M, Tscherne DM, McCullough C, Caffrey M, García-Sastre A, Rong L. 2012. Residue Y161 of
influenza virus hemagglutinin is involved in viral recognition of sialylated complexes from different hosts.
Journal of Virology 86:4455-4462.
35. Nemanichvili N, Tomris I, Turner HL, McBride R, Grant OC, van der Woude R, Aldosari MH, Pieters RJ,
Woods RJ, Paulson JC, Boons GJ, Ward AB, Verheije MH, de Vries RP. 2019. Fluorescent Trimeric
Hemagglutinins Reveal Multivalent Receptor Binding Properties. J Mol Biol 431:842-856.
36. Nicholls JM, Chan RW, Russell RJ, Air GM, Peiris JS. 2008. Evolving complexities of influenza virus and
its receptors. Trends Microbiol 16:149-57.
37. Zhang H. 2009. Tissue and host tropism of influenza viruses: importance of quantitative analysis. Sci
China C Life Sci 52:1101-10.
38. Chutinimitkul S, Herfst S, Steel J, Lowen AC, Ye J, van Riel D, Schrauwen EJ, Bestebroer TM, Koel B,
Burke DF, Sutherland-Cash KH, Whittleston CS, Russell CA, Wales DJ, Smith DJ, Jonges M, Meijer A,
Koopmans M, Rimmelzwaan GF, Kuiken T, Osterhaus AD, García-Sastre A, Perez DR, Fouchier RA.
2010. Virulence-associated substitution D222G in the hemagglutinin of 2009 pandemic influenza
A(H1N1) virus affects receptor binding. J Virol 84:11802-13.
39. CDC. CDC Reports Second Human Case of H5 Bird Flu Tied to Dairy Cow Outbreak.
doi:https://www.cdc.gov/media/releases/2024/s0522-human-case-h5.html.
40. Fouchier RA, Schneeberger PM, Rozendaal FW, Broekman JM, Kemink SA, Munster V, Kuiken T,
Rimmelzwaan GF, Schutten M, Van Doornum GJ, Koch G, Bosman A, Koopmans M, Osterhaus AD.
2004. Avian influenza A virus (H7N7) associated with human conjunctivitis and a fatal case of acute
respiratory distress syndrome. Proc Natl Acad Sci U S A 101:1356-61.
41. Belser JA, Gustin KM, Maines TR, Pantin-Jackwood MJ, Katz JM, Tumpey TM. 2012. Influenza virus
respiratory infection and transmission following ocular inoculation in ferrets. PLoS Pathog 8:e1002569.
42. Anonymous. doi:https://www.canr.msu.edu/news/hpai-dairy-herd-infection-case-report.
43. Graaf A, Piesche R, Sehl-Ewert J, Grund C, Pohlmann A, Beer M, Harder T. 2023. Low Susceptibility of
Pigs against Experimental Infection with HPAI Virus H5N1 Clade 2.3.4.4b. Emerg Infect Dis 29:1492-
1495.
44. Kwon T, Trujillo JD, Carossino M, Lyoo EL, McDowell CD, Cool K, Matias-Ferreyra FS, Jeevan T,
Morozov I, Gaudreault NN, Balasuriya UBR, Webby RJ, Osterrieder N, Richt JA. 2024. Pigs are highly
susceptible to but do not transmit mink-derived highly pathogenic avian influenza virus H5N1 clade
2.3.4.4b. Emerg Microbes Infect doi:10.1080/22221751.2024.2353292:2353292.
45. Shaner NC, Lin MZ, McKeown MR, Steinbach PA, Hazelwood KL, Davidson MW, Tsien RY. 2008.
Improving the photostability of bright monomeric orange and red fluorescent proteins. Nat Methods
5:545-51.
46. Bouwman KM, Parsons LM, Berends AJ, de Vries RP, Cipollo JF, Verheije MH. 2020. Three amino acid
changes in avian coronavirus spike protein allow binding to kidney tissue. J Virol 94.
47. Wickramasinghe IN, de Vries RP, Grone A, de Haan CA, Verheije MH. 2011. Binding of avian
coronavirus spike proteins to host factors reflects virus tropism and pathogenicity. J Virol 85:8903-12.

Figure S1. Overview of lectin specificity to differentially linked un- and modified SIA. Black boxes indicate
core specificities, gray boxes indicate possible ligands.

Figure S2. Hemagglutination assay with chicken erythrocytes with the lectins used in this study.

379