The Impact of Glutathione Administration on Body Weight and Lipid Metabolism in Mice Following Exposure to Mobile Phone Radiation of Frequency 850-1900

Article · May 2024

8 authors, including:

Abdulhakeem Binhambali
Tuskegee University
18 PUBLICATIONS 5 CITATIONS

SEE PROFILE

Rabiu AbduSSALAM Magaji
Ahmadu Bello University
85 PUBLICATIONS 588 CITATIONS

SEE PROFILE

Amat Abdoulie Tekanyi
Ahmadu Bello University
4 PUBLICATIONS 0 CITATIONS

SEE PROFILE

Tolulope Ipinlaiye
2 PUBLICATIONS 0 CITATIONS

SEE PROFILE

PLOS ONE

The Impact of Glutathione Administration on Body Weight and Lipid Metabolism in Mice Following Exposure to Mobile Phone Radiation of Frequency 850-1900
--Manuscript Draft--

Manuscript Number:	
Article Type:	Research Article
Full Title:	The Impact of Glutathione Administration on Body Weight and Lipid Metabolism in Mice Following Exposure to Mobile Phone Radiation of Frequency 850-1900
Short Title:	None
Corresponding Author:	Abdulhakeem Binhambali, DVM North Carolina State University DOCS: NC State University Department of Clinical Sciences Raleigh, NC State UNITED STATES
Keywords:	Mobile phone radiation, Glutathione, Serum lipid profiles, High density lipoprotein, Radiation effec
Abstract:	In an era where electronic advancements surge forward at an unprecedented pace, public apprehension surrounding the health ramifications of radio-frequency (RF) radiation intensifies. This study looked into the impact of chronic exposure to mobile phone radiation, coupled with glutathione supplementation, on serum lipid profiles and body weight in mice—a subject vital for understanding potential health risks in humans and animals respectively. Thirty-five male mice were divided into seven groups, each subjected to various mobile phone modes and glutathione treatments. Over five weeks, these mice were exposed to 300 missed calls daily, simulating real-world exposure scenarios. Different serum lipid parameters—triglycerides, cholesterol, high-density lipoprotein (HDL), and low-density lipoprotein (LDL)—were monitored and observed throughout the study period. Remarkably, triglycerides, cholesterol, and LDL levels exhibited no noteworthy statistical variances compared to the control group or among experimental cohorts exposed to diverse range of mobile phone radiation modes or glutathione supplements. However, we noted some little deviations in HDL levels, particularly in the silent and silent+glutathione groups. Moreover, we also noted an unexpected result in weight loss across all groups by week three, which was pronounced by week five, and notably pronounced in the glutathione-administered cohorts. This outcome hints at glutathione's complex role in mitigating some complications of mobile phone radiation exposure. In conclusion, the study suggests a potential interplay between glutathione supplementation, mobile phone radiation exposure, and HDL levels in mice—an avenue ripe for further exploration into the understanding of cellular response to modern technological exposures.
Order of Authors:	Aliyah Temitayo Ahmed, Bsc.
	Abdulhakeem Binhambali, DVM
	Abdullahi Hussein Umar, PhD
	Amat Abdoulie Tekanyi, BSc.
	Aisha Pate Ahmad, Bsc.
	Amina Nuhu, Bsc.
	Tolulope Ipinlaiye, Bsc.
	Rabiu AbdusSALAM Magaji, PhD
Opposed Reviewers:	
Additional Information:	
Question	Response

Financial Disclosure Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the submission guidelines for detailed requirements. View published research articles from *PLOS ONE* for specific examples. This statement is required for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate. **Funded studies** Enter a statement with the following details: • Initials of the authors who received each award • Grant numbers awarded to each author • The full name of each funder • URL of each funder website • Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript? Did you receive funding for this work?	The author(s) received no specific funding for this work.
Competing Interests Use the instructions below to enter a competing interest statement for this submission. On behalf of all authors, disclose any competing interests that could be perceived to bias this work—acknowledging all financial support and any other relevant financial or non-financial competing interests. This statement is required for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate and that any funding sources listed in your Funding Information later in the submission form are also declared in your Financial Disclosure statement.	The authors have declared that no competing interests exist

View published research articles from
PLOS ONE for specific examples.

NO authors have competing interests

Enter: *The authors have declared that no competing interests exist*.

Authors with competing interests

Enter competing interest details beginning with this statement:

I have read the journal's policy and the authors of this manuscript have the following competing interests: [insert competing interests here]

* typeset

Ethics Statement

Enter an ethics statement for this submission. This statement is required if the study involved:

• Human participants
• Human specimens or tissue
• Vertebrate animals or cephalopods
• Vertebrate embryos or tissues
• Field research

Write "N/A" if the submission does not require an ethics statement.

General guidance is provided below. Consult the submission guidelines for detailed instructions. **Make sure that all information entered here is included in the Methods section of the manuscript.**

AHMADU BELLO UNIVESRITY COMMITTEE ON ANIMAL USE AND CARE

Approval No: ABUCAUC/2021/005

Format for specific study types

Human Subject Research (involving human participants and/or tissue)
- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

Animal Research (involving vertebrate animals, embryos or tissues)
- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering
- If anesthesia, euthanasia, or any kind of animal sacrifice is part of the study, include briefly which substances and/or methods were applied

Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:
- Field permit number
- Name of the institution or relevant body that granted permission

Data Availability

Authors are required to make all data underlying the findings described fully available, without restriction, and from the time of publication. PLOS allows rare exceptions to address legal and ethical concerns. See the PLOS Data Policy and FAQ for detailed information.

Yes - all data are fully available without restriction

A Data Availability Statement describing where the data can be found is required at submission. Your answers to this question constitute the Data Availability Statement and **will be published in the article**, if accepted.

Important: Stating 'data available on request from the author' is not sufficient. If your data are only available upon request, select 'No' for the first question and explain your exceptional situation in the text box.

Do the authors confirm that all data underlying the findings described in their manuscript are fully available without restriction?

Describe where the data may be found in full sentences. If you are copying our sample text, replace any instances of XXX with the appropriate details.

- If the data are **held or will be held in a public repository**, include URLs, accession numbers or DOIs. If this information will only be available after acceptance, indicate this by ticking the box below. For example: *All XXX files are available from the XXX database (accession number(s) XXX, XXX.).*
- If the data are all contained **within the manuscript and/or Supporting Information files**, enter the following: *All relevant data are within the manuscript and its Supporting Information files.*
- If neither of these applies but you are able to provide **details of access elsewhere**, with or without limitations, please do so. For example:

  Data cannot be shared publicly because of [XXX]. Data are available from the XXX Institutional Data Access / Ethics Committee (contact via XXX) for researchers who meet the criteria for access to confidential data.

  The data underlying the results presented in the study are available from (include the name of the third party

N/A

and contact information or URL). • This text is appropriate if the data are owned by a third party and authors do not have permission to share the data. * typeset	
Additional data availability information:	Tick here if the URLs/accession numbers/DOIs will be available only after acceptance of the manuscript for publication so that we can ensure their inclusion before publication.

Abdulhakeem Binhambali, DVM

Department of Clinical Sciences,

College of Veterinary Medicine,

North Carolina State University,

1060, William Moore drive, 27606, Raleigh, USA.

02/25/2024

Dear Editor,

We are writing to submit our original research article, titled "**The Impact of Glutathione Administration on Body Weight and Lipid Metabolism in Mice Following Exposure to Mobile Phone Radiation of Frequency 850-1900**," for consideration by PLOS ONE.

We humbly confirm that this work is original and has not been published elsewhere, nor is it currently under consideration for publication elsewhere and was approved by Ahmadu Bello University Animal Use and Care Committee prior to the commencement of the work with the approval number of ABUCAUC/2021/005.

In this paper, we present findings on the effect of glutathione on the mice body weight and serum lipid after exposure to GSM radiation of frequency 850-1900. Our research outcomes offer compelling insights into the interplay between animal health and human well-being, underscoring the interconnectedness of medical research across species boundaries. Importantly, our findings diverge from previous studies, highlighting the dynamic nature of animal research and the multifaceted factors influencing diseases initiation and outcomes in both humans and animals.

We believe that our research represents a foundational step towards a deeper understanding of radiaton exposure at the cellular and molecular level, by giving us the privilege to understand the diagnostics and disease dynamics, particularly within pathophysiological contest. We are enthusiastic about the potential avenues for future research and remain dedicated to advancing scientific knowledge in this domain.

Thank you for considering our research findings, and we eagerly anticipate the opportunity for continued collaboration and exchange of ideas in the field of medicine in general.

We have no conflicts of interest to disclose.

Please direct all correspondence regarding this manuscript to me at abinham@ncsu.edu. Thank you for your consideration.

Sincerely,

Abdulhakeem Binhambali, DVM

The Impact of Glutathione Administration on Body Weight and Lipid Metabolism in Mice

Following Exposure to Mobile Phone Radiation of Frequency 850-1900

AUTHORS:

Aliyah Temitayo Ahmed[1], Abdulhakeem Binhambali [2,3] * Abdullahi Hussein Umar [1], Amat

Abdoulie Tekanyi[1], Aisha Ahmad Pate[1], Amina Nuhu[1],Tolulope Ipinlaiye[1], Rabiu

AbduSSALAM Magaji[1]

[1]Department of Human Physiology, Faculty of Basic Medical Sciences, College of Medical

Sciences, Ahmadu Bello University, Zaria, Nigeria

[2]Department of Clinical Science, College of Veterinary Medicine, North Carolina State

University, Raleigh, USA

[3]Translation Research in Pain, College of Veterinary Medicine, NCSU, Raleigh, USA

Corresponding author: Abdulhakeem Binhambali, DVM,

abinham@ncsu.edu/email2hambali@gmail.com, Department of Clinical Sciences, College of

Veterinary Medicine, NCSU, Raleigh, USA.

ABSTRACT

In an era where electronic advancements surge forward at an unprecedented pace, public

apprehension surrounding the health ramifications of radio-frequency (RF) radiation intensifies.

This study looked into the impact of chronic exposure to mobile phone radiation, coupled with

glutathione supplementation, on serum lipid profiles and body weight in mice—a subject vital

for understanding potential health risks in humans and animals respectively.

Thirty-five male mice were divided into seven groups, each subjected to various mobile phone

modes and glutathione treatments. Over five weeks, these mice were exposed to 300 missed calls

daily, simulating real-world exposure scenarios. Different serum lipid parameters—triglycerides,

cholesterol, high-density lipoprotein (HDL), and low-density lipoprotein (LDL)—were

monitored and observed throughout the study period.

Remarkably, triglycerides, cholesterol, and LDL levels exhibited no noteworthy statistical

variances compared to the control group or among experimental cohorts exposed to diverse

range of mobile phone radiation modes or glutathione supplements. However, we noted some

little deviations in HDL levels, particularly in the silent and silent+glutathione groups. Moreover,

we also noted an unexpected result in weight loss across all groups by week three, which was

pronounced by week five, and notably pronounced in the glutathione-administered cohorts. This

outcome hints at glutathione's complex role in mitigating some complications of mobile phone

radiation exposure.

In conclusion, the study suggests a potential interplay between glutathione supplementation,

mobile phone radiation exposure, and HDL levels in mice—an avenue ripe for further

exploration into the understanding of cellular response to modern technological exposures.

Keywords: Mobile phone radiation, Glutathione, Serum lipid profiles, High density lipoprotein, Radiation effect

INTRODUCTION

The ubiquitous presence of mobile phones, embraced by approximately 91% of the younger population globally, has revolutionized our lifestyles [1]. These devices have seamlessly integrated into daily routines, becoming indispensable tools for communication and information access. However, as mobile phone usage continues to soar, concerns about their potential impact on human health have intensified [2, 3]. With mobile phones pervading every corner of the globe, exposure to electromagnetic field radiation (EMR) has become a constant presence in modern society [4].

Radiations consist of both electric and magnetic fields, originating from natural and man-made sources. EMR is present in various scenarios of everyone's life, with some of the most common sources being solar radiation, household electric currents (e.g., mobile phones, television sets, Wi-Fi, microwave ovens, computers), and telecommunication antennas [5-11].

Electromagnetic frequency (EMF) has numerous biochemical consequences, including the degradation of large cell molecules and imbalances in physiological ionic equilibrium. EMF exposure is associated with increased formation of reactive oxygen species (ROS) [12], which can cause injury to cellular constituents such as lipids, proteins, and DNA. It has been shown that electromagnetic radiation influences living organisms by producing or increasing ROS, leading to various physiological and biochemical effects, including DNA damage [13]. Analog telephones typically use frequencies between 800 and 900 MHz, while digital telephones use frequencies between 1850 and 1990 MHz, and microwave ovens operate at a frequency of 2450 MHz. The 900 MHz EMF emitted by mobile phones is commonly used in low-income countries.

However, cell phone radiation, a form of electromagnetic radiation known as microwaves,

comprises electric and magnetic waves that travel through space at the speed of light [14].

Concerns have been raised by researchers regarding the release of EMF from mobile phones into

the atmosphere, suggesting potential harmful effects on human health [15, 16]. Indeed, the

adverse impacts of EMF have become a pressing global issue, with reports of detrimental effects

on various organs such as the brain, ear, and reproductive organs [17, 18, 5, 19, 20].In today's

digital era, the proliferation of cell phone usage is on the rise, contributing to a growing addiction

among individuals [21]. Over the past two decades, mobile phone subscriptions have surged

from 12.4 million to over 5.6 billion globally, reaching approximately 70% of the world's

population [74]. However, alongside this widespread adoption, concerns regarding its impact on

public health have emerged. There are numerous reports of both mental and physical health

hazards associated with cell phone use across all age groups, with some effects proving to be

severe, including the heightened risk of cancers [22].

Electromagnetic Radiation (EMR) is a dynamic force of energy propagation, characterized by

photons exhibiting both particle and wave properties, traveling at the speed of light. These waves

bear energy and transmit it upon interaction with matter. The energy content of EMR is directly

linked to its frequency and inversely related to its wavelength, meaning that waves with shorter

wavelengths possess higher energy levels. EMR encompasses a spectrum ranging from radio

waves to microwaves, infrared, visible light, UV, and radiographs, each with varying energy

levels [23]. Both EMR and particulate radiation have the ability to generate ion pairs through

their interaction with matter. Ionizing radiation, including alpha particles, neutrons,

beta rays, charged nuclei, and positron radiation, aids in producing high-quality images for

medical diagnostics and is utilized in cancer treatment as well [24].

The impact of EMF extends to biochemical processes, including the degradation of large cellular

molecules and disruptions in ionic equilibrium. Exposure to EMF has been linked to heightened

formation of reactive oxygen species (ROS) [12] as stated earlier, ROS generation can also occur

aside from the previously mentioned EMF, including, radioactivity, stress, cigarette smoke, and

redox processes within the body [5]. It is also important to know that the adverse effects of

mobile phone radiation exposure on vital organs like the kidney and testis remain poorly

understood, as does its impact on serum lipids, which play crucial roles in diseases such as

cardiovascular disorders [13]. While interest in evaluating the health effects of non-ionizing

radiation, particularly radiofrequency (RF) electromagnetic fields, has grown significantly in

recent decades, fueled by the widespread use of mobile telecommunications worldwide, the exact

mechanisms and consequences of such exposure are still being unraveled. Mobile networks

operate on varying frequencies across different regions, with Global System for Mobile

Communications (GSM) networks typically utilizing the 900 and 1800 MHz bands [30].

Among the potential effects of radiofrequency radiation on the human body, research has

predominantly focused on its impact on the brain, cancer incidence, and fertility. This includes

investigations into enzyme induction, neurological symptoms, toxicological effects, genotoxicity,

carcinogenicity, and alterations in sperm cell fertility. The literature extensively examines the

influence of radiofrequency radiation on various cellular processes, including mitochondrial

function, apoptosis pathways, heat shock proteins, free radical metabolism, cell proliferation,

differentiation, DNA damage, and plasma membrane integrity [31-36]. Despite these efforts,

gaps remain in our understanding of the full spectrum of health effects associated with

radiofrequency radiation exposure, necessitating further research to elucidate its potential risks

and inform regulatory measures to protect public health.On the other hand, non-ionizing

radiation, though lacking in ionizing power, can still pose risks with prolonged exposure, a

crucial factor when considering its interaction with the human body [26]. The consensus among

many researchers is that non-ionizing radiation carries significant hazards for public health

[27].Termed electromagnetic pollution, the potential health impacts of EMF exposure are

increasingly recognized [28]. Non-ionizing radiation spans from extremely low frequency (ELF)

to visible light, while ionizing radiation ranges from higher frequencies beyond visible

light[29].The proliferation of electric EMF from modern technologies, including everyday

appliances and mobile devices, poses a tangible risk to human well-being as explained earlier.

Glutathione, on the other hand, is a vital tripeptide composed of cysteine, glycine, and glutamic

acid, serving as a cornerstone in numerous bodily functions, from combating oxidative stress to

bolstering immune responses [37]. Its significance extends to mitigating toxin accumulation by

aiding in the detoxification process, particularly in the liver's conversion and elimination of

harmful compounds like mercury and persistent organic pollutants (POPs). Functioning as a

potent antioxidant, glutathione is ubiquitous across various organisms and cell types, primarily

existing in either its reduced (GSH) or oxidized (GSSG) form [38, 39]. Under normal

circumstances, the majority of glutathione remains in its reduced state, pivotal for maintaining

cellular health and redox balance. Studies by Markov *et al.* [40], have explored the impact of

radio frequency electromagnetic waves (RF-EMW) on human health, noting a potential

correlation between long-term exposure and reduced body weight. While RF radiation, falling

within the spectrum of 0.5 MHz to 100 GHz, may affect certain organs and systems, the precise

health risks from various sources of non-ionizing radiation, including those emitted by mobile

phones, remain less understood [41].Research by NRC and Roberts *et al.* [42] has indicated that

exposure to RF/MW radiation can influence several hematological variables, both in animal

models and humans. This underscores the need for further investigation into the potential health

implications of non-ionizing radiation exposure, particularly in light of its pervasiveness in

modern society.

In animal research, the welfare of experimental subjects is of paramount importance. Criteria

such as a 20 percent reduction in body weight are often designated as indicators of severe

suffering, guiding humane endpoint decisions [43]. Previous studies have underscored the

potential impact of low electric fields on mice, revealing diminished body weight and reduced

water intake following one month of exposure to harsh conditions [44].

Mobile phone users in various countries and continents are exposed to different frequencies.

EMF exposure depends on the frequency of the cell phone [45]. Furthermore, there are

indications that mobile phone radiation may also influence body weight in experimental animals.

In our contemporary era, where mobile phone usage is common and its potential health

implications are a subject of concern, we endeavor to explore the effect of such radiation on

serum lipid and the interplay between glutathione and body mass in mice subjected to chronic

mobile phone radiation exposure. Through this investigation, we aim to shed light on the

complex relationship between glutathione, mobile phone radiation, and serum lipid in

experimental models. We evaluated the effects of EMF on the serum lipid profiles and the body

weight and the potentially harmful effects of EMF on these parameters of mice model.

MATERIALS AND METHOD

METHODOLOGY

Animals and grouping

The animals involved in the study were maintained and used in accordance with the Animal

Welfare Act and the Guide for the Care and Use of Laboratory animals prepared by the Ahmadu

Bello University Committee on Animal Use and Care (ABUCAUC). And ethical approval was

obtained from the ABUCAUC.

At the start of the experiment, thirty five (35) apparently healthy adult male Mice between the

ages of 8-12 weeks were obtained from the Department of Human Physiology, College of Basic

Medical Sciences, Ahmadu Bello University, Zaria, Nigeria animal breeding section. They were

housed in standard polypropylene cages and were maintained and allowed free access to feed and

water 24/7 and 12h alternate day light and darkness. The animals were allowed to acclimatize to

the environment of the behavioral laboratory for two weeks before the commencement of the

experiment and they were provided with feeds and water adlibitum.

The animals were randomly selected and divided into seven (7) groups of five (n=5) mice per

group as follows: Group I: Normal control group, Group II: Group exposed to mobile phone

radiation in ringtone mode, Group III: Group exposed to mobile phone radiation in vibration

mode , Group IV: Group exposed to mobile phone radiation in silent mode, Group V: Group

exposed to mobile phone radiation in ringtone mode plus glutathione , Group VI: Group exposed

to mobile phone radiation in vibration mode plus glutathione and the last group, Group VII were

exposed to mobile phone radiation in silent mode plus glutathione.

The animals in groups II to VII were all exposed to 4hour mobile phone radiation of 900 hertz

and an average of 300 missed calls per day for 5 weeks. The group I as the normal control is not

exposed to any form of radiation or glutathione.

Radiation exposure

Using Itel mobile phone (it2160) with specific absorption rate (SAR) of 2W/kg and 2G Network

Bands, of frequency 850-1900 MHz, the animals in groups II to VII were all exposed to 4 hours

mobile phone radiation of an average of 300 missed calls per day for six (6) weeks. Animals in

groups V, VI and VII were administered glutathione (250 mg/kg) thirty minutes prior to mobile

phone radiation daily.

Determination of body weight

The body weight is taken early in the morning (between 7am-9am) prior to their feeding.

Animals were individually weighed on a weekly basis (Wednesdays) in order to detect any

changes in their body weights. Before administration of glutathione and exposure to radiation,

the animals were weighed with a sensitive weighing balance (Uline Balance Scale – H-9886,

7.5X5.7 square, 2,200 g x 0.1 g)

Glutathione at a dose of 250 mg/Kg, was administered orally to each Mice daily to group V-VII

before exposure to radiation for 6 weeks. Animals in group II - IV were exposed to mobile phone

radiation only, for 4 hours daily throughout the six weeks radiation without Glutathione

Determination of lipoproteins

Triglyceride

0.5ml of working reagent was added to a set of clean labeled test tubes and two of the test tubes

were labeled standard 1 and 2 while another was labeled "blank".0.1ml of the sample was added

to the approximately labeled test tube .mixed and incubated for 30 minutes at 37º, the absorbance

of the sample and standard were read against the reagent blank at the wavelength of 546nm.

Total Cholesterol (TC)

0.5ml of working reagent was added to a set of clean labeled test tubes and two of the test tubes

were labeled standard 1 and 2 while another was labeled "blank".0.1ml of homogenate was

added to the approximately labeled test tube .mixed and incubated for 30 minutes at 37º, the

absorbance of the sample and standard were read against the reagent blank at the wavelength of

546nm.

High-density lipoproteins (HDL)

0.1 ml of homogenate sample was mixed with 0.5ml of working reagent (R1), it was mixed and

incubated for exactly 30 minutes for 37º, 0.5ml of the second reagent (R2) was mixed and

allowed to stand for 20 minutes at 20-25ºC, then 5.0ml of sodium hydroxide was added and

mixed, the absorbance of the sample (A sample) against the reagent blank was read after 5

minutes at the wavelength of 546nm 21

Low-density lipoproteins (LDL)

LDL cholesterol was calculated from measured values of total cholesterol, triglycerides and

HDL-cholesterol according to the relationship:

[LDL-chol]= [total chol] – [HDL-chol] – [TG]/5

Where [TG]/5 is an estimate of VLDL-cholesterol and all values are expressed in mg/dl

219
220   **RESULTS**

221   *WEEKLY WEIGHT PROFILES*

In the first week, the impact of mobile phone radiation and glutathione on mean body weight in mice is illustrated in [Figure 1]. The number of open entries recorded were as follows: control (19.2400 ± 1.00080), ringtone (17.8600 ± .56798), vibration (18.8200 ± .55624), silent (16.7000 ± .92304), ringtone + glutathione (20.9200 ± .1.01114), vibration + glutathione (19.3000 ± 1.31567), and silent + glutathione modes (20.9000 ± 1.52381). No statistically significant difference ($p > 0.05$) was observed when comparing the number of entries in the control group to all other groups. The impact of mobile phone radiation and glutathione on body weight in mice, as depicted in [Figure 1], was also examined. The number of open entries recorded were: control (21.0400 ± .66076), ringtone (18.4800 ± .65985), vibration (19.8400 ± .63135), silent (17.3200 ± .49739), ringtone + glutathione (21.3000 ± .86487), vibration + glutathione (20.5600± 1.10526), and silent + glutathione modes (20.5800 ± 1.33768). There was no statistically significant difference ($p > 0.05$) in the number of entries between the control group and all other groups, except for the silent, ringtone+GSH, and silent=GSH. Similarly, the effects of mobile phone radiation and glutathione on body weight in mice are displayed in [Figure 1]. The number of open entries were: control (24.5600 ± .46648), ringtone (22.6200 ± .40423), vibration (22.1800 ± .69957), silent (21.8200 ± .45869), ringtone + glutathione (20.3600 ± .86487), vibration + glutathione (19.4200 ± 1.15039), and silent + glutathione modes (22.9000 ± 1.10770). There was no statistically significant difference ($p > 0.05$) in the number of entries between the control group and all other groups, except for ringtone + GSH, vibration + GSH, and silent+GSH. Furthermore, the impact of mobile phone radiation and glutathione on body weight in mice is showcased in [Figure 1]. The number of open entries recorded were: control (26.1000 ± .90277), ringtone (22.1600 ± .59380), vibration (22.5200 ± .91291), silent (21.7400 ± .58703), ringtone + glutathione (23.3800 ± .75525), vibration + glutathione (22.0340 ± 1.14471), and silent +

glutathione modes (20.5200 ± 1.26270). A statistically significant difference (p<0.05, ANOVA)

was observed when comparing the number of entries in the control group to all other groups,

except for vibration and ringtone+GSH. Lastly, the impact of mobile phone radiation and

glutathione on body weight in mice, illustrated in [Figure 1], indicates the following number of

open entries: control (27.2200 ± .34264), ringtone (22.8600 ± .63372), vibration (25.1200 ±

.58856), silent (23.4000 ± .70071), ringtone + glutathione (23.2400± .83403), vibration +

glutathione (22.3400± 1.01371), and silent + glutathione modes (21.3600 ± 1.31095). A

statistically significant difference (p<0.05, ANOVA) was noted when comparing the number of

entries in the control group to all other groups.

Figure 1: Effects of mobile phone radiation and glutathione on body weight in mice

SERUM LIPID PROFILES

Triglycerides

The effect of mobile phone radiation and glutathione on triglycerides level in mice is shown in

[Figure 2]. The concentration level were; control (109.5 ± 18.9), ringtone (110.4 ± 6.8), vibration

(111.7 ± 7.2), silent (85.5 ± 5.3), ringtone + glutathione (115.6 ± 10.3), vibration + glutathione

(108.5 ± 6.0) and silent +glutathione modes (81.5 ± 6.0). There was no statistical significance (p

> 0.05) difference when the concentration level of control group was compared to all the groups.

Figure 2: Effects of mobile phone radiation and glutathione on triglycerides level in mice.

(Mean ± S.E.M, n = 7). Glut=Glutathione, Sil= Silent, Vib=Vibration, Ring=Ringtone

There were no statistical significance differences between the groups ($P > 0.05$ ANOVA).

Cholesterol

The effects of mobile phone radiation and glutathione on cholesterol level in mice are shown in

[Figure 3]. The concentration level were; control (147.4 ± 14.8), ringtone (122.7 ± 2.5), vibration

(166.0 ± 9.3), silent (140.3 ± 13.0), ringtone + glutathione (115.8 ± 20.5), vibration + glutathione

(134.3 ± 14.4) and silent +glutathione modes (123.7 ± 17.1). There was no statistical significance

($p > 0.05$) difference when the concentration level of control group was compared to all the

groups.

Figure 3: Effects of mobile phone radiation and glutathione on cholesterol level in mice. (Mean

± S.E.M, n = 7). Glut=Glutathione, Sil= Silent, Vib=Vibration, Ring=Ringtone

There were no statistical significance differences between the groups ($P > 0.05$ ANOVA).

High-density lipoproteins

The effects of mobile phone radiation and glutathione on high-density lipoproteins (HDL) level

in Mice are shown in [Figure 4]. The concentration level were; control (111.2 ± 5.5), ringtone

(79.3 ± 3.4), vibration (108.8 ± 6.2), silent (116.5 ± 7.4), ringtone + glutathione (117.0 ± 8.7),

vibration + glutathione (124.8 ± 6.2) and silent +glutathione modes (86.9 ± 4.5). There was

statistical significance (*p* < 0.05) difference when the concentration level of control group was

compared to ringtone and silent+glutathione groups and no significant differences was seen in

the remaining groups.

Figure 4: Effects of mobile phone radiation and glutathione on high-density lipoproteins level in

mice. Bars with different letters are significant (Mean ± S.E.M, n = 7). Glut=Glutathione, Sil=

Silent, Vib=Vibration, Ring=Ringtone

There was statistical significance differences between the groups (*P* <0.05 ANOVA).

Low-density Lipoprotein

The effects of mobile phone radiation and glutathione on low-density lipoproteins (LDL) level in

mice are shown in [Figure 5]. The concentration level were; control (65.2 ± 16.7), ringtone (23.3

± 2.9), vibration (46.8 ± 6.5), silent (28.7 ± 3.0), ringtone + glutathione (23.0 ± 7.4), vibration +

glutathione (35.1 ± 10.3) and silent +glutathione modes (32.7 ± 16.4). There was no statistical

significance ($p > 0.05$) difference when the concentration level of control group was compared to

all the groups.

Figure 5: The effects of mobile phone radiation and glutathione on low-density lipoproteins

level in mice. (Mean ± S.E.M, n = 7). Glut=Glutathione, Sil= Silent, Vib=Vibration,

Ring=Ringtone

There were no statistically significance differences between the groups ($P > 0.05$ ANOVA)

DISCUSSION

Many studies conducted in recent years have reported various biological effects, spanning epidemiological and clinical investigations in humans, as well as experimental research involving rodents, flies, and cell cultures exposed to phone radiation[46]. However, despite this extensive research, little attention has been paid to serum lipid parameters and body weight reduction to radiation exposure. In today's era, mobile phones have become indispensable tools in our daily lives, serving critical roles in business, commerce, education, and emergency communication. Nevertheless, the proliferation of EMR from sources such as cell phone towers, TV, and FM towers has raised concerns about potential health hazards [47]. Studies by Hocking *et al.* in Australia have linked the proximity of individuals to such towers with higher incidences of leukemia, tumors, and cancer, highlighting the need for further investigation into these health risks [7, 48].

Lipid peroxidation, a consequence of oxidative stress induced by radiation, entails the degradation of lipids when reactive oxygen species (ROS) generated by radiation interact with polyunsaturated fatty acids in the phospholipid membrane bilayer [49]. Glutathione, a vital cellular antioxidant, plays a crucial role in maintaining cellular health. Notably, it has been observed to significantly impact high-density lipoprotein levels [49]. Found abundantly in foods like onions, garlic, avocados, and cruciferous vegetables, glutathione and its precursors, such as l-cysteine and lipoic acid, function as potent scavengers of free radicals, thereby protecting cells from oxidative damage. Moreover, glutathione aids in preserving proteolytic enzymes in the cortical lens fibers, with studies dating back to the late 1960s highlighting its diminished levels in mature cataracts, suggesting its involvement in cataract formation [50].

As a critical antioxidant in various organisms, glutathione prevents damage to cellular

components caused by ROS, including free radicals and peroxides. Its sulfur compounds confer a

lipid-lowering effect, with lower levels of glutathione associated with an increased risk of

cardiovascular disease [51]. Consequently, supplementation with glutathione precursors or

engaging in regular exercise to stimulate glutathione production may confer cardiovascular

benefits during different pathophysiological conditions [52].

While numerous studies have demonstrated the antioxidant effects of glutathione in various

metabolic contexts, its role in mitigating oxidative stress induced by electromagnetic field (EMF)

exposure remains unexplored in the literature. In this study, we investigated the effects of 900-

MHz radiofrequency EMF exposure on oxidative stress in adult mice by assessing parameters

such as triglycerides, cholesterol, HDL, LDL and body weight.

Numerous studies employing robust methodologies have underscored the potential of

electromagnetic fields (EMF) to induce adverse health effects. In the present study, we reported

the repercussions of exposure to mobile phone radiation, specifically exploring the impact of

different modes (ringtone, vibration, silent, etc.) on lipid profiles and weight in mice. We

investigated potential interactions between radiation and its effects on various groups of mice,

aiming to discern any discernible patterns or differences. Our findings unveiled new insights into

the relationship between mobile phone radiation and lipid parameters. While triglycerides

(TGCL), cholesterol (CHL), and low-density lipoprotein (LDL) levels did not exhibit statistical

differences compared to the control group, high-density lipoprotein (HDL) levels showed a

significant difference [Figure 3]. Additionally, we scrutinized the effect of radiation on body

weight, observing notable changes across all groups at week 5, with variations also evident at

weeks 3 and 4 [Figures 1]. These observations align with previous studies by Wilson *et al*.[54]

and Sokolovic *et al*.[55], highlighting the short-term effects of electromagnetic fields on body

mass [54, 55, 56].

Findings reported by Borek *et al*. [65], noted that when rats are exposed to 900-MHz EMF, they

exhibited increased protein oxidation and lipid peroxidation in the brain. However,

administration of 500 mg/kg/day dissolved sulfur compound (garlic) mitigated these effects,

bringing the levels back to normal. In our investigation using glutathione, which is also found in

garlic(as reported in their study), we found that glutathione supplementation did not attenuate the

oxidative effects of EMF exposure on body weight at any point in time; rather the group with

glutathione showed more weight loss from week two and more pronounced at week five than the

control group and the groups that were not given glutathione. This was a surprising thing to us,

with the known fact that glutathione reduces oxidative stress, but in our case the result was

contradictory.

Also interesting to know, our study's duration and methodology yielded results consistent with

the notion that electromagnetic field exposure may induce short-term fluctuations in body

weight, as observed in similar experiments with rats [56]. Also noted, our findings was similar

with the report by Shabat *et al*[57]*,* which noted a decrease in initial mean weight in groups

exposed to high electromagnetic radiation [57]. Likewise the findings from the studies by Hassan

et al. [13] and Lee *et al*. [58], which found no significant weight differences in rats exposed to

magnetic fields [13, 58] was contradicting to our findings which noticed a significant weight loss

in some groups from week 3 and more pronounced at week five and even more especially from

the groups that were given glutathione. Others have also reported an increase in body weight

with long-term exposure to electromagnetic radiation [59] and our finding does not agree with

this point. The reduction in body weight post-EMF exposure may be attributed to direct damage

and excessive production of reactive oxygen species (ROS) induced by electromagnetic

radiation, which glutathione may not adequately scavenge effectively during the period of

radiation exposure [60].

Our study also shed light on the significance of HDL-associated proteins, highlighting the

interplay between serum lipid levels and tumor genesis [64]. While our findings regarding serum

HDL-c levels contradicted those of Wang *et al*.[62], who reported no statistical significance,

although it's essential to note that their focus was primarily on breast cancer, which differs from

our study's scope [62].Moreover, our study did not find statistical significance in cholesterol and

triglyceride levels, aligning with findings from other studies suggesting that these parameters

may not be directly associated with disease initiation or progression from radiation exposure [63,

64, 65].

In summary, our findings revealed that exposure to a 900-MHz EMF led to increased serum

HDL in mice and prompted a loss of body weight by week 3. Glutathione administration

mitigated EMF-induced weight loss in mice, particularly evident at week 5. However, we

advocate for further research, particularly focusing on these parameters with further studies on

liver enzymes and serum electrolytes, to deepen our understanding of the complexities

surrounding EMF exposure and its potential health implications.

ACKNOWLEDGEMENT

This case was carried out without direct funding, and the authors would like to thank the Head and the

entire staffs of the Department of Human Physiology, College of Medical Sciences, Ahmadu Bello

University, Zaria, for their support and provision of necessary facilities. The authors also acknowledge

the indispensable contributions of the Technicians at the Department, whose dedication greatly

contributed to the success of the study.

AUTHOR'S CONTRIBUTIONS

All authors contributed greatly to the success of this work

CONFLICT OF INTEREST

No conflict of interest to declare in this manuscript

REFERENCES

1. Ranie L. Cell phone ownership hits 91% of adult. Pew Research Center, Washington; 2013.

2. Kumar P, Kalonia H, Kumar A. Nitric oxide mechanism in the protective effect of antidepressants against 3-nitropropionic acid-induced cognitive deficit, glutathione and mitochondrial alterations in animal model of Huntington's disease. Behav Pharmacol. 2010;21:217–30. https://doi.org/10.1097/FBP.0b013e32833a5bf4.

3. Thomée S. Mobile Phone Use and Mental Health. A Review of the Research That Takes a Psychological Perspective on Exposure. Int J Environ Res Public Health. 2018;15(12):2692. doi: 10.3390/ijerph15122692. PMID: 30501032; PMCID: PMC6314044.

4. Feytchting M, Ahlbom A, Kheifets L. Electromagnetic field and health. Annu Rev Public Health. 2005;26:165-189.

5. Erogul O, Oztas E, Yildirim I, Kir T, Aydur E, Komesli G, et al. Effects of

electromagnetic radiation from a cellular phone on human sperm motility: An in vitro

study. Arch Med Res. 2006;37(7):840-843.

6. Diem E, Schwarz C, Adlkofer F, Jahn O, Rüdiger H. Non-thermal DNA breakage by

mobile-phone radiation (1800MHz) in human fibroblasts and in transformed GFSH-R17

rat granulosa cells in vitro. Mutat Res. 2005;583:178-183.

7. Hocking B, Gordon IR, Grain HL, Hatfield GE. Cancer incidence and mortality and

proximity to TV towers. Med J Aust. 1996;165:601-605.

8. Avendano C, Mata A, Sarmiento CA, Doncel GF. Use of laptop computers connected to

internet through Wi-Fi decreases human sperm motility and increases sperm DNA

fragmentation. Fertil Steril. 2012;97:39-45.

9. Bernardi P, Cavagnaro M, Pisa S, Piuzzi E. SAR distribution and temperature increase in

an anatomical model of the human eye exposed to the field radiated by the user antenna

in a wireless LAN. IEEE Trans Microw Theory Tech. 1998;46:2074-2082.

10. Azizi MS, Mohamad NR, Salleh A, Othman A, Aris A, Hassan N. Ultra-Wideband

(UWB) antenna radiation on action potentials in human nerve fibers. J Telecom Elec

Comp Eng. 2017;9:59-64.

11. Dasdag S, Yavuz I, Akdag MZ. Effects of radiofrequencies emitted from mobile phones

on teeth and oral tissues. J Int Dent Med Res. 2017;10:176-189.

12. Kivrak E, Yurt K, Kaplan A, Alkan I, Altun G. Effects of electromagnetic fields exposure

on the antioxidant defense system. J Microsc Ultrastruct. 2017;5:167–176. doi:

10.1016/j.jmau.2017.07.003.

13. Hasan I, Rubayet Jahan M, Nabiul Islam M, Rafiqul Islam M. Effect of 2400 MHz

mobile phone radiation exposure on the behavior and hippocampus morphology in Swiss

mouse model. Saudi J Biol Sci. 2022;29(1):102-110. doi: 10.1016/j.sjbs.2021.08.063.

14. behavior and hippocampus morphology in Swiss mouse model. Saudi J Biol Sci. 2022

Jan;29(1):102-110. doi: 10.1016/j.sjbs.2021.08.063. Epub 2021 Aug 25. PMID:

35002399; PMCID: PMC8716897.

15. Azab AE, Ebrahim SA. Exposure to electromagnetic fields induces oxidative stress and

pathophysiological changes in the cardiovascular system. J Appl Biotechnol Bioeng.

2017;4(2):00096.

16. Elmas O. Effect of electromagnetic field exposure on the heart: A systemic review.

Toxicol Ind Health. 2016;32(1):76-82.

17. Herdell L, Carlberg M, Mild KH. Pooled analysis of case-control studies on malignant

brain tumors and the use of mobile and cordless phones including living and deceased

objects. J Oncol. 2011;35(5):1465-1474.

18. Colleti V, Mandala M, Manganotti P, Ramat S, Sacchetto L, Colleti L. Intraoperative

observation of changes in cochlear nerve action potentials during exposure to

electromagnetic fields generated by mobile phone. J Neurol Neurosurg Psychiatry.

2011;82(7):766-771.

19. Falzone N, Huyser C, Becker P, Leszczynski D, Franken DR. The effect of pulsed 900-

MHz GSM mobile phone radiation on the acrosome reaction, head morphometry and

zona binding of human spermatozoa. Int J Androl. 2011;34:20-26.

20. Khillare B, Behari J. Effect of amplitude modulated radiofrequency radiation on

reproduction pattern in rats. Electro Magnetobiol Med. 1998;17:43-45.

21. Nour MM, Chen J, Allman-Farinelli M. Efficacy and external validity of electronic and

mobile phone-based interventions promoting vegetable intake in young adults: A

Systematic Review Protocol. J Med Internet Res. 2015;4(3):e4665.

22. Subhi Y, Bube SH, Rolskov BS, Skou TAS, Konge L. Expert involvement and adherence

to medical evidence in medical mobile phone apps: A systematic review. J Med Internet

Res. 2015;3(3):e79. doi: 10.2196/mobilehealth.4169.

23. Funk RK, Stockham AL, Laack NNI. Basics of Radiation Therapy. In: Herrmann J,

editor. Clinical Cardio-Oncology. Elsevier; 2016. p. 39-60.

24. Anand U, Dey A, Chandel AKS, Sanyal R, Mishra A, Pandey DK, et al. Cancer

chemotherapy and beyond: Current status, drug candidates, associated risks and progress

in targeted therapeutics. Genes Dis. 2022;10(4):1367-1401. doi:

10.1016/j.gendis.2022.02.007. Erratum in: Genes Dis. 2024;11(4):101211.

25. Okano H. Effects of static magnetic fields in biology: role of free radicals. Front Biosci.

2008;13:610–625. doi: 10.2741/3141.

26. International Agency for Research on Cancer. Ionizing Radiation, Part 1: X- and Gamma

(γ)-Radiation, and Neutrons. IARC Monographs on the Evaluation of Carcinogenic Risks

to Humans, No. 75. Lyon (FR): International Agency for Research on Cancer; 2000.

27. Omer H. Radiobiological effects and medical applications of non-ionizing radiation.

Saudi J Biol Sci. 2021 Oct;28(10):5585-5592. doi: 10.1016/j.sjbs.2021.05.071. Epub

2021 Jun 1. PMID: 34588869; PMCID: PMC8459055.

28. Bortkiewicz A. Health effects of Radiofrequency Electromagnetic Fields (RF EMF).

Industrial Health. 2019;57(4):403-405. Available from:

https://doi.org/10.2486/indhealth.57_400. Accessed April 24, 2024.

29. Syaza SKF, Umar R, Hazmin SN, Kamarudin MKA, Hassan A, Juahir H. J Fundam Appl Sci. 2017;9(2S):308-316

30. Kazemi E, Mortazavi SM, Ali-Ghanbari A, Sharifzadeh S, Ranjbaran R, Mostafavi-Pour Z, Zal F, Haghani M. Effect of 900 MHz Electromagnetic Radiation on the Induction of ROS in Human Peripheral Blood Mononuclear Cells. J Biomed Phys Eng. 2015;5(3):105-114

31. ICNIRP. ICNIRP Guidelines for limiting exposure to electromagnetic fields (100 kHz to 300 GHz). Health Phys. 2020;118:483–524.

32. SCENIHR. Opinion on Potential Health Effects of Exposure to Electromagnetic Fields (EMF). 2015. Available from: https://ec.europa.eu/health/scientific_committees/emerging/docs/scenihr_o_041.pDf (accessed on 25 November 2021).

33. Desai NR, Kesari KK, Agarwal A. Pathophysiology of cell phone radiation: Oxidative stress and carcinogenesis with focus on male reproductive system. Reprod Biol Endocrinol. 2009;7:114.

34. Vornoli A, Falcioni L, Mandrioli D, Bua L, Belpoggi F. The contribution of in vivo mammalian studies to the knowledge of adverse effects of radiofrequency radiation on human health. Int J Environ Res Public Health. 2019;16:3379.

35. Kesari KK, Kumar S, Nirala J, Siddiqui MH, Behari J. Biophysical evaluation of radiofrequency electromagnetic field effects on male reproductive pattern. Cell Biochem Biophys. 2013;65:85–96.

36. American Cancer Society. Available from: https://www.cancer.org/cancer/cancer-causes/

37. Deanna MM, Benjamin IB. A review of dietary (phyto)nutrients for glutathione support.

Nutrients. 2019;11(9):2073.

38. Aquilano K, Baldelli S, Ciriolo MR. Glutathione: new roles in redox signaling for an old

antioxidant. Front Pharmacol. 2014 Aug 26;5:196. doi: 10.3389/fphar.2014.00196.

PMID: 25206336; PMCID: PMC4144092.

39. Langbein K, Hesse J, Gussew A, Milleit B, Lavoie S, Amminger GP, Gaser C, Wagner

G, Reichenbach JR, Hipler UC, Winter D, Smesny S. Disturbed glutathione antioxidative

defense is associated with structural brain changes in neuroleptic-naïve first-episode

psychosis patients. Prostaglandins Leukot Essent Fatty Acids. 2018;136:103-110.

40. Markov MS. Effects of electromagnetic fields on biological systems. Electromagn Biol

Med. 2013;32:121-122. doi: 10.3109/15368378.2013.776333.

41. Goldoni J. Zastita od neionizirajucih zracenja u medicini [Non-ionizing radiation

protection in medicine]. Arh Hig Rada Toksikol. 1994 Jun;45(2):175-187. Croatian.

PMID: 7980026.

42. National Research Council (US) Committee on Assessment of the Possible Health

Effects of Ground Wave Emergency Network (GWEN). Assessment of the Possible

Health Effects of Ground Wave Emergency Network. Washington (DC): National

Academies Press (US); 1993. Effects of Electromagnetic Fields on Organs and Tissues.

Available from: https://www.ncbi.nlm.nih.gov/books/NBK208983/.

43. Talbot SR, Biernot S, Bleich A, van Dijk RM, Ernst L, Häger C, Helgers SOA, Koegel B,

Koska I, Kuhla A, Miljanovic N, Müller-Graff FT, Schwabe K, Tolba R, Vollmar B,

Weegh N, Wölk T, Wolf F, Wree A, Zieglowski L, Potschka H, Zechner D. Defining

body-weight reduction as a humane endpoint: a critical appraisal. Lab Anim. 2020

Feb;54(1):99-110. doi: 10.1177/0023677219883319. Epub 2019 Oct 30. PMID:

31665969.

44. Marino AA, Berger TJ, Austin BP, Becker RO, Hart FX. Evaluation of electrochemical

information transfer system: I. Effect of electric fields on living organisms. J Electrochem

Soc. 1976;123:1199. doi: 10.1149/1.2133035.

45. Meo SA, Arif M, Rashied S, Husain S, Khan MM, Vohra MS, Usmani AM, Imran MB,

Al-Drees AM. Hypospermatogenesis and spermatozoa maturation arrest in rats induced

by mobile phone radiation. J Coll Physicians Surg Pak. 2011;21:262–265.

46. Chavdoula ED, Panagopoulos DJ, Margaritis LH. Comparison of biological effects

between continuous and intermittent exposure to GSM-900-MHz mobile phone radiation:

Detection of apoptotic cell-death features. Mutat Res. 2010;700(1-2):51-61.

47. Tanil CT, Yong MH. Mobile phones: The effect of its presence on learning and memory.

PLoS One. 2020;15(8):e0219233.

48. Batool S, Bibi A, Frezza F, Mangini F. Benefits and hazards of electromagnetic waves,

telecommunication, physical and biomedical: a review. Eur Rev Med Pharmacol Sci.

2019;23(7):3121-3128.

49. Abouelsayed A, Hezma AM, El-Bahy GS, Abdelrazzak AB. Modification of protein

secondary structure as an indicator of radiation-induced abscopal effect: A spectroscopic

investigation. Spectrochim Acta A Mol Biomol Spectrosc. 2023;287:122093.

50. Abel R. Cataracts. In: Rakel D, editor. Integrative Medicine. Elsevier; 2018. p. 830-

837.e1.

51. Aaseth J, Gerhardsson L, Skaug MA, Alexander J. General Chemistry of Metal Toxicity

and Basis for Metal Complexation. In: Aaseth J, Crisponi G, Andersen O, editors.

Chelation Therapy in the Treatment of Metal Intoxication. Academic Press; 2016. p. 1-33.

52. Ballatori N, Krance SM, Notenboom S, Shi S, Tieu K, Hammond CL. Glutathione dysregulation and the etiology and progression of human diseases. Biol Chem. 2009;390(3):191-214.

53. Moustafa YM, Moustafa RM, Belacy A, Abou-El-Ela SH, Ali FM. Effects of acute exposure to the radiofrequency fields of cellular phones on plasma lipid peroxide and antioxidase activities in human erythrocytes. J Pharm Biomed Anal. 2001;26(4):605-8.

54. Wilson BW, Matt KS, Morris JE, Sasser LB, Miller DL, Anderson LE. Effects of 60 Hz magnetic field exposure on the pineal and hypothalamic pituitary-gonadal axis in the Siberian hamster (Phodopus sungorus). Bioelectromagnetics. 1999;20:224–232.

55. Sokolovic D, Djordjevic B, Kocic G, Babovic P, Ristic G, Stanojkovic Z, Sokolovic DM, Veljkovic A, Jankovic A, Radovanovic Z. The effect of melatonin on body mass and behaviour of rats during an exposure to microwave radiation from mobile phone. Bratisl Lek Listy. 2012;113(5):265-9.

56. Sandrey MA, Vesper DN, Johnson MT, Nindl G, Swez JA, Chamberlain J, Balcavage WX. Effect of Short Duration Electromagnetic Field Exposures on Rat Mass. Bioelectromagnetics. 2002;23(1):2–6.

57. Shabat MM, Shahwan OA. Analysis of the biochemical parameters of liver, kidney functions and thyroid stimulated hormone in children after exposure to mobile phone base station radiation and therapeutic action of olive oil. IUG J Nat Stud. 2017;79–84.

58. Lee JS, Ahn SS, Jung KC, Kim YW, Lee SK. Effects of 60 Hz electromagnetic field exposure on testicular germ cell apoptosis in mice. Asian J Androl. 2004;6:29–34.

59. Gerardi G, De Ninno A, Prosdocimi M, Ferrari V, Barbaro F, Mazzariol S, Bernardini D,

Talpo G. Effects of electromagnetic fields of low frequency and low intensity on rat

metabolism. Biomagn Res Technol. 2008;6:3.

60. Eid FA, El-Gendy AM, Zahkouk SA, El-Tahway NA, El-Shamy SA. Ameliorative effect

of two antioxidants on the liver of male albino rats exposed to electromagnetic field.

Egypt J Hosp Med. 2015;58:74–93.

61. Ridtitid W, DeWitt JM, Schmidt CM, Roch A, Stuart JS, Sherman S, et al.

62. Wang X, Wang Y, Wang M, et al. The association between serum lipid levels and

histological type of breast cancer. Eur J Med Res. 2022;27:154

63. Rodrigues Dos Santos C, Fonseca I, Dias S, Mendes de Almeida JC. Plasma level of

LDL-cholesterol at diagnosis is a predictor factor of breast tumor progression. BMC

Cancer. 2014;14:132–42.

64. Orho-Melander M, Hindy G, Borgquist S, Schulz CA, Manjer J, Melander O, et al. Blood

lipid genetic scores, the HMGCR gene and cancer risk: a Mendelian randomization study.

Int J Epidemiol. 2018;47(2):495–505.

65. Borek C. Antioxidant Health Effects of Aged Garlic Extract. J Nutr. 2001;131(3):1010S-

1015S.

Supporting Information

Click here to access/download
Supporting Information
ETHICAL CLEARANCE.pdf