Spatiotemporal distribution of neural crest cells in the

common wall lizard *Podarcis muralis*

Robin Pranter[1] (https://orcid.org/0000-0003-3615-0281) and Nathalie

Feiner[1]

[1]Department of Biology, Lund University, Sweden

Corresponding authors: robin.pranter@biol.lu.se, +46738413877 and nathalie.feiner@biol.lu.se

Running title: Neural crest cells in common wall lizards

Key words: Gene expression patterns, cell migratory routes, squamate reptiles, HNK-1

Data availability statement: Not applicable.

Funding statement: This work was supported by Kungliga Fysiografiska Sällskapet i Lund

(42010), Jörgen Lindströms stipendiefond to RP, and by the European Research Council (948126),

and Swedish Research Council (2020-03650) to NF.

Conflict of interest disclosure: The authors of this study declare no conflicts of interest.

Ethics approval statement: The embryos were collected from eggs laid by animals kept under

breading permit 5.2.18-10993/18 granted by the Swedish ministry of agriculture.

Patient consent statement: Not applicable

Permission to reproduce material from other sources: Not applicable.

Abstract

Background: Neural crest cells (NCCs) are migratory embryonic stem cells that give rise to a diverse set of cell types. Here we describe the dynamic distribution of NCCs in developing embryos of the common wall lizard *Podarcis muralis* inferred from ten markers. Our aim is to provide insights into the NCC development of lacertid lizards and to infer evolutionary modifications by comparisons to other tetrapods.

Results: NCC migration is ongoing at oviposition, following three streams in the head and multiple in the trunk. From 21ss, we observe expression patterns indicating the beginning of differentiation towards mesenchymal and neuronal fates. By 35ss, migration is restricted to caudal levels, and fully differentiated chromaffin cells are observed.

Conclusions: We find that some markers show patterns that differ from other tetrapods. For example, the antibody HNK-1 labels three NCC streams from the hindbrain while some comparable reptile studies describe four. However, the information emerging from all markers combined shows that the overall spatiotemporal distribution of NCCs in the common wall lizard is largely conserved with that of other tetrapods. Our study highlights the dynamic nature of seemingly canonical marker genes and provides the first description of spatiotemporal NCC dynamics in a lacertid lizard.

Introduction

Neural crest cells (NCCs) are transient, migratory stem cells that are unique to vertebrates and that give rise to many different cell types, thus contributing to a diversity of traits. NCCs are specified in the dorsal neural tube from where they delaminate in an epithelial-to-mesenchymal transition (EMT) and subsequently migrate throughout the embryo and differentiate. The cell types they give rise to include chromatophores responsible for skin color, osteocytes and chondrocytes that constitute the facial skeleton, neurons and glia of the peripheral nervous system and endocrinal chromaffin cells in the adrenal gland.[1]

NCCs originated at the dawn of vertebrates, and many of the vertebrate key innovations (most notably the jaw and other craniofacial structures) develop from NCCs.[2] For this reason, a substantial body of research has targeted the evolutionary origin of NCCs by investigating NCCs in early branching vertebrates and NCC-like cells in non-vertebrate chordates.[3–5] In addition to these attempts to elucidate the evolutionary origin of NCCs in a macro-evolutionary context, much research has focused on investigating their function and development in a number of model organisms, most notably mouse (*Mus musculus*), chicken (*Gallus gallus*), African clawed frog (*Xenopus laevis*) and zebrafish (*Danio rerio*). Motivation for this research partially stems from the fact that failure of NCC functions cause neurocristopathies, a highly heterogeneous family of congenital diseases,[6] which makes them relevant from a medical perspective.

In squamate reptiles (lizards and snakes), NCC biology has been investigated in three species: the California kingsnake (*Lampropeltis getula californiae*[7]), the Egyptian cobra (*Naja haje haje*[8]) and the veiled chameleon (*Chamaeleo calyptratus*[9]). The reports on the two snake species mainly focus on the migration of NCCs in the trunk, while the report in chameleon provides

a comprehensive description of both cranial and trunk NCCs. Both snakes and chameleons have

highly derived body plans including in NCC-derived traits such as skull morphology.

In the chameleon, cranial NCC migration starts between somite stage 4 and 6 (4ss and

6ss),[9] which is similar to mouse (5ss[10–12]), human (4ss[13]) and chicken (6ss[14(p66)]). In general, cranial

NCC migration typically follows three main dorsolateral streams.[15] The NCCs in the anterior-most

stream are specified in the caudal midbrain and the first three rhombomeres of the hindbrain and

contribute cells to the rostrum, maxillary process and first pharyngeal arch.[15] The second stream

is constituted by NCCs that are specified in rhombomeres three through five (anterior to the otic

vesicle) and migrate to the second pharyngeal arch.[15] And lastly, the NCCs in the third stream are

specified in rhombomeres five and six (posterior to the otic vesicle) and migrate to the third

pharyngeal arch.[15] In the chameleon, however, Diaz et al.[9] identified a fourth stream of cranial

NCCs starting posteriorly to the third stream and migrating to the fourth pharyngeal arch. An

equivalent stream has also been described in alligator (*Alligator mississippiensis*) and ostrich

(*Struthio camelus*).[16]

NCC migration in the trunk follows multiple, periodic streams starting in the dorsal neural

tube. Viewed in cross-sections in a transverse plane, these streams typically follow two different

routes.[17–19] An early wave of migrating cells follows a ventromedial route traversing between

somites and through the anterior portion of somites and giving rise to, for example, peripheral

nervous system and chromaffin tissue.[17–19] A later wave follows a dorsolateral route and gives rise

to chromatophores.[17–19] While this pattern seems to be largely conserved across tetrapods,[7–9] it has

been reported that some NCCs in the second wave follow a ventromedial rather than dorsolateral

route in the California kingsnake.[7] The equivalent was not reported for chameleon or cobra.[8,9]

In between and partially overlapping with the cranial and trunk NCCs, there is a small

subpopulation of NCCs called vagal NCCs.[15] Similar to trunk NCCs, vagal NCCs migrate along

both dorsolateral and ventromedial routes.[20] The dorsolateral migrating NCCs reach the

pharyngeal arches and the heart.[20] Some vagal NCCs of both the dorsoventral and ventromedial

streams reach the foregut which they then follow caudally.[20] Vagal NCCs have not been described

in squamates.

While efforts in squamates have increased our understanding of NCC specification,

migration and differentiation across vertebrates, evolutionary changes in NCC behavior and

potentially associated phenotypic changes are currently underexplored.[21] Since NCCs give rise to

functionally diverse and ecologically relevant traits ranging from coloration (e.g., pigment cells)

and morphology (e.g., skull and jaw bones) to physiology and behavior (e.g., adrenal gland

influencing aggression), their coupling in a shared developmental origin can potentially have

evolutionary consequences. Indeed, this coupling has been suggested to explain the domestication

syndrome, which is based on the observation that domesticated animals share a set of changes in

NCC-derived traits.[22,23] In general, due to their central role during development, modifications in

the regulation of NCCs are expected to have the potential to generate biases in micro-evolutionary

variation in natural populations of vertebrates and therefore contribute to adaptation and

diversification.[21,24,25] To evaluate hypotheses like this, establishing patterns of NCC migration in

species where NCCs may be implied in micro-evolutionary adaptation is crucial.

Here we present a description of NCCs in the common wall lizard (*Podarcis muralis*),

which is a small (approx. 50–70 mm snout-to-vent length[26]), egg-laying lacertid. This species

belongs to a genus of lizards distributed in the Mediterranean and Southern/Central Europe that

shows high variation in coloration and color patterns.[27,28] Within the common wall lizard, there is

substantial variation in predominantly NCC-derived traits, namely coloration (ranging from

brown/tan to black/green),[29–32] social behavior (in particular aggression in male-male

interactions),[33,34] and morphology (e.g., body size, head size and shape).[29,32] In Italy, the strength

of expression of these traits is tightly correlated across the landscape, culminating in the

nigriventris syndrome at one end of the extreme and the ancestral phenotype at the other end.[32]

The genetic basis of these trait differences has been identified as being polygenic, with more than

half of all identified genes having a known association with the regulation of NCCs.[32] This

prompted us to investigate NCC biology in this species. Females typically lay 2-3 clutches per

breeding season with 2-10 eggs per clutch.[26(p189)] We use a set of ten NCC markers that target

different aspects of NCC biology to investigate the spatial distribution and infer migration patterns

of NCCs from the earliest stage at oviposition (~13ss) until the latest stages of NCC activity (~

64ss). To identify evolutionary changes to the spatiotemporal distribution of NCCs , we compare

our observations to equivalent patterns described in other squamates and more distantly related

tetrapods. We find broad conservation in general NCC specification, migration, and

differentiation, but also taxonomic differences in the expression patterns of individual markers.

Taken together, our description of spatial and temporal patterns of NCCs during embryonic

development expands our understanding of the taxonomic diversity in NCC biology and presents

a first step towards investigating the role of NCCs in shaping micro-evolutionary patterns.

Results and Discussion

Given that NCCs are a heterogeneous and highly dynamic cell population, there is no single marker

that distinguishes them from other cell types throughout development. We therefore use a two-

pronged approach to resolve spatiotemporal dynamics of putative NCCs. Firstly, we use

immunostaining with antibodies against Human Natural Killer 1 (HNK-1) to label putative NCCs

throughout development. HNK-1 is a carbohydrate on the cell surface of NCCs that is involved in

cell-cell communication during migration[35,36] and it is a widely used marker of newly formed and

migrating NCCs[16,17,37–43] (note however that it is not labelling NCCs in mouse)[44]. This antibody

has been successfully applied in other squamate species,[7–9] which allows a straightforward

comparison between our observations in the common wall lizard and other squamates.

Since HNK-1 is known to also label other cell types, in particular at late developmental

stages,[45] we complement HNK-1 immunostainings with gene expression analyses of NCC marker

genes. For this second approach, we select a panel of nine marker genes whose expression patterns

are visualized using *in situ* hybridizations. Collectively, these marker genes label the majority of

NCC subpopulations and target various stages of NCC development.[46] Note however that some

markers do not exclusively label NCCs since they fulfil pleiotropic functions (e.g., *Twist1* stains

mesodermal cells in addition to NCCs[47]). Specifically, we select *Wnt1* and *Snai2* to label

predominantly early NCCs, *Snai1*, *Twist1,* and *Sox9* to mark NCCs biased towards mesenchymal

fates and *FoxD3*, *Sox10 and Phox2b* to mark NCCs biased towards neuronal and glial fates.[46,48,49]

In addition, we use *Tyrosine hydroxylase* (*Th*) to label NCCs differentiating into chromaffin cells.[50]

While HNK-1 immunostaining and *in situ* hybridizations provide snapshots of the location of

NCCs throughout development, we infer migratory routes from the series of observed imaging

data aided by comparisons to relevant literature. For clarity, results for cranial, vagal (where

applicable) and trunk NCCs are presented separately for each marker.

HNK-1 stains migratory and early differentiating NCCs

HNK-1-positive cells are present in all investigated stages (five stages ranging from 13ss to 35ss;

Figure 1A-P). Cranial NCC migration can be inferred along three routes. Each stream originates

in the dorsal midline and extends ventrally in the developing head (Figure 1 A, B, D, E and J;

dashed arrows indicate inferred streams). The first stream starts at the level of the midbrain-

hindbrain boundary and reaches towards the rostrum, maxillary process and the first pharyngeal

arch (Figure 1A). The second and third inferred streams originate in the hindbrain on either side

of the otic vesicle, extending towards the second and third pharyngeal arches respectively (Figure

1D and E). While the first stream is visible already at the first investigated stage (13ss), the second

and third streams are seen more clearly from 17ss. The three streams of migrating cranial NCCs

remain recognizable until 22ss (Figure 1S), after which migration of cranial NCCs presumably

seizes. By then, HNK-1 signal is widely spread in the developing head including structure that will

give rise to the facial skeleton such as the rostrum, maxillary process, and the first three pharyngeal

arches (Figure 1J), and some cranial placodes (Figure 1N), which are co-derived from NCCs and

placodal cells.[56,57] HNK-1 signal in the cranial peripheral nervous system is maintained as the

cranial placodes give rise to cranial nerves (Figure 1O; cranial nerve nomenclature follows Diaz

et al. (2019),[9] Lee et al. (2003)[57] and Streeter (1905)[58]). There is also strong segmental staining in

the rhombomeres of the hindbrain (Figure 1K and L).

At vagal levels, NCC migration can first be inferred by 13ss along a dorsolateral route to

the heart (open arrows in Figure 1A and C), which is also strongly stained by HNK-1 (indicated

with "H" in Figure 1A, C and D). At later developmental stages, there is a branch from the third

cranial stream extending caudally into the trunk, which potentially indicates a stream of vagal

NCCs on their way to populate the gut and give rise to the enteric nervous system (indicated with

"Va" in Figure 1D). Consistently, cross-sections of the trunk reveal staining around the gut (Figure

1H-I); a similar staining has been shown in chicken.[38]

In the trunk, HNK-1 signal in NCCs can be roughly divided into four stages: premigratory,

early migratory, migratory, and early neural differentiation. HNK-1 signal is present in a stripe

along the dorsal midline where premigratory NCCs are expected (Figure 1B, E and G).

Immediately lateral to the dorsal midline, HNK-1 stains early migratory NCCs in loose cell

aggregates, not yet forming migratory streams (Figure 1A, B, F, G and M). The staining of

premigratory and early migratory NCCs is found at progressively more posterior positions along

the trunk as development proceeds. At more anterior levels, multiple streams of NCCs can be

inferred between the somites (arrows in Figure 1F, G and M), and, to a lesser extent, at the level

of the somites (open arrow in Figure 1M). Cross-sections reveal that the streams of migratory

NCCs take a ventral route between somites (inter-somitic; Figure 1H), while they take a

dorsolateral route at the levels of the somites (Figure 1I). NCC migration in the most caudal region

of the trunk is still ongoing in the oldest investigated stages (28ss and 35ss). While HNK-1 signal

is known to mark the NCC-derived dorsal root ganglia in chicken,[38] and also a turtle (*Trachemys*

scripta)[59], an alligator (*Alligator mississippiensis*)[16] and the Californian kingsnake (*Lampropeltis*

getula calilforniae),[7] this staining is absent in common wall lizards (Figure 1P). The reported

HNK-1 staining in dorsal root ganglia in chameleon is also less distinct than in other organisms.[9]

This emphasizes that HNK-1 is neither specific, nor general enough to figure as a sole NCC

marker, and therefore requires complementation with other markers. In later stages, there is some

HNK-1 staining in the mesonephros (Figure P). This is also seen in other reptiles, but it is unclear

if it is due to NCCs contributing to this organ.[7,9] At later stages (35ss), we detect HNK-1 signal in

blood cells inside the aorta and heart (Figure 1O). Presumably, this is caused by the fact that blood

cells possess HNK-1 on their cell surface, despite not being NCC-derived.[45]

Wnt1 and Snai2 are expressed in premigratory NCCs

Besides being a crucial organizer of the midbrain-hindbrain region,[60] *Wnt1* is a classic marker of

the neural border (including the rhombic lips[61]) and inducer of NCC specification.[46,62] And the

Wnt1-Cre transgenic lines have been instrumental in numerous experimental studies of NCCs in

mice,[48,63,64] highlighting its crucial role as a NCC regulator.

Wnt1 is expressed in all investigated stages (10 stages ranging from 13ss to 60ss), mostly

along the dorsal midline (Figure 2A-E). Through most of the investigated developmental stages

(16-30ss), there is a noticeable gap of expression in the most anterior part of the hindbrain (the

metencephalon; asterisks in Figure 2A-D), separating the expression into an anterior and a

posterior domain. This pattern has been described in chicken.[60] The gap of expression overlaps

with the origin of the first cranial stream as inferred from HNK-1 staining.

The anterior domain runs along the dorsal midline of the head from the level of the

diencephalon to the midbrain-hindbrain boundary (Figure 2A, C). It has some lateral extensions

that are presumably not NCC related and will not be described in further detail. The posterior

domain consists of two parallel lines running along the rhombic lips in the hindbrain (Figure 2D)

and continuing caudally along the dorsal midline of the trunk (Figure 2B), extending caudally

approximately as far as to the last formed somite pair, and continuously extends posteriorly as

development progresses (Figure 2A, C and E). The caudal extension of *Wnt1* expression is

concordant with premigratory NCCs as inferred from HNK-1 staining.

Snai1 and *-2* are classical NCC markers previously known as *Snail* and *Slug*.[65] They are

transcription factors that control cadherin-transcription and thereby EMT.[46] There was some

uncertainty in regards to the homology relationships between these genes, and EMT is induced by

paralogous genes in chicken (*Snai2*) and mouse (*Snai1*).[66] We confirm the expected orthology

relationships between *Snai1* and *-2* of the common wall lizard and their homologs in human, mouse

and chicken (see Methods) and investigate both of them using *in situ* hybridization. *Snai2*

expression is investigated in eight stages ranging from 13ss to 51ss.

Snai2 expression in the head is found in putative premigratory NCCs along the dorsal

midline and the rhombic lips in the hindbrain (Figure 3A and B), and in the three streams of cranial

NCC migration previously inferred with HNK-1 staining (dashed arrows in Figure 3A and C). In

cross-section through the second stream, *Snai2* expression can be seen from the dorsal tip of the

neural crest to the pharyngeal arches (Figure 3C). While a caudally migrating vagal stream of

NCCs is only weakly suggested by HNK-1, it is strongly marked by *Snai2* expression, visible as a

branch from the third cranial stream (Figure 3A). In cross-section at the level of the heart, strong

Snai2 expression is located lateral to the gut, presumably indicating caudally migrating vagal

NCCs (Figure 3D).[20]

Similar to HNK-1, *Snai2* expression in putative premigratory NCCs along the dorsal

midline of the trunk progressively extends caudally as development progresses (arrows in Figure

3A and E). *Snai2* expression is also evident in multiple streams of migratory NCCs traversing the

somites (Figure A).

Taken together, the expression patterns of *Wnt1* and *Snai2* are compatible with the expected

location of premigratory NCCs. As in chicken, but not mouse, *Snai2* seems to be expressed where

NCCs are expected to undergo EMT.[66] Similar to HNK-1, *Snai2*, but not *Wnt1*, is expressed in

migratory NCCs. In addition, *Snai2* expression labels putative vagal NCCs more clearly than

HNK-1.

Twist1, Snai1 and Sox9 are expressed in NCCs biased towards mesenchymal fate

Twist1 is a well-known NCC marker and transcription factor that has been studied in various other organisms. It has been found to be involved in EMT,[67] maintenance of a migratory state, repression of neuronal differentiation,[68] and induction of chondrocyte differentiation.[48]

Expression of *Twist1* is observed in all investigated stages (7 stages ranging from 13 through 51ss; Figure 4A-D). Cranial *Twist1* expression is missing along the dorsal midline but is consistent with the expected location of migratory NCCs (including the vagal branch) observed with HNK-1 and *Snai2* (dashed arrows in Figure 4A and C). The lack of *Twist1* expression in the presumed premigratory NCCs in the dorsal midline is consistent with findings in mouse and chicken,[68,69] but different from frog and zebrafish.[67] Later expression is mostly restricted to the maxillary process and pharyngeal arches (arrow and arrowhead in Figure 4D). Given the described role of *Twist1* as an inducer of chondrocytes,[43] this expression likely label the formation of craniofacial structures in these regions. At early stages, there is also broad but transient expression of *Twist1* in the head ectoderm. While *Twist1* is also extensively expressed in the trunk, it is restricted to putative mesodermal cells in the somites and limb buds and will therefore be disregarded here.

Snai1 is expressed in all investigated stages (7 stages ranging from 16 to 51ss) and its expression patterns resemble that of *Twist1* (Figure 5A-C). In contrast to *Snai2*, *Snai1* is not expressed where premigratory NCCs are expected to undergo EMT in the common wall lizard (Figure 5A and B), which is different from findings in mouse but consistent with findings in chicken.[66] Compared to Twist1, the different streams of cranial migration are not as easily discernible for *Snai1* since the

intensity of the expression signal is generally high (Figure 5A). Given this high intensity, it is

possible that not only NCCs contribute to the signal.

Together with *Snai1, -2* and *Twist1*, *Sox9* is known to be a crucial regulator of EMT[46,70] and it is

also known to regulate migration and chondrocyte differentiation of cranial NCCs.[46] However, in

the trunk, *Sox9* is expressed in premigratory but not migratory NCCs in chicken.[71]

In the common wall lizard, *Sox9* is expressed in all investigated stages (8 stages ranging

from 13ss through 51ss; Figure 6A-D). In the head, *Sox9* appears to label both premigratory

(arrows in Figure 6A) and migratory (dotted arrows in Figure 6A) NCCs. In the third stream, it is

mainly the caudally directed putative vagal branch that can be inferred. Later in development, the

expression in the inferred streams extends to much of the surface of the head (Figure 6C). At this

point, the expression pattern resembles that of late *Twist1* expression.

Sox9 expression in the trunk is consistent with premigratory NCCs along the dorsal

midline, resembling that of *Snai2*. There is also expression in what appears to be early migratory

NCCs just lateral to the dorsal midline that have not yet condensed into migratory streams (Figure

6B and C). Otherwise, *Sox9* does not appear to be expressed in migratory NCCs in the trunk, which

is consistent with findings in chicken.[71] There is expression in non-NCC related structures (Figures

6B and C), which will not be described in more detail here. Some *Sox9* expression can be seen in

the heart at 39ss (Figure 6D) where it stains a structure that may be the outflow tract, which is

partially derived from NCCs in mouse.[68]

In summary, the expression patterns of *Snai1*, *Twist1* and *Sox9* generally support the

spatiotemporal location of migratory cranial neural crest cells inferred from HNK-1. In addition,

they distinctly label regions such as the maxillary process and pharyngeal arches where facial

skeleton will develop. *Sox9* is also expressed in the heart and in putative premigratory NCCs along

the dorsal midline of both head and trunk.

***Sox10*, *FoxD3* and *Phox2b* are expressed in NCCs biased towards neuronal and glial fates**

After HNK-1, *Sox10* is arguably the most canonical NCC marker, and together with *FoxD3* and

Ets1 it is considered a NCC specifier gene.[46] It is a marker of migratory NCCs,[72] and it is necessary

for NCC differentiation into melanocytes, neurons and glia.[46]

Sox10 is expressed in all investigated stages (11 stages ranging from 13 to 64ss; Figure 7

A-P). All three streams of migratory cranial NCCs inferred from the HNK-1 staining are clearly

expressing *Sox10* (Figure 7A and B). However, *Sox10* expression, in contrast to HNK-1, *Snai2,*

Snai1, Twist1 and *Sox9* does not stain the streams all the way to the rostrum, maxillary process

and pharyngeal arches (Figure7A). As development progresses, cranial *Sox10* expression, more

distinctly than HNK-1 staining, localizes in some cranial placodes (arrow in Figure 7H-J) and

subsequently the corresponding nerves of the cranial peripheral nervous system (Figure 7L, M and

P). A putative vagal stream migrating caudally in the trunk from the third cranial stream is marked

by *Sox10* expression (arrowhead in Figure 7H and J). Later in development, the vagus nerve is

clearly discernable through *Sox10* expression (Figure 7P).

Early *Sox10* expression in the trunk is largely similar to the spatiotemporal distribution of

NCCs inferred from HNK-1 staining including putative premigratory and migratory NCCs (Figure

7A, C, H and K). However, the expression in the putative premigratory NCCs along the dorsal

midline is rather weak (asterisks in Figure 7C, H and K). In cross-section at the level of the heart

at 13ss, the inferred streams are revealed to migrate ventrally along the edge of the neural tube

(Figure 7D-F). By looking at different levels along the anterior-posterior axis, two different routes

of migration can be distinguished; one turns laterally and enters the somite (Figure 7D and E),

presumably give rise to dorsal root ganglia.[18] The other one continues ventrally (Figure 7D and F),

likely giving rise to sympathetic ganglia.[18] The expression in the migratory streams subsequently

condenses to a row of dorsal root ganglia in each somite with ventral projections to the much

smaller ganglia of the sympathetic chain (Figure 7L). The dorsal root ganglia, including their

dorsal and ventral roots and peripheral projections, are clearly visible in cross-section, while

sympathetic ganglia are evident more ventrally, though less strongly expressing *Sox10* (Figure

7O).

FoxD3 is an important transcription factor in NCC development.[17] Early in NCC development,

FoxD3 acts as an activator, allowing NCCs to delaminate from the neural tube, but later its role

changes and it maintains the multipotency of NCCs by repressing differentiation.[18,73,74]

In the common wall lizard, *FoxD3* is expressed in all investigated stages (9 stages from 13

to 64ss), and its expression patterns are similar to those of *Sox10*, although mostly not as strong.

Like *Sox10, FoxD3* stain premigratory NCCs along the dorsal midline (Figure 8A-C), migratory

NCCs including three cranial streams (Figure 8A and B), and NCCs condensing in the peripheral

nervous system of both the head and trunk (Figure 8 C and D). Unlike *Sox10*, *FoxD3* staining in

premigratory NCCs along the dorsal midline is strong. It also stains in the geniculate ganglion

(open arrow in Figure 8D) but not the otic vesicle.

Downstream of *Sox10, Phox2b* is a regulator of NCC differentiation into autonomic and especially

sympathetic neurons and chromaffin cell progenitors.[46,75–77] In the common wall lizard, *Phox2b* is

expressed in all investigated stages (7 stages ranging from 17-51ss) except the youngest (17ss;

data not shown). However, the expression is most distinct at 39ss (Figure 9A-C), and expression

before and after this stage is generally restricted and weak. *Phox2b* expression localizes to several

of the cranial ganglia, which are co-derived from the cranial placodes and NCCs.[78] There is also

significant expression in the hindbrain in the form of three longitudinal stripes on each side (Figure

9A), similar to previous results in mouse.[79] Contrary to previous results in mouse and chicken

there is no distinct expression of *Phox2b* in the dorsal root ganglia in any of the investigated

stages.[78,79]

Taken together, gene expression profiles of NCCs biased towards neuronal and glial fates (*Sox10*,

FoxD3 and *Phox2b*) largely confirm and expand the spatiotemporal distribution of NCCs inferred

from HNK-1 staining. Together their expression patterns clearly visualize the contribution of

NCCs to the development of the peripheral nervous system, including the sympathetic nervous

system.

Th expression identifies NCCs differentiating into chromaffin cells

In mouse, *Tyrosine hydroxylase* (*Th*) has been identified as a suitable marker for separating the

chromaffin cells from the sympathetic neurons.[76,80] It is an enzyme catalyzing one of the reactions

in the metabolic pathway that result in the production of adrenalins.[81]

Th expression is first detected at 35ss and is maintained at least until 51ss. The expression

is located internally in the trunk, at the level of the forelimb bud (Figure 10A-F). *Th* expression is

localized in two lateral parallel lines (Figure 10B) in the dorsomedial corner of the mesonephros

(Figure 10D). This is putatively the position where the sympathetic ganglia including the adrenal

glands are developing. The areas of expression are discontinuous (see for example arrowheads in

Figure 10E) concordant with chromaffin cells being restricted to the sympathetic ganglia including

the adrenal glands.

Conclusions

Taken together, the insights gained from HNK-1 immunostainings and gene expression patterns

of nine marker genes provide a detailed profile of the spatiotemporal distribution of NCCs in

developing wall lizard embryos (Figure 11). By the time of egg-laying, the embryo has already

reached at least 13ss at which extensive migration is ongoing in the head. Around 35ss, most

migration is completed and differentiation towards various fates is well underway. In the

following, we briefly summarize the major patterns of NCC development at four different stages.

At 14ss, premigratory NCCs are present along the dorsal neural tube all the way from the

midbrain-hindbrain boundary to the last formed somite pair (orange in Figure 11A). Streams of

cranial NCCs are already migratory. In the caudal trunk, there is no migration observed at this

stage. At more anterior regions in the trunk, a field of NCCs lateral to the dorsal midline indicates

that the first wave of NCC migration has started but these cells have not yet entered the somites

(green in Figure 11A). In the anterior-most part of the trunk, migration has proceeded further with

the first migratory streams traversing the anterior parts of the somites (turquoise in Figure 11A).

At 21ss, NCC specification and migration in the trunk has proceeded further posteriorly

and migration now occurs through several of the anterior-most somites (approximately the first

13-15). In the head, the streams of migratory cranial NCCs have reached further and are bifurcating

more. NCCs with mesenchymal fate are located in the first pharyngeal arch (purple in Figure 11A).

and NCCs with neural fate are located in the otic vesicle and trigeminal nerve placode (magenta

in Figure 11A). Vagal NCCs appear as a branch extending caudally through the anterior trunk

from the third cranial stream.

By 28ss, NCC specification and migration in the trunk has proceeded even further

posteriorly and the first differentiated dorsal root ganglia occur in the anterior-most somites

(approximately the first 3; magenta in Figure 11A). The cranial streams are now more advanced.

NCCs with mesenchymal fate appear in the maxillary process and the rostrum and NCCs with

neural fate are expanding in cranial placodes and the first neural projections (magenta in Figure

11A).

At 35ss, specification and migration of NCCs is restricted to the caudal-most trunk region.

Neuronal differentiation is well underway in both the dorsal root ganglia and the sympathetic

ganglia of the trunk. At the level of the forelimb, some NCCs have differentiated into chromaffin

cells (yellow in Figure 11A). The mesenchymal NCCs are now distributed broadly in the head

including the first three pharyngeal arches, maxillary process, rostrum and close to the olfactory

pit. Cranial nerves have developed further, the olfactory (I), oculomotor (III), trigeminal (V),

glossopharyngeal (IX), vagus (X) and spinal accessory nerves are now all clearly distinguishable.

This synthesis is derived from the collective information gathered from ten markers (Figure

11B), which proved to be crucial to overcome incomplete information stemming from individual

markers. For example, HNK-1 immunostaining and *Phox2b* expression signals are not located in

the dorsal root ganglia, yet other markers, for example *Sox10* and *FoxD3*, clearly identify this

tissue as NCC-derived. The HNK-1 signal in dorsal root ganglia reported for the veiled chameleon[9]

is also less distinct than typically seen in other species. Thus, the lack or reduction of HNK-1 on

dorsal root ganglia may be a more widespread feature of lizards. In addition, since there are no

distinct NCC markers that identify exclusively NCCs, off-target staining necessarily needs to be

taken into account. Examples include the expression of *Snai1* and *Twist1* in trunk mesoderm or

the presence of HNK-1 on blood cells.

Comparing our synthesis of NCCs in the common wall lizard to those reported for other

squamates (the Californian kingsnake[7], Egyptian cobra[82] and veiled chameleon[9]) reveals that NCC

biology is largely conserved. One caveat for this comparison is that our analysis is restricted to

embryos older than 13ss, and we therefore lack observations of the earliest stages of NCC

specification and migration, particularly in the head. It was therefore not possible to investigate if

common wall lizards, similar to chameleons[9] and crocodiles[76], possess an early rostral stream of

migrating cranial NCCs stemming from the level of the forebrain. In future studies, efforts could

be made to obtain earlier embryos by, for example, treating gravid females hormonally to induce

premature egg laying,[84] or by selecting lizard species that lay eggs at earlier developmental stages.

The two main exceptions of NCC regulation in squamates relative to other tetrapods that

have been reported to date are the fourth stream of migrating NCCs from the hindbrain observed

in the veiled chameleon[9] (also found in an alligator and an ostrich)[16] and the late wave of trunk

NCCs taking a ventromedial (instead of the expected dorsolateral route), which was found in the

Californian kingsnake.[7] Our observations in the common wall lizard do not clearly identify a fourth

stream of NCCs from the hindbrain. However, to confidently rule out the existence of this stream,

further analyses including cell lineage tracing may be required. Regarding the ventromedial route

of late wave trunk NCCs, we do find ventromedial intra-somitic migration that is coinciding with

dorsolateral migration, but it is not clear that these correspond to the late wave ventromedial

migration reported for the California kingsnake.[7] On a more technical note, it would also be

desirable to confirm and cross-validate results in various species with the same antibody given that

observed species-level differences might be partially attributed to the use of different antibodies
targeting HNK-1.

In summary, our study provides the first complete description of NCCs, including cranial,
vagal and trunk NCCs, in a lacertid and contributes to our understanding of taxon-specific
differences in the dynamic distribution of this cell type. Thus, it lays the foundation for future work
targeting the role of this cell type in evolutionary diversifications.

Experimental procedures

Embryo collection

All embryos stem from captive breeding colonies of common wall lizards that were wild caught
in central Italy. Parents showed variation in phenotypes (coloration, morphology, and behavior),
including in traits derived from the neural crest. In the present study, we describe broad
developmental patterns that we do not expect to differ on a population level, and we therefore do
not discriminate between embryos of different parental origins.

Adult wall lizards were kept as described in Feiner et al., (2018).[85] Eggs were collected
during the breeding seasons (April-July) of 2021 to 2023. Each cage had a pot filled with moist
sand in which females laid their clutches. The eggs were collected each morning, and one egg per
clutch was dissected immediately to determine the developmental stage (somite count) of the
embryos in the clutch. The rest of the eggs were incubated at 24°C in moist vermiculite until the
appropriate stage was reached. We aimed to sample a distribution of stages from as young as
possible up to around 60ss. Based on 563 eggs, the mean embryonic stage on the day of oviposition
was estimated to be 21.4ss (sd = 5.2 somites) and the earliest embryos were at 11ss. The pace of

somite formation is constant and has previously been estimated to be four somites per day when

incubated at 24°C.[85]

Egg dissections were performed in cold, nuclease-free PBS (10mmol/L phosphate buffer,

2.7mmol/L KCl and 137mmol/L NaCl, pH 7.4) after which each embryo was immediately

transferred to 4% paraformaldehyde in PBS for overnight fixation at 4°C on a rotating platform.

After fixation, embryos were transferred to gradually increasing concentrations of methanol and

stored in 100% methanol at -20°C. A small subset of embryos intended to be used in probe

synthesis (see below) was dissected as described above, incubated in RNAlater (Invitrogen

AM7020) at 4°C for 24h and then transferred to -80°C for storage.

Immunohistochemistry

Paraformaldehyde-fixed embryos were rehydrated in PBS-T with decreasing concentrations of

methanol. Embryos were blocked using TN-blocking buffer with bovine serum albumin (1%

bovine serum albumin, 0.1mol/L Tris pH7.5, 0.15 mol/L NaCl in H_2O) followed by overnight

incubation with primary antibodies against HNK-1 (1:500; Anti-Hu CD57 eBioscience 11-

057742) in TN-blocking buffer. Embryos were washed with PBS and incubated overnight with

secondary antibodies (1:200; Alexa Flour 488 goat anti-mouse IgM (μ chain) Invitrogen A21042)

in TN-blocking buffer. Embryos were washed in PBS and stained with DAPI (1:500;

ThermoScientific 62248). Lastly, embryos were cleared by overnight incubation in RapiClear 1.49

(SUNJin Lab) and then mounted in fresh RapiClear medium between two 0.17μm cover slips using

iSpacers (SUNJin Lab). The stained and cleared embryos were imaged using a confocal

microscope (Leica TCS SP8) using a 20X/0.75 IMM objective (plan apochromatic multi-

immersion). Tiles were merged and 2D images were rendered using the LAS X software version

0.5.7.23225.

Probe synthesis for *in situ* hybridization

Total RNA was extracted from the RNAlater-stored embryos using the RNeasy Mini Kit (Qiagen)

including DNase digestion and stored in nuclease-free water at -80°C. Extracted RNA was used

as template in reverse transcription into cDNA using SuperScript III Reverse Transcriptase and

oligo-dT primers following the manufacturer's instructions. The selected marker genes were

amplified by polymerase chain reaction (PCR) from the cDNA using GoTaq G2 Hot Start Green

Master Mix (Promega), and primers designed using the PodMur_1.0 reference genome.[86] Primers

were designed to amplify 800-1200 bp of the gene of interest with approximately 500 bps in the

3´UTR and T7 and SP6 promoters attached to the reverse and forward primers, respectively (see

Table 1 for primer sequences). Amplified gene fragments were Sanger-sequenced to confirm each

amplicon's identity. PCR products were purified using the MinElute PCR Purification Kit

(Qiagen) and *in vitro* transcribed to produce antisense RNA-probes using the DIG Probe Synthesis

Kit (Roche). The resulting RNA-probes were purified using the RNeasy Mini Kit (Qiagen) and

stored in nuclease-free water at -80°C.

Whole mount *in situ* hybridization

Following the methodology in Feiner (2019)[87] paraformaldehyde-fixed embryos were rehydrated

in nuclease-free PBS-T (1x PBS with 0.1% Tween-20) using gradually lower concentrations of

methanol. Embryos were permeabilized using Proteinase K, followed by refixation using 4%

paraformaldehyde in PBS. Hybridization was done with 100 ng/mL probe at 68°C in hybridization

buffer (5x SSC pH 7.5, 50% deionized formamide, 50 µg/ml heparin, 0.3 mg/ml torula RNA, 2%

Tween-20, 2% blocking reagent and 10 mmol/L EDTA). Hybridized embryos were blocked in

blocking buffer (1.5% blocking reagent in PBS-T), followed by incubation with 0.1875 U/mL anti-

DIG antibody in blocking buffer. This was followed by 1.5 days of washes in maleic acid buffer

(15 mmol/L maleic acid pH7.5, 300 mmol/L NaCl and 0.1% Tween-20) before color development

with 2% NBT/BCIP in alkaline phosphatase buffer (0.1 mol/L Tris Base pH 9.5, 0.1mol/L NaCl,

1 mmol/L $MgCl_2$ and 0.1% Tween-20). Once a good signal-to-noise ratio was achieved, the

reaction was stopped by extensive washes in PBS-T and re-fixing in 4% paraformaldehyde. Some

stained embryos were cross-sectioned by hand using a scalpel. Stained embryos and sections were

photographed in glycerol using a Nikon SMZ18 stereo microscope with a Nikon SHR Plan Apo

2X WD: 20 objective and Nikon DS-Ri1 camera.

Gene orthology

Since the orthology relationships of *Snai1* and *Snai2* to other tetrapod genes was unclear[66] a gene

tree was constructed to confirm their identities. Both sequences from the common wall lizard were

obtained from Ensemble and blasted (protein blast) against the Human, mouse, and chicken

databases on NCBI, and the top hits were retrieved. Alignment (using the muscle algorithm),

model selection (JTT+G) and maximum likelihood tree construction was performed in MEGA

(Figure 12).[88]

Acknowledgements

We are grateful to Stanley Heinze, Michael Bok, Marjorie Lienard and Carina Rasmussen for

technical advice and Ola Gustafsson for help with confocal microscopy. We thank Tobias Uller

for valuable discussion and help with animal husbandry. We thank two anonymous reviewers for

insightful comments.

References

1.   Barresi MJF, Gilbert SF. *Developmental Biology*. 12th ed. Oxford, New York: Oxford
522   University Press; 2020.

523   2.   Alkobtawi M, Monsoro-Burq AH. The neural crest, a vertebrate invention. In: *Evolving*
524   *Neural Crest Cells*. Evolutionary cell biology. Boca Raton: CRC Press; 2020:5-66. DOI:
525   10.1201/b22096

526   3.   Abitua PB. The Hunt for Neural Crest in Invertebrate Chordates. In: *Evolving Neural Crest*
527   *Cells*. Evolutionary cell biology. Boca Raton: CRC Press; 2020:137-156.
528   https://doi.org/10.1201/b22096

529   4.   Kuratani S. Neural Crest and Craniofacial Evolution of Early Vertebrates. In: *Evolving*
530   *Neural Crest Cells*. Evolutionary cell biology. Boca Raton: CRC Press; 2020:185-218.
531   https://doi.org/10.1201/b22096

532   5.   Yu JK, Su YH. The Evolution of the Neural Border and Peripheral Nervous System—Insights
533   from Invertebrate Deuterostome Animals. In: *Evolving Nerual Crest Cells*. Evolutionary cell
534   biology. Boca Raton: CRC Press; 2020:103-136. https://doi.org/10.1201/b22096

535   6.   Cerrizuela S, Vega-Lopez GA, Méndez-Maldonado K, Velasco I, Aybar MJ. The crucial role
536   of model systems in understanding the complexity of cell signaling in human
537   neurocristopathies. *WIREs Mechanisms of Disease*. 2022;14(1):e1537.
538   https://doi.org/10.1002/wsbm.1537

539   7.   Reyes M, Zandberg K, Desmawati I, De Bellard ME. Emergence and migration of trunk
540   neural crest cells in a snake, the California Kingsnake (Lampropeltis getula californiae).
541   *BMC Developmental Biology*. 2010;10. https://doi.org/10.1186/1471-213X-10-52

542   8.   Khannoon ER, Alvarado C, Poveda R, de Bellard ME. Description of trunk neural crest
543   migration and peripheral nervous system formation in the Egyptian cobra Naja haje haje.
544   *Differentiation*. 2023;133:40-50. https://doi.org/10.1016/j.diff.2023.06.002

545   9.   Diaz RE, Shylo NA, Roellig D, Bronner M, Trainor PA. Filling in the phylogenetic gaps:
546   Induction, migration, and differentiation of neural crest cells in a squamate reptile, the
547   veiled chameleon (Chamaeleo calyptratus). *Developmental Dynamics*. 2019;248(8):709-
548   727. https://doi.org/10.1002/dvdy.38

549   10.   Nichols DH. Neural crest formation in the head of the mouse embryo as observed using a
550   new histological technique. *Development*. 1981;64(1):105-120.
551   https://doi.org/10.1242/dev.64.1.105

552   11.   Nichols DH. Formation and distribution of neural crest mesenchyme to the first pharyngeal
553   arch region of the mouse embryo. *American Journal of Anatomy*. 1986;176(2):221-231.
554   https://doi.org/10.1002/aja.1001760210

555   12.   Serbedzija GN, Bronner-Fraser M, Fraser SE. Vital dye analysis of cranial neural crest cell
556   migration in the mouse embryo. *Development*. 1992;116(2):297-307.
557   https://doi.org/10.1242/dev.116.2.297

558   13.   O'Rahilly R, Müller F. The development of the neural crest in the human. *Journal of
559   Anatomy*. 2007;211(3):335-351. https://doi.org/10.1111/j.1469-7580.2007.00773.x

560   14.   Le Douarin N, Kalcheim C. *The Neural Crest*. 2nd ed.; 1999. Accessed May 2, 2024.
561   https://www.cambridge.org/se/academic/subjects/life-sciences/cell-biology-and-
562   developmental-biology/neural-crest-2nd-edition,
563   https://www.cambridge.org/se/academic/subjects/life-sciences/cell-biology-and-
564   developmental-biology

565   15.   Le Douarin NM. The avian embryo as a model to study the development of the neural
566   crest: a long and still ongoing story. *Mechanisms of Development*. 2004;121(9):1089-1102.
567   https://doi.org/10.1016/j.mod.2004.06.003

568   16.   Kundrát M. Heterochronic shift between early organogenesis and migration of cephalic
569   neural crest cells in two divergent evolutionary phenotypes of archosaurs: Crocodile and
570   ostrich. *Evolution and Development*. 2009;11(5):535-546. https://doi.org/10.1111/j.1525-
571   142X.2009.00352.x

572   17.   Bronner-Fraser M. Analysis of the early stages of trunk neural crest migration in avian
573   embryos using monoclonal antibody HNK-1. *Developmental Biology*. 1986;115(1):44-55.
574   https://doi.org/10.1016/0012-1606(86)90226-5

575   18.   Harris ML, Erickson CA. Lineage specification in neural crest cell pathfinding.
576   *Developmental Dynamics*. 2007;236(1):1-19. https://doi.org/10.1002/dvdy.20919

577   19.   Weston JA. 6 Sequential Segregation and Fate of Developmentally Restricted Intermediate
578   Cell Populations in the Neural Crest Lineage. In: Bode HR, ed. *Current Topics in
579   Developmental Biology*. Vol 25. Academic Press; 1991:133-153.
580   https://doi.org/10.1016/S0070-2153(08)60414-7

581   20.   Kuo BR, Erickson CA. Vagal neural crest cell migratory behavior: A transition between the
582   cranial and trunk crest. *Developmental Dynamics*. 2011;240(9):2084-2100.
583   https://doi.org/10.1002/dvdy.22715

21. Brandon AA, Almeida D, Powder KE. Neural crest cells as a source of microevolutionary
variation. *Seminars in Cell & Developmental Biology*. Published online June 16, 2022.
https://doi.org/10.1016/j.semcdb.2022.06.001

22. Darwin C. *The Variation of Animals and Plants under Domestication*. Vol 2. 2nd ed.
London: John Murray; 1888.

23. Wilkins AS, Wrangham RW, Tecumseh Fitch W. The "domestication syndrome" in
mammals: A unified explanation based on neural crest cell behavior and genetics.
Genetics. 2014;197(3):795-808. https://doi.org/10.1534/genetics.114.165423

24. San-Jose LM, Roulin A. On the Potential Role of the Neural Crest Cells in Integrating
Pigmentation Into Behavioral and Physiological Syndromes. *Frontiers in Ecology and
Evolution*. 2020;8:278. https://doi.org/10.3389/fevo.2020.00278

25. Uller T, Moczek AP, Watson RA, Brakefield PM, Laland KN. Developmental Bias and
Evolution: A Regulatory Network Perspective. *Genetics*. 2018;209(4):949-966.
https://doi.org/10.1534/genetics.118.300995

26. Böhme W. *Handbuch der Reptilien und Amphibien Europas Band2/II Echsen III (Podarcis)*.
Wiesbaden: AULA-Verlag; 1986.

27. Eimer T, Eimer A. *Untersuchungen Über Das Variiren Der Mauereidechse : Ein Beitrag Zur
Theorie von Der Entwicklung Aus Constitutionellen Ursachen, Sowie Zum Darwinismus*.
Berlin: Nicolaische Verlags-Buchhandlung; 1881. https://doi.org/10.5962/bhl.title.49215

28. Kammerer P. *Der Artenwandel Auf Inseln Und Seine Ursachen Ermittelt Durch Vergleich
Und Versuch an Den Eidechsen Der Dalmatinischen Eilande*. Leipzig und Wien: Verlag von
Franz Deuticke; 1926.

29. While GM, Michaelides S, Heathcote RJP, et al. Sexual selection drives asymmetric
introgression in wall lizards. *Ecology Letters*. 2015;18(12):1366-1375.
https://doi.org/10.1111/ele.12531

30. Miñano MR, While GM, Yang W, et al. Climate Shapes the Geographic Distribution and
Introgressive Spread of Color Ornamentation in Common Wall Lizards. *The American
Naturalist*. 2021;198(3):379-393. https://doi.org/10.1086/715186

31. Yang W, While GM, Laakkonen H, et al. Genomic evidence for asymmetric introgression by
sexual selection in the common wall lizard. *Molecular Ecology*. 2018;27(21):4213-4224.
https://doi.org/10.1111/mec.14861

32. Feiner N, Yang W, Bunikis I, While GM, Uller T. Adaptive introgression reveals the genetic
basis of a sexually selected syndrome in wall lizards. *Science Advances*.
2024;10(14):eadk9315. https://doi.org/10.1126/sciadv.adk9315

33. Heathcote RJP, While GM, MacGregor HEA, et al. Male behaviour drives assortative
reproduction during the initial stage of secondary contact. *Journal of Evolutionary Biology*.
2016;29(5):1003-1015. https://doi.org/10.1111/jeb.12840

34. MacGregor HEA, While GM, Barrett J, et al. Experimental contact zones reveal causes and
targets of sexual selection in hybridizing lizards. *Functional Ecology*. 2017;31(3):742-752.
https://doi.org/10.1111/1365-2435.12767

35. Nagase T, Sanai Y, Nakamura S, Asato H, Harii K, Osumi N. Roles of HNK-1 carbohydrate
epitope and its synthetic glucuronyltransferase genes on migration of rat neural crest
cells. *Journal of Anatomy*. 2003;203(1):77-88. https://doi.org/10.1046/j.1469-
7580.2003.00205.x

36. Bronner-Fraser M. Perturbation of cranial neural crest migration by the HNK-1 antibody.
Developmental Biology. 1987;123(2):321-331. https://doi.org/10.1016/0012-
1606(87)90390-3

37. Clark K, Bender G, Murray BP, et al. Evidence for the neural crest origin of turtle plastron
bones. *genesis*. 2001;31(3):111-117. https://doi.org/10.1002/gene.10012

38. Giovannone D, Ortega B, Reyes M, et al. Chicken trunk neural crest migration visualized
with HNK1. *Acta Histochemica*. 2015;117(3):255-266.
https://doi.org/10.1016/j.acthis.2015.03.002

39. Hirata M, Ito K, Tsuneki K. Migration and Colonization Patterns of HNK-1-Immunoreactive
Neural Crest Cells in Lamprey and Swordtail Embryos. *jzoo*. 1997;14(2):305-312.
https://doi.org/10.2108/zsj.14.305

40. Juarez M, Reyes M, Coleman T, et al. Characterization of the trunk neural crest in the
bamboo shark, Chiloscyllium punctatum. *Journal of Comparative Neurology*.
2013;521(14):3303-3320. https://doi.org/10.1002/cne.23351

41. Kalcheim C, Le Douarin NM. Requirement of a neural tube signal for the differentiation of
neural crest cells into dorsal root ganglia. *Developmental Biology*. 1986;116(2):451-466.
https://doi.org/10.1016/0012-1606(86)90146-6

42. Olsson L, Moury JD, Carl TF, Håstad O, Hanken J. Cranial neural crest-cell migration in the
direct-developing frog, Eleutherodactylus coqui: Molecular heterogeneity within and
among migratory streams. *Zoology*. 2002;105(1):3-13. https://doi.org/10.1078/0944-
2006-00051

43. Erickson CA, Loring JF, Lester SM. Migratory pathways of HNK-1-immunoreactive neural
crest cells in the rat embryo. *Developmental Biology*. 1989;134(1):112-118.
https://doi.org/10.1016/0012-1606(89)90082-1

44.   Tucker GC, Delarue M, Zada S, Boucaut JC, Thiery JP. Expression of the HNK-1/NC-1
epitope in early vertebrate neurogenesis. *Cell Tissue Res*. 1988;251(2):457-465.
https://doi.org/10.1007/BF00215855

45.   Abo T, Balch CM. A differentiation antigen of human NK and K cells identified by a
monoclonal antibody (HNK-1). *The Journal of Immunology*. 1981;127(3):1024-1029.
https://doi.org/10.4049/jimmunol.127.3.1024

46.   Simões-Costa M, Bronner ME. Establishing neural crest identity: a gene regulatory recipe.
Development (Cambridge, England). 2015;142(2):242-257.
https://doi.org/10.1242/dev.105445

47.   Qin Q, Xu Y, He T, Qin C, Xu J. Normal and disease-related biological functions of Twist1
and underlying molecular mechanisms. *Cell Res*. 2012;22(1):90-106.
https://doi.org/10.1038/cr.2011.144

48.   Soldatov R, Kaucka M, Kastriti ME, et al. Spatiotemporal structure of cell fate decisions in
murine neural crest. *Science*. 2019;364(6444):eaas9536.
https://doi.org/10.1126/science.aas9536

49.   Howard AG, Baker PA, Ibarra-García-Padilla R, et al. An atlas of neural crest lineages along
the posterior developing zebrafish at single-cell resolution. *eLife*. 2021;10:1-31.
https://doi.org/10.7554/eLife.60005

50.   Hanemaaijer ES, Margaritis T, Sanders K, et al. Single-cell atlas of developing murine
adrenal gland reveals relation of Schwann cell precursor signature to neuroblastoma
phenotype. *PNAS*. 2021;118(5). https://doi.org/10.1073/pnas.2022350118

51.   Kuratani SC, Kirby ML. Initial migration and distribution of the cardiac neural crest in the
avian embryo: An introduction to the concept of the circumpharyngeal crest. *American
Journal of Anatomy*. 1991;191(3):215-227. https://doi.org/10.1002/aja.1001910302

52.   Kvasilova A, Gregorovicova M, Kundrat M, Sedmera D. HNK-1 in Morphological Study of
Development of the Cardiac Conduction System in Selected Groups of Sauropsida. *The
Anatomical Record*. 2019;302(1):69-82. https://doi.org/10.1002/ar.23925

53.   Nakagawa M, Thompson RP, Terracio L, Borg TK. Developmental anatomy of HNK-1
immunoreactivity in the embryonic rat heart: co-distribution with early conduction tissue.
Anat Embryol. 1993;187(5):445-460. https://doi.org/10.1007/BF00174420

54.   Chuck ET, Watanabe M. Differential expression of PSA-NCAM and HNK-1 epitopes in the
developing cardiac conduction system of the chick. *Developmental Dynamics*.
1997;209(2):182-195. https://doi.org/10.1002/(SICI)1097-0177(199706)209:2<182::AID-
AJA4>3.0.CO;2-E

55.   Ikeda T, Iwasaki K, Shimokawa I, Sakai H, Ito H, Matsuo T. Leu-7 immunoreactivity in
687   human and rat embryonic hearts, with special reference to the development of the
688   conduction tissue. *Anat Embryol*. 1990;182(6):553-562.
689   https://doi.org/10.1007/BF00186462

690   56.   Freyer L, Aggarwal V, Morrow BE. Dual embryonic origin of the mammalian otic vesicle
691   forming the inner ear. *Development*. 2011;138(24):5403-5414.
692   https://doi.org/10.1242/dev.069849

693   57.   Lee VM, Sechrist JW, Luetolf S, Bronner-Fraser M. Both neural crest and placode
694   contribute to the ciliary ganglion and oculomotor nerve. *Developmental Biology*.
695   2003;263(2):176-190. https://doi.org/10.1016/j.ydbio.2003.07.004

696   58.   Streeter GL. The development of the cranial and spinal nerves in the occipital region of the
697   human embryo. *American Journal of Anatomy*. 1905;4(1):83-116.
698   https://doi.org/10.1002/aja.1000040106

699   59.   Goldberg S, Venkatesh A, Martinez J, et al. The development of the trunk neural crest in
700   the turtle Trachemys scripta. *Developmental Dynamics*. 2020;249(1):125-140.
701   https://doi.org/10.1002/dvdy.119

702   60.   Canning CA, Lee L, Irving C, Mason I, Jones CM. Sustained interactive Wnt and FGF
703   signaling is required to maintain isthmic identity. *Developmental Biology*. 2007;305(1):276-
704   286. https://doi.org/10.1016/j.ydbio.2007.02.009

705   61.   Nothwang HG, Ebbers L, Schlüter T, Willaredt MA. The emerging framework of mammalian
706   auditory hindbrain development. *Cell Tissue Res*. 2015;361(1):33-48.
707   https://doi.org/10.1007/s00441-014-2110-7

708   62.   Azambuja AP, Simoes-Costa M. A regulatory sub-circuit downstream of Wnt signaling
709   controls developmental transitions in neural crest formation. *PLOS Genetics*.
710   2021;17(1):e1009296. https://doi.org/10.1371/journal.pgen.1009296

711   63.   Yoshida T, Vivatbutsiri P, Morriss-Kay G, Saga Y, Iseki S. Cell lineage in mammalian
712   craniofacial mesenchyme. *Mechanisms of Development*. 2008;125(9):797-808.
713   https://doi.org/10.1016/j.mod.2008.06.007

714   64.   Chai Y, Jiang X, Ito Y, et al. Fate of the mammalian cranial neural crest during tooth and
715   mandibular morphogenesis. *Development*. 2000;127(8):1671-1679.
716   https://doi.org/10.1242/dev.127.8.1671

717   65.   Barrallo-Gimeno A, Nieto MA. The Snail genes as inducers of cell movement and survival:
718   implications in development and cancer. *Development*. 2005;132(14):3151-3161.
719   https://doi.org/10.1242/dev.01907

720   66.   Sefton M, Sanchez S, Nieto MA. Conserved and divergent roles for members of the Snail
721   family of transcription factors in the chick and mouse embryo. *Development*.
722   1998;125(16):3111-3121. https://doi.org/10.1242/dev.125.16.3111

723   67.   Barriga EH, Maxwell PH, Reyes AE, Mayor R. The hypoxia factor Hif-1α controls neural
724   crest chemotaxis and epithelial to mesenchymal transition. *Journal of Cell Biology*.
725   2013;201(5):759-776. https://doi.org/10.1083/jcb.201212100

726   68.   Vincentz JW, Firulli BA, Lin A, Spicer DB, Howard MJ, Firulli AB. Twist1 Controls a Cell-
727   Specification Switch Governing Cell Fate Decisions within the Cardiac Neural Crest. *PLOS*
728   *Genetics*. 2013;9(3):e1003405. https://doi.org/10.1371/journal.pgen.1003405

729   69.   Tavares AT, Izpisuja-Belmonte JC, Rodriguez-Leon J. Developmental expression of chick
730   twist and its regulation during limb patterning. *Int J Dev Biol*. 2004;45(5-6):707-713.
731   https://doi.org/10.1387/ijdb.11669372

732   70.   Liu JAJ, Wu MH, Yan CH, et al. Phosphorylation of Sox9 is required for neural crest
733   delamination and is regulated downstream of BMP and canonical Wnt signaling.
734   *Proceedings of the National Academy of Sciences*. 2013;110(8):2882-2887.
735   https://doi.org/10.1073/pnas.1211747110

736   71.   McKeown SJ, Lee VM, Bronner-Fraser M, Newgreen DF, Farlie PG. Sox10 overexpression
737   induces neural crest-like cells from all dorsoventral levels of the neural tube but inhibits
738   differentiation. *Developmental Dynamics*. 2005;233(2):430-444.
739   https://doi.org/10.1002/dvdy.20341

740   72.   Cheng YC, Cheung M, Abu-Elmagd MM, Orme A, Scotting PJ. Chick Sox10, a transcription
741   factor expressed in both early neural crest cells and central nervous system.
742   *Developmental Brain Research*. 2000;121(2):233-241. https://doi.org/10.1016/S0165-
743   3806(00)00049-3

744   73.   Kos R, Reedy MV, Johnson RL, Erickson CA. The winged-helix transcription factor FoxD3 is
745   important for establishing the neural crest lineage and repressing melanogenesis in avian
746   embryos. *Development*. 2001;128(8):1467-1479. https://doi.org/10.1242/dev.128.8.1467

747   74.   Lukoseviciute M, Gavriouchkina D, Williams RM, et al. From Pioneer to Repressor: Bimodal
748   foxd3 Activity Dynamically Remodels Neural Crest Regulatory Landscape In Vivo.
749   *Developmental Cell*. 2018;47(5):608-628.e6. https://doi.org/10.1016/j.devcel.2018.11.009

750   75.   D'Autréaux F, Coppola E, Hirsch MR, Birchmeier C, Brunet JF. Homeoprotein Phox2b
751   commands a somatic-to-visceral switch in cranial sensory pathways. *Proceedings of the*
752   *National Academy of Sciences*. 2011;108(50):20018-20023.
753   https://doi.org/10.1073/pnas.1110416108

76. Furlan A, Dyachuk V, Kastriti ME, et al. Multipotent peripheral glial cells generate
neuroendocrine cells of the adrenal medulla. *Science*. 2017;357(6346):eaal3753.
https://doi.org/10.1126/science.aal3753

77. Pattyn A, Morin X, Cremer H, Goridis C, Brunet JF. The homeobox gene Phox2b is essential
for the development of autonomic neural crest derivatives. *Nature*. 1999;399(6734):366-
370. https://doi.org/10.1038/20700

78. Patthey C, Clifford H, Haerty W, Ponting CP, Shimeld SM, Begbie J. Identification of
molecular signatures specific for distinct cranial sensory ganglia in the developing chick.
Neural Development. 2016;11(3). https://doi.org/10.1186/s13064-016-0057-y

79. Pattyn A, Morin X, Cremer H, Goridis C, Brunet JF. Expression and interactions of the two
closely related homeobox genes Phox2a and Phox2b during neurogenesis. *Development*.
1997;124(20):4065-4075. https://doi.org/10.1242/dev.124.20.4065

80. Chan WH, Komada M, Fukushima T, Southard-Smith EM, Anderson CR, Wakefield MJ.
RNA-seq of Isolated Chromaffin Cells Highlights the Role of Sex-Linked and Imprinted
Genes in Adrenal Medulla Development. *Sci Rep*. 2019;9(1):3929.
https://doi.org/10.1038/s41598-019-40501-0

81. Langley K, Grant NJ. Molecular markers of sympathoadrenal cells. *Cell Tissue Res*.
1999;298(2):185-206. https://doi.org/10.1007/PL00008810

82. Khannoon T, Alvarado C, De Bellard ME. Early Development of The Trunk Neural Crest
Cells in the Egyptian Cobra Naja H. Haje. Published online May 26, 2021.
https://doi.org/10.21203/rs.3.rs-547365/v1

83. Kundrát M. HNK-1 immunoreactivity during early morphogenesis of the head region in a
nonmodel vertebrate, crocodile embryo. *Naturwissenschaften*. 2008;95(11):1063-1072.
https://doi.org/10.1007/s00114-008-0426-4

84. Atkins N, Jones SM, Guillette LJ. Timing of parturition in two species of viviparous lizard:
influences of β-adrenergic stimulation and temperature upon uterine responses to
arginine vasotocin (AVT). *J Comp Physiol B*. 2006;176(8):783-792.
https://doi.org/10.1007/s00360-006-0100-0

85. Feiner N, Rago A, While GM, Uller T. Signatures of selection in embryonic transcriptomes
of lizards adapting in parallel to cool climate. *Evolution*. 2018;72(1):67-81.
https://doi.org/10.1111/evo.13397

86. Andrade P, Pinho C, Pérez i de Lanuza G, et al. Regulatory changes in pterin and carotenoid
genes underlie balanced color polymorphisms in the wall lizard. *Proceedings of the
National Academy of Sciences*. 2019;116(12):5633-5642.
https://doi.org/10.1073/pnas.1820320116

87. Feiner N. Evolutionary lability in Hox cluster structure and gene expression in Anolis
lizards. *Evolution Letters*. 2019;3(5):474-484. https://doi.org/10.1002/evl3.131

88. Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis Version
11. *Molecular Biology and Evolution*. 2021;38(7):3022-3027.
https://doi.org/10.1093/molbev/msab120

Figure legends

Figure 1. Immunohistochemistry of HNK-1 stains migratory and some early differentiating NCCs in

common wall lizard embryos. Dashed arrows indicate inferred cranial NCC migration. **A-C** 13ss embryo

in lateral (**A**) and dorsal (**B**) view, and in optical cross-section (**C**). Asterisks indicate expression in

premigratory and early migratory NCCs, and open arrows indicate vagal NCCs. Staining in the heart is best

seen in **A** and **C. D-E** Head of 17ss embryo in lateral (**D**) and dorsal (**E**) view. **F-I** Trunk of 17ss embryo

in lateral (**F**) and dorsal (**G**) view, and in optical cross-sections (**H** and **I**). Arrows indicate inter-somitic

NCC migration and asterisks indicate premigratory and early migratory NCCs. **J-L** Head of 22ss embryo

in lateral (**J**) and dorsal (**K**) view, and in optical cross-section (**L**). **M** Trunk of 22ss embryo in lateral view.

Arrow indicates inter-somitic NCC migration and open arrow indicates migration at the level of a somite.

N and **O** Heads of 28 (**N**) and 35ss (**O**) embryos in lateral view. **P** Trunk of 35ss embryo in lateral view. **O**

and **P** were rendered with lower DAPI-opacity. The locations of the optical cross sections **C**, **H**, **I** and **L**

are indicated in **A**, **F** and **J**. **Abbreviations:** E, eye; Fb, forebrain; Fl, Forelimb; G, gut; H, heart; HB,

hindbrain; Mn, mesonephros; MP, maxillary process; NT neural tube; OV, otic vesicle; PA, pharyngeal

arch; Ph, pharynx; S, somite; Va, vagal. **Cranial nerves:** I, olfactory nerve (also known as cranial nerve

1); V, trigeminal nerve (also known as cranial nerve 5); Vmx, maxillary branch of the trigeminal nerve;

Vopth, ophthalmic branch of the trigeminal nerve. **Scale bars** in **B**-**D**, **F**-**L** and **O**-**P** are 0.5 mm, all other

scale bars are 1 mm.

813
Figure 2. *Wnt1* is expressed along the dorsal midline. Asterisks indicate a gap with no expression in the

anterior hindbrain. **A, B** 16ss embryo in lateral (**A**) and dorsolateral (**B**) view. **C** 22ss embryos in lateral

view. **D** Hindbrain of 30ss embryo in dorsolateral view. **E** 30ss embryos in lateral view. **Scale bars** in **B**

and **D** are 0.5 mm, all other scale bars are 1 mm.

Figure 3. *Snai2* is expressed in both premigratory and migratory NCCs. Arrows indicates expression

along the dorsal midline. **A-D** 17ss embryo in lateral (**A**) and dorsal (**B**) view and in cross-sections (**C** and

D) at the levels indicated in **A**. Arrowhead indicates migration through the anterior half of the somites and

dashed arrows indicate the inferred streams of migratory cranial NCCs. **E** 22ss embryo in lateral view.

Abbreviations: Ao, aorta; E, eye; G, gut; H, heart; HB, hind brasin; MP, maxillary process; Nc, notochord;

NT, neural tube; OV, otic vesicle; PA, pharyngeal arch; Ph, pharynx; R, rostrum; Va, vagal NCCs. **Scale**

bar in **B** is 0.5 mm and scale bars in **C** and **D** are 0.1 mm. All other scale bars are 1 mm.

Figure 4. *Twist1* is expressed in migratory cranial NCCs. **A-C** Heads of 13, 17 and 22ss embryos in

lateral view. Dashed arrows indicate the inferred streams of cranial NCC migration. **D** Head of 35ss embryo

in lateral view. Arrow indicates the maxillary process and arrowhead indicates the first pharyngeal arch.

Abbreviations: E, eye; MP, maxillary process; OV, otic vesicle; PA1-3, pharyngeal arch 1-3; R, rostrum;

Va, posterior stream of vagal NCCs. **Scale bar** in **A** is 0.5 mm, all others are 1 mm.

Figure 5. *Snai1* is expressed in migratory cranial NCCs. **A** Head of 16ss embryo in lateral view. Dashed

lines indicate inferred streams of cranial NCC migration. **B** 22ss embryo in lateral view. Arrow indicates a

bifurcation in the inferred caudal stream of vagal NCCs. Expression in the trunk is in tissues of non-NCC

origin. **C** Head of 28ss embryo in lateral view. The dashed arrows indicate the third stream of inferred

cranial NCC migration. **Abbreviations:** E, eye; MP, maxillary process; OV, otic vesicle; PA1-3,

pharyngeal arch 1-3; R, rostrum; Va, posterior stream of vagal NCCs. **Scale bars** are 1 mm.

Figure 6. *Sox9* expression labels both premigratory NCCs and migratory cranial NCCs. A, B Head

(**A**) and trunk (**B**) of 17ss embryo in lateral view. Dashed arrows indicate inferred streams of cranial NCC

migration. Arrows indicate expression in the dorsal midline and open arrow indicates early migratory NCCs. **C** 22ss embryo in lateral view. **D** Trunk of 39ss embryo in lateral view. Arrow indicates expression in the heart. **Abbreviations:** E, eye; MP, maxillary process; OV, otic vesicle; PA, pharyngeal arch; R, rostrum; Va, posterior stream of vagal NCCs. Scale bars in **A** and **B** are 0.5 mm, all others are 1 mm.

Figure 7. *Sox10* expression identifies migratory NCCs and the developing peripheral nervous system. Asterisks indicate premigratory and early migratory NCCs along the dorsal midline, arrows indicate the trigeminal nerve placode, open arrows indicate migratory trunk NCCs and arrowheads indicate caudally migrating vagal NCCs. **A-F** 13ss embryo in lateral (**A, C**) and dorsal (**B**) view and in cross-section (**D-F**). The positions of **C-F** are indicated in **A**. Dashed arrows indicate inferred streams of NCC migration. **D** is a composite of many focal planes, while **E** and **F** are single focal planes. **G** 17ss embryo in lateral view. Notice the weak signal, a result replicated in three separate experiments. **H-I** 22ss embryo in lateral (**H**) and dorsal (**I**) view. **J-K** Head (**J**) and trunk (**K**) of 30ss embryo in lateral view. **L** 37ss embryo in lateral view. Asterisk indicates weak expression along the dorsal midline of the tail. **M-O** 39ss embryo in dorsal (**M**) and lateral (**N**) view, and in cross-section (**O**). The approximate level of the cross-section in **O** is indicated by a dashed line in **N**. **P** 51ss embryo in lateral view. **Abbreviations:** Ao, aorta; CG, ciliary ganglion; DR, dorsal root; DRG, dorsal root ganglion; G, gut; Mn, mesonephros; Nc, notochord; NT, neural tube; OV, otic vesicle; PN, peripheral nerve; SG, sympathetic ganglion; VR, ventral root. **Cranial nerves:** I, olfactory nerve (also known as cranial nerve 1); III, oculomotor nerve (also known as cranial nerve 3); V, trigeminal nerve (also known as cranial nerve 5); Vopth, ophthalmic branch of the trigeminal nerve; Vmd, mandibular branch of the trigeminal nerve; Vmx, maxillary branch of the trigeminal nerve; IX, glossopharyngeal nerve (also known as cranial nerve 9); X, vagus nerve (also known as cranial nerve 10); XI, spinal accessory nerve (also known as cranial nerve 11); XII, hypoglossal nerve (also known as cranial nerve 12). **Scale bars** in **B** and **O** are 0.5 mm and **C**, **D**, **E** and **F** are 0.1 mm. All other scale bars are 1 mm.

Figure 8. *FoxD3* expression labels both premigratory and migratory NCCs as well as the developing

peripheral nervous system. A and **B** 13 and 16ss embryos in lateral view. Dashed arrows indicate streams

of migratory cranial NCCs, arrows indicate expression in premigratory NCCs and open arrows indicate

early migratory NCCs. **C** and **D** 30ss embryo in lateral view. Arrow indicates the trigeminal nerve placode

and open arrow indicates the geniculate ganglion (also known as cranial nerve ganglion 7) The position of

D is indicated in **A**. **Scale bars** are 1 mm.

Figure 9. *Phox2b* expression identifies cranial ganglia. A-C Head of 39ss embryo in lateral (**A**) and

dorsal (**B** and **C**) view. Expression signals in several cranial nerve ganglia are indicated. **Abbreviations:**

CG, ciliary ganglion; III, oculomotor nerve (also known as cranial nerve 3); gVII, geniculate ganglion (also

known as cranial nerve ganglion 7); IX, glossopharyngeal ganglion (also known as cranial nerve 9); X,

vagus nerve (also known as cranial nerve 10); OV, otic vesicle. **Scale bars** are 1 mm.

Figure 10. *Th* expression identifies the chromaffin cells of the putative adrenal gland. A-D 39ss

embryo in lateral (**A, C**) and dorsal (**B**) view and in cross-section (**D**). Dashed rectangle and line in **A**

indicate the approximate position of panels **C** and **D** respectively. **E, F** Trunk of 35 and 51ss embryos in

lateral view. Arrowheads indicate discontinuity in the expression. **Abbreviations:** Ao, aorta; Mn,

mesonephros; Nc, notochord; NT, neural tube; S, somite. **Scale bar** in **D** is 0.5 mm, all others are 1 mm.

Figure 11. Synthesis of the spatiotemporal distribution of NCCs in developing common wall lizards.

A Summary of the inferred distribution and migratory streams of NCCs in four developmental stages of the

common wall lizards. Different groups of NCCs are color-coded according to their state or expected fate.

B Schematic overview of the state of NCCs (premigratory, migratory or differentiating) and their respective

domain (cranial, vagal or trunk) with respect to their expression of individual marker genes and the presence

of HNK-1 positive staining. **Abbreviations**: Fb, forebrain; Fl, forelimb; Hb, hindbrain; Hl, hindlimb; Mb,

midbrain.

Figure 12 Gene tree of *Snai1*, *-2* and *-3*. Orthology relations of the transcription factors *Snai1* and *-2* in

the common wall lizard was confirmed with regard to human, mouse and chicken. The *Snai3* gene appears

to be absent from the common wall lizard genome. Accession numbers from Ensemble (wall lizard) and

NCBI (mouse, human and chicken) are given next to each branch tip.

Tables

Table 1 Sequences of primers used in this study (5' to 3').

Gene	Forward primer	Reverse primer
Snai2	ATAAGCAGTTGCACTGTG	GTGTGTATCAAGCATGTG
Eya2	GATGGTGTTGAAGAGGAA	AGGTAACTCTTTCTCCTG
FoxD3	AAAGCCTTCAACTCCCAG	CTCCCAATCCTTTGCTTG
Mitf	ACCATCACTTTCAACGAC	TTAGAAGGTTGTCCACTC
Snai1	AACTGTGAGAAGGAGTAC	CAAGAGAGAATTGAGTGG
Sox9	AGTTTGACCAGTATCTCC	TAAGGCATGCCATTTTGC
Sox10	AAAGACAGAACTGCAGTC	GAGGGTCGTATATACAG
Th	ATAATATCCCTCAGCTGG	TGGATAGCAACACAAACC
Twist1	TACATCGACTTCCTCTAC	AAATGAAAGAGGAGCAGG
Wnt1	CTCTGCAAGAATAGGAAC	TCATCTACGGCAACAAAG

902 **Reagent list**

Reagent	Company	Catalog#	Cas#
PBS	Sigma	P4417	

Water, nucl. free, Mol. Biol. Grade, Ultrapure	ThermoScientific	J71786.K8	
Paraformaldehyde	Sigma-Aldrich	P6148	30525-89-4
Methanol	Merck Millipore	1.06009.1000	67-56-1
RNAlater	Invitrogen	AM7020	
Tween-20	Sigma	P9416	
Tris Base for molecular biology white crystals or powder	Fisher Scientific	BP152	77-86-1
Sodium Chloride, molecular biology grade (NaCl)	MP	194848	7647-14-5
Bovine Serum Albumin (BSA)	Sigma	A9418	
Primary antibody: CD57 Monoclonal Antibody (TB01 (TBO1)), FITC	eBioscience	11-0577-42	
Secondary antibody: Alexa Fluor 488 goat anti-mouse IgM (u chain)	Invitrogen	A21042	
DAPI	ThermoScientific	62248	
RapiClear 1.49	SUNJin Lab	RC149001	
Cover Glass 1.5 (~0.17μm)	VWR	631-0134	
iSpacer	SUNJin Lab		
Rneasy Mini Kit	Qiagen	74104	
SupersScript III Reverse Transcriptase	Invitrogen	18080	
GoTaq G2 Hot Start Green Master Mix	Promega	M742A	
MinElute PCR Purification Kit	Qiagen	28004	
DIG-labeled dNTPs	Roche	1.16E+10	
Proteinase K, recombinant PCR Grade	Roche	3.12E+09	
20x SSC	Fisher BioReagents	BP13254	
Formamide deionized for mol. Biology	PanReac AppliChem	A2156	1975-12-07

Heparin sodium salt from porcine intestinal mucosa	Sigma-Aldrich	H4784	9041-08-01
Riobonucleic acid from torula yeast	Sigma	R6625	63231-63-0
Blocking Reagent	Roche	1.11E+10	
EDTA	Invitrogen	AM9260G	
Anti-Digoxenin-AP Fab fragments	Roche	1.11E+10	
Maleic acid	Sigma-Aldrich	M0375	110-16-7
NBT/BCIP	Roche	1.17E+10	
Glycerol	VWR	24397.3	56-81-5

	Premigratory			Migratory			Differentiating		
	Cranial	Vagal	Trunk	Cranial	Vagal	Trunk	Cranial	Vagal	Trunk
HNK1	■	■	■	■	■	■	■■		■
Wnt1	■	■	■						
Snai2	■	■	■	■	■	■	■		
Snai1				■	■		■		
Twist1				■	■		■		
Sox9	■	■	■	■	■		■	■	
FoxD3	■	■	■	■	■	■	■	■	■
Sox10				■	■	■	■	■	■
Phox2b							■	■	■
Th									■