The second messenger signaling molecule cyclic di-AMP drives developmental cycle progression in *Chlamydia trachomatis*

Junghoon Lee[1] and Scot P. Ouellette[1*]

[1]Department of Pathology, Microbiology, and Immunology, College of Medicine, University of

Nebraska Medical Center, Omaha, NE, USA.

Running Head: Cyclic di-AMP drives secondary differentiation in *Chlamydia*

*Corresponding author:

Scot P. Ouellette

Department of Pathology, Microbiology, and Immunology

University of Nebraska Medical Center

985900 Nebraska Medical Center, DRC2 5029

Omaha, NE 68198

Phone: 402-559-0763

E-mail: scot.ouellette@unmc.edu

Abstract

The obligate intracellular bacterium *Chlamydia* alternates between two functional forms during its developmental cycle: elementary body (EB) and reticulate body (RB). However, the molecular mechanisms governing the transitions between these forms are unknown. Here, we present evidence cyclic di-AMP (c-di-AMP) is a key factor in triggering the transition from RB to EB (i.e., secondary differentiation) in the chlamydial developmental cycle. By overexpressing or knocking down expression of c-di-AMP synthase genes, we made strains producing different levels of c-di-AMP, which we linked to changes in secondary differentiation status. Increases in c-di-AMP resulted in an earlier increase in transcription of EB-associated genes, and this was further manifested in earlier production of EBs. In contrast, when c-di-AMP levels were decreased, EB production was reduced. Based on these data, we conclude there is a threshold level of c-di-AMP needed to trigger secondary differentiation in *Chlamydia*. This study identifies a mechanism by which secondary differentiation is initiated in *Chlamydia* and reveals a critical role for the second messenger signaling molecule c-di-AMP in this process.

Word count: 169

Keywords: Chlamydia, cyclic di-AMP, c-di-AMP, cyclic dinucleotides, differentiation, development, second messenger, signaling

Introduction

  Chlamydia species are major pathogens of humans and animals. These obligate intracellular bacteria share one key feature: their unique developmental cycle (see ref. (1)). During this cycle, *Chlamydia* transitions between two different functional and morphological forms: the elementary body (EB), an infectious but non-dividing cell, and the reticulate body (RB), a dividing but non-infectious cell (Fig. 1A). A third form, the intermediate body (IB) is a transitional form from the RB to EB. Besides these characteristics, EBs and RBs differ in other ways. For example, EBs are small (~0.3 μm), have a highly disulfide-crosslinked outer membrane (2), and have DNA condensed by histone-like proteins (3). In contrast, RBs are larger (~1 μm), have a Gram-negative cell envelope that lacks peptidoglycan (4-6), and have a dispersed chromosome. RBs divide by an asymmetric polarized division mechanism dependent on MreB-directed peptidoglycan synthesis specifically at the septum (7-10). Not surprisingly, *Chlamydia* expresses genes in a temporally defined manner that corresponds broadly with its developmental cycle (11, 12). "Early" genes (e.g., *euo*) are expressed immediately upon entry into a target host cell and are likely involved in establishing the intracellular niche of *Chlamydia*, the inclusion, and mediating primary differentiation from EB to RB. "Mid" cycle genes (e.g., *mreB*, *clpPX*) facilitate RB replication and division and inclusion growth. "Late" genes (e.g., *hctA*, *omcB*) are expressed when secondary differentiation is initiated to trigger EB formation. Although developmental gene expression has been characterized for decades, the signals or events that initiate differentiation from one form to the other are not known.

  In a 2013 study, another defining characteristic of EBs and RBs was identified: their relative levels of the second messenger signaling molecule cyclic di-AMP (c-di-AMP) (see ref. (13)). Barker et al. studied how IFNβ production is activated in cells infected with *C. trachomatis*. The authors identified a role for the innate immune response protein, STING, which recognizes c-di-AMP. *Chlamydia* encodes a diadenylate cyclase enzyme, DacA, associated with c-di-AMP production, that had not been characterized. As part of their study, the authors determined that c-di-AMP accumulates over the course of the chlamydial developmental cycle, that EBs have high levels of this molecule whereas RBs have low levels, and that DacA is a diadenylate cyclase. It is unlikely that *Chlamydia* produces c-di-AMP only to signal host immune responses. Rather, the parsimonious interpretation is that *Chlamydia* uses this signaling molecule to regulate some aspect of its physiology and that activation of host signaling is "accidental" – similar to activation of NOD2 by chlamydial peptidoglycan (14, 15). However, no direct function of c-di-AMP in chlamydial biology has been defined.

  Diverse functions of c-di-AMP have been reported in Gram-positive bacteria. For example, c-di-AMP is implicated in the response to changes in osmotic pressure. When the extracellular solute level is high, bacteria prevent dehydration by importing both extracellular solute molecules and cations (16). In these mechanisms, c-di-AMP binds to proteins associated with K^+ uptake systems such as Ktr/Trk, KimA, Kup, and Kdp and inhibits their activities in *Bacillus subtilis*, *Staphylococcus aureus*, and *Lactococcus lactis* (17-20). Similarly, K^+ export mechanisms are also regulated by c-di-AMP. In *S. aureus*, the cation/proton antiporter A (CpaA) is activated by binding c-di-AMP (21). Moreover, c-di-AMP binds to the riboswitch upstream of the genes encoding the K^+ transporters and controls their transcriptional levels (22). With these mechanisms, osmotic stress is controlled by c-di-AMP. Of note, *Chlamydia* lacks annotated orthologs of K^+ transporters. In addition to osmotic homeostasis, c-di-AMP

has been reported to affect DNA replication and sporulation (23). When DNA damage is detected, DisA, a diadenylate cyclase, forms a DNA repair complex with RecA and RadA, resulting in inhibition of diadenylate cyclase activity in *B. subtilis*. Subsequently, c-di-AMP levels decrease, and DNA replication and sporulation are arrested (23-26).

Given the differences in c-di-AMP levels between EBs and RBs and its function as a diffusible second messenger signaling molecule, we hypothesized that the accumulation of c-di-AMP during the developmental cycle might be a trigger for secondary differentiation in *Chlamydia* (Fig. 1B). Once a threshold concentration of c-di-AMP has been reached in a given RB, it will begin the differentiation process to an EB. Importantly, we do not propose that c-di-AMP is necessarily required for RB replication or growth. We predicted that, if our hypothesis were correct, then we should be able to alter the levels of c-di-AMP in the organism and affect its differentiation kinetics accordingly. For example, if we increase c-di-AMP production, then we anticipate prematurely triggering RB-to-EB conversion with a concomitant reduction in overall growth and replication (since only RBs divide). Conversely, if we prevent c-di-AMP production, then we anticipate delaying EB production without impacting growth rate (i.e., normal replication with reduced RB-to-EB conversion).

There are three principal mechanisms to regulate c-di-AMP levels: through synthesis by diadenylate cyclase, through degradation by a phosphodiesterase (PDE), and through secretion by a transporter (27). Interestingly, *C. trachomatis* only encodes the synthesis mechanism as it possesses the genes for diadenylate cyclase (*dacA*) and its regulator (*ybbR*) within a bicistronic operon, but no annotated PDEs or c-di-AMP transporters (28). Both DacA and YbbR are predicted to contain transmembrane domains (3 for DacA and 1 for YbbR) (Fig. S1, (13)). To test our hypothesis, we genetically manipulated the levels of DacA and/or YbbR using recently developed strategies in the field. In the present study, we characterized the growth and developmental cycle state of chlamydial strains producing high or low levels of c-di-AMP. Our data show that higher levels of c-di-AMP are directly linked to increased transcript levels of late genes associated with secondary differentiation as well as the concomitant production of EBs at an earlier stage in the developmental cycle. In contrast, in cells with reduced c-di-AMP levels, chlamydial growth was impaired and the developmental cycle was significantly delayed. Based on these data, we conclude that there is a threshold level of c-di-AMP necessary to trigger secondary differentiation in *C. trachomatis*. This is the first study to identify a physiological function for c-di-AMP in *Chlamydia* as well as a signaling mechanism by which secondary differentiation is initiated in these unique bacteria.

Results

Cyclic di-AMP levels increase at later stages of the developmental cycle and in bacteria that overexpress both DacA and YbbR.

To verify that EBs have higher levels of c-di-AMP compared to RBs, we infected HeLa cells at a multiplicity of infection (MOI) of 1 with a transformant of *C. trachomatis* L2 carrying a shuttle plasmid with inducible mCherry. This strain serves as a control for subsequent overexpression experiments, and we induced mCherry expression with 5 nM aTc at 10 hours post-infection (hpi). We collected uninfected and infected HeLa cell lysates at 16 and 24 hpi and measured c-di-AMP levels by ELISA (Fig. 1C). As expected, only basal amounts (<100 pg/mL) of c-di-AMP were detected in

uninfected HeLa cells at either timepoint (Fig. 1C). In infected cells, the 16hpi timepoint is characterized by predominantly an RB population whereas the 24 hpi timepoint is characterized by ongoing secondary differentiation and a mixture of RBs, IBs, and EBs. We observed that c-di-AMP levels were significantly higher at 24hpi (~1,700 pg/mL) compared to 16hpi (~250 pg/mL) and increased approximately 7-fold over this timeframe (Fig. 1C). These results are in agreement with the Barker et al. study indicating higher levels of c-di-AMP in EBs (13).

To test the link between c-di-AMP levels and chlamydial developmental cycle progression, we made a collection of *C. trachomatis* strains carrying anhydrotetracycline (aTc)-inducible constructs (Fig. 1D). These included strains to overexpress a wild-type or a catalytically dead (D164N) DacA isoform (14), wild-type or mutant isoforms of DacA lacking transmembrane domains (ΔTM), YbbR_6xH, or both wild-type or mutant DacA and YbbR_6xH (*dacA*op or *dacA*opMut). For all of these overexpression strains, we emphasize that the chromosomal expression of wild-type *dacA* and *ybbR* is maintained. In addition to the overexpression constructs, we also made a conditional knockdown construct for *dacA* (*dacA*-KD), targeting its promoter region, to decrease the expression of *dacA-ybbR* using a dCas12/crRNA-based system that our group developed for *Chlamydia* (29). Finally, we made a complementation construct for *dacA*-KD by introducing *dacA-ybbR*_6xH (*dacA*-KDcom) 3' to dCas12 such that induction of dCas12 results in the coexpression of DacA and YbbR_6xH during knockdown of endogenous *dacA-ybbR* transcripts.

To assess whether we could alter c-di-AMP levels in our various strains, we first measured c-di-AMP levels from infected cell lysates at 16 and 24hpi after inducing overexpression or knockdown of the target genes at 10 hpi (Fig. 1E). Overexpressing DacA resulted in ~4 fold increase in c-di-AMP levels as compared to the mCherry-expressing strain at 16hpi (~1,000 pg/mL) and ~3-fold increase at 24 hpi (~5,000 pg/mL). Overexpressing YbbR_6xH did not impact c-di-AMP production. When overexpressing the catalytically inactive mutant of DacA(D164N), c-di-AMP levels were ~2-fold lower at 16 hpi (~115 pg/mL) and, at 24 hpi (~320 pg/mL), was similar to the 16hpi mCherry control, indicating a severe reduction in c-di-AMP accumulation. Overexpressing ΔTMDacA phenocopied the effect of overexpressing full-length DacA on c-di-AMP production. Interestingly, the negative effects of overexpressing full-length DacA(D164N) associated with reduced c-di-AMP levels were lost when expressing ΔTMDacA(D164N), which phenocopied the mCherry-expressing control. When DacA and YbbR_6xH were co-overexpressed, c-di-AMP levels increased by approximately 30-fold at 16 hpi (~8,000 pg/mL) and 120-fold at 24 hpi (~340,000 pg/mL) compared to that of the mCherry-expressing control strain. Of note, the levels of c-di-AMP at 16hpi in the *dacA*op strain (~8,000 pg/mL) were higher even than the control strain at 24 hpi (~1,700 pg/mL). To confirm whether DacA enzyme activity is critical for the high levels of c-di-AMP in the *dacA*op strain, we co-overexpressed DacA(D164N) and YbbR_6xH (*dacA*opMut). Here, c-di-AMP levels were reduced compared to the control at 16 (~100 pg/mL) and 24 hpi (~400 pg/mL), suggesting that c-di-AMP synthase activity was blocked in this strain similar to overexpressing the DacA(D164N) alone. For the *dacA*-KD strain, the c-di-AMP level was unchanged at 16 hpi and reduced a statistically significant ~5-fold at 24 hpi (~500 pg/mL; roughly twice the level of the 16h mCherry-expressing control). The loss of c-di-AMP production in this strain was restored in the complemented *dacA*-KD_*dacA*op strain, which showed a phenotype similar to the *dacA*op overexpression strain with c-di-AMP levels above the 24hpi mCherry-expressing strain (~3,700 pg/mL at 16hpi; ~10,000 pg/mL at 24hpi). Based on these data, we conclude that both DacA and YbbR

164 are necessary for optimal c-di-AMP synthesis in *Chlamydia*. Importantly, our collection of strains that
165 are high or low producers of c-di-AMP give us an opportunity to test our overarching hypothesis (Fig.
166 1B).

Overexpression of membrane-localized DacA isoforms, but not YbbR, disrupts chlamydial growth
and development.

To begin exploring the impact of altering *dacA* and/or *ybbR* expression on chlamydial growth,
we performed a series of experiments to assess their protein localization in chlamydiae. We also
measured impacts of overexpressing each individually on chlamydial growth and development. To
observe the localization of DacA and YbbR, we infected HeLa cells with transformants encoding *dacA*
or *ybbR_6xH* alone and induced expression of the constructs at 10 hpi with 5 nM aTc. At 24 hpi,
infected cells were fixed, and we performed an indirect immunofluorescence assay (IFA) by labeling the
chlamydial major outer membrane protein (MOMP) and DacA or 6xH. We observed that both DacA and
YbbR localized at the bacterial membrane as expected (Fig. 2A&S2). Interestingly, when DacA was
overexpressed, the inclusion size was significantly reduced in area (~6 fold) with larger individual
organisms (~2 fold) than those in the uninduced control (Fig. S3A and B). Similarly, overexpressing the
inactive DacA(D164N) mutant resulted in smaller inclusions similar to overexpressing wild-type DacA
(Fig. 2B). We did not quantify effects of YbbR_6xH overexpression on inclusion or bacterial size since
the measured phenotypes indicated no differences from the uninduced control (see below; Fig. S4A).

To investigate the effects of DacA isoforms or YbbR_6xH overexpression on chlamydial
growth, we measured EB progeny production using an inclusion forming unit (IFU) assay and genomic
DNA copy number (a proxy for total bacteria, i.e., EB+RB) by qPCR. To quantify IFUs, a lysate from a
primary infection is prepared and used to infect a fresh monolayer of cells. Any inclusions in the
secondary infection are derived from viable EBs present in the primary infection. The vector control
strain showed no differences in IFUs when overexpressing mCherry (Fig. S5). At 24 hpi, both IFUs and
genome copy numbers were significantly decreased in the DacA overexpression strain (Fig. 2C and D).
We obtained similar results when overexpressing wild-type DacA with a 6xHis tag, indicating that a C-
terminal tag does not impact these phenotypes in *Chlamydia* (Fig. S6). We also performed the same
experiments with the inactive isoform of DacA, DacA(D164N). Like overexpression of the wild-type
DacA, overexpression of DacA(D164N) also negatively affected IFU production and resulted in reduced
genome copy numbers even though c-di-AMP levels were reduced under these conditions (Fig. 2E and
F). YbbR_6xH overexpression did result in a statistically significant, ~2-fold decrease in IFU production
(Fig. S4B). This reflects one division cycle difference from the uninduced control and, in the absence of
any other phenotypic discrepancy, is not considered biologically relevant. No differences were noted in
genome copy number when overexpressing YbbR_6xH (Fig. S4C).

To further investigate impacts of overexpression of these proteins on developmental cycle
progression, we quantified transcripts by RT-qPCR for a canonical early-cycle gene, *euo*, and two late-
cycle genes, *hctA* (an "early" late gene (30)) and *omcB* (a canonical late gene (11)). Consistent with the
genome copy numbers and IFU data, overexpression of DacA resulted in elevated *euo* transcripts and a
reduction in the amounts of the late gene transcripts at 24hpi (Fig. 2G). Similar effects on these

transcripts were noted when overexpressing DacA(D164N) (Fig. 2H) We measured no effect of
YbbR_6xH overexpression on transcript levels for *euo*, *hctA*, or *omcB* (Fig. S4D).

Given that overexpression of either the wild-type or mutant isoform of DacA gave the same
phenotype yet yielded differences in c-di-AMP levels, we next explored the need for DacA or
DacA(D164N) to be membrane-localized to effect these changes. Therefore, we evaluated the effects of
overexpression of wild-type or mutant DacA lacking its transmembrane domains (ΔTM). As noted in
Figure 1, overexpression of wild-type ΔTM DacA yielded the same level of c-di-AMP as overexpression
of the full-length wild-type DacA. In contrast, c-di-AMP levels measured from chlamydiae
overexpressing the mutant ΔTM DacA(D164N) were the same as the mCherry-expressing strain but
reduced in chlamydiae overexpressing the full-length DacA(D164N). By IFA, both ΔTM isoforms
localized to the cytosol, and there were no observable differences in inclusion or bacterial morphology
as compared to the uninduced control (Fig. 3A and B). We did not quantify effects of overexpression on
inclusion or bacterial size since the measured phenotypes indicated no differences from the uninduced
control. Overexpression of the ΔTM isoforms resulted in no statistical differences in IFU production or
genome copy numbers, and, similarly, there were no differences in transcript levels for *ybbR, euo, hctA,*
or *omcB* as compared to the uninduced controls (Fig. 3C-H). These data indicate that the negative
impacts of DacA overexpression are linked to its membrane localization and are independent of c-di-
AMP production.

A low level of c-di-AMP decreases the transcript levels of late genes.

As mentioned above, the *dacA*-KD strain displayed a lower c-di-AMP level compared to that of
the control strain at 24 hpi whereas the *dacA*-KDcom complemented strain exhibited high levels of c-di-
AMP at this timepoint (Fig. 1E). The complemented strain encodes not only the *dacA*-knockdown
system but also the *dacA-ybbR_6xH* operon as a transcriptional fusion with dCas12. To observe effects
of knockdown or complementation on the developmental cycle, we first examined inclusion and
bacterial morphology by IFA. When the knockdown system was induced, bacterial cell size was
enlarged (~2 fold), but we observed no change in inclusion area as compared to that of the uninduced
control (Fig. 3A, S7A&B). In the complemented strain, induction of dCas12 and YbbR_6xH was
confirmed by IFA under inducing conditions (Fig. 4B). We expected that co-expressing DacA and
YbbR_6xH would complement the phenotype of *dacA*-KD. However, the inclusion area was reduced
(~3 fold) with slightly larger organisms (~1.4 fold) as compared to the uninduced control (Fig. S7C&D).

In addition, we also measured the amount of EB progeny (IFUs) and genome copy numbers.
Although genome copy numbers were the same, IFUs decreased by approximately 80% compared to
that of the uninduced sample at 24 hpi in the knockdown strain (Fig. 4C&D). IFUs and genome copies
showed an approximate 2-fold reduction at 24 hpi in the complemented strain (Fig. 4E&F). The data for
the knockdown strain suggest that low levels of c-di-AMP are detrimental for secondary differentiation.
To further explore this, we performed RT-qPCR to quantify transcripts of relevant gene targets. When
dCas12 was induced, both *dacA* and *ybbR* transcript levels decreased (Fig. 4G). Since *dacA* and *ybbR*
are transcribed in an operon, this result is not surprising. Consistent with reduced EB yields, transcripts
of the late genes *omcB* and *hctA* were decreased at 24 hpi. In contrast, transcripts of the early gene *euo*

were slightly elevated at 24 hpi. We have previously observed no effects on genome levels or
transcription of these genes when overexpressing the dCas12 gene alone (29), indicating these effects
are specific to *dacA*-KD. For the complemented strain, both *dacA* and *ybbR* transcripts were increased
compared to the *dacA*-KD strain (Fig. 4H), and transcript levels for these genes were increased beyond
the "wild-type" uninduced control levels. Nonetheless, transcripts for the early gene *euo* were
indistinguishable to levels measured in the uninduced strain (Fig. 4H). Surprisingly, *hctA* transcripts
were increased ~10-fold at 14 hpi compared to that of the uninduced sample, and *omcB* transcripts were
slightly, but not significantly, increased at 14 and 24 hpi (Fig. 4H). These data suggest that
overexpressing DacA and YbbR_6xH may alter the timing of secondary differentiation.

High levels of c-di-AMP induce late gene expression.

To clarify the effect of DacA and YbbR_6xH overexpression on secondary differentiation, we
next evaluated the phenotype of the *dacA*op strain in contrast with the *dacA*opMut strain, in which the
active site residue of DacA has been mutated. When we induced expression of the wild-type constructs,
we confirmed the induction of DacA and YbbR_6xH and their colocalization at the membrane (Fig. 5A;
Pearson correlation coefficient of 0.713 ± 0.109 from 20 inclusions measured by JACoP Plugin of
ImageJ; values near 1 indicate colocalization (31, 32)). This is not surprising as both proteins are critical
for c-di-AMP synthesis based on our c-di-AMP measurements (Fig. 1E). Organism and inclusion
morphology were similar to the *dacA*-KDcom complemented strain (Fig. 4B&5A; Fig. S8A&B). The
bacterial morphology of the *dacA*opMut strain indicated larger organisms (~1.8 fold) in smaller
inclusions (~2.5 fold) after inducing expression (Fig. 5B; S8C&D). We next assessed whether the
overexpressed DacA and YbbR_6xH affected IFU production and replication. IFUs were reduced
roughly 2-fold for the *dacA*op strain and 1,000-fold in the *dacA*opMut strain after inducing
overexpression (Fig. 5C&E). Again, the genome copy data for *dacA*op overexpression closely
phenocopied the complemented knockdown strain, showing a decrease at 24 hpi (Fig. 5D). Similarly,
genome copy numbers for the *dacA*opMut strain showed a significant drop at 24 hpi after inducing
expression (Fig. 5F).

We then quantified transcripts for the developmentally regulated genes *euo*, *hctA*, and *omcB* as
well as for *dacA* and *ybbR* (Fig. 5G&H). Not surprisingly, *dacA* and *ybbR* transcripts were elevated at
the timepoints assessed under inducing conditions for both the *dacA*op and *dacA*opMut strains.
Transcripts for *euo* were not statistically changed but trended higher at the 24 hpi timepoint whereas
omcB transcripts were slightly, but not significantly, increased at 14 and 24 hpi during *dacA*op
overexpression. Once again, we observed that *hctA* transcripts were increased over 10-fold at 14hpi in
the developmental cycle (Fig. 5G), similar to what we measured for the complemented knockdown
strain (Fig. 4H). These data reinforce that elevated c-di-AMP levels (Fig. 1E) in these strains lead to
increased expression of the late gene *hctA* at an earlier timepoint (14 hpi) in the developmental cycle. As
further validation of this *dacA*op overexpression strain that was constructed in a beta-lactamase
producing background, we also generated a spectinomycin-resistant *dacA*op overexpression strain and
validated that induction of the *dacA* operon resulted in earlier accumulation of *hctA* transcripts (Fig. S9).
In assessing developmentally-regulated transcripts for the *dacA*opMut strain, *euo* levels were maintained,
albeit not significantly so, at 24 hpi (Fig. 5H). In contrast to the c-di-AMP overproducing strains,

transcripts for *hctA* and *omcB* were decreased at 24 hpi (Fig. 5H). The transcriptional results of the
developmentally regulated genes in the *dacA*opMut strain were very similar to the *dacA*-KD strain (Fig.
4G), suggesting that blocking c-di-AMP accumulation interferes with developmental cycle progression.

Elevated c-di-AMP levels result in increased transcript levels of genes necessary for secondary
differentiation.

Given the surprising finding that, in strains overproducing c-di-AMP, *hctA* transcripts were 10-
fold higher at a timepoint not associated with secondary differentiation, we asked the question whether
all genes related to secondary differentiation were increased after inducing production of c-di-AMP.
Conversely, we wanted to explore whether reducing c-di-AMP levels would delay expression of genes
related to secondary differentiation. Thus, to further investigate how c-di-AMP affects transcription of
such genes, we performed RNA sequencing on both the *dacA*op overexpression and *dacA*-KD strains
and compared the transcriptome between uninduced and induced samples within the given strain at the
given timepoint. HeLa cells were infected with these transformants, and overexpression or knockdown
was induced or not at 10 hpi with 5 nM aTc. For the *dacA*op strain, RNA was collected at 16 hpi, a time
at which late genes are beginning to be expressed (as opposed to 14hpi) but remain near a basal level of
transcription(11). The rationale for this was to determine whether high c-di-AMP levels result in
increased late gene transcripts at this timepoint *above and beyond* the levels of the control, uninduced
condition. For the *dacA*-KD strain, RNA was collected at 24 hpi, a time at which late genes are peaking
in their expression. The rationale for this was to determine whether late gene transcription was decreased,
which could not otherwise be assessed at the 16 hpi timepoint.

RNA sequencing results were statistically analyzed by the UNMC Bioinformatics Core (Table
S2&3). Figure 6 shows a volcano plot of the results for the *dacA*op overexpression and *dacA*-KD strains.
Of note, many late genes were evident in the upregulated quadrant for the *dacAop* strain whereas these
genes were present in the downregulated quadrant for the *dacA*-KD strain. We further characterized the
upregulated or downregulated gene sets for the *dacA*op overexpression and *dacA*-KD strains,
respectively, based on significant difference ($p<0.05$) and fold-change (>1.5). A summary of these
results is presented in Table 1 (all data are presented in Table S3). We grouped the differentially
expressed genes into five categories: 1) canonical late genes for which the literature has associated them
with EB function, 2) outer membrane associated, 3) gene regulation associated, 4) glycogen synthesis
associated, and 5) type III secretion system associated. Recent work from our group and the Hefty group
to define the regulons of the alternative sigma factors in *Chlamydia* demonstrated that these sigma
factors regulate some late gene expression associated with outer membrane remodeling, type III
secretion, and other processes (33, 34). Consistent with our RT-qPCR data, we observed that all
canonical late genes as well as all the other genes listed in these categories showed an increase in
expression after c-di-AMP production was induced. In contrast, all the genes, and particularly the
canonical late genes, showed a decrease in expression under conditions where c-di-AMP production was
impaired. Overall, these RNA sequencing data confirm the direct influence of c-di-AMP on expression
of genes related to secondary differentiation.

Alterations in c-di-AMP levels impact the timing of EB production

The phenotypic effect of increased late gene transcription should be an increase in EB
production. However, if EB production is initiated at an earlier time in the developmental cycle when
there are fewer non-infectious RBs to convert, then overall EB production should be decreased with a
concomitant decrease in genome copies since only the RB replicates DNA. Conversely, delayed
expression of late genes should be associated with a delay in EB production. Therefore, to test these
predictions, we quantified EB production at 2h intervals from 18 to 24hpi to assess EB production
during earlier phases of the developmental cycle, as well as at 32 and 48hpi to assess overall EB yields
(Fig. 7). We did not detect any EBs at 16hpi or earlier (not shown). Cells were infected with the *dacA*op,
*dacA*opMut, and *dacA*-KD strains and induced or not at 10 hpi with 5 nM aTc as previously noted.
Consistent with the transcriptional data, we measured higher EB yields at 18 and 20 hpi during *dacA*op
overexpression that quickly plateaued by 24 hpi (Fig. 7A). Conversely, when c-di-AMP accumulation
was blocked by reducing the expression of DacA and YbbR (*dacA*-KD), we observed delayed EB
production (Fig. 7B). We also measured lower EB yields when c-di-AMP levels were decreased by
overexpression of the DacA(D164N) isoform and YbbR (*dacA*opMut; Fig. 7C). The uninduced control
conditions for each strain showed a steady accumulation of EBs throughout the course of the experiment,
as expected. The vector control strain expressing mCherry showed no differences as previously noted
(Fig. S5). From these data, we conclude that earlier production of c-di-AMP levels results in earlier
production of EBs.

Discussion

Secondary differentiation is essential for the propagation and survival of *Chlamydia* species.
However, the mechanisms governing this essential process are largely unknown. It is possible that both
internal and external environmental changes might serve as signals to trigger the shift from the non-
infectious RB to the infectious EB. It is also possible, and even likely, that *Chlamydia* integrates
multiple signal inputs during the differentiation process. For example, we recently described two post-
translational processes that impact secondary differentiation: the activity of the unfoldases ClpX and
ClpC(35, 36) and the redox status of *Chlamydia*(37). As secondary differentiation is asynchronous, any
mechanism must account for the stochasticity within the population of RBs "considering" differentiating
to EBs. Here, we provide the first evidence of a signaling molecule that can directly activate EB-
associated gene expression with a concomitant early production of EBs.

Several reports have proposed mechanisms by which *Chlamydia* triggers secondary
differentiation. For example, Thompson et al. investigated the non-canonical regulation of the major
sigma factor, σ^{66}, by the Rsb phosphoregulatory system(38). The authors provided evidence that altering
the levels of RsbW or RsbV1 impacted the expression of σ^{66}-controlled genes. This led them to propose
a model whereby sensing of ATP by the Rsb system controls the availability of σ^{66}, implying that low
levels of ATP lead to initiation of secondary differentiation. Further support from this came from work
by Kuwabara et al. who showed effects of ATP, GTP, and glucose on the activity of different Rsb
components(39). Finally, work from Soules et al. identified TCA intermediates as ligands for RsbU,
linking the TCA cycle and ATP synthesis to the Rsb system(40). However, no effect was noted on
premature late gene expression in these experimental systems.

Sequestration/inactivation of the major sigma factor under low ATP conditions would
presumably render RNA polymerase free to interact with the alternative sigma factors, σ^{54} and σ^{28}.

These alternative sigma factors have been linked to late gene expression with studies from Soules et al. and Hatch and Ouellette indicating that σ^{54} is associated with increased transcription of outer membrane components, type III secretion (T3S) system components, and other genes typically expressed late in development (33, 34). However, some of these changes are likely indirect and downstream to the σ^{54} regulon itself since some affected genes have been shown to have σ^{66} promoter sequence elements (41, 42). Significant remodeling of the outer membrane occurs as the RB transitions to the EB (43, 44), and, similarly, the EB is prepackaged with T3S effectors, like TarP (45), that facilitate EB invasion into a target host cell. σ^{28} was shown to control expression of two canonical late genes: *hctB* and *tsp* (33). Each of the encoded proteins is highly toxic if overexpressed (46, 47), suggesting that the additional layer of regulation by σ^{28} is necessary to ensure EBs are formed at the correct time. Hatch and Ouellette proposed that these genes may be the last to be activated precisely because of this (33). Overall, these data clearly link transcriptional regulation, via the sigma factors, to developmental cycle progression and secondary differentiation.

Despite the clear changes in transcription that occur during the chlamydial developmental cycle, activation of late gene expression alone does not guarantee secondary differentiation. We recently explored the function of the Clp protease systems in chlamydial growth and development (36). Interestingly, we observed that the inability to degrade SsrA-tagged products prevented functional secondary differentiation with a severe defect in EB production. This was demonstrated in strains unable to degrade SsrA products either by preventing their recognition by ClpX or by altering the SsrA tag to a version that is not degraded efficiently. Nonetheless, late gene transcription was activated in these strains. Therefore, post-translational regulation of secondary differentiation is also critical to the process.

Assuming a post-translational gene regulation model for secondary differentiation, we were interested in exploring known differences between EBs and RBs as a clue to what factors could be serving as a signal for this process. EBs and RBs differ in a number of characteristics, from their function to their morphology to their gene expression. In 2013, a study from Barker et al. added another difference – their relative levels of c-di-AMP with RBs having lower levels than EBs. Although the study from Barker et al. was focused on understanding how IFNβ is activated in *Chlamydia*-infected cells, we were intrigued by this relative difference in c-di-AMP levels. We hypothesized that c-di-AMP production might be a signal for secondary differentiation. We compared chlamydial growth in normal HeLa or STING-KO HeLa cells but could not detect any differences in growth in these cell types (Fig. 4G&S6 vs Fig. S10), indicating that c-di-AMP production by *Chlamydia* is unlikely used as means of activating the host cell. Rather, STING activation is an "accidental" outcome during chlamydial developmental cycle progression. Using recently developed genetic tools for *Chlamydia*, we demonstrated that we could manipulate the levels of c-di-AMP in the bacterium to either block or stimulate its production. Excitingly, we observed that elevated c-di-AMP was linked to earlier secondary differentiation whereas blocking c-di-AMP production prevented EB production (Fig. 7). Importantly, c-di-AMP production resulted in an increase in late gene transcripts as noted above for the function of the alternative sigma factors (Fig. 6). Therefore, c-di-AMP may act as a post-translational mechanism to trigger secondary differentiation. We can also directly conclude from our experiments that the threshold level of c-di-AMP necessary to trigger secondary differentiation in our culture conditions is greater than 1,000 pg/mL and less than 2,000 pg/mL. We base this on the fact that the 16hpi level of c-di-AMP in the DacA-overexpressing strain (1,000 pg/mL) did not trigger secondary differentiation whereas, during a

normal infection (e.g., mCherry expressing strain) when EBs are being produced at 24hpi, the c-di-AMP level is ~2,000pg/mL. To our knowledge, this is the first study to identify and provide experimental evidence for a signaling factor involved in differentiation of an obligate intracellular bacterium.

How might c-di-AMP function in *Chlamydia*? This is not readily apparent because *Chlamydia* lacks orthologs of proteins that have been characterized in other bacterial systems to bind c-di-AMP. For example, *Chlamydia* lacks annotated K$^+$ transporters, and it is unknown how these bacteria maintain their osmolarity. However, a recent study suggested that changes in K$^+$ levels affect the chlamydial developmental cycle (48). Therefore, c-di-AMP may function as a K$^+$ homeostasis coordinator in *Chlamydia* through as-yet unknown pathways. We did observe altered chlamydial morphology in some of our strains in which we altered c-di-AMP levels, and this may indicate changes in osmostability or effects on cell division. In regard to the latter possibility, in *S. aureus*, DacA activity is inhibited by interacting with GlmM (phosphoglucosamine mutase) (49). GlmM produces UDP-GlcNAc that is a precursor in both peptidoglycan and LPS synthesis. Given that *Chlamydia* uses peptidoglycan for cell division, we tested whether DacA and YbbR were localized to the division septum. However, their localization was not associated with the division septum (Fig. S10). In addition, we did not detect an interaction between DacA and GlmM using the Bacterial Adenylate Cyclase-based Two Hybrid (BACTH) system (data not shown), and the chlamydial GlmM lacks the residues required for DacA interaction. Therefore, we conclude that DacA and YbbR are unlikely to directly impact cell division. However, based on the cell morphological changes in DacA overexpression, we cannot exclude a function for c-di-AMP on cell wall metabolism. Future studies will focus on identifying c-di-AMP binding proteins and characterizing their function in *Chlamydia*. We also cannot exclude a function for c-di-AMP in directly manipulating gene expression by riboswitches (50), and further work is necessary to understand how c-di-AMP directly controls chlamydial development.

The use of c-di-AMP as a signal for secondary differentiation presents a "chicken-or-egg" quandary. How is the activity and expression of DacA, the diadenylate cyclase, regulated? Our transcript analyses indicate that *dacA-ybbR* transcripts peak during the RB phase of growth, similar to most genes in *Chlamydia*. One possibility not easily tested due to the obligate intracellular nature of *Chlamydia*, is that c-di-AMP production depletes ATP levels, which then reduce σ^{66} activity as described above. Our data indicate YbbR is necessary to activate DacA function since expressing membrane-localized or ΔTM isoforms of DacA alone only modestly increased c-di-AMP levels (Fig. 1E). However, overexpressing DacA alone did negatively impact chlamydial growth, suggesting that the balance between DacA and YbbR levels is important, with YbbR being the limiting factor for c-di-AMP production. For example, too much DacA insertion into the membrane without binding to YbbR may disrupt the membrane biology of the RB. This is supported by the fact that we observed no detrimental effects of overexpressing ΔTM isoforms of either wild-type or mutant (D164N) DacA. Indeed, overexpressing ΔTMDacA(D164N) did not alter c-di-AMP levels. Therefore, YbbR itself may be a target for regulation. As a monotopic transmembrane protein, YbbR may be degraded by proteases such as the inner membrane-associated FtsH or the periplasmic proteases HtrA or Tsp. This would presumably shut down c-di-AMP synthesis. Alternatively, DacA may interact with other membrane proteins, and overexpressing it alone may impair the function of this binding partner(s), resulting in the observed phenotypes. However, this possibility contradicts the effects of overexpressing the D164N isoform, which blocked c-di-AMP production. These experiments were conducted in the presence of the

chromosomally expressed copy of DacA, and we suggest that the mutant isoform acts as a dominant
negative by binding and/or interfering with the wild-type DacA or by titrating away YbbR from the
wild-type DacA. This is also supported by the fact that overexpressing the ΔTMDacA(D164N) isoform
had no impact on c-di-AMP levels or chlamydial growth. In contrast, knocking down *dacA-ybbR*
transcripts will reduce, but not completely eliminate, DacA and YbbR proteins, which may result in
residual c-di-AMP levels as compared to overexpressing the DacA mutant (either alone or with YbbR in
*dacA*opMut) as well as the phenotypic differences between these strains. Further work is needed to test
these possibilities. However, our data are clear in linking c-di-AMP levels to chlamydial developmental
progression.

Even though diverse functions of c-di-AMP in other bacteria have been reported previously, this
is the first time c-di-AMP has been described as a checkpoint in chlamydial development. It is possible
that the levels of c-di-AMP act as a de facto means for monitoring the bacterial population within the
inclusion or for their overall developmental status. Canonical bacteria can use quorum sensing to detect
bacteria at the population level and to communicate with other species (51). However, *C. trachomatis*
lacks homologues of genes related to quorum sensing (28). In addition, secondary differentiation occurs
asynchronously, and this feature is different from a quorum sensing mechanism. Our model accounts for
the asynchronicity of secondary differentiation. As RBs accumulate c-di-AMP and divide, it is likely the
c-di-AMP will be distributed unevenly between the mother and daughter cell since *Chlamydia* divides
through an asymmetric budding mechanism (7) (Fig. 8). This will lead to two cells with different levels
of c-di-AMP, one of which may then accumulate enough c-di-AMP to trigger secondary differentiation
while the other continues to divide (Fig. 8). This is consistent with a recent model proposing that a
population of division-competent RBs is maintained throughout the developmental cycle (52). Similar
uneven distribution of post-translational factors between asymmetrically dividing mother and daughter
cells would also impact other recently described signals for differentiation, including the levels of ClpC
protein or overall redox via AhpC antioxidant levels(35, 37).

Materials and Methods

Organisms and Cell Culture

Wild-type (ATCC, Manassas, VA) and STING KO (Dr. Frank van Kuppeveld; Utrecht University)
HeLa, human cervical epithelial-derived, and McCoy (kind gift of Dr. Harlan Caldwell, NIH), mouse
fibroblast-derived, cell lines were cultured at 37˚C with 5% CO_2 in Dulbecco's Modified Eagle Medium
(DMEM; Invitrogen, Waltham, MA) containing 10% fetal bovine serum (FBS; Hyclone, Logan, UT)
and 10 μg/mL gentamicin (Gibco, Waltham, MA). *C. trachomatis* serovar L2 (434/Bu) lacking the
endogenous plasmid (-pL2; kind gift of Dr. Ian Clarke, Univ. Southampton) was used for transformation.
All cell cultures and chlamydial stocks were routinely tested for Mycoplasma contamination using the
Mycoplasma PCR detection kit (Sigma, St. Louis, MO). For *E. coli*, NEB10β competent cells (New
England Biolabs, Ipswich, MA) were used for the amplification of pBOMB-derivative vectors. *E. coli*
was grown at 30˚C in LB media. All chemicals and antibiotics were obtained from Sigma unless
otherwise noted.

Cloning

The list of the vectors and primers used in this study is detailed in Supplemental Table 1. Target genes
were amplified by PCR with Phusion DNA polymerase (NEB) using 10 ng *C. trachomatis* L2 genomic
DNA or appropriate vectors as a template. Some DNA segments were directly synthesized as a gBlock
fragment (Integrated DNA Technologies, Coralville, IA). If plasmids were used as a template, then we
treated the PCR product with DpnI enzyme to remove templates. The PCR products were purified using
a PCR purification kit (Qiagen, Hilden, Germany). The HiFi Assembly reaction master mix (NEB) was
used following the manufacturer's manual in conjunction with plasmids pBOMBL(29) linearized with
EagI and KpnI or pBOMBL12CRia (empty vector) linearized with BamHI. The linearized plasmids
were also dephosphorylated with FastAP (ThermoFisher). The products of the HiFi reaction were
transformed into NEB10β competent cells (NEB) and plated on LB agar with appropriate antibiotics.
Plasmids were subsequently isolated using a mini-prep kit (Qiagen) and verified by plasmid digest and
sequencing from individual colonies grown overnight in LB broth with appropriate antibiotic selection.

Transformation of *Chlamydia trachomatis*

McCoy cells were plated in a six-well plate the day before beginning the transformation procedure. *C.*
trachomatis serovar L2 without plasmid (-pL2) resuspended with Tris-CaCl$_2$ buffer (10 mM Tris-Cl pH
7.5, 50 mM CaCl$_2$) was incubated with 2 μg plasmid at room temperature for 30 min. During this step,
McCoy cells were washed with 2 mL Hank's Balanced Salt Solution (HBSS) media containing Ca^{2+} and
Mg^{2+} (Gibco). After that, McCoy cells were infected with the transformants in 2 mL HBSS per well.
The plate was centrifuged at 400 x g for 15 min at room temperature and incubated at 37˚C for 15 min.
The inoculum was aspirated, and 2 mL 1X DMEM containing 10% FBS and 10 μg/mL gentamicin was
added per well. At 8 hours post infection (hpi), 1 μg/mL cycloheximide and either 1 or 2 U/mL
penicillin G or 500 μg/mL spectinomycin were added, and the plate was incubated at 37˚C until 48 hpi.
At 48 hpi, the transformants were harvested and infected onto a new McCoy cell monolayer. These
harvest and infection steps were repeated every 44-48 hpi until fluorescent, antibiotic-resistant
inclusions were observed.

(RT-)qPCR

HeLa cells were infected with chlamydial transformants at an MOI of 0.5. At 10 hpi, 5 nM
anhydrotetracycline (aTc) was added or not to the culture medium. Total RNA and DNA were harvested
at this time point from duplicate wells not treated with aTc. At 14 and 24 hpi, total RNA and DNA were
collected using Trizol (Invitrogen) and DNeasy Tissue (Qiagen) kit, respectively, as described
elsewhere(33). After DNase treatment of total RNA, cDNA was synthesized using Superscript III
reverse transcriptase (Invitrogen). After diluting the cDNA 10-fold, 5 μl of the diluted cDNA was used
as a template for qPCR. Equal masses of genomic DNA were used from each of the samples to quantify
chlamydial genomes, which were used to normalize transcript data as described(41). For both cDNA and
gDNA samples, qPCR reactions were prepared using 2X SYBR Green (ThermoFisher) in a total volume
of 25 uL per well. Standard cycling conditions were used with a melting curve analysis to verify
products. Transcripts and genome copies were assessed from at least three biological replicates.

Indirect immunofluorescence assay (IFA)

HeLa cells were infected with chlamydial transformants as above. At 10 hpi, 5 nM aTc was added or not,
and the infected cells were fixed with fixing solution (3.2% formaldehyde and 0.022% glutaraldehyde in

1X DPBS) for 2 min and permeabilized with 90% MeOH for 1 min at 10.5 hpi or 24 hpi. The fixed cells
were labeled with primary antibodies including rabbit anti-DacA (custom anti-peptide antibody targeting
the C-terminal TRNERKTNPIISWMRKK prepared by Pacific Immunology, Ramona, CA), goat anti-
major outer-membrane protein (MOMP; Meridian, Memphis, TN), and rabbit anti-six histidine tag
(Genscript, Piscataway, NJ and Abcam, Cambridge, UK, respectively). To visualize the primary
antibodies, donkey anti-goat antibody (488), donkey anti-mouse (405), or donkey anti-rabbit antibody
(594) were used as secondary antibodies. The secondary antibodies were obtained from Invitrogen or
Jackson Immunology (West Grove, PA). Coverslips were observed using a Zeiss AxioImager.Z2 with
Apotome2 as noted in the figure legends.

Inclusion forming unit (IFU) measurement

HeLa cells were infected with chlamydial transformants as above. At 10 hpi, 5 nM aTc was added or not
to the culture medium. At the indicated times, infected cells were harvested in 1 mL 2SP media then
frozen at -80˚C. After thawing the lysates, the samples were serially 1:10 diluted and used to infect
HeLa cells seeded in 24-well plates. At 24 hpi, the number of GFP expressing inclusions was counted
from 30 fields of view to calculate the IFUs from the original sample. Three biological replicates were
performed.

Cyclic di-AMP measurement

HeLa cells were infected with the indicated chlamydial transformants as above. At 10 hpi, expression of
the constructs was induced or not with 5 nM aTc, and, at 16 and 24 hpi, samples were prepared for
measuring the level of cyclic di-AMP. After aspirating the media, the infected cell monolayers were
washed with 1X PBS and resuspended with B-Per Bacterial Cell Lysis Buffer (Thermo). Samples were
vortexed for 1 min and then centrifuged at 13,300xg for 15 min at 4□. The levels of cyclic di-AMP from
the supernatants of the cell lysates were quantified using the Cyclic di-AMP ELISA kit (Cayman)
following the manufacturer's instructions(53).

Preparation of RNA sequencing samples

Samples were prepared as reported previously(33). Briefly, the transformants of *dacA*op and *dacA*-KD
were infected into HeLa cells. At 10 hpi, the constructs were induced or not with 5 nM aTc. RNA
samples were prepared from *dacA*op-infected cells and *dacA*-KD-infected cells at 16 hpi and 24 hpi,
respectively. 20 μg RNA samples were treated with DNase to remove DNA contamination using DNA-
free Turbo kit (Thermo) according to the manufacturer's instruction. Ribosomal RNA was depleted from
samples using the MICROBEnrich (Thermo) and MICROBExpress kits (Thermo) following the
manufacturer's instructions. RNA samples were processed for sequencing by the UNMC Genomics
Core Facility. The resultant libraries from the individual samples were multiplexed and subjected to
100-bp paired-read sequencing to generate approximately 60 million pairs of reads per sample on an
Illumina NovaSeq6000 sequencer in the UNMC Genomics Core facility. The original fastq format reads
were trimmed by fqtirm tool (https://ccb.jhu.edu/software/fqtrim) to remove adapters, terminal unknown
bases (Ns) and low quality 3' regions (Phred score < 30). The trimmed fastq files were processed by
FastQC (54) for quality control . *Chlamydia trachomatis* 434/Bu bacterial reference genome and
annotation files were downloaded from Ensembl
(http://bacteria.ensembl.org/Chlamydia_trachomatis_434_bu/Info/Index). Sequencing data were

analyzed by the Bioinformatics and Systems Biology Core (BSBC). The trimmed fastq files were mapped to *Chlamydia trachomatis* 434/Bu by CLC Genomics Workbench 23 for RNAseq analyses.

Data availability

The raw and processed RNA sequencing reads in fastq format have been deposited in the Gene Expression Omnibus (GEO; www.ncbi.nlm.nih.gov/geo/) under accession no. GSE252732. All other data are available in the main text or the supplementary materials.

Statistical Analysis

To analyze the statistical significance between uninduced and induced samples of qPCR and IFU data, we used two sample equal variance Student's t-test. For the levels of cyclic di-AMP data, we used two sample equal variance Student's t-test.

Acknowledgements

This study was supported in part by an NIH/NIGMS award (R35GM124798) and in part by an NIH/NIAID award (R21AI180574) to SPO and by start-up funds from UNMC. The UNMC Genomics Core Facility receives partial support from the National Institute for General Medical Science (NIGMS) INBRE - P20GM103427-19, as well as the National Cancer Institute the Fred & Pamela Buffett Cancer Center Support Grant- P30CA036727. This publication's contents are the sole responsibility of the authors and do not necessarily represent the official views of the NIH or NIGMS. The authors would like to thank Dr. Frank van Kuppeveld (Utrecht University) for STING-KO HeLa cells, Dr. Harlan Caldwell (NIAID/NIH) for McCoy cells, and Dr. Ian Clarke (University of Southampton) for the plasmidless *C. trachomatis* serovar L2 strain. In addition, we thank the UNMC Genomics Core Facility and the Bioinformatics and Systems Biology Core facility for RNA sequencing and analysis services, respectively.

Author contributions:

Junghoon Lee, Methodology, Investigation, Visualization, Writing – original draft, and Writing-review & editing

Scot P. Ouellette, Conceptualization, Methodology, Investigation, Visualization, Supervision, and Writing-review & editing

Competing interests: Authors declare that they have no competing interests.

Figure Legends

Figure 1. Cyclic di-AMP accumulation is linked to secondary differentiation in *C. trachomatis*. A. Defining characteristics of chlamydial EBs and RBs. **B.** A hypothetical model representing the correlation between c-di-AMP levels and the timing of secondary differentiation. The dashed line represents a threshold level of c-di-AMP needed to drive secondary differentiation in a given RB. hpi = hours post-infection. **C.** Measurement of c-di-AMP concentrations in uninfected (-Ctr) and infected (+Ctr) HeLa cell lysates. For infected HeLa cells, *C. trachomatis* serovar L2 (434/Bu) transformed with an mCherry-encoding construct was infected into HeLa cells, and expression of mCherry was induced at 10 hpi with 5 nM anhydrotetracycline (aTc). All samples were harvested at 16 and 24 hpi. **D.** A schematic diagram of the constructs used in this study. All constructs used are aTc-inducible as shown by the P_{tet} promoter. The location of the 6xH tag, the approximate location of the crRNA for the CRISPRi vectors are shown as well as the transmembrane domain of DacA. Diagram is not to scale. **E.** Measurement of c-di-AMP concentrations in infected cell lysates from the strains shown in panel D. *C. trachomatis* serovar L2 (434/Bu) transformed with the indicated constructs was infected into HeLa cells. At 10 hpi, expression of the construct was induced with 5 nM aTc, and the infected cells were harvested at 16 or 24 hpi. Levels of c-di-AMP in the supernatant were measured using ELISA. The left and right panels show the levels of c-di-AMP in the indicated strains at 16 and 24 hpi, respectively, on a log2 scale. For reference, $2^{10} = 1,024$, $2^{15} = 32,768$, and $2^{20} = 1,048,576$. The dashed line in both graphs represents the c-di-AMP level of the 24hpi mCherry-expressing control that is associated with EB production. *: p<0.05, **: p<0.001, NS: Not significant via two sample equal variance t-test compared to the mCherry control.

Figure 2. Overexpression of DacA or DacA(D164N) are detrimental to the chlamydial developmental cycle. HeLa cells were infected with *C. trachomatis* transformed with a plasmid encoding an anhydrotetracycline (aTc)-inducible DacA or DacA(D164N) (i.e., *dacA* and *dacA*(D164N) respectively; see Fig. 1D). At 10 hpi, expression of the construct was induced or not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi. Immunofluorescence analysis (IFA) and inclusion forming units (IFU) samples were collected at 24hpi. For IFA images, the green color represents chlamydial major outer membrane protein (MOMP), which shows the chlamydial cell morphology, and the red color represents DacA or DacA(D164N). **A&B.** IFA images of the *dacA*(A) and *dacA*(D164N)(B) strains at 24 hpi. Shown are individual panels of a representative inclusion for the strains with DacA and MOMP labeling as well as the merged image. IFA images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 1 µm (A) or 2 µm (B). **C&D.** Quantification of IFUs (C) and genomic DNA copy number (D) from uninduced and induced samples of *dacA* at 24 hpi. **E&F.** Quantification of IFUs (E) and genomic DNA copy number (F) from uninduced and induced samples of *dacA*(D164N) at 24 hpi. **G&H.** Quantification of transcripts by RT-qPCR for *dacA, ybbR*, *euo*, *hctA*, and *omcB* from uninduced and induced samples of *dacA* (G) and *dacA*(D164N) (H). UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: p<0.05; **: p<0.001 via two sample equal variance t-test.

Figure 3. Overexpression of ΔTMDacA or ΔTMDacA(D164N) does not alter the chlamydial developmental cycle. HeLa cells were infected with *C. trachomatis* transformed with a plasmid encoding an anhydrotetracycline (aTc)-inducible ΔTMDacA or ΔTMDacA(D164N) (i.e., ΔTM*dacA* and

ΔTM*dacA*(D164N); see Fig. 1D). At 10 hpi, expression of the construct was induced or not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi. Immunofluorescence analysis (IFA) and inclusion forming units (IFU) samples were collected at 24 hpi. For IFA images, the green color represents chlamydial major outer membrane protein (MOMP), which shows the chlamydial cell morphology, and the red color represents ΔTMDacA or ΔTMDacA(D164N). **A&B.** IFA images of ΔTM*dacA*(A) and ΔTM*dacA*(D164N)(B) strains at 24 hpi. Shown are individual panels of a representative inclusion for the strains with DacA and MOMP labeling as well as the merged image. IFA images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 2 µm. **C&D.** Quantification of IFUs (C) and genomic DNA copy number (D) from uninduced and induced samples of ΔTM*dacA* at 24 hpi. **E&F.** Quantification of IFUs (E) and genomic DNA copy number (F) from uninduced and induced samples of ΔTM*dacA*(D164N). **G&H.** Quantification of transcripts by RT-qPCR for *dacA, ybbR*, *euo*, *hctA*, and *omcB* from uninduced and induced samples of ΔTM*dacA* (G) and ΔTM*dacA*(D164N) (H). UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: p<0.05 via two sample equal variance t-test.

Figure 4. CRISPRi-mediated *dacA-ybbR* knockdown displays reduced levels of transcripts for late genes. HeLa cells were infected with *C. trachomatis* transformed with a plasmid encoding an anhydrotetracycline (aTc)-inducible CRISPRi-dCas12 system targeting the *dacA* promoter (*dacA*-KD) or *dacA*-KD system and DacA/YbbR_6xH (i.e., *dacA*-KDcom; see Fig. 1D). At 10 hpi, knockdown was induced or not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi. Immunofluorescence analysis (IFA) and inclusion forming units (IFU) samples were collected at 24hpi. **A&B.** IFA images of the *dacA*-KD (A) and *dacA*-KDcom (B) strains at 24 hpi. Shown are individual panels of a representative inclusion for the strains for dCas12, YbbR_6xH, and major outer membrane protein (MOMP) labeling as well as the merged image. IFA images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 2(A) or 1(B) µm. **C&D.** Quantification of IFUs (C) and genomic DNA copy number (D) from uninduced and induced samples of *dacA*-KD at 24 hpi. **E&F.** Quantification of IFUs (E) and genomic DNA copy number (F) in uninduced and induced samples of *dacA*-KDcom. **G&H.** Quantification of transcripts by RT-qPCR for *dacA, ybbR*, *euo*, *hctA*, and *omcB* from uninduced and induced samples of *dacA*-KD (G) and *dacA*-KDcom (H). UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: p<0.05; **: p<0.001 via two sample equal variance t-test.

Figure 5. Overexpression of DacA and YbbR_6xH prematurely increases *hctA* transcript levels. HeLa cells were infected with *C. trachomatis* transformed with an aTc-inducible plasmid encoding wild-type DacA/YbbR_6xH or DacA(D164N)/YbbR_6xH (i.e., *dacA*op and *dacA*opMut, respectively; see Fig. 1D). At 10 hpi, expression of the constructs were induced or not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi. Immunofluorescence analysis (IFA) samples were collected at 24 hpi. **A&B.** IFA images of the *dacA*op (A) and *dacA*opMut (B) at 24 hpi. Shown are individual panels of a representative inclusion for the strains for DacA and YbbR_6xH as well as the merged image with major outer membrane protein (MOMP) labeling. The arrowheads represent the co-localization of DacA and YbbR. IFA images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 2 µm. **C&D** Quantification of IFUs (C) and genomic DNA copy number (D) from uninduced and induced samples of *dacA*op at 24 hpi. **E&F.** Quantification of IFUs (E) and genomic DNA copy number (F) from uninduced and induced samples of *dacA*opMut.

G&H. Quantification of transcripts by RT-qPCR for *dacA, ybbR, euo, hctA,* and *omcB* from uninduced and induced samples of *dacA*op (G) and *dacA*opMut (H). UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: p<0.05 two sample equal variance t-test.

Figure 6. The levels of c-di-AMP are correlated with late gene transcripts. RNA sequencing was performed from HeLa cells infected with either the *dacA*op or *dacA*-KD strains. RNA samples were collected at 16hpi for *dacA*op and 24hpi for *dacA*-KD after inducing expression of the relevant constructs at 10hpi. Shown is a volcano plot of the RNA sequencing results with the vertical dashed lines indicating a two-fold change in transcript levels as compared to the respective uninduced control for the given strain and the horizontal lines indicating a p-value of 0.05. The plot was made using GraphPad Prism software. Green spots represent genes demonstrating a statistically significant two-fold change in transcription levels between uninduced and induced samples. Red dots represent genes with significant changes in transcript levels less than two-fold. Blue dots represent genes not significantly different but more than two-fold changed between the conditions. Black dots represent genes not significantly different and less than two-fold changed between the conditions. See also Table 1 and Supplemental Tables 2 and 3 for more details.

Figure 7. EB production is induced by high levels of c-di-AMP. (A-C) HeLa cells were infected with the chlamydial transformants *dacA*op (A), *dacA*-KD (B), and *dacA*opMut (C). Expression of the constructs was induced or not with 5 nM aTc at 10hpi. At 18, 20, 22, 24, 32, and 48 hpi, infected cell lysates were harvested for IFU quantification. UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. #: Non-detected from the induced samples. *: p<0.05; **: p<0.001 via two sample equal variance t-test.

Figure 8. A working model illustrating how c-di-AMP impacts the onset of secondary differentiation in *C. trachomatis*. A. During RB division, there is a gradient of c-di-AMP that forms between the mother and daughter cell. The lighter blue represents lower levels of c-di-AMP, the darker blue represents higher levels of c-di-AMP. In the depicted scenario, the daughter cell is able to divide again, having not reached a critical threshold of c-di-AMP. In contrast, the mother cell accumulates sufficient c-di-AMP to trigger secondary differentiation to an EB. **B.** A model illustrating how c-di-AMP levels impact secondary differentiation progression. Shown is the "normal" condition of wild-type bacteria, bacteria overexpressing DacA and YbbR (higher cyclic di-AMP), and bacteria expressing the CRISPRi system targeting the *dacA* promoter (lower cyclic di-AMP). Depending on the amount of c-di-AMP produced, secondary differentiation can either be triggered earlier or later as shown. The black dots represent EBs; the bigger and blue-colored circles show the RB cells. The brightness of blue reflects the c-di-AMP level.

716 **Table 1. Genes impacted by cyclic di-AMP levels.**

Gene ID	Ctr D ORF	Name	Protein Names	Fold change		Ref.
			Canonical late genes	*dacA*op_OE	*dacA*-KD	
CTL0112	CT743	*hctA*	Histone H1-like protein HC1	6.70	-3.47	(11, 56)
CTL0302	CT046	*hct2*	Histone H1-like protein HC2	3.52	-5.96	(11, 57)
CTL0336	CT080	*ltuB*	Late transcription unit B protein	4.00	-3.61	(11, 56)
CTL0700	CT441	*tsp*	Carboxy-terminal processing protease	6.18	-6.62	(57)
CTL0702	CT443	*omcB*	Large cysteine-rich periplasmic protein	5.00	-4.04	(11, 34)
CTL0703	CT444	*omcA*	Small cysteine-rich outer membrane protein	5.93	-3.71	(11, 34)
CTL0716	CT456	*tarp*	Translocated actin-recruiting phosphoprotein	6.06	-5.97	(33, 34)
			Membrane organization associated	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.
CTL0082	CT713	*ompB*	Outer membrane protein B	2.73	-2.15	(58)
CTL0248	CT869	*pmpE*	Polymorphic outer membrane protein	1.99	-2.04	(33, 58)
CTL0249	CT870	*pmpF*	Polymorphic outer membrane protein	3.03	-2.23	(33, 58)
CTL0250	CT871	*pmpG*	Polymorphic outer membrane protein	2.03	-1.55	(11, 58)
CTL0429	CT177	*dsbA*	Disulfide bond chaperone	2.79	-2.47	(33)
CTL0610	CT356	*dsbH*	Thioredox_DsbH domain-containing protein	3.07	-2.11	(11, 58)
CTL0670	CT413	*pmpB*	Polymorphic outer membrane protein	4.17	-1.74	(33, 58)
			Gene Regulation associated	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.
CTL0044	CT675	*mcsB*	Protein-arginine kinase	3.52	-2.21	(33)
CTL0045	CT676	*mcsA*	UVR domain-containing protein	2.53	-2.92	(33)
CTL0727	CT467	*atoS*	Two component regulator, histidine kinase	3.33	-1.80	(33)
CTL0728	CT468	*atoC*	Two component system response regulator	7.95	-2.78	
CTL0894	CT630	*chxR*	Atypical response regulator protein ChxR	3.68	-2.93	(59)
			Glycogen synthesis	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.

Gene ID	Ctr D ORF	Name	Protein Names	dacAop_OE	dacA-KD	Ref.
CTL0167	CT798	*glgA*	Glycogen synthase	9.58	-2.86	(11, 58)
CTL0342	CT087	*malQ*	4-alpha-glucanotransferase	9.43	-4.29	(42)
CTL0500	CT248	*glgP*	Alpha-1,4 glucan phosphorylase	2.70	-1.71	(58)
			Type III Secretion System	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.
CTL0041	CT672	*sctQ*	Type III secretion component, basal body	1.93	-1.76	(42)
CTL0043	CT674	*cdsC*	Type III secretion structural protein	1.79	-1.56	(42)
CTL0343	CT088	*scc1*	Type III secretion chaperone	4.48	-3.78	(42)
CTL0824	CT561	*sctL*	Type III secretion system protein	2.18	-1.75	(33)

OE = overexpression; KD = knockdown

References

1.   Abdelrahman YM, Belland RJ. 2005. The chlamydial developmental cycle. FEMS Microbiol

Rev 29:949-59.

2.   Everett KD, Hatch TP. 1991. Sequence analysis and lipid modification of the cysteine-rich

envelope proteins of Chlamydia psittaci 6BC. J Bacteriol 173:3821-30.

3.   Hackstadt T, Baehr W, Ying Y. 1991. Chlamydia trachomatis developmentally regulated protein

is homologous to eukaryotic histone H1. Proc Natl Acad Sci U S A 88:3937-41.

4.   Moulder JW. 1993. Why is Chlamydia sensitive to penicillin in the absence of peptidoglycan?

Infect Agents Dis 2:87-99.

5.   Fox A, Rogers JC, Gilbart J, Morgan S, Davis CH, Knight S, Wyrick PB. 1990. Muramic acid is

not detectable in Chlamydia psittaci or Chlamydia trachomatis by gas chromatography-mass

spectrometry. Infect Immun 58:835-7.

6.   Barbour AG, Amano K, Hackstadt T, Perry L, Caldwell HD. 1982. Chlamydia trachomatis has

penicillin-binding proteins but not detectable muramic acid. J Bacteriol 151:420-8.

7.   Abdelrahman Y, Ouellette SP, Belland RJ, Cox JV. 2016. Polarized Cell Division of Chlamydia

trachomatis. PLoS Pathog 12:e1005822.

8.   Liechti G, Kuru E, Packiam M, Hsu YP, Tekkam S, Hall E, Rittichier JT, VanNieuwenhze M,

Brun YV, Maurelli AT. 2016. Pathogenic Chlamydia Lack a Classical Sacculus but Synthesize a

Narrow, Mid-cell Peptidoglycan Ring, Regulated by MreB, for Cell Division. PLoS Pathog

12:e1005590.

9.   Ouellette SP, Karimova G, Subtil A, Ladant D. 2012. Chlamydia co-opts the rod shape-

determining proteins MreB and Pbp2 for cell division. Mol Microbiol 85:164-78.

10.   Lee J, Cox JV, Ouellette SP. 2020. Critical Role for the Extended N-Terminus of Chlamydial

MreB in Directing Its Membrane Association and Potential Interaction with Divisome Proteins. J

Bacteriol doi:10.1128/JB.00034-20.

11.   Belland RJ, Zhong G, Crane DD, Hogan D, Sturdevant D, Sharma J, Beatty WL, Caldwell HD.
        2003. Genomic transcriptional profiling of the developmental cycle of Chlamydia trachomatis.
        Proc Natl Acad Sci U S A 100:8478-83.

12.   Ouellette SP, Hatch TP, AbdelRahman YM, Rose LA, Belland RJ, Byrne GI. 2006. Global
        transcriptional upregulation in the absence of increased translation in Chlamydia during
        IFNgamma-mediated host cell tryptophan starvation. Mol Microbiol 62:1387-401.

13.   Barker JR, Koestler BJ, Carpenter VK, Burdette DL, Waters CM, Vance RE, Valdivia RH. 2013.
        STING-dependent recognition of cyclic di-AMP mediates type I interferon responses during
        Chlamydia trachomatis infection. mBio 4:e00018-13.

14.   Girardin SE, Boneca IG, Viala J, Chamaillard M, Labigne A, Thomas G, Philpott DJ, Sansonetti
        PJ. 2003. Nod2 is a general sensor of peptidoglycan through muramyl dipeptide (MDP) detection.
        J Biol Chem 278:8869-72.

15.   Packiam M, Weinrick B, Jacobs WR, Jr., Maurelli AT. 2015. Structural characterization of
        muropeptides from Chlamydia trachomatis peptidoglycan by mass spectrometry resolves
        "chlamydial anomaly". Proc Natl Acad Sci U S A 112:11660-5.

16.   Booth IR, Higgins CF. 1990. Enteric bacteria and osmotic stress: intracellular potassium
        glutamate as a secondary signal of osmotic stress? FEMS Microbiol Rev 6:239-46.

17.   Kim H, Youn SJ, Kim SO, Ko J, Lee JO, Choi BS. 2015. Structural Studies of Potassium
        Transport Protein KtrA Regulator of Conductance of K+ (RCK) C Domain in Complex with
        Cyclic Diadenosine Monophosphate (c-di-AMP). J Biol Chem 290:16393-402.

18.   Gibhardt J, Hoffmann G, Turdiev A, Wang M, Lee VT, Commichau FM. 2019. c-di-AMP assists
        osmoadaptation by regulating the Listeria monocytogenes potassium transporters KimA and

766   KtrCD. J Biol Chem 294:16020-16033.

767   19.   Quintana IM, Gibhardt J, Turdiev A, Hammer E, Commichau FM, Lee VT, Magni C, Stulke J.

768   2019. The KupA and KupB Proteins of Lactococcus lactis IL1403 Are Novel c-di-AMP

769   Receptor Proteins Responsible for Potassium Uptake. J Bacteriol 201.

770   20.   Moscoso JA, Schramke H, Zhang Y, Tosi T, Dehbi A, Jung K, Grundling A. 2016. Binding of

771   Cyclic Di-AMP to the Staphylococcus aureus Sensor Kinase KdpD Occurs via the Universal

772   Stress Protein Domain and Downregulates the Expression of the Kdp Potassium Transporter. J

773   Bacteriol 198:98-110.

774   21.   Corrigan RM, Campeotto I, Jeganathan T, Roelofs KG, Lee VT, Grundling A. 2013. Systematic

775   identification of conserved bacterial c-di-AMP receptor proteins. Proc Natl Acad Sci U S A

776   110:9084-9.

777   22.   Wang X, Cai X, Ma H, Yin W, Zhu L, Li X, Lim HM, Chou SH, He J. 2019. A c-di-AMP

778   riboswitch controlling kdpFABC operon transcription regulates the potassium transporter system

779   in Bacillus thuringiensis. Commun Biol 2:151.

780   23.   Bejerano-Sagie M, Oppenheimer-Shaanan Y, Berlatzky I, Rouvinski A, Meyerovich M, Ben-

781   Yehuda S. 2006. A checkpoint protein that scans the chromosome for damage at the start of

782   sporulation in Bacillus subtilis. Cell 125:679-90.

783   24.   Oppenheimer-Shaanan Y, Wexselblatt E, Katzhendler J, Yavin E, Ben-Yehuda S. 2011. c-di-

784   AMP reports DNA integrity during sporulation in Bacillus subtilis. EMBO Rep 12:594-601.

785   25.   Zhang L, He ZG. 2013. Radiation-sensitive gene A (RadA) targets DisA, DNA integrity

786   scanning protein A, to negatively affect cyclic Di-AMP synthesis activity in Mycobacterium

787   smegmatis. J Biol Chem 288:22426-36.

26.   Gandara C, de Lucena DKC, Torres R, Serrano E, Altenburger S, Graumann PL, Alonso JC.

2017. Activity and in vivo dynamics of Bacillus subtilis DisA are affected by RadA/Sms and by

Holliday junction-processing proteins. DNA Repair (Amst) 55:17-30.

27. Corrigan RM, Grundling A. 2013. Cyclic di-AMP: another second messenger enters the fray.

Nat Rev Microbiol 11:513-24.

28. Stephens RS, Kalman S, Lammel C, Fan J, Marathe R, Aravind L, Mitchell W, Olinger L,

Tatusov RL, Zhao Q, Koonin EV, Davis RW. 1998. Genome sequence of an obligate

intracellular pathogen of humans: Chlamydia trachomatis. Science 282:754-9.

29. Ouellette SP, Blay EA, Hatch ND, Fisher-Marvin LA. 2021. CRISPR Interference to Inducibly

Repress Gene Expression in Chlamydia trachomatis. Infection and Immunity *in press*.

30. Chiarelli TJ, Grieshaber NA, Omsland A, Remien CH, Grieshaber SS. 2020. Single-Inclusion

Kinetics of Chlamydia trachomatis Development. mSystems 5.

31. Bolte S, Cordelieres FP. 2006. A guided tour into subcellular colocalization analysis in light

microscopy. J Microsc 224:213-32.

32. Dunn KW, Kamocka MM, McDonald JH. 2011. A practical guide to evaluating colocalization in

biological microscopy. Am J Physiol Cell Physiol 300:C723-42.

33. Hatch ND, Ouellette SP. 2023. Identification of the alternative sigma factor regulons of

Chlamydia trachomatis using multiplexed CRISPR interference. mSphere 8:e0039123.

34. Soules KR, LaBrie SD, May BH, Hefty PS. 2020. Sigma 54-Regulated Transcription Is

Associated with Membrane Reorganization and Type III Secretion Effectors during Conversion

to Infectious Forms of Chlamydia trachomatis. mBio 11.

35. Jensen AA, Firdous S, Lei L, Fisher DJ, Ouellette SP. 2025. Overexpressing the ClpC AAA+

unfoldase accelerates developmental cycle progression in Chlamydia trachomatis. mBio

16:e0287024.

36. Wood NA, Swoboda AR, Blocker AM, Fisher DJ, Ouellette SP. 2022. Tag-Dependent Substrate

Selection of ClpX Underlies Secondary Differentiation of Chlamydia trachomatis. mBio

13:e0185822.

37. Singh V, Ouellette SP. 2025. Altering the redox status of Chlamydia trachomatis directly impacts

its developmental cycle progression. Elife 13.

38. Thompson CC, Griffiths C, Nicod SS, Lowden NM, Wigneshweraraj S, Fisher DJ, McClure MO.

2015. The Rsb Phosphoregulatory Network Controls Availability of the Primary Sigma Factor in

Chlamydia trachomatis and Influences the Kinetics of Growth and Development. PLoS Pathog

11:e1005125.

39. Kuwabara S, Landers ER, Fisher DJ. 2022. Impact of nutrients on the function of the chlamydial

Rsb partner switching mechanism. Pathog Dis 80.

40. Soules KR, Dmitriev A, LaBrie SD, Dimond ZE, May BH, Johnson DK, Zhang Y, Battaile KP,

Lovell S, Hefty PS. 2020. Structural and ligand binding analyses of the periplasmic sensor

domain of RsbU in Chlamydia trachomatis support a role in TCA cycle regulation. Mol

Microbiol 113:68-88.

41. Ouellette SP, Abdelrahman YM, Belland RJ, Byrne GI. 2005. The Chlamydia pneumoniae type

III secretion-related lcrH gene clusters are developmentally expressed operons. J Bacteriol

187:7853-6.

42. Hefty PS, Stephens RS. 2007. Chlamydial type III secretion system is encoded on ten operons

preceded by sigma 70-like promoter elements. J Bacteriol 189:198-206.

43. Tamura A, Matsumoto A, Manire GP, Higashi N. 1971. Electron microscopic observations on

the structure of the envelopes of mature elementary bodies and developmental reticulate forms of

Chlamydia psittaci. J Bacteriol 105:355-60.

44.   Tamura A, Manire GP. 1967. Preparation and chemical composition of the cell membranes of

developmental reticulate forms of meningopneumonitis organisms. J Bacteriol 94:1184-8.

45.   Clifton DR, Fields KA, Grieshaber SS, Dooley CA, Fischer ER, Mead DJ, Carabeo RA,

Hackstadt T. 2004. A chlamydial type III translocated protein is tyrosine-phosphorylated at the

site of entry and associated with recruitment of actin. Proc Natl Acad Sci U S A 101:10166-71.

46.   Swoboda AR, Wood NA, Saery EA, Fisher DJ, Ouellette SP. 2023. The Periplasmic Tail-

Specific Protease, Tsp, Is Essential for Secondary Differentiation in Chlamydia trachomatis. J

Bacteriol 205:e0009923.

47.   Grieshaber NA, Chiarelli TJ, Appa CR, Neiswanger G, Peretti K, Grieshaber SS. 2022.

Translational gene expression control in Chlamydia trachomatis. PLoS One 17:e0257259.

48.   Andrew SC, Dumoux M, Hayward RD. 2021. Chlamydia Uses K(+) Electrical Signalling to

Orchestrate Host Sensing, Inter-Bacterial Communication and Differentiation. Microorganisms 9.

49.   Tosi T, Hoshiga F, Millership C, Singh R, Eldrid C, Patin D, Mengin-Lecreulx D, Thalassinos K,

Freemont P, Grundling A. 2019. Inhibition of the Staphylococcus aureus c-di-AMP cyclase

DacA by direct interaction with the phosphoglucosamine mutase GlmM. PLoS Pathog

15:e1007537.

50.   Nelson JW, Sudarsan N, Furukawa K, Weinberg Z, Wang JX, Breaker RR. 2013. Riboswitches

in eubacteria sense the second messenger c-di-AMP. Nat Chem Biol 9:834-9.

51.   Miller MB, Bassler BL. 2001. Quorum sensing in bacteria. Annu Rev Microbiol 55:165-99.

52.   Chiarelli TJ, Grieshaber NA, Appa C, Grieshaber SS. 2023. Computational Modeling of the

Chlamydial Developmental Cycle Reveals a Potential Role for Asymmetric Division. mSystems

8:e0005323.

53.   Underwood AJ, Zhang Y, Metzger DW, Bai G. 2014. Detection of cyclic di-AMP using a

competitive ELISA with a unique pneumococcal cyclic di-AMP binding protein. J Microbiol

Methods 107:58-62.

54.   Andrews SC. 2010. FastQC: a quality control tool for high throughput sequence data.

55.   Tsirigos KD, Peters C, Shu N, Kall L, Elofsson A. 2015. The TOPCONS web server for

consensus prediction of membrane protein topology and signal peptides. Nucleic Acids Res

43:W401-7.

56.   Fahr MJ, Douglas AL, Xia W, Hatch TP. 1995. Characterization of late gene promoters of

Chlamydia trachomatis. J Bacteriol 177:4252-60.

57.   Yu HH, Kibler D, Tan M. 2006. In silico prediction and functional validation of sigma28-

regulated genes in Chlamydia and Escherichia coli. J Bacteriol 188:8206-12.

58.   Nicholson TL, Olinger L, Chong K, Schoolnik G, Stephens RS. 2003. Global stage-specific gene

regulation during the developmental cycle of Chlamydia trachomatis. J Bacteriol 185:3179-89.

59.   Yang C, Kari L, Sturdevant GL, Song L, Patton MJ, Couch CE, Ilgenfritz JM, Southern TR,

Whitmire WM, Briones M, Bonner C, Grant C, Hu P, McClarty G, Caldwell HD. 2017.

Chlamydia trachomatis ChxR is a transcriptional regulator of virulence factors that function in in

vivo host-pathogen interactions. Pathog Dis 75.

Fig.1

Fig.2

Fig.3

Fig.4

A. *dacA*-KD

B. *dacA*-KDcom

Fig.5

Fig.6

Fig.7

Fig.8