Cyclic di-AMP drives developmental cycle progression in *Chlamydia trachomatis*

Junghoon Lee[1] and Scot P. Ouellette[1*]

[1]Department of Pathology, Microbiology, and Immunology, College of Medicine, University of

Nebraska Medical Center, Omaha, NE, USA.

Running Head: Cyclic di-AMP drives secondary differentiation in *Chlamydia*

*Corresponding author:

Scot P. Ouellette

Department of Pathology, Microbiology, and Immunology

University of Nebraska Medical Center

985900 Nebraska Medical Center, DRC2 5029

Omaha, NE 68198

Phone: 402-559-0763

E-mail: scot.ouellette@unmc.edu

Abstract

The obligate intracellular bacterium *Chlamydia* alternates between two functional forms during its developmental cycle: elementary body (EB) and reticulate body (RB). However, the molecular mechanisms governing the transitions between these forms are unknown. Here, we present evidence cyclic di-AMP (c-di-AMP) is a key factor in triggering the transition from RB to EB (i.e., secondary differentiation) in the chlamydial developmental cycle. We made strains producing different levels of c-di-AMP, which we linked to changes in secondary differentiation status. Increases in c-di-AMP resulted in an earlier increase in transcription of EB-associated genes, and this was further manifested in earlier production of EBs. In contrast, when c-di-AMP levels were decreased, secondary differentiation was delayed. Based on these data, we conclude there is a threshold level of c-di-AMP needed to trigger secondary differentiation in *Chlamydia*. This is the first study to identify a physiological function for c-di-AMP production in *Chlamydia* as well as a mechanism by which secondary differentiation is initiated in these pathogens.

(160/250)

Importance

The second messenger molecule, cyclic di-AMP, shows diverse functions in bacteria. This molecule is usually detected in Gram-positive bacteria and is related to the osmotic stress response, DNA replication, and sporulation. *Chlamydia trachomatis*, a Gram-negative bacterium, encodes genes related to cyclic di-AMP synthesis. Cyclic di-AMP has been detected in *C. trachomatis*, where it has been shown to trigger a STING-dependent immune response in host cells. However, its physiological function in *C. trachomatis* is unknown. In this study, we identify a function for cyclic di-AMP in triggering gene expression linked to secondary differentiation in chlamydial developmental cycle. Our findings are important in understanding the molecular mechanism of the chlamydial developmental cycle and contribute to providing new therapeutic strategies for chlamydial infectious diseases.

(121/150)

Keywords: Chlamydia, cyclic di-AMP, c-di-AMP, cyclic dinucleotides, differentiation, development, second messenger, signaling

Introduction

Chlamydia species are major pathogens of humans and animals. These obligate intracellular bacteria share one key feature: their unique developmental cycle (See ref.(1)). During this cycle, *Chlamydia* transitions between two different functional and morphological forms: the elementary body (EB), an infectious but non-dividing cell, and the reticulate body (RB), a dividing but non-infectious cell (Fig. 1A). A third form, the intermediate body (IB) is a transitional form from the RB to EB. Besides these characteristics, EBs and RBs differ in other ways. For example, EBs are small (~0.3 μm), have a highly disulfide-crosslinked outer membrane(2) and have DNA condensed by histone-like proteins(3). In contrast, RBs are larger (~1 μm), have a Gram-negative cell envelope that lacks peptidoglycan(4-6), and have a "relaxed" chromosome. RBs divide by an asymmetric polarized division mechanism dependent on MreB-directed peptidoglycan synthesis specifically at the septum(7-10). Not surprisingly, *Chlamydia* expresses genes in a temporally defined manner that correspond broadly with its developmental cycle(11-13). "Early" genes are expressed immediately upon entry into a target host cell and are likely involved in establishing the intracellular niche of *Chlamydia*, the inclusion, and mediating primary differentiation from EB to RB. "Mid" cycle genes facilitate RB replication and division and inclusion growth. "Late" genes are expressed when secondary differentiation is initiated to trigger EB formation. Although developmental gene expression has been characterized for decades, the signals or events that initiate differentiation from one form to the other are not known.

In a 2013 study, another defining characteristic of EBs and RBs was identified: their relative levels of the second messenger signaling molecule cyclic di-AMP (c-di-AMP) (See ref. (14)). Barker et al. studied how IFNβ production is activated in cells infected with *C. trachomatis*. The authors identified a role for the innate immune response protein, STING, which recognizes c-di-AMP. *Chlamydia* encodes a diadenylate cyclase enzyme, DacA, associated with c-di-AMP production, that had not been characterized. As part of their study, the authors determined that c-di-AMP accumulates over the course of the chlamydial developmental cycle, that EBs have high levels of this molecule whereas RBs have low levels, and that DacA is a diadenylate cyclase. It is unlikely that *Chlamydia* produces c-di-AMP only to signal host immune responses. Rather, the parsimonious interpretation is that *Chlamydia* uses this signaling molecule to regulate some aspect of its physiology and that activation of host signaling is "accidental" – similar to activation of NOD2 by chlamydial peptidoglycan(15). However, no direct function of c-di-AMP in chlamydial biology has been defined.

Diverse functions of c-di-AMP have been reported in Gram-positive bacteria. For example, c-di-AMP is implicated in the response to changes in osmotic pressure. When the extracellular solute level is high, bacteria prevent dehydration by importing both extracellular solute molecules and cations(16). In these mechanisms, c-di-AMP binds to proteins associated with K^+ uptake systems such as Ktr/Trk, KimA, Kup, and Kdp and inhibits their activities in *Bacillus subtilis*, *Staphylococcus aureus*, and *Lactococcus lactis(17-20)*. Similarly, K^+ export mechanisms are also regulated by c-di-AMP. In *S. aureus*, the cation/proton antiporter A (CpaA) is activated by binding c-di-AMP(21). Moreover, c-di-AMP binds to the riboswitch upstream of the genes encoding the K^+ transporters and controls their transcriptional levels(22). With these mechanisms, osmotic stress is controlled by c-di-AMP. Of note, *Chlamydia* lacks annotated orthologs of K^+ transporters. In addition to osmotic homeostasis, c-di-AMP has been reported to affect DNA replication and sporulation(23). When DNA damage is detected, DisA,

a diadenylate cyclase, forms a DNA repair complex with RecA and RadA, resulting in inhibition of diadenylate cyclase activity in *B. subtilis*. Subsequently, c-di-AMP levels decrease, and DNA replication and sporulation are arrested(23-26).

Given the differences in c-di-AMP levels between EBs and RBs and its function as a diffusible second messenger signaling molecule, we hypothesized that the accumulation of c-di-AMP during the developmental cycle might be a trigger for secondary differentiation in *Chlamydia* (Fig. 1B). Once a threshold concentration of c-di-AMP has been reached in a given RB, it will begin the differentiation process to an EB. We predicted that, if our hypothesis were correct, then we should be able to alter the levels of c-di-AMP in the organism and affect its differentiation status accordingly. For example, if we increase c-di-AMP production, then we anticipate prematurely triggering EB production. Conversely, if we prevent c-di-AMP production, then we anticipate delaying EB production.

There are two principal mechanisms to regulate c-di-AMP levels: through synthesis by diadenylate cyclase and through degradation by a phosphodiesterase (PDE)(27). Interestingly, *C. trachomatis* only encodes the synthesis mechanism as it possesses the genes for diadenylate cyclase (*dacA*) and its regulator (*ybbR*) within a bicistronic operon, but no annotated PDEs(28). Therefore, to test our hypothesis, we genetically manipulated the levels of DacA and/or YbbR using recently developed strategies in the field. In the present study, we characterized the growth and developmental cycle state of chlamydial strains producing high or low levels of c-di-AMP. Our data show that higher levels of c-di-AMP are directly linked to increases in transcription of late genes associated with secondary differentiation as well as the concomitant production of EBs at an earlier stage in the developmental cycle. In contrast, in cells with reduced c-di-AMP levels, the developmental cycle was significantly delayed. Based on these data, we conclude that there is a threshold level of c-di-AMP necessary to trigger secondary differentiation in *C. trachomatis*. This is the first study to identify a physiological function for c-di-AMP in *Chlamydia* as well as a mechanism by which secondary differentiation is initiated in these unique bacteria.

Results

Both DacA and YbbR are crucial for cyclic di-AMP synthesis in *C. trachomatis*.

To test the link between c-di-AMP levels and chlamydial developmental cycle progression, we made a collection of *C. trachomatis* strains carrying anhydrotetracycline (aTc)-inducible constructs (Fig. 1C). These included strains to overexpress DacA_6xH, YbbR_6xH, both DacA and YbbR_6xH (*dacA*op), or both the catalytically dead DacA(D164N) and YbbR_6xH (*dacA*opMut). As a control for overexpression, we used a strain carrying the same plasmid backbone with aTc-inducible mCherry. In addition to the overexpression constructs, we also made a conditional knockdown construct for *dacA* (*dacA*-KD), targeting its promoter region, to decrease the level of c-di-AMP using a dCas12/crRNA-based system that our group developed for *Chlamydia* (29, 30). Finally, we made a complementation construct for *dacA*-KD by introducing *dacA-ybbR*_6xH (*dacA*-KD_*dacA*op) 3' to dCas12 such that induction of dCas12 results in the coexpression of DacA and YbbR_6xH as well.

Each of the transformant strains were used to infect HeLa cells at a multiplicity of infection (MOI) of 1 and subsequently induced or not with 5 nM aTc at 10 hours post-infection (hpi). STING KO cells were used to eliminate any potentially confounding effects of IFNβ production(31). We collected

cell lysates at 16 and 24 hpi and measured c-di-AMP levels by ELISA (Fig. 1D). The former timepoint is characterized by predominantly an RB population whereas the latter timepoint is characterized by ongoing secondary differentiation and a mixture of RBs, IBs, and EBs. We initially assessed the "empty vector" mCherry expressing control strain for its c-di-AMP levels and observed that c-di-AMP levels were significantly higher at 24hpi compared to 16hpi (Fig. 1D), in agreement with the Barker et al. study indicating higher levels of c-di-AMP in EBs(14).

Interestingly, c-di-AMP levels were comparable between the empty vector control strain and the DacA_6xH or YbbR_6xH overexpression strains, implying that expression of either alone does not alter endogenous c-di-AMP levels. However, when DacA and YbbR_6xH were co-overexpressed, c-di-AMP levels increased by approximately 30-fold at 16 hpi and 120-fold at 24 hpi compared to that of the control strain (Fig. 1D). These data indicate DacA activity requires YbbR as a positive regulator. Of note, the levels of c-di-AMP at 16hpi in the *dacA*op strain were higher even than the control strain at 24hpi. To confirm whether DacA enzyme activity is critical for the high levels of c-di-AMP in the *dacA*op strain, we substituted the catalytic residue of DacA with a mutation to render it inactive (D164N)(14). When DacA(D164N) and YbbR_6xH were co-overexpressed (*dacA*opMut), c-di-AMP levels were reduced compared to the control and did not increase between 16 and 24hpi, suggesting that c-di-AMP synthase activity was blocked in this strain. For the *dacA*-KD strain, the c-di-AMP level decreased about 1.2 fold at 16 hpi and ~10 fold at 24 hpi. The loss of c-di-AMP production in this strain was restored in the complemented *dacA*-KD_*dacA*op strain, which showed a phenotype similar to the *dacA*op overexpression strain. Based on these data, we conclude that both DacA and YbbR are crucial for c-di-AMP synthesis. Importantly, our collection of strains that are high or low producers of c-di-AMP give us an opportunity to test our overarching hypothesis (Fig. 1B).

Overexpression of DacA isoforms, but not YbbR, is detrimental to the chlamydial developmental cycle.

DacA and YbbR are both predicted to contain transmembrane domains (3 for DacA and 1 for YbbR)(Fig. S1, (14)). To observe the localization of DacA and YbbR, we infected HeLa cells with transformants encoding *dacA*_6xH or *ybbR*_6xH and induced the constructs at 10 hpi with 5 nM aTc. At 24 hpi, infected cells were fixed, and we performed an indirect immunofluorescence assay (IFA) by labeling the major outer membrane protein (MOMP) and 6xH. We observed that both DacA and YbbR localize at the membrane as expected (Fig. S2A and E). Interestingly, when DacA_6xH was overexpressed, the inclusion size appeared reduced with apparently larger individual organisms than those in the uninduced control. This suggests that overexpressing DacA_6xH alone may negatively impact chlamydial replication.

To investigate this in more detail, we measured EB progeny production and genomic DNA levels (a proxy for total bacteria, i.e., EB+RB). At 24 hpi, both EB progeny and genomic DNA levels were significantly decreased in the DacA_6xH overexpression strain compared to that of the uninduced control (Fig. S2B and C), indicating DacA_6xH overexpression disrupts developmental cycle progression. We performed the same experiments with the inactive isoform of DacA, DacA(D164N)_6xH. Like overexpression of the wild-type DacA_6xH, overexpression of DacA(D164N)_6xH also negatively affected the developmental cycle (Fig. S3). Based on these data, we conclude overexpressed DacA_6xH is detrimental to the chlamydial developmental cycle progression

even though it does not impact overall c-di-AMP production (Fig. 1D). To further investigate this, we
quantified transcripts by RT-qPCR for a canonical early-cycle gene, *euo*, and two late-cycle genes, *hctA*
and *omcB*. Consistent with the gDNA and EB data, overexpression of DacA_6xH resulted in elevated
euo transcripts and a reduction in the amounts of the late gene transcripts at 24hpi (Fig. S2D).

We also measured the effect of YbbR_6xH overexpression on the developmental cycle using
the same parameters. Unlike DacA_6xH, we measured no difference between uninduced and induced
samples for genomic DNA level, indicating no impact of YbbR_6xH overexpression on RB replication
or total bacterial numbers (Fig. S2G). YbbR_6xH overexpression did result in a statistically significant,
but not biologically significant, ~2-fold decrease in EB production (Fig. S2F). Therefore, we conclude
YbbR_6xH overexpression alone does not affect the chlamydial developmental cycle. This was further
supported by the absence of any effect of YbbR_6xH overexpression on transcript levels for *euo*, *hctA*,
or *omcB* (Fig. S2H).

A low level of c-di-AMP decreases the transcript levels of late genes.

As mentioned above, the *dacA*-KD strain showed a lower c-di-AMP level compared to that of
the control strain (Fig. 1D). To observe effects of knockdown on the developmental cycle, we examined
if there were defects in inclusion or bacterial morphology by IFA. When the knockdown system was
induced, bacterial cell size appeared somewhat enlarged, and we observed the inclusion size was smaller
compared to that of the uninduced control (Fig. 2A). In addition, we also measured the amount of EB
progeny and genomic DNA. Although genomic DNA levels were the same, EB progeny production
decreased by approximately 80% compared to that of the uninduced sample at 24 hpi (Fig. 2B and C).
Based on these data, we conclude that low levels of c-di-AMP are detrimental for secondary
differentiation. To further explore this, we performed RT-qPCR to quantify transcripts of relevant gene
targets. When dCas12 was induced, both *dacA* and *ybbR* transcript levels decreased (Fig. 2D). Since
dacA and *ybbR* are transcribed in an operon, this result is not surprising. Consistent with reduced EB
yields, transcripts of the late genes *omcB* and *hctA* were decreased at 24 hpi. In contrast, transcripts of
the early gene *euo* were slightly elevated at 24 hpi. We have previously observed no effects on gDNA
levels or transcription of these genes when overexpressing the dCas12 gene alone(30), indicating these
effects are specific to *dacA*-KD. These data support the conclusion that reducing the levels of c-di-AMP
delays developmental cycle progression.

We next tested whether the *dacA*-KD strain's phenotype is complemented by ectopic expression
of *dacA-ybbR_6xH* (*dacA*-KD_*dacA*op) (Fig. 1B). This construct encodes not only the *dacA*-knockdown
system but also the *dacA-ybbR_6xH* operon as a transcriptional fusion with dCas12. Induction of
dCas12 and YbbR_6xH was confirmed by IFA under inducing conditions (Fig. 2E). We expected that
co-expressing DacA and YbbR_6xH would complement the phenotype of *dacA*-KD. However, the
inclusion and bacterial size phenotypes were not fully restored to the uninduced control. We then
measured gDNA levels and EB progeny production and observed that the complemented strain showed
an approximate 2-fold reduction in both gDNA and EBs at 24 hpi (Fig. 2F&G). We next measured
transcript levels as described previously. Both *dacA* and *ybbR* transcripts were increased compared to
the *dacA*-KD strain (Fig. 2H), and transcript levels for these genes were increased beyond the "wild-type"
uninduced control levels, suggesting that the complemented strain is more similar to the *dacA*op strain
(Fig. 1B). Nonetheless, transcripts for the early gene *euo* were identical to those in the uninduced strain

(Fig. 2H). Surprisingly, *hctA* transcripts were increased ~10-fold at 14 hpi compared to that of the
uninduced sample, and *omcB* transcripts were slightly, but not significantly, increased at 14 and 24 hpi
(Fig. 2H). Based on these unusual findings of elevated late gene transcripts earlier in the developmental
cycle, we conclude that overexpressed DacA and YbbR_6xH may alter the timing of secondary
differentiation.

High levels of c-di-AMP induce late gene expression.

To clarify the effect of DacA and YbbR_6xH overexpression on secondary differentiation, we
next evaluated the phenotype of the *dacA*op strain. When we induced expression of this construct, we
confirmed the induction of DacA and YbbR_6xH and their colocalization at the membrane (Fig. 3A).
This is not surprising as both proteins are critical for c-di-AMP synthesis. Organism and inclusion
morphology was similar to the *dacA*-KD_*dacA*op complemented strain. We next assessed whether the
overexpressed DacA and YbbR_6xH affected replication by measuring gDNA levels. Again, the gDNA
data closely phenocopied the complemented knockdown strain, showing a decrease at 24 hpi (Fig. 3B).
We then quantified transcripts for the developmentally regulated genes *euo*, *hctA*, and *omcB* as well as
for *dacA* and *ybbR* (Fig. 3C). Not surprisingly, *dacA* and *ybbR* transcripts were elevated at the
timepoints assessed. Transcripts for *euo* were not statistically changed but trended higher at the 24 hpi
timepoint whereas *omcB* transcripts were slightly, but not significantly, increased at 14 and 24 hpi. Once
again, we observed that *hctA* transcripts were increased over 10-fold at 14hpi in the developmental cycle
(Fig. 3C), similar to what we measured for the complemented knockdown strain (Fig. 2H). These data
reinforce that elevated c-di-AMP levels (Fig. 1D) in these strains lead to increased expression of late
genes at an earlier timepoint in the developmental cycle.

To confirm whether these phenotypes are caused by a higher level of c-di-AMP, we compared
our results for the *dacA*op strain to the *dacA*opMut strain, in which the active site residue of DacA has
been mutated. The c-di-AMP levels in this strain remain low (Fig. 1D). The bacterial morphology
indicated slightly larger organisms without an effect on overall inclusion size (Fig. 3D). Genomic DNA
levels showed a significant drop at 24 hpi (Fig. 3E). In assessing transcripts, *dacA* and *ybbR* were
increased whereas *euo* levels were maintained, albeit not significantly so, at 24 hpi (Fig. 3F). In contrast
to the c-di-AMP overproducing strains, transcripts for *hctA* and *omcB* were unchanged or decreased at
14 and 24 hpi, respectively (Fig. 3F). The transcriptional results of the developmentally regulated genes
in the *dacA*opMut strain were very similar to the *dacA*-KD strain (Fig. 2D). These data suggest that the
inability to produce c-di-AMP delays developmental cycle progression.

Elevated c-di-AMP activates expression of genes necessary for secondary differentiation.

Given the surprising finding that, in strains overproducing c-di-AMP, *hctA* transcripts were 10-
fold higher at a timepoint not associated with secondary differentiation, we asked the question whether
all genes related to secondary differentiation were activated after inducing production of c-di-AMP.
Conversely, we wanted to explore whether reducing c-di-AMP levels would delay expression of genes
related to secondary differentiation. Thus, to further investigate how c-di-AMP affects transcription of
such genes, we performed RNA sequencing on both the *dacA*op overexpression and *dacA*-KD strains.
HeLa cells were infected with these transformants, and overexpression or knockdown was induced or
not at 10 hpi with 5 nM aTc. For the *dacA*op strain, RNA was collected at 16 hpi, a time at which late

genes are beginning to be expressed (as opposed to 14hpi) but remain near a basal level of
transcription(12). The rationale for this was to determine whether high c-di-AMP levels result in
increased late gene transcripts at this timepoint beyond the levels of the control, uninduced condition.
For the *dacA*-KD strain, RNA was collected at 24 hpi, a time at which late genes are peaking in their
expression. The rationale for this was to determine whether late gene transcription was repressed, which
could not otherwise be assessed at the 16 hpi timepoint.

RNA sequencing results were statistically analyzed by the UNMC Bioinformatics Core (Table
S2). Figure 4 shows a volcano plot of the results for the *dacA*op overexpression and *dacA*-KD strains.
Of note, many late genes were evident in the upregulated quadrant for the *dacAop* strain whereas the
converse was true for the *dacA*-KD strain. We further characterized the upregulated or downregulated
gene sets for the *dacA*op overexpression and *dacA*-KD strains, respectively, based on significant
difference (p<0.05) and fold-change (>1.5). A summary of these results is presented in Table 1 (all data
are presented in Table S3). We grouped the differentially expressed genes into five categories: 1)
canonical late genes for which the literature has associated them with EB function, 2) outer membrane
associated, 3) glycogen synthesis associated, 4) type III secretion system associated, and 5) gene
regulation associated. Recent work from our group and the Hefty group to define the regulons of the
alternative sigma factors in *Chlamydia* demonstrated that these sigma factors regulate some late gene
expression associated with outer membrane remodeling, type III secretion, and other processes (32, 33).
Consistent with our RT-qPCR data, we observed that all canonical late genes as well as all the other
genes listed in these categories showed an increase in expression after c-di-AMP production was
induced. In contrast, all the genes, and particularly the canonical late genes, showed a decrease in
expression under conditions where c-di-AMP production was impaired. Overall, these RNA sequencing
data confirm the direct influence of c-di-AMP on expression of genes related to secondary
differentiation.

Increased c-di-AMP production activates late gene transcription during penicillin treatment

We next asked whether late gene transcription could be activated by elevated c-di-AMP levels
even under conditions when developmental progression is blocked. Penicillin treatment of *Chlamydia*
blocks cell division and results in aberrantly enlarged RBs that fail to activate late gene transcription
with a concomitant reduction in EB progeny production. As our previously described *dacA*op
overexpression strain was constructed in a beta-lactamase producing background, we first generated a
spectinomycin-resistant *dacA*op overexpression strain and validated that induction of the *dacA* operon
resulted in earlier accumulation of *hctA* transcripts (Fig. S5). We next used this strain to infect cells and
induced overexpression or not with 5 nM aTc at 10 hpi. We also treated cells with 1 U/mL penicillin at
this timepoint. Penicillin treatment resulted in aberrantly enlarged RBs, as expected (Fig. 5A). However,
we did observe that overexpressing *dacA*op in the presence of penicillin negatively impacted overall
inclusion size as compared to the uninduced but penicillin-treated condition. Genomic DNA levels
similarly were decreased at 24hpi as compared to the uninduced control (Fig. 5B). Transcripts for *dacA*
and *ybbR* were increased after inducing their expression, and *euo* transcripts remained elevated at 24hpi
under these conditions. Interestingly, *hctA* transcripts in the uninduced but penicillin-treated cultures
remained low at 14hpi but increased by 24hpi whereas *omcB* transcripts remained low at all timepoints
under these conditions (Fig. 5C). Conversely, when *dacA*op is overexpressed in the presence of

penicillin, *hctA* transcripts increased from 10 to 24hpi, albeit not to the same extent as measured in the absence of penicillin. Transcripts for *omcB* similarly increased at 24hpi but were unchanged at 14hpi. These data suggest that, even under conditions when the developmental cycle is blocked, c-di-AMP can activate late gene transcription.

Alterations in c-di-AMP levels impact the timing of EB production

The phenotypic effect of increased late gene transcription should be an increase in EB production. However, if EB production is initiated at an earlier time in the developmental cycle when there are fewer non-infectious RBs to convert, then overall EB production should be decreased. Conversely, delayed expression of late genes should be associated with a delay in EB production. Therefore, to test these predictions, we quantified EB production at 2 h intervals from 18 to 24hpi, to assess EB production during earlier phases of the developmental cycle, as well as at 32 and 48hpi, to assess overall EB yields (Fig. 6). We did not detect any EBs at 16hpi or earlier (not shown). Cells were infected with the *dacA*op, *dacA*opMut, and *dacA*-KD strains and induced or not at 10 hpi with 5 nM aTc as previously noted. Consistent with the transcriptional data, we measured higher EB yields at 18 and 20 hpi during *dacA*op overexpression that quickly plateaued by 24 hpi (Fig. 6A). Conversely, when c-di-AMP production was blocked by reducing the expression of DacA and YbbR (*dacA*-KD), we observed delayed EB production (Fig. 6B). We also measured lower EB yields when c-di-AMP levels were decreased by overexpression of the DacA(D164N) isoform and YbbR (*dacA*opMut; Fig. 6C). The uninduced control conditions for each strain showed a steady accumulation of EBs throughout the course of the experiment, as expected. From these data, we conclude that earlier production of c-di-AMP levels results in earlier production of EBs.

Discussion

Secondary differentiation is essential for the propagation and survival of *Chlamydia* species. However, the mechanisms governing this essential process are largely unknown. It is possible that both internal and external environmental changes might serve as signals to trigger the shift from the non-infectious RB to the infectious EB. It is also possible, and even likely, that *Chlamydia* integrates multiple signal inputs during the differentiation process. However, as secondary differentiation is asynchronous, any mechanism must account for the stochasticity within the population of RBs "considering" differentiating to EBs. Here, we provide the first evidence of a signal that can directly activate EB-associated gene expression with a concomitant early production of EBs.

Several reports have proposed mechanisms by which *Chlamydia* triggers secondary differentiation. For example, Thompson et al. (2015) investigated the non-canonical regulation of the major sigma factor, σ^{66}, by the Rsb phosphoregulatory system(34). The authors provided evidence that altering the levels of RsbW or RsbV1 impacted the expression of σ^{66}-controlled genes. This led them to propose a model whereby sensing of ATP by the Rsb system controls the availability of σ^{66}, implying that low levels of ATP lead to initiation of secondary differentiation. Further support from this came from work by Kuwabara et al. (2022) who showed effects of ATP, GTP, and glucose on the activity of different Rsb components(35). Finally, work from Soules et al. (2020a) identified TCA intermediates as ligands for RsbU, linking the TCA cycle and ATP synthesis to the Rsb system(36). However, no effect was noted on premature late gene expression in these experimental systems.

Sequestration/inactivation of the major sigma factor under low ATP conditions would presumably render RNA polymerase free to interact with the alternative sigma factors, σ^{54} and σ^{28}. These alternative sigma factors have been linked to late gene expression with studies from Soules et al. (2020b) and Hatch and Ouellette indicating that σ^{54} is associated with increased transcription of outer membrane components, type III secretion (T3S) system components, and other genes typically expressed late in development(32, 33). However, some of these changes are likely indirect and downstream to the σ^{54} regulon itself since some affected genes have been shown to have σ^{66} promoter sequence elements(37, 38). Significant remodeling of the outer membrane occurs as the RB transitions to the EB(39), and, similarly, the EB is prepackaged with T3S effectors, like TarP(40), that facilitate EB invasion into a target host cell. σ^{28} was shown to control expression of two canonical late genes: *hctB* and *tsp*(32, 41). Each of the encoded proteins is highly toxic if overexpressed(42, 43), suggesting that the additional layer of regulation by σ^{28} is necessary to ensure EBs are formed at the correct time. Hatch and Ouellette proposed that these genes may be the last to be activated precisely because of this(32). Overall, these data clearly link transcriptional regulation, via the sigma factors, to developmental cycle progression and secondary differentiation.

Despite the clear changes in transcription that occur during the chlamydial developmental cycle, activation of late gene expression alone does not guarantee secondary differentiation. We recently explored the function of the Clp protease systems in chlamydial growth and development(44). Interestingly, we observed that the inability to degrade SsrA-tagged products prevented functional secondary differentiation with a severe defect in EB production. This was demonstrated in strains unable to degrade SsrA products either by preventing their recognition by ClpX or by altering the SsrA tag to a version that is not degraded efficiently. Nonetheless, late gene transcription was activated in these strains. Therefore, post-translational regulation of secondary differentiation is also critical to the process.

Assuming a post-translational gene regulation model for secondary differentiation, we were interested in exploring known differences between EBs and RBs as a clue to what factors could be serving as a signal for this process. EBs and RBs differ in a number of characteristics, from their function to their morphology to their gene expression. In 2013, a study from Barker et al. added another difference – their relative levels of c-di-AMP with RBs having lower levels than EBs. Although the study from Barker et al. was focused on understanding how IFNβ is activated in *Chlamydia*-infected cells, we were intrigued by this relative difference in c-di-AMP levels. We hypothesized that c-di-AMP production might be a signal for secondary differentiation. We compared chlamydial growth in normal HeLa or STING-KO HeLa cells but could not detect any differences in growth in these cell types (Fig. 2 and S2 vs Fig. S6), indicating that c-di-AMP production by *Chlamydia* is unlikely used as means of activating the host cell. Rather, STING activation is "accidental". Using recently developed genetic tools for *Chlamydia*, we demonstrated that we could manipulate the levels of c-di-AMP in the bacterium to either block or stimulate its production. Excitingly, we observed that elevated c-di-AMP was linked to earlier secondary differentiation whereas blocking c-di-AMP production prevented EB production (Fig. 6). Importantly, c-di-AMP production resulted in the activation of late gene transcripts as noted above for the alternative sigma factors. Therefore, c-di-AMP acts as a post-translational mechanism to trigger secondary differentiation. To our knowledge, this is the first study to identify and provide experimental evidence for a signaling factor involved in this process.

How might c-di-AMP function in *Chlamydia*? This is not readily apparent because *Chlamydia* lacks orthologs of proteins that have been characterized in other bacterial systems to bind c-di-AMP. For example, *Chlamydia* lacks annotated K^+ transporters, and it is unknown how these bacteria maintain their osmolarity. However, a recent study suggested that changes in K^+ levels affect the chlamydial developmental cycle(45). Therefore, c-di-AMP may function as a K^+ homeostasis coordinator in *Chlamydia* through as-yet unknown pathways. We did observe altered chlamydial morphology in some of our strains in which we altered c-di-AMP levels, and this may indicate changes in osmostability or effects on cell division. In regard to the latter possibility, in *S. aureus*, DacA activity is inhibited by interacting with GlmM (phosphoglucosamine mutase)(46). GlmM produces UDP-GlcNAc that is a precursor in both peptidoglycan and LPS synthesis. Given that *Chlamydia* uses peptidoglycan for cell division, we tested whether DacA and YbbR were localized to the division septum. However, their localization was not associated with the division septum (Fig. S4). In addition, we did not detect an interaction between DacA and GlmM using the Bacterial Adenylate Cyclase-based Two Hybrid (BACTH) system (data not shown), and the chlamydial GlmM lacks the residues required for DacA interaction. Therefore, we conclude that DacA and YbbR are unlikely to directly impact cell division. However, based on the cell morphological changes in DacA or YbbR overexpression, we cannot exclude a function for c-di-AMP on cell wall metabolism. Future studies will focus on identifying c-di-AMP binding proteins and characterizing their function in *Chlamydia*.

The use of c-di-AMP as a signal for secondary differentiation presents a "chicken-or-egg" quandary. How is the activity and expression of DacA, the diadenylate cyclase, regulated? Our transcript analyses indicate that *dacA-ybbR* transcripts peak during the RB phase of growth, similar to most genes in *Chlamydia*. One possibility not easily tested due to the obligate intracellular nature of *Chlamydia*, is that c-di-AMP production depletes ATP levels, which then reduce σ^{66} activity as described above. Our data indicate that YbbR is a positive regulator of DacA activity since expressing DacA alone had no impact on overall c-di-AMP levels (Fig. 1D). Therefore, YbbR itself may be a target for regulation. As a monotopic transmembrane protein, YbbR may be degraded by proteases such as the inner membrane-associated FtsH or the periplasmic proteases HtrA or Tsp. This would presumably shutdown c-di-AMP synthesis. Further work is needed to test these possibilities.

Even though diverse functions of c-di-AMP in other bacteria have been reported previously, this is the first time c-di-AMP has been described as a checkpoint in chlamydial development. It is possible that the levels of c-di-AMP act as a de facto means for monitoring the bacterial population within the inclusion or for their overall developmental status. Canonical bacteria can use quorum sensing to detect bacteria at the population level and to communicate with other species(47). However, *C. trachomatis* lacks homologues of genes related to quorum sensing(28). In addition, secondary differentiation occurs asynchronously, and this feature is different from a quorum sensing mechanism. Our model accounts for the asynchronicity of secondary differentiation. As RBs accumulate c-di-AMP and divide, it is likely the c-di-AMP will be distributed unevenly between the mother and daughter cell since *Chlamydia* divides through an asymmetric budding mechanism(7) (Fig. 7). This will lead to two cells with different levels of c-di-AMP, one of which may then accumulate enough c-di-AMP to trigger secondary differentiation while the other continues to divide (Fig. 7). This is consistent with a recent model proposing that a population of division-competent RBs is maintained throughout the developmental cycle(48).

Other ongoing studies in the lab have revealed additional post-translational mechanisms that drive secondary differentiation (Jensen et al. *in revision*; (49)), thus it seems likely that *Chlamydia* integrates multiple signal inputs to ensure that all conditions are met before undergoing this essential step in their biology.

Materials and Methods

Organisms and Cell Culture

Wild-type (ATCC, Manassas, VA) and STING KO (Dr. Frank van Kuppeveld; Utrecht University) HeLa, human cervical epithelial-derived, and McCoy (kind gift of Dr. Harlan Caldwell, NIH), mouse fibroblast-derived, cell lines were cultured at 37°C with 5% CO_2 in Dulbecco's Modified Eagle Medium (DMEM; Invitrogen, Waltham, MA) containing 10% fetal bovine serum (FBS; Hyclone, Logan, UT) and 10 μg/mL gentamicin (Gibco, Waltham, MA). *C. trachomatis* serovar L2 (434/Bu) lacking the endogenous plasmid (-pL2; kind gift of Dr. Ian Clarke, Univ. Southampton) was used for transformation. All cell cultures and chlamydial stocks were routinely tested for Mycoplasma contamination using the Mycoplasma PCR detection kit (Sigma, St. Louis, MO). For *E. coli*, NEB10β competent cells (New England Biolabs, Ipswich, MA) were used for the amplification of pBOMB-derivative vectors. *E. coli* was grown at 30°C in LB media. All chemicals and antibiotics were obtained from Sigma unless otherwise noted.

Cloning

The list of the vectors and primers used in this study is detailed in Supplemental Table 1. Target genes were amplified by PCR with Phusion DNA polymerase (NEB) using 10 ng *C. trachomatis* L2 genomic DNA or appropriate vectors as a template. Some DNA segments were directly synthesized as a gBlock fragment (Integrated DNA Technologies, Coralville, IA). If plasmids were used as a template, then we treated the PCR product with DpnI enzyme to remove templates. The PCR products were purified using a PCR purification kit (Qiagen, Hilden, Germany). The HiFi Assembly reaction master mix (New England Biolabs (NEB), Ipswich, MA) was used following the manufacturer's manual in conjunction with plasmids pBOMBL(30) linearized with EagI and KpnI or pBOMBL12CRia (empty vector) linearized with BamHI. The linearized plasmids were also dephosphorylated with FastAP (ThermoFisher). The products of the HiFi reaction were transformed into NEB10β competent cells (NEB) and plated on LB agar with appropriate antibiotics. Plasmids were subsequently isolated using a mini-prep kit (Qiagen) and verified by plasmid digest and sequencing from individual colonies grown overnight in LB broth with appropriate antibiotic selection.

Transformation of *Chlamydia trachomatis*

McCoy cells were plated in a six-well plate the day before beginning the transformation procedure. *C. trachomatis* serovar L2 without plasmid (-pL2) resuspended with Tris-CaCl$_2$ buffer (10 mM Tris-Cl pH 7.5, 50 mM CaCl$_2$) was incubated with 2 μg plasmid at room temperature for 30 min. During this step, McCoy cells were washed with 2 mL Hank's Balanced Salt Solution (HBSS) media containing Ca^{2+} and Mg^{2+} (Gibco). After that, McCoy cells were infected with the transformants in 2 mL HBSS per well. The plate was centrifuged at 400 x g for 15 min at room temperature and incubated at 37°C for 15 min. The inoculum was aspirated, and 2 mL 1X DMEM containing 10% FBS and 10 μg/mL gentamicin was

added per well. At 8 hours post infection (hpi), 1 μg/mL cycloheximide and either 1 or 2 U/mL penicillin G were added, and the plate was incubated at 37˚C until 48 hpi. At 48 hpi, the transformants were harvested and infected onto a new McCoy cell monolayer. These harvest and infection steps were repeated every 44-48 hpi until fluorescent, penicillin-resistant inclusions were observed.

(RT-)qPCR

HeLa cells were infected with chlamydial transformants at an MOI of 0.5. At 10 hpi, 5 nM anhydrotetracycline (aTc) was added or not to the culture medium. Total RNA and DNA was harvested at this time point from duplicate wells not treated with aTc. At 14 and 24 hpi, total RNA and DNA were collected using Trizol (Invitrogen) and DNeasy Tissue (Qiagen) kit, respectively, as described elsewhere(32). After DNase treatment of total RNA, cDNA was synthesized using Superscript III reverse transcriptase (Invitrogen). After diluting the cDNA 10-fold, 5 μl of the diluted cDNA was used as a template for qPCR. Equal masses of genomic DNA were used from each of the samples to quantify chlamydial genomes, which were used to normalize transcript data as described(37). For both cDNA and gDNA samples, qPCR reactions were prepared using 2X SYBR Green (ThermoFisher) in a total volume of 25uL per well. Standard cycling conditions were used with a melting curve analysis to verify products. Transcripts and genome copies were assessed from at least three biological replicates.

Indirect immunofluorescence assay (IFA)

HeLa cells were infected with chlamydial transformants as above. At 10 hpi, 5 nM aTc was added or not, and the infected cells were fixed with fixing solution (3.2% formaldehyde and 0.022% glutaraldehyde in 1X DPBS) for 2 min and permeabilized with 90% MeOH for 1 min at 10.5 hpi or 24 hpi. The fixed cells were labeled with primary antibodies including rabbit anti-DacA (custom anti-peptide antibody prepared by Pacific Immunology, Ramona, CA), goat anti-major outer-membrane protein (MOMP; Meridian, Memphis, TN), and rabbit anti-six histidine tag (Genscript, Piscataway, NJ and Abcam, Cambridge, UK, respectively). To visualize the primary antibodies, donkey anti-goat antibody (488), donkey anti-mouse (405), or donkey anti-rabbit antibody (594) were used as secondary antibodies. The secondary antibodies were obtained from Invitrogen or Jackson Immunology (West Grove, PA). Coverslips were observed using a Zeiss AxioImager.Z2 with Apotome2 as noted in the figure legends.

Inclusion forming unit (IFU) measurement

HeLa cells were infected with chlamydial transformants as above. At 10 hpi, 5 nM aTc was added or not to the culture medium. At the indicated times, infected cells were harvested in 1 mL 2SP media then frozen at -80˚C. After thawing the lysates, the samples were serially 1:10 diluted and used to infect HeLa cells seeded in 24-well plates. At 24 hpi, the number of GFP expressing inclusions were counted from 30 fields of view to calculate the IFUs from the original sample. Three biological replicates were performed.

Cyclic di-AMP measurement

HeLa cells were infected with the indicated chlamydial transformants as above. At 10 hpi, expression of the constructs was induced or not with 5 nM aTc, and, at 16 and 24 hpi, samples were prepared for measuring the level of cyclic di-AMP. After aspirating the media, the infected cell monolayers were washed with 1X PBS and resuspended with B-Per Bacterial Cell Lysis Buffer (Thermo). Samples were

495 vortexed for 1 min and then centrifuged at 13,300xg for 15 min at 4℃. The levels of cyclic di-AMP
496 from the supernatants of the cell lysates were quantified using the Cyclic di-AMP ELISA kit (Cayman)
497 following the manufacturer's instructions(50).

Preparation of RNA sequencing samples

We prepared the samples as reported previously(32). Briefly, the transformants of *dacA*op and *dacA*-KD
were infected into HeLa cells. At 10 hpi, the constructs were induced or not with 5 nM aTc. RNA
samples were prepared from *dacA*op-infected cells and *dacA*-KD-infected cells at 16 hpi and 24 hpi,
respectively. 20 μg RNA samples were treated with DNase to remove DNA contamination by using
DNA-free Turbo kit (Thermo) according to the manufacturer's instruction. Ribosomal RNA was
depleted from samples using the MICROBEnrich (Thermo) and MICROBExpress kits (Thermo)
following the manufacturer's instructions. RNA samples were processed for sequencing by the UNMC
Genomics Core Facility. The resultant libraries from the individual samples were multiplexed and
subjected to 100-bp paired-read sequencing to generate approximately 60 million pairs of reads per
sample on an Illumina NovaSeq6000 sequencer in the UNMC Genomics Core facility. The original
fastq format reads were trimmed by fqtirm tool (https://ccb.jhu.edu/software/fqtrim) to remove adapters,
terminal unknown bases (Ns) and low quality 3' regions (Phred score < 30). The trimmed fastq files
were processed by FastQC (Andrews S. (2010). FastQC: a quality control tool for high throughput
sequence data; available online at http://www.bioinformatics.babraham.ac.uk/projects/fastqc for quality
control. *Chlamydia trachomatis* 434/Bu bacterial reference genome and annotation files were
downloaded from Ensembl (http://bacteria.ensembl.org/Chlamydia_trachomatis_434_bu/Info/Index).
Sequencing data were analyzed by the Bioinformatics and Systems Biology Core (BSBC). The trimmed
fastq files were mapped to *Chlamydia trachomatis* 434/Bu by CLC Genomics Workbench 23 for
RNAseq analyses.

Data availability

519 The raw and processed RNA sequencing reads in fastq format have been deposited in the Gene
520 Expression Omnibus (GEO; www.ncbi.nlm.nih.gov/geo/) under accession no. GSE252732.

Statistical Analysis

522 To analyze the statistical significance between uninduced and induced samples of qPCR and IFU data,
523 we used two sample equal variance t-test. For the levels of cyclic di-AMP data, we used a ratio paired t-
524 test.

Data and materials availability

526 All data are available in the main text or the supplementary materials.

Acknowledgements

529 This study was supported in part by an NIH/NIGMS award (R35GM124798) to SPO and by start-up
530 funds from UNMC. The UNMC Genomics Core Facility receives partial support from the National
531 Institute for General Medical Science (NIGMS) INBRE - P20GM103427-19, as well as the National

Cancer Institute the Fred & Pamela Buffett Cancer Center Support Grant- P30CA036727. This publication's contents are the sole responsibility of the authors and do not necessarily represent the official views of the NIH or NIGMS. The authors would like to thank Dr. Frank van Kuppeveld (Utrecht University) for STING-KO HeLa cells, Dr. Harlan Caldwell (NIAID/NIH) for McCoy cells, and Dr. Ian Clarke (University of Southampton) for the plasmidless *C. trachomatis* serovar L2 strain. In addition, we thank the UNMC Genomics Core Facility and the Bioinformatics and Systems Biology Core facility for RNA sequencing and analysis services, respectively.

Author contributions:

Junghoon Lee, Methodology, Investigation, Visualization, Writing – original draft, and Writing-review & editing

Scot P. Ouellette, Conceptualization, Methodology, Investigation, Visualization, Supervision, and Writing-review & editing

Competing interests: Authors declare that they have no competing interests.

References

1. Abdelrahman YM, Belland RJ. 2005. The chlamydial developmental cycle. FEMS Microbiol Rev 29:949-59.
2. Everett KD, Hatch TP. 1991. Sequence analysis and lipid modification of the cysteine-rich envelope proteins of Chlamydia psittaci 6BC. J Bacteriol 173:3821-30.
3. Hackstadt T, Baehr W, Ying Y. 1991. Chlamydia trachomatis developmentally regulated protein is homologous to eukaryotic histone H1. Proc Natl Acad Sci U S A 88:3937-41.
4. Moulder JW. 1993. Why is Chlamydia sensitive to penicillin in the absence of peptidoglycan? Infect Agents Dis 2:87-99.
5. Fox A, Rogers JC, Gilbart J, Morgan S, Davis CH, Knight S, Wyrick PB. 1990. Muramic acid is not detectable in Chlamydia psittaci or Chlamydia trachomatis by gas chromatography-mass spectrometry. Infect Immun 58:835-7.
6. Barbour AG, Amano K, Hackstadt T, Perry L, Caldwell HD. 1982. Chlamydia trachomatis has penicillin-binding proteins but not detectable muramic acid. J Bacteriol 151:420-8.
7. Abdelrahman Y, Ouellette SP, Belland RJ, Cox JV. 2016. Polarized Cell Division of Chlamydia trachomatis. PLoS Pathog 12:e1005822.
8. Liechti G, Kuru E, Packiam M, Hsu YP, Tekkam S, Hall E, Rittichier JT, VanNieuwenhze M, Brun YV, Maurelli AT. 2016. Pathogenic Chlamydia Lack a Classical Sacculus but Synthesize a Narrow, Mid-cell Peptidoglycan Ring, Regulated by MreB, for Cell Division. PLoS Pathog 12:e1005590.
9. Ouellette SP, Karimova G, Subtil A, Ladant D. 2012. Chlamydia co-opts the rod shape-determining proteins MreB and Pbp2 for cell division. Mol Microbiol 85:164-78.
10. Lee J, Cox JV, Ouellette SP. 2020. Critical Role for the Extended N-Terminus of Chlamydial MreB in Directing Its Membrane Association and Potential Interaction with Divisome Proteins. J Bacteriol doi:10.1128/JB.00034-20.
11. Shaw JH, Grund VR, Durling L, Caldwell HD. 2001. Expression of genes encoding Th1 cell-activating cytokines and lymphoid homing chemokines by chlamydia-pulsed dendritic cells

correlates with protective immunizing efficacy. Infect Immun 69:4667-72.

12. Belland RJ, Zhong G, Crane DD, Hogan D, Sturdevant D, Sharma J, Beatty WL, Caldwell HD.
2003. Genomic transcriptional profiling of the developmental cycle of Chlamydia trachomatis.
Proc Natl Acad Sci U S A 100:8478-83.

13. Ouellette SP, Hatch TP, AbdelRahman YM, Rose LA, Belland RJ, Byrne GI. 2006. Global
transcriptional upregulation in the absence of increased translation in Chlamydia during
IFNgamma-mediated host cell tryptophan starvation. Mol Microbiol 62:1387-401.

14. Barker JR, Koestler BJ, Carpenter VK, Burdette DL, Waters CM, Vance RE, Valdivia RH. 2013.
STING-dependent recognition of cyclic di-AMP mediates type I interferon responses during
Chlamydia trachomatis infection. mBio 4:e00018-13.

15. Girardin SE, Boneca IG, Viala J, Chamaillard M, Labigne A, Thomas G, Philpott DJ, Sansonetti
PJ. 2003. Nod2 is a general sensor of peptidoglycan through muramyl dipeptide (MDP) detection.
J Biol Chem 278:8869-72.

16. Booth IR, Higgins CF. 1990. Enteric bacteria and osmotic stress: intracellular potassium
glutamate as a secondary signal of osmotic stress? FEMS Microbiol Rev 6:239-46.

17. Kim H, Youn SJ, Kim SO, Ko J, Lee JO, Choi BS. 2015. Structural Studies of Potassium
Transport Protein KtrA Regulator of Conductance of K+ (RCK) C Domain in Complex with
Cyclic Diadenosine Monophosphate (c-di-AMP). J Biol Chem 290:16393-402.

18. Gibhardt J, Hoffmann G, Turdiev A, Wang M, Lee VT, Commichau FM. 2019. c-di-AMP assists
osmoadaptation by regulating the Listeria monocytogenes potassium transporters KimA and
KtrCD. J Biol Chem 294:16020-16033.

19. Quintana IM, Gibhardt J, Turdiev A, Hammer E, Commichau FM, Lee VT, Magni C, Stulke J.
2019. The KupA and KupB Proteins of Lactococcus lactis IL1403 Are Novel c-di-AMP
Receptor Proteins Responsible for Potassium Uptake. J Bacteriol 201.

20. Moscoso JA, Schramke H, Zhang Y, Tosi T, Dehbi A, Jung K, Grundling A. 2016. Binding of
Cyclic Di-AMP to the Staphylococcus aureus Sensor Kinase KdpD Occurs via the Universal
Stress Protein Domain and Downregulates the Expression of the Kdp Potassium Transporter. J
Bacteriol 198:98-110.

21. Corrigan RM, Campeotto I, Jeganathan T, Roelofs KG, Lee VT, Grundling A. 2013. Systematic
identification of conserved bacterial c-di-AMP receptor proteins. Proc Natl Acad Sci U S A
110:9084-9.

22. Wang X, Cai X, Ma H, Yin W, Zhu L, Li X, Lim HM, Chou SH, He J. 2019. A c-di-AMP
riboswitch controlling kdpFABC operon transcription regulates the potassium transporter system
in Bacillus thuringiensis. Commun Biol 2:151.

23. Bejerano-Sagie M, Oppenheimer-Shaanan Y, Berlatzky I, Rouvinski A, Meyerovich M, Ben-
Yehuda S. 2006. A checkpoint protein that scans the chromosome for damage at the start of
sporulation in Bacillus subtilis. Cell 125:679-90.

24. Oppenheimer-Shaanan Y, Wexselblatt E, Katzhendler J, Yavin E, Ben-Yehuda S. 2011. c-di-
AMP reports DNA integrity during sporulation in Bacillus subtilis. EMBO Rep 12:594-601.

25. Zhang L, He ZG. 2013. Radiation-sensitive gene A (RadA) targets DisA, DNA integrity
scanning protein A, to negatively affect cyclic Di-AMP synthesis activity in Mycobacterium
smegmatis. J Biol Chem 288:22426-36.

26. Gandara C, de Lucena DKC, Torres R, Serrano E, Altenburger S, Graumann PL, Alonso JC.
2017. Activity and in vivo dynamics of Bacillus subtilis DisA are affected by RadA/Sms and by
Holliday junction-processing proteins. DNA Repair (Amst) 55:17-30.

27. Corrigan RM, Grundling A. 2013. Cyclic di-AMP: another second messenger enters the fray.

Nat Rev Microbiol 11:513-24.

28. Stephens RS, Kalman S, Lammel C, Fan J, Marathe R, Aravind L, Mitchell W, Olinger L, Tatusov RL, Zhao Q, Koonin EV, Davis RW. 1998. Genome sequence of an obligate intracellular pathogen of humans: Chlamydia trachomatis. Science 282:754-9.

29. Ouellette SP. 2018. Feasibility of a Conditional Knockout System for Chlamydia Based on CRISPR Interference. Front Cell Infect Microbiol 8:59.

30. Ouellette SP, Blay EA, Hatch ND, Fisher-Marvin LA. 2021. CRISPR Interference to Inducibly Repress Gene Expression in Chlamydia trachomatis. Infection and Immunity *in press*.

31. Sui H, Zhou M, Imamichi H, Jiao X, Sherman BT, Lane HC, Imamichi T. 2017. STING is an essential mediator of the Ku70-mediated production of IFN-lambda1 in response to exogenous DNA. Sci Signal 10.

32. Hatch ND, Ouellette SP. 2023. Identification of the alternative sigma factor regulons of Chlamydia trachomatis using multiplexed CRISPR interference. mSphere 8:e0039123.

33. Soules KR, LaBrie SD, May BH, Hefty PS. 2020. Sigma 54-Regulated Transcription Is Associated with Membrane Reorganization and Type III Secretion Effectors during Conversion to Infectious Forms of Chlamydia trachomatis. mBio 11.

34. Thompson CC, Griffiths C, Nicod SS, Lowden NM, Wigneshweraraj S, Fisher DJ, McClure MO. 2015. The Rsb Phosphoregulatory Network Controls Availability of the Primary Sigma Factor in Chlamydia trachomatis and Influences the Kinetics of Growth and Development. PLoS Pathog 11:e1005125.

35. Kuwabara S, Landers ER, Fisher DJ. 2022. Impact of nutrients on the function of the chlamydial Rsb partner switching mechanism. Pathog Dis 80.

36. Soules KR, Dmitriev A, LaBrie SD, Dimond ZE, May BH, Johnson DK, Zhang Y, Battaile KP, Lovell S, Hefty PS. 2020. Structural and ligand binding analyses of the periplasmic sensor domain of RsbU in Chlamydia trachomatis support a role in TCA cycle regulation. Mol Microbiol 113:68-88.

37. Ouellette SP, Abdelrahman YM, Belland RJ, Byrne GI. 2005. The Chlamydia pneumoniae type III secretion-related lcrH gene clusters are developmentally expressed operons. J Bacteriol 187:7853-6.

38. Hefty PS, Stephens RS. 2007. Chlamydial type III secretion system is encoded on ten operons preceded by sigma 70-like promoter elements. J Bacteriol 189:198-206.

39. Zhong J, Douglas AL, Hatch TP. 2001. Characterization of integration host factor (IHF) binding upstream of the cysteine-rich protein operon (omcAB) promoter of Chlamydia trachomatis LGV serovar L2. Mol Microbiol 41:451-62.

40. Clifton DR, Fields KA, Grieshaber SS, Dooley CA, Fischer ER, Mead DJ, Carabeo RA, Hackstadt T. 2004. A chlamydial type III translocated protein is tyrosine-phosphorylated at the site of entry and associated with recruitment of actin. Proc Natl Acad Sci U S A 101:10166-71.

41. Yu HH, Kibler D, Tan M. 2006. In silico prediction and functional validation of sigma28-regulated genes in Chlamydia and Escherichia coli. J Bacteriol 188:8206-12.

42. Swoboda AR, Wood NA, Saery EA, Fisher DJ, Ouellette SP. 2023. The Periplasmic Tail-Specific Protease, Tsp, Is Essential for Secondary Differentiation in Chlamydia trachomatis. J Bacteriol 205:e0009923.

43. Grieshaber NA, Chiarelli TJ, Appa CR, Neiswanger G, Peretti K, Grieshaber SS. 2022. Translational gene expression control in Chlamydia trachomatis. PLoS One 17:e0257259.

44. Wood NA, Swoboda AR, Blocker AM, Fisher DJ, Ouellette SP. 2022. Tag-Dependent Substrate Selection of ClpX Underlies Secondary Differentiation of Chlamydia trachomatis. mBio

13:e0185822.

45. Andrew SC, Dumoux M, Hayward RD. 2021. Chlamydia Uses K(+) Electrical Signalling to Orchestrate Host Sensing, Inter-Bacterial Communication and Differentiation. Microorganisms 9.

46. Tosi T, Hoshiga F, Millership C, Singh R, Eldrid C, Patin D, Mengin-Lecreulx D, Thalassinos K, Freemont P, Grundling A. 2019. Inhibition of the Staphylococcus aureus c-di-AMP cyclase DacA by direct interaction with the phosphoglucosamine mutase GlmM. PLoS Pathog 15:e1007537.

47. Miller MB, Bassler BL. 2001. Quorum sensing in bacteria. Annu Rev Microbiol 55:165-99.

48. Chiarelli TJ, Grieshaber NA, Appa C, Grieshaber SS. 2023. Computational Modeling of the Chlamydial Developmental Cycle Reveals a Potential Role for Asymmetric Division. mSystems 8:e0005323.

49. Singh V, and Ouellette, S. P. . 2024. Altering the redox status of Chlamydia trachomatis directly impacts its developmental cycle progression. BioRxiv doi:https://doi.org/10.1101/2024.04.26.591247.

50. Underwood AJ, Zhang Y, Metzger DW, Bai G. 2014. Detection of cyclic di-AMP using a competitive ELISA with a unique pneumococcal cyclic di-AMP binding protein. J Microbiol Methods 107:58-62.

51. Tsirigos KD, Peters C, Shu N, Kall L, Elofsson A. 2015. The TOPCONS web server for consensus prediction of membrane protein topology and signal peptides. Nucleic Acids Res 43:W401-7.

52. Fahr MJ, Douglas AL, Xia W, Hatch TP. 1995. Characterization of late gene promoters of Chlamydia trachomatis. J Bacteriol 177:4252-60.

53. Jury B, Fleming C, Huston WM, Luu LDW. 2023. Molecular pathogenesis of Chlamydia trachomatis. Front Cell Infect Microbiol 13:1281823.

54. Yang C, Kari L, Sturdevant GL, Song L, Patton MJ, Couch CE, Ilgenfritz JM, Southern TR, Whitmire WM, Briones M, Bonner C, Grant C, Hu P, McClarty G, Caldwell HD. 2017. Chlamydia trachomatis ChxR is a transcriptional regulator of virulence factors that function in in vivo host-pathogen interactions. Pathog Dis 75.

55. Nicholson TL, Olinger L, Chong K, Schoolnik G, Stephens RS. 2003. Global stage-specific gene regulation during the developmental cycle of Chlamydia trachomatis. J Bacteriol 185:3179-89.

697   Table 1. Late genes regulated by cyclic di-AMP levels.

			Canonical late genes	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.
CTL0112	CT743	*hctA*	Histone H1-like protein HC1	6.70	-3.47	(12, 52)
CTL0302	CT046	*hct2*	Histone H1-like protein HC2	3.52	-5.96	(12, 41)
CTL0336	CT080	*ltuB*	Late transcription unit B protein	4.00	-3.61	(12, 52)
CTL0700	CT441	*tsp*	Carboxy-terminal processing protease	6.18	-6.62	(41)
CTL0702	CT443	*omcB*	Large cysteine-rich periplasmic protein	5.00	-4.04	(12, 33)
CTL0703	CT444	*omcA*	Small cysteine-rich outer membrane protein	5.93	-3.71	(12, 33)
CTL0716	CT456	*tarp*	Translocated actin-recruiting phosphoprotein	6.06	-5.97	(33)
			Membrane organization associated	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.
CTL0082	CT713	*ompB*	Outer membrane protein B	2.73	-2.15	(12)
CTL0248	CT869	*pmpE*	Polymorphic outer membrane protein	1.99	-2.04	(32)
CTL0249	CT870	*pmpF*	Polymorphic outer membrane protein	3.03	-2.23	(32)
CTL0250	CT871	*pmpG*	Polymorphic outer membrane protein	2.03	-1.55	(12)
CTL0429	CT177	*dsbG*	Disulfide bond chaperone	2.79	-2.47	(32)
CTL0610	CT356	*dsbH*	Thioredox_DsbH domain-containing protein	3.07	-2.11	(12)
CTL0670	CT413	*pmpB*	Polymorphic outer membrane protein	4.17	-1.74	(32)
			Gene Regulation associated	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.
CTL0044	CT675	*mcsB*	Protein-arginine kinase	3.52	-2.21	(32)
CTL0045	CT676	*mcsA*	UVR domain-containing protein	2.53	-2.92	(32)
CTL0727	CT467	*atoS*	Two component regulator, histidine kinase	3.33	-1.80	(32)
CTL0728	CT468	*atoC*	Two component system response regulator	7.95	-2.78	(53)
CTL0894	CT630	*chxR*	Atypical response regulator protein ChxR	3.68	-2.93	(54)
			Glycogen synthesis	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.
CTL0167	CT798	*glgA*	Glycogen synthase	9.58	-2.86	(12, 55)
CTL0342	CT087	*malQ*	4-alpha-glucanotransferase	9.43	-4.29	(38)
CTL0500	CT248	*glgP*	Alpha-1,4 glucan phosphorylase	2.70	-1.71	(55)
			Type III Secretion System	Fold change		
Gene ID	Ctr D ORF	Name	Protein Names	*dacA*op_OE	*dacA*-KD	Ref.
CTL0041	CT672	*sctQ*	Type III secretion component, basal body	1.93	-1.76	(38)
CTL0043	CT674	*cdsC*	Type III secretion structural protein	1.79	-1.56	(38)
CTL0343	CT088	*scc1*	Type III secretion chaperone	4.48	-3.78	(38)
CTL0824	CT561	*sctL*	Type III secretion system protein	2.18	-1.75	(32)

Figure Legends

Figure 1. Cyclic di-AMP accumulation is linked to secondary differentiation in *C. trachomatis*. A. Defining characteristics of chlamydial EBs and RBs. **B.** A hypothetical model representing the correlation between c-di-AMP levels and secondary differentiation. The dashed line represents a threshold level of c-di-AMP needed to drive secondary differentiation in a given RB. **C.** A schematic diagram of the constructs used in this study. All constructs used are anhydrotetracycline (aTc)-inducible as shown by the P_{tet} promoter. The location of the 6xH tag is shown as well as the approximate location of the crRNA for the CRISPRi vectors. Diagram is not to scale. **D.** Measurement of c-di-AMP concentrations in infected cell lysates from the strains shown in panel C. *C. trachomatis* serovar L2 (434/Bu) transformed with the indicated constructs were infected into HeLa cells. At 10 hpi, expression of the construct was induced or not with 5 nM aTc, and the infected cells were harvested at 16 or 24hpi. Levels of c-di-AMP in the supernatant were measured using by ELISA. The left panel shows the increase from 16 to 24hpi during overexpression of the control mCherry protein. The middle and right panels show the levels of c-di-AMP in the indicated strains at 16 and 24hpi, respectively. *: $p<0.05$, **: $p<0.001$ via ratio paired t-test as compared to the mCherry control.

Figure 2. CRISPRi-mediated *dacA-ybbR* knockdown displays reduced levels of transcripts for late genes. (A-D) HeLa cells were infected with *C. trachomatis* transformed with a plasmid encoding an anhydrotetracycline (aTc)-inducible CRISPRi-dCas12 system targeting the *dacA* promoter (i.e., *dacA*-KD; see Fig. 1C). At 10 hpi, knockdown was induced or not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi. Immunofluorescence analysis (IFA) and inclusion forming units (IFU) samples were collected at 24hpi. **A.** IFA images of the *dacA*-KD strain at 24 hpi. Shown are individual panels for dCas12 and major outer membrane protein (MOMP) labeling as well as the merged image. **B.** Quantification of IFUs from uninduced and induced samples at 24 hpi. **C.** Quantification of genomic DNA by qPCR in uninduced and induced samples. **D.** Quantification of transcripts by RT-qPCR for *dacA, ybbR*, *euo*, *hctA*, and *omcB* from uninduced and induced samples. **(E-H)** HeLa cells were infected with *C. trachomatis* transformed with a plasmid encoding an aTc-inducible CRISPRi-dCas12 system targeting the *dacA* promoter and the complementing *dacA and ybbR_6xH* alleles (i.e., *dacA*-KD_*dacAop*; see Fig. 1C). At 10 hpi, knockdown was induced or not with 5 nM aTc, and immunofluorescence analysis (IFA), inclusion forming units (IFU), DNA, and RNA samples were collected as above. **E.** IFA images of *dacA*-KD-*dacA*op strain at 24 hpi. **F.** Quantification of IFU from uninduced and induced samples at 24 hpi. **G.** Quantification of genomic DNA by qPCR in uninduced and induced samples. **H.** Quantification of transcripts by RT-qPCR for *dacA, ybbR*, *euo*, *hctA*, and *omcB*. IFA images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 1 μm. UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: $p<0.05$; **: $p<0.001$ via two sample equal variance t-test.

Figure 3. Overexpression of DacA and YbbR_6xH prematurely increases *hctA* transcript levels. (A-C) HeLa cells were infected with *C. trachomatis* transformed with an aTc-inducible plasmid encoding wild-type DacA and YbbR_6xH (i.e., *dacA*op; see Fig. 1C). At 10 hpi, expression of the construct was induced or not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi. Immunofluorescence analysis (IFA) samples were collected at 24hpi. **A.** IFA images of the *dacA*op strain at 24 hpi. Shown are individual panels for DacA, YbbR_6xH, and major outer membrane

protein (MOMP) labeling as well as the merged image. **B.** Quantification of genomic DNA by qPCR in uninduced and induced samples. **C.** Quantification of transcripts by RT-qPCR for *dacA, ybbR, euo, hctA, and omcB*. **(D-F)** HeLa cells were infected with *C. trachomatis* transformed with an aTc-inducible plasmid encoding DacA(D164N) and YbbR_6xH (i.e., *dacA*opMut; see Fig. 1C). At 10 hpi, expression of the construct was induced or not with 5 nM aTc, and DNA,and RNA samples were collected at 10, 14, and 24 hpi. IFA samples were collected at 24hpi. **D.** IFA images of *dacA*opMut strain at 24 hpi. Shown are individual panels for DacA, YbbR_6xH, and MOMP labeling as well as the merged image. **E.** Quantification of genomic DNA by qPCR in uninduced and induced samples. **F.** Quantification of transcripts by RT-qPCR for *dacA, ybbR, euo, hctA*, and *omcB* from uninduced and induced samples. IFA images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 1 μm. UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: p<0.05; **: p<0.001 via two sample equal variance t-test.

Figure 4. The levels of c-di-AMP are correlated with late gene transcripts. RNA sequencing was performed from HeLa cells infected with either the *dacA*op or *dacA*-KD strains. RNA samples were collected at 16hpi for *dacA*op and 24hpi for *dacA*-KD after inducing expression of the relevant constructs at 10hpi. Shown is a volcano plot of the RNA sequencing results with the vertical dashed lines indicating a two-fold change in transcript levels and the horizontal lines indicating a p-value of 0.05. The plot was made using GraphPad Prism software. Green spots represent genes demonstrating a statistically significant two-fold change in transcription levels between uninduced and induced samples. See also Table 2 and Supplemental Table 2 for more details.

Figure 5. Late genes transcripts are increased by high level of c-di-AMP in penicillin-treated conditions. HeLa cells were infected with a *C. trachomatis* strain transformed with a *dacA*op construct encoding spectinomycin resistance. At 10 hpi, expression of the construct was induced or not with 5 nM aTc and treated with 1 U/mL of penicillin G (PenG). DNA and RNA samples were collected at 10, 14, and 24 hpi, and immunofluorescence analysis (IFA) samples were collected at 24hpi. **A.** IFA images of the *dacA*op strain in 1 U/mL PenG at 24 hpi. Shown are individual panels for DacA, YbbR_6xH, and major outer membrane protein (MOMP) labeling as well as the merged image. IFA images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 2 μm. **B.** Quantification of genomic DNA by qPCR in uninduced and induced samples. **C.** Quantification of transcripts by RT-qPCR for *dacA, ybbR, euo, hctA,* and *omcB*. UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: p<0.05; **: p<0.001 via two sample equal variance t-test.

Figure 6. EB production is induced by high levels of c-di-AMP. (A-C) HeLa cells were infected with the chlamydial transformants *dacA*op (A), *dacA*-KD (B), and *dacA*opMut (C). Expression of the constructs was induced or not with 5 nM aTc at 10hpi. At 18, 20, 22, 24, 32, and 48 hpi, infected cell lysates were harvested for IFU quantification. UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. #: Non-detected from the induced samples. *: p<0.05; **: p<0.001 via two sample equal variance t-test.

Figure 7. A working model illustrating how c-di-AMP impacts the onset of secondary differentiation in *C. trachomatis*. A. During RB division, there is a gradient of c-di-AMP that forms between the mother and daughter cell. The lighter blue represents lower levels of c-di-AMP, the darker

blue represents higher levels of c-di-AMP. In the depicted scenario, the daughter cell is able to divide
again, having not reached a critical threshold of c-di-AMP. In contrast, the mother cell accumulates
sufficient c-di-AMP to trigger secondary differentiation to an EB. **B.** A model illustrating how c-di-
AMP levels impact secondary differentiation progression. Shown is the "normal" condition of wild-type
bacteria, bacteria overexpressing DacA and YbbR, and bacteria expressing the CRISPRi system
targeting the *dacA* promoter. Depending on the amount of c-di-AMP produced, secondary differentiation
can either be triggered earlier or later as shown. The black dots represent EBs; the bigger and blue-
colored circles show the RB cells. The brightness of blue reflects the c-di-AMP level.

Supplemental Information

Figure S1. The predicted transmembrane domains in DacA and YbbR. Transmembrane domains in
DacA (A) and YbbR (B) were predicted with TOPCONS (https://topcons.cbr.su.se/)(51). The red and
blue lines represent cytosolic and periplasmic domains, respectively.

Figure S2. Overexpression of DacA, but not YbbR, is detrimental to the chlamydial developmental
cycle. (A-D) HeLa cells were infected with *C. trachomatis* transformed with a plasmid encoding an
anhydrotetracycline (aTc)-inducible DacA_6xH. At 10 hpi, expression of the construct was induced or
not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi.
Immunofluorescence analysis (IFA) and inclusion forming units (IFU) samples were collected at 24hpi.
A. IFA images of the *dacA*_6xH strain at 24 hpi. Shown are individual panels for DacA_6xH and major
outer membrane protein (MOMP) labeling as well as the merged image. **B.** Quantification of IFUs from
uninduced and induced samples at 24 hpi. **C.** Quantification of genomic DNA by qPCR in uninduced
and induced samples. **D.** Quantification of transcripts by RT-qPCR for *dacA, ybbR*, *euo*, *hctA*, and *omcB*
from uninduced and induced samples. **(E-H)** HeLa cells were infected with *C. trachomatis* transformed
with a plasmid encoding an aTc-inducible YbbR_6xH. At 10 hpi, the construct was induced or not with
5 nM aTc, and IFA, IFU, DNA, and RNA samples were collected as above. **E.** IFA images of *ybbR*_6xH
strain at 24 hpi. **F.** Quantification of IFU from uninduced and induced samples at 24 hpi. **G.**
Quantification of genomic DNA by qPCR in uninduced and induced samples. **H.** Quantification of
transcripts by RT-qPCR for *dacA, ybbR*, *euo*, *hctA*, and *omcB*. IFA images were acquired on a Zeiss
AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 1 μm. UI =
uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: p<0.05; **: p<0.001 via two
sample equal variance t-test.

Figure S3. The phenotype of the transformants encoding DacA(D164N)_6xH. HeLa cells were
infected with *C. trachomatis* transformed with a plasmid encoding an anhydrotetracycline (aTc)-
inducible DacA(D164N)_6xH. At 10 hpi, expression of the construct was induced or not with 5 nM aTc,
and IFA and IFU samples were collected at 24 hpi. **A.** IFA images of the dacA_6xH strain at 24 hpi.
Shown are individual panels for DacA_6xH and major outer membrane protein (MOMP) labeling as
well as the merged image. **B.** Quantification of IFUs from uninduced and induced samples at 24 hpi. The
images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens
objective. Scale bar: 1 μm

Figure S4. The localization of DacA_6xH and YbbR_6xH during the first RB division. HeLa cells
were infected with *C. trachomatis* transformed with a plasmid encoding an anhydrotetracycline (aTc)-
inducible DacA_6xH (Upper) or YbbR_6xH (Lower). At 4 hpi, expression of the constructs was
induced with 5 nM aTc. At 10.5 hpi, the infected cells were fixed and permeabilized. The images were
acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar:
1 μm

Figure S5. Phenotypic characterization of a *dacA*op overexpression construct encoding
spectinomycin resistance (Spc^R). HeLa cells were infected with *C. trachomatis* transformed with a
*dacA*op overexpression construct encoding Spc^R. At 10 hpi, expression of the construct was induced or
not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi.

Immunofluorescence analysis (IFA) samples were collected at 24hpi. **A.** IFA images of the *dacA*op strain at 24 hpi. Shown are individual panels for DacA, YbbR_6xH, and major outer membrane protein (MOMP) labeling as well as the merged image. **B.** Quantification of genomic DNA by qPCR in uninduced and induced samples. **C.** Quantification of transcripts by RT-qPCR for *dacA, ybbR, euo, hctA,* and *omcB*. IFA images were acquired on a Zeiss AxioImager.Z2 equipped with an Apotome2 using a 100X lens objective. Scale bar: 2 μm. UI = uninduced (i.e., -aTc); I = induced (i.e., +aTc) for all sample types. *: $p<0.05$; **: $p<0.001$ via two sample equal variance t-test.

Figure S6. Quantification of transcripts by RT-qPCR for *dacA*, *ybbR*, *euo*, *hctA*, and *omcB* from *dacA*-KD and dacA_6xH strains infected into STING-KO HeLa cells. STING-KO HeLa cells were infected with *C. trachomatis* transformed with a plasmid encoding an anhydrotetracycline (aTc)-inducible **A.** CRISPRi-dCas12 system targeting the *dacA* promoter and/or **B.** DacA_6xH. At 10 hpi, expression of the construct was induced or not with 5 nM aTc, and DNA and RNA samples were collected at 10, 14, and 24 hpi. See also Figures 2 in the main text and Supplemental Figure 2.

Table S1. List of plasmids and primers used in the study.

Table S2. Summary of RNA sequencing reads and coverage.

Table S3&4. RNA sequencing results for *dacA*op overexpression (S3) and *dacA* knockdown (KD) (S4).

A. DacA_{WT}-YbbR6xH

B. dacA-KD

C. DacA_{D164N}-YbbR6xH

A.

Cell division

EB

B.

Normal Condition

Higher cyclic di-AMP

Lower cyclic di-AMP

1   8   12   16   24   32   48   hpi

Cyclic di-AMP level

EB
Threshold
RB

Normal

Cyclic di-AMP level

EB
Threshold
RB

Higher c-di-AMP

Earlier

Cyclic di-AMP level

EB
Threshold
RB

Lower c-di-AMP

Delayed

Time (hpi)