Emergent patterns of patchiness reflect decoupling between

ocean physics and biology

Patrick Clifton Gray*[,1,2], Yoav Lehahn[2], Emmanuel Boss[1]

1. School of Marine Sciences, University of Maine, Orono, ME, USA

2. Department of Marine Geosciences, Charney School of Marine Sciences, University

of Haifa, Haifa, Israel

*Corresponding author: Patrick Gray (patrick.gray@maine.edu)

Abstract

While a rich history of patchiness research has explored spatial structure in the ocean,

there is still no consensus over the controls on biological patchiness and how

biogeochemical processes and patchiness relate. The prevailing thought is that physics

structures biology, but this has not been tested at the basin scale with consistent *in situ*

measurements. Here we show that the patchiness of physics and biology are decoupled

at the global scale through analysis of a global dataset of in situ surface optical

properties from the R/V *Tara* and using the slope of spatial scale vs variance to quantify

patchiness. Based on analysis of ~650,000 nearly continuous (dx~150m)

measurements from an underway sampling system - representing five years of

consistently collected data across the Atlantic, Pacific, and Southern Oceans - we find

the patchiness of physical and biological parameters are uncorrelated. We show that

variance slope is an emergent property with unique patterns in biogeochemical

properties that are distinct from physical tracers, yet connected to other biological

tracers. These results provide context for decades of discrepancy between *in situ*

studies, could support new tests of biogeochemical model parameterizations, and open

the way for new insight into processes regulating the observed patterns.

Introduction

Most phytoplankton species near the ocean's surface have a doubling time on the order

of a day[1]. This rapid doubling time means phytoplankton are extremely responsive to

nutrient fluxes and external forcing[2]. Coupled with the dynamic physical nature of the

marine environment and grazing pressures with resulting loss rates similar to growth

rates for phytoplankton[3,4], this results in considerable spatial heterogeneity, or

patchiness, at all scales[5–8]. This spatial heterogeneity supports increased diversity by

creating different environmental conditions within close proximity, ecosystem stability

through diversity and maintenance of seed populations, and efficient transfer of carbon

up trophic levels[9]. The link between patchiness and the efficient transfer of carbon in the

marine ecosystem is largely due to dense aggregations that make it energy efficient for

zooplankton to graze, and patches of zooplankton make it feasible for planktivorous fish

to meet their own energy requirements[10,11]. This spatial variability of phytoplankton and

zooplankton, and the generally increasing patchiness as one moves up the trophic

chain, has important implications for food availability and productivity of the entire ocean

food web[12,13].

The spatial patterns of the plankton ecosystem and their drivers have long been

investigated. An early outline by Hutchinson discusses nutrient and temperature

gradients, stochastic events such as storms, intra-species signaling, competition, and

predation among processes that likely control these patterns[14]. An updated synthesis by

Levin described pattern and scale as *the* central problem in ecology, arguing for cross-

scale investigations to understand the mechanisms and consequences of marine

patchiness[5]. The scale-dependent spatial heterogeneity exhibited by both biology and

physics has been a central focus of studies into pattern and scale in the ocean[6], with

much of this work attempting to divide spatial scales into domains of scale-

independence and then investigating the dominant processes influencing biology in

those domains[15,16]. Power spectral analysis, a Fourier decomposition where variance is

partitioned into the contribution of bands of specific frequencies or wavenumbers (see [17]

for an early review on the implementation of the technique in ecology), is often used to

describe phytoplankton patchiness and quantify this scale-dependence. The scaling

behavior of a geophysical variable is typically represented as a log-log plot of variance

against length scale. If that relationship is approximately linear (in log space) this

represents a power law relation between variance and spatial scale where the exponent

of that power law relationship is the spectral slope[18]. A flatter spectral slope indicates

less of a decrease of variance with scale, i.e. more variability at smaller scales relative

to larger scales, when compared to a steeper slope. This parameter has often been

used synonymously with patchiness, but can also be thought of as quantifying the

cascade of variance across scales. While there are a range of issues with this

decomposition, such as discarding of phase information[9,19], it is commonly used as an

indicator of the distribution of spatial variance.

There is over a century of inquiry into the spatial heterogeneity of ocean physics.

Turbulence in the ocean generally cascades down spatial scales. Turbulent mesoscale

eddies spin off of large-scale circulation features. Energy then transfers from these

major eddies to smaller turbulent whorls and finally to viscous dissipation into heat[20,21].

This turbulent cascade generally has a spectral slope of -5/3 in the inertial subrange (a

spatial range smaller than the major energy containing eddies, but larger than viscous

eddies). While Kolomogrov initially proposed this cascade of variance for fully

developed 3D turbulence in the ocean's inertial subrange, O(10-0.001) meters

depending on flow characteristics[22], the -5/3 spectral slope has been shown to hold

generally true for the inertial subrange of 2D ocean turbulence O(100-1) km as well[23].

Temperature generally follows the same cascade as turbulent kinetic energy. It has a

source of variance at large scales, a dissipation of variance at small scales, and it is

typically being mixed by similar turbulent processes[24].

The patchiness of phytoplankton was initially attributed simply to turbulent stirring by

physical processes[25], with a general conclusion of consistent scaling behavior between

physical and biological features[16]. This conclusion was drawn largely from its spectral

slope that appeared similar to the -5/3 of 2D turbulence[25,26]. Modeling indicated that the

spectral slope of phytoplankton was also influenced by the phytoplankton reproductive

rate, diagnosed by a scaling break at larger spatial scales where the spectral slope

became flatter than that of a passive tracer[27,28]. This led to a substantial amount of

work, with many conflicting results, investigating the time and length scales where

biologically generated spatial variability (growth rates, grazing, etc) is relevant and

important to plankton versus where physics dominates[6,7,9,22,29].

Abrahams[7] suggested and others have corroborated in modeling[30,31] that the spatial

patterns in plankton are a product of the timescales of their response to physically

driven changes. For example, the equilibration of temperature to a new heat flux is

considerably slower than the response of phytoplankton to a nutrient flux. A small burst

of cold nutrient-rich water thus leads to increased spatial heterogeneity in chlorophyll-a

(chl-a) vs temperature, but as nutrients are exhausted and the heat is mixed over time

chl-a and temperature would return to covarying[29]. Modeling work shows a steeper

spectral slope in temperature compared to chl-a, attributing some of the slope variability

to low frequency physical variations and the correlation of physical and biological

variables to "the dominant role of physics in the spatial variability of plankton

distribution"[30].

Despite these theoretical advances, actual observations remain inconsistent. Some

work concludes that at <1km scales biological processes are important for patch

dynamics[10,32], while other work asserts that at this scale spatial distributions of

phytoplankton are dominated by turbulent diffusion. For example this latter conclusion is

advanced in work that found a strong correlation between chl-a and temperature

patchiness below 5km yet not so above 5 km to 80km[17]. Instances spring up where

phytoplankton are patchier than zooplankton[33], directly conflicting generally accepted

findings that the patchiness of phytoplankton is between that of zooplankton and

physical factors[7,25,34]. Work assessing the spectral slope of chl-a in the same region for

multiple years found completely contrasting measurements from month to month[22,35].

After decades of work, it is not clear at which spatiotemporal scales biological

processes are important, or even which processes are dominant in controlling

patchiness in the ocean.

While biophysical ocean patchiness has been relatively well documented on local

scales, its global characteristics have not yet been analyzed. Therefore, there is little

data to test new theories, constrain impacts on productivity and diversity, and simply

describe the patchiness itself. Importantly, this limits our ability to parameterize global

biogeochemical models that are computationally limited from explicitly modeling these

scales in our global models of the carbon cycle.

Here we address this gap, and explore the global characteristics of biophysical

patchiness using a unique dataset of *in situ* surface plankton-related optical properties

from the R/V *Tara* program. All data comes from the Tara Pacific[36] and Tara

Microbiome missions which represent five years of observations spanning the Atlantic,

Pacific, and Southern Oceans. This data was collected and processed consistently.

After conservative quality control this dataset amounts to N=661,552 (dt=1 minute,

dx~150m) measurements used in this analysis (Figure 1). As we aim to discern

between patterns of physical and biological patchiness, the dataset variables are

divided into two corresponding groups, with the physical domain being represented by

temperature, salinity and density, and the biological domain by chl-a, particulate

attenuation of light at 443nm (c_p(443)), and γ an optical proxy for mean particle size[37].

The latter two, because they may include non-living particles, are referred to as

biogeochemical properties.

Figure 1: The spatial extent of data used in this study and the geographic distribution of

chlorophyll-a (a) and sea surface temperature (b). The data includes 661,552 minute

binned measurements. Insets show the histograms of both variables. The four numbers

in panel (a) correspond to the example data shown in Figure 2.

Results and discussion

While the Fourier-based spectral slope is an extremely useful and commonly used tool

for characterization of patchiness, it is not suitable for analyzing unevenly sampled

datasets. To overcome this limitation here we quantify patchiness via the dependence

of the variance, V, on length scale, Γ following [31]. As found in previous work, V varies

mostly linearly with Γ in log space (Figure 2), indicating a power law relationship of the

form: (1) $V = L^{\Gamma}$, where Γ describes the slope (exponent) of the relationship in log

space. In this work we focus on Γ, the "variance slope". As with a power spectral density

slope, a transect with a lower Γ would have relatively more variance retained at smaller

length scales relative to larger scales compared to a transect with higher Γ, i.e. a lower

Γ is patchier. This metric can be thought of qualitatively as a ratio of large scale to small

scale variance (Figure S1), with Γ increasing as the large scale variance dominates the

small. This parameter is similar and strongly correlated with the conventional Fourier-

based spectral slope exponent (Figure S2 and S3), but our approach (from [31]) has the

advantage of being applicable to data with gaps, as long as they are smaller than 20%

of a transect.

Figure 2. Examples of the chlorophyll-a (green) and temperature (red) absolute values along a leg (left panel in each pair) and the scale-dependent variance used to calculate the variance slope (Γ) derived from these legs (right panel in each pair). The numbers of each panel (1-4) correspond to the locations in Figure 1a.

Specifically in this work, variance was calculated on legs of 500 samples over a set of 11 log-distributed windows from 3 to 500 samples, spanning approximately 0.5km to 75km. Both variance and window size were log-transformed, and we ran a least-squares fit to find the slope of this line, yielding the Γ of $V = L^\Gamma$. In our analysis of global patterns of oceanic patchiness, we distinguish between Γ of the biological and biogeochemical variables chl-a, $c_p(443)$ and γ (Γ_{chl-a}, Γ_{cp} and Γ_γ, respectively), and of the physical variables temperature, salinity and density ($\Gamma_{temperature}$, $\Gamma_{salinity}$ and $\Gamma_{density}$, respectively).

Comparison between the geographic distributions of Γ_{chla} and $\Gamma_{temperature}$ reveals distinct differences between global patterns of physical and biological patchiness (Figure 3). Notably, the geographic distributions of $\Gamma_{temperature}$ are fairly random, while Γ_{chla} is consistently lower in oligotrophic regions, particularly the subtropical gyres, and higher in physically energetic regions such as western boundary currents and eastern boundary upwelling zones. In agreement with most previous observations, Γ_{chla} is, on average, lower than $\Gamma_{temperature}$, indicating chl-a is patchier than temperature. $\Gamma_{temperature}$ is fairly normally distributed around 1, while Γ_{chla} is slightly right-skewed with a mean around 0.5 (see inserts in Figure 3). Based on the observed relationship between Γ and

the power spectral density slope (Figure S3) the classic -5/3 slope found in previous

studies for the inertial subrange would correspond to a   value of 0.84 which is near,

but slightly lower than the mean of our global   temperature calculations, 0.97.

Figure 3. Geographic distributions of   chla (a) and   temperature (b). Insets show the

histograms for these two variables.

Interestingly, running the same analysis on satellite-based chl-a and temperature

reveals contradicting results. Although satellite imagery has been successfully used to

track individual chl-a patches[38,39], and to quantify patchiness at local scales[25], in the

satellite-based analysis we find $\Gamma_{temperature}$ is lower than Γ_{chla} (i.e. the opposite of our *in*

situ results), and the geographic patterns for both chl-a and temperature are different

(Figures S4 and S5, respectively). We attribute this discrepancy primarily to issues of

measurement sensitivity and atmospheric correction of ocean color remote sensing

(from which chl-a is derived) which is applied as a pixel level correction, artificially

injecting variance at small scales. This effect is likely to have the strongest impact in

oligotrophic regions, where chlorophyll-a levels are extremely low, resulting in minimal

absorption and reduced particulate scattering, and thus a low signal to noise ratio in the

satellite retrieval. This could cause issues in both spatial and temporal analyses[40]. This

corroborates results from previous work that used a semivariogram approach to analyze

variance in global satellite chl-a patterns from SeaWIFS[41] which showed that

unresolved variance was negatively correlated with mean chl-a and the authors

suggested the high fraction of unresolved variance in the subtropical gyres might stem

from atmospheric correction, though they did not differentiate between instrument noise

and submesoscale variability. A more recent, but similar study compared the unresolved

variance between MODIS-Aqua and SeaWIFS and found it was reduced, but not

eliminated with MODIS[42].

The global patterns characterizing our observations point to decoupling between the

governing processes underlying the formation of physical and biological ocean

patchiness. This decoupling is emphasized when plotting the correlation between the

different types of patchiness (Figure 4), which we show both for all individual legs and

averaged by Longhurst provinces[43] (Figure 5). The Γ of biological variables is correlated

globally, even between chl-a and γ, the mean particle size (Figure 4a,b). Similarly, Γ of

physical variables are well correlated with each other (Figure 4c,d). Yet, there is little to

no correlation between the Γ of biological variables and physical variables (Figure 4e,f).

This suggests that ecosystem patchiness is an emergent property due to biological

processes rather than driven by physical forcings.

If we decrease the top end length scale that is used for the calculation of Γ, for example

from 75km to 5km, Γ_{chla} and $\Gamma_{temperature}$ become only marginally more correlated

($R^2=0.04$ for 0.5km to 5km vs $R^2=0.03$ for 0.5km to 75km). This is in contrast with older

work which showed that chl-a and temperature patchiness were strongly correlated

below 5km yet not so above 5 km to 80km[17].

Similarly to the differences between geographic distributions patterns (Figures 1 and 3)

the correlations between Γ of different physical variables differ substantially from the

correlations between absolute values (Figure S6). We show chl-a and gamma have

minimal correlation in absolute value yet a correlation in Γ, chl-a and temperature have

a strong correlation in absolute value yet minimal in Γ, and temperature and salinity

have minimal correlation in absolute value yet a strong correlation in Γ.

Figure 4. Variance slope ( ) comparisons between different variables. $_{chla}$ correlates

with the biogeochemical parameters such as attenuation (a) and   (mean particle size)

(b), and   of different physical variables are correlated (d, c), yet biogeochemical vs

physical variables don't have a notable correlation (e, f). Larger colored markers

correspond with the Longhurst provinces from Figure 5. N.b. correlations are shown for

all data, not the Longhurst province means, and all plots have a p-value < 0.001.

Figure 5. Longhurst provinces that contain N=> 25 legs shown geographically along

with the Tara legs within each.

Another fundamental difference between global patterns of physical and biological

ocean patchiness emerges when looking at the data averaged by biogeochemical

provinces and plotting the relationship between   and the absolute value of each

variable (Figure 6). The biological variables that are concentration dependent (i.e. chl-a

and cp(443)) have a positive relationship between their   and their absolute value, such

that high biomass provinces are less patchy (i.e. associated with higher   ) than low

biomass provinces, (Figure 6a-c). In contrast, there is no relationship between   and the

absolute value of the physical variables (Figure 6d-e). In other words, biomass and

biological patchiness are linked (inversely) while heat and salt content does not

correlate with their patchiness.

Figure 6. The top (a-c) and bottom (d-f) rows show the absolute value vs variance slope

( ) for a range of biological and physical variables, respectively. The average of each

province is shown in the larger markers and the colors correspond to the provinces on

the map. N.b. correlations are shown for all data, not the Longhurst province means,

and all plots have a p-value < 0.001.

The linkage between biological patchiness and biological productivity is emphasized

when taking into account nutrient availability averaged by Longhurst Province (Figure

7). When mapped onto the absolute value of chl-a and Γ_{chla}, a range of nutrients have a

clear increasing relationship with both (Figure 7), with the highest nutrient values also

having high chl-a and Γ_{chla}.

Figure 7. Chlorophyll-a (chl-a) concentration vs Γ_{chla} colored by average nutrient

concentrations in each Longhurst province. Nutrient data is from the World Ocean Atlas

2018.

The nature of the small scale variability in biological signals, and the way they differ

from the physical signals, is emphasized when zooming on segments of the transects

(Figure 8). Careful examination reveals that while the high frequency changes in our

underway observations of biological variables may appear to be associated with

instrumental noise, the individual measurements are robust, and the changes reflect

small scale (<1km) biological variability patterns (Figure 8). This robustness can be

seen both in the covariation between different sensors (Figure 8a) and the raw

absorption and attenuation spectra (Figure 8b,c). This small scale variability lowers

and can make the variance spectra nearly flat in regions without low frequency

variation, such as the Sargasso Sea and Subtropical South Pacific.

Figure 8. Example of measured variables along a single leg (a), showing data "spikes"

that may initially be considered instrumental noise, but when inspecting the individual

spectra (b and c), used to derive the various proxies, they are consistent with

expectation and across the absorption and attenuation sensors. The two bottom rows

show a 30 minute subset of particulate absorption spectra (b) and particulate

attenuation spectra (c) representing the period between the vertical black lines in panel

a. Each spectra is the average of one minute of sampling, is colored by time (from

brown to blue), and spans wavelengths from 410nm to 750nm. This transect is from the

Coral Sea.

Our work does not tie the variance slope to any particular process. Possibly Γ

represents the integrated impact of physical and biogeochemical characteristics within a

region. The lower Γ_{chla} in oligotriohic regions may be attributed to a sustained injection

of variance from ecosystem interactions (e.g. growth, grazing). Recent work

investigating zooplankton patchiness found coherent small scales patches across many

taxa dominated by length scales of 10-30 m[40]. Even highly idealized models have been

shown to generate patchiness via grazing[44] which could help explain aspects of our

results. While previous work using this same metric suggested that the response time of

phytoplankton (i.e. growth rate) is faster than the response time of temperature (i.e.

equilibration time) as the reason chl-a is patchier[31], this doesn't agree with our results,

where the regions with higher assumed growth rates are less patchy (e.g. eastern

upwelling zones). This may suggest that certain elements of the ecosystem are

responding faster in these highly patchy oligotrophic regions, possibly grazers or growth

following fine-scale vertical nutrient fluxes. Stratification may play a role with a more

stratified water column leading to more intermittent linkages between the surface and

the nutrient-rich deep ocean, resulting in turn in a patchier expression in the highly

stratified subtropical regions. While the drivers are not clear from our work, Mahadevan

& Campbell[31] suggest that an increasingly patchy distribution requires finer scale

sampling or grid spacing, and if these processes are important, this indicates fine-scale

grid spacing is in fact still important across broad swaths of the ocean generally

considered "homogenous".

While the focus of this work is on scaling relationships, the fractal-like patchiness of

phytoplankton is not infinite. At a fundamental level, patches are formed by individual

phytoplankton living and dying[45]. Currently it is unclear how to connect the scales being

investigated here with the individual organisms at the heart of the matter; Franks[22] notes

that it is not likely we can diagnose the dynamics underlying the observations from

spectral slopes alone. Towards mechanistic understanding of the processes underlying

the spatial patterns we expect Lagrangian approaches will be key, particularly to parse

out spatial and temporal dynamics. Future work should examine joint spatial patterning

across trophic levels, combining the methods here, rapid methods for a wide size range

of plankton[46], and methods for quantifying zooplankton patchiness[10]. Additionally, it may

be that improved atmosphere correction of satellite products could enable a new source

of information via spatial analyses across the global ocean. Higher signal to noise

regions, such as coastal waters or higher latitudes during the spring bloom, may be

more amenable to satellite analyses of spatial patterns under existing atmospheric

correction schemes.

Summary and conclusion

We identified two fundamental differences between global patterns of physical and
biological patchiness: globally biological patchiness at the scales analyzed here
(~0.5km to 75km) is itself spatially organized while physical patchiness is more random,
and patchiness of concentration-dependent biological variables is correlated with the
absolute values of these variables, while no such correlation is found for physical
variables. The statistical distributions of Γ_{chla} and $\Gamma_{temperature}$ agree with previous work,
where Γ_{chla} has a generally lower value, i.e. a patchier spatial pattern, compared to
$\Gamma_{temperature}$. We find global patterns of biological patchiness to coincide with
biogeochemical provinces and with nutrients levels. While our analysis shows the
patchiness of biological variables are intercorrelated, and patchiness of physical
variables are intercorrelated, contrary to common view, we find no interdependence
between physical and biological patchiness. Our results point to decoupling between the
governing processes underlying the formation of physical and biological ocean
patchiness. This suggests that biological processes regulate the spatial patterning of the
oceanic ecosystem independently of the physical processes or, rather than true
independence, that they modulate the initial physical-driven spatial pattern baseline
sufficiently to manifest as a decoupling. We emphasize here that the absolute value of
variance itself is correlated at all scales between physical and biological variables
(Figure S7). Moreover, while it is well known that there is a relationship between
temperature and chl-a, largely attributed to nutrient fluxes[47], our findings show this does
not extend to variance slope. Our results contradict results from model studies (e.g. [30])

and from satellite analyses (e.g. [48]), including our own (Figure S4 and S5). This

discrepancy holds even when comparing to other work on the same range of scales as

those examined here, which had concluded that biological patchiness is controlled by

mesoscale mixing[25]. This suggests that these two important tools for understanding

ocean ecosystems (models and satellite observations), fail to capture fundamental

characteristics of the marine ecosystem. Our results may also help explain the

disagreement shown across decades of patchiness research including many *in situ*

studies. It may be that there is no generalizable patchiness relationship between

physics and biology in the ocean and no consistent spatiotemporal scale where biology

or physics dominates. Accordingly, basing general theories on the patchiness of a given

tracer type from a specific site and over short time scales could be misguided. Instead,

the ocean appears to consist of myriad physical and biogeochemical processes

manifesting as a diverse set of spatial patterns. Thus, while no consistent relationships

emerge these observable spatial patterns may provide a window into the processes

creating them.

These results support the use of low dimensional spatial encodings such as Γ to study

these upper-ocean processes. We speculate these spatial metrics may be sensitive to

biogeochemical parameters not represented by the absolute value of chl-a. Importantly,

we cannot resolve marine ecosystems at the mesoscale and below in basin-scale

biogeochemical models, thus we must increase our ability to observe, invert, and

understand what our models are missing[49]. Spatial patchiness patterns may help drive

our models to reality and serve as an underutilized source of insight into the underlying

processes generating the observed patchiness.

Methods

All *in situ* measurements are from the R/V *Tara* underway sampling system. This pulls

seawater from 2m depth. Temperature and salinity data is from a Seabird Scientific

thermosalinograph (SBE 45) and all optical data is derived from particulate absorption

and attenuation measured with a Seabird Scientific ACs.

We quantify patchiness using variance slope, Γ, following [31] which is similar to spectral

analysis, but able to ingest vectors (i.e. transects) with small gaps within them. Γ is

calculated on groups of 500 samples using a set of 11 log-distributed windows from

0.5km to 75km. Variance and window size were log-transformed, and we ran a least-

squares fit to find the slope of this line, yielding the Γ of $V = L^{\Gamma}$. It is worth noting a few

characteristics of Γ to gain some intuition into the metric. The variance slope of a

randomly distributed variable is centered around 0. The calculation is not sensitive to

whether a variable is log-normally distributed (such as chl-a). For example, Γ_{chla} is

identical whether raw chl-a values are input or log-transformed chl-a values. The R/V

Tara is a sailing vessel and thus does not always travel in a straight line or with the

same speed so here we have used time (minute bins of data) instead of distance to

calculate the window, but filtered out legs that have a maximum distance traveled under

30km or over 150km. We have required at least 80% of the transect to contain data and

filtered out transects where the maximum time elapsed is greater than 16 hours.

Hyperspectral absorption and attenuation (400 to 735 nm at ~4 nm spectral resolution,

AC-s, Seabird Sci.) were measured continuously. A 0.2 μm filter cartridge was

connected to the system and we automatically redirected the flow to measure the

properties of filtered seawater for 10 minutes every hour. Total ("normal") seawater was

flowing the rest of the time. Absorption and attenuation measurements recorded during

the filtered periods were interpolated across the transect using CDOM and subtracted

from the total seawater measurements to obtain an estimate of the particulate

absorption (a_p), attenuation (c_p) spectra. This workflow permits retrieval of particulate

optical properties independently from the instrument drift and biofouling[50]. For each

minute of total seawater measurements (sampled at 4Hz) the signal between the 2.5th

and 97.5th percentiles were averaged, and their standard deviation was used to

quantify uncertainty. Dropping the 2.5th to 97.5th percentiles filters out noisy spikes

from bubbles which can be a major problem in optical measurements. All spectra were

manually checked and quality controlled for obviously bad measurements (e.g. bubbles,

bad filtered seawater measurements). These inherent optical properties were used as

proxies for a range of particulate properties, primarily chl-a line height, a chl-a estimate

derived from the absorption peak at 676 nm[51,52], and γ, a proxy for mean particle size[37].

Underway data was collected using Inlinino[53] an open-source logging and visualization

program, and processed using InlineAnalysis

(https://github.com/OceanOptics/InLineAnalysis) following best practices[54].

Longhurst provinces [43] (downloaded from

https://www.marineregions.org/sources.php#longhurst) were used as approximate

delineations of the global ocean into biogeochemical regions. Nutrient data came from

the World Ocean Atlas 2018 (https://www.ncei.noaa.gov/data/oceans/woa/WOA18).

Data Availability

442   All data is available in raw form at NASA's SeaBASS archive. All data prepared and

443   formatted for this study is available on GitHub

444   https://github.com/patrickcgray/spatial_patchiness_tara, easily ingestable as a single

445   geofeather or a csv, and archived on Zenodo [DOI].

Code Availability

All code necessary to generate the figures is on GitHub

https://github.com/patrickcgray/spatial_patchiness_tara and archived at Zenodo [DOI]

To obtain a Docker image for running this in an identical environment to the one used in

this study simply run "*docker pull pangeo/pangeo-notebook:2024.04.05*" from the

command line with Docker Desktop running. A series of Jupyter Notebooks are

provided in the above Github repo for an exact reproduction of all work in this study.

Acknowledgements

We deeply appreciate the R/V *Tara* crew for their invaluable work in collecting this

dataset. We also thank the Tara Ocean Foundation and the Tara Pacific and

Microbiome Expedition participants. Our thanks to Guillaume Bourdin for Tara underway

system processing support. We acknowledge support from the Zuckerman STEM

Leadership Program to PCG. NSF EarthCube program award #2026932 supported the

Pangeo Cloud platform where all analysis was conducted. The optical inline dataset

was collected and analyzed with support from NASA Ocean Biology and

Biogeochemistry program under grants NNX13AE58G, NNX15AC08G and

80NSSC21K0783 to the University of Maine.

Author Contributions

The study was conceived by all authors. The analysis approach was architected by Y.L.

The specific methods and analysis were conducted by P.G. All authors interpreted the

results, wrote the paper, and supported data analysis.

Corresponding author

Correspondence to Patrick Gray

Competing interests

The authors declare no competing interests.

References

1. Laws, E. A. Evaluation of In Situ Phytoplankton Growth Rates: A Synthesis of Data from
Varied Approaches. *Ann Rev Mar Sci* **5**, 247–268 (2013).
2. Falkowski, P. G., Barber, R. T. & Smetacek, V. Biogeochemical controls and feedbacks
on ocean primary production. *Science (1979)* **281**, 200–206 (1998).
3. Behrenfeld, M. J. & Boss, E. S. Resurrecting the ecological underpinnings of ocean
plankton blooms. *Ann Rev Mar Sci* **6**, 167–194 (2014).

4.   Banse, K. Grazing and Zooplankton Production as Key Controls of Phytoplankton
Production in the Open Ocean. *Oceanography* **7**, 13–20 (1994).

5.   Levin, S. A. The problem of pattern and scale in ecology. *Ecology* **73**, 1943–1967 (1992).

6.   Mackas, D. L., Denman, K. L. & Abbott, M. R. Plankton patchiness: biology in the
physical vernacular. *Bull Mar Sci* **37**, 653–674 (1985).

7.   Abraham, E. R. The generation of plankton patchiness by turbulent stirring. *Nature* **391**,
577–580 (1998).

8.   Martin, A. P. Phytoplankton patchiness: the role of lateral stirring and mixing. *Prog*
Oceanogr **57**, 125–174 (2003).

9.   Lévy, M., Franks, P. J. S. & Smith, K. S. The role of submesoscale currents in structuring
marine ecosystems. *Nature communications* vol. 9 4758 Preprint at
https://doi.org/10.1038/s41467-018-07059-3 (2018).

10.   Robinson, K. L., Sponaugle, S., Luo, J. Y., Gleiber, M. R. & Cowen, R. K. Big or small,
patchy all: Resolution of marine plankton patch structure at micro- to submesoscales for
36 taxa. *Sci Adv* **7**, (2021).

11.   Mahadevan, A. The Impact of Submesoscale Physics on Primary Productivity of
Plankton. *Ann Rev Mar Sci* **8**, 161–184 (2016).

12.   Estapa, M. L. *et al.* Decoupling of net community and export production on
submesoscales in the Sargasso Sea. *Global Biogeochem Cycles* **29**, 1266–1282 (2015).

13.   Levy, M. & Martin, A. P. The influence of mesoscale and submesoscale heterogeneity on
ocean biogeochemical reactions. *Global Biogeochem Cycles* **27**, 1139–1150 (2013).

14.   Hutchinson, G. E. The Concept of Pattern in Ecology. *Proceedings of the Academy of*
Natural Sciences of Philadelphia 1–12 (1953).

15.   Haury, L. R., McGowan, J. A. & Wiebe, P. H. Patterns and Processes in the Time-Space
Scales of Plankton Distributions. in *Spatial Pattern in Plankton Communities* (1978).
doi:10.1007/978-1-4899-2195-6_12.

16.   Wiens, J. A. Spatial Scaling in Ecology. *Funct Ecol* **3**, 385 (1989).

17.   Platt, T. & Denman, K. L. Spectral Analysis in Ecology. *Annu Rev Ecol Syst* **6**, 189–210
(1975).

18.   West, G. *Scale: The Universal Laws of Growth, Innovation, Sustainability, and the Pace*
of Life in Organisms, Cities, Economies, and Companies. (Penguin Random House,
2017).

19.   Armi, L. & Flament, P. Cautionary remarks on the spectral interpretation of turbulent
flows. *J Geophys Res* **90**, 11779 (1985).

20.   Kolmogorov, A. N. The local structure of turbulence in incompressible viscous fluid for
very large Reynolds numbers. *Proc R Soc Lond A Math Phys Sci* **434**, 9–13 (1941).

21.   Richardson, L. F. The supply of energy from and to atmospheric eddies. *Proceedings of*
the Royal Society of London. Series A, Containing Papers of a Mathematical and
Physical Character **97**, 354–373 (1920).

22.   Franks, P. J. S. Plankton patchiness, turbulent transport and spatial spectra. *Mar Ecol*
Prog Ser **294**, 295–309 (2005).

23.   Kraichnan, R. H. Inertial ranges in two-dimensional turbulence. *Physics of Fluids* **10**,
1417–1423 (1967).

24.   Batchelor, G. K. Small-scale variation of convected quantities like temperature in
turbulent fluid: Part 1. General discussion and the case of small conductivity. *J Fluid*
Mech **5**, 113–133 (1959).
25.   Gower, J. F. R., Denman, K. L. & Holyer, R. J. Phytoplankton patchiness indicates the
fluctuation spectrum of mesoscale oceanic structure. *Nature* **288**, 157–159 (1980).
26.   Platt, T. Local phytoplankton abundance and turbulence. *Deep-Sea Research and*
Oceanographic Abstracts **19**, 183–187 (1972).
27.   Denman, K., Okubo, A. & Platt, T. The chlorophyll fluctuation spectrum in the sea,. *Limnol*
Oceanogr (1977) doi:10.4319/lo.1977.22.6.1033.
28.   Denman, K. & Platt, T. The variance spectrum of phytoplankton in a turbulent ocean. *J*
Mar Res **34**, 593–601 (1976).
29.   Bracco, A., Clayton, S. & Pasquero, C. Horizontal advection, diffusion, and plankton
spectra at the sea surface. *J Geophys Res Oceans* (2009) doi:10.1029/2007JC004671.
30.   Lévy, M. & Klein, P. Does the low frequency variability of mesoscale dynamics explain a
part of the phytoplankton and zooplankton spectral variability? *Proceedings of the Royal*
Society A: Mathematical, Physical and Engineering Sciences **460**, 1673–1687 (2004).
31.   Mahadevan, A. & Campbell, J. W. Biogeochemical patchiness at the sea surface.
Geophys Res Lett **29**, 1–4 (2002).
32.   Prairie, J. C., Sutherland, K. R., Nickols, K. J. & Kaltenberg, A. M. Biophysical
interactions in the plankton: A cross-scale review. *Limnology and Oceanography: Fluids*
and Environments **2**, 121–145 (2012).
33.   Martin, A. P. & Srokosz, M. A. Plankton distribution spectra: Inter-size class variability
and the relative slopes for phytoplankton and zooplankton. *Geophys Res Lett* **29**, 1998–
2001 (2002).
34.   Mackas, D. L. & Boyd, C. M. Spectral analysis of zooplankton spatial heterogeneity.
Science (1979) **204**, 62–64 (1979).
35.   van Gennip, S. Understanding the Extent of Universality in Phytoplankton Spatial
Properties. *Dissertation* (2014).
36.   Gorsky, G. *et al.* Expanding Tara Oceans Protocols for Underway, Ecosystemic Sampling
of the Ocean-Atmosphere Interface During Tara Pacific Expedition (2016–2018). *Front*
Mar Sci **6**, (2019).
37.   Boss, Twardowski, M. S. & Herring, S. Shape of the particulate beam attenuation
spectrum and its inversion to obtain the shape of the particulate size distribution. *Appl*
Opt **40**, 4885 (2001).
38.   Lehahn, Y., d'Ovidio, F. & Koren, I. A Satellite-Based Lagrangian View on Phytoplankton
Dynamics. *Ann Rev Mar Sci* **10**, 99–119 (2018).
39.   Lehahn, Y. *et al.* Dispersion/dilution enhances phytoplankton blooms in low-nutrient
waters. *Nat Commun* **8**, 14868 (2017).
40.   Agarwal, V., Chávez-Casillas, J., Inomura, K. & Mouw, C. B. Patterns in the temporal
complexity of global chlorophyll concentration. *Nat Commun* **15**, 1522 (2024).
41.   Doney, S. C., Glover, D. M., McCue, S. J. & Fuentes, M. Mesoscale variability of
Sea-viewing Wide Field-of-view Sensor (SeaWiFS) satellite ocean color: Global patterns
and spatial scales. *J Geophys Res Oceans* **108**, (2003).

42. Glover, D. M., Doney, S. C., Oestreich, W. K. & Tullo, A. W. Geostatistical Analysis of
Mesoscale Spatial Variability and Error in SeaWiFS and MODIS/Aqua Global Ocean
Color Data. *J Geophys Res Oceans* **123**, 22–39 (2018).

43. Longhurst, A. R. *Ecological Geography of the Sea. Ecological Geography of the Sea*
(2007). doi:10.1016/B978-0-12-455521-1.X5000-1.

44. LEVIN, S. A. & SEGEL, L. A. Hypothesis for origin of planktonic patchiness. *Nature* **259**,
659–659 (1976).

45. Kiørboe, T. *A Mechanistic Approach to Plankton Ecology.* (Princeton University Press,
2008).

46. Lombard, F. *et al.* Globally consistent quantitative observations of planktonic ecosystems.
Front Mar Sci **6**, (2019).

47. Sunagawa, S. *et al.* Structure and function of the global ocean microbiome. *Science*
(1979) **348**, (2015).

48. Denman, K. L. & Abbott, M. R. Time scales of pattern evolution from cross‐spectrum
analysis of advanced very high resolution radiometer and coastal zone color scanner
imagery. *J Geophys Res Oceans* **99**, 7433–7442 (1994).

49. Martin, A. P., Richards, K. J., Bracco, A. & Provenzale, A. Patchy productivity in the open
ocean. *Global Biogeochem Cycles* **16**, 9-1-9–9 (2002).

50. Slade, W. H. *et al.* Underway and moored methods for improving accuracy in
measurement of spectral particulate absorption and attenuation. *J Atmos Ocean Technol*
27, 1733–1746 (2010).

51. Boss, E. S., Collier, R., Pegau, W. S., Larson, G. & Fennel, K. Measurements of spectral
optical properties and their relation to biogeochemical variables and processes in Crater
Lake, Crater Lake National Park, OR. in *Long-term Limnological Research and*
Monitoring at Crater Lake, Oregon 149–159 (Springer Netherlands, Dordrecht, 2007).
doi:10.1007/978-1-4020-5824-0_9.

52. Roesler, C. S. & Barnard, A. H. Optical proxy for phytoplankton biomass in the absence
of photophysiology: Rethinking the absorption line height. *Methods in Oceanography* **7**,
79–94 (2013).

53. Haëntjens, N. & Boss, E. Inlinino: A Modular Software Data Logger for Oceanography.
Oceanography **33**, 80–84 (2020).

54. Boss *et al.* Inherent Optical Property Measurements and Protocols: Best practices for the
collection and processing of ship-based underway flow-through optical data. *IOCCG*
Ocean Optics and Biogeochemistry Protocols for Satellite Ocean Colour Sensor
Validation 1–23 (2019).