Lumped Parameter Simulations of Cervical Lymphatic Vessels Reveal Dynamics of Cerebrospinal Fluid Efflux from the Skull

Daehyun Kim[1] and Jeffrey Tithof[1*]

[1]Department of Mechanical Engineering, University of Minnesota, 111 Church St SE, Minneapolis, MN, 55455, United States.

*Corresponding author(s). E-mail(s): tithof@umn.edu;

Abstract

Background: Growing evidence suggests that for rodents, a substantial fraction of cerebrospinal fluid (CSF) drains by crossing the cribriform plate into the nasopharyngeal lymphatics, eventually reaching the cervical lymphatic vessels (CLVs). Disruption of this drainage pathway is associated with various neurological disorders.

Methods: We employ a lumped parameter method to numerically model CSF drainage across the cribriform plate to CLVs. Our model uses intracranial pressure as an inlet pressure and central venous blood pressure as an outlet pressure. The model incorporates initial lymphatic vessels (modeling those in the nasal region) that absorb the CSF and collecting lymphatic vessels (modeling CLVs) to transport the CSF against an adverse pressure gradient. To determine unknown parameters such as wall stiffness and valve properties, we utilize a Monte Carlo approach and validate our simulation against recent *in vivo* experimental measurements.

Results: Our parameter analysis reveals the physical characteristics of CLVs. Our results suggest that the stiffness of the vessel wall and the closing state of the valve are crucial for maintaining the vessel size and volume flow rate observed *in vivo*. Furthermore, we find that a decreased contraction amplitude and frequency leads to a reduction in volume flow rate. Additionally, we provide evidence that branching of initial lymphatic vessels may deviate from Murray's law to reduce sensitivity to elevated intracranial pressure.

Conclusions: This is the first numerical study of CSF drainage through CLVs. Our comprehensive parameter analysis offers guidance for future numerical modeling of CLVs. This study also provides a foundation for understanding physiology of CSF drainage, helping guide future experimental studies aimed at identifying causal mechanisms of reduction in CLV transport and potential therapeutic approaches to enhance flow.

Keywords: cervical lymphatic vessels, lymphatic system, cerebrospinal fluid, traumatic brain injury

Background

Cerebrospinal fluid (CSF) is a clear and colorless fluid that circulates around the brain and spinal cord, offering physical protection, acting as a "water cushion" to the central nervous system [1]. Additionally, it serves as a medium for supplying nutrients to the brain [2]. Beyond these traditional functions, CSF has gained growing recognition for its significant role in clearing metabolic wastes from the brain. This is

largely a consequence of two significant discoveries in neuroscience from the past decade: the glymphatic system [3] and meningeal lymphatic system [4]. The glymphatic system operates as a dynamic pathway where CSF enveloping the brain engages in a solute exchange with the interstitial fluid (ISF) deep within the brain's parenchyma. This intricate process involves CSF entering the brain through perivascular spaces and subsequently flushing waste from the parenchyma [5, 6]. This solute exchange between the CSF and ISF plays a significant role in clearing cerebral metabolic waste, including amyloid-β. Notably, the glymphatic system is most active during sleep [7]. In parallel, demonstrations of the importance of meningeal lymphatic vessels have illuminated this CSF drainage pathway as critical for clearing metabolic wastes from the brain. The meningeal lymphatic network pervades nearly the entire meninges (layers of tissue surrounding the brain) in mammals, serving as the conduit for draining immune cells, small molecules, and fluids (including ISF and CSF) from the central nervous system into deep cervical lymph nodes [8]. Together, these two transport pathways are integral to the brain's waste clearance system, and disruption has been closely tied to various neurological disorders, including Alzheimer's disease [5].

CSF is produced at choroid plexus, found within the ventricles of the brain [9]. Following its production, CSF traverses the intricate ventricular system and subsequently reaches the subarachnoid space (SAS) surrounding the brain and spinal cord. Traditionally, the prevailing belief was that CSF is absorbed by arachnoid granulations. However, advancements in experimental techniques have revealed that substantial CSF drains from the skull, eventually reaching the cervical lymphatic vessels (CLVs) [10–15]. The specific details of the route through which CSF reaches the CLVs is currently an area of active research and debate [16]. However, a substantial body of research – especially in mouse models – suggests that a substantial fraction of fluid follows a pathway through the cribriform plate along the cranial nerves, draining into lymphatic vessels that eventually reach the CLVs [14, 17–20].

Understanding fundamental aspects of CSF drainage to CLVs is significant due to its intricate association with various neurological disorders [21, 22]. Traumatic brain injury (TBI) also has been linked to CSF drainage disruption. In mouse models, research has revealed a significant reduction in glymphatic pathway activity following TBI [23]. Additionally, it has been demonstrated that CSF flow serves as a conduit for transporting biomarkers associated with neurodegenerative diseases into the bloodstream via the cervical lymphatics [24]. More recently, TBI was linked to weakening of CLV activity, impaired glymphatic transport, and exacerbation of brain edema, which the authors argued arose as a consequence of a post-TBI surge in norepinephrine [20]. Notably, administration of a norepinephrine antagonist cocktail led to restoration of CLV function, reduction of edema, and improved cognitive outcomes. These findings suggest TBI may directly impact CLV function, disrupting CSF drainage from the skull.

Despite the significant implications for advancing disease treatments, precise quantification of CSF drainage routes has proven challenging. Spatial and temporal limitations in imaging pose one of the

biggest challenges. Since lymphatic vessels are narrow and long, and span a wide region that can be difficult to optically access [25, 26], *in vivo* visualization of a large fraction of the lymphatic network that drains CSF is tremendously challenging. Moreover, the alteration of biological variables, such as intracranial pressure (ICP), during experimental testing has the potential to disrupt this delicate biological system [27]. As a consequence, these inherent constraints in experiments leave many open questions which can perhaps be addressed via numerical simulation.

Numerical simulations serve as efficient, safe, and ethically desirable alternatives to experiments; simulations also enable manipulation of parameters and the ability to prescribe/determine precise conditions that may be difficult or impossible to achieve experimentally, such as pressures inside vessels. Numerous prior studies have used a numerical approach to model lymphatic vessels. Reddy et al. pioneered this field by creating the initial mathematical model for branching lymphatic vessels [28]. Their one-dimensional (1D) model simulated laminar lymph flow across 35 lymphangions, representing the thoracic duct. Quick et al. employed a circuit-theory approach, commonly employed in the simulation of cardiovascular systems [29], to simulate both individual lymphangions and a chain of lymphangions. Bertram et al. extended this methodology and focused on a series of lymphangions, highlighting the efficiency of sequential contractions compared to synchronized contractions for fluid transport [30]. These foundational models lay the groundwork for simulating a large number of interconnected lymphangions, with incorporation of compliant walls and valves. Notably, these models were primarily based on parameters from the thoracic duct and mesenteric lymphatic vessels.

In parallel, two-dimensional (2D) or three-dimensional (3D) models have typically focused on a single lymphangion and often utilize fluid-structure interaction techniques [31–33]. Such studies have investigated the influence of active contractions driven by nitric oxide and calcium ions on fluid drainage. Validation of Poiseuille flow under substantial radial contractions was also studied. Computational cost has largely confined such studies to simulation of a single lymphangion, although some 2D/3D simulations for a series of lymphatic vessels, leveraging fluid-structure interaction approaches, have been achieved [34–36]. These studies tackled a range of factors, including the chain's length and its influence on mean flow rates, the effects of adverse pressure differences on flow rates, contraction timing, dynamics related to valve geometry impacting backflow, and the effects of valve elasticity. Again, it is worth highlighting that these simulations were based on lymphatic vessels in the other regions, such as mesenteric lymphatic vessels, and were not specifically tailored to CSF drainage through CLVs.

Currently, there is an absence of numerical models that predict fluid transport through CLVs. The goal of this study is to construct a comprehensive lumped parameter model that captures the intricate CSF drainage process in mice. In the absence of knowledge of the exact geometry and physical properties

of CLVs, we use a Monte Carlo approach to identify a parameter regime that matches closely with experimental data. Simultaneously, we perform a parameter sensitivity analysis to examine the characteristics of CLVs and how changes in these parameters affect the CLV diameter and net volume flow rate. Using our comprehensive model, we explore strategies to enhance CSF flow rate through CLVs by manipulating certain variables. Finally, we explore how the variation of ICP affects CSF efflux through the CLVs. This study is the first numerical model of CSF outflow through CLVs, providing novel insights into potential strategies to enhance CSF drainage. These findings provide guidance for future experimental research and serve as a valuable resource for optimizing CSF drainage strategies.

Methods

We use the lumped parameter method to model the nasal and cervical lymphatic system, which are treated as initial and collecting lymphatics. The versatility and computational efficiency makes this approach suitable for simulating the complexities and capturing the impact of uncertainties associated with this lymphatic network that connects the SAS to the end of the CLVs. Given the lack of detailed parameter values for CLVs in the literature (e.g., stiffness of the vessel wall, active tension due to smooth muscle cells, valve properties), this approach allows for computationally-efficient iterative determination of parameters to obtain agreement with experimental observations.

Modelling CSF drainage via CLVs

Lymphatic vessels typically consist of initial lymphatic vessels (lymphatic capillaries) and collecting lymphatic vessels [37]. The initial lymphatic vessels form the starting point of the lymphatic system and are responsible for absorbing ISF. These vessels consist of a single layer of endothelial cells, and their button-like junctions structure allows ISF and small particles to enter. As these interstitial lymphatic vessels merge, they give rise to larger lymphatic vessels known as collecting lymphatic vessels. Collecting lymphatic vessels consist of interconnected compartments called lymphangions which serve as the primary transporters of lymph within the lymphatic system. Unlike initial lymphatic vessels, collecting lymphatic vessels are covered by smooth muscle cells. There are two types of pumping mechanism in the collecting lymphatic vessels: extrinsic and intrinsic. Extrinsic pumping occurs by compression due to skeletal muscle contractions (external force), while intrinsic pumping refers to the transport arising from natural contractile activity of smooth muscle cells lining the lymphatic vessel lumen. In the periphery, a combination of extrinsic and intrinsic forces facilitate the movement of lymph against an adverse pressure gradient. The presence of valves between lymphangions ensures net transport of lymph by preventing backflow. Thus intrinsic and extrinsic pumping, along with the presence of valves enables collecting lymphatic vessels to efficiently transport lymph to lymph nodes.

Fig. 1: Comparative illustration and mathematical approach to modeling CSF drainage. (A) Illustration of the biological pathway for CSF drainage through CLVs. CSF in the SAS is directly absorbed by lymphatic capillaries that cross the cribriform plate. These capillaries merge to form larger vessels, which eventually connect to the CLVs. The CLVs then transport the CSF back into the central venous blood. (B) Schematic of the mathematical model of this CSF drainage pathway. The inlet pressure is the ICP, and the lymphatic capillaries are modeled using a net equivalent hydraulic resistance. The outlet pressure is assumed to be a constant, equal to that of central venous blood. CLVs are modeled to incorporate contractions and expansions of lymphangions, as well as opening and closing of valves in between; this portion is inspired by Bertram et al. [30].

We used this general lymphatic structure to model CSF outflow through CLVs. A schematic of the model is illustrated in Fig. 1A. Due to limited anatomical data and flow measurements for humans, we instead developed our model to capture murine anatomy. Even though the exact route connecting the CSF in SAS to the lymphatic capillaries is not fully established [16], recent experiments suggest that tiny lymphatics have direct connection to the SAS across the cribriform plate [17, 38–40]. Based on these findings, we assume there are direct connections through which the lymphatic capillaries absorb CSF from the SAS. These capillaries then merge to form larger vessels, adopting the principle that each parent vessel bifurcates into two daughter vessels of equal diameter in accordance with Murray's Law but with a different exponent value ($n = 1.45$) [41]. For simplicity, we assume that initial lymphatics are impermeable and absorption of CSF occurs at the tip of the vessels. CSF in the initial lymphatics is transported to the CLVs, then eventually pumped to the central venous blood. Fig. 1B summarizes the mathematical model that describes CSF efflux from the SAS to CLVs then venous blood. Our model is based on the lumped parameter method, which is analogous to an electrical circuit, so our simulation diagram includes circuit elements, including an equivalent resistor R_{initial} which captures the hydraulic resistance of the initial lymphatic vessels.

The size of the initial lymphatics located near the cribriform plate was estimated by analyzing images from Spera et al. [42] (tip diameter: 7.5 μm) and Norwood et al. [17] (length of total branches: 370 μm). By applying the modified Murray's Law ($d_i^{1.45} = d_{i+1}^{1.45} + d_{i+1}^{1.45}$), we calculated the diameters of parent vessels and number of branching generations until their diameter reached the median diameter of CLVs ($D = 84.1$ μm) as reported in Hussain et al. [20]. We assume that the branching of the CLVs begins at the first lymphangion, which bifurcates into two branche of equal diameter. We obtained $N = 5$ for the number of generations (compared to $N = 11$ for standard Murray's law with exponent 3). The modified exponent in our model results in a smaller generation ($N = 5$), leading to smaller diameters of the branches. Also, each branch becomes longer, as the total length of the entire branching structure is fixed. Subsequently, we computed the hydraulic resistance of this branching network under the assumption of Hagen-Poiseuille flow. Finally, the total hydraulic resistance was calculated using computation of series of parallel resistances as,

$$R_{\text{initial}} = \sum_{i=1}^{N} \frac{128\mu L/\pi d_i^4}{2^i} = \quad 2.64 \times 10^9 \quad \text{dyne s/mL cm}^2 \tag{1}$$

where i is the specific generation, μ is dynamic viscosity of the CSF, L is length of each generation of daughter capillaries, and d_i is the initial lymphatic vessel diameter at generation i.

Bertram et al's works are the inspiration for our approach to modeling CLVs [30, 43, 44]. We assume a collecting lymphatic vessel consists of five lymphangions, based on our estimates of 10 mm from the nasal mucosa to superficial cervical lymph node and a 2 mm lymphangion length [18, 20]. Each lymphangion, i, is characterized by time-dependent variables, namely, CLV diameter D_i, and pressures $P_{i,1}$, $P_{i,m}$, and $P_{i,2}$ representing the pressures at the inlet, midpoint, and outlet respectively. Additionally, the model includes the volumetric flow rates Q_i and Q_{i+1}, representing the inflow and outflow rates at each valve of a given lymphangion, respectively. The fluid flow from one chamber to the next chamber is described by the conservation of mass and momentum,

$$\frac{dD_i}{dt} = \frac{2(Q_i - Q_{i+1})}{\pi L D_i} \tag{2}$$

$$\frac{P_{i,1} - P_{i,m}}{L} = \frac{64\mu Q_i}{\pi D_i^4}, \quad \frac{P_{i,m} - P_{i,2}}{L} = \frac{64\mu Q_{i+1}}{\pi D_i^4} \tag{3}$$

under the quasi-steady assumption of fully developed Hagen-Poiseuille flow. This assumption is justified by the small Reynolds number ($Re = UD_0/\nu \approx 7 \times 10^{-3}$) and small Womersley number ($Wo = D_0\sqrt{2\pi f/\nu} \approx 0.05$) of flow in the CLVs, where $U = 64.8$ μm/s is the mean flow speed, $D_0 = 84.1$ μm is the mean CLV diameter, $\nu = 0.7 \times 10^{-6}$ m^2/s, and $f \approx 0.04$ Hz is the contraction frequency. Each

lymphangion is separated by valves with hydraulic resistance,

$$R_{V_{i+1}} = R_{V_{\min}} + R_{V_{\max}} \left(\frac{1}{1 + \exp(-s_{open}((P_{i,2} - P_{i+1,1}) - P_{open}))} \right), \tag{4}$$

where $R_{V_{\min}}$ and $R_{V_{\max}}$ are the respective minimum and maximum hydraulic resistance of the valves when open or closed, P_{open} is the minimum pressure required to open the valve, and s_{open} is the slope of the valve opening. A large s_{open} means a more rapid response of the valve opening to changes in pressure difference across the valve, while a smaller s_{open} indicates a more gradual response. The volume flow rate through valves is calculated based on the momentum equation,

$$Q_{i+1} = \frac{P_{i,2} - P_{i+1,1}}{R_{V_{i+1}}}. \tag{5}$$

The associated sets of differential algebraic equations are then closed by specifying the transmural pressure (i.e., pressure difference between inside and outside the lymphangion) using an equation that contains both passive and periodic contractile components:

$$P_{im} - P_e = P_d \left(\exp \left(\frac{D_i}{D_d} \right) - \left(\frac{D_d}{D_i} \right)^3 \right) + \frac{M}{D_i} \left(1 - \cos \left(2\pi f \left(t - t_{pi} \right) \right) \right), \tag{6}$$

where P_e is a uniform pressure external to the lymphangiongs, P_d and D_d are passive properties that account for mechanical characteristics of lymphanion's contractions and expansions, M is the active tension, and t_d is temporal phase between contractions of adjacent lymphangions (these parameters are discussed in more detail below). We assume the external pressure P_e is constant and slightly lower than the inlet and outlet pressures to make the transmural pressure operate more dynamically, similar to Bertram et al's work [30].

Parameter estimation

We first discuss parameters in the equations that are well-known based on literature and experimental observations, which are detailed in Table 1. The frequency of lymphangions' contractions, f, and the length of each lymphangion, L, are determined directly from in vivo images [20] using our recently published image analysis techniques [45]. The dynamic viscosity of CSF was used in our model [46]. We used the same values for the temporal phase between adjacent lymphangion contractions, t_d, and the slope of valve opening, s_{open}, as in previous studies [30, 44].

Estimating other parameters poses a substantial challenge, particularly for those not directly measurable, such as properties of the cervical lymphatic vessel walls (P_d, D_d), active tension generated by

Parameters	Definitions	Value	Unit	Reference
f	Contraction frequency	2.4	min^{-1}	[20]
L	Length of lymphangion	2	mm	[20]
n	Number of lymphangions	5		Assumed
t_d	Temporal phase between adjacent lymphangion contraction	90	degree	[44]
s_{open}	Slope of valve opening	0.2	cm^2/dyne	[43]
μ	Dynamic viscosity of CSF	0.007	Pa s	[46]
P_e	External pressure	3.75	mmHg	Assumed
ICP	Intracranial pressure	4	mmHg	[48, 49]
P_{venous}	Central venous pressure	6	mmHg	[50]
$R_{initial}$	Equivalent hydraulic resistance of lymphatic capillaries	3.21×10^8 (n = 3) 2.653×10^9 (n = 1.45)	dyne s/mL cm^2	[41]

Table 1: Reasonably well-known parameters for modeling CLVs.

smooth muscle cells (M), the minimum pressure required to open the valve (P_{open}), and hydraulic resistance of the valves ($R_{V_{\min}}, R_{V_{\max}}$). While more detailed explanation of these constants is available in the original works [30, 43, 44, 47], it is worthwhile to provide a brief explanation of these unknown properties here. P_d represents the vessel wall stiffness. A higher P_d value means that the transmural pressure reacts more sharply to any change in vessel diameter. D_d is the threshold diameter at which the vessel begins to experience positive or negative transmural pressure. When the lymphangion's diameter (D_i) surpasses D_d, the transmural pressure becomes positive, with the shape of an exponential curve. Conversely, when the lymphangion's diameter is less than D_d, the transmural pressure becomes negative, described by a cubic function. M represents the tension generated by smooth muscle cells around the vessel. The transmural pressure incorporates this active tension (M) as well as the vessel's passive properties (P_d and D_d) to account for the mechanical characteristics of lymphangion's contractions and expansions. P_{open} is the threshold pressure for the opening of valves between adjacent lymphangion. The valves are analogous to a diode in an electrical circuit. When the pressure difference across the valve exceeds P_{open}, the resistance of the valve becomes approximately $R_{V_{\min}}$. When the upstream pressure (pressure before the valve) is lower than the downstream pressure (pressure after the valve), the hydraulic resistance of the valve becomes approximately $R_{V_{\max}}$, meaning that if this value is very large, the valve is practically closed.

To overcome the challenge of not knowing the mechanical properties of CLVs, we adopt an inverse problem-solving approach based on a Monte Carlo method in which we vary parameters then compare model outputs to experimental measurements of the median vessel diameter (84.072 μm) and mean volume flow rate (0.0226 μl/min) [20]. This approach allows us to iteratively adjust and optimize our parameters by validating against the known outputs, aiming to achieve a more accurate and comprehensive model of the CSF drainage dynamics.

We set bounds for unknown parameters based on prior research on rat mesenteric lymphatic vessels [30, 43, 44, 47, 51]. The lower and upper bounds we set come from the minimum and maximum of reported values. We then expanded the bounds by subtracting or adding twice the standard deviation σ.

Parameters	Definitions	Lower Bound	Upper Bound	Unit
M	Active Tension	3.6	24.3	dyne/cm
R_{Vmin}	Minimum valve resistance	600	2.83×10^6	dyne s/mL cm^2
R_{Vmax}	Maximum valve resistance	1.2×10^7	2.82×10^{10}	dyne s/mL cm^2
p_{open}	Valve opening pressure	-70	-15	dyne/cm^2
P_d	Constitutive-relation constant	20	732	dyne/cm^2
D_d	Constitutive-relation constant	0.00845	0.0613	cm

Table 2: Bounds on unknown parameters, used in our Monte Carlo approach to modeling CLVs to determine parameter sets that lead to simulations that closely match average experimental measurements.

Our resulting bounds for unknown parameters are summarized in Table 2. We next generated a random uniform distribution for each of the unknown parameters in accordance with each range. 10,000 simulations were conducted based on these randomly sampled parameters, and results from the model were compared to the median diameter and mean volumetric flow rate of the CLVs as measured *in vivo*.

Diameters, pressures, and volumetric flow rates in the differential algebraic equations (1)-(3) coupled with valve and transmural pressure equations (4)-(6) were solved numerically. MATLAB's built-in solvers, *fsolve* and *fzero*, were used to find the transmural pressures of each lymphangion. Diameters were calculated by employing the 4th order Runge-Kutta method. These calculations continued until converged periodicity of variables was achieved. Finally, the volume flow rates at the valves were calculated based on the hydraulic resistance of the valve and pressure difference across it. We computed the median diameter $med(D_{sim})$ and mean volumetric flow rate $\overline{Q}_{sim}$ for each simulation, which we used as criteria for the search. If the comparison fell within 5% of the experimental value (denoted with subscript "exp"), such that,

$$\frac{\mid med(D_{sim}) - med(D_{exp}) \mid}{med(D_{exp})} \leq 0.05 \quad , \quad \frac{\mid \overline{Q}_{sim} - \overline{Q}_{exp} \mid}{\overline{Q}_{exp}} \leq 0.05 \tag{7}$$

we noted that particular parameter set. We continued this process for every randomly chosen parameter set and repeated the same numerical procedure. A summary of this process, including the Monte Carlo parameter selection, is depicted in Fig. 2. Note that we stored the results of all simulations (including those that did not meet our 5% criteria) for the sake of performing sensitivity analysis.

Fig. 2: Algorithm flow chart

Results

Inverse problem: Monte Carlo approach

We conducted a Monte Carlo simulation to estimate parameters that are consistent with experimental observations. Out of 10,000 simulations, only 585 successfully converged. The converged simulations indicate that M, D_d, and P_d must fall within a narrow range for simulations to successfully converge. To further explore the sensitivity, we narrowed the values of M, D_d, and P_d and performed an additional 2,000 simulations, randomly sampling parameters from within these revised limits (Fig. 3).

The parameter sensitivity analysis revealed that P_{open} and $R_{V_{min}}$ do not significantly impact the vessel diameter, as no clear trend in median diameter value emerged as each of these parameters were varied (Fig. 3A, C and Supplementary Fig. S1B, D). Values close to the experimental value we sought to match (84.1 μm), in a color indicated by the black arrow on the color bar, are randomly distributed throughout the parameter space. $R_{V_{max}}$ had a barely discernible influence on the diameter when $R_{V_{max}}$ was very small; this led to increases in diameter over 120 μm (see red circles at left in Fig. 3A). On the other hand, the parameters D_d, P_d, and M were found to influence vessel size, with a strong tendency for the diameter to increase with lower M, higher P_d, and higher D_d (Fig. 3B and Supplementary Fig. S1A, E, F). This indicates that the balance between the stiffness of the vessel wall and the active tension exerted by smooth muscle cells significantly affects the diameter of the CLV.

The mean volumetric flow rate through the CLVs was not greatly affected by P_{open} or $R_{V_{min}}$, but showed significant sensitivity to $R_{V_{max}}$ (Fig. 3C and Supplementary Fig. S2B, C, D). Very small $R_{V_{max}}$, in particular, resulted in a negative mean volume flow rate (Supplementary Fig. S2C) meaning that backflow exceeded forward flow. This suggests that the closure of the valve is crucial for prograde net transport,

Fig. 3: Results from the Monte Carlo search and analysis of parameter sensitivity. The target values for each criteria are indicated by black arrows on the color bars. (A) 3D scatter plots showing the variation of median diameter with changes in P_{open}, $R_{V_{min}}$, and $R_{V_{max}}$. The hydraulic resistance of the valves, $R_{V_{min}}$ and $R_{V_{max}}$ are normalized by their maximum values such that $[R_{V_{min}}] = R_{V_{min}}/max(R_{V_{min}})$ and $[R_{V_{max}}] = R_{V_{max}}/max(R_{V_{max}})$ (see text for values). (B) 3D scatter plots showing the variation of median diameter with changes in D_d, P_d, and M. (C) 3D scatter plot showing the variation of mean volume flow rate $\overline{Q}$ with changes in P_{open}, $R_{V_{min}}$, and $R_{V_{max}}$. (D) 3D scatter plot showing the variation of mean volume flow rate with changes in D_d, P_d, and M. For (C) and (D), volume flow rates below 0 are plotted as 0 to emphasize the sensitivity among positive values.

even more so than the pressure difference required to open the valve or the hydraulic resistance of the open valve (for the parameter ranges considered here, anyway). We also found the mean volume flow rate increases with higher M, lower P_d, and is insensitive to changes in D_d (Fig. 3D and Supplementary Fig. S2A, E, F). Hence, higher active tension along the vessel wall and reduced vessel stiffness increase net transport. This suggests that active properties are more important than passive properties for enhancing transport, and this is achieved by increasing vessel wall compliance with respect to the action of smooth muscle cells. A more detailed analysis of intrinsic pumping is discussed in the next section.

From this analysis, we identified the following parameters that lead to simulation results that align closely with experimental data: $M = 20.86$ dyne/cm, $R_{V_{min}} = 7.43 \times 10^5$ dyne s/mL cm^2, $R_{V_{max}} = 2.03 \times 10^{10}$ dyne s/mL cm^2, $P_{open} = -45$ dyne/cm^2, $P_d = 521$ dyne/cm^2, and $D_d = 0.015$ cm. This parameter set will be used below for further analysis.

CSF efflux through CLVs

Fig. 4: CSF drainage through CLVs. (A) Illustration of the geometry of the simulation. A red and a royal blue X indicate the locations at which pressure upstream (ICP) and downstream (venous blood), respectively, are plotted in (B). Additional X symbols inside each lymphangion is color-coded to indicate the locations of pressure and diameter in (B)-(D); similarly, arrows inside each valve are color-coded to indicate volume flow rate in (B)-(D). (B) Plot of pressure versus time. We assume constant pressure for the inlet (red dashed line) and outlet (royal blue dashed line). (C) Plot of diameters versus time. (D) Plot of volumetric flow rate versus time. Note the large cyan peak corresponds to flow through the last valve, furthest right in (A).

The adverse pressure difference that CLVs must overcome is higher than other lymphatic regions. The CSF must be transported against this adverse pressure, and this is enabled through intrinsic pumping in the presence of bileaflet valves in the lymphangions. Fig. 4A-D illustrate this process. The ICP is indicated in red, while central venous blood pressure is plotted in royal blue.

Let's examine the first (blue in Fig. 4A) and second (orange in Fig. 4A) lymphangions. Initially, the pressure in the first lymphangion is lower than the second lymphangion (Fig. 4B). As the pressure inside the first lymphangion begins to exceed that of the second lymphangion (Fig. 4B) and surpasses the additional threshold pressure needed to open the valve (p_{open}), CSF is transported from the first to the

Fig. 5: Influence of intrinsic pumping, including the active tension of smooth muscle cells (M) and the contraction frequency (f). (A) Contraction amplitude (left y-axis) and mean volume flow rate $\overline{Q}$ (right y-axis) as a function of M. (B) Volume flow rate $\overline{Q}$ as a function of f.

second lymphangion (Fig. 4D) as the first lymphangion contracts (Fig. 4C) and opens the valve between them. This results in the expansion of the second lymphangion as the CSF flows in. However, this forward flow is not continuous. The pressure in the second lymphangion exceeds that of the first lymphangion again (Fig. 4B), causing the valve between them close, leading to a small but non-negligible backflow from the second to the first lymphangion. The cycle of expansion and contraction of each lymphangion, driven by oscillatory active tension from smooth muscle cells (intrinsic pumping) and the opening and closing of the valves, enable CSF transport against a relatively large adverse pressure difference ($\sim$2 mmHg), ultimately facilitating fluid drainage into the central venous blood.

An interesting flow is observed at the inlet and outlet of the CLV (Fig. 4D). The volumetric flow rate at the first lymphangion is positive over a wide region (bold blue curve in Fig. 4D), due to the presence of large initial lymphatics attached to it. The addition of the net hydraulic resistance of the initial lymphatics results in a gradual change in the volumetric flow rate. The impact of the net hydraulic resistance of the initial lymphatic vessels will be discussed in more detail in the following section. The outflow from the last lymphangion to the central venous blood is narrow but high (light blue curve in Fig. 4D). This is due to the higher contraction ratio of the last lymphangion compared to the other lymphangions. It is important to note that the time-averaged volume flow rate across all lymphangions remains consistent, in accordance with the mass conservation.

Effect of intrinsic pumping

Lymphatic vessels rely on both intrinsic and extrinsic pumping mechanisms to transport fluid. We focus only on the impact of intrinsic pumping for CSF efflux from the brain, as our prior simulations suggest experimentally observed volume flow rates can be explained through intrinsic pumping alone [20]. Active tension generated by smooth muscle cells in response to stimuli, such as adrenergic and nitric

oxide signals, is the primary driver of intrinsic pumping, regulating lymph flow. As active tension (M) increases, lymphangion contraction amplitude, which we define as the difference between the maximum and minimum diameter divided by the maximum diameter ($\frac{D_{max}-D_{min}}{D_{max}}$), also increases (blue curve in Fig. 5A). The orange curve in Fig. 5A illustrates the relationship between mean volumetric flow rate and M, indicating that a higher value of M (i.e., higher contraction amplitude) results in greater net transport.

In Fig. 5A, two key points stand out. First, at $M = 14$ dyne/cm, as M increases in the $10 - 14$ range, the contraction amplitude increases slowly, and the mean volumetric flow rate remains nearly steady. This is because the active tension is too low to significantly affect contraction compared to the lymphangion's passive properties. In the $15 - 20$ range for M, the contraction amplitude increases dramatically as active tension dominates the lymphangion's passive properties. Beyond M of about 21, the increase in contraction amplitude slows down, suggesting volume flow rate may stabilize beyond this point. This is because the vessel diameter remains small when active tension is large (and hydraulic resistance depends sensitively on diameter, scaling with D^4). This indicates that excessively high tension does not necessary promote CSF transport, and the the optimal range is likely around 20 $dyne/cm$.

Next, we fixed $M = 20.86$ dyne/cm and adjusted the contraction frequency, f. Initially, the mean volume flow rate rose sharply for frequencies in the approximate range $2 - 8$ min^{-1}. For frequencies in the range $8 - 22$ min^{-1}, the volume flow rate increased more gradually. However, at about 27 min^{-1}, the volume flow rate began to decrease slightly. This suggests that increased contraction frequency only increases net transport over a finite range. Detailed inspection of these results suggests that when the lymphangions contract too quickly, the valves don't have enough time to close completely at these high frequency.

Effect of lymphatic capillary branching assumptions

In our final test, we examined how lymphatic capillary branching affects net volumete flow rate. We applied an inlet pressure corresponding to either normal ICP (4 mmHg) or elevated ICP (10 mmHg), as indicated in Fig. 6A. We then tested two different branching scenarios: one adhering to the typical Murray's law (with exponent 3) and another corresponding to Murray's law with exponent 1.45 [41]. The model with the smaller exponent led to smaller vessel diameters, but longer length, across each bifurcation compared to the normal case, ultimately generating a higher hydraulic resistance $R_{initial}$.

At normal ICP, both the diameter and volume flow rate for the two cases of different exponents operate within a physiological range. The diameter and volume flow rate for typical Murray's law are only slightly higher than those for the modified Murray's law (Fig.6B,C). However, at elevated ICP, the differences for the two different exponent cases become substantial. Both the diameter and volume

Fig. 6: Impact of lymphatic capillary branching on diameter and volume flow rate under different ICP conditions. (A) Plots of inlet pressure (ICP) for a normal (left) and elevated (right) case. (B) CLV diameter versus time for different ICP. The exponent in Murray's law is $n = 1.45$ (blue curve) or $n = 3$ (orange curve). (C) Volumete flow rate versus time for different ICP.

flow rate increase for both exponent cases (Fig.6B and C), but the model with modified Murray's law ($n = 1.45$, blue curve) shows only a small increase in diameter which spans a range of about 100 to 130 µm. On the other hand, the model with normal Murray's law ($n = 3$, orange curve) exhibits large amplitude fluctuations, ranging from about 130 to 250 µm. Physiologically, such a large peak diameter and range in diameter is unrealistic and could likely lead to vessel rupture when ICP increases. The volume flow rate also increases with the elevated ICP, as expected when the upstream pressure exceeds the downstream pressure. However, the model with the modified Murray's law ($n = 1.45$, blue curve) shows a gradual increase in volume flow rate, while the model with normal Murray's law ($n = 3$, orange curve) exhibits an extreme increase. Excessive outflow due to elevated ICP would likely have deleterious effects. These results suggest that lymphatic capillaries that absorb the CSF in the SAS likely do not follow normal Murray's law but may instead adhere to a modified version with a smaller exponent ($n < 3$) so that the associated increase in hydraulic resistance will buffer transport rates with the respect to changes in ICP.

Discussion

Lymphatic vessels transport fluid against an adverse pressure gradient, and CLVs are no exception. In fact, CLVs must overcome a higher adverse pressure gradient compared to other lymphatics [48, 49], which is a consequence of their anatomy which bridges the skull and cervical lymph nodes. Our numerical model suggests that the both active and passive properties of the vessel wall and the hydraulic resistance of the valve in its closed state are crucial for maintaining the vessel's size and volume flow rate observed via *in vivo* measurements [20]. Compared to mesenteric lymphatic vessels, CLVs exhibit a stronger sensitivity to wall stiffening (D_d and P_d) with changes in diameter at both positive and negative transmural pressures [30]. Due to this stiffer wall, the smooth muscle cells need to generate stronger active tension (M) to transport CSF. It was also found that the system is more sensitive to maximum valve resistance ($R_{V_{max}}$) than to minimum valve resistance ($R_{V_{min}}$). This implies that the valves must close completely to overcome the high pressure gradient and prevent backflow. Our model suggests that increased vessel/valve tissue rigidity, as occurs with aging [52], may impair CLV net drainage, perhaps reducing clearance of protein waste such as amyloid-β and tau, contributing to progression of neurodegenerative diseases.

As a lymphangion expands, the pressure inside it decreases, allowing cerebrospinal fluid (CSF) to enter. As it contracts, the CSF is transported to the next lymphangion, opening the valve in between and eventually draining to the central venous blood. It is noteworthy that the average diameters of each of the five lymphangions we model increase as one moves in the downstream direction. This is because the same parameters were used for all five lymphangions and the transmural pressure difference increases along the streamwise direction. Parameters that control the passive properties of the wall, such as distensibility (D_d) and the pressure at which the vessel is fully distended (P_d), can be modeled to increase along the downstream direction [30, 43] in order to maintain the same average diameters.

In the study by Hussain et al. it was observed that the CSF outflow through CLVs decreases following TBI [20]. This study also found that the level of norepinephrine increased after TBI, which was associated with a decrease in the contraction amplitude and frequency of the CLVs. Our simulations support the idea that decreased contraction amplitude and frequency lead to a decreased volume flow rate. This suggests that enhancing lymphatic activity could be a target for restoring CSF drainage. Lymphatic vessels transport fluid using both intrinsic and extrinsic pumping mechanisms. Currently, our model only includes intrinsic pumping which captures net volume flow rates that are in good agreement with experimental measurements. However, future numerical and/or experimental work may investigate the role of external pressures, such as those exerted by skeletal muscles or arising from neck massage, which may enhance drainage.

The ICP, which was assumed as the inlet pressure in our model, can increase due to gross swelling or intracranial bleeding after TBI [53]. One may hypothesize that this increased ICP could lead to rapid drainage of CSF from the skull. However, our simulations indicate that if the initial lymphatics that absorb the CSF bifurcate according to a modified version of Murray's law (with exponent substantially less than the typical $n = 3$), the associated increase in hydraulic resistance will act as a buffer that inhibits rapid CSF drainage. Murray's law, with an exponent of 3, was formulated to describe the bifurcations of arteries, aiming to explain how arterial branching minimizes hydraulic resistance. Our simulations suggest that minimizing hydraulic resistance in the initial lymphatic branches connecting the skull and CLVs may not be necessary. Future experiments could test for evidence of this safety mechanism by directly measuring the diameters of sequential initial lymphatic vessels or even by counting the number of branching generations. Our calculations suggest that approximately 11 versus 5 generations exist for Murray's law exponents of $n = 3$ versus $n = 1.45$.

Substantial evidence indicates that CSF drains through the cribriform plate to nasal lymphatics in rodents [16]. However, this pathway appears less important in humans [54], highlighting critical differences that may exist in CSF pathways for rodents versus humans. Meningeal lymphatics may potentially serve as a primary efflux route in humans. In our model, the initial lymphatic vessels are currently based on the nasopharyngeal lymphatic vessels of rodents. However, our model can be adapted to capture efflux to any other lymphatic vessels with different dimensions and properties, including meningeal lymphatic vessels.

In the recent study by Yoon et al., a complex network of lymphatics draining CSF through the nasopharynx to the cervical lymph nodes was identified [55]. This plexus contains tiny lymphatic vasculature at upstream locations, which merge into larger vessels. These larger vessels are considered pre-collector lymphatic vessels due to the presence of valves and somewhat loose covering of smooth muscle cells. Currently, our model does not include pre-collector lymphatic vessels. Future studies could incorporate a model of the nasopharyngeal plexus and test its hydraulic resistance, which may further buffer changes in ICP. Additionally, the presence of valves in the plexus supports the idea that hydraulic resistance of the valve in its closed state is critical for CLVs when the pressure difference is high. Although the closure of lymphatic valves is not perfect and does allows a small amount of backflow, the series of valves located in the plexus likely help reduce backflow, allowing the CLVs to maintain an optimal diameter and volume flow rate.

Several important limitations of this study should be acknowledged. Spatial resolution is limited in our simulation which prevents, for example, detailed simulation of variation in contraction amplitude along the length of each lymphangion. As a consequence of using lumped parameter modeling, our predictions only include volume flow rates and pressures for three points per lymphangion (one at the center, one at

each valve). However, a more detailed treatment is precluded by the current scarcity of quantitative data characterizing CLV geometry and physical properties. A further limitation is that the lumped parameter approach assumes that contractions of adjacent lymphangions occur with some fixed temporal phase (parameter t_d in Table 1). *In vitro* observations of CLV contractions following exposure to norepinephine reveal loss of contraction entrainment [20], which cannot be readily modeled using the lumped parameter approach. Nonetheless, the lumped parameter method is appealing in that simulations are fast and our Monte Carlo approach could be conducted efficiently. Overall, our study has uncovered valuable insights into not only fundamentals of the physical properties of CLVs, but also potential therapeutic strategies.

Conclusions

In this study, we performed comprehensive simulations of CSF efflux to cervical lymphatic vessels using a lumped parameter model. This model captures the CSF drainage from the subarachnoid space, absorbed by the initial lymphatics embedded in the nasal region, which merge over several generations and drain to the cervical lymphatic vessels, eventually reaching the central venous blood. We identified parameters that are unknown and difficult to experimentally measure using a Monte Carlo search in which we matched simulation predictions to *in vivo* measurements. Simultaneously, we explored how the unknown parameters in the governing equations affect the median diameter of the vessels and the mean volumetric flow rate, concluding that magnitude of active tension along the wall, passive properties of the vessel wall, and hydraulic resistance of the valve in its closed state have the greatest effect. We also demonstrated that increasing the active tension along the vessel wall and/or the contraction frequency increases the overall volume flow rate, but only to a limited extent. We also tested how the bifurcations of the upstream lymphatic capillaries affect the overall flow in response to elevated ICP. Smaller and longer branches (arising from a modified form of Murray's law with exponent 1.45) increase the net hydraulic resistance, increasing the robustness of the system to elevated ICP.

This is the first numerical study of CSF drainage through CLVs. We anticipate that our rigorous parameter analysis using our proposed model will help guide future numerical studies aimed at modeling CLVs under physiological and pathological conditions. Additionally, our results form a foundation for future experiments in this field, contributing to our understanding of CSF drainage and its potential therapeutic applications.

Abbreviations

CSF: cerebrospinal fluid; CLV: cervical lymphatic vessel; TBI: traumatic brian injury; ICP: intracranial pressure; SAS: subarachnoid space

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Data and materials are available from the authors following any reasonable request.

Competing interests

The authors declare no competing interests.

Funding

This work is supported by a Career Award at the Scientific Interface from Burroughs Wellcome Fund and the Research & Innovation Office at University of Minnesota.

Authors' contributions

Conceptualization: DK, JT; Methodology: DK, JT; Formal analysis and investigation: DK; Writing - original draft preparation: DK; Writing - review and editing: JT; Funding acquisition: JT; Resources: JT; Supervision: JT

Acknowledgements

Not applicable.

Authors' information

Authors and Affiliations

Daehyun Kim and Jeffrey Tithof

Mechanical Engineering, University of Minnesota

Corresponding author

Correspondence to Jeffrey Tithof

References

[1] Spector, R., Snodgrass, S.R., Johanson, C.E.: A balanced view of the cerebrospinal fluid composition and functions: Focus on adult humans. Exp. Neurol. **273**, 57–68 (2015)

[2] Veening, J.G., Barendregt, H.P.: The regulation of brain states by neuroactive substances distributed via the cerebrospinal fluid; a review. Cerebrospinal Fluid Res. **7**(1), 1–16 (2010)

[3] Iliff, J.J., Wang, M., Liao, Y., Plogg, B.A., Peng, W., Gundersen, G.A., Benveniste, H., Vates, G.E., Deane, R., Goldman, S.A., *et al.*: A paravascular pathway facilitates csf flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid β. Sci. Transl. Med. **4**(147), 147–111147111 (2012)

[4] Louveau, A., Smirnov, I., Keyes, T.J., Eccles, J.D., Rouhani, S.J., Peske, J.D., Derecki, N.C., Castle, D., Mandell, J.W., Lee, K.S., *et al.*: Structural and functional features of central nervous system lymphatic vessels. Nature **523**(7560), 337–341 (2015)

[5] Jessen, N.A., Munk, A.S.F., Lundgaard, I., Nedergaard, M.: The glymphatic system: a beginner's guide. Neurochem. Res. **40**, 2583–2599 (2015)

[6] Mestre, H., Mori, Y., Nedergaard, M.: The brain's glymphatic system: current controversies. Trends Neurosci. **43**(7), 458–466 (2020)

[7] Xie, L., Kang, H., Xu, Q., Chen, M.J., Liao, Y., Thiyagarajan, M., O'Donnell, J., Christensen, D.J., Nicholson, C., Iliff, J.J., *et al.*: Sleep drives metabolite clearance from the adult brain. Science **342**(6156), 373–377 (2013)

[8] Louveau, A., Plog, B.A., Antila, S., Alitalo, K., Nedergaard, M., Kipnis, J., *et al.*: Understanding the functions and relationships of the glymphatic system and meningeal lymphatics. J. Clin. Invest.

127(9), 3210–3219 (2017)

[9] Brown, P., Davies, S., Speake, T., Millar, I.: Molecular mechanisms of cerebrospinal fluid production. Neuroscience **129**(4), 955–968 (2004)

[10] Bradbury, M., Cole, D.: The role of the lymphatic system in drainage of cerebrospinal fluid and aqueous humour. J. Physiol. **299**(1), 353–365 (1980)

[11] Brinker, T., Lüdemann, W., Rautenfeld, D., Samii, M.: Dynamic properties of lymphatic pathways for the absorption of cerebrospinal fluid. Acta Neuropath. **94**, 493–498 (1997)

[12] Cserr, H.F., Harling-Berg, C.J., Knopf, P.M.: Drainage of brain extracellular fluid into blood and deep cervical lymph and its immunological significance. Brain Pathol. **2**(4), 269–276 (1992)

[13] Walter, B.A., Valera, V., Takahashi, S., Ushiki, T.: The olfactory route for cerebrospinal fluid drainage into the peripheral lymphatic system. Neuropath. Appl. Neuro. **32**(4), 388–396 (2006)

[14] Ma, Q., Ineichen, B.V., Detmar, M., Proulx, S.T.: Outflow of cerebrospinal fluid is predominantly through lymphatic vessels and is reduced in aged mice. Nat. Commun. **8**(1), 1434 (2017)

[15] Ahn, J.H., Cho, H., Kim, J.-H., Kim, S.H., Ham, J.-S., Park, I., Suh, S.H., Hong, S.P., Song, J.-H., Hong, Y.-K., *et al.*: Meningeal lymphatic vessels at the skull base drain cerebrospinal fluid. Nature **572**(7767), 62–66 (2019)

[16] Proulx, S.T.: Cerebrospinal fluid outflow: a review of the historical and contemporary evidence for arachnoid villi, perineural routes, and dural lymphatics. Cell. Mole. Life Sci. **78**(6), 2429–2457 (2021) https://doi.org/10.1007/s00018-020-03706-5

[17] Norwood, J.N., Zhang, Q., Card, D., Craine, A., Ryan, T.M., Drew, P.J.: Anatomical basis and physiological role of cerebrospinal fluid transport through the murine cribriform plate. eLife **8**, 44278 (2019) https://doi.org/10.7554/eLife.44278 . Publisher: eLife Sciences Publications, Ltd

[18] Decker, Y., Krämer, J., Xin, L., Müller, A., Scheller, A., Fassbender, K., Proulx, S.T.: Magnetic resonance imaging of cerebrospinal fluid outflow after low-rate lateral ventricle infusion in mice. JCI Insight **7**(3) (2022)

[19] Spera, I., Cousin, N., Ries, M., Kedracka, A., Castillo, A., Aleandri, S., Vladymyrov, M., Mapunda, J.A., Engelhardt, B., Luciani, P., et al.: Open pathways for cerebrospinal fluid outflow at the cribriform plate along the olfactory nerves. EBioMedicine **91** (2023)

[20] Hussain, R., Tithof, J., Wang, W., Cheetham-West, A., Song, W., Peng, W., Sigurdsson, B., Kim, D., Sun, Q., Peng, S., *et al.*: Potentiating glymphatic drainage minimizes post-traumatic cerebral oedema. Nature **623**(7989), 992–1000 (2023)

[21] Olsson, B., Lautner, R., Andreasson, U., Öhrfelt, A., Portelius, E., Bjerke, M., Hölttä, M., Rosén, C., Olsson, C., Strobel, G., *et al.*: Csf and blood biomarkers for the diagnosis of alzheimer's disease: a systematic review and meta-analysis. Lancet Neurol. **15**(7), 673–684 (2016)

[22] Engelhardt, B., Carare, R.O., Bechmann, I., Flügel, A., Laman, J.D., Weller, R.O.: Vascular, glial, and lymphatic immune gateways of the central nervous system. Acta Neuropathol. **132**, 317–338 (2016)

[23] Iliff, J.J., Chen, M.J., Plog, B.A., Zeppenfeld, D.M., Soltero, M., Yang, L., Singh, I., Deane, R., Nedergaard, M.: Impairment of glymphatic pathway function promotes tau pathology after traumatic brain injury. J. Neurosci. **34**(49), 16180–16193 (2014)

[24] Plog, B.A., Dashnaw, M.L., Hitomi, E., Peng, W., Liao, Y., Lou, N., Deane, R., Nedergaard, M.: Biomarkers of traumatic injury are transported from brain to blood via the glymphatic system. J. Neurosci. **35**(2), 518–526 (2015)

[25] Lohrberg, M., Wilting, J.: The lymphatic vascular system of the mouse head. Cell Tissue Res. **366**(3), 667–677 (2016)

[26] Aspelund, A., Antila, S., Proulx, S.T., Karlsen, T.V., Karaman, S., Detmar, M., Wiig, H., Alitalo, K.: A dural lymphatic vascular system that drains brain interstitial fluid and macromolecules. J. Exp. Med. **212**(7), 991–999 (2015) https://doi.org/10.1084/jem.20142290

[27] Xiang, T., Feng, D., Zhang, X., Chen, Y., Wang, H., Liu, X., Gong, Z., Yuan, J., Liu, M., Sha, Z., *et al.*: Effects of increased intracranial pressure on cerebrospinal fluid influx, cerebral vascular hemodynamic indexes, and cerebrospinal fluid lymphatic efflux. Journal Cerebr. Blood F. Met. **42**(12), 2287–2302 (2022)

[28] Reddy, N.P., Krouskop, T.A., Newell Jr, P.H.: A computer model of the lymphatic system. Comput. Biol. Med. **7**(3), 181–197 (1977)

[29] Quick, C.M., Venugopal, A.M., Dongaonkar, R.M., Laine, G.A., Stewart, R.H.: First-order approximation for the pressure-flow relationship of spontaneously contracting lymphangions. Am. J. Physiol.-Heart C. **294**(5), 2144–2149 (2008)

[30] Bertram, C.D., Macaskill, C., Moore, J.E.: Simulation of a Chain of Collapsible Contracting Lymphangions With Progressive Valve Closure. J. Biomech. Eng. **133**(1), 011008 (2011) https://doi.org/10.1115/1.4002799

[31] Li, H., Mei, Y., Maimon, N., Padera, T.P., Baish, J.W., Munn, L.L.: The effects of valve leaflet mechanics on lymphatic pumping assessed using numerical simulations. Sci. Rep. **9**(1), 10649 (2019)

[32] Baish, J.W., Kunert, C., Padera, T.P., Munn, L.L.: Synchronization and random triggering of lymphatic vessel contractions. PLoS Comput. Biol. **12**(12), 1005231 (2016)

[33] Rahbar, E., Moore Jr, J.E.: A model of a radially expanding and contracting lymphangion. J. Biomech. **44**(6), 1001–1007 (2011)

[34] Elich, H., Barrett, A., Shankar, V., Fogelson, A.L.: Pump efficacy in a two-dimensional, fluid–structure interaction model of a chain of contracting lymphangions. Biomech. Model. Mechan. **20**, 1941–1968 (2021)

[35] Ballard, M., Wolf, K.T., Nepiyushchikh, Z., Dixon, J.B., Alexeev, A.: Probing the effect of morphology on lymphatic valve dynamic function. Biomech. Model. Mechan. **17**, 1343–1356 (2018)

[36] Wolf, K.T., Dixon, J.B., Alexeev, A.: Fluid pumping of peristaltic vessel fitted with elastic valves. J. Fluid Mech. **918**, 28 (2021)

[37] Moore Jr, J.E., Bertram, C.D.: Lymphatic system flows. Annu. Rev. Fluid Mech. **50**, 459–482 (2018)

[38] Hsu, M., Rayasam, A., Kijak, J.A., Choi, Y.H., Harding, J.S., Marcus, S.A., Karpus, W.J., Sandor, M., Fabry, Z.: Neuroinflammation-induced lymphangiogenesis near the cribriform plate contributes to drainage of CNS-derived antigens and immune cells. Nat. Commun. **10**(1), 229 (2019) https://doi.org/10.1038/s41467-018-08163-0 . Number: 1 Publisher: Nature Publishing Group

[39] Furukawa, M., Shimoda, H., Kajiwara, T., Kato, S., Yanagisawa, S.: Topographic study on nerve-associated lymphatic vessels in the murine craniofacial region by immunohistochemistry and electron microscopy. Biomed. Res. **29**(6), 289–296 (2008)

[40] Antila, S., Karaman, S., Nurmi, H., Airavaara, M., Voutilainen, M.H., Mathivet, T., Chilov, D., Li, Z., Koppinen, T., Park, J.-H., *et al.*: Development and plasticity of meningeal lymphatic vessels. J. Exp. Med. **214**(12), 3645–3667 (2017)

[41] Talkington, A.M., Davis, R.B., Datto, N.C., Goodwin, E.R., Miller, L.A., Caron, K.M.: Dermal Lymphatic Capillaries Do Not Obey Murray's Law. Front. Cardiovasc. Med. **9**, 840305 (2022) https:

//doi.org/10.3389/fcvm.2022.840305

[42] Spera, I., Cousin, N., Ries, M., Kedracka, A., Castillo, A., Aleandri, S., Vladymyrov, M., Mapunda, J.A., Engelhardt, B., Luciani, P., Detmar, M., Proulx, S.T.: Open pathways for cerebrospinal fluid outflow at the cribriform plate along the olfactory nerves. eBioMedicine **91**, 104558 (2023) https://doi.org/10.1016/j.ebiom.2023.104558

[43] Bertram, C.D., Macaskill, C., Davis, M.J., Moore, J.E.: Development of a model of a multi-lymphangion lymphatic vessel incorporating realistic and measured parameter values. Biomech. Model. Mechan. **13**(2), 401–416 (2014) https://doi.org/10.1007/s10237-013-0505-0

[44] Jamalian, S., Bertram, C.D., Richardson, W.J., Moore, J.E.: Parameter sensitivity analysis of a lumped-parameter model of a chain of lymphangions in series. Am. J. Physiol.-Heart C. **305**(12), 1709–1717 (2013) https://doi.org/10.1152/ajpheart.00403.2013

[45] Kim, D., Gan, Y., Nedergaard, M., Kelley, D.H., Tithof, J.: Image analysis techniques for in vivo quantification of cerebrospinal fluid flow. Exp. Fluids **64**(11), 181 (2023)

[46] Tithof, J., Boster, K.A., Bork, P.A., Nedergaard, M., Thomas, J.H., Kelley, D.H.: A network model of glymphatic flow under different experimentally-motivated parametric scenarios. iScience **25**(5) (2022)

[47] Bertram, C.D., Macaskill, C., Davis, M.J., Moore, J.E.: Consequences of intravascular lymphatic valve properties: a study of contraction timing in a multi-lymphangion model. Am. J. Physiol.-Heart C. **310**(7), 847–860 (2016) https://doi.org/10.1152/ajpheart.00669.2015

[48] Yang, B., Zador, Z., Verkman, A.: Glial cell aquaporin-4 overexpression in transgenic mice accelerates cytotoxic brain swelling. J. Biol. Chem. **283**(22), 15280–15286 (2008)

[49] Moazen, M., Alazmani, A., Rafferty, K., Liu, Z.-J., Gustafson, J., Cunningham, M.L., Fagan, M.J., Herring, S.W.: Intracranial pressure changes during mouse development. J. Biomech. **49**(1), 123–126 (2016)

[50] Xie, C., Wei, W., Zhang, T., Dirsch, O., Dahmen, U.: Monitoring of systemic and hepatic hemodynamic parameters in mice. JoVE-J. Vis. Exp. (92), 51955 (2014)

[51] Bertram, C.D., Macaskill, C., Davis, M.J., Moore, J.E.: Contraction of collecting lymphatics: organization of pressure-dependent rate for multiple lymphangions. Biomech. Model. Mechan. **17**(5), 1513–1532 (2018) https://doi.org/10.1007/s10237-018-1042-7

[52] González-Loyola, A., Petrova, T.V.: Development and aging of the lymphatic vascular system. Adv. Drug Deliver. Rev. **169**, 63–78 (2021)

[53] Dunn, L.T.: Raised intracranial pressure. J. Neurol. Neurosur. Ps. **73**(suppl 1), 23–27 (2002)

[54] Ringstad, G., Eide, P.K.: Glymphatic-lymphatic coupling: assessment of the evidence from magnetic resonance imaging of humans. Cell. Mol. Life Sci. **81**(1), 1–14 (2024)

[55] Yoon, J.-H., Jin, H., Kim, H.J., Hong, S.P., Yang, M.J., Ahn, J.H., Kim, Y.-C., Seo, J., Lee, Y., McDonald, D.M., et al.: Nasopharyngeal lymphatic plexus is a hub for cerebrospinal fluid drainage. Nature, 1–10 (2024)

Supplementary information

Fig. S1: Supplementary analysis of parameter sensitivity in median diameter. All data points are represented as light grey scattered points. The mean value is plotted in blue, while mean plus/minus one standard deviation is plotted in black (and the region in between is shaded light blue). (A-F) Median diameter as a function of (A) active tension, (B) normalized hydraulic resistance of the open valve, (C) normalized hydraulic resistance of the closed valve, (D) threshold pressure to open the valve, (E) threshold diameter for positive or negative transmural pressure in the vessel, and (F) vessel wall stiffness.

Fig. S2: Supplementary analysis of parameter sensitivity in mean volume flow rate. All data points are represented as light grey scatter points. The mean value is plotted in orange, while mean plus/minus one standard deviation is plotted in black (and the region in between is shaded light orange). (A-F) Mean volume flow rate as a function of (A) active tension, (B) normalized hydraulic resistance of the open valve, (C) normalized hydraulic resistance of the closed valve, (D) threshold pressure to open the valve, (E) threshold diameter for positive or negative transmural pressure in the vessel, and (F) vessel wall stiffness.