Driving proteomic imbalance in malignancy provokes proteomic catastrophe

and confers tumor suppression

Babul Moni Ram[1], Omprakash Shriwas[1], Meng Xu[1], Kun-Han Chuang[1,2], and Chengkai Dai[1,*]

[1]Mouse Cancer Genetics Program, Center for Cancer Research, National Cancer Institute,

Frederick, MD 21702, USA

[2]Present address: School of Basic Medical Sciences, Fujian Medical University, Fuzhou, Fujian

350108, China

*Correspondence: Chengkai.dai@nih.gov

ABSTRACT

Unlike genomic instability, the role of proteomic instability in cancer remains poorly defined.

Heat shock factor 1 (HSF1), a master regulator of the proteotoxic stress response, preserves

proteome integrity under stress and is increasingly recognized as an oncogenic enabler. In

Neurofibromatosis type I (*NF1*)-deficient malignant peripheral nerve sheath tumor (MPNST)

cells, *HSF1* loss induces widespread protein polyubiquitination, aggregation, and tumor-

suppressive amyloidogenesis, yet is dispensable in non-transformed Schwann cells.

Mechanistically, HSF1 protects the mitochondrial chaperone HSP60 from toxic soluble amyloid

oligomers. To adapt to compromised protein quality, *HSF1*-deficient MPNST cells activate JNK

to repress mTORC1-dependent translation, reducing protein load. Stimulating mTORC1 in these

HSF1-deficient cells drives catastrophic proteomic imbalance, triggering pronounced cell death,

in part through unchecked amyloidogenesis, and suppressing tumor growth *in vivo*. Thus, HSF1

safeguards the cancer proteome, enabling the oncogenic capacity of mTORC1. This proof-of-

principle study establishes the induction of proteomic catastrophe as a new paradigm for

combating malignancy.

INTRODUCTION

Compared to genomic instability, proteomic instability in cancer is underappreciated. Malignant transformation triggers proteotoxic stress and destabilizes the proteome.[1, 2] Multiple mechanisms, including reactive oxygen species, genetic mutations, and dysregulated protein translation, contribute to increased protein damage, misfolding, and aggregation.[1-3] Notably, amyloidogenesis, a hallmark of neurodegenerative disorders,[4] also occurs in cancer.[5] Unlike neurons, however, cancer cells can contain amyloid formation and mitigate its cytotoxic effects. A key component of this defense is heat shock factor 1 (HSF1), which drives the heat shock response/proteotoxic stress response (HSR/PSR), a potent cytoprotective transcriptional program.[6-8] Consistent with its role in repressing proteomic instability, HSF1 has emerged as a powerful oncogenic enabler independent of driver mutations.[9-15]

In contrast to its dispensability in normal and non-transformed cells, HSF1 is essential in many cancer cell lines,[9,13,16-18] due to elevated intrinsic proteotoxic stress. This dependency, termed "non-oncogene addiction" to HSF1,[2,19] highlights its potential as a therapeutic target.

Neurofibromatosis type 1 (NF1) is a cancer predisposition syndrome affecting approximately 1 in 3000 individuals worldwide.[20] About 30-50% of NF1 patients develop benign plexiform neurofibromas, which can progress into highly aggressive malignant peripheral nerve sheath tumors (MPNSTs) with poor prognosis.[21] The *NF1* gene encodes neurofibromin, a 250-280 kDa RAS GTPase activating protein (GAP) that negatively regulates RAS signaling.[22] Loss of *NF1* in MPNST cells leads to hyperactive RAS/MAPK signaling, which not only promotes cellular growth but also drives constitutive HSF1 activation.[13] Accordingly, HSF1 overexpression and activation

are observed in both experimental MPNST models and primary NF1 tumors.[13] Significantly, *Hsf1*

deficiency impaired NF1-associated tumorigenesis in mice, partly through attenuation of

oncogenic RAS signaling.[13]

Utilizing the NF1 tumor model system, we herein identify HSF1 as a central regulator that

counteracts proteomic instability in cancer. We demonstrate that amyloidogenesis occurs in *NF1-*

deficient MPNST cells, driven in part by hyperactive mTORC1 signaling and increased protein

translation. Inhibition of HSF1 destabilizes the cancer proteome, leading to widespread protein

polyubiquitination, aggregation, and amyloid formation. This proteomic instability, ultimately,

causes cytotoxicity. To survive HSF1 inhibition, MPNST cells activate JNK to suppress

mTORC1 and reduce protein synthesis. Disruption of this adaptative response via JNK inhibition

exacerbates amyloidogenesis and cell death. Similarly, mTORC1 stimulation—via genetic *TSC2*

depletion or supplementation with leucine or its analog—synergizes with HSF1 inhibition to

induce profound proteomic chaos and widespread non-apoptotic cell death. Notably, blocking

amyloidogenesis significantly rescues cell viability. Together, these findings reveal a critical role

for HSF1 in antagonizing proteomic instability in cancer cells, representing a general pro-

oncogenic mechanism. This work provides proof of principle that inducing proteomic

catastrophe may serve as a next-generation anti-cancer therapeutic strategy.

RESULTS

HSF1 counters proteomic instability in *NF1*-deficient MPNST cells

We previously demonstrated that *NF1* loss leads to constitutive HSF1 activation and that *Hsf1* deficiency impairs tumorigenesis in *Nf1*[+/-]; *p53*[+/-] (NPcis) mice.[13] Consistent with these findings, analysis of human cancers revealed that *NF1* deletion is associated with elevated *HSF1* gene expression (Figure 1A), and *NF1* and *HSF1* expression levels are inversely correlated (Figure S1A). Importantly, among patients with *NF1*-deleted tumors, higher *HSF1* expression correlates with reduced survival (Figure 1B). Together, these data support a pro-oncogenic role for HSF1 in *NF1*-deficient malignancies.

Given the critical role of HSF1 in protein homeostasis, we examined how its depletion impacts the proteome of *NF1*-deficient MPNST cells compared to *NF1*-proficient, non-transformed human Schwann cells (HSCs), both primary and immortalized. HSCs are the presumed cell of origin for MPNSTs.[20,22] In three different MPNST cell lines (90-8TL, S462, and SNF96.2), transient *HSF1* depletion, using two previously validated lentiviral shRNAs,[9,13] led to a marked increase in global protein Lys48-specific polyubiquitination, a molecular mark for proteasomal degradation,[23] in both detergent-soluble and -insoluble fractions (Figures 1C, 1D, S1B, and S1C), indicative of widespread protein misfolding and aggregation. In contrast, no such increase was observed in HSCs (Figures 1C and 1D). These results indicate that HSF1 is specifically required to maintain proteomic stability in malignant Schwann cells.

While most cellular protein aggregates are amorphous, a subset adopts highly ordered β-sheet-rich structures known as amyloids, which are classically associated with neurodegenerative disorders,

especially Alzheimer's disease.[24] To determine whether *HSF1* depletion induces amyloidogenesis,

we stained cells with Congo red (CR), a widely applied amyloid-binding fluorescent dye.[25]

MPNST cells exhibited stronger CR staining than HSCs at baseline (Figure 1E), suggesting higher

amyloid burden in malignant cells. Notably, *HSF1* depletion further increased CR staining in

MPNST cells but not in HSCs (Figures 1E, 1F, S1D, and S1E). Similar results were obtained using

Thioflavin T staining (Figure S1F), another popular amyloid dye.[25] We further validated amyloid

accumulation using ELISA with two conformation-specific antibodies (A11 and OC), which

recognize soluble amyloid oligomers (AOs) and insoluble amyloid fibrils (AFs), respectively.[26,27]

HSF1 depletion significantly increased both $A11^+$ AOs and OC^+ AFs across all three MPNST cell

lines (Figures 1G, 1H, S1G-S1J), whereas no significant changes were detected in HSCs (Figures

1I and 1J). Collectively, these findings indicate that HSF1 is essential for maintaining proteome

integrity in malignant Schwann cells, particularly by suppressing amyloidogenesis, while being

largely dispensable in non-transformed Schwann cells.

Unchecked proteomic instability is cytotoxic

It has become evident that HSF1 acts as a generic oncogenic enabler; nevertheless, a universal

underlying mechanism has yet to be fully defined.

A healthy proteome enables the translation of genotypes into phenotypes. Thus, we reasoned that

preservation of proteomic stability represents a fundamental pro-oncogenic mechanism of HSF1.

To test this, we first determined whether proteomic instability—particularly amyloidogenesis—is

causally related to the cytotoxicity induced by *HSF1* depletion in cancer cells. To inhibit amyloid

formation, we employed CR, which exerts anti-amyloid effects through its amyloid-binding

activity.[5,28,29] Consistent with the concept of "non-oncogene addiction" in cancer, *HSF1*

depletion markedly impaired the viability of MPNST cells. Significantly, CR treatment partially

rescued this impaired viability (Figures 2A, S2A, and S2B), indicating that amyloidogenesis

contributes to the cytotoxic effects of HSF1 inhibition. In contrast, neither *HSF1* depletion nor

CR treatment affected the viability of HSCs (Figure 2B), congruent with their lack of proteomic

instability and amyloidogenesis. Together, these findings demonstrate that unchecked proteomic

instability—particularly amyloidogenesis—is cytotoxic. Accordingly, HSF1 enables

oncogenesis, at least in part, by constraining proteomic instability.

HSF1 neutralizes soluble amyloid oligomers to protect MPNST cells

How does HSF1 counter amyloid toxicity to enable the growth and survival of MPNST cells? In

previous research using tissue overgrowth models, we discovered that HSF1, unexpectedly,

physically neutralizes soluble AOs, thereby preventing their assault on HSP60.[30] As an essential

mitochondrial chaperone, HSP60 safeguards the mitochondrial proteome.[30-32]

We next asked whether this mechanism operates in cancer cells. Co-IP experiments indicated

that A11$^+$ AOs physically interacted with HSF1, but not major HSPs, in MPNST cells (Figure

2C), consistent with our previous findings.[30] This AO–HSF1 interaction was further validated

using an independent technique, *in situ* Proximity Ligation Assay (PLA) (Figures 2D-2F).

Significantly, *HSF1* depletion promoted the AO–HSP60 interaction (Figure 2G and 2H). Given

that loss of *HSP60* destabilized the mitochondrial proteome,[30,32] this AO–HSP60 interaction

signifies the induction of mitochondrial damage and subsequent cytotoxicity. Collectively, our

findings reveal that HSF1 protects MPNST cells, at least in part, by neutralizing soluble AOs,

thereby preserving the mitochondrial chaperone HSP60 function (Figure 2I).

MPNST cells react to HSF1 deficiency by activating JNK to attenuate translation

Given their dependency on HSF1, cancer cells are expected to engage adaptive responses to

survive its inhibition. Previously, we uncovered that in *Hsf1*-deficient mice, the stress-responsive

JNK kinase becomes constitutively activated, leading to disassembly of mTORC1 through

phosphorylation of its core components, RAPTOR and mTOR.[33] This inhibition of mTORC1

diminishes protein translation, leading to reduced cell size and whole-body lean mass in *Hsf1*-

deficient mice.[33]

We therefore hypothesized that cancer cells might exploit this same molecular regulation for

survival and adaptation. In MPNST cells, *HSF1* knockdown activated JNK, as indicated by

increased phosphorylation at T183/Y185. This activation was accompanied by suppression of

mTORC1 signaling, evidenced by diminished p70S6K T389 and 4EBP1 S65/T70

phosphorylation (Figures 3A, 3B, and S3A-S3D).[34,35] In contrast, *HSF*1 knockdown had minimal

effects on JNK and mTORC1 signaling in non-transformed HSCs (Figures 3A and 3B). Detected

by PLA, in MPNST cells HSF1 interacted with JNK. Upon *HSF1* depletion, JNK–RAPTOR

interactions increased, while mTOR–RAPTOR interactions decreased (Figures 3C-3E). Because

the mTOR-RAPTOR interaction is essential for mTORC1 assembly and activity,[36] these findings

indicate that JNK senses HSF1 deficiency to repress mTORC1. Notably, this sensing mechanism

appears more prominent in malignant cells than in non-transformed counterparts.

Consistent with mTORC1 inhibition, *HSF1* depletion reduced global protein translation rate in

MPNST cells (Figures 3F, S3E, and S3F), as measured by puromycin labeling.[33,37] This

regulatory relationship was conserved across diverse human cancer cell lines (Figure 3G).

Importantly, treatment with JNK-IN-8, an irreversible JNK inhibitor,[38] largely restored protein

translation following *HSF1* depletion and increased basal translation rates (Figures 3F, S3E, and

S3F). Consistent with these observations, in human cancers higher *HSF1* expression correlates

with lower JNK activation (Figure S3G), which in turn is associated with elevated mTORC1

activity, as indicated by increased 4EBP1 Ser65 phosphorylation (Figure S3H). In aggregate, our

findings demonstrate that JNK activation mediates mTORC1 inhibition and translational

suppression in response to *HSF1* deficiency. Thus, by restraining JNK signaling, HSF1 permits

robust protein translation fundamental to malignant growth (Figure 3H).

Lowering protein quantity is an adaptive response to alleviate proteomic instability

We next asked why cells attenuate protein translation in response to HSF1 deficiency. While

HSF1 maintains protein quality through HSPs, mTORC1 regulates protein quantity by

controlling translation. Because it is beneficial to lower protein quantity when protein quality is

compromised, we reasoned that this regulation represents a cellular adaptive response to the

imbalance between protein quantity and quality induced by HSF1 deficiency.

Proteomic imbalance disrupts proteome homeostasis and promotes proteomic instability, which

may ultimately trigger amyloidogenesis. To induce proteomic imbalance, we first stimulated

protein translation in MPNST cells by inhibiting JNK. As expected, treatment with JNK-IN-8

effectively reduced phosphorylation of c-JUN, a canonical JNK substrate,[34] in MPNST cells

(Figure 4A). Consistent with enhanced protein translation, JNK-IN-8 further activated mTORC1

signaling in these cells (Figures 4A and S4A). In contrast, only minimal effects were observed in

HSCs (Figure 4B), in line with their low basal JNK activity, as evidenced by unchanged c-JUN

phosphorylation. Intriguingly, the heightened protein translation following JNK inhibition was

accompanied by elevated amyloid accumulation in MPNST cells (Figures 4C and S4B), whereas

no significant changes were observed in HSCs (Figure 4D).

To directly assess the role of protein translation in amyloidogenesis, we inhibited translation

using two pharmacological agents with distinct mechanisms of action: 4EGI-1, which disrupts

eIF4E–eIF4G interactions,[39] and LY2584702, an inhibitor of p70S6K.[40] Both compounds

reduced basal amyloid levels in MPNST cells (Figures 4E,4F, and S4C), establishing a causal

link between protein translation and amyloidogenesis. Furthermore, translation inhibition

abrogated the induction of amyloids by JNK-IN-8 in MPNST cells (Figures 4G, 4H, S4D, and

S4E), indicating that JNK suppresses amyloidogenesis by inhibiting translation. No significant

effects were observed in HSCs (Figures 4G and 4H).

Given this relationship, we reasoned that preventing translation attenuation would aggravate

proteomic imbalance and enhance cytotoxicity in malignant cells. Consistent with this notion,

JNK inhibition by JNK-IN-8 further elevated amyloid levels in *HSF1*-depleted MPNST cells,

accompanied by exacerbated apoptosis as evidenced by caspase 3 cleavage (Figures 4I, 4J, and

S4F).[41] Notably, CR treatment suppressed this apoptosis, as indicated by reduced PARP cleavage

(Figure S4G),[42] causally implicating amyloidogenesis in this cell death. In contrast, *HSF1*

depletion and JNK inhibition had minimal effects on non-transformed Schwann cells (Figures

4K and 4L), consistent with their balanced proteome and low propensity for amyloid formation.

Together, these findings demonstrate that proteomic imbalance in malignant cells drives

proteomic instability and amyloidogenesis, leading to cytotoxicity. In non-transformed cells,

balanced protein quantity and quality prevent such instability. Thus, attenuation of protein

translation represents an adaptive mechanism by which cancer cells mitigate proteotoxic stress

and survive proteomic imbalance (Figure 4M).

mTORC1 stimulation aggravates proteomic imbalance and cytotoxicity induced by HSF1

inhibition

Because JNK regulates multiple targets beyond mTORC1, we sought to stimulate mTORC1 and

protein translation specifically. To this end, we genetically inhibited the tumor suppressor

Tuberous Sclerosis Complex subunit 2 (*TSC2*), a key negative regulator of mTORC1,[43] in

MPNST cells. As expected, *TSC2* knockdown in MPNST cells increased phosphorylation of

both p70S6K and 4EBP1, indicative of mTORC1 activation, which was accompanied by

elevated amyloid levels (Figures S5A-S5F). Like genetic *HSF1* depletion, pharmacological

inhibition of HSF1 with DTHIB— a small-molecule inhibitor that binds the HSF1 DNA-binding

domain to block transcription[44] —elevated global protein polyubiquitination and amyloids in

MPNST cells (Figures 5A, 5B, and S5G), indicating proteomic instability. Notably, HSF1

inhibition induced considerably greater amyloid accumulation in *TSC2*-deficient cells than in

control cells (Figure 5B), suggesting that elevated protein translation exacerbates proteotoxic

stress. *TSC2*-deficient cells also exhibited heightened JNK activation and mTORC1 inhibition

following DTHIB treatment, reflecting an amplified cellular response to severe proteotoxic

stress. Correspondingly, whereas control MPNST cells showed only modest cytotoxicity upon

HSF1 inhibition, measured by plasma membrane permeabilization, *TSC2*-deficient cells were

markedly more sensitive (Figures 5C, 5D, and S5H-S5K). Importantly, either inhibition of

protein translation or blockade of amyloid formation with CR largely rescued this cytotoxicity

(Figures 5C and 5D), causally implicating proteomic instability and amyloidogenesis in this cell

death. In support of a role for HSF1 in enabling *TSC2* deficiency, in tumors harboring *TSC2*

deletion higher *HSF1* gene expression correlates with poorer patient survival (Figure S5L).

To exclude potential mTORC1-independent effects of *TSC2* deficiency, we stimulated mTORC1

using NV-5138, a leucine analog currently in clinical trials as an antidepressant.[45,46]Amino acids,

particularly leucine, are potent activators of mTORC1.[47] Of note, although NV-5138 activates

mTORC1, it cannot be incorporated into protein synthesis.[45,46] To maximize mTORC1 activation

ex vivo, cells were cultured in leucine-free medium. Similar to *TSC2* knockdown, NV-5138

stimulated mTORC1 and elevated amyloid levels in MPNST cells (Figures S5M and 5E).

However, it elicited pronounced cytotoxicity only when combined with HSF1 inhibition, which

was largely rescued by CR treatment (Figure 5F). In contrast, combined NV-5138 and DTHIB

treatment had minimal effects on non-transformed Schwann cells (Figure 5G), consistent with

their reduced responsiveness to mTORC1 stimulation (Figures S5N). We next reasoned that by

directly contributing to protein quantity, L-leucine might impact the cancer proteome more

profoundly. In both S462 and 90-8TL cells, L-leucine synergized with DTHIB to induce severe

cytotoxicity (Figures 5H and S5O), whereas no cytotoxicity was induced in HSCs (Figure 5I).

To further corroborate the HSF1-dependent effects of DTHIB, we tested an independent small-

molecule inhibitor, KRIBB11[48], which binds the HSF1 transactivation domain to block the

recruitment of pTEFb. KRIBB11, consistent with DTHIB, synergized with L-leucine to promote

amyloid formation and trigger severe cytotoxicity in MPNST cells; importantly, CR treatment

both diminished amyloid levels and blocked cell death (Figures 5J and 5K). Finally, genetic

depletion of *HSF1* using shRNAs similarly synergized with L-leucine stimulation to provoke

prominent cytotoxicity (Figure 5L), providing orthogonal validation.

Collectively, these findings demonstrate that mTORC1 stimulation synergizes with HSF1

inhibition to drive severe cytotoxicity in cancer cells. This effect is mediated, at least in part, by

unconstrained proteomic imbalance and amyloidogenesis.

Unconstrained proteomic imbalance incites pronounced cell death

HSF1 inhibition alone induced both apoptotic and non-apoptotic death in MPNST cells, as

indicated by the presence or absence of caspase 3 cleavage. Intriguingly, concurrent mTORC1

stimulation shifted cell death predominantly toward a non-apoptotic mode, characterized by

plasma membrane permeabilization without caspase 3 cleavage (Figures 6A and 6B).

Multiple forms of non-apoptotic cell death, including pyroptosis, necroptosis, ferroptosis, and

autophagic cell death, can lead to loss of membrane integrity. To define the mechanism, we first

tested the role of caspases. Corroborating a caspase-independent mechanism, the pan-caspase

inhibitor Q-VD-OPh failed to prevent membrane permeabilization in *TSC2*-deficient MPNST

cells treated with the HSF1 inhibitor DTHIB (Figures 6A and 6B).[49] In contrast, Q-VD-OPh

effectively blocked apoptosis in control MPNST cells (Figures 6A and S6A), confirming its

activity. Because Q-VD-OPh also inhibits caspase 1 and no caspase 1 cleavage was detected

(Figure S6B),[50] a key mediator of pyroptosis,[51] these results argue against pyroptotic cell death.

Similarly, HSF1 inhibition did not induce necroptosis in *TSC2*-deficient MPNST cells, as

indicated by the absence of MLKL Ser358 phosphorylation (Figure S6B), a hallmark of

necroptotic execution.[52] Congruently, the necroptosis inhibitor necrostatin-1 had no effects on

membrane permeabilization (Figures 6C and S6C).[52]

Intriguingly, HSF1 inhibition in *TSC2*-deficient MPNST cells decreased glutathione peroxidase

4 (GPX4) but increased transferrin receptor 1 (TFR1), accompanied by markedly elevated lipid

peroxidation (Figures 6D and S6D), features typically associated with ferroptosis.[53,54] However,

the ferroptosis inhibitor liproxstatin-1,[53,54] while effectively reversing TFR1 elevation induced by

the ferroptosis inducer erastin, failed to suppress either TFR1 upregulation or membrane

permeabilization triggered by DTHIB (Figures 6C, 6D, and S6C). These results are not due to

ineffective inhibition, as both necrostatin-1 and liproxstatin-1 robustly blocked necroptosis and

ferroptosis, respectively, in control HT-29 cells (Figure S6E). HSF1 inhibition also elicited

autophagy, as evidenced by diminished p62/SQSTM1 and increased LC3B-II levels (Figure

6E).[55] To assess whether autophagy contributes to cell death, we blocked autophagy using the

class III PI3K inhibitor wortmannin.[56] Wortmannin effectively blocked autophagic flux, as

shown by reversal of rapamycin-induced changes in p62/SQSTM1 and LC3B-II (Figure 6E),

confirming successful inhibition of autophagy. Nevertheless, it failed to prevent membrane

permeabilization in *TSC2*-deficient cells treated with HSF1 inhibitors (Figures 6F and S6F).

These findings indicate that ferroptosis- and autophagy-associated changes occur downstream of

severe proteotoxic stress and are not the primary causes of cell death. Instead, our data support a

model in which unrestrained proteomic instability and consequent amyloidogenesis trigger a

unique form of cell death that is independent of apoptosis, autophagy, and canonical

programmed necrosis pathways (Figure 6G). We term this process proteomic catastrophe-

induced death (PCID). While its precise molecular execution remains to be fully defined, PCID

represents a mechanistically distinct vulnerability arising from the collapse of proteome

homeostasis in cancer cells.

Unconstrained proteomic imbalance suppresses *in vivo* tumor growth

Due to the limited *in vivo* bioavailability of DTHIB in our mouse models, we employed

KRIBB11 for *in vivo* studies. In agreement with our *ex vivo* findings, combined L-leucine

supplementation and KRIBB11 administration significantly suppressed the growth of

xenografted human S462 MPNST tumors in immunocompromised mice (Figure 7A). Despite

this overall reduction in tumor burden, 2 out of 8 tumors did not respond to the combination

treatment (Figure S7A), resulting in notable variations and a right-skewed distribution. Notably,

KRIBB11 alone only marginally affected tumor growth; conversely, in the presence of leucine it

markedly slowed tumor exponential growth rates and delayed tumor volume doubling time

(Figures S7B). This combination treatment also trended toward improved survival (Figure S7C).

To evaluate the broader applicability of this therapeutic strategy, we established a syngeneic

mouse melanoma model using B16-F10 cells in C57BL/6J mice. Similar to the MPNST model,

combined KRIBB11 and L-leucine supplementation significantly impeded tumor growth in this

aggressive model (Figures 7B and S7D). As expected, L-leucine activated mTORC1 signaling,

as indicated by increased S6RP phosphorylation (Figure 7C). Successful inhibition of HSF1 by

KRIBB11 was confirmed by the reduced transcription of *Hspa1a/Hsp72* and *Hspb1/Hsp25*

within the tumors (Figure 7D). Significantly, the combination treatment markedly induced

amyloid accumulation within tumors (Figure 7E). In stark contrast, no amyloids were induced in

normal tissues, including the brain and spleen (Figures 7F and 7G). Moreover, these treatments

did not cause significant loss of body weight (Figure S7E). In aggregate, these *in vivo* findings

from two distinct mouse models demonstrate that enforced proteomic imbalance—achieved

through combined HSF1 inhibition and mTORC1 stimulation—induces proteomic catastrophe

that suppresses tumor growth while sparing normal tissues (Figure 7H).

DISCUSSIONS

Genomic instability is widely recognized as a fundamental driver of oncogenesis. In contrast, the

role of proteomic instability in cancer remains obscure.

Proteomic instability is tumor suppressive yet inevitable

Proteomic instability does occur in cancer, as evidenced by elevated protein polyubiquitination,

aggregation, and even amyloidogenesis. Multiple factors contribute to this instability, including

protein oxidation by reactive oxygen species, mutation-induced protein misfolding, and

increased protein dosage resulting from aneuploidy.[1,3] Our recent findings further identify

uncontrolled protein translation as a key contributor to amyloidogenesis. This is not entirely

surprising, as it is estimated that up to 30% of newly synthesized polypeptides are misfolded and

targeted for proteasomal degradation.[57] Aberrant protein synthesis is therefore likely to increase

the burden of misfolded proteins, promoting protein aggregation and amyloid formation—

particularly in cancer cells, which harobor numerous mutations affecting protein folding.

Heightened protein translation, paradoxically, is also essential for malignant growth, rendering

proteomic instability and amyloidogenesis an unavoidable consequence of oncogenesis.

At the cellular level, amyloids are toxic to both neurons and cancer cells. At the organismal

level, while amyloidogenesis contributes to neurodegeneration, it appears to exert tumor-

suppressive effects. Mechanistically, our findings demonstrate that soluble AOs can directly

target HSP60, a key guardian of the mitochondrial proteome. It is likely that additional targets

and mechanisms contribute to amyloid toxicity in cancer. Moreover, the composition of cancer-

associated amyloids remains largely undefined. Beyond well-characterized species such as

Alzheimer's disease-associated Aβ, identifying the proteins that form amyloids in cancer will be

critical for understanding their functional roles and therapeutic potential.

HSF1 contains proteomic instability to enable malignancy

The diverse pro-oncogenic functions of HSF1 have been attributed to a wide array of

mechanisms; nonetheless, a unifying principle has remained elusive.

As an inherent consequence of malignant transformation, proteomic instability must be

restrained to avert tumor suppression and sustain cancer cell fitness. Now, our findings identify

containment of proteomic instability—particularly amyloidogenesis—as a central, unifying pro-

oncogenic function of HSF1. Notably, HSF1 does not eliminate amyloid formation but instead

limits its extent and toxicity. This is consistent with the fact that uncontrolled protein translation,

a hallmark of cancer, continuously drives amyloidogenesis. Consequently, cancer cells become

dependent on HSF1 to buffer this intrinsic proteotoxic burden, providing a mechanistic basis for

"HSF1 addiction".

Mechanistically, HSF1 contains amyloidogenesis through both indirect and direct means.

Indirectly, HSF1 induces HSPs to ensures proper protein folding and maintain proteome

integrity, thereby limiting the generation of misfolded proteins and amyloids. This is supported

by the observation that HSF1 inhibition leads to increased protein ubiquitination, aggregation,

and amyloid accumulation. Directly, HSF1 neutralizes soluble AOs once they form. With HSF1

inhibition, AOs are free to target HSP60, compromising the mitochondrial proteome and

triggering cytotoxicity. Thus, in cancer cells HSF1 inhibition not only increases the burden of

amyloids but, critically, exposes their intrinsic toxicity. Together, these findings establish that HSF1 enables malignancy by buffering proteomic instability, positioning it as a central safeguard against proteotoxic collapse in cancer cells.

mTORC1 signaling becomes tumor suppressive in the absence of HSF1

Given the fundamental dependency of malignant growth on protein translation, mTORC1 signaling is frequently hyperactivated in human cancers and is traditionally viewed as oncogenic.[58,59] However, our findings reveal a striking context-dependent shift: in the absence of HSF1, mTORC1 signaling becomes tumor suppressive. This functional reversal arises from a profound imbalance between protein quantity and quality. Without HSF1 to maintain proteome integrity, continued mTORC1-driven protein synthesis exacerbates proteomic instability, leading to enhanced amyloidogenesis and cytotoxicity. The observation that cytotoxicity is alleviated by either translation inhibitors or blockade of amyloid formation with CR strongly supports this model.

Conceptually, HSF1 enables the oncogenic potential of mTORC1 signaling by buffering the proteotoxic stress associated with elevated protein synthesis. In its absence, mTORC1-driven translation becomes detrimental, converting a canonical growth-promoting pathway into a tumor-suppressive force. These findings further establish HSF1 as a critical oncogenic enabler that permits cancer cells to tolerate the proteomic burden of malignant growth.

Inducing proteomic catastrophe represents a next-generation therapeutic paradigm

Conventional chemo- and radiotherapies primarily target the cancer genome by inducing DNA

damage. An emerging paradigm in oncology is to instead target the cancer proteome. To date,

most efforts have focused on inhibiting individual HSP families, such as HSP70 and HSP90.[60,61]

However, this strategy faces intrinsic limitations: cells harbor both constitutive HSPs, which

maintain essential basal proteostasis, and inducible HSPs, which provide additional chaperoning

capacity that is dispensable for basal proteostasis but required under stress. The structural and

functional similarity between these HSPs complicates selective targeting, and redundancy across

multiple HSP families further limits the efficacy of targeting any single chaperone system.

Our findings, together with prior studies, suggest that malignant cells operate under chronic

proteotoxic stress, resulting in constitutive HSF1 activation. Consequently, cancer cells become

reliant on the inducible chaperoning capacity governed by HSF1 for growth and survival.

Targeting HSF1 therefore represents a strategy to selectively disable this stress-adaptive reserve

while preserving basal proteostasis in normal cells. These insights provide a strong rationale for

therapeutic targeting of HSF1, an approach that is now beginning to enter clinical investigation.[62]

In response to HSF1 inhibition, cancer cells attenuate protein synthesis as an adaptive

mechanism, highlighting the importance of maintaining a balance between protein quantity and

quality. This observation led us to explore whether deliberately driving proteomic imbalance

could be therapeutically exploited.

Although mTORC1 hyperactivation is widespread in human cancers and has long been targeted

with inhibitors,[59, 63] these agents have largely produced cytostatic effects and limited clinical

benefit.[63,64] Moreover, because protein synthesis is essential for normal cellular function,

systemic mTORC1 inhibition carries significant toxicity, including immunosuppression.[63,65]

In contrast to conventional approaches, we pursued a counterintuitive strategy: stimulating

mTORC1 to drive proteomic imbalance. When combined with HSF1 inhibition, mTORC1

stimulation engenders catastrophic proteomic instability in cancer, culminating in cell death and

tumor suppression (Figure 7G). This PCID offers a wider therapeutic window, as normal cells,

with intact proteostasis and low proteotoxic stress, are largely spared. Mechanistically, mTORC1

activation increases protein synthesis, which can be accommodated in normal cells but

overwhelms the already stressed proteome of cancer cells, particularly given their burden of

mutant and misfolded proteins. Although cancer cells attempt to adapt via JNK-mediated

suppression of mTORC1, further specific blockade of this adaptive response is expected to

amplify proteomic imbalance and enhance therapeutic efficacy.

Transcription factors, traditionally, have been considered "undruggable" targets for small

molecules. While prototype direct HSF1 inhibitors such as DTHIB and KRIBB11 provide

important proof-of-concept, their limited *in vivo* potency underscores the need for more

clinically viable compounds. Nonetheless, our study establishes a new therapeutic paradigm: the

deliberate induction of proteomic catastrophe in cancer cells. We demonstrate a feasible strategy

to achieve this by simultaneously disrupting proteome quality control and enhancing protein

synthesis, thereby overwhelming the proteostatic capacity of malignant cells. This approach may

be broadly applicable across tumor types and represents a promising next-generation strategy for

cancer therapy.

ACKNOWLEDGMENTS

We would like to thank the Optical Microscopy and Image Analysis lab (OMAL) for their assistance with the confocal microscopy studies and Holly Morris for her assistance in animal studies. Schematic illustrations are created with BioRender.com. This work was supported by the grant from Department of Defense Congressionally Directed Medical Research Programs to C.D. (NF150028) and by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research (1ZIABC011767). The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. government.

AUTHOR CONTRIBUTIONS: B.M.R., O. S., and K-H.C. designed and conducted the experiments. C.D. conceptualized and supervised this study and analyzed the results. B.M.R. and C.D. wrote the manuscript.

DECLARATION OF INTERESTS: The authors declare no competing interests.

EXPERIMENTAL MATERIALS AND METHODS

Cells, Chemicals, and reagents

S462 (female) and 90-8TL (female) cells were kindly provided by Dr. Karen Cichowski.

sNF96.2 cells (male) and immortalized human Schwann cells (hTERT ipn02.3 2λ, female) were

purchased from ATCC. Primary human Schwann cells (Cat#1700, ScienCell Research

Laboratories) were grown on poly-L-lysine coated T-75 flasks and maintained in complete

Schwann Cell Medium (Cat#1701, ScienCell Research Laboratories). B16-F10-luc2 (male)

cells were purchased from ATCC (cat#CRL-6475-LUC2). All cell lines, except 90-8TL and

HSCs (no known STR profiles available), were authenticated by ATCC. All cell lines, except

primary human Schwann cells, were maintained in DMEM supplemented with 10% HyClone

bovine growth serum and 1% penicillin–streptomycin. Cells were maintained in an incubator

with 5% CO2 at 37 °C.

The following chemicals were purchased from commercial sources: JNK-IN-8 (Cat#S4901,

Selleckchem), 6-FAM-dC-puromycin (cat#NU-925-6FM, Jena Bioscience), Congo red

(cat#B2431014, Thermo Scientific Chemicals), Thioflavin T (cat#211760050, Thermo Scientific

Chemicals), LY2584702 (cat#S7698, Selleckchem), 4EGI-1 (cat#S7369, Selleckchem), and

DTHIB (cat# HY-138280, MedChemExpress), KRIBB11 (cat#HY-100872, MedChemExpress),

KRIBB11 (cat#T3652, TargetMol Chemicals), Q-VD-OPh (cat#HY-12305, MedChemExpress),

necrostatin-1 (cat#HY-15760, MedChemExpress), Liproxstatin-1 (cat#HY-12726,

MedChemExpress), wortmannin (cat#HY-10197, MedChemExpress), rapamycin (cat#HY-

10219, MedChemExpress), erastin (cat#HY-15763, MedChemExpress), MHY1485 (cat#HY-

B0795, MedChemExpress), NV-5138 HCl (cat#HY-114384B, MedChemExpress), and Liperfluo

(cat#L248, Dojindo Laboratories), dimethylacetamide (cat#T19439, TargetMol Chemicals

Inc.), PEG300 (cat#T7022, TargetMol Chemicals Inc.), and Captisol (cat#RC-0C7-100, CyDex

Pharmaceuticals).

Rabbit phospho-4EBP1 S65/T70 antibody (cat#sc-12884-R), mouse monoclonal HSF1 (E-4,

cat#sc-17757) antibody, rabbit HSF1 antibody (H-311, cat#sc-9144), mouse monoclonal JNK

(D-2, cat#sc-7345), rabbit JNK1/3 (C-17, cat#sc-474) antibody were purchased from Santa

Cruz Biotechnology; antibodies for phosphor-4EBP1 Thr70 (cat#9455), phospho-4EBP1

Thr37/46 (236B4, cat#2855), phospho-p70S6K T389 (108D2, cat# 9234), phosphor-S6RP

Ser235/236 (D57.2.2E, cat#4858), p70 S6K (49D7, cat#2708), 4EBP1 (53H11, cat#9644),

S6RP (5G10, cat#2217), mTOR (7C10, cat#2983), phospho-JNK1/2 T183/Y185 (81E11,

cat#4668), JNK2 (56G8, cat#9258), c-Jun (60A8, cat#9165), phospho-c-Jun Ser63 II

(cat#9261), PARP (cat#9542), Tuberin/TSC2 (D93F12, cat#4308), SQSTM1/p62 (D1Q5S,

cat#39749), cleaved caspase 3 (Asp175) (5A1E, cat#9664), caspase 1(D7F10, cat#3866),

gasdermin D (E9S1X, cat#39754), phosphor-MLKL Ser358 (D6H3V, cat#91689), GPX4

(E5Y8K, cat#59735), and transferrin receptor/CD71 (H68.4, cat#46222) antibody were

purchased from Cell Signaling Technologies; mouse β-actin antibody-HRP (Cat#A00730)

conjugates were from GenScript. Anti-K48-specific ubiquitin antibody (Apu2, cat#05-1307)

and mouse monoclonal anti-RAPTOR antibody (1H6.2, cat#05-1470) were purchased from

EMD Millipore. Mouse monoclonal anti-JNK1 (C-terminal region) antibody (cat#JM2671)

was purchased from ECM Biosciences. Anti-amyloid oligomers (A11, cat#SPC-506D) and

anti-amyloid fibrils (OC, cat#SPC-507D) antibodies were purchased from StressMarq

Biosciences Inc. Mouse monoclonal anti-HSP60 antibody (LK-1, cat#ADI-SPA-806) was

purchased from Enzo Life Sciences Inc. β-Actin (cat#GTX629630), GAPDH

(cat#GTX627408), α-tubulin (GT114, cat#GTX628802), and LC3B antibody

(cat#GTX127375) was purchased from GeneTex Inc.

METHOD DETAILS

Lentiviral Particle Production and Transduction of lentiviral shRNAs

Lentiviral pLKO.1 shRNA plasmids targeting human *HSF1* were described previously[10]:

TRCN0000007480 (hA6) and TRCN0000007483 (hA9). The sequence of control hairpins

targeting a scrambled sequence with no known homology to any human genes is 5′-

CCTAAGGTTAAGTCGCCCTCG-3′. The pLKO.1 shRNA plasmid targeting human *TSC2* was

a gift from Do-Hyung Kim (Cat#15478, Addgene). Lentiviral pLKO shRNA plasmids were

transiently co-transfected individually along with plasmids encoding ΔVPR and VSV-G into

HEK293T cells using TurboFect™ (Cat#R0531, Thermo Scientific). Viral supernatants were

collected after centrifugation at 1,000 rpm for 15 min. Viral transduction was achieved by

incubating target cells with viral supernatants containing 10μg/ml polybrene (Cat#TR-1003-G,

Millipore-Sigma) for 24h. For stable knockdown of *TSC2,* lentivirus particles were prepared using

pLKO shRNA plasmids targeting human *TSC2*. Scrambled shRNA was used for stable control

cells. After 4 days of transduction, cells were gradually selected using 1-4 μg/ml puromycin for 2

weeks.

Soluble and Insoluble Protein Fractionation

Equal numbers of cells were incubated with cold cell-lysis buffer (100 mM NaCl, 30 mM Tris-

HCl pH 7.6, 1% Triton X-100, 20 mM sodium fluoride, 1mM EDTA, 1mM sodium

orthovanadate, and 1x Halt™ protease inhibitor cocktail) on ice for 20 min. The crude lysates

were first centrifuged at 500xg for 5 min at 4°C. The supernatants from low-speed (500xg)

centrifugation were transferred to a new Eppendorf tube. The first pellets were resuspended in

DNA digestion solution (2 units DNase I, 1% Triton X-100, 10 mM Tris-HCl, 2.5 mM

MgCl2, 0.5 mM CaCl2, pH 7.6) for 30 min at RT. An equal volume of 1xPBS with 2% SDS

was then added to resuspend the first pellets for additional 30 min at RT. Following the

centrifugation at 17,000xg for 10 min at RT, the supernatants were removed. The supernatants

from low-speed centrifuge were added back to the second pellets, followed by centrifugation

at 17,000xg for 15 min at 4°C. The final supernatants and pellets from high-speed (17,000xg)

centrifugation were collected as detergent-soluble and -insoluble fractions, respectively.

Insoluble fractions were further sonicated in 1xPBS with 2% SDS at high intensity using a

Bioruptor® Sonication Device (Diagenode) for SDS-PAGE.

ThT and CR Staining for Flow Cytometry

After washing with 1xPBS, cells were fixed with 4% formaldehyde at RT for 10 min.

Following fixation and 1xPBS washing, cells were resuspended in 1 ml permeabilization

buffer (1xPBS with 0.5% Triton X-100 and 3 mM EDTA) and incubated on ice for 30 min.

Following washing with 1xPBS, Cells were stained with 10 μM ThT or CR dissolved in

1xPBS for 5 min. Cells were washed and resuspended in PBS and fluorescence signals were

measured by a FACSCalibur, LSRFortessa X-20 or LSR II flow cytometer (BD Biosciences).

Amyloid Oligomer and Fibril Quantitation by Direct ELISA and Flow Cytometry

To quantitate soluble amyloid prefibrillar oligomers, soluble cell lysates diluted in 1xPBS

from equal numbers of cells were coated on each well in a 96-well ELISA plate at 4°C

overnight followed by blocking (5% non-fat milk in TBS-0.05% Tween® 20) at RT for 1 hr.

Each well was incubated with 100 μl amyloid oligomer antibodies (A11, 1:1,000 diluted in

the blocking buffer) at RT for 2 hr. After washing with TBS-T, goat anti-rabbit secondary Ab

HRP conjugates (1:1,000 diluted in blocking buffer) were added to each well and incubated at

RT for 1 hr. Following washing, 100 μl 1-Step Ultra TMB-ELISA substrates (Cat#34029,

Thermo Scientific) were added to each well. After adding 100μl Stop solution, absorbance was

measured at 450 nm using a SpectraMax iD5 microplate reader (Molecular Devices).

To quantitate amyloid fibrils, detergent-insoluble fractions from equal numbers of cells were

extracted as described in Soluble and Insoluble Protein Fractionation. The final pellets were

collected as detergent-insoluble fractions and solubilized by sonication for 15 min in 1xPBS

with 2% SDS. Solubilized proteins diluted in 1xPBS were added to each well and incubated at

37°C without cover overnight to air dry the wells. The following steps were identical to the

oligomer detection except for the use of amyloid fibril antibodies (OC) as the primary Ab.

For amyloid oligomer and fibril quantification by flow cytometry, cells were collected by

trypsinization and fixed in 4% Formaldehyde at RT for 10 min. Following permeabilization using

0.3% Triton-X-100, samples were incubated with primary antibodies A11-FITC (1:100 in 5%

BSA, StressMarq Biosciences, cat# SPC-506D-FITC) or OC-PerCP (1:100 in 5% BSA,

StressMarq Biosciences, cat# SPC-507D-PCP) at RT for 3 hrs. The cells were washed with PBS

and analyzed using flow cytometry.

Measurement of Cell Viability

Cell viability was determined using the CellTiter-Blue® Cell Viability Assay (cat#G8080, Promega). Cells were grown in 96 well plates (1×10^4 per well) for 24h. A 20µl CellTiter-Blue® reagent was added to 100µl of media in each well. The plates were incubated for 2h at 37°C and the fluorescence was recorded at 560/590 nm. The reagents were discarded, and the wells were washed with PBS before adding fresh media. Similarly, subsequent readings were recorded at 24h intervals till 96h time point.

Proximity Ligation Assay

Cells were fixed with 4% formaldehyde in PBS for 15 min at RT. After blocking with 5% normal goat serum in PBS with 0.3% Triton X-100, primary antibodies 1:100 diluted in blocking buffer were incubated with fixed cells overnight at 4°C. Following washing with PLA wash Buffer A, samples were incubated with 1:5 diluted Duolink® in Situ PLA Probe anti-rabbit Plus (cat#DUO92002, Sigma-Aldrich) and anti-mouse Minus (cat#DUO92004, Sigma-Aldrich) at 37°C for 1 hr. Subsequently, ligation (30 minutes, 37°C), rolling circle amplification, and detection (120 minutes, 37°C) were performed using the Duolink® In Situ Detection Reagents Red (cat#DUO92008, Sigma-Aldrich). Nuclei were counterstained with Hoechst 33342. PLA signals were documented by a Zeiss LSM780 confocal microscope (Carl Zeiss). For PLA flow cytometry, an equal number of cells were collected in microcentrifuge tubes by trypsinization and fixed with 4% formaldehyde in PBS for 10 min at RT. Subsequent PLA steps were followed as described earlier. The PLA signals were quantified by a FACSCalibur flow cytometer (BD Biosciences).

Mouse anti-JNK1 (JM2671) and rabbit anti-HSF1 (H-311) antibodies were used for JNK-HSF1

PLA. Rabbit anti-JNK1/3 (C-17) and mouse anti-RAPTOR (1H6.2) antibodies were used for

JNK-RAPTOR PLA. Rabbit anti-mTOR (7C10) and mouse anti-RAPTOR (1H6.2) antibodies

were used for mTOR-RAPTOR PLA. Mouse anti-HSF1 (E-4) and rabbit anti-AO (A11)

antibodies were used for HSF1-AO PLA. Rabbit anti-AO (A11) and mouse anti-HSP60 (LK1)

antibodies were used for AO-HSP60 PLA.

Measurement of global protein translation

Live cells were incubated with 100 nM 6-FAM-dC-puromycin in complete culture medium

for 30 min and analyzed by a FACSCalibur flow cytometer (BD Biosciences).

Immunoblotting and Immunoprecipitation

Whole-cell protein extracts were prepared in cold cell-lysis buffer (100 mM NaCl, 30 mM Tris-

HCl pH 7.6, 1% Triton X-100, 20 mM sodium fluoride, 1mM EDTA, 1mM sodium

orthovanadate, and 1x Halt™ protease inhibitor cocktail). Proteins were transferred to

nitrocellulose membranes. Following incubation with the blocking buffer (5% non-fat milk in 1x

TBS-T) for 1 hour at RT, membranes were incubated with primary antibodies (1:1,000 dilution in

the blocking buffer) overnight at 4 °C. After washing with 1xTBS-T for 3 times, membranes

were incubated with peroxidase-conjugated secondary antibodies (1: 3,000 diluted in the

blocking buffer) at RT for 1 hr. Signals were detected using SuperSignal™ West

chemiluminescent substrates. For normal Co-IP, proteins were extracted using a QSonica Q125

sonicator (total process time: 15S, pulse-on time: 5S, pulse-off time: 10S, output intensity: 30%)

in 1x sonication buffer (20 mM Tris, 20 mM NaCl, 1 mM EDTA, 20 mM β-glycerol-phosphate,

20 mM sodium fluoride, 4 mM sodium orthovanadate, and 1 mM DTT pH7.4, supplemented

with Halt protease inhibitor cocktail). Total 1 mg whole cell lysates were incubated with A11

antibodies at 4 °C overnight. Normal rabbit IgG were used as the negative controls. The primary

antibody was precipitated using Protein G magnetic beads. After washing with the lysis buffer

for 3 times, beads were boiled in 1x loading buffer for 5 min before loading on SDS-PAGE.

EasyBlocker (Cat#GTX425858, GeneTex) was used for blocking and anti-HSF1 (H-311)

primary antibody (cat#sc-9144, Santa Cruz Biotechnology) incubation (1:1,000, O/N at 4°C).

The blot was incubated with EasyBlot anti-Rabbit IgG HRP conjugated antibody

(cat#GTX221666-01, GeneTex) for 1 hour. Chemiluminescent signals were captured by an

iBright™ FL1000 imaging system (ThermoFisher Scientific).

Live-Or-Dye™ Dead Cell Staining

Cells were cultured in DMEM for 48 hours before being treated with inhibitors for the required

periods and collected by trypsinization. 1×10^6 cells/ml cells were stained with Live-or-Dye™

fixable dead cell dyes (1:1,000 dilution, cat#32003, Biotium) at RT for 30 min, protected from

light. Stained cells were washed once with PBS and analyzed by flow cytometry. For co-staining

with cleaved caspase 3 antibody, stained cells were fixed with 4% formaldehyde (10 min, RT)

and permeabilized with 0.3% Triton-X-100 in PBS (10 min, RT). The cells were stained by

Cleaved Caspase 3 antibody (Asp175) (5A1E) (1:1,000, cat#9664, Cell Signaling Technology)

followed by goat anti-Rabbit IgG Alexa Fluor™ 488 conjugated (1:1,000, cat#A-11034,

ThermoFisher Scientific) secondary antibodies. The samples were analyzed by flow cytometry.

Inhibitor treatments

Treatments with DTHIB, Q-VD-OPh, Liproxstatin-1, Necrostatin-1, and Wortmannin were done for 72 hours. For Congo Red and LY-2584702 treatments, cells were pretreated for 24h followed by DTHIB for an additional 72h. For treatments with NV-5138, cells were pretreated for 48 hours in a leucine-free DMEM (cat#226-024, Crystalgen) before adding DTHIB for an additional 72 hours.

Animal studies

All animal experiments described here were approved by the institutional Animal Care and Use Committee (ACUC) of the National Cancer Institute, Frederick. Experiments on all animals were conducted following the recommendations of the Guide for the Care and Use of Laboratory Animals (National Academies Press, 2011). Animals were housed and maintained on a 12-hour light/12-hour dark cycle at temperatures between 20°C and 27°C and humidity levels of 30% to 70%, with food and water *available ad libitum*.

To establish the MPNST xenograft model, 9×10^6 S462 MPNST cells (female) suspended with 30 % Matrigel were subcutaneously implanted the right flanks of 8-week-old female NIH-III nude mice (Charles River Laboratories). After 7 days, xenografted mice were assigned to 4 groups, eight each group, by stratified randomization based on tumor sizes (between 72 and 199 mm³). Mice were treated with KRIBB11 (65mg/kg, i.p. injection daily, cat#HY-100872, Lot#317606, Medchemexpress LLC) with and without 2% L-leucine dissolved in drinking water stored in amber-colored bottles. KRIBB11 was freshly dissolved daily in the vehicle (final concentration:10% dimethylacetamide, 50% PEG300, 20% Captisol, and 20% sterile saline).

To establish the syngeneic melanoma model, 2×10^6 B16-F10-luc2 cells (male) mixed with

Matrigel (1:1 ratio) in a 200μL volume were subcutaneously implanted into the right flanks of

8-week-old C57BL/6J male mice (The Jackson Laboratory). After 5 days, xenografted mice were

assigned to 4 groups, eight each group, by stratified randomization based on tumor sizes (between

75 and 284 mm³). Mice were treated with KRIBB11 (65 mg/kg, i.p. injection daily, cat#T3652,

Lot#132988, TargetMol Chemicals Inc.) with and without 2% L-leucine dissolved in drinking

water stored in amber-colored bottles. KRIBB11 was freshly dissolved daily in the vehicle (final

concentration:10% dimethylacetamide, 50% PEG300, 16% Captisol, and 24% sterile saline).

To formulate the KRIBB11 solution, KRIBB11 was first dissolved in dimethylacetamide,

followed by addition of PEG300. Lastly, 40 or 50% Capitol dissolved in sterile saline was added.

Tumor sizes were measured using a caliper by an animal technician blind to the study. To

assess the compound's toxicity, the body weight of tumor-bearing animals was also recorded.

Tumor volumes were estimated using the formula [length (mm) $\times$ width^2 (mm^2)]/2.

Quantification And Statistical Analysis

Statistical analyses were performed using Prism GraphPad 10.0 (GraphPad Software). The

detailed statistical methods and sample sizes are provided in the figure legends. All results are

expressed as mean±SD, mean±SEM, or median and IQR. The statistical significance is defined

as: *$p<0.05$, **$p<0.01$; ***$p<0.001$; ****$p<0.0001$; n.s.: not significant. For *in vitro*

experiments, sample size required was not determined a priori. The experiments were not

randomized. For *in vivo* experiments, sample size required was determined by pilot experiments.

REFERENCES

1.   Dai C, Dai S, Cao J. Proteotoxic stress of cancer: implication of the heat-shock response in

oncogenesis. J Cell Physiol. 2012;227(8):2982-7. doi: 10.1002/jcp.24017. PubMed PMID:

22105155; PMCID: PMC3431153.

2.   Dai C. The heat-shock, or HSF1-mediated proteotoxic stress, response in cancer: from

proteomic stability to oncogenesis. Philos Trans R Soc Lond B Biol Sci. 2018;373(1738). doi:

10.1098/rstb.2016.0525. PubMed PMID: 29203710; PMCID: PMC5717525.

3.   Guang MHZ, Kavanagh EL, Dunne LP, Dowling P, Zhang L, Lindsay S, Bazou D, Goh CY,

Hanley C, Bianchi G, Anderson KC, O'Gorman P, McCann A. Targeting Proteotoxic Stress in

Cancer: A Review of the Role that Protein Quality Control Pathways Play in Oncogenesis.

Cancers (Basel). 2019;11(1). Epub 20190109. doi: 10.3390/cancers11010066. PubMed

PMID: 30634515; PMCID: PMC6356294.

4.   Wilson DM, 3rd, Cookson MR, Van Den Bosch L, Zetterberg H, Holtzman DM, Dewachter I.

Hallmarks of neurodegenerative diseases. Cell. 2023;186(4):693-714. doi:

10.1016/j.cell.2022.12.032. PubMed PMID: 36803602.

5.   Tang Z, Dai S, He Y, Doty RA, Shultz LD, Sampson SB, Dai C. MEK guards proteome

stability and inhibits tumor-suppressive amyloidogenesis via HSF1. Cell. 2015;160(4):729-44.

doi: 10.1016/j.cell.2015.01.028. PubMed PMID: 25679764; PMCID: PMC4564124.

6.   Li J, Labbadia J, Morimoto RI. Rethinking HSF1 in Stress, Development, and Organismal

Health. Trends Cell Biol. 2017;27(12):895-905. Epub 20170907. doi:

10.1016/j.tcb.2017.08.002. PubMed PMID: 28890254; PMCID: PMC5696061.

7.   Anckar J, Sistonen L. Regulation of HSF1 function in the heat stress response: implications in

aging and disease. Annu Rev Biochem. 2011;80:1089-115. doi: 10.1146/annurev-biochem-

060809-095203. PubMed PMID: 21417720.

8. Gomez-Pastor R, Burchfiel ET, Thiele DJ. Regulation of heat shock transcription factors and

their roles in physiology and disease. Nat Rev Mol Cell Biol. 2018;19(1):4-19. Epub

20170830. doi: 10.1038/nrm.2017.73. PubMed PMID: 28852220; PMCID: PMC5794010.

9. Min JN, Huang L, Zimonjic DB, Moskophidis D, Mivechi NF. Selective suppression of

lymphomas by functional loss of Hsf1 in a p53-deficient mouse model for spontaneous

tumors. Oncogene. 2007;26(35):5086-97. Epub 20070219. doi: 10.1038/sj.onc.1210317.

PubMed PMID: 17310987.

10. Dai C, Whitesell L, Rogers AB, Lindquist S. Heat shock factor 1 is a powerful multifaceted

modifier of carcinogenesis. Cell. 2007;130(6):1005-18. doi: 10.1016/j.cell.2007.07.020.

PubMed PMID: 17889646; PMCID: PMC2586609.

11. Meng L, Gabai VL, Sherman MY. Heat-shock transcription factor HSF1 has a critical role in

human epidermal growth factor receptor-2-induced cellular transformation and

tumorigenesis. Oncogene. 2010;29(37):5204-13. Epub 20100712. doi:

10.1038/onc.2010.277. PubMed PMID: 20622894; PMCID: PMC2940982.

12. Jin X, Moskophidis D, Mivechi NF. Heat shock transcription factor 1 is a key determinant of

HCC development by regulating hepatic steatosis and metabolic syndrome. Cell Metab.

2011;14(1):91-103. doi: 10.1016/j.cmet.2011.03.025. PubMed PMID: 21723507; PMCID:

PMC3214631.

13. Dai C, Santagata S, Tang Z, Shi J, Cao J, Kwon H, Bronson RT, Whitesell L, Lindquist S.

Loss of tumor suppressor NF1 activates HSF1 to promote carcinogenesis. J Clin Invest.

2012;122(10):3742-54. Epub 20120904. doi: 10.1172/JCI62727. PubMed PMID: 22945628;

PMCID: PMC3461912.

14. Kourtis N, Lazaris C, Hockemeyer K, Balandran JC, Jimenez AR, Mullenders J, Gong Y,

Trimarchi T, Bhatt K, Hu H, Shrestha L, Ambesi-Impiombato A, Kelliher M, Paietta E,

Chiosis G, Guzman ML, Ferrando AA, Tsirigos A, Aifantis I. Oncogenic hijacking of the

stress response machinery in T cell acute lymphoblastic leukemia. Nat Med.

2018;24(8):1157-66. Epub 20180723. doi: 10.1038/s41591-018-0105-8. PubMed PMID:

30038221; PMCID: PMC6082694.

15. Fujimoto M, Takii R, Matsumoto M, Okada M, Nakayama KI, Nakato R, Fujiki K, Shirahige

K, Nakai A. HSF1 phosphorylation establishes an active chromatin state via the TRRAP-

TIP60 complex and promotes tumorigenesis. Nat Commun. 2022;13(1):4355. Epub

20220729. doi: 10.1038/s41467-022-32034-4. PubMed PMID: 35906200; PMCID:

PMC9338313.

16. Heimberger T, Andrulis M, Riedel S, Stuhmer T, Schraud H, Beilhack A, Bumm T, Bogen

B, Einsele H, Bargou RC, Chatterjee M. The heat shock transcription factor 1 as a potential

new therapeutic target in multiple myeloma. Br J Haematol. 2013;160(4):465-76. Epub

20121218. doi: 10.1111/bjh.12164. PubMed PMID: 23252346.

17. Chuma M, Sakamoto N, Nakai A, Hige S, Nakanishi M, Natsuizaka M, Suda G, Sho T,

Hatanaka K, Matsuno Y, Yokoo H, Kamiyama T, Taketomi A, Fujii G, Tashiro K, Hikiba Y,

Fujimoto M, Asaka M, Maeda S. Heat shock factor 1 accelerates hepatocellular carcinoma

development by activating nuclear factor-kappaB/mitogen-activated protein kinase.

Carcinogenesis. 2014;35(2):272-81. Epub 20131015. doi: 10.1093/carcin/bgt343. PubMed

PMID: 24130164.

18. Fujimoto M, Takaki E, Takii R, Tan K, Prakasam R, Hayashida N, Iemura S, Natsume T,

Nakai A. RPA assists HSF1 access to nucleosomal DNA by recruiting histone chaperone

FACT. Mol Cell. 2012;48(2):182-94. Epub 20120830. doi: 10.1016/j.molcel.2012.07.026.

PubMed PMID: 22940245.

19. Solimini NL, Luo J, Elledge SJ. Non-oncogene addiction and the stress phenotype of cancer

cells. Cell. 2007;130(6):986-8. doi: 10.1016/j.cell.2007.09.007. PubMed PMID: 17889643.

20. Gutmann DH, Ferner RE, Listernick RH, Korf BR, Wolters PL, Johnson KJ.

Neurofibromatosis type 1. Nat Rev Dis Primers. 2017;3:17004. Epub 20170223. doi:

10.1038/nrdp.2017.4. PubMed PMID: 28230061.

21. Hirbe AC, Gutmann DH. Neurofibromatosis type 1: a multidisciplinary approach to care.

Lancet Neurol. 2014;13(8):834-43. doi: 10.1016/S1474-4422(14)70063-8. PubMed PMID:

25030515.

22. Cichowski K, Jacks T. NF1 tumor suppressor gene function: narrowing the GAP. Cell.

2001;104(4):593-604. doi: 10.1016/s0092-8674(01)00245-8. PubMed PMID: 11239415.

23. Grice GL, Nathan JA. The recognition of ubiquitinated proteins by the proteasome. Cell Mol

Life Sci. 2016;73(18):3497-506. Epub 20160502. doi: 10.1007/s00018-016-2255-5. PubMed

PMID: 27137187; PMCID: PMC4980412.

24. Knowles TP, Vendruscolo M, Dobson CM. The amyloid state and its association with protein

misfolding diseases. Nat Rev Mol Cell Biol. 2014;15(6):384-96. doi: 10.1038/nrm3810.

PubMed PMID: 24854788.

25. Yakupova EI, Bobyleva LG, Vikhlyantsev IM, Bobylev AG. Congo Red and amyloids:

history and relationship. Biosci Rep. 2019;39(1). Epub 20190115. doi:

10.1042/BSR20181415. PubMed PMID: 30567726; PMCID: PMC6331669.

26. Kayed R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, Glabe CG.

Common structure of soluble amyloid oligomers implies common mechanism of

pathogenesis. Science. 2003;300(5618):486-9. doi: 10.1126/science.1079469. PubMed

PMID: 12702875.

27. Kayed R, Head E, Sarsoza F, Saing T, Cotman CW, Necula M, Margol L, Wu J, Breydo L,

Thompson JL, Rasool S, Gurlo T, Butler P, Glabe CG. Fibril specific, conformation

dependent antibodies recognize a generic epitope common to amyloid fibrils and fibrillar

oligomers that is absent in prefibrillar oligomers. Mol Neurodegener. 2007;2:18. Epub

20070926. doi: 10.1186/1750-1326-2-18. PubMed PMID: 17897471; PMCID:

PMC2100048.

28. Heiser V, Scherzinger E, Boeddrich A, Nordhoff E, Lurz R, Schugardt N, Lehrach H,

Wanker EE. Inhibition of huntingtin fibrillogenesis by specific antibodies and small

molecules: implications for Huntington's disease therapy. Proc Natl Acad Sci U S A.

2000;97(12):6739-44. doi: 10.1073/pnas.110138997. PubMed PMID: 10829068; PMCID:

PMC18723.

29. Sanchez I, Mahlke C, Yuan J. Pivotal role of oligomerization in expanded polyglutamine

neurodegenerative disorders. Nature. 2003;421(6921):373-9. doi: 10.1038/nature01301.

PubMed PMID: 12540902.

30. Tang Z, Su KH, Xu M, Dai C. HSF1 physically neutralizes amyloid oligomers to empower

overgrowth and bestow neuroprotection. Sci Adv. 2020;6(46). Epub 20201111. doi:

10.1126/sciadv.abc6871. PubMed PMID: 33177089; PMCID: PMC7673739.

31. Christensen JH, Nielsen MN, Hansen J, Fuchtbauer A, Fuchtbauer EM, West M, Corydon

TJ, Gregersen N, Bross P. Inactivation of the hereditary spastic paraplegia-associated Hspd1

gene encoding the Hsp60 chaperone results in early embryonic lethality in mice. Cell Stress

Chaperones. 2010;15(6):851-63. doi: 10.1007/s12192-010-0194-x. PubMed PMID:

20393889; PMCID: PMC3024079.

32. Fan F, Duan Y, Yang F, Trexler C, Wang H, Huang L, Li Y, Tang H, Wang G, Fang X, Liu

J, Jia N, Chen J, Ouyang K. Deletion of heat shock protein 60 in adult mouse cardiomyocytes

perturbs mitochondrial protein homeostasis and causes heart failure. Cell Death Differ.

2020;27(2):587-600. Epub 20190617. doi: 10.1038/s41418-019-0374-x. PubMed PMID:

31209364; PMCID: PMC7205885.

33. Su KH, Cao J, Tang Z, Dai S, He Y, Sampson SB, Benjamin IJ, Dai C. HSF1 critically

attunes proteotoxic stress sensing by mTORC1 to combat stress and promote growth. Nat

Cell Biol. 2016;18(5):527-39. Epub 20160404. doi: 10.1038/ncb3335. PubMed PMID:

27043084; PMCID: PMC5341796.

34. Johnson GL, Nakamura K. The c-jun kinase/stress-activated pathway: regulation, function

and role in human disease. Biochim Biophys Acta. 2007;1773(8):1341-8. Epub 20070104.

doi: 10.1016/j.bbamcr.2006.12.009. PubMed PMID: 17306896; PMCID: PMC1995559.

35. Battaglioni S, Benjamin D, Walchli M, Maier T, Hall MN. mTOR substrate phosphorylation

in growth control. Cell. 2022;185(11):1814-36. Epub 20220516. doi:

10.1016/j.cell.2022.04.013. PubMed PMID: 35580586.

36. Kim DH, Sarbassov DD, Ali SM, King JE, Latek RR, Erdjument-Bromage H, Tempst P,

Sabatini DM. mTOR interacts with raptor to form a nutrient-sensitive complex that signals to

the cell growth machinery. Cell. 2002;110(2):163-75. doi: 10.1016/s0092-8674(02)00808-5.

PubMed PMID: 12150925.

37. Schmidt EK, Clavarino G, Ceppi M, Pierre P. SUnSET, a nonradioactive method to monitor

protein synthesis. Nat Methods. 2009;6(4):275-7. Epub 20090322. doi: 10.1038/nmeth.1314.

PubMed PMID: 19305406.

38. Zhang T, Inesta-Vaquera F, Niepel M, Zhang J, Ficarro SB, Machleidt T, Xie T, Marto JA,

Kim N, Sim T, Laughlin JD, Park H, LoGrasso PV, Patricelli M, Nomanbhoy TK, Sorger

PK, Alessi DR, Gray NS. Discovery of potent and selective covalent inhibitors of JNK.

Chem Biol. 2012;19(1):140-54. doi: 10.1016/j.chembiol.2011.11.010. PubMed PMID:

22284361; PMCID: PMC3270411.

39. Moerke NJ, Aktas H, Chen H, Cantel S, Reibarkh MY, Fahmy A, Gross JD, Degterev A,

Yuan J, Chorev M, Halperin JA, Wagner G. Small-molecule inhibition of the interaction

between the translation initiation factors eIF4E and eIF4G. Cell. 2007;128(2):257-67. doi:

10.1016/j.cell.2006.11.046. PubMed PMID: 17254965.

40. Tolcher A, Goldman J, Patnaik A, Papadopoulos KP, Westwood P, Kelly CS, Bumgardner

W, Sams L, Geeganage S, Wang T, Capen AR, Huang J, Joseph S, Miller J, Benhadji KA,

Brail LH, Rosen LS. A phase I trial of LY2584702 tosylate, a p70 S6 kinase inhibitor, in

patients with advanced solid tumours. Eur J Cancer. 2014;50(5):867-75. Epub 20140115.

doi: 10.1016/j.ejca.2013.11.039. PubMed PMID: 24440085.

41. Porter AG, Janicke RU. Emerging roles of caspase-3 in apoptosis. Cell Death Differ.

1999;6(2):99-104. doi: 10.1038/sj.cdd.4400476. PubMed PMID: 10200555.

42. Boulares AH, Yakovlev AG, Ivanova V, Stoica BA, Wang G, Iyer S, Smulson M. Role of

poly(ADP-ribose) polymerase (PARP) cleavage in apoptosis. Caspase 3-resistant PARP

mutant increases rates of apoptosis in transfected cells. J Biol Chem. 1999;274(33):22932-40.

doi: 10.1074/jbc.274.33.22932. PubMed PMID: 10438458.

43. Huang J, Manning BD. The TSC1-TSC2 complex: a molecular switchboard controlling cell

growth. Biochem J. 2008;412(2):179-90. doi: 10.1042/BJ20080281. PubMed PMID:

18466115; PMCID: PMC2735030.

44. Dong B, Jaeger AM, Hughes PF, Loiselle DR, Hauck JS, Fu Y, Haystead TA, Huang J,

Thiele DJ. Targeting therapy-resistant prostate cancer via a direct inhibitor of the human heat

shock transcription factor 1. Sci Transl Med. 2020;12(574). doi:

10.1126/scitranslmed.abb5647. PubMed PMID: 33328331; PMCID: PMC10571035.

45. Sengupta S, Giaime E, Narayan S, Hahm S, Howell J, O'Neill D, Vlasuk GP, Saiah E.

Discovery of NV-5138, the first selective Brain mTORC1 activator. Sci Rep.

2019;9(1):4107. Epub 20190311. doi: 10.1038/s41598-019-40693-5. PubMed PMID:

30858438; PMCID: PMC6412019.

46. Kato T, Pothula S, Liu RJ, Duman CH, Terwilliger R, Vlasuk GP, Saiah E, Hahm S, Duman

RS. Sestrin modulator NV-5138 produces rapid antidepressant effects via direct mTORC1

activation. J Clin Invest. 2019;129(6):2542-54. Epub 20190416. doi: 10.1172/JCI126859.

PubMed PMID: 30990795; PMCID: PMC6546461.

47. Wolfson RL, Chantranupong L, Saxton RA, Shen K, Scaria SM, Cantor JR, Sabatini DM.

Sestrin2 is a leucine sensor for the mTORC1 pathway. Science. 2016;351(6268):43-8. Epub

20151008. doi: 10.1126/science.aab2674. PubMed PMID: 26449471; PMCID:

PMC4698017.

48. Yoon YJ, Kim JA, Shin KD, Shin DS, Han YM, Lee YJ, Lee JS, Kwon BM, Han DC.

KRIBB11 inhibits HSP70 synthesis through inhibition of heat shock factor 1 function by

impairing the recruitment of positive transcription elongation factor b to the hsp70 promoter.

J Biol Chem. 2011;286(3):1737-47. Epub 20101115. doi: 10.1074/jbc.M110.179440.

PubMed PMID: 21078672; PMCID: PMC3023468.

49. Caserta TM, Smith AN, Gultice AD, Reedy MA, Brown TL. Q-VD-OPh, a broad spectrum

caspase inhibitor with potent antiapoptotic properties. Apoptosis. 2003;8(4):345-52. doi:

10.1023/a:1024116916932. PubMed PMID: 12815277.

50. Chauvier D, Ankri S, Charriaut-Marlangue C, Casimir R, Jacotot E. Broad-spectrum caspase

inhibitors: from myth to reality? Cell Death Differ. 2007;14(2):387-91. Epub 20060929. doi:

10.1038/sj.cdd.4402044. PubMed PMID: 17008913.

51. Miao EA, Rajan JV, Aderem A. Caspase-1-induced pyroptotic cell death. Immunol Rev.

2011;243(1):206-14. doi: 10.1111/j.1600-065X.2011.01044.x. PubMed PMID: 21884178;

PMCID: PMC3609431.

52. Shan B, Pan H, Najafov A, Yuan J. Necroptosis in development and diseases. Genes Dev.

2018;32(5-6):327-40. doi: 10.1101/gad.312561.118. PubMed PMID: 29593066; PMCID:

PMC5900707.

53. Chen X, Comish PB, Tang D, Kang R. Characteristics and Biomarkers of Ferroptosis. Front

Cell Dev Biol. 2021;9:637162. Epub 20210121. doi: 10.3389/fcell.2021.637162. PubMed

PMID: 33553189; PMCID: PMC7859349.

54. Stockwell BR. Ferroptosis turns 10: Emerging mechanisms, physiological functions, and

therapeutic applications. Cell. 2022;185(14):2401-21. doi: 10.1016/j.cell.2022.06.003.

PubMed PMID: 35803244; PMCID: PMC9273022.

55. Yoshii SR, Mizushima N. Monitoring and Measuring Autophagy. Int J Mol Sci. 2017;18(9).

Epub 20170828. doi: 10.3390/ijms18091865. PubMed PMID: 28846632; PMCID:

PMC5618514.

56. Blommaart EF, Krause U, Schellens JP, Vreeling-Sindelarova H, Meijer AJ. The

phosphatidylinositol 3-kinase inhibitors wortmannin and LY294002 inhibit autophagy in

isolated rat hepatocytes. Eur J Biochem. 1997;243(1-2):240-6. doi: 10.1111/j.1432-

1033.1997.0240a.x. PubMed PMID: 9030745.

57. Schubert U, Anton LC, Gibbs J, Norbury CC, Yewdell JW, Bennink JR. Rapid degradation

of a large fraction of newly synthesized proteins by proteasomes. Nature.

2000;404(6779):770-4. doi: 10.1038/35008096. PubMed PMID: 10783891.

58. Kim LC, Cook RS, Chen J. mTORC1 and mTORC2 in cancer and the tumor

microenvironment. Oncogene. 2017;36(16):2191-201. Epub 20161017. doi:

10.1038/onc.2016.363. PubMed PMID: 27748764; PMCID: PMC5393956.

59. Zou Z, Tao T, Li H, Zhu X. mTOR signaling pathway and mTOR inhibitors in cancer:

progress and challenges. Cell Biosci. 2020;10:31. Epub 20200310. doi: 10.1186/s13578-020-

00396-1. PubMed PMID: 32175074; PMCID: PMC7063815.

60. Goloudina AR, Demidov ON, Garrido C. Inhibition of HSP70: a challenging anti-cancer

strategy. Cancer Lett. 2012;325(2):117-24. Epub 20120628. doi:

10.1016/j.canlet.2012.06.003. PubMed PMID: 22750096.

61. Jhaveri K, Taldone T, Modi S, Chiosis G. Advances in the clinical development of heat

shock protein 90 (Hsp90) inhibitors in cancers. Biochim Biophys Acta. 2012;1823(3):742-

55. Epub 20111029. doi: 10.1016/j.bbamcr.2011.10.008. PubMed PMID: 22062686;

PMCID: PMC3288123.

62. Pasqua AE, Sharp SY, Chessum NEA, Hayes A, Pellegrino L, Tucker MJ, Miah A, Wilding

B, Evans LE, Rye CS, Mok NY, Liu M, Henley AT, Gowan S, De Billy E, Te Poele R,

Powers M, Eccles SA, Clarke PA, Raynaud FI, Workman P, Jones K, Cheeseman MD. HSF1

Pathway Inhibitor Clinical Candidate (CCT361814/NXP800) Developed from a Phenotypic

Screen as a Potential Treatment for Refractory Ovarian Cancer and Other Malignancies. J

Med Chem. 2023;66(8):5907-36. Epub 20230405. doi: 10.1021/acs.jmedchem.3c00156.

PubMed PMID: 37017629; PMCID: PMC10150365.

63. Ali ES, Mitra K, Akter S, Ramproshad S, Mondal B, Khan IN, Islam MT, Sharifi-Rad J,

Calina D, Cho WC. Recent advances and limitations of mTOR inhibitors in the treatment of

cancer. Cancer Cell Int. 2022;22(1):284. Epub 20220915. doi: 10.1186/s12935-022-02706-8.

PubMed PMID: 36109789; PMCID: PMC9476305.

64. Wood KC, Gutkind JS. Challenges and Emerging Opportunities for Targeting mTOR in

Cancer. Cancer Res. 2022;82(21):3884-7. doi: 10.1158/0008-5472.CAN-22-0602. PubMed

PMID: 36321262; PMCID: PMC9634680.

65. Thomson AW, Turnquist HR, Raimondi G. Immunoregulatory functions of mTOR

inhibition. Nat Rev Immunol. 2009;9(5):324-37. doi: 10.1038/nri2546. PubMed PMID:

19390566; PMCID: PMC2847476.

Figure 1

  FIGURE LEGENDS:

Figure 1: HSF1 is required to suppress proteomic instability and amyloidogenesis in

MPNST cells. (A) Violin plots of *HSF1* expression in human cancer tissues showing *NF1*

deletion (median and IQR, n=6424, 1598, or 88, Kruskal-Wallis test). Data are generated by the

TGCA Research Network (https://www.cancer.gov/tcga). (B) Kaplan-Meier survival curves of

patients whose tumors harbor *NF1* deletion (n=420 or 419, Log-rank test). Data are generated by

the TGCA Research Network (https://www.cancer.gov/tcga). (C) and (D) The impacts of *HSF1*

depletion on protein polyubiquitination in both 90-8TL cells and primary human Schwann cells.

Cells were transduced with lentiviral shRNAs for 4 days and global protein Lys48-specific

ubiquitination was detected by immunoblotting in both detergent-soluble and -insoluble fractions

of whole cell lysates (C). Protein ubiquitination was quantitated using Fiji imaging software (D)

(mean ± SD, n=3 independent experiments, Two-way ANOVA). (E) and (F) Congo Red (CR)

staining of 90-8TL cells and primary human Schwann cells with and without *HSF1* depletion,

analyzed by flow cytometry (E). Quantitation of CR staining using the median fluorescence

intensity (F) (mean ± SD, n=3 independent experiments, One-way ANOVA). (G)-(J)

Quantitation of soluble amyloid oligomers and insoluble amyloid fibrils by ELISA using A11

and OC antibodies, respectively (mean ± SD, n=3 independent experiments, One-way ANOVA).

1000

1001

1002

Figure 2

Figure 2: HSF1 counters the tumor-suppressive effects of amyloids in MPNST cells.

(A) and (B) The growth curves of 90-8TL and primary human Schwann cells with and without

HSF1 depletion and 20 μM CR treatment (mean ± SD, n=3 independent experiments, Two-way

ANOVA). (C) Co-IP of amyloid oligomers (AOs) with HSF1 in 90-8TL cells. (D)-(F) Detection

of AO–HSF1 interactions by PLA flow cytometry in 90-8TL and S462 cells. Quantitation of the

PLA signals using the median fluorescence intensity (mean ± SD, n=3 or 4 independent

experiments, Student's t test). (G) and (H) Detection of AO–HSP60 interactions by PLA flow

cytometry in S462 cells following *HSF1* depletion (G). Quantitation of the PLA signals using the

median fluorescence intensity (H) (mean ± SD, n=3 independent experiments, Student's t test).

(I) Schematic depiction of AO neuralization by HSF1 and HSP60 attack by AO following *HSF1*

depletion in mitochondria.

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

Figure 3

Figure 3: JNK senses *HSF1* deficiency to repress mTORC1 and protein translation in

MPNST cells. (A) Detection of JNK activation and mTORC1 inhibition following *HSF1*

depletion in S462 and primary human Schwann cells by immunoblotting. (B) Quantitation of (A)

using Fiji imaging software (mean ± SD, n=3 independent experiments, One-way ANOVA). (C)

Detection of HSF1-JNK, JNK-RAPTOR, and mTOR-RAPTOR interactions by PLA in S462

cells with and without *HSF1* depletion. (D) and (E) Detection and quantitation of JNK-RAPTOR

and mTOR-RAPTOR interactions by PLA flow cytometry in 90-8TL cells (mean ± SD, n=3

independent experiments, One-way ANOVA). (F) Measurement of protein translation rates in

90-8TL and primary human Schwann cells with and without *HSF1* depletion by puromycin

labeling, analyzed by flow cytometry. Cells were treated with 1 μM JNK-IN-8 for 24 hrs before

labeling with 100 nM 6-FAM-dc-puromycin for 30 min (mean ± SD, n=3 independent

experiments, Two-way ANOVA). (G) Heatmap of protein translation rates measured by

puromycin labeling in diverse human cancer cell lines with and without *HSF1* depletion. (H)

Schematic depiction of regulation of protein translation by HSF1.

1040

1041

1042

1043

1044

1045

1046

1047

1048

Figure 4

Figure 4: Protein translation is causally related to amyloidogenesis in MPNST cells.

(A) and (B) Detection of mTORC1 stimulation following JNK inhibition in 90-8TL and immortalized human Schwann cells by immunoblotting. Cells were treated with JNK-IN-8 for 2 days. (C) and (D) Quantitation of soluble AOs and insoluble AFs by ELISA in 90-8TL and human Schwann cells with and without 3 μM JNK-IN-8 treatment for 3 days (mean ± SD, n=3 independent experiments, Student's t test). (E) Detection of LY2584702-mediated S6K inhibition in 90-8TL cells by immunoblotting. Cells were treated with 20 μM LY2584702 for 2 days. (F) Quantitation of soluble AOs and insoluble AFs by ELISA in 90-8TL cells with and without 20 μM LY2584702 treatment for 2 days (mean ± SD, n=3 independent experiments, Student's t test). (G) CR staining of 90-8TL and immortalized human Schwann cells with and without JNK-IN-8 and 4EGI-1 treatment, detected by flow cytometry. (H) Quantitation of (G) using the median fluorescence intensity (mean ± SD, n=3 independent experiments, One-way ANOVA). (I) Quantitation of amyloids by CR staining in 90-8TL cells with and without *HSF1* depletion and 3 μM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (J) Quantitation of caspase 3 cleavage by ELISA in 90-8TL cells with and without *HSF1* depletion and 3 μM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (K) Quantitation of amyloids by CR staining in immortalized human Schwann cells with and without *HSF1* depletion and 3 μM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (L) Quantitation of caspase 3 cleavage by ELISA in immortalized human Schwann cells with and without *HSF1* depletion and 3 μM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (M) Schematic depiction of the adaptation of MPNST cells to *HSF1* depletion by activating JNK to suppress mTORC1 and protein translation, thereby alleviating amyloidogenesis and cytotoxicity.

Figure 5

Figure 5: mTORC1 stimulation aggravates the amyloidogenesis and cytotoxicity elicited by

HSF1 inhibition. (A) Immunoblotting detection of JNK activation, mTORC1 activity, and

protein polyubiquitination in S462 cells following stable *TSC2* knockdown and/or HSF1

inhibition by 10 µM DTHIB for 3 days. (B) Quantitation of soluble AOs by flow cytometry

using A11 antibody staining in S462 cells with and without *TSC2* knockdown and 10 µM

DTHIB treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (C)

Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

without stable *TSC2* knockdown and DTHIB (10 µM) or combined DTHIB and LY2584702 (20

µM) treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). Cells were pre-

treated with LY2584702 for 1 day followed by DTHIB treatment for another 3 days. (D)

Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

without stable *TSC2* knockdown and DTHIB (10 µM) or combined DTHIB and CR (30 µM)

treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). Cells were pre-

treated with CR for 1 day followed by DTHIB treatment for another 3 days. (E) Quantitation of

soluble AOs by flow cytometry using A11 antibody staining in S462 cells with and without 500

µM NV-5138 stimulation and 10 µM DTHIB treatment (mean ± SD, n=3 independent

experiments, One-way ANOVA). (F) Quantitation of cytotoxicity by flow cytometry using Live-

or-Dye stains in S462 cells with and without 500 µM NV-5138 stimulation and 10 µM DTHIB

or combined DTHIB and CR treatment (mean ± SD, n=5 independent experiments, One-way

ANOVA). (G) Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in

immortalized human Schwann cells with and without 500 µM NV-5138 stimulation and 10 µM

DTHIB treatment (mean ± SD, n=3 independent experiments, One-way ANOVA). (H)

Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

without 500 μM L-leucine stimulation and 10 μM DTHIB treatment (mean ± SD, n=4

independent experiments, One-way ANOVA). (I) Quantitation of cytotoxicity by flow cytometry

using Live-or-Dye stains in immortalized human Schwann cells with and without 500 μM L-

leucine stimulation and 10 μM DTHIB treatment (mean ± SD, n=3 independent experiments,

One-way ANOVA). (J) Quantitation of soluble AOs by ELISA in S462 cells treated with and

without 800 μM L-leucine and 20 μM KRIBB11 or 30 μM CR (mean ± SD, n=3 independent

experiments, One-way ANOVA). (K) Quantitation of cytotoxicity by flow cytometry using Live-

or-Dye stains in S462 cells treated as above (mean ± SD, n=4 independent experiments, One-

way ANOVA). (L) Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in

S462 cells stably expressing inducible shRNAs with and without L-leucine stimulation. Cells

were first cultured with or without 1.2 mM L-leucine for 2 days, followed by addition of 20

ng/ml doxycycline for 7 days (mean ± SD, n=3 independent experiments, One-way ANOVA).

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

Figure 6

Figure 6: Concurrent HSF1 inhibition and mTORC1 stimulation instigate widespread non-apoptotic cell death. (A) and (B) Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains combined with cleaved caspase 3 (Asp175) antibody staining. S462 cells with and without stable *TSC2* knockdown were treated with DMSO, 10 μM DTHIB, or combined DTHIB and 30 μM Q-VD-OPh (mean ± SD, n=3 independent experiments, Two-way ANOVA). (C) Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and without stable *TSC2* knockdown. Cells were treated with 10 μM DTHIB alone or co-treated with 10 μM DTHIB and 30 μM Necrostatin-1 or 20 μM Liproxstatin-1 for 3 days (mean ± SD, n=3 independent experiments, Two-way ANOVA). (D) Immunoblotting detection of ferroptosis markers in S462 cells with and without stable *TSC2* knockdown treated with 10 μM DTHIB alone or co-treated with 10 μM DTHIB and 20 μM liproxstatin-1 for 3 days. Erastin was included as a positive control to induce canonical ferroptosis. (E) Immunoblotting detection of autophagy markers in S462 cells with stable *TSC2* knockdown treated with 10 μM DTHIB alone or co-treated with 10 μM DTHIB and 3 μM wortmannin for 3 days. Rapamycin was included as a positive control to induce autophagy. (F) Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and without stable *TSC2* knockdown. Cells were treated with 10 μM DTHIB alone or co-treated with 10 μM DTHIB and 3 μM wortmannin for 3 days (mean ± SD, n=3 independent experiments, Two-way ANOVA). (G) Schematic depiction of instigation of cell death by severe proteomic imbalance, owing to simultaneous mTORC1 stimulation (❶) and HSF1 inhibition (❷). In cancer cells, constitutive HSF1 activation provides extra chaperoning capacity to cope with elevated protein misfolding, partly due to enhanced protein synthesis and widespread genetic mutations. Nevertheless, amyloids still emerge, although at low levels. Importantly, HSF1 can neutralize highly toxic amyloid

oligomers, averting lethal consequences. By contrast, HSF1 inhibition diminishes chaperoning

capacity, insufficient to counterbalance the robust protein translation. mTORC1 stimulation

further aggravates this proteomic imbalance, which, in turn, promotes amyloidogenesis and

proteomic catastrophe. In consequence, the amounts of amyloid oligomers surpass the

neutralizing capacity of HSF1, leading to cell death; nonetheless, it remains unclear how this

proteomic catastrophe-induced death (PCID) occurs.

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

1163

Figure 7: Concurrent HSF1 inhibition and mTORC1 stimulation suppress *in vivo* tumor growth. (A) Combined KRIBB11 treatment and L-leucine stimulation impaired the *in vivo* growth of xenografted S462 human MPNST cells in NIH-III nude mice (mean ± SEM, n=8 mice per group, $\log_2$ transformation for Two-way ANOVA). (B) Combined KRIBB11 treatment and L-leucine stimulation impeded the *in vivo* growth of xenografted B16-F10 murine melanoma cells in C57BL/6J mice (mean ± SEM, n=8 mice per group, $\log_2$ transformation for Two-way ANOVA). (C) Quantitation of phospho-S6RP by immunoblotting in treated melanomas. Images show representative 3 tumors each group (mean ± SEM, n=5 tumors per group, One-way ANOVA). (D) Quantitation of *Hsp72* and *Hsp25* mRNAs in treated melanomas by RT-qPCR (mean ± SD, n=5 tumors per group, One-way ANOVA). (E) Quantitation of soluble AOs in treated B16F10 melanomas by ELISA ((mean ± SD, n=5 tumors per group, One-way ANOVA). (F) and (G) Quantitation of soluble AOs in brains and spleens from treated C57BL/6J mice by ELISA (mean ± SD, n=5 brains or spleens per group, One-way ANOVA). (H) Schematic depiction of the concept of provoking proteomic catastrophe to combat malignancy. On the one hand, in cancer cells, mTORC1 is inevitably activated to stimulate protein translation, markedly augmenting protein quantity. On the other hand, the extra chaperoning capacity governed by HSF1, albeit dispensable for normal life, becomes necessary to ensure sufficient protein quality in cancer cells, thereby counterbalancing augmented protein quantity and suppressing proteomic instability. Thus, proteomic balance promotes malignant growth. By contrast, disrupting proteomic balance, through HSF1 inhibition, is sufficient to elicit proteomic instability and tumor suppression. However, simultaneous mTORC1 stimulation can remarkably drive proteomic imbalance, causing proteomic catastrophe that leads to tumor suppression.