Driving proteomic imbalance to combat Neurofibromatosis type I (NF1)-

associated malignancy

Babul Moni Ram[1], Kun-Han Chuang[1,2], and Chengkai Dai[1,*]

[1]Mouse Cancer Genetics Program, Center for Cancer Research, National Cancer Institute,

Frederick, MD 21702, USA

[2]Present address: School of Basic Medical Sciences, Fujian Medical University, Fuzhou, Fujian

350108, China

*Correspondence: Chengkai.dai@nih.gov

ABSTRACT

Unlike genomic instability, the implications of proteomic instability in cancer remain

ambiguous. By governing the proteotoxic stress response, heat shock factor 1 (HSF1) sustains

proteomic stability upon environmental insults. Apart from its importance to stress resistance and

survival, HSF1 is emerging as a powerful, generic oncogenic enabler. In the Neurofibromatosis

type I (*NF1*)-deficient malignant peripheral nerve sheath tumor (MPNST) cells, *HSF1* depletion

prompted protein polyubiquitination, aggregation, and even tumor-suppressive amyloidogenesis.

In stark contrast, *HSF1* is dispensable for the proteome of non-transformed human Schwann

cells. Mechanistically, in MPNST cells HSF1 defends the essential mitochondrial chaperone

HSP60 against the direct assault from soluble amyloid oligomers. To survive and adapt to

impaired protein quality, owing to *HSF1* deficiency, MPNST cells mobilized JNK to repress

mTORC1 and protein translation, thereby attenuating protein quantity to alleviate proteomic

imbalance. mTORC1 stimulation, via either pharmacological JNK blockade, genetic *TSC2*

depletion, or supplement with the leucine analog NV-5138, markedly aggravated the proteomic

imbalance elicited by *HSF1* deficiency. This profound proteomic imbalance instigated non-

apoptotic cell death partly through unchecked amyloidogenesis. Thus, HSF1 safeguards the

cancer proteome to enable the oncogenic potential of mTORC1. Counterintuitively, this proof-

of-concept study highlights driving proteomic imbalance as a new therapeutic concept to combat

malignancies.

INTRODUCTION

Compared to genomic instability, proteomic instability in cancer is much less appreciated. Malignant transformation provokes proteotoxic stress and destabilizes the proteome.[1, 2] Several mechanisms, including reactive oxygen species, genetic mutations, and uncontrolled protein translation, all lead to increased protein damage, misfolding, as well as aggregation.[1-3] Of great interest, amyloidogenesis, a hallmark of neurodegenerative disorders,[4] also occurs in cancer.[5] Unlike neurons, cancerous cells manage to contain amyloidogenesis and counteract the cytotoxic effects of amyloids. Among various cellular anti-amyloid mechanisms is heat shock factor 1 (HSF1) and its mediated heat shock response or proteotoxic stress response (HSR/PSR), a powerful cytoprotective transcriptional program.[6-8] In accordance with its role in repressing proteomic instability, HSF1 has emerged as a potent oncogenic enabler, independently of driver mutations.[9-15]

Sharply contrasting with its dispensability for primary and non-transformed cells, HSF1 becomes essential to numerous cancer cell lines,[9,13,16-18] owing to their elevated intrinsic proteotoxic stress. This phenomenon is referred to as "HSF1 or non-oncogene addiction of cancer",[2,19] justifying HSF1 as a promising therapeutic target.

Neurofibromatosis type 1 (NF1) is a cancer predisposition syndrome afflicting approximately 1 in 3000 individuals worldwide.[20] About 30-50% of NF1 individuals develop benign plexiform neurofibromas, which, unlike dermal neurofibromas, can transform into highly aggressive malignant peripheral nerve sheath tumors (MPNSTs) with poor prognosis and survival.[21] The *NF1* gene encodes neurofibromin, a 250-280 kDa RAS GTPase activating protein (GAP) that

negatively regulates RAS activity.[22] *NF1* inactivation in MPNST cells dysregulates RAS/MAPK

signaling which, in addition to stimulating cell growth, leads to constitutive HSF1 activation.[13]

Concomitantly, HSF1 overexpression and activation are seen in both *in vitro* and *in vivo* MPNST

models as well as primary NF1 tumors.[13] Importantly, *Hsf1* deficiency impaired the NF1-

associated tumorigenesis in mice in part by attenuating oncogenic RAS signaling.[13]

Utilizing the NF1 tumor model system, we herein demonstrate HSF1 as a key player in

countering proteomic instability in cancer. We discover that amyloidogenesis occurs in *NF1-*

deficient MPNST cells, partly due to hyperactive mTORC1 and augmented protein translation.

HSF1 inhibition destabilizes the cancer proteome, promoting global protein polyubiquitination,

aggregation, as well as amyloidogenesis. Importantly, this proteomic instability causes

cytotoxicity. To adapt and survive HSF1 inhibition, MPNST cells mobilize JNK to suppress

mTORC1 and thereby mitigate protein translation. Disrupting this adaptative response through

JNK inhibition aggravates amyloidogenesis and cytotoxicity. Similarly, mTORC1 stimulation,

via either genetic *TSC2* depletion or supplement with the leucine analog NV-5138, synergizes

with HSF1 inhibition to inflict profound proteomic chaos and widespread non-apoptotic cell

death. Of note, impairment of amyloidogenesis markedly blocks cell death. Thus, our findings

illuminate a prominent role of HSF1 in antagonizing proteomic instability in cancerous cells,

constituting a generic pro-oncogenic mechanism. Moreover, this study exemplifies a proof of

concept that highlights driving proteomic imbalance as a counterintuitive therapeutic strategy.

RESULTS

HSF1 counters proteomic instability in *NF1*-deficient MPNST cells

Previously, we found that loss of *NF1* leads to constitutive activation of HSF1 and that *Hsf1* deficiency impairs the tumorigenesis in *Nf1*[+/-]; *p53*[+/-] (NPcis) mice.[13] In human cancers, *NF1* deletion is associated with elevated *HSF1* gene expression (Figure 1A). Moreover, the expression of *NF1* and *HSF1* genes are inversely correlated in human cancers (Figure S1A). Importantly, among patients whose tumors harbor *NF1* deletion higher *HSF1* expression is correlated with shortened survival (Figure 1B). Collectively, these findings support a pro-oncogenic role of HSF1 in *NF1*-deficient malignancies.

Given its critical role in governing protein folding, we were curious about how *HSF1* depletion impacts the proteomes of *NF1*-deficient MPNST cells as well as *NF1*-proficient non-transformed human Schwann cells (HSCs), both primary and immortalized. HSCs are considered as the cell-of-origin of MPNSTs.[20,22] In three different MPNST cell lines (90-8TL, S462, and SNF96.2), transient depletion of *HSF1*, using two previously validated lentiviral shRNAs,[9,13] all caused an increase in global protein Lys48-specific polyubiquitination, a molecular mark for proteasomal degradation,[23] in both detergent-soluble and -insoluble fractions (Figures 1C, 1D, S1B, and S1C). This increased polyubiquitination signifies proteome-wide misfolding and aggregation. In stark contrast, this change was not observed in HSCs (Figures 1C and 1D). Thus, *HSF1* depletion provokes proteomic instability in MPNST cells, but not in HSCs.

Most cellular protein aggregates are amorphous; however, a tiny fraction becomes highly ordered and enriched with β-sheet structures. These special protein aggregates, namely amyloids, are

closely associated with neurodegenerative disorders in humans, especially Alzheimer's disease.[24]

To assess whether *HSF1* depletion could elicit amyloidogenesis, we stained both MPNST cells and

HSCs with Congo red (CR), a widely applied amyloid-binding fluorescent dye.[25] Compared to

HSCs, MPNST cells displayed more intense CR staining (Figure 1E), suggesting increased

amyloids in malignant Schwann cells. Importantly, *HSF1* depletion further intensified CR staining

in MPNST cells, but not in HSCs (Figures 1E, 1F, S1D, and S1E). Similar results were obtained

by Thioflavin T staining (Figure S1F), another popular amyloid dye.[25] Importantly, the induction

of amyloids was further confirmed by ELISA using two conformation-specific antibodies (A11

and OC), which recognize soluble amyloid oligomers (AOs) and insoluble amyloid fibrils (AFs),

respectively.[26,27] Upon *HSF1* depletion, all three MPNST cell lines displayed elevated levels of

A11$^+$ AOs and OC$^+$ AFs (Figures 1G, 1H, S1G-S1J). Again, no evident increase in amyloids was

detected in HSCs (Figures 1I and 1J). Together, these findings indicate that HSF1 is required to

safeguard the proteome of malignant Schwann cells and particularly to repress amyloidogenesis;

on the contrary, it is dispensable for the proteome of non-transformed Schwann cells.

Unchecked proteomic instability is tumor suppressive

It has become evident that HSF1 acts as a generic oncogenic enabler; nevertheless, a universal

underlying mechanism remains undefined.

A healthy proteome enables the translation of genotypes into phenotypes. Thus, we reasoned that

preservation of proteomic stability is most likely a generic pro-oncogenic mechanism of HSF1.

To test this notion, we first determined whether proteomic instability, particularly

amyloidogenesis, is causally related to the cytotoxicity induced by *HSF1* depletion in cancer

cells. To block amyloidogenesis we employed CR, which is known to exert anti-amyloid effects

through its amyloid-binding activity.[5,28,29] Consistent with "HSF1 addiction of cancer", *HSF1*

depletion markedly impaired the viability of MPNST cells; importantly, CR treatment partially

rescued this impaired viability (Figures 2A, S2A, and S2B). In contrast, both *HSF1* depletion and

CR treatment did not affect HSCs (Figure 2B), congruent with a lack of proteomic instability and

amyloidogenesis in these non-transformed cells. Together, these findings support that unchecked

proteomic instability, particularly amyloidogenesis, is cytotoxic and tumor suppressive.

Moreover, HSF1 enables oncogenesis in part by constraining proteomic instability.

HSF1 neutralizes soluble amyloid oligomers to protect MPNST cells

Then, how does HSF1 counter amyloids to enable the growth and survival of MPNST cells?

Recently, by researching tissue overgrowth models we discovered that HSF1, surprisingly,

physically neutralizes soluble AOs, thereby preventing their assault on HSP60.[30] As an essential

mitochondrial chaperone, HSP60 functions as a guardian of the mitochondrial proteome.[30-32]

We wondered whether HSF1 acts the same way in both cancer and tissue overgrowth. Co-IP

experiments indicated that A11[+] AOs physically interacted with HSF1, but not major HSPs, in

MPNST cells (Figure 2C), consistent with our findings in tissue overgrowth models.[30] This AO-

HSF1 interaction was further validated by another independent technique, *in situ* Proximity

Ligation Assay (PLA) (Figures 2D-2F). Importantly, *HSF1* depletion promoted the AO-HSP60

interaction (Figure 2G and 2H). Loss of *HSP60* destabilized the mitochondrial proteome.[30,32]

Thus, this AO-HSP60 interaction signifies the instigation of mitochondrial damage and

cytotoxicity. Taken together, our findings reveal that HSF1 protects MPNST cells in part by

neutralizing soluble AOs to safeguard HSP60 and the mitochondrial proteome (Figure 2I).

MPNST cells react to HSF1 deficiency by activating JNK to attenuate translation

Given their dependency on HSF1, cancer cells are expected to mount certain responses to

survive and adapt to the lethal HSF1 inhibition. Previously, we uncovered that in *Hsf1*-deficient

mice, the stress-responsive JNK kinase becomes constitutively activated, which, in turn,

disassembles mTORC1.[33] JNK does so by phosphorylating both RAPTOR and mTOR, two core

components of mTORC1. This mTORC1 inhibition diminishes protein translation, leading to

reduced cell size and whole-body lean mass in *Hsf1*-deficient mice.[33]

We reasoned that cancer cells might exploit this molecular regulation for their survival and

adaptation. In MPNST cells *HSF1* knockdown activated JNK, indicated by elevated JNK

T183/Y185 phosphorylation; in parallel to JNK activation, there was mTORC1 inhibition,

evidenced by diminished p70S6K T389 phosphorylation and 4EBP1 S65/T70 phosphorylation

(Figures 3A, 3B, and S3A-S3D).[34,35] In contrast, *HSF*1 knockdown minimally affected JNK and

mTORC1 in HSCs (Figures 3A and 3B). Detected by PLA, in MPNST cells HSF1 interacted

with JNK; however, *HSF1* depletion promoted JNK-RAPTOR interactions but diminished

mTOR-RAPTOR interactions simultaneously (Figures 3C-3E). The mTOR-RAPTOR interaction

is essential to the assembly and activity of mTORC1.[36] Thus, these findings support that JNK

senses HSF1 deficiency to repress mTORC1. Of interest, compared to non-transformed cells,

this sensing mechanism appears more prominent in malignant cells.

As expected for mTORC1 inhibition, *HSF1* depletion reduced global protein translation rate in

MPNST cells (Figures 3F, S3E, and S3F), detected by puromycin labeling as described

previously.[33,37] Beyond MPNST cells, this regulation of translation by HSF1 was conserved in

diverse human cancer cell lines (Figure 3G). Importantly, treatment with JNK-IN-8, an

irreversible JNK inhibitor,[38] largely rescued this impaired protein translation owing to *HSF1*

depletion (Figures 3F, S3E, and S3F). Interestingly, JNK inhibition augmented the basal

translation rate as well. Consistent with these findings, in human cancers higher *HSF1*

expression is associated with lower JNK activation (Figure S3G); in turn, lower JNK activation

is accompanied by higher mTORC1 activity, reflected by higher 4EBP1 Ser65 phosphorylation

(Figure S3H). In aggregate, our findings demonstrate that JNK activation is causally linked to

inhibition of mTORC1 and translation. Therefore, through JNK suppression HSF1 permits

robust protein translation, which is fundamental to malignant growth (Figure 3H).

Lessening protein quantity is an adaptive response to alleviate proteomic instability

We next asked why cells respond to HSF1 deficiency by attenuating translation. While HSF1

controls protein quality via HSPs, mTORC1 governs protein quantity via translation. It would be

beneficial to lower protein quantity when protein quality is jeopardized. Thus, we reasoned that

this regulation might represent a cellular adaptive response to imbalanced protein quantity and

quality engendered by HSF1 deficiency.

Proteomic imbalance disrupts proteome homeostasis and contributes to proteomic instability,

which may ultimately trigger amyloidogenesis. To inflict proteomic imbalance, we first

stimulated protein translation in MPNST cells by inhibiting JNK. As expected, JNK-IN-8

reduced the phosphorylation of c-JUN, a canonical substrate of JNK,[34] in MPNST cells (Figure

4A). In support of enhanced protein translation, JNK-IN-8 further activated mTORC1 in these

cells (Figures 4A and S4A); in contrast, marginal effects were observed in HSCs (Figure 4B).

Congruently, JNK-IN-8 did not significantly reduce c-JUN phosphorylation in HSCs, supporting

a lack of JNK activation in these non-transformed Schwann cells. Intriguingly, the heightened

protein translation following JNK inhibition was accompanied by elevated amyloid levels in

MPNST cells (Figures 4C and S4B). On the contrary, there were no significant changes in

amyloid levels in HSCs (Figure 4D).

To determine the role of protein translation in amyloidogenesis, we chose to block translation

using two pharmacological inhibitors, 4EGI-1 and LY2584702, with distinct mechanisms of

action. While 4EGI-1 disrupts eIF4E-eIF4G interactions,[39] LY2584702 inhibits p70S6K.[40]

Interestingly, both inhibitors diminished the basal amyloid levels in MPNST cells (Figures

4E,4F, and S4C), revealing a causal relationship between protein translation and

amyloidogenesis. Importantly, translation inhibition also blocked the induction of amyloids by

JNK-IN-8 in MPNST cells (Figures 4G, 4H, S4D, and S4E), indicating that JNK suppresses

amyloidogenesis via translation inhibition. No significant effects were observed in HSCs

(Figures 4G and 4H).

Considering the role of translation in amyloidogenesis, we reasoned that preventing translation

attenuation would aggravate proteomic imbalance and enhance cytotoxicity in malignant cells.

As expected, JNK-IN-8 further elevated the amyloid levels in MPNST cells following *HSF1*

depletion, which was accompanied by exacerbated apoptosis, detected by caspase 3 cleavage

(Figures 4I, 4J, and S4F).[41] Importantly, CR treatment blocked apoptosis, detected by PARP

cleavage (Figure S4G),[42] causally implicating amyloidogenesis. In contrast, *HSF1* depletion and

JNK inhibition exerted no significant impacts on non-transformed Schwann cells (Figures 4K

and 4L). Thus, our findings support that proteomic imbalance in malignant cells contributes to

proteomic instability and amyloidogenesis. In stark contrast, there is no or little amyloidogenesis

in non-transformed cells, owing to their balanced protein quantity and quality. Moreover,

through translation attenuation, malignant cells attempt to survive and adapt to proteomic

imbalance (Figure 4M).

mTORC1 stimulation aggravates proteomic imbalance and cytotoxicity induced by HSF1

inhibition

Apart from mTORC1, JNK regulates many other targets. To stimulate mTORC1 and protein

translation specifically, we next genetically inhibited the tumor suppressor Tuberous Sclerosis

Complex subunit 2 (*TSC2*), a key negative regulator of mTORC1,[43] in MPNST cells. As

expected, knocking down *TSC2* in MPNST cells promoted phosphorylation of both p70S6K and

4EBP1, which was accompanied by elevated amyloids (Figures S5A-S5F). Like genetic *HSF1*

depletion, pharmacological inhibition of HSF1 by DTHIB elevated global protein

polyubiquitination and amyloids in MPNST cells (Figures 5A, 5B, and S5G), eliciting proteomic

instability. The newly developed small-molecule inhibitor DTHIB directly binds to the DNA-

binding domain of HSF1 and potently blocks the HSF1-mediated transcription.[44] Compared to

control cells, HSF1 inhibition by DTHIB induced considerably higher levels of amyloids in

TSC2-deficient MPNST cells (Figure 5B). Of note, *TSC2*-deficient cells displayed more

profound JNK activation and mTORC1 inhibition following DTHIB treatment, reflecting a

cellular response to severe proteotoxic stress. Consistently, compared to slight or intermediate

cytotoxicity in control MPNST cells, detected by plasma membrane permeabilization, *TSC2*-

deficient cells were highly sensitive to HSF1 inhibition (Figures 5C, 5D, and S5H-S5K).

Importantly, either translation inhibition or CR treatment largely blocked this cytotoxicity

(Figures 5C and 5D), causally implicating amyloidogenesis. In support of a role of HSF1 in

enabling *TSC2* deficiency, among patients whose tumors harbor *TSC2* deletion higher *HSF1*

gene expression is correlated with shortened survival (Figure S5L).

To exclude possible mTORC1-independent effects of *TSC2* deficiency, we alternatively fed cells

with NV-5138, a leucine analog currently under clinical trials as an antidepressant.[45,46]Amino

acids, particularly leucine, are potent activators of mTORC1.[47] Of interest, whereas NV-5138 is

capable of activating mTORC1, it cannot be utilized for protein synthesis.[45,46] To maximize

mTORC1 stimulation *ex vivo*, cells were cultured in leucine-free medium. Like *TSC2*

knockdown, NV-5138 stimulated mTORC1 and heightened amyloid levels in MPNST cells

(Figures 5E and 5F); however, it only elicited pronounced cytotoxicity with concurrent HSF1

inhibition (Figure 5G). Again, CR treatment largely blocked this cytotoxicity (Figure 5G). In

sharp contrast, combined NV-5138 and DTHIB treatment did not significantly impact non-

transformed Schwann cells (Figure 5H). Of interest, their mTORC1 signaling was less

responsive to NV-5138 stimulation (Figures 5I). Collectively, these findings indicate that

mTORC1 stimulation synergizes with HSF1 inhibition to rapidly trigger severe cytotoxicity in

MPNST cells, in part by prompting unconstrained proteomic imbalance and amyloidogenesis.

Unconstrained proteomic imbalance incites non-apoptotic, non-autophagic death

HSF1 inhibition alone induced both apoptotic and non-apoptotic death in MPNST cells,

distinguished by caspase 3 cleavage; intriguingly, concurrent mTORC1 stimulation caused a

striking shift towards non-apoptotic cell death, characterized by cell membrane permeabilization

without caspase 3 cleavage (Figures 6A and 6B).

To date, several forms of non-apoptotic cell death have been described, including pyroptosis,

necroptosis, ferroptosis, and autophagic death, all of which could result in loss of cell membrane

integrity. Corroborating a non-apoptotic mechanism, the pan-caspase inhibitor Q-VD-OPh did

not prevent the membrane permeabilization in *TSC2*-deficient MPNST cells treated with DTHIB

(Figures 6A and 6B).[48] However, it did block the apoptosis in control MPNST cells treated with

DTHIB (Figures 6A and S6A). As Q-VD-OPh inhibits caspase 1 and no caspase 1 cleavage was

detected (Figure S6B),[49] a key driver of pyroptosis,[50] this result does not support a pyroptotic cell

death either. Similarly, HSF1 inhibition did not induce necroptosis in *TSC2*-deficient MPNST

cells, evidenced by the lack of MLKL Ser358 phosphorylation (Figure S6B), an essential event

executing necroptosis.[51] Congruently, the necroptosis inhibitor necrostatin-1 exerted no effects

on the cell membrane disruption (Figures 6C and S6C).[51]

Intriguingly, HSF1 inhibition diminished glutathione peroxidase 4 (GPX4) but increased

transferrin receptor 1 (TFR1), which was accompanied with markedly elevated lipid

peroxidation, in *TSC2*-deficient MPNST cells (Figures 6D and S6D). All these changes are

indicative of ferroptosis.[52,53] Although in *TSC2*-deificent cells liproxstatin-1, a potent ferroptosis

inhibitor,[52,53] reversed the elevation of TFR1 elicited by the ferroptosis inducer erastin, it neither

blocked the marked increase in TFR1 nor stopped the membrane disruption induced by DTHIB

(Figures 6C, 6D, and S6C). Similarly, HSF1 inhibition also elicited autophagy, evidenced by

diminished p62/SQSTM1 and elevated LC3B-II (Figure 6E).[54] To determine whether autophagic

cell death was implicated, we employed the class III PI3K inhibitor wortmannin to block

autophagy.[55] As expected, wortmannin effectively prevented the decreases in p62/SQSTM1 and

LC3B-II induced by rapamycin in *TSC2*-deficient MPNST cells (Figure 6E), indicating

successful blockade of autophagy. Nevertheless, wortmannin failed to block the cell membrane

disruption in *TSC2*-deficient cells treated with HSF1 inhibitors (Figures 6F and S6E). These

findings suggest that both ferroptosis and autophagy are the consequences, rather than the

causes, of cell membrane disruption. Collectively, our findings support that uncontained

amyloids potently trigger cell death, which is neither apoptotic nor autophagic (Figure 6G).

DISCUSSIONS

Genomic instability has been broadly recognized as a driver of oncogenesis. In contrast, the implications of proteomic instability in cancer remain obscure.

Proteomic instability is tumor suppressive but inevitable

Proteomic instability does occur in cancer, evidenced by elevated protein polyubiquitination, aggregation, and even amyloidogenesis. In cancer, proteomic instability can be ascribed to several mechanisms, including protein oxidation by reactive oxygen species, protein misfolding owing to genetic mutations, and augmentation of protein dosage caused by aneuploidy.[1,3] Revealed by our recent studies, uncontrolled protein translation also contributes to amyloidogenesis. This is not entirely surprising, as it has been estimated that up to 30% of all newly synthesized polypeptides are misfolded and subjected to proteasomal degradation.[56] Imaginably, aberrant protein synthesis would increase the chance of producing protein aggregates and even amyloids, especially considering numerous genetic mutations affecting protein folding in cancer cells. Ironically, dysregulated protein translation is fundamental to malignant growth, making proteomic instability and amyloidogenesis inevitable.

At the cellular level, amyloids are toxic to both neurons and cancer cells. At the organismal level, amyloidogenesis is neurodegenerative but tumor suppressive. Mechanistically, our studies demonstrate that soluble AOs can directly attack the guardian of the mitochondrial proteome HSP60. Of course, mechanisms other than targeting HSP60 exist as well. Moreover, it remains to be elucidated what other proteins comprise cancer-associated amyloids, in addition to the Alzheimer's disease-associated Aβ.

HSF1 contains proteomic instability to enable malignancy

The multifaceted pro-oncogenic roles of HSF1 have been ascribed to a wide array of underlying

mechanisms; nonetheless, a universal one remains to be defined.

As an inevitable price of malignancy, proteomic instability must be contained by cancer cells to

avert tumor suppression and sustain malignant phenotypes. Now, our findings reveal that HSF1

contains proteomic instability, particularly amyloidogenesis, constituting a generic pro-

oncogenic mechanism. It is worth noting that HSF1 only contains and mitigates, but cannot

eradicate, amyloidogenesis, as uncontrolled protein translation is a direct cause of

amyloidogenesis in cancer cells. This dependency on HSF1 to contain proteomic instability

partly underlies the phenomenon of "HSF1 addiction of cancer".

Our studies further support that HSF1 contains amyloidogenesis via at least two mechanisms,

direct and indirect. On the one hand, through induction of HSPs, HSF1 ensures appropriate

protein folding to maintain a healthy proteome, thereby preventing the emergence of amyloids.

This is supported by the finding that HSF1 inhibition heightens protein ubiquitination,

aggregation, and amyloidogenesis. On the other hand, HSF1 can neutralize soluble AOs directly

once they emerge. Under this context, HSF1 inhibition would unleash AOs to assault HSP60,

inflicting mitochondrial damage and toxicity. Thus, in cancer cells HSF1 inhibition both enlarges

the quantity of amyloids and, more importantly, renders them toxic.

mTORC1 signaling becomes tumor suppressive in the absence of HSF1

Given the dependency of malignant growth on protein translation, unsurprisingly, mTORC1

signaling is hyperactive in human cancers, which is, canonically, considered oncogenic.[57,58]

Consistently, 4EBP1 Ser65 phosphorylation, indicative of mTORC1 activation, in tumor tissues

is correlated with shortened patients' survival (Figure S6F).

Unexpectedly, our current studies reveal that mTORC1 signaling becomes tumor suppressive in

the absence of HSF1, owing to profound disproportion between protein quantity and quality.

This unconstrained proteomic imbalance, in turn, exacerbates proteomic instability and tumor-

suppressive amyloidogenesis. The notable blockage of cytotoxicity by either translation

inhibitors or CR lends strong support to this model. Conceptually, HSF1 empowers the

oncogenic potential of mTORC1 signaling, reinforcing HSF1 as an oncogenic enabler. In line

with this notion, co-upregulation of *HSF1* mRNA and mTORC1 activity is significantly

associated with shortened patients' survival (Figure S6G).

Driving proteomic imbalance represents a new therapeutic concept

Currently popular chemo-and radiotherapies mainly target the cancer genome, causing DNA

damage. A new horizon in cancer therapy is targeting the cancer proteome. To date, much effort

has been devoted to inhibiting individual HSP families, such as HSP70 and HSP90.[59,60]

However, every cell is equipped with both constitutive and inducible HSPs. While the former

comprise the basal chaperoning capacity that is essential to life, the latter supply the extra

chaperoning capacity, which is dispensable for normal life and only required under stressful

conditions. Their structural and functional resemblances pose the biggest challenge for small

molecules to distinguish these cognate HSPs. Moreover, multiple families of HSPs, including

HSP100, HSP90, HSP70, HSP60, HSP40, and small HSPs, exist inside cells. Targeting single

HSP family would have limited therapeutic effects. Thus, new approaches or directions are

warranted to overcome these drawbacks.

Our and others' studies collectively support that malignant cells constantly endure intrinsic

proteotoxic stress, rendering HSF1 constitutively active. In turn, cancer cells become reliant on

the extra chaperoning capacity, conferred by HSF1, for their growth and survival. Thus, only

eliminating this extra chaperoning capacity but sparing the basal one would be an ideal strategy

to combat malignancy. Together, these concepts provide strong rationales for targeting HSF1 in

cancer. It is exciting to witness the recent launch of HSF1 inhibitors into clinical trials.[61]

In reaction to HSF1 inhibition, unsurprisingly, cancer cells attenuate protein synthesis. This

adaptive response underscores the importance of balanced protein quantity and quality to cellular

fitness, a key concept inspiring us to explore driving proteomic imbalance for cancer therapy. Of

note, breaking this adaptation may impede tumor evolution.

The widespread hyperactivation of mTORC1 in human cancers makes it an attractive therapeutic

target. Hence, an array of mTOR inhibitors have been developed, several of which are actively

under clinical evaluation.[58,62] Thus far, these inhibitors have displayed mostly cytostatic effects

and less than expected therapeutic efficacies.[62,63] Considering the essentiality of protein

translation to life, profound mTORC1 inhibition would inflict undesirable toxicities in normal

cells and tissues. In fact, mTORC1 inhibitors have long been known as immunosuppressants.[62,64]

Counterintuitively, we stimulated mTORC1 to drive proteomic imbalance, opposite to the

prevailing strategy in cancer therapy. mTORC1 stimulation, when combined with HSF1

inhibition, engendered prominent cytotoxic effects (Figure 6H), representing a key therapeutic

benefit. Moreover, this amyloid-dependent cell death offers a large therapeutic window, owing to

the lack of amyloidogenesis in primary cells. Thus, using the *NF1*-deficient MPNST model

system, we herein accomplished the proof-of-concept experiment to demonstrate the feasibility

of driving proteomic imbalance in cancer. Noteworthily, this therapeutic concept should be

applicable to a broad spectrum of human cancers, beyond this specific NF1-associated

malignancy.

ACKNOWLEDGMENTS

We would like to thank the Optical Microscopy and Image Analysis lab (OMAL) for their

assistance with the confocal microscopy studies. Schematic illustrations are created with

BioRender.com. This work was supported by the grant from Department of Defense

Congressionally Directed Medical Research Programs to C.D. (NF150028) and by the Intramural

Research Program of the NIH, National Cancer Institute, Center for Cancer Research

(1ZIABC011767). The content of this publication does not necessarily reflect the views or

policies of the Department of Health and Human Services, nor does mention of trade names,

commercial products, or organizations imply endorsement by the U.S. government.

AUTHOR CONTRIBUTIONS: B.M.R. and K-H.C. designed and conducted the experiments.

C.D. conceptualized and supervised this study and analyzed the results. B.M.R. and C.D. wrote

the manuscript.

DECLARATION OF INTERESTS: The authors declare no competing interests.

EXPERIMENTAL MATERIALS AND METHODS

Cells, Chemicals, and reagents

S462 (female) and 90-8TL (female) cells were kindly provided by Dr. Karen Cichowski.

sNF96.2 cells (male) and immortalized human Schwann cells (hTERT ipn02.3 2λ, female) were

purchased from ATCC. Primary human Schwann cells (Cat#1700, ScienCell Research

Laboratories) were grown on poly-L-lysine coated T-75 flasks and maintained in complete

Schwann Cell Medium (Cat#1701, ScienCell Research Laboratories). All cell lines, except 90-

8TL and HSCs (no known STR profiles available), were authenticated by ATCC. All cell lines,

except primary human Schwann cells, were maintained in DMEM supplemented with 10%

HyClone bovine growth serum and 1% penicillin–streptomycin. Cells were maintained in an

incubator with 5% CO_2 at 37 °C.

The following chemicals were purchased from commercial sources: JNK-IN-8 (Cat#S4901,

Selleckchem), 6-FAM-dC-puromycin (cat#NU-925-6FM, Jena Bioscience), Congo red

(cat#B2431014, Thermo Scientific Chemicals), Thioflavin T (cat#211760050, Thermo Scientific

Chemicals), LY2584702 (cat#S7698, Selleckchem), 4EGI-1 (cat#S7369, Selleckchem), and

DTHIB (cat# HY-138280, MedChemExpress), Q-VD-OPh (cat#HY-12305, MedChemExpress),

necrostatin-1 (cat#HY-15760, MedChemExpress), Liproxstatin-1 (cat#HY-12726,

MedChemExpress), wortmannin (cat#HY-10197, MedChemExpress), rapamycin (cat#HY-

10219, MedChemExpress), erastin (cat#HY-15763, MedChemExpress), MHY1485 (cat#HY-

B0795, MedChemExpress), NV-5138 HCl (cat#HY-114384B, MedChemExpress), and Liperfluo

(cat#L248, Dojindo Laboratories).

Rabbit phospho-4EBP1 S65/T70 antibody (cat#sc-12884-R), mouse monoclonal HSF1 (E-4,

cat#sc-17757) antibody, rabbit HSF1 antibody (H-311, cat#sc-9144), mouse monoclonal JNK

(D-2, cat#sc-7345), rabbit JNK1/3 (C-17, cat#sc-474) antibody were purchased from Santa

Cruz Biotechnology; antibodies for phosphor-4EBP1 Thr70 (cat#9455), phospho-4EBP1

Thr37/46 (236B4, cat#2855), phospho-p70S6K T389 (108D2, cat# 9234), phosphor-S6RP

Ser235/236 (D57.2.2E, cat#4858), p70 S6K (49D7, cat#2708), 4EBP1 (53H11, cat#9644),

S6RP (5G10, cat#2217), mTOR (7C10, cat#2983), phospho-JNK1/2 T183/Y185 (81E11,

cat#4668), JNK2 (56G8, cat#9258), c-Jun (60A8, cat#9165), phospho-c-Jun Ser63 II

(cat#9261), PARP (cat#9542), Tuberin/TSC2 (D93F12, cat#4308), SQSTM1/p62 (D1Q5S,

cat#39749), cleaved caspase 3 (Asp175) (5A1E, cat#9664), caspase 1(D7F10, cat#3866),

gasdermin D (E9S1X, cat#39754), phosphor-MLKL Ser358 (D6H3V, cat#91689), GPX4

(E5Y8K, cat#59735), and transferrin receptor/CD71 (H68.4, cat#46222) antibody were

purchased from Cell Signaling Technologies; mouse β-actin antibody-HRP (Cat#A00730)

conjugates were from GenScript. Anti-K48-specific ubiquitin antibody (Apu2, cat#05-1307)

and mouse monoclonal anti-RAPTOR antibody (1H6.2, cat#05-1470) were purchased from

EMD Millipore. Mouse monoclonal anti-JNK1 (C-terminal region) antibody (cat#JM2671)

was purchased from ECM Biosciences. Anti-amyloid oligomers (A11, cat#SPC-506D) and

anti-amyloid fibrils (OC, cat#SPC-507D) antibodies were purchased from StressMarq

Biosciences Inc. Mouse monoclonal anti-HSP60 antibody (LK-1, cat#ADI-SPA-806) was

purchased from Enzo Life Sciences Inc. β-Actin (cat#GTX629630), GAPDH

(cat#GTX627408), α-tubulin (GT114, cat#GTX628802), and LC3B antibody

(cat#GTX127375) was purchased from GeneTex Inc.

METHOD DETAILS

Lentiviral Particle Production and Transduction of lentiviral shRNAs

Lentiviral pLKO.1 shRNA plasmids targeting human *HSF1* were described previously[10]: TRCN0000007480 (hA6) and TRCN0000007483 (hA9). The sequence of control hairpins targeting a scrambled sequence with no known homology to any human genes is 5′-CCTAAGGTTAAGTCGCCCTCG-3′. The pLKO.1 shRNA plasmid targeting human *TSC2* was a gift from Do-Hyung Kim (Cat#15478, Addgene). Lentiviral pLKO shRNA plasmids were transiently co-transfected individually along with plasmids encoding ΔVPR and VSV-G into HEK293T cells using TurboFect™ (Cat#R0531, Thermo Scientific). Viral supernatants were collected after centrifugation at 1,000 rpm for 15 min. Viral transduction was achieved by incubating target cells with viral supernatants containing 10μg/ml polybrene (Cat#TR-1003-G, Millipore-Sigma) for 24h. For stable knockdown of *TSC2,* lentivirus particles were prepared using pLKO shRNA plasmids targeting human *TSC2*. Scrambled shRNA was used for stable control cells. After 4 days of transduction, cells were gradually selected using 1-4 μg/ml puromycin for 2 weeks.

Soluble and Insoluble Protein Fractionation

Equal numbers of cells were incubated with cold cell-lysis buffer (100 mM NaCl, 30 mM Tris-HCl pH 7.6, 1% Triton X-100, 20 mM sodium fluoride, 1mM EDTA, 1mM sodium orthovanadate, and 1x Halt™ protease inhibitor cocktail) on ice for 20 min. The crude lysates were first centrifuged at 500xg for 5 min at 4°C. The supernatants from low-speed (500xg) centrifugation were transferred to a new Eppendorf tube. The first pellets were resuspended in

DNA digestion solution (2 units DNase I, 1% Triton X-100, 10 mM Tris-HCl, 2.5 mM

MgCl2, 0.5 mM CaCl2, pH 7.6) for 30 min at RT. An equal volume of 1xPBS with 2% SDS

was then added to resuspend the first pellets for additional 30 min at RT. Following the

centrifugation at 17,000xg for 10 min at RT, the supernatants were removed. The supernatants

from low-speed centrifuge were added back to the second pellets, followed by centrifugation

at 17,000xg for 15 min at 4°C. The final supernatants and pellets from high-speed (17,000xg)

centrifugation were collected as detergent-soluble and -insoluble fractions, respectively.

Insoluble fractions were further sonicated in 1xPBS with 2% SDS at high intensity using a

Bioruptor® Sonication Device (Diagenode) for SDS-PAGE.

ThT and CR Staining for Flow Cytometry

After washing with 1xPBS, cells were fixed with 4% formaldehyde at RT for 10 min.

Following fixation and 1xPBS washing, cells were resuspended in 1 ml permeabilization

buffer (1xPBS with 0.5% Triton X-100 and 3 mM EDTA) and incubated on ice for 30 min.

Following washing with 1xPBS, Cells were stained with 10 μM ThT or CR dissolved in

1xPBS for 5 min. Cells were washed and resuspended in PBS and fluorescence signals were

measured by a FACSCalibur, LSRFortessa X-20 or LSR II flow cytometer (BD Biosciences).

Amyloid Oligomer and Fibril Quantitation by Direct ELISA and Flow Cytometry

To quantitate soluble amyloid prefibrillar oligomers, soluble cell lysates diluted in 1xPBS

from equal numbers of cells were coated on each well in a 96-well ELISA plate at 4°C

overnight followed by blocking (5% non-fat milk in TBS-0.05% Tween® 20) at RT for 1 hr.

Each well was incubated with 100 μl amyloid oligomer antibodies (A11, 1:1,000 diluted in

the blocking buffer) at RT for 2 hr. After washing with TBS-T, goat anti-rabbit secondary Ab

HRP conjugates (1:1,000 diluted in blocking buffer) were added to each well and incubated at

RT for 1 hr. Following washing, 100 μl 1-Step Ultra TMB-ELISA substrates (Cat#34029,

Thermo Scientific) were added to each well. After adding 100μl Stop solution, absorbance was

measured at 450 nm using a SpectraMax iD5 microplate reader (Molecular Devices).

To quantitate amyloid fibrils, detergent-insoluble fractions from equal numbers of cells were

extracted as described in Soluble and Insoluble Protein Fractionation. The final pellets were

collected as detergent-insoluble fractions and solubilized by sonication for 15 min in 1xPBS

with 2% SDS. Solubilized proteins diluted in 1xPBS were added to each well and incubated at

37°C without cover overnight to air dry the wells. The following steps were identical to the

oligomer detection except for the use of amyloid fibril antibodies (OC) as the primary Ab.

For amyloid oligomer and fibril quantification by flow cytometry, cells were collected by

trypsinization and fixed in 4% Formaldehyde at RT for 10 min. Following permeabilization using

0.3% Triton-X-100, samples were incubated with primary antibodies A11-FITC (1:100 in 5%

BSA, StressMarq Biosciences, cat# SPC-506D-FITC) or OC-PerCP (1:100 in 5% BSA,

StressMarq Biosciences, cat# SPC-507D-PCP) at RT for 3 hrs. The cells were washed with PBS

and analyzed using flow cytometry.

Measurement of Cell Viability

Cell viability was determined using the CellTiter-Blue® Cell Viability Assay (cat#G8080,

Promega). Cells were grown in 96 well plates (1×10^4 per well) for 24h. A 20μl CellTiter-Blue®

reagent was added to 100μl of media in each well. The plates were incubated for 2h at 37°C and

the fluorescence was recorded at 560/590 nm. The reagents were discarded, and the wells were

washed with PBS before adding fresh media. Similarly, subsequent readings were recorded at

24h intervals till 96h time point.

Proximity Ligation Assay

Cells were fixed with 4% formaldehyde in PBS for 15 min at RT. After blocking with 5% normal

goat serum in PBS with 0.3% Triton X-100, primary antibodies 1:100 diluted in blocking buffer

were incubated with fixed cells overnight at 4°C. Following washing with PLA wash Buffer A,

samples were incubated with 1:5 diluted Duolink® in Situ PLA Probe anti-rabbit Plus

(cat#DUO92002, Sigma-Aldrich) and anti-mouse Minus (cat#DUO92004, Sigma-Aldrich) at

37°C for 1 hr. Subsequently, ligation (30 minutes, 37°C), rolling circle amplification, and detection

(120 minutes, 37°C) were performed using the Duolink® In Situ Detection Reagents Red

(cat#DUO92008, Sigma-Aldrich). Nuclei were counterstained with Hoechst 33342. PLA

signals were documented by a Zeiss LSM780 confocal microscope (Carl Zeiss). For PLA flow

cytometry, an equal number of cells were collected in microcentrifuge tubes by trypsinization and

fixed with 4% formaldehyde in PBS for 10 min at RT. Subsequent PLA steps were followed as

described earlier. The PLA signals were quantified by a FACSCalibur flow cytometer (BD

Biosciences).

Mouse anti-JNK1 (JM2671) and rabbit anti-HSF1 (H-311) antibodies were used for JNK-HSF1

PLA. Rabbit anti-JNK1/3 (C-17) and mouse anti-RAPTOR (1H6.2) antibodies were used for

JNK-RAPTOR PLA. Rabbit anti-mTOR (7C10) and mouse anti-RAPTOR (1H6.2) antibodies

were used for mTOR-RAPTOR PLA. Mouse anti-HSF1 (E-4) and rabbit anti-AO (A11)

antibodies were used for HSF1-AO PLA. Rabbit anti-AO (A11) and mouse anti-HSP60 (LK1)

antibodies were used for AO-HSP60 PLA.

Measurement of global protein translation

Live cells were incubated with 100 nM 6-FAM-dC-puromycin in complete culture medium

for 30 min and analyzed by a FACSCalibur flow cytometer (BD Biosciences).

Immunoblotting and Immunoprecipitation

Whole-cell protein extracts were prepared in cold cell-lysis buffer (100 mM NaCl, 30 mM Tris-

HCl pH 7.6, 1% Triton X-100, 20 mM sodium fluoride, 1mM EDTA, 1mM sodium

orthovanadate, and 1x Halt™ protease inhibitor cocktail). Proteins were transferred to

nitrocellulose membranes. Following incubation with the blocking buffer (5% non-fat milk in 1x

TBS-T) for 1 hour at RT, membranes were incubated with primary antibodies (1:1,000 dilution in

the blocking buffer) overnight at 4 °C. After washing with 1xTBS-T for 3 times, membranes

were incubated with peroxidase-conjugated secondary antibodies (1: 3,000 diluted in the

blocking buffer) at RT for 1 hr. Signals were detected using SuperSignal™ West

chemiluminescent substrates. For normal Co-IP, proteins were extracted using a QSonica Q125

sonicator (total process time: 15S, pulse-on time: 5S, pulse-off time: 10S, output intensity: 30%)

in 1x sonication buffer (20 mM Tris, 20 mM NaCl, 1 mM EDTA, 20 mM β-glycerol-phosphate,

20 mM sodium fluoride, 4 mM sodium orthovanadate, and 1 mM DTT pH7.4, supplemented

with Halt protease inhibitor cocktail). Total 1 mg whole cell lysates were incubated with A11

antibodies at 4 °C overnight. Normal rabbit IgG were used as the negative controls. The primary

antibody was precipitated using Protein G magnetic beads. After washing with the lysis buffer

for 3 times, beads were boiled in 1x loading buffer for 5 min before loading on SDS-PAGE.

EasyBlocker (Cat#GTX425858, GeneTex) was used for blocking and anti-HSF1 (H-311)

primary antibody (cat#sc-9144, Santa Cruz Biotechnology) incubation (1:1,000, O/N at 4°C).

The blot was incubated with EasyBlot anti-Rabbit IgG HRP conjugated antibody

(cat#GTX221666-01, GeneTex) for 1 hour. Chemiluminescent signals were captured by an

iBright™ FL1000 imaging system (ThermoFisher Scientific).

Live-Or-Dye™ Dead Cell Staining

Cells were cultured in DMEM for 48 hours before being treated with inhibitors for the required

periods and collected by trypsinization. $1x10^6$ cells/ml cells were stained with Live-or-Dye™

fixable dead cell dyes (1:1,000 dilution, cat#32016, Biotium) at RT for 30 min, protected from

light. Stained cells were washed once with PBS and analyzed by flow cytometry. For co-staining

with cleaved caspase 3 antibody, stained cells were fixed with 4% formaldehyde (10 min, RT)

and permeabilized with 0.3% Triton-X-100 in PBS (10 min, RT). The cells were stained by

Cleaved Caspase 3 antibody (Asp175) (5A1E) (1:1,000, cat#9664, Cell Signaling Technology)

followed by goat anti-Rabbit IgG Alexa Fluor™ 488 conjugated (1:1,000, cat#A-11034,

ThermoFisher Scientific) secondary antibodies. The samples were analyzed by flow cytometry.

Inhibitor treatments

Treatments with DTHIB, Q-VD-OPh, Liproxstatin-1, Necrostatin-1, and Wortmannin were

done for 72 hours. For Congo Red and LY-2584702 treatments, cells were pretreated for 24h

followed by DTHIB for an additional 72h. For treatments with NV-5138, cells were pretreated

for 48 hours in a leucine-free DMEM (cat#226-024, Crystalgen) before adding DTHIB for an

additional 72 hours.

Quantification And Statistical Analysis

Statistical analyses were performed using Prism GraphPad 10.0 (GraphPad Software). The

detailed statistical methods and sample sizes are provided in the figure legends. All results are

expressed as mean±SD, mean±SEM, or median and IQR. The statistical significance is defined

as: *p<0.05, **p<0.01; ***p<0.001; ****p<0.0001; n.s.: not significant. For *in vitro*

experiments, sample size required was not determined a priori. The experiments were not

randomized**.**

REFERENCES

1. Dai C, Dai S, Cao J. Proteotoxic stress of cancer: implication of the heat-shock response in oncogenesis. J Cell Physiol. 2012;227(8):2982-7. doi: 10.1002/jcp.24017. PubMed PMID: 22105155; PMCID: PMC3431153.

2. Dai C. The heat-shock, or HSF1-mediated proteotoxic stress, response in cancer: from proteomic stability to oncogenesis. Philos Trans R Soc Lond B Biol Sci. 2018;373(1738). doi: 10.1098/rstb.2016.0525. PubMed PMID: 29203710; PMCID: PMC5717525.

3. Guang MHZ, Kavanagh EL, Dunne LP, Dowling P, Zhang L, Lindsay S, Bazou D, Goh CY, Hanley C, Bianchi G, Anderson KC, O'Gorman P, McCann A. Targeting Proteotoxic Stress in Cancer: A Review of the Role that Protein Quality Control Pathways Play in Oncogenesis. Cancers (Basel). 2019;11(1). Epub 20190109. doi: 10.3390/cancers11010066. PubMed PMID: 30634515; PMCID: PMC6356294.

4. Wilson DM, 3rd, Cookson MR, Van Den Bosch L, Zetterberg H, Holtzman DM, Dewachter I. Hallmarks of neurodegenerative diseases. Cell. 2023;186(4):693-714. doi: 10.1016/j.cell.2022.12.032. PubMed PMID: 36803602.

5. Tang Z, Dai S, He Y, Doty RA, Shultz LD, Sampson SB, Dai C. MEK guards proteome stability and inhibits tumor-suppressive amyloidogenesis via HSF1. Cell. 2015;160(4):729-44. doi: 10.1016/j.cell.2015.01.028. PubMed PMID: 25679764; PMCID: PMC4564124.

6. Li J, Labbadia J, Morimoto RI. Rethinking HSF1 in Stress, Development, and Organismal Health. Trends Cell Biol. 2017;27(12):895-905. Epub 20170907. doi: 10.1016/j.tcb.2017.08.002. PubMed PMID: 28890254; PMCID: PMC5696061.

7. Anckar J, Sistonen L. Regulation of HSF1 function in the heat stress response: implications in aging and disease. Annu Rev Biochem. 2011;80:1089-115. doi: 10.1146/annurev-

biochem-060809-095203. PubMed PMID: 21417720.

8. Gomez-Pastor R, Burchfiel ET, Thiele DJ. Regulation of heat shock transcription factors and

their roles in physiology and disease. Nat Rev Mol Cell Biol. 2018;19(1):4-19. Epub

20170830. doi: 10.1038/nrm.2017.73. PubMed PMID: 28852220; PMCID: PMC5794010.

9. Min JN, Huang L, Zimonjic DB, Moskophidis D, Mivechi NF. Selective suppression of

lymphomas by functional loss of Hsf1 in a p53-deficient mouse model for spontaneous

tumors. Oncogene. 2007;26(35):5086-97. Epub 20070219. doi: 10.1038/sj.onc.1210317.

PubMed PMID: 17310987.

10. Dai C, Whitesell L, Rogers AB, Lindquist S. Heat shock factor 1 is a powerful multifaceted

modifier of carcinogenesis. Cell. 2007;130(6):1005-18. doi: 10.1016/j.cell.2007.07.020.

PubMed PMID: 17889646; PMCID: PMC2586609.

11. Meng L, Gabai VL, Sherman MY. Heat-shock transcription factor HSF1 has a critical role in

human epidermal growth factor receptor-2-induced cellular transformation and

tumorigenesis. Oncogene. 2010;29(37):5204-13. Epub 20100712. doi:

10.1038/onc.2010.277. PubMed PMID: 20622894; PMCID: PMC2940982.

12. Jin X, Moskophidis D, Mivechi NF. Heat shock transcription factor 1 is a key determinant of

HCC development by regulating hepatic steatosis and metabolic syndrome. Cell Metab.

2011;14(1):91-103. doi: 10.1016/j.cmet.2011.03.025. PubMed PMID: 21723507; PMCID:

PMC3214631.

13. Dai C, Santagata S, Tang Z, Shi J, Cao J, Kwon H, Bronson RT, Whitesell L, Lindquist S.

Loss of tumor suppressor NF1 activates HSF1 to promote carcinogenesis. J Clin Invest.

2012;122(10):3742-54. Epub 20120904. doi: 10.1172/JCI62727. PubMed PMID: 22945628;

PMCID: PMC3461912.

14. Kourtis N, Lazaris C, Hockemeyer K, Balandran JC, Jimenez AR, Mullenders J, Gong Y,

Trimarchi T, Bhatt K, Hu H, Shrestha L, Ambesi-Impiombato A, Kelliher M, Paietta E,

Chiosis G, Guzman ML, Ferrando AA, Tsirigos A, Aifantis I. Oncogenic hijacking of the

stress response machinery in T cell acute lymphoblastic leukemia. Nat Med.

2018;24(8):1157-66. Epub 20180723. doi: 10.1038/s41591-018-0105-8. PubMed PMID:

30038221; PMCID: PMC6082694.

15. Fujimoto M, Takii R, Matsumoto M, Okada M, Nakayama KI, Nakato R, Fujiki K, Shirahige

K, Nakai A. HSF1 phosphorylation establishes an active chromatin state via the TRRAP-

TIP60 complex and promotes tumorigenesis. Nat Commun. 2022;13(1):4355. Epub

20220729. doi: 10.1038/s41467-022-32034-4. PubMed PMID: 35906200; PMCID:

PMC9338313.

16. Heimberger T, Andrulis M, Riedel S, Stuhmer T, Schraud H, Beilhack A, Bumm T, Bogen

B, Einsele H, Bargou RC, Chatterjee M. The heat shock transcription factor 1 as a potential

new therapeutic target in multiple myeloma. Br J Haematol. 2013;160(4):465-76. Epub

20121218. doi: 10.1111/bjh.12164. PubMed PMID: 23252346.

17. Chuma M, Sakamoto N, Nakai A, Hige S, Nakanishi M, Natsuizaka M, Suda G, Sho T,

Hatanaka K, Matsuno Y, Yokoo H, Kamiyama T, Taketomi A, Fujii G, Tashiro K, Hikiba Y,

Fujimoto M, Asaka M, Maeda S. Heat shock factor 1 accelerates hepatocellular carcinoma

development by activating nuclear factor-kappaB/mitogen-activated protein kinase.

Carcinogenesis. 2014;35(2):272-81. Epub 20131015. doi: 10.1093/carcin/bgt343. PubMed

PMID: 24130164.

18. Fujimoto M, Takaki E, Takii R, Tan K, Prakasam R, Hayashida N, Iemura S, Natsume T,

Nakai A. RPA assists HSF1 access to nucleosomal DNA by recruiting histone chaperone

FACT. Mol Cell. 2012;48(2):182-94. Epub 20120830. doi: 10.1016/j.molcel.2012.07.026.

PubMed PMID: 22940245.

19. Solimini NL, Luo J, Elledge SJ. Non-oncogene addiction and the stress phenotype of cancer

cells. Cell. 2007;130(6):986-8. doi: 10.1016/j.cell.2007.09.007. PubMed PMID: 17889643.

20. Gutmann DH, Ferner RE, Listernick RH, Korf BR, Wolters PL, Johnson KJ.

Neurofibromatosis type 1. Nat Rev Dis Primers. 2017;3:17004. Epub 20170223. doi:

10.1038/nrdp.2017.4. PubMed PMID: 28230061.

21. Hirbe AC, Gutmann DH. Neurofibromatosis type 1: a multidisciplinary approach to care.

Lancet Neurol. 2014;13(8):834-43. doi: 10.1016/S1474-4422(14)70063-8. PubMed PMID:

25030515.

22. Cichowski K, Jacks T. NF1 tumor suppressor gene function: narrowing the GAP. Cell.

2001;104(4):593-604. doi: 10.1016/s0092-8674(01)00245-8. PubMed PMID: 11239415.

23. Grice GL, Nathan JA. The recognition of ubiquitinated proteins by the proteasome. Cell Mol

Life Sci. 2016;73(18):3497-506. Epub 20160502. doi: 10.1007/s00018-016-2255-5. PubMed

PMID: 27137187; PMCID: PMC4980412.

24. Knowles TP, Vendruscolo M, Dobson CM. The amyloid state and its association with protein

misfolding diseases. Nat Rev Mol Cell Biol. 2014;15(6):384-96. doi: 10.1038/nrm3810.

PubMed PMID: 24854788.

25. Yakupova EI, Bobyleva LG, Vikhlyantsev IM, Bobylev AG. Congo Red and amyloids:

history and relationship. Biosci Rep. 2019;39(1). Epub 20190115. doi:

10.1042/BSR20181415. PubMed PMID: 30567726; PMCID: PMC6331669.

26. Kayed R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, Glabe CG.

Common structure of soluble amyloid oligomers implies common mechanism of

pathogenesis. Science. 2003;300(5618):486-9. doi: 10.1126/science.1079469. PubMed

PMID: 12702875.

27. Kayed R, Head E, Sarsoza F, Saing T, Cotman CW, Necula M, Margol L, Wu J, Breydo L,

Thompson JL, Rasool S, Gurlo T, Butler P, Glabe CG. Fibril specific, conformation

dependent antibodies recognize a generic epitope common to amyloid fibrils and fibrillar

oligomers that is absent in prefibrillar oligomers. Mol Neurodegener. 2007;2:18. Epub

20070926. doi: 10.1186/1750-1326-2-18. PubMed PMID: 17897471; PMCID:

PMC2100048.

28. Heiser V, Scherzinger E, Boeddrich A, Nordhoff E, Lurz R, Schugardt N, Lehrach H,

Wanker EE. Inhibition of huntingtin fibrillogenesis by specific antibodies and small

molecules: implications for Huntington's disease therapy. Proc Natl Acad Sci U S A.

2000;97(12):6739-44. doi: 10.1073/pnas.110138997. PubMed PMID: 10829068; PMCID:

PMC18723.

29. Sanchez I, Mahlke C, Yuan J. Pivotal role of oligomerization in expanded polyglutamine

neurodegenerative disorders. Nature. 2003;421(6921):373-9. doi: 10.1038/nature01301.

PubMed PMID: 12540902.

30. Tang Z, Su KH, Xu M, Dai C. HSF1 physically neutralizes amyloid oligomers to empower

overgrowth and bestow neuroprotection. Sci Adv. 2020;6(46). Epub 20201111. doi:

10.1126/sciadv.abc6871. PubMed PMID: 33177089; PMCID: PMC7673739.

31. Christensen JH, Nielsen MN, Hansen J, Fuchtbauer A, Fuchtbauer EM, West M, Corydon

TJ, Gregersen N, Bross P. Inactivation of the hereditary spastic paraplegia-associated Hspd1

gene encoding the Hsp60 chaperone results in early embryonic lethality in mice. Cell Stress

Chaperones. 2010;15(6):851-63. doi: 10.1007/s12192-010-0194-x. PubMed PMID:

20393889; PMCID: PMC3024079.

32. Fan F, Duan Y, Yang F, Trexler C, Wang H, Huang L, Li Y, Tang H, Wang G, Fang X, Liu

J, Jia N, Chen J, Ouyang K. Deletion of heat shock protein 60 in adult mouse cardiomyocytes

perturbs mitochondrial protein homeostasis and causes heart failure. Cell Death Differ.

2020;27(2):587-600. Epub 20190617. doi: 10.1038/s41418-019-0374-x. PubMed PMID:

31209364; PMCID: PMC7205885.

33. Su KH, Cao J, Tang Z, Dai S, He Y, Sampson SB, Benjamin IJ, Dai C. HSF1 critically

attunes proteotoxic stress sensing by mTORC1 to combat stress and promote growth. Nat

Cell Biol. 2016;18(5):527-39. Epub 20160404. doi: 10.1038/ncb3335. PubMed PMID:

27043084; PMCID: PMC5341796.

34. Johnson GL, Nakamura K. The c-jun kinase/stress-activated pathway: regulation, function

and role in human disease. Biochim Biophys Acta. 2007;1773(8):1341-8. Epub 20070104.

doi: 10.1016/j.bbamcr.2006.12.009. PubMed PMID: 17306896; PMCID: PMC1995559.

35. Battaglioni S, Benjamin D, Walchli M, Maier T, Hall MN. mTOR substrate phosphorylation

in growth control. Cell. 2022;185(11):1814-36. Epub 20220516. doi:

10.1016/j.cell.2022.04.013. PubMed PMID: 35580586.

36. Kim DH, Sarbassov DD, Ali SM, King JE, Latek RR, Erdjument-Bromage H, Tempst P,

Sabatini DM. mTOR interacts with raptor to form a nutrient-sensitive complex that signals to

the cell growth machinery. Cell. 2002;110(2):163-75. doi: 10.1016/s0092-8674(02)00808-5.

PubMed PMID: 12150925.

37. Schmidt EK, Clavarino G, Ceppi M, Pierre P. SUnSET, a nonradioactive method to monitor

protein synthesis. Nat Methods. 2009;6(4):275-7. Epub 20090322. doi: 10.1038/nmeth.1314.

PubMed PMID: 19305406.

38. Zhang T, Inesta-Vaquera F, Niepel M, Zhang J, Ficarro SB, Machleidt T, Xie T, Marto JA,

Kim N, Sim T, Laughlin JD, Park H, LoGrasso PV, Patricelli M, Nomanbhoy TK, Sorger

PK, Alessi DR, Gray NS. Discovery of potent and selective covalent inhibitors of JNK.

Chem Biol. 2012;19(1):140-54. doi: 10.1016/j.chembiol.2011.11.010. PubMed PMID:

22284361; PMCID: PMC3270411.

39. Moerke NJ, Aktas H, Chen H, Cantel S, Reibarkh MY, Fahmy A, Gross JD, Degterev A,

Yuan J, Chorev M, Halperin JA, Wagner G. Small-molecule inhibition of the interaction

between the translation initiation factors eIF4E and eIF4G. Cell. 2007;128(2):257-67. doi:

10.1016/j.cell.2006.11.046. PubMed PMID: 17254965.

40. Tolcher A, Goldman J, Patnaik A, Papadopoulos KP, Westwood P, Kelly CS, Bumgardner

W, Sams L, Geeganage S, Wang T, Capen AR, Huang J, Joseph S, Miller J, Benhadji KA,

Brail LH, Rosen LS. A phase I trial of LY2584702 tosylate, a p70 S6 kinase inhibitor, in

patients with advanced solid tumours. Eur J Cancer. 2014;50(5):867-75. Epub 20140115.

doi: 10.1016/j.ejca.2013.11.039. PubMed PMID: 24440085.

41. Porter AG, Janicke RU. Emerging roles of caspase-3 in apoptosis. Cell Death Differ.

1999;6(2):99-104. doi: 10.1038/sj.cdd.4400476. PubMed PMID: 10200555.

42. Boulares AH, Yakovlev AG, Ivanova V, Stoica BA, Wang G, Iyer S, Smulson M. Role of

poly(ADP-ribose) polymerase (PARP) cleavage in apoptosis. Caspase 3-resistant PARP

mutant increases rates of apoptosis in transfected cells. J Biol Chem. 1999;274(33):22932-40.

doi: 10.1074/jbc.274.33.22932. PubMed PMID: 10438458.

43. Huang J, Manning BD. The TSC1-TSC2 complex: a molecular switchboard controlling cell

growth. Biochem J. 2008;412(2):179-90. doi: 10.1042/BJ20080281. PubMed PMID:

18466115; PMCID: PMC2735030.

44. Dong B, Jaeger AM, Hughes PF, Loiselle DR, Hauck JS, Fu Y, Haystead TA, Huang J,

Thiele DJ. Targeting therapy-resistant prostate cancer via a direct inhibitor of the human heat

shock transcription factor 1. Sci Transl Med. 2020;12(574). doi:

10.1126/scitranslmed.abb5647. PubMed PMID: 33328331; PMCID: PMC10571035.

45. Sengupta S, Giaime E, Narayan S, Hahm S, Howell J, O'Neill D, Vlasuk GP, Saiah E.

Discovery of NV-5138, the first selective Brain mTORC1 activator. Sci Rep.

2019;9(1):4107. Epub 20190311. doi: 10.1038/s41598-019-40693-5. PubMed PMID:

30858438; PMCID: PMC6412019.

46. Kato T, Pothula S, Liu RJ, Duman CH, Terwilliger R, Vlasuk GP, Saiah E, Hahm S, Duman

RS. Sestrin modulator NV-5138 produces rapid antidepressant effects via direct mTORC1

activation. J Clin Invest. 2019;129(6):2542-54. Epub 20190416. doi: 10.1172/JCI126859.

PubMed PMID: 30990795; PMCID: PMC6546461.

47. Wolfson RL, Chantranupong L, Saxton RA, Shen K, Scaria SM, Cantor JR, Sabatini DM.

Sestrin2 is a leucine sensor for the mTORC1 pathway. Science. 2016;351(6268):43-8. Epub

20151008. doi: 10.1126/science.aab2674. PubMed PMID: 26449471; PMCID:

PMC4698017.

48. Caserta TM, Smith AN, Gultice AD, Reedy MA, Brown TL. Q-VD-OPh, a broad spectrum

caspase inhibitor with potent antiapoptotic properties. Apoptosis. 2003;8(4):345-52. doi:

10.1023/a:1024116916932. PubMed PMID: 12815277.

49. Chauvier D, Ankri S, Charriaut-Marlangue C, Casimir R, Jacotot E. Broad-spectrum caspase

inhibitors: from myth to reality? Cell Death Differ. 2007;14(2):387-91. Epub 20060929. doi:

10.1038/sj.cdd.4402044. PubMed PMID: 17008913.

50. Miao EA, Rajan JV, Aderem A. Caspase-1-induced pyroptotic cell death. Immunol Rev.

2011;243(1):206-14. doi: 10.1111/j.1600-065X.2011.01044.x. PubMed PMID: 21884178;

PMCID: PMC3609431.

51. Shan B, Pan H, Najafov A, Yuan J. Necroptosis in development and diseases. Genes Dev.

2018;32(5-6):327-40. doi: 10.1101/gad.312561.118. PubMed PMID: 29593066; PMCID:

PMC5900707.

52. Chen X, Comish PB, Tang D, Kang R. Characteristics and Biomarkers of Ferroptosis. Front

Cell Dev Biol. 2021;9:637162. Epub 20210121. doi: 10.3389/fcell.2021.637162. PubMed

PMID: 33553189; PMCID: PMC7859349.

53. Stockwell BR. Ferroptosis turns 10: Emerging mechanisms, physiological functions, and

therapeutic applications. Cell. 2022;185(14):2401-21. doi: 10.1016/j.cell.2022.06.003.

PubMed PMID: 35803244; PMCID: PMC9273022.

54. Yoshii SR, Mizushima N. Monitoring and Measuring Autophagy. Int J Mol Sci. 2017;18(9).

Epub 20170828. doi: 10.3390/ijms18091865. PubMed PMID: 28846632; PMCID:

PMC5618514.

55. Blommaart EF, Krause U, Schellens JP, Vreeling-Sindelarova H, Meijer AJ. The

phosphatidylinositol 3-kinase inhibitors wortmannin and LY294002 inhibit autophagy in

isolated rat hepatocytes. Eur J Biochem. 1997;243(1-2):240-6. doi: 10.1111/j.1432-

1033.1997.0240a.x. PubMed PMID: 9030745.

56. Schubert U, Anton LC, Gibbs J, Norbury CC, Yewdell JW, Bennink JR. Rapid degradation

of a large fraction of newly synthesized proteins by proteasomes. Nature.

2000;404(6779):770-4. doi: 10.1038/35008096. PubMed PMID: 10783891.

57. Kim LC, Cook RS, Chen J. mTORC1 and mTORC2 in cancer and the tumor

microenvironment. Oncogene. 2017;36(16):2191-201. Epub 20161017. doi:

10.1038/onc.2016.363. PubMed PMID: 27748764; PMCID: PMC5393956.

58. Zou Z, Tao T, Li H, Zhu X. mTOR signaling pathway and mTOR inhibitors in cancer:

progress and challenges. Cell Biosci. 2020;10:31. Epub 20200310. doi: 10.1186/s13578-020-

00396-1. PubMed PMID: 32175074; PMCID: PMC7063815.

59. Goloudina AR, Demidov ON, Garrido C. Inhibition of HSP70: a challenging anti-cancer

strategy. Cancer Lett. 2012;325(2):117-24. Epub 20120628. doi:

10.1016/j.canlet.2012.06.003. PubMed PMID: 22750096.

60. Jhaveri K, Taldone T, Modi S, Chiosis G. Advances in the clinical development of heat

shock protein 90 (Hsp90) inhibitors in cancers. Biochim Biophys Acta. 2012;1823(3):742-

55. Epub 20111029. doi: 10.1016/j.bbamcr.2011.10.008. PubMed PMID: 22062686;

PMCID: PMC3288123.

61. Pasqua AE, Sharp SY, Chessum NEA, Hayes A, Pellegrino L, Tucker MJ, Miah A, Wilding

B, Evans LE, Rye CS, Mok NY, Liu M, Henley AT, Gowan S, De Billy E, Te Poele R,

Powers M, Eccles SA, Clarke PA, Raynaud FI, Workman P, Jones K, Cheeseman MD. HSF1

Pathway Inhibitor Clinical Candidate (CCT361814/NXP800) Developed from a Phenotypic

Screen as a Potential Treatment for Refractory Ovarian Cancer and Other Malignancies. J

Med Chem. 2023;66(8):5907-36. Epub 20230405. doi: 10.1021/acs.jmedchem.3c00156.

PubMed PMID: 37017629; PMCID: PMC10150365.

62. Ali ES, Mitra K, Akter S, Ramproshad S, Mondal B, Khan IN, Islam MT, Sharifi-Rad J,

Calina D, Cho WC. Recent advances and limitations of mTOR inhibitors in the treatment of

cancer. Cancer Cell Int. 2022;22(1):284. Epub 20220915. doi: 10.1186/s12935-022-02706-8.

PubMed PMID: 36109789; PMCID: PMC9476305.

63. Wood KC, Gutkind JS. Challenges and Emerging Opportunities for Targeting mTOR in

Cancer. Cancer Res. 2022;82(21):3884-7. doi: 10.1158/0008-5472.CAN-22-0602. PubMed

PMID: 36321262; PMCID: PMC9634680.

64. Thomson AW, Turnquist HR, Raimondi G. Immunoregulatory functions of mTOR

inhibition. Nat Rev Immunol. 2009;9(5):324-37. doi: 10.1038/nri2546. PubMed PMID:

19390566; PMCID: PMC2847476.

Figure 1

FIGURE LEGENDS:

Figure 1: HSF1 is required to suppress proteomic instability and amyloidogenesis in

MPNST cells. (A) Violin plots of *HSF1* expression in human cancer tissues showing *NF1*

deletion (median and IQR, n=6424, 1598, or 88, Kruskal-Wallis test). Data are generated by the

TGCA Research Network (https://www.cancer.gov/tcga). (B) Kaplan-Meier survival curves of

patients whose tumors harbor *NF1* deletion (n=420 or 419, Log-rank test). Data are generated by

the TGCA Research Network (https://www.cancer.gov/tcga). (C) and (D) The impacts of *HSF1*

depletion on protein polyubiquitination in both 90-8TL cells and primary human Schwann cells.

Cells were transduced with lentiviral shRNAs for 4 days and global protein Lys48-specific

ubiquitination was detected by immunoblotting in both detergent-soluble and -insoluble fractions

of whole cell lysates (C). Protein ubiquitination was quantitated using Fiji imaging software (D)

(mean ± SD, n=3 independent experiments, Two-way ANOVA). (E) and (F) Congo Red (CR)

staining of 90-8TL cells and primary human Schwann cells with and without *HSF1* depletion,

analyzed by flow cytometry (E). Quantitation of CR staining using the median fluorescence

intensity (F) (mean ± SD, n=3 independent experiments, One-way ANOVA). (G)-(J)

Quantitation of soluble amyloid oligomers and insoluble amyloid fibrils by ELISA using A11

and OC antibodies, respectively (mean ± SD, n=3 independent experiments, One-way ANOVA).

Figure 2

Figure 2: HSF1 counters the tumor-suppressive effects of amyloids in MPNST cells.

(A) and (B) The growth curves of 90-8TL and primary human Schwann cells with and without

HSF1 depletion and 20 μM CR treatment (mean ± SD, n=3 independent experiments, Two-way

ANOVA). (C) Co-IP of amyloid oligomers (AOs) with HSF1 in 90-8TL cells. (D)-(F) Detection

of AO-HSF1 interactions by PLA flow cytometry (D). Quantitation of AO-HSF1 PLA using the

median fluorescence intensity (F) (mean ± SD, n=3 or 4 independent experiments, Student's t

test). (G) and (H) Detection of AO-HSP60 interactions by PLA flow cytometry in S462 cells

following *HSF1* depletion (G). Quantitation of AO-HSP60 PLA using the median fluorescence

intensity (F) (mean ± SD, n=3 independent experiments, Student's t test). (I) Schematic depiction

of AO neuralization by HSF1 and HSP60 attack by AO following *HSF1* depletion in

mitochondria.

Figure 3

Figure 3: JNK senses *HSF1* deficiency to repress mTORC1 and protein translation in

MPNST cells. (A) Detection of JNK activation and mTORC1 inhibition following *HSF1*

depletion in S462 and primary human Schwann cells by immunoblotting. (B) Quantitation of (A)

using Fiji imaging software (mean ± SD, n=3 independent experiments, One-way ANOVA). (C)

Detection of HSF1-JNK, JNK-RAPTOR, and mTOR-RAPTOR interactions by PLA in S462

cells with and without *HSF1* depletion. (D) and (E) Detection and quantitation of JNK-RAPTOR

and mTOR-RAPTOR interactions by PLA flow cytometry in 90-8TL cells (mean ± SD, n=3

independent experiments, One-way ANOVA). (F) Measurement of protein translation rates in

90-8TL and primary human Schwann cells with and without *HSF1* depletion by puromycin

labeling, analyzed by flow cytometry. Cells were treated with 1 μM JNK-IN-8 for 24 hrs before

labeling with 100 nM 6-FAM-dc-puromycin for 30 min (mean ± SD, n=3 independent

experiments, Two-way ANOVA). (G) Heatmap of protein translation rates measured by

puromycin labeling in diverse human cancer cell lines with and without *HSF1* depletion. (H)

Schematic depiction of regulation of protein translation by HSF1.

Figure 4

Figure 4: Protein translation is causally related to amyloidogenesis in MPNST cells.

(A) and (B) Detection of mTORC1 stimulation following JNK inhibition in 90-8TL and immortalized human Schwann cells by immunoblotting. Cells were treated with JNK-IN-8 for 2 days. (C) and (D) Quantitation of soluble AOs and insoluble AFs by ELISA in 90-8TL and human Schwann cells with and without 3 µM JNK-IN-8 treatment for 3 days (mean ± SD, n=3 independent experiments, Student's t test). (E) Detection of LY2584702-mediated S6K inhibition in 90-8TL cells by immunoblotting. Cells were treated with 20 µM LY2584702 for 2 days. (F) Quantitation of soluble AOs and insoluble AFs by ELISA in 90-8TL cells with and without 20 µM LY2584702 treatment for 2 days (mean ± SD, n=3 independent experiments, Student's t test). (G) CR staining of 90-8TL and immortalized human Schwann cells with and without JNK-IN-8 and 4EGI-1 treatment, detected by flow cytometry. (H) Quantitation of (G) using the median fluorescence intensity (mean ± SD, n=3 independent experiments, One-way ANOVA). (I) Quantitation of amyloids by CR staining in 90-8TL cells with and without *HSF1* depletion and 3 µM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (J) Quantitation of caspase 3 cleavage by ELISA in 90-8TL cells with and without *HSF1* depletion and 3 µM JNK-IN-8 treatment (mean ± SD, n=2 independent experiments, Two-way ANOVA). (K) Quantitation of amyloids by CR staining in immortalized human Schwann cells with and without *HSF1* depletion and 3 µM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (L) Quantitation of caspase 3 cleavage by ELISA in immortalized human Schwann cells with and without *HSF1* depletion and 3 µM JNK-IN-8 treatment (mean ± SD, n=2 independent experiments, Two-way ANOVA). (M) Schematic depiction of the adaptation of MPNST cells to *HSF1* depletion by activating JNK to suppress mTORC1 and protein translation, thereby alleviating amyloidogenesis and cytotoxicity.

Figure 5

Figure 5: mTORC1 stimulation aggravates the amyloidogenesis and cytotoxicity elicited by

HSF1 inhibition. (A) Immunoblotting detection of JNK activation, mTORC1 activity, and

protein polyubiquitination in S462 cells following stable *TSC2* knockdown and/or HSF1

inhibition by 10 μM DTHIB for 3 days. (B) Quantitation of soluble AOs by flow cytometry

using A11 antibody staining in S462 cells with and without *TSC2* knockdown and 10 μM

DTHIB treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (C)

Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

without stable *TSC2* knockdown and DTHIB (10 μM) or combined DTHIB and LY2584702 (20

μM) treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). Cells were pre-

treated with LY2584702 for 1 day followed by DTHIB treatment for another 3 days. (D)

Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

without stable *TSC2* knockdown and DTHIB (10 μM) or combined DTHIB and CR (30 μM)

treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). Cells were pre-

treated with CR for 1 day followed by DTHIB treatment for another 3 days. (E) Immunoblotting

detection of mTORC1 stimulation by 500 μM NV-5138 in S462 cells with and without 10 μM

DTHIB treatment. (F) Quantitation of soluble AOs by flow cytometry using A11 antibody

staining in S462 cells with and without 500 μM NV-5138 stimulation and 10 μM DTHIB

treatment (mean ± SD, n=3 independent experiments, One-way ANOVA). (G) Quantitation of

cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and without 500 μM

NV-5138 stimulation and 10 μM DTHIB or combined DTHIB and CR treatment (mean ± SD,

n=5 independent experiments, One-way ANOVA). (H) Quantitation of cytotoxicity by flow

cytometry using Live-or-Dye stains in immortalized human Schwann cells with and without 500

μM NV-5138 stimulation and 10 μM DTHIB treatment (mean ± SD, n=3 independent

experiments, One-way ANOVA). (I) Immunoblotting detection of mTORC1 stimulation by 500

µM NV-5138 in immortalized human Schwann cells with and without 10 µM DTHIB treatment.

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

Figure 6

Figure 6: Concurrent HSF1 inhibition and mTORC1 stimulation instigate widespread non-

apoptotic cell death. (A) and (B) Quantitation of cytotoxicity by flow cytometry using Live-or-

Dye stains combined with cleaved caspase 3 (Asp175) antibody staining. S462 cells with and

without stable *TSC2* knockdown were treated with DMSO, 10 μM DTHIB, or combined DTHIB

and 30 μM Q-VD-OPh (mean ± SD, n=3 independent experiments, Two-way ANOVA). (C)

Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

without stable *TSC2* knockdown. Cells were treated with 10 μM DTHIB alone or co-treated with

10 μM DTHIB and 30 μM Necrostatin-1 or 20 μM Liproxstatin-1 for 3 days (mean ± SD, n=3

independent experiments, Two-way ANOVA). (D) Immunoblotting detection of ferroptosis

markers in S462 cells with and without stable *TSC2* knockdown treated with 10 μM DTHIB

alone or co-treated with 10 μM DTHIB and 20 μM liproxstatin-1 for 3 days. Erastin was

included as a positive control to induce canonical ferroptosis. (E) Immunoblotting detection of

autophagy markers in S462 cells with stable *TSC2* knockdown treated with 10 μM DTHIB alone

or co-treated with 10 μM DTHIB and 3 μM wortmannin for 3 days. Rapamycin was included as

a positive control to induce autophagy. (F) Quantitation of cytotoxicity by flow cytometry using

Live-or-Dye stains in S462 cells with and without stable *TSC2* knockdown. Cells were treated

with 10 μM DTHIB alone or co-treated with 10 μM DTHIB and 3 μM wortmannin for 3 days

(mean ± SD, n=3 independent experiments, Two-way ANOVA). (G) Schematic depiction of

instigation of cell death by severe proteomic imbalance, owing to simultaneous mTORC1

stimulation (❶) and HSF1 inhibition (❷). In cancer cells, constitutive HSF1 activation

provides extra chaperoning capacity to cope with elevated protein misfolding, partly due to

enhanced protein synthesis and widespread genetic mutations. Nevertheless, amyloids still

emerge, although at low levels. Importantly, HSF1 can neutralize highly toxic amyloid

oligomers, averting lethal consequences. By contrast, HSF1 inhibition diminishes chaperoning

capacity, insufficient to counterbalance the robust protein translation. mTORC1 stimulation

further aggravates this proteomic imbalance, which, in turn, strongly promotes amyloidogenesis.

In consequence, the amounts of amyloid oligomers exceed the neutralizing capacity of HSF1,

leading to cell death; nonetheless, it remains unclear how this non-apoptotic, non-autophagic

death occurs. (H) Schematic depiction of the concept of driving proteomic imbalance to combat

malignancy. On the one hand, in cancer cells, mTORC1 is inevitably activated to stimulate

protein translation, markedly augmenting protein quantity. On the other hand, the extra

chaperoning capacity governed by HSF1, albeit dispensable for normal life, becomes necessary

to ensure sufficient protein quality in cancer cells, thereby counterbalancing augmented protein

quantity and suppressing proteomic instability. Thus, proteomic balance promotes malignant

growth. By contrast, disrupting proteomic balance, through HSF1 inhibition, is sufficient to

provoke proteomic instability and elicit tumor suppression. However, simultaneous mTORC1

stimulation can remarkably drive proteomic imbalance, causing severe proteomic instability and

profound tumor suppression.

1089

1090

1091

1092

1093

1094

1095