Driving proteomic imbalance in malignancy provokes proteomic catastrophe

and confers tumor suppression

Babul Moni Ram[1], Omprakash Shriwas[1], Meng Xu[1], Kun-Han Chuang[1,2], and Chengkai Dai[1,*]

[1]Mouse Cancer Genetics Program, Center for Cancer Research, National Cancer Institute,

Frederick, MD 21702, USA

[2]Present address: School of Basic Medical Sciences, Fujian Medical University, Fuzhou, Fujian

350108, China

*Correspondence: Chengkai.dai@nih.gov

ABSTRACT

Unlike genomic instability, the implications of proteomic instability in cancer remain
ambiguous. By governing the proteotoxic stress response, heat shock factor 1 (HSF1) sustains
proteomic stability upon environmental insults. Apart from its importance to stress resistance and
survival, HSF1 is emerging as a powerful oncogenic enabler. In the Neurofibromatosis type I
(*NF1*)-deficient malignant peripheral nerve sheath tumor (MPNST) cells, *HSF1* depletion
triggered protein polyubiquitination, aggregation, and even tumor-suppressive amyloidogenesis.
In contrast, *HSF1* is dispensable for the proteome of non-transformed human Schwann cells.
Mechanistically, HSF1 defends the essential mitochondrial chaperone HSP60 against the direct
assault from soluble amyloid oligomers. To survive and adapt to compromised protein quality,
owing to *HSF1* deficiency, MPNST cells mobilized JNK to repress mTORC1 and protein
translation, thereby attenuating protein quantity to alleviate proteomic imbalance. mTORC1
stimulation, via either pharmacological JNK blockade, genetic *TSC2* depletion, or leucine
supplementation, markedly aggravated the proteomic imbalance elicited by *HSF1* deficiency.
This catastrophic imbalance instigated pronounced cell death partly through unchecked
amyloidogenesis, thereby exerting tumor suppression in both MPNST and melanoma models *in
vivo*. Thus, HSF1 safeguards the cancer proteome to enable the oncogenic potential of mTORC1.
This proof-of-principle study highlights provoking proteomic catastrophe as a next-generation
therapeutic concept for combating malignancy.

INTRODUCTION

Compared to genomic instability, proteomic instability in cancer is underappreciated. Malignant

transformation prompts proteotoxic stress and destabilizes the proteome.[1, 2] Several mechanisms,

including reactive oxygen species, genetic mutations, and uncontrolled protein translation, all

lead to increased protein damage, misfolding, as well as aggregation.[1-3] Of great interest,

amyloidogenesis, a hallmark of neurodegenerative disorders,[4] also occurs in cancer.[5] Unlike

neurons, cancerous cells manage to contain amyloidogenesis and counteract the cytotoxic effects

of amyloids. Among various cellular anti-amyloid mechanisms is heat shock factor 1 (HSF1) and

its mediated heat shock response or proteotoxic stress response (HSR/PSR), a powerful

cytoprotective transcriptional program.[6-8] In accordance with its role in repressing proteomic

instability, HSF1 has emerged as a potent oncogenic enabler, independently of driver

mutations.[9-15]

Sharply contrasting with its dispensability for primary and non-transformed cells, HSF1 becomes

essential to numerous cancer cell lines,[9,13,16-18] owing to their elevated intrinsic proteotoxic stress.

This phenomenon is referred to as "HSF1 or non-oncogene addiction of cancer",[2,19] justifying

HSF1 as a promising therapeutic target.

Neurofibromatosis type 1 (NF1) is a cancer predisposition syndrome afflicting approximately 1 in

3000 individuals worldwide.[20] About 30-50% of NF1 individuals develop benign plexiform

neurofibromas, which, unlike dermal neurofibromas, can transform into highly aggressive

malignant peripheral nerve sheath tumors (MPNSTs) with poor prognosis and survival.[21] The *NF1*

gene encodes neurofibromin, a 250-280 kDa RAS GTPase activating protein (GAP) that

negatively regulates RAS activity.[22] *NF1* inactivation in MPNST cells dysregulates RAS/MAPK

signaling which, in addition to stimulating cell growth, leads to constitutive HSF1 activation.[13]

Concomitantly, HSF1 overexpression and activation are seen in both *in vitro* and *in vivo* MPNST

models as well as primary NF1 tumors.[13] Importantly, *Hsf1* deficiency impaired the NF1-

associated tumorigenesis in mice in part by attenuating oncogenic RAS signaling.[13]

Utilizing the NF1 tumor model system, we herein demonstrate HSF1 as a key player in

countering proteomic instability in cancer. We discover that amyloidogenesis occurs in *NF1*-

deficient MPNST cells, partly due to hyperactive mTORC1 and augmented protein translation.

HSF1 inhibition destabilizes the cancer proteome, promoting global protein polyubiquitination,

aggregation, as well as amyloidogenesis. Importantly, this proteomic instability causes

cytotoxicity. To adapt and survive HSF1 inhibition, MPNST cells mobilize JNK to suppress

mTORC1 and thereby mitigate protein translation. Disrupting this adaptive response through

JNK inhibition aggravates amyloidogenesis and cytotoxicity. Similarly, mTORC1 stimulation,

via either genetic *TSC2* depletion or supplementation with leucine or its analog, synergizes with

HSF1 inhibition to inflict profound proteomic chaos and widespread non-apoptotic cell death. Of

note, blockade of amyloidogenesis markedly alleviates cell death. Thus, our findings illuminate a

prominent role of HSF1 in antagonizing proteomic instability in cancerous cells, constituting a

generic pro-oncogenic mechanism. This study exemplifies a proof of principle and, importantly,

institutes provoking proteomic catastrophe as a next-generation anti-cancer therapeutic concept.

RESULTS

HSF1 counters proteomic instability in *NF1*-deficient MPNST cells

Previously, we found that loss of *NF1* leads to constitutive activation of HSF1 and that *Hsf1*

deficiency impairs the tumorigenesis in *Nf1*$^{+/-}$; *p53*$^{+/-}$ (NPcis) mice.[13] In human cancers, *NF1*

deletion is associated with elevated *HSF1* gene expression (Figure 1A). Moreover, the expression

of *NF1* and *HSF1* genes are inversely correlated in human cancers (Figure S1A). Importantly,

among patients whose tumors harbor *NF1* deletion higher *HSF1* expression is correlated with

shortened survival (Figure 1B). Collectively, these findings support a pro-oncogenic role of HSF1

in *NF1*-deficient malignancies.

Given its critical role in governing protein folding, we were curious about how *HSF1* depletion

impacts the proteomes of *NF1*-deficient MPNST cells as well as *NF1*-proficient non-transformed

human Schwann cells (HSCs), both primary and immortalized. HSCs are considered as the cell-

of-origin of MPNSTs.[20,22] In three different MPNST cell lines (90-8TL, S462, and SNF96.2),

transient depletion of *HSF1*, using two previously validated lentiviral shRNAs,[9,13] all caused an

increase in global protein Lys48-specific polyubiquitination, a molecular mark for proteasomal

degradation,[23] in both detergent-soluble and -insoluble fractions (Figures 1C, 1D, S1B, and S1C).

This increased polyubiquitination signifies proteome-wide misfolding and aggregation. In stark

contrast, this change was not observed in HSCs (Figures 1C and 1D). Thus, *HSF1* depletion

provokes proteomic instability in MPNST cells, but not in HSCs.

Most cellular protein aggregates are amorphous; however, a tiny fraction becomes highly ordered

and enriched with β-sheet structures. These special protein aggregates, namely amyloids, are

closely associated with neurodegenerative disorders in humans, especially Alzheimer's disease.[24]

To assess whether *HSF1* depletion could elicit amyloidogenesis, we stained both MPNST cells and

HSCs with Congo red (CR), a widely applied amyloid-binding fluorescent dye.[25] Compared to

HSCs, MPNST cells displayed more intense CR staining (Figure 1E), suggesting increased

amyloids in malignant Schwann cells. Importantly, *HSF1* depletion further intensified CR staining

in MPNST cells, but not in HSCs (Figures 1E, 1F, S1D, and S1E). Similar results were obtained

by Thioflavin T staining (Figure S1F), another popular amyloid dye.[25] Importantly, the induction

of amyloids was further confirmed by ELISA using two conformation-specific antibodies (A11

and OC), which recognize soluble amyloid oligomers (AOs) and insoluble amyloid fibrils (AFs),

respectively.[26,27] Upon *HSF1* depletion, all three MPNST cell lines displayed elevated levels of

$A11^+$ AOs and OC^+ AFs (Figures 1G, 1H, S1G-S1J). Again, no evident increase in amyloids was

detected in HSCs (Figures 1I and 1J). Together, these findings indicate that HSF1 is required to

safeguard the proteome of malignant Schwann cells and particularly to repress amyloidogenesis;

on the contrary, it is dispensable for the proteome of non-transformed Schwann cells.

Unchecked proteomic instability is tumor suppressive

It has become evident that HSF1 acts as a generic oncogenic enabler; nevertheless, a universal

underlying mechanism remains undefined.

A healthy proteome enables the translation of genotypes into phenotypes. Thus, we reasoned that

preservation of proteomic stability is most likely a generic pro-oncogenic mechanism of HSF1.

To test this notion, we first determined whether proteomic instability, particularly

amyloidogenesis, is causally related to the cytotoxicity induced by *HSF1* depletion in cancer

cells. To block amyloidogenesis we employed CR, which is known to exert anti-amyloid effects

through its amyloid-binding activity.[5,28,29] Consistent with "HSF1 addiction of cancer", *HSF1*

depletion markedly impaired the viability of MPNST cells; importantly, CR treatment partially

rescued this impaired viability (Figures 2A, S2A, and S2B). In contrast, both *HSF1* depletion and

CR treatment did not affect HSCs (Figure 2B), congruent with a lack of proteomic instability and

amyloidogenesis in these non-transformed cells. Together, these findings support that unchecked

proteomic instability, particularly amyloidogenesis, is cytotoxic and tumor suppressive.

Moreover, HSF1 enables oncogenesis in part by constraining proteomic instability.

HSF1 neutralizes soluble amyloid oligomers to protect MPNST cells

Then, how does HSF1 counter amyloids to enable the growth and survival of MPNST cells?

Recently, by researching tissue overgrowth models we discovered that HSF1, surprisingly,

physically neutralizes soluble AOs, thereby preventing their assault on HSP60.[30] As an essential

mitochondrial chaperone, HSP60 functions as a guardian of the mitochondrial proteome.[30-32]

We wondered whether HSF1 acts the same way in both cancer and tissue overgrowth. Co-IP

experiments indicated that A11[+] AOs physically interacted with HSF1, but not major HSPs, in

MPNST cells (Figure 2C), consistent with our findings in tissue overgrowth models.[30] This AO-

HSF1 interaction was further validated by another independent technique, *in situ* Proximity

Ligation Assay (PLA) (Figures 2D-2F). Importantly, *HSF1* depletion promoted the AO-HSP60

interaction (Figure 2G and 2H). Loss of *HSP60* destabilized the mitochondrial proteome.[30,32]

Thus, this AO-HSP60 interaction signifies the instigation of mitochondrial damage and

cytotoxicity. Taken together, our findings reveal that HSF1 protects MPNST cells in part by

neutralizing soluble AOs to safeguard HSP60 and the mitochondrial proteome (Figure 2I).

MPNST cells react to HSF1 deficiency by activating JNK to attenuate translation

Given their dependency on HSF1, cancer cells are expected to mount certain responses to

survive and adapt to the lethal HSF1 inhibition. Previously, we uncovered that in *Hsf1*-deficient

mice, the stress-responsive JNK kinase becomes constitutively activated, which, in turn,

disassembles mTORC1.[33] JNK does so by phosphorylating both RAPTOR and mTOR, two core

components of mTORC1. This mTORC1 inhibition diminishes protein translation, leading to

reduced cell size and whole-body lean mass in *Hsf1*-deficient mice.[33]

We reasoned that cancer cells might exploit this molecular regulation for their survival and

adaptation. In MPNST cells *HSF1* knockdown activated JNK, indicated by elevated JNK

T183/Y185 phosphorylation; in parallel to JNK activation, there was mTORC1 inhibition,

evidenced by diminished p70S6K T389 phosphorylation and 4EBP1 S65/T70 phosphorylation

(Figures 3A, 3B, and S3A-S3D).[34,35] In contrast, *HSF*1 knockdown minimally affected JNK and

mTORC1 in HSCs (Figures 3A and 3B). Detected by PLA, in MPNST cells HSF1 interacted

with JNK; however, *HSF1* depletion promoted JNK-RAPTOR interactions but diminished

mTOR-RAPTOR interactions simultaneously (Figures 3C-3E). The mTOR-RAPTOR interaction

is essential to the assembly and activity of mTORC1.[36] Thus, these findings support that JNK

senses HSF1 deficiency to repress mTORC1. Of interest, compared to non-transformed cells,

this sensing mechanism appears more prominent in malignant cells.

As expected for mTORC1 inhibition, *HSF1* depletion reduced global protein translation rate in

MPNST cells (Figures 3F, S3E, and S3F), detected by puromycin labeling as described

previously.[33,37] Beyond MPNST cells, this regulation of translation by HSF1 was conserved in

diverse human cancer cell lines (Figure 3G). Importantly, treatment with JNK-IN-8, an

irreversible JNK inhibitor,[38] largely rescued this impaired protein translation owing to *HSF1*

depletion (Figures 3F, S3E, and S3F). Interestingly, JNK inhibition augmented the basal

translation rate as well. Consistent with these findings, in human cancers higher *HSF1*

expression is associated with lower JNK activation (Figure S3G); in turn, lower JNK activation

is accompanied by higher mTORC1 activity, reflected by higher 4EBP1 Ser65 phosphorylation

(Figure S3H). In aggregate, our findings demonstrate that JNK activation is causally linked to

inhibition of mTORC1 and translation. Therefore, through JNK suppression HSF1 permits

robust protein translation, which is fundamental to malignant growth (Figure 3H).

Lessening protein quantity is an adaptive response to alleviate proteomic instability

We next asked why cells respond to HSF1 deficiency by attenuating translation. While HSF1

controls protein quality via HSPs, mTORC1 governs protein quantity via translation. It would be

beneficial to lower protein quantity when protein quality is jeopardized. Thus, we reasoned that

this regulation might represent a cellular adaptive response to imbalanced protein quantity and

quality engendered by HSF1 deficiency.

Proteomic imbalance disrupts proteome homeostasis and contributes to proteomic instability,

which may ultimately trigger amyloidogenesis. To inflict proteomic imbalance, we first

stimulated protein translation in MPNST cells by inhibiting JNK. As expected, JNK-IN-8

reduced the phosphorylation of c-JUN, a canonical substrate of JNK,[34] in MPNST cells (Figure

4A). In support of enhanced protein translation, JNK-IN-8 further activated mTORC1 in these

cells (Figures 4A and S4A); in contrast, marginal effects were observed in HSCs (Figure 4B).

Congruently, JNK-IN-8 did not significantly reduce c-JUN phosphorylation in HSCs, supporting

a lack of JNK activation in these non-transformed Schwann cells. Intriguingly, the heightened

protein translation following JNK inhibition was accompanied by elevated amyloid levels in

MPNST cells (Figures 4C and S4B). On the contrary, there were no significant changes in

amyloid levels in HSCs (Figure 4D).

To determine the role of protein translation in amyloidogenesis, we chose to block translation

using two pharmacological inhibitors, 4EGI-1 and LY2584702, with distinct mechanisms of

action. While 4EGI-1 disrupts eIF4E-eIF4G interactions,[39] LY2584702 inhibits p70S6K.[40]

Interestingly, both inhibitors diminished the basal amyloid levels in MPNST cells (Figures

4E,4F, and S4C), revealing a causal relationship between protein translation and

amyloidogenesis. Importantly, translation inhibition also blocked the induction of amyloids by

JNK-IN-8 in MPNST cells (Figures 4G, 4H, S4D, and S4E), indicating that JNK suppresses

amyloidogenesis via translation inhibition. No significant effects were observed in HSCs

(Figures 4G and 4H).

Considering the role of translation in amyloidogenesis, we reasoned that preventing translation

attenuation would aggravate proteomic imbalance and enhance cytotoxicity in malignant cells.

As expected, JNK-IN-8 further elevated the amyloid levels in MPNST cells following *HSF1*

depletion, which was accompanied by exacerbated apoptosis, detected by caspase 3 cleavage

(Figures 4I, 4J, and S4F).[41] Importantly, CR treatment blocked apoptosis, detected by PARP

cleavage (Figure S4G),[42] causally implicating amyloidogenesis. In contrast, *HSF1* depletion and

JNK inhibition exerted no significant impacts on non-transformed Schwann cells (Figures 4K

and 4L). Thus, our findings support that proteomic imbalance in malignant cells contributes to

proteomic instability and amyloidogenesis. In stark contrast, there is no or little amyloidogenesis

in non-transformed cells, owing to their balanced protein quantity and quality. Moreover,

through translation attenuation, malignant cells attempt to survive and adapt to proteomic

imbalance (Figure 4M).

mTORC1 stimulation aggravates proteomic imbalance and cytotoxicity induced by HSF1

inhibition

JNK regulates numerous targets other than mTORC1. To stimulate mTORC1 and protein

translation specifically, we next genetically inhibited the tumor suppressor Tuberous Sclerosis

Complex subunit 2 (*TSC2*), a key negative regulator of mTORC1,[43] in MPNST cells. As

expected, knocking down *TSC2* in MPNST cells promoted phosphorylation of both p70S6K and

4EBP1, which was accompanied by elevated amyloids (Figures S5A-S5F). Like genetic *HSF1*

depletion, pharmacological inhibition of HSF1 by DTHIB elevated global protein

polyubiquitination and amyloids in MPNST cells (Figures 5A, 5B, and S5G), eliciting proteomic

instability. The newly developed small-molecule inhibitor DTHIB directly binds to the DNA-

binding domain of HSF1 and potently blocks the HSF1-mediated transcription.[44] Compared to

control cells, HSF1 inhibition by DTHIB induced considerably higher levels of amyloids in

TSC2-deficient MPNST cells (Figure 5B). Of note, *TSC2*-deficient cells displayed more

profound JNK activation and mTORC1 inhibition following DTHIB treatment, reflecting an

intensified cellular response to severe proteotoxic stress. Consistently, compared to slight or

intermediate cytotoxicity in control MPNST cells, detected by plasma membrane

permeabilization, *TSC2*-deficient cells were highly sensitive to HSF1 inhibition (Figures 5C, 5D,

and S5H-S5K). Importantly, either translation inhibition or CR treatment largely blocked this

cytotoxicity (Figures 5C and 5D), causally implicating proteomic instability and

amyloidogenesis. In support of a role of HSF1 in enabling *TSC2* deficiency, among patients

whose tumors harbor *TSC2* deletion higher *HSF1* gene expression is correlated with shortened

survival (Figure S5L).

To exclude possible mTORC1-independent effects of *TSC2* deficiency, we alternatively fed cells

with NV-5138, a leucine analog currently under clinical trials as an antidepressant.[45,46]Amino

acids, particularly leucine, are potent activators of mTORC1.[47] Of interest, whereas NV-5138 is

capable of activating mTORC1, it cannot be utilized for protein synthesis.[45,46] To maximize

mTORC1 stimulation *ex vivo*, cells were cultured in leucine-free medium. Like *TSC2*

knockdown, NV-5138 stimulated mTORC1 and heightened amyloid levels in MPNST cells

(Figures S5M and 5E); however, it only elicited pronounced cytotoxicity with concurrent HSF1

inhibition (Figure 5F). Again, CR treatment largely blocked this cytotoxicity (Figure 5F). In

sharp contrast, combined NV-5138 and DTHIB treatment did not significantly impact non-

transformed Schwann cells (Figure 5G). Of interest, their mTORC1 signaling was less

responsive to NV-5138 stimulation (Figures S5N). We next reasoned that by directly

contributing to protein quantity, L-leucine might impact the cancer proteome more profoundly.

In both S462 and 90-8TL cells, L-leucine synergized with DTHIB to induce severe cytotoxicity

(Figures 5H and S5O). On the contrary, no cytotoxicity was induced in non-transformed

Schwann cells (Figure 5I).

To further corroborate the HSF1-dependent effects of DTHIB, we evaluated another direct

small-molecule HSF1 inhibitor KRIBB11[48], which blocks the recruitment of pTEFb likely

through binding to the transactivation domain of HSF1. KRIBB11, like DTHIB, synergized with

L-leucine to induce amyloids and trigger severe cytotoxicity in MPNST cells; importantly, CR

treatment both diminished amyloids and blocked cell death (Figures 5J and 5K).

Furthermore, in S462 cells shRNA-mediated *HSF1* knockdown and L-leucine stimulation

synergistically provoked prominent cytotoxicity as well (Figure 5L), providing a genetic

validation. Thus, these findings collectively indicate that mTORC1 stimulation synergizes with

HSF1 inhibition to rapidly trigger severe cytotoxicity in cancerous cells, in part by prompting

unconstrained proteomic imbalance and amyloidogenesis.

Unconstrained proteomic imbalance incites non-apoptotic, non-autophagic death

HSF1 inhibition alone induced both apoptotic and non-apoptotic death in MPNST cells,

distinguished by caspase 3 cleavage; intriguingly, concurrent mTORC1 stimulation caused a

striking shift towards non-apoptotic cell death, characterized by cell membrane permeabilization

without caspase 3 cleavage (Figures 6A and 6B).

To date, several modes of non-apoptotic cell death have been described, including pyroptosis,

necroptosis, ferroptosis, and autophagic death, all of which could result in loss of cell membrane

integrity. Corroborating a non-apoptotic mechanism, the pan-caspase inhibitor Q-VD-OPh did

not prevent the membrane permeabilization in *TSC2*-deficient MPNST cells treated with DTHIB

(Figures 6A and 6B).[49] However, it did block the apoptosis in control MPNST cells treated with

DTHIB (Figures 6A and S6A). As Q-VD-OPh inhibits caspase 1 and no caspase 1 cleavage was

detected (Figure S6B),[50] a key driver of pyroptosis,[51] this result does not support a pyroptotic cell

death either. Similarly, HSF1 inhibition did not induce necroptosis in *TSC2*-deficient MPNST

cells, evidenced by the lack of MLKL Ser358 phosphorylation (Figure S6B), an essential event

executing necroptosis.[52] Congruently, the necroptosis inhibitor necrostatin-1 exerted no effects

on the cell membrane disruption (Figures 6C and S6C).[52]

Intriguingly, HSF1 inhibition diminished glutathione peroxidase 4 (GPX4) but increased

transferrin receptor 1 (TFR1), which was accompanied with markedly elevated lipid

peroxidation, in *TSC2*-deficient MPNST cells (Figures 6D and S6D). All these changes are

indicative of ferroptosis.[53,54] Although in *TSC2*-deficient cells liproxstatin-1, a potent ferroptosis

inhibitor,[53,54] reversed the elevation of TFR1 elicited by the ferroptosis inducer erastin, it neither

blocked the marked increase in TFR1 nor stopped the membrane disruption induced by DTHIB

(Figures 6C, 6D, and S6C). This is not due to the defectiveness of necrostatin-1 and liproxstatin-

1, as both inhibitors potently blocked necroptosis and ferroptosis induced in HT-29 cells,

respectively (Figure S6E). HSF1 inhibition also elicited autophagy, evidenced by diminished

p62/SQSTM1 and elevated LC3B-II (Figure 6E).[55] To determine whether autophagic cell death

was implicated, we employed the class III PI3K inhibitor wortmannin to block autophagy.[56] As

expected, wortmannin effectively prevented the decreases in p62/SQSTM1 and LC3B-II induced

by rapamycin in *TSC2*-deficient MPNST cells (Figure 6E), indicating successful blockade of

autophagy. Nevertheless, wortmannin failed to block the cell membrane disruption in *TSC2*-

deficient cells treated with HSF1 inhibitors (Figures 6F and S6F). These findings suggest that

both ferroptosis and autophagy are the consequences, rather than the causes, of cell membrane

disruption. Collectively, our findings support that uncontained proteomic instability and ensuing

amyloidogenesis potently trigger cell death, which is neither apoptotic nor autophagic (Figure

6G). Moreover, several common modes of programmed necrosis are not deeply implicated as

well. The precise nature of this proteomic catastrophe-induced death (PCID), nonetheless,

remains to be fully delineated.

Unconstrained proteomic imbalance suppresses *in vivo* tumor growth

Due to the limited *in vivo* bioavailability of DTHIB in our mouse models, we focused on

KRIBB11 for *in vivo* studies. Consistent with *ex vivo* findings, combined L-leucine stimulation

and KRIBB11 administration significantly impaired the *in vivo* growth of xenografted human

S462 MPNST cells in immunocompromised mice (Figure 7A). Despite this overall impact, 2 out

of 8 tumors did not respond to the combination treatment (Figure S7A), creating large variations

and right-skewed distribution. Of note, without leucine stimulation KRIBB11 only marginally

affected tumor growth; by contrast, in the presence of leucine KRIBB11 did evidently slow

tumor exponential growth rates and delay tumor volume doubling time (Figures S7B). In

addition, mice receiving this combination treatment tended to survive better as well (Figure

S7C).

Next, we asked if this therapeutic strategy could be applied broadly. To this end, we established

a syngeneic mouse xenograft model of melanoma, where the murine melanoma B16-F10 cells

were transplanted into C57BL/6J mice. Like in the MPNST model, KRIBB11 combined with L-

leucine stimulation also impeded the tumor growth in this fast-growing tumor model (Figures 7B

and S7D). As expected, L-leucine stimulated mTORC1 signaling, evidenced by increased S6RP

phosphorylation (Figure 7C). KRIBB11 reduced the transcription of *Hspa1a/Hsp72* and

Hspb1/Hsp25 in melanomas (Figure 7D), confirming a successful inhibition of HSF1.

Importantly, combined KRIBB11 administration and leucine stimulation markedly induced

amyloids in tumors (Figure 7E). By stark contrast, no amyloids were induced in normal tissues

including brains and spleens (Figures 7F and 7G). Treatments did not cause loss of body weights

in mice (Figure S7E). In aggregate, these *in vivo* findings from two distinct mouse models

support that unconstrained proteomic imbalance, provoked by combined HSF1 inhibition and

mTORC1 stimulation, causes proteomic catastrophe that is tumor suppressive (Figure 7H).

DISCUSSIONS

Genomic instability has been broadly recognized as a driver of oncogenesis. Contrastingly, the implications of proteomic instability in cancer remain obscure.

Proteomic instability is tumor suppressive but inevitable

Proteomic instability does occur in cancer, evidenced by elevated protein polyubiquitination, aggregation, and even amyloidogenesis. In cancer, proteomic instability can be ascribed to several mechanisms, including protein oxidation by reactive oxygen species, protein misfolding owing to genetic mutations, and augmentation of protein dosage caused by aneuploidy.[1,3] Revealed by our recent studies, uncontrolled protein translation also contributes to amyloidogenesis. This is not entirely surprising, as it has been estimated that up to 30% of all newly synthesized polypeptides are misfolded and subjected to proteasomal degradation.[57] Imaginably, aberrant protein synthesis would increase the chance of producing protein aggregates and even amyloids, especially considering numerous genetic mutations affecting protein folding in cancer cells. Dysregulated protein translation, ironically, is fundamental to malignant growth, making proteomic instability and amyloidogenesis inevitable.

At the cellular level, amyloids are toxic to both neurons and cancer cells. At the organismal level, amyloidogenesis is neurodegenerative but tumor suppressive. Mechanistically, our studies demonstrate that soluble AOs can directly attack the guardian of the mitochondrial proteome HSP60. Of course, mechanisms other than targeting HSP60 exist as well. Moreover, it remains to be elucidated what other proteins comprise cancer-associated amyloids, in addition to the Alzheimer's disease-associated Aβ.

HSF1 contains proteomic instability to enable malignancy

The multifaceted pro-oncogenic roles of HSF1 have been ascribed to a wide array of underlying

mechanisms; nonetheless, a universal one remains to be defined.

As an inevitable price of malignancy, proteomic instability must be contained by cancer cells to

avert tumor suppression and sustain malignant phenotypes. Now, our findings reveal that HSF1

contains proteomic instability, particularly amyloidogenesis, constituting a generic pro-

oncogenic mechanism. It is worth noting that HSF1 only contains and mitigates, but cannot

eradicate, amyloidogenesis, as uncontrolled protein translation is a direct cause of

amyloidogenesis in cancer cells. This dependency on HSF1 to contain proteomic instability

partly underlies the phenomenon of "HSF1 addiction of cancer".

Our studies further support that HSF1 contains amyloidogenesis via at least two mechanisms,

direct and indirect. On the one hand, through induction of HSPs, HSF1 ensures appropriate

protein folding to maintain a healthy proteome, thereby preventing the emergence of amyloids.

This is supported by the finding that HSF1 inhibition heightens protein ubiquitination,

aggregation, and amyloidogenesis. On the other hand, HSF1 can neutralize soluble AOs directly

once they emerge. Under this context, HSF1 inhibition would unleash AOs to assault HSP60,

inflicting mitochondrial damage and toxicity. Thus, in cancer cells HSF1 inhibition both enlarges

the quantity of amyloids and, more importantly, renders them toxic.

mTORC1 signaling becomes tumor suppressive in the absence of HSF1

Given the dependency of malignant growth on protein translation, unsurprisingly, mTORC1

signaling is hyperactive in human cancers, which is, canonically, considered oncogenic.[58,59]

Unexpectedly, our current studies reveal that mTORC1 signaling becomes tumor suppressive in

the absence of HSF1, owing to profound disproportion between protein quantity and quality.

This unconstrained proteomic imbalance, in turn, exacerbates proteomic instability and tumor-

suppressive amyloidogenesis. The notable blockage of cytotoxicity by either translation

inhibitors or CR lends strong support to this model. Conceptually, HSF1 empowers the

oncogenic potential of mTORC1 signaling, reinforcing HSF1 as an oncogenic enabler.

Provoking proteomic catastrophe exemplifies a next-generation therapeutic concept

Currently popular chemo-and radiotherapies mainly target the cancer genome, causing DNA

damage. An emerging horizon in cancer therapy is targeting the cancer proteome. To date, much

effort has been devoted to inhibiting individual HSP families, such as HSP70 and HSP90.[60,61]

However, every cell is equipped with both constitutive and inducible HSPs. While the former

constitute the basal chaperoning capacity that is essential to life, the latter supply the extra

chaperoning capacity, which is dispensable for normal life but required under stressful

conditions. Their structural and functional resemblances pose the biggest challenge for small

molecules to distinguish these cognate HSPs. Moreover, multiple families of HSPs, including

HSP100, HSP90, HSP70, HSP60, HSP40, and small HSPs, exist inside cells. Targeting single

HSP family would have limited therapeutic effects. Thus, new approaches or directions are

warranted to overcome these drawbacks.

Our and others' studies collectively support that malignant cells constantly endure intrinsic

proteotoxic stress, rendering HSF1 constitutively active. In turn, cancer cells become reliant on

the extra chaperoning capacity, conferred by HSF1, for their growth and survival. Thus, only

eliminating this extra chaperoning capacity but sparing the basal one would be an ideal strategy

to combat malignancy. Together, these concepts provide strong rationales for targeting HSF1 in

cancer. It is exciting to witness the recent launch of HSF1 inhibitors into clinical trials.[62]

In reaction to HSF1 inhibition, unsurprisingly, cancer cells attenuate their protein synthesis. This

adaptive response underscores the importance of balanced protein quantity and quality to cellular

fitness, a key concept inspiring us to explore driving proteomic imbalance for cancer therapy. Of

note, breaking this adaptation may impede tumor evolution.

The widespread hyperactivation of mTORC1 in human cancers makes it an attractive therapeutic

target. Hence, an array of mTOR inhibitors have been developed, several of which are actively

under clinical evaluation.[59,63] Thus far, these inhibitors have displayed mostly cytostatic effects

and less than expected therapeutic efficacies.[63,64] Considering the essentiality of protein

translation to life, profound mTORC1 inhibition would inflict undesirable toxicities in normal

cells and tissues. In fact, mTORC1 inhibitors have long been known as immunosuppressants.[63,65]

Opposite to the prevailing tactic in cancer therapy, we stimulated mTORC1 to drive proteomic

imbalance. Counterintuitive mTORC1 stimulation, when combined with HSF1 inhibition,

engendered catastrophic proteomic instability in cancer and elicited tumor suppression (Figure

7G). This PCID offers a wide therapeutic window, largely due to the lack of proteomic

instability and amyloidogenesis in primary cells and tissues. Conceivably, mTORC1 stimulation

boosts production of wild-type proteins in primary cells, which can be easily managed without

impacting well-preserved proteomic stability; in sharp contrast, it creates an enormous problem

for the already stressed, fragile cancer proteome through synthesis of copious mutant, misfolded

proteins. The adaptive response, noteworthily, still arises from concurrent mTORC1 stimulation

and HSF1 inhibition in cancer. It is expected that additional, specific blockade of the JNK-

mediated mTORC1 repression would further exacerbate proteomic imbalance, impelling

cancerous cells to extinction. It remains possible that leucine stimulation could exert further

mTORC1-independent effects, contributing to tumor suppression. Transcription factors,

traditionally, have been regarded as "undruggable" targets for small molecules. While both

DTHIB and KRIBB11 epitomize the prototypes of direct HSF1 inhibitors, their inadequate *in*

vivo bioactivities render them unsuitable for clinical applications at the current state.

Nevertheless, we herein accomplished the proof-of-concept experiments to demonstrate the

feasibility of propelling proteomic catastrophe in cancer. This next-generation therapeutic

concept may be applicable to a broad spectrum of human cancers and, with the advent of clinic-

relevant molecules, holds promise for combating malignancy.

ACKNOWLEDGMENTS

We would like to thank the Optical Microscopy and Image Analysis lab (OMAL) for their assistance with the confocal microscopy studies and Holly Morris for her assistance in animal studies. Schematic illustrations are created with BioRender.com. This work was supported by the grant from Department of Defense Congressionally Directed Medical Research Programs to C.D. (NF150028) and by the Intramural Research Program of the NIH, National Cancer Institute, Center for Cancer Research (1ZIABC011767). The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. government.

AUTHOR CONTRIBUTIONS: B.M.R., O. S., and K-H.C. designed and conducted the experiments. C.D. conceptualized and supervised this study and analyzed the results. B.M.R. and C.D. wrote the manuscript.

DECLARATION OF INTERESTS: The authors declare no competing interests.

EXPERIMENTAL MATERIALS AND METHODS

Cells, Chemicals, and reagents

S462 (female) and 90-8TL (female) cells were kindly provided by Dr. Karen Cichowski.

sNF96.2 cells (male) and immortalized human Schwann cells (hTERT ipn02.3 2λ, female) were

purchased from ATCC. Primary human Schwann cells (Cat#1700, ScienCell Research

Laboratories) were grown on poly-L-lysine coated T-75 flasks and maintained in complete

Schwann Cell Medium (Cat#1701, ScienCell Research Laboratories). B16-F10-luc2 (male)

cells were purchased from ATCC (cat#CRL-6475-LUC2). All cell lines, except 90-8TL and

HSCs (no known STR profiles available), were authenticated by ATCC. All cell lines, except

primary human Schwann cells, were maintained in DMEM supplemented with 10% HyClone

bovine growth serum and 1% penicillin–streptomycin. Cells were maintained in an incubator

with 5% CO_2 at 37 °C.

The following chemicals were purchased from commercial sources: JNK-IN-8 (Cat#S4901,

Selleckchem), 6-FAM-dC-puromycin (cat#NU-925-6FM, Jena Bioscience), Congo red

(cat#B2431014, Thermo Scientific Chemicals), Thioflavin T (cat#211760050, Thermo Scientific

Chemicals), LY2584702 (cat#S7698, Selleckchem), 4EGI-1 (cat#S7369, Selleckchem), and

DTHIB (cat# HY-138280, MedChemExpress), KRIBB11 (cat#HY-100872, MedChemExpress),

KRIBB11 (cat#T3652, TargetMol Chemicals), Q-VD-OPh (cat#HY-12305, MedChemExpress),

necrostatin-1 (cat#HY-15760, MedChemExpress), Liproxstatin-1 (cat#HY-12726,

MedChemExpress), wortmannin (cat#HY-10197, MedChemExpress), rapamycin (cat#HY-

10219, MedChemExpress), erastin (cat#HY-15763, MedChemExpress), MHY1485 (cat#HY-

B0795, MedChemExpress), NV-5138 HCl (cat#HY-114384B, MedChemExpress), and Liperfluo

(cat#L248, Dojindo Laboratories), dimethylacetamide (cat#T19439, TargetMol Chemicals

Inc.), PEG300 (cat#T7022, TargetMol Chemicals Inc.), and Captisol (cat#RC-0C7-100, CyDex

Pharmaceuticals).

Rabbit phospho-4EBP1 S65/T70 antibody (cat#sc-12884-R), mouse monoclonal HSF1 (E-4,

cat#sc-17757) antibody, rabbit HSF1 antibody (H-311, cat#sc-9144), mouse monoclonal JNK

(D-2, cat#sc-7345), rabbit JNK1/3 (C-17, cat#sc-474) antibody were purchased from Santa

Cruz Biotechnology; antibodies for phosphor-4EBP1 Thr70 (cat#9455), phospho-4EBP1

Thr37/46 (236B4, cat#2855), phospho-p70S6K T389 (108D2, cat# 9234), phosphor-S6RP

Ser235/236 (D57.2.2E, cat#4858), p70 S6K (49D7, cat#2708), 4EBP1 (53H11, cat#9644),

S6RP (5G10, cat#2217), mTOR (7C10, cat#2983), phospho-JNK1/2 T183/Y185 (81E11,

cat#4668), JNK2 (56G8, cat#9258), c-Jun (60A8, cat#9165), phospho-c-Jun Ser63 II

(cat#9261), PARP (cat#9542), Tuberin/TSC2 (D93F12, cat#4308), SQSTM1/p62 (D1Q5S,

cat#39749), cleaved caspase 3 (Asp175) (5A1E, cat#9664), caspase 1(D7F10, cat#3866),

gasdermin D (E9S1X, cat#39754), phosphor-MLKL Ser358 (D6H3V, cat#91689), GPX4

(E5Y8K, cat#59735), and transferrin receptor/CD71 (H68.4, cat#46222) antibody were

purchased from Cell Signaling Technologies; mouse β-actin antibody-HRP (Cat#A00730)

conjugates were from GenScript. Anti-K48-specific ubiquitin antibody (Apu2, cat#05-1307)

and mouse monoclonal anti-RAPTOR antibody (1H6.2, cat#05-1470) were purchased from

EMD Millipore. Mouse monoclonal anti-JNK1 (C-terminal region) antibody (cat#JM2671)

was purchased from ECM Biosciences. Anti-amyloid oligomers (A11, cat#SPC-506D) and

anti-amyloid fibrils (OC, cat#SPC-507D) antibodies were purchased from StressMarq

Biosciences Inc. Mouse monoclonal anti-HSP60 antibody (LK-1, cat#ADI-SPA-806) was

purchased from Enzo Life Sciences Inc. β-Actin (cat#GTX629630), GAPDH

(cat#GTX627408), α-tubulin (GT114, cat#GTX628802), and LC3B antibody

(cat#GTX127375) was purchased from GeneTex Inc.

METHOD DETAILS

Lentiviral Particle Production and Transduction of lentiviral shRNAs

Lentiviral pLKO.1 shRNA plasmids targeting human *HSF1* were described previously[10]:

TRCN0000007480 (hA6) and TRCN0000007483 (hA9). The sequence of control hairpins

targeting a scrambled sequence with no known homology to any human genes is 5′-

CCTAAGGTTAAGTCGCCCTCG-3′. The pLKO.1 shRNA plasmid targeting human *TSC2* was

a gift from Do-Hyung Kim (Cat#15478, Addgene). Lentiviral pLKO shRNA plasmids were

transiently co-transfected individually along with plasmids encoding ΔVPR and VSV-G into

HEK293T cells using TurboFect™ (Cat#R0531, Thermo Scientific). Viral supernatants were

collected after centrifugation at 1,000 rpm for 15 min. Viral transduction was achieved by

incubating target cells with viral supernatants containing 10μg/ml polybrene (Cat#TR-1003-G,

Millipore-Sigma) for 24h. For stable knockdown of *TSC2,* lentivirus particles were prepared using

pLKO shRNA plasmids targeting human *TSC2*. Scrambled shRNA was used for stable control

cells. After 4 days of transduction, cells were gradually selected using 1-4 μg/ml puromycin for 2

weeks.

Soluble and Insoluble Protein Fractionation

Equal numbers of cells were incubated with cold cell-lysis buffer (100 mM NaCl, 30 mM Tris-

HCl pH 7.6, 1% Triton X-100, 20 mM sodium fluoride, 1mM EDTA, 1mM sodium

orthovanadate, and 1x Halt™ protease inhibitor cocktail) on ice for 20 min. The crude lysates

were first centrifuged at 500xg for 5 min at 4°C. The supernatants from low-speed (500xg)

centrifugation were transferred to a new Eppendorf tube. The first pellets were resuspended in

DNA digestion solution (2 units DNase I, 1% Triton X-100, 10 mM Tris-HCl, 2.5 mM

$MgCl_2$, 0.5 mM $CaCl_2$, pH 7.6) for 30 min at RT. An equal volume of 1xPBS with 2% SDS

was then added to resuspend the first pellets for additional 30 min at RT. Following the

centrifugation at 17,000xg for 10 min at RT, the supernatants were removed. The supernatants

from low-speed centrifuge were added back to the second pellets, followed by centrifugation

at 17,000xg for 15 min at 4°C. The final supernatants and pellets from high-speed (17,000xg)

centrifugation were collected as detergent-soluble and -insoluble fractions, respectively.

Insoluble fractions were further sonicated in 1xPBS with 2% SDS at high intensity using a

Bioruptor® Sonication Device (Diagenode) for SDS-PAGE.

ThT and CR Staining for Flow Cytometry

After washing with 1xPBS, cells were fixed with 4% formaldehyde at RT for 10 min.

Following fixation and 1xPBS washing, cells were resuspended in 1 ml permeabilization

buffer (1xPBS with 0.5% Triton X-100 and 3 mM EDTA) and incubated on ice for 30 min.

Following washing with 1xPBS, Cells were stained with 10 μM ThT or CR dissolved in

1xPBS for 5 min. Cells were washed and resuspended in PBS and fluorescence signals were

measured by a FACSCalibur, LSRFortessa X-20 or LSR II flow cytometer (BD Biosciences).

Amyloid Oligomer and Fibril Quantitation by Direct ELISA and Flow Cytometry

To quantitate soluble amyloid prefibrillar oligomers, soluble cell lysates diluted in 1xPBS

from equal numbers of cells were coated on each well in a 96-well ELISA plate at 4°C

overnight followed by blocking (5% non-fat milk in TBS-0.05% Tween® 20) at RT for 1 hr.

Each well was incubated with 100 µl amyloid oligomer antibodies (A11, 1:1,000 diluted in

the blocking buffer) at RT for 2 hr. After washing with TBS-T, goat anti-rabbit secondary Ab

HRP conjugates (1:1,000 diluted in blocking buffer) were added to each well and incubated at

RT for 1 hr. Following washing, 100 µl 1-Step Ultra TMB-ELISA substrates (Cat#34029,

Thermo Scientific) were added to each well. After adding 100µl Stop solution, absorbance was

measured at 450 nm using a SpectraMax iD5 microplate reader (Molecular Devices).

To quantitate amyloid fibrils, detergent-insoluble fractions from equal numbers of cells were

extracted as described in Soluble and Insoluble Protein Fractionation. The final pellets were

collected as detergent-insoluble fractions and solubilized by sonication for 15 min in 1xPBS

with 2% SDS. Solubilized proteins diluted in 1xPBS were added to each well and incubated at

37°C without cover overnight to air dry the wells. The following steps were identical to the

oligomer detection except for the use of amyloid fibril antibodies (OC) as the primary Ab.

For amyloid oligomer and fibril quantification by flow cytometry, cells were collected by

trypsinization and fixed in 4% Formaldehyde at RT for 10 min. Following permeabilization using

0.3% Triton-X-100, samples were incubated with primary antibodies A11-FITC (1:100 in 5%

BSA, StressMarq Biosciences, cat# SPC-506D-FITC) or OC-PerCP (1:100 in 5% BSA,

StressMarq Biosciences, cat# SPC-507D-PCP) at RT for 3 hrs. The cells were washed with PBS

and analyzed using flow cytometry.

Measurement of Cell Viability

Cell viability was determined using the CellTiter-Blue® Cell Viability Assay (cat#G8080, Promega). Cells were grown in 96 well plates (1×10^4 per well) for 24h. A 20µl CellTiter-Blue® reagent was added to 100µl of media in each well. The plates were incubated for 2h at 37°C and the fluorescence was recorded at 560/590 nm. The reagents were discarded, and the wells were washed with PBS before adding fresh media. Similarly, subsequent readings were recorded at 24h intervals till 96h time point.

Proximity Ligation Assay

Cells were fixed with 4% formaldehyde in PBS for 15 min at RT. After blocking with 5% normal goat serum in PBS with 0.3% Triton X-100, primary antibodies 1:100 diluted in blocking buffer were incubated with fixed cells overnight at 4°C. Following washing with PLA wash Buffer A, samples were incubated with 1:5 diluted Duolink® in Situ PLA Probe anti-rabbit Plus (cat#DUO92002, Sigma-Aldrich) and anti-mouse Minus (cat#DUO92004, Sigma-Aldrich) at 37°C for 1 hr. Subsequently, ligation (30 minutes, 37°C), rolling circle amplification, and detection (120 minutes, 37°C) were performed using the Duolink® In Situ Detection Reagents Red (cat#DUO92008, Sigma-Aldrich). Nuclei were counterstained with Hoechst 33342. PLA signals were documented by a Zeiss LSM780 confocal microscope (Carl Zeiss). For PLA flow cytometry, an equal number of cells were collected in microcentrifuge tubes by trypsinization and fixed with 4% formaldehyde in PBS for 10 min at RT. Subsequent PLA steps were followed as described earlier. The PLA signals were quantified by a FACSCalibur flow cytometer (BD Biosciences).

Mouse anti-JNK1 (JM2671) and rabbit anti-HSF1 (H-311) antibodies were used for JNK-HSF1

PLA. Rabbit anti-JNK1/3 (C-17) and mouse anti-RAPTOR (1H6.2) antibodies were used for

JNK-RAPTOR PLA. Rabbit anti-mTOR (7C10) and mouse anti-RAPTOR (1H6.2) antibodies

were used for mTOR-RAPTOR PLA. Mouse anti-HSF1 (E-4) and rabbit anti-AO (A11)

antibodies were used for HSF1-AO PLA. Rabbit anti-AO (A11) and mouse anti-HSP60 (LK1)

antibodies were used for AO-HSP60 PLA.

Measurement of global protein translation

Live cells were incubated with 100 nM 6-FAM-dC-puromycin in complete culture medium

for 30 min and analyzed by a FACSCalibur flow cytometer (BD Biosciences).

Immunoblotting and Immunoprecipitation

Whole-cell protein extracts were prepared in cold cell-lysis buffer (100 mM NaCl, 30 mM Tris-

HCl pH 7.6, 1% Triton X-100, 20 mM sodium fluoride, 1mM EDTA, 1mM sodium

orthovanadate, and 1x Halt™ protease inhibitor cocktail). Proteins were transferred to

nitrocellulose membranes. Following incubation with the blocking buffer (5% non-fat milk in 1x

TBS-T) for 1 hour at RT, membranes were incubated with primary antibodies (1:1,000 dilution in

the blocking buffer) overnight at 4 °C. After washing with 1xTBS-T for 3 times, membranes

were incubated with peroxidase-conjugated secondary antibodies (1: 3,000 diluted in the

blocking buffer) at RT for 1 hr. Signals were detected using SuperSignal™ West

chemiluminescent substrates. For normal Co-IP, proteins were extracted using a QSonica Q125

sonicator (total process time: 15S, pulse-on time: 5S, pulse-off time: 10S, output intensity: 30%)

in 1x sonication buffer (20 mM Tris, 20 mM NaCl, 1 mM EDTA, 20 mM β-glycerol-phosphate,

20 mM sodium fluoride, 4 mM sodium orthovanadate, and 1 mM DTT pH7.4, supplemented

with Halt protease inhibitor cocktail). Total 1 mg whole cell lysates were incubated with A11

antibodies at 4 °C overnight. Normal rabbit IgG were used as the negative controls. The primary

antibody was precipitated using Protein G magnetic beads. After washing with the lysis buffer

for 3 times, beads were boiled in 1x loading buffer for 5 min before loading on SDS-PAGE.

EasyBlocker (Cat#GTX425858, GeneTex) was used for blocking and anti-HSF1 (H-311)

primary antibody (cat#sc-9144, Santa Cruz Biotechnology) incubation (1:1,000, O/N at 4°C).

The blot was incubated with EasyBlot anti-Rabbit IgG HRP conjugated antibody

(cat#GTX221666-01, GeneTex) for 1 hour. Chemiluminescent signals were captured by an

iBright™ FL1000 imaging system (ThermoFisher Scientific).

Live-Or-Dye™ Dead Cell Staining

Cells were cultured in DMEM for 48 hours before being treated with inhibitors for the required

periods and collected by trypsinization. $1x10^6$ cells/ml cells were stained with Live-or-Dye™

fixable dead cell dyes (1:1,000 dilution, cat#32003, Biotium) at RT for 30 min, protected from

light. Stained cells were washed once with PBS and analyzed by flow cytometry. For co-staining

with cleaved caspase 3 antibody, stained cells were fixed with 4% formaldehyde (10 min, RT)

and permeabilized with 0.3% Triton-X-100 in PBS (10 min, RT). The cells were stained by

Cleaved Caspase 3 antibody (Asp175) (5A1E) (1:1,000, cat#9664, Cell Signaling Technology)

followed by goat anti-Rabbit IgG Alexa Fluor™ 488 conjugated (1:1,000, cat#A-11034,

ThermoFisher Scientific) secondary antibodies. The samples were analyzed by flow cytometry.

Inhibitor treatments

Treatments with DTHIB, Q-VD-OPh, Liproxstatin-1, Necrostatin-1, and Wortmannin were done for 72 hours. For Congo Red and LY-2584702 treatments, cells were pretreated for 24h followed by DTHIB for an additional 72h. For treatments with NV-5138, cells were pretreated for 48 hours in a leucine-free DMEM (cat#226-024, Crystalgen) before adding DTHIB for an additional 72 hours.

Animal studies

All animal experiments described here were approved by the institutional Animal Care and Use Committee (ACUC) of the National Cancer Institute, Frederick. Experiments on all animals were conducted following the recommendations of the Guide for the Care and Use of Laboratory Animals (National Academies Press, 2011). Animals were housed and maintained on a 12-hour light/12-hour dark cycle at temperatures between 20°C and 27°C and humidity levels of 30% to 70%, with food and water *available ad libitum*.

To establish the MPNST xenograft model, $9x10^6$ S462 MPNST cells (female) suspended with 30 % Matrigel were subcutaneously implanted the right flanks of 8-week-old female NIH-III nude mice (Charles River Laboratories). After 7 days, xenografted mice were assigned to 4 groups, eight each group, by stratified randomization based on tumor sizes (between 72 and 199 mm³). Mice were treated with KRIBB11 (65mg/kg, i.p. injection daily, cat#HY-100872, Lot#317606, Medchemexpress LLC) with and without 2% L-leucine dissolved in drinking water stored in amber-colored bottles. KRIBB11 was freshly dissolved daily in the vehicle (final concentration:10% dimethylacetamide, 50% PEG300, 20% Captisol, and 20% sterile saline).

To establish the syngeneic melanoma model, 2×10^6 B16-F10-luc2 cells (male) mixed with

Matrigel (1:1 ratio) in a 200μL volume were subcutaneously implanted into the right flanks of

8-week-old C57BL/6J male mice (The Jackson Laboratory). After 5 days, xenografted mice were

assigned to 4 groups, eight each group, by stratified randomization based on tumor sizes (between

75 and 284 mm^3). Mice were treated with KRIBB11 (65 mg/kg, i.p. injection daily, cat#T3652,

Lot#132988, TargetMol Chemicals Inc.) with and without 2% L-leucine dissolved in drinking

water stored in amber-colored bottles. KRIBB11 was freshly dissolved daily in the vehicle (final

concentration:10% dimethylacetamide, 50% PEG300, 16% Captisol, and 24% sterile saline).

To formulate the KRIBB11 solution, KRIBB11 was first dissolved in dimethylacetamide,

followed by addition of PEG300. Lastly, 40 or 50% Capitol dissolved in sterile saline was added.

Tumor sizes were measured using a caliper by an animal technician blind to the study. To

assess the compound's toxicity, the body weight of tumor-bearing animals was also recorded.

Tumor volumes were estimated using the formula [length (mm) × width^2 (mm^2)]/2.

Quantification And Statistical Analysis

Statistical analyses were performed using Prism GraphPad 10.0 (GraphPad Software). The

detailed statistical methods and sample sizes are provided in the figure legends. All results are

expressed as mean±SD, mean±SEM, or median and IQR. The statistical significance is defined

as: *p<0.05, **p<0.01; ***p<0.001; ****p<0.0001; n.s.: not significant. For *in vitro*

experiments, sample size required was not determined a priori. The experiments were not

randomized. For *in vivo* experiments, sample size required was determined by pilot experiments.

REFERENCES

1. Dai C, Dai S, Cao J. Proteotoxic stress of cancer: implication of the heat-shock response in oncogenesis. J Cell Physiol. 2012;227(8):2982-7. doi: 10.1002/jcp.24017. PubMed PMID: 22105155; PMCID: PMC3431153.

2. Dai C. The heat-shock, or HSF1-mediated proteotoxic stress, response in cancer: from proteomic stability to oncogenesis. Philos Trans R Soc Lond B Biol Sci. 2018;373(1738). doi: 10.1098/rstb.2016.0525. PubMed PMID: 29203710; PMCID: PMC5717525.

3. Guang MHZ, Kavanagh EL, Dunne LP, Dowling P, Zhang L, Lindsay S, Bazou D, Goh CY, Hanley C, Bianchi G, Anderson KC, O'Gorman P, McCann A. Targeting Proteotoxic Stress in Cancer: A Review of the Role that Protein Quality Control Pathways Play in Oncogenesis. Cancers (Basel). 2019;11(1). Epub 20190109. doi: 10.3390/cancers11010066. PubMed PMID: 30634515; PMCID: PMC6356294.

4. Wilson DM, 3rd, Cookson MR, Van Den Bosch L, Zetterberg H, Holtzman DM, Dewachter I. Hallmarks of neurodegenerative diseases. Cell. 2023;186(4):693-714. doi: 10.1016/j.cell.2022.12.032. PubMed PMID: 36803602.

5. Tang Z, Dai S, He Y, Doty RA, Shultz LD, Sampson SB, Dai C. MEK guards proteome stability and inhibits tumor-suppressive amyloidogenesis via HSF1. Cell. 2015;160(4):729-44. doi: 10.1016/j.cell.2015.01.028. PubMed PMID: 25679764; PMCID: PMC4564124.

6. Li J, Labbadia J, Morimoto RI. Rethinking HSF1 in Stress, Development, and Organismal Health. Trends Cell Biol. 2017;27(12):895-905. Epub 20170907. doi: 10.1016/j.tcb.2017.08.002. PubMed PMID: 28890254; PMCID: PMC5696061.

7. Anckar J, Sistonen L. Regulation of HSF1 function in the heat stress response: implications in aging and disease. Annu Rev Biochem. 2011;80:1089-115. doi: 10.1146/annurev-biochem-

060809-095203. PubMed PMID: 21417720.

8. Gomez-Pastor R, Burchfiel ET, Thiele DJ. Regulation of heat shock transcription factors and

their roles in physiology and disease. Nat Rev Mol Cell Biol. 2018;19(1):4-19. Epub

20170830. doi: 10.1038/nrm.2017.73. PubMed PMID: 28852220; PMCID: PMC5794010.

9. Min JN, Huang L, Zimonjic DB, Moskophidis D, Mivechi NF. Selective suppression of

lymphomas by functional loss of Hsf1 in a p53-deficient mouse model for spontaneous

tumors. Oncogene. 2007;26(35):5086-97. Epub 20070219. doi: 10.1038/sj.onc.1210317.

PubMed PMID: 17310987.

10. Dai C, Whitesell L, Rogers AB, Lindquist S. Heat shock factor 1 is a powerful multifaceted

modifier of carcinogenesis. Cell. 2007;130(6):1005-18. doi: 10.1016/j.cell.2007.07.020.

PubMed PMID: 17889646; PMCID: PMC2586609.

11. Meng L, Gabai VL, Sherman MY. Heat-shock transcription factor HSF1 has a critical role in

human epidermal growth factor receptor-2-induced cellular transformation and

tumorigenesis. Oncogene. 2010;29(37):5204-13. Epub 20100712. doi:

10.1038/onc.2010.277. PubMed PMID: 20622894; PMCID: PMC2940982.

12. Jin X, Moskophidis D, Mivechi NF. Heat shock transcription factor 1 is a key determinant of

HCC development by regulating hepatic steatosis and metabolic syndrome. Cell Metab.

2011;14(1):91-103. doi: 10.1016/j.cmet.2011.03.025. PubMed PMID: 21723507; PMCID:

PMC3214631.

13. Dai C, Santagata S, Tang Z, Shi J, Cao J, Kwon H, Bronson RT, Whitesell L, Lindquist S.

Loss of tumor suppressor NF1 activates HSF1 to promote carcinogenesis. J Clin Invest.

2012;122(10):3742-54. Epub 20120904. doi: 10.1172/JCI62727. PubMed PMID: 22945628;

PMCID: PMC3461912.

14. Kourtis N, Lazaris C, Hockemeyer K, Balandran JC, Jimenez AR, Mullenders J, Gong Y,

Trimarchi T, Bhatt K, Hu H, Shrestha L, Ambesi-Impiombato A, Kelliher M, Paietta E,

Chiosis G, Guzman ML, Ferrando AA, Tsirigos A, Aifantis I. Oncogenic hijacking of the

stress response machinery in T cell acute lymphoblastic leukemia. Nat Med.

2018;24(8):1157-66. Epub 20180723. doi: 10.1038/s41591-018-0105-8. PubMed PMID:

30038221; PMCID: PMC6082694.

15. Fujimoto M, Takii R, Matsumoto M, Okada M, Nakayama KI, Nakato R, Fujiki K, Shirahige

K, Nakai A. HSF1 phosphorylation establishes an active chromatin state via the TRRAP-

TIP60 complex and promotes tumorigenesis. Nat Commun. 2022;13(1):4355. Epub

20220729. doi: 10.1038/s41467-022-32034-4. PubMed PMID: 35906200; PMCID:

PMC9338313.

16. Heimberger T, Andrulis M, Riedel S, Stuhmer T, Schraud H, Beilhack A, Bumm T, Bogen

B, Einsele H, Bargou RC, Chatterjee M. The heat shock transcription factor 1 as a potential

new therapeutic target in multiple myeloma. Br J Haematol. 2013;160(4):465-76. Epub

20121218. doi: 10.1111/bjh.12164. PubMed PMID: 23252346.

17. Chuma M, Sakamoto N, Nakai A, Hige S, Nakanishi M, Natsuizaka M, Suda G, Sho T,

Hatanaka K, Matsuno Y, Yokoo H, Kamiyama T, Taketomi A, Fujii G, Tashiro K, Hikiba Y,

Fujimoto M, Asaka M, Maeda S. Heat shock factor 1 accelerates hepatocellular carcinoma

development by activating nuclear factor-kappaB/mitogen-activated protein kinase.

Carcinogenesis. 2014;35(2):272-81. Epub 20131015. doi: 10.1093/carcin/bgt343. PubMed

PMID: 24130164.

18. Fujimoto M, Takaki E, Takii R, Tan K, Prakasam R, Hayashida N, Iemura S, Natsume T,

Nakai A. RPA assists HSF1 access to nucleosomal DNA by recruiting histone chaperone

FACT. Mol Cell. 2012;48(2):182-94. Epub 20120830. doi: 10.1016/j.molcel.2012.07.026.

PubMed PMID: 22940245.

19. Solimini NL, Luo J, Elledge SJ. Non-oncogene addiction and the stress phenotype of cancer

cells. Cell. 2007;130(6):986-8. doi: 10.1016/j.cell.2007.09.007. PubMed PMID: 17889643.

20. Gutmann DH, Ferner RE, Listernick RH, Korf BR, Wolters PL, Johnson KJ.

Neurofibromatosis type 1. Nat Rev Dis Primers. 2017;3:17004. Epub 20170223. doi:

10.1038/nrdp.2017.4. PubMed PMID: 28230061.

21. Hirbe AC, Gutmann DH. Neurofibromatosis type 1: a multidisciplinary approach to care.

Lancet Neurol. 2014;13(8):834-43. doi: 10.1016/S1474-4422(14)70063-8. PubMed PMID:

25030515.

22. Cichowski K, Jacks T. NF1 tumor suppressor gene function: narrowing the GAP. Cell.

2001;104(4):593-604. doi: 10.1016/s0092-8674(01)00245-8. PubMed PMID: 11239415.

23. Grice GL, Nathan JA. The recognition of ubiquitinated proteins by the proteasome. Cell Mol

Life Sci. 2016;73(18):3497-506. Epub 20160502. doi: 10.1007/s00018-016-2255-5. PubMed

PMID: 27137187; PMCID: PMC4980412.

24. Knowles TP, Vendruscolo M, Dobson CM. The amyloid state and its association with protein

misfolding diseases. Nat Rev Mol Cell Biol. 2014;15(6):384-96. doi: 10.1038/nrm3810.

PubMed PMID: 24854788.

25. Yakupova EI, Bobyleva LG, Vikhlyantsev IM, Bobylev AG. Congo Red and amyloids:

history and relationship. Biosci Rep. 2019;39(1). Epub 20190115. doi:

10.1042/BSR20181415. PubMed PMID: 30567726; PMCID: PMC6331669.

26. Kayed R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, Glabe CG.

Common structure of soluble amyloid oligomers implies common mechanism of

pathogenesis. Science. 2003;300(5618):486-9. doi: 10.1126/science.1079469. PubMed

PMID: 12702875.

27. Kayed R, Head E, Sarsoza F, Saing T, Cotman CW, Necula M, Margol L, Wu J, Breydo L,

Thompson JL, Rasool S, Gurlo T, Butler P, Glabe CG. Fibril specific, conformation

dependent antibodies recognize a generic epitope common to amyloid fibrils and fibrillar

oligomers that is absent in prefibrillar oligomers. Mol Neurodegener. 2007;2:18. Epub

20070926. doi: 10.1186/1750-1326-2-18. PubMed PMID: 17897471; PMCID:

PMC2100048.

28. Heiser V, Scherzinger E, Boeddrich A, Nordhoff E, Lurz R, Schugardt N, Lehrach H,

Wanker EE. Inhibition of huntingtin fibrillogenesis by specific antibodies and small

molecules: implications for Huntington's disease therapy. Proc Natl Acad Sci U S A.

2000;97(12):6739-44. doi: 10.1073/pnas.110138997. PubMed PMID: 10829068; PMCID:

PMC18723.

29. Sanchez I, Mahlke C, Yuan J. Pivotal role of oligomerization in expanded polyglutamine

neurodegenerative disorders. Nature. 2003;421(6921):373-9. doi: 10.1038/nature01301.

PubMed PMID: 12540902.

30. Tang Z, Su KH, Xu M, Dai C. HSF1 physically neutralizes amyloid oligomers to empower

overgrowth and bestow neuroprotection. Sci Adv. 2020;6(46). Epub 20201111. doi:

10.1126/sciadv.abc6871. PubMed PMID: 33177089; PMCID: PMC7673739.

31. Christensen JH, Nielsen MN, Hansen J, Fuchtbauer A, Fuchtbauer EM, West M, Corydon

TJ, Gregersen N, Bross P. Inactivation of the hereditary spastic paraplegia-associated Hspd1

gene encoding the Hsp60 chaperone results in early embryonic lethality in mice. Cell Stress

Chaperones. 2010;15(6):851-63. doi: 10.1007/s12192-010-0194-x. PubMed PMID:

20393889; PMCID: PMC3024079.

32. Fan F, Duan Y, Yang F, Trexler C, Wang H, Huang L, Li Y, Tang H, Wang G, Fang X, Liu

J, Jia N, Chen J, Ouyang K. Deletion of heat shock protein 60 in adult mouse cardiomyocytes

perturbs mitochondrial protein homeostasis and causes heart failure. Cell Death Differ.

2020;27(2):587-600. Epub 20190617. doi: 10.1038/s41418-019-0374-x. PubMed PMID:

31209364; PMCID: PMC7205885.

33. Su KH, Cao J, Tang Z, Dai S, He Y, Sampson SB, Benjamin IJ, Dai C. HSF1 critically

attunes proteotoxic stress sensing by mTORC1 to combat stress and promote growth. Nat

Cell Biol. 2016;18(5):527-39. Epub 20160404. doi: 10.1038/ncb3335. PubMed PMID:

27043084; PMCID: PMC5341796.

34. Johnson GL, Nakamura K. The c-jun kinase/stress-activated pathway: regulation, function

and role in human disease. Biochim Biophys Acta. 2007;1773(8):1341-8. Epub 20070104.

doi: 10.1016/j.bbamcr.2006.12.009. PubMed PMID: 17306896; PMCID: PMC1995559.

35. Battaglioni S, Benjamin D, Walchli M, Maier T, Hall MN. mTOR substrate phosphorylation

in growth control. Cell. 2022;185(11):1814-36. Epub 20220516. doi:

10.1016/j.cell.2022.04.013. PubMed PMID: 35580586.

36. Kim DH, Sarbassov DD, Ali SM, King JE, Latek RR, Erdjument-Bromage H, Tempst P,

Sabatini DM. mTOR interacts with raptor to form a nutrient-sensitive complex that signals to

the cell growth machinery. Cell. 2002;110(2):163-75. doi: 10.1016/s0092-8674(02)00808-5.

PubMed PMID: 12150925.

37. Schmidt EK, Clavarino G, Ceppi M, Pierre P. SUnSET, a nonradioactive method to monitor

protein synthesis. Nat Methods. 2009;6(4):275-7. Epub 20090322. doi: 10.1038/nmeth.1314.

PubMed PMID: 19305406.

38. Zhang T, Inesta-Vaquera F, Niepel M, Zhang J, Ficarro SB, Machleidt T, Xie T, Marto JA,

Kim N, Sim T, Laughlin JD, Park H, LoGrasso PV, Patricelli M, Nomanbhoy TK, Sorger

PK, Alessi DR, Gray NS. Discovery of potent and selective covalent inhibitors of JNK.

Chem Biol. 2012;19(1):140-54. doi: 10.1016/j.chembiol.2011.11.010. PubMed PMID:

22284361; PMCID: PMC3270411.

39. Moerke NJ, Aktas H, Chen H, Cantel S, Reibarkh MY, Fahmy A, Gross JD, Degterev A,

Yuan J, Chorev M, Halperin JA, Wagner G. Small-molecule inhibition of the interaction

between the translation initiation factors eIF4E and eIF4G. Cell. 2007;128(2):257-67. doi:

10.1016/j.cell.2006.11.046. PubMed PMID: 17254965.

40. Tolcher A, Goldman J, Patnaik A, Papadopoulos KP, Westwood P, Kelly CS, Bumgardner

W, Sams L, Geeganage S, Wang T, Capen AR, Huang J, Joseph S, Miller J, Benhadji KA,

Brail LH, Rosen LS. A phase I trial of LY2584702 tosylate, a p70 S6 kinase inhibitor, in

patients with advanced solid tumours. Eur J Cancer. 2014;50(5):867-75. Epub 20140115.

doi: 10.1016/j.ejca.2013.11.039. PubMed PMID: 24440085.

41. Porter AG, Janicke RU. Emerging roles of caspase-3 in apoptosis. Cell Death Differ.

1999;6(2):99-104. doi: 10.1038/sj.cdd.4400476. PubMed PMID: 10200555.

42. Boulares AH, Yakovlev AG, Ivanova V, Stoica BA, Wang G, Iyer S, Smulson M. Role of

poly(ADP-ribose) polymerase (PARP) cleavage in apoptosis. Caspase 3-resistant PARP

mutant increases rates of apoptosis in transfected cells. J Biol Chem. 1999;274(33):22932-40.

doi: 10.1074/jbc.274.33.22932. PubMed PMID: 10438458.

43. Huang J, Manning BD. The TSC1-TSC2 complex: a molecular switchboard controlling cell

growth. Biochem J. 2008;412(2):179-90. doi: 10.1042/BJ20080281. PubMed PMID:

18466115; PMCID: PMC2735030.

44. Dong B, Jaeger AM, Hughes PF, Loiselle DR, Hauck JS, Fu Y, Haystead TA, Huang J,

Thiele DJ. Targeting therapy-resistant prostate cancer via a direct inhibitor of the human heat

shock transcription factor 1. Sci Transl Med. 2020;12(574). doi:

10.1126/scitranslmed.abb5647. PubMed PMID: 33328331; PMCID: PMC10571035.

45. Sengupta S, Giaime E, Narayan S, Hahm S, Howell J, O'Neill D, Vlasuk GP, Saiah E.

Discovery of NV-5138, the first selective Brain mTORC1 activator. Sci Rep.

2019;9(1):4107. Epub 20190311. doi: 10.1038/s41598-019-40693-5. PubMed PMID:

30858438; PMCID: PMC6412019.

46. Kato T, Pothula S, Liu RJ, Duman CH, Terwilliger R, Vlasuk GP, Saiah E, Hahm S, Duman

RS. Sestrin modulator NV-5138 produces rapid antidepressant effects via direct mTORC1

activation. J Clin Invest. 2019;129(6):2542-54. Epub 20190416. doi: 10.1172/JCI126859.

PubMed PMID: 30990795; PMCID: PMC6546461.

47. Wolfson RL, Chantranupong L, Saxton RA, Shen K, Scaria SM, Cantor JR, Sabatini DM.

Sestrin2 is a leucine sensor for the mTORC1 pathway. Science. 2016;351(6268):43-8. Epub

20151008. doi: 10.1126/science.aab2674. PubMed PMID: 26449471; PMCID:

PMC4698017.

48. Yoon YJ, Kim JA, Shin KD, Shin DS, Han YM, Lee YJ, Lee JS, Kwon BM, Han DC.

KRIBB11 inhibits HSP70 synthesis through inhibition of heat shock factor 1 function by

impairing the recruitment of positive transcription elongation factor b to the hsp70 promoter.

J Biol Chem. 2011;286(3):1737-47. Epub 20101115. doi: 10.1074/jbc.M110.179440.

PubMed PMID: 21078672; PMCID: PMC3023468.

49. Caserta TM, Smith AN, Gultice AD, Reedy MA, Brown TL. Q-VD-OPh, a broad spectrum

caspase inhibitor with potent antiapoptotic properties. Apoptosis. 2003;8(4):345-52. doi:

10.1023/a:1024116916932. PubMed PMID: 12815277.

50. Chauvier D, Ankri S, Charriaut-Marlangue C, Casimir R, Jacotot E. Broad-spectrum caspase

inhibitors: from myth to reality? Cell Death Differ. 2007;14(2):387-91. Epub 20060929. doi:

10.1038/sj.cdd.4402044. PubMed PMID: 17008913.

51. Miao EA, Rajan JV, Aderem A. Caspase-1-induced pyroptotic cell death. Immunol Rev.

2011;243(1):206-14. doi: 10.1111/j.1600-065X.2011.01044.x. PubMed PMID: 21884178;

PMCID: PMC3609431.

52. Shan B, Pan H, Najafov A, Yuan J. Necroptosis in development and diseases. Genes Dev.

2018;32(5-6):327-40. doi: 10.1101/gad.312561.118. PubMed PMID: 29593066; PMCID:

PMC5900707.

53. Chen X, Comish PB, Tang D, Kang R. Characteristics and Biomarkers of Ferroptosis. Front

Cell Dev Biol. 2021;9:637162. Epub 20210121. doi: 10.3389/fcell.2021.637162. PubMed

PMID: 33553189; PMCID: PMC7859349.

54. Stockwell BR. Ferroptosis turns 10: Emerging mechanisms, physiological functions, and

therapeutic applications. Cell. 2022;185(14):2401-21. doi: 10.1016/j.cell.2022.06.003.

PubMed PMID: 35803244; PMCID: PMC9273022.

55. Yoshii SR, Mizushima N. Monitoring and Measuring Autophagy. Int J Mol Sci. 2017;18(9).

Epub 20170828. doi: 10.3390/ijms18091865. PubMed PMID: 28846632; PMCID:

PMC5618514.

56. Blommaart EF, Krause U, Schellens JP, Vreeling-Sindelarova H, Meijer AJ. The

phosphatidylinositol 3-kinase inhibitors wortmannin and LY294002 inhibit autophagy in

isolated rat hepatocytes. Eur J Biochem. 1997;243(1-2):240-6. doi: 10.1111/j.1432-

1033.1997.0240a.x. PubMed PMID: 9030745.

57. Schubert U, Anton LC, Gibbs J, Norbury CC, Yewdell JW, Bennink JR. Rapid degradation

of a large fraction of newly synthesized proteins by proteasomes. Nature.

2000;404(6779):770-4. doi: 10.1038/35008096. PubMed PMID: 10783891.

58. Kim LC, Cook RS, Chen J. mTORC1 and mTORC2 in cancer and the tumor

microenvironment. Oncogene. 2017;36(16):2191-201. Epub 20161017. doi:

10.1038/onc.2016.363. PubMed PMID: 27748764; PMCID: PMC5393956.

59. Zou Z, Tao T, Li H, Zhu X. mTOR signaling pathway and mTOR inhibitors in cancer:

progress and challenges. Cell Biosci. 2020;10:31. Epub 20200310. doi: 10.1186/s13578-020-

00396-1. PubMed PMID: 32175074; PMCID: PMC7063815.

60. Goloudina AR, Demidov ON, Garrido C. Inhibition of HSP70: a challenging anti-cancer

strategy. Cancer Lett. 2012;325(2):117-24. Epub 20120628. doi:

10.1016/j.canlet.2012.06.003. PubMed PMID: 22750096.

61. Jhaveri K, Taldone T, Modi S, Chiosis G. Advances in the clinical development of heat

shock protein 90 (Hsp90) inhibitors in cancers. Biochim Biophys Acta. 2012;1823(3):742-

55. Epub 20111029. doi: 10.1016/j.bbamcr.2011.10.008. PubMed PMID: 22062686;

PMCID: PMC3288123.

62. Pasqua AE, Sharp SY, Chessum NEA, Hayes A, Pellegrino L, Tucker MJ, Miah A, Wilding

B, Evans LE, Rye CS, Mok NY, Liu M, Henley AT, Gowan S, De Billy E, Te Poele R,

Powers M, Eccles SA, Clarke PA, Raynaud FI, Workman P, Jones K, Cheeseman MD. HSF1

Pathway Inhibitor Clinical Candidate (CCT361814/NXP800) Developed from a Phenotypic

Screen as a Potential Treatment for Refractory Ovarian Cancer and Other Malignancies. J

Med Chem. 2023;66(8):5907-36. Epub 20230405. doi: 10.1021/acs.jmedchem.3c00156.

PubMed PMID: 37017629; PMCID: PMC10150365.

63. Ali ES, Mitra K, Akter S, Ramproshad S, Mondal B, Khan IN, Islam MT, Sharifi-Rad J,

Calina D, Cho WC. Recent advances and limitations of mTOR inhibitors in the treatment of

cancer. Cancer Cell Int. 2022;22(1):284. Epub 20220915. doi: 10.1186/s12935-022-02706-8.

PubMed PMID: 36109789; PMCID: PMC9476305.

64. Wood KC, Gutkind JS. Challenges and Emerging Opportunities for Targeting mTOR in

Cancer. Cancer Res. 2022;82(21):3884-7. doi: 10.1158/0008-5472.CAN-22-0602. PubMed

PMID: 36321262; PMCID: PMC9634680.

65. Thomson AW, Turnquist HR, Raimondi G. Immunoregulatory functions of mTOR

inhibition. Nat Rev Immunol. 2009;9(5):324-37. doi: 10.1038/nri2546. PubMed PMID:

19390566; PMCID: PMC2847476.

Figure 1

FIGURE LEGENDS:

Figure 1: HSF1 is required to suppress proteomic instability and amyloidogenesis in

MPNST cells. (A) Violin plots of *HSF1* expression in human cancer tissues showing *NF1*

deletion (median and IQR, n=6424, 1598, or 88, Kruskal-Wallis test). Data are generated by the

TGCA Research Network (https://www.cancer.gov/tcga). (B) Kaplan-Meier survival curves of

patients whose tumors harbor *NF1* deletion (n=420 or 419, Log-rank test). Data are generated by

the TGCA Research Network (https://www.cancer.gov/tcga). (C) and (D) The impacts of *HSF1*

depletion on protein polyubiquitination in both 90-8TL cells and primary human Schwann cells.

Cells were transduced with lentiviral shRNAs for 4 days and global protein Lys48-specific

ubiquitination was detected by immunoblotting in both detergent-soluble and -insoluble fractions

of whole cell lysates (C). Protein ubiquitination was quantitated using Fiji imaging software (D)

(mean ± SD, n=3 independent experiments, Two-way ANOVA). (E) and (F) Congo Red (CR)

staining of 90-8TL cells and primary human Schwann cells with and without *HSF1* depletion,

analyzed by flow cytometry (E). Quantitation of CR staining using the median fluorescence

intensity (F) (mean ± SD, n=3 independent experiments, One-way ANOVA). (G)-(J)

Quantitation of soluble amyloid oligomers and insoluble amyloid fibrils by ELISA using A11

and OC antibodies, respectively (mean ± SD, n=3 independent experiments, One-way ANOVA).

1000

1001

Figure 2

Figure 2: HSF1 counters the tumor-suppressive effects of amyloids in MPNST cells.

(A) and (B) The growth curves of 90-8TL and primary human Schwann cells with and without

HSF1 depletion and 20 μM CR treatment (mean ± SD, n=3 independent experiments, Two-way

ANOVA). (C) Co-IP of amyloid oligomers (AOs) with HSF1 in 90-8TL cells. (D)-(F) Detection

of AO–HSF1 interactions by PLA flow cytometry in 90-8TL and S462 cells. Quantitation of the

PLA signals using the median fluorescence intensity (mean ± SD, n=3 or 4 independent

experiments, Student's t test). (G) and (H) Detection of AO–HSP60 interactions by PLA flow

cytometry in S462 cells following *HSF1* depletion (G). Quantitation of the PLA signals using the

median fluorescence intensity (H) (mean ± SD, n=3 independent experiments, Student's t test).

(I) Schematic depiction of AO neuralization by HSF1 and HSP60 attack by AO following *HSF1*

depletion in mitochondria.

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

Figure 3

Figure 3: JNK senses *HSF1* deficiency to repress mTORC1 and protein translation in MPNST cells. (A) Detection of JNK activation and mTORC1 inhibition following *HSF1* depletion in S462 and primary human Schwann cells by immunoblotting. (B) Quantitation of (A) using Fiji imaging software (mean ± SD, n=3 independent experiments, One-way ANOVA). (C) Detection of HSF1-JNK, JNK-RAPTOR, and mTOR-RAPTOR interactions by PLA in S462 cells with and without *HSF1* depletion. (D) and (E) Detection and quantitation of JNK-RAPTOR and mTOR-RAPTOR interactions by PLA flow cytometry in 90-8TL cells (mean ± SD, n=3 independent experiments, One-way ANOVA). (F) Measurement of protein translation rates in 90-8TL and primary human Schwann cells with and without *HSF1* depletion by puromycin labeling, analyzed by flow cytometry. Cells were treated with 1 μM JNK-IN-8 for 24 hrs before labeling with 100 nM 6-FAM-dc-puromycin for 30 min (mean ± SD, n=3 independent experiments, Two-way ANOVA). (G) Heatmap of protein translation rates measured by puromycin labeling in diverse human cancer cell lines with and without *HSF1* depletion. (H) Schematic depiction of regulation of protein translation by HSF1.

Figure 4

Figure 4: Protein translation is causally related to amyloidogenesis in MPNST cells.

(A) and (B) Detection of mTORC1 stimulation following JNK inhibition in 90-8TL and immortalized human Schwann cells by immunoblotting. Cells were treated with JNK-IN-8 for 2 days. (C) and (D) Quantitation of soluble AOs and insoluble AFs by ELISA in 90-8TL and human Schwann cells with and without 3 μM JNK-IN-8 treatment for 3 days (mean ± SD, n=3 independent experiments, Student's t test). (E) Detection of LY2584702-mediated S6K inhibition in 90-8TL cells by immunoblotting. Cells were treated with 20 μM LY2584702 for 2 days. (F) Quantitation of soluble AOs and insoluble AFs by ELISA in 90-8TL cells with and without 20 μM LY2584702 treatment for 2 days (mean ± SD, n=3 independent experiments, Student's t test). (G) CR staining of 90-8TL and immortalized human Schwann cells with and without JNK-IN-8 and 4EGI-1 treatment, detected by flow cytometry. (H) Quantitation of (G) using the median fluorescence intensity (mean ± SD, n=3 independent experiments, One-way ANOVA). (I) Quantitation of amyloids by CR staining in 90-8TL cells with and without *HSF1* depletion and 3 μM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (J) Quantitation of caspase 3 cleavage by ELISA in 90-8TL cells with and without *HSF1* depletion and 3 μM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (K) Quantitation of amyloids by CR staining in immortalized human Schwann cells with and without *HSF1* depletion and 3 μM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (L) Quantitation of caspase 3 cleavage by ELISA in immortalized human Schwann cells with and without *HSF1* depletion and 3 μM JNK-IN-8 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (M) Schematic depiction of the adaptation of MPNST cells to *HSF1* depletion by activating JNK to suppress mTORC1 and protein translation, thereby alleviating amyloidogenesis and cytotoxicity.

Figure 5

 Figure 5: mTORC1 stimulation aggravates the amyloidogenesis and cytotoxicity elicited by

 HSF1 inhibition. (A) Immunoblotting detection of JNK activation, mTORC1 activity, and

 protein polyubiquitination in S462 cells following stable *TSC2* knockdown and/or HSF1

 inhibition by 10 μM DTHIB for 3 days. (B) Quantitation of soluble AOs by flow cytometry

 using A11 antibody staining in S462 cells with and without *TSC2* knockdown and 10 μM

 DTHIB treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). (C)

 Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

 without stable *TSC2* knockdown and DTHIB (10 μM) or combined DTHIB and LY2584702 (20

 μM) treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). Cells were pre-

 treated with LY2584702 for 1 day followed by DTHIB treatment for another 3 days. (D)

 Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

 without stable *TSC2* knockdown and DTHIB (10 μM) or combined DTHIB and CR (30 μM)

 treatment (mean ± SD, n=3 independent experiments, Two-way ANOVA). Cells were pre-

 treated with CR for 1 day followed by DTHIB treatment for another 3 days. (E) Quantitation of

 soluble AOs by flow cytometry using A11 antibody staining in S462 cells with and without 500

 μM NV-5138 stimulation and 10 μM DTHIB treatment (mean ± SD, n=3 independent

 experiments, One-way ANOVA). (F) Quantitation of cytotoxicity by flow cytometry using Live-

 or-Dye stains in S462 cells with and without 500 μM NV-5138 stimulation and 10 μM DTHIB

 or combined DTHIB and CR treatment (mean ± SD, n=5 independent experiments, One-way

 ANOVA). (G) Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in

 immortalized human Schwann cells with and without 500 μM NV-5138 stimulation and 10 μM

 DTHIB treatment (mean ± SD, n=3 independent experiments, One-way ANOVA). (H)

 Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

without 500 μM L-leucine stimulation and 10 μM DTHIB treatment (mean ± SD, n=4

independent experiments, One-way ANOVA). (I) Quantitation of cytotoxicity by flow cytometry

using Live-or-Dye stains in immortalized human Schwann cells with and without 500 μM L-

leucine stimulation and 10 μM DTHIB treatment (mean ± SD, n=3 independent experiments,

One-way ANOVA). (J) Quantitation of soluble AOs by ELISA in S462 cells treated with and

without 800 μM L-leucine and 20 μM KRIBB11 or 30 μM CR (mean ± SD, n=3 independent

experiments, One-way ANOVA). (K) Quantitation of cytotoxicity by flow cytometry using Live-

or-Dye stains in S462 cells treated as above (mean ± SD, n=4 independent experiments, One-

way ANOVA). (L) Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in

S462 cells stably expressing inducible shRNAs with and without L-leucine stimulation. Cells

were first cultured with or without 1.2 mM L-leucine for 2 days, followed by addition of 20

ng/ml doxycycline for 7 days (mean ± SD, n=3 independent experiments, One-way ANOVA).

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

Figure 6

Figure 6: Concurrent HSF1 inhibition and mTORC1 stimulation instigate widespread non-

apoptotic cell death. (A) and (B) Quantitation of cytotoxicity by flow cytometry using Live-or-

Dye stains combined with cleaved caspase 3 (Asp175) antibody staining. S462 cells with and

without stable *TSC2* knockdown were treated with DMSO, 10 µM DTHIB, or combined DTHIB

and 30 µM Q-VD-OPh (mean ± SD, n=3 independent experiments, Two-way ANOVA). (C)

Quantitation of cytotoxicity by flow cytometry using Live-or-Dye stains in S462 cells with and

without stable *TSC2* knockdown. Cells were treated with 10 µM DTHIB alone or co-treated with

10 µM DTHIB and 30 µM Necrostatin-1 or 20 µM Liproxstatin-1 for 3 days (mean ± SD, n=3

independent experiments, Two-way ANOVA). (D) Immunoblotting detection of ferroptosis

markers in S462 cells with and without stable *TSC2* knockdown treated with 10 µM DTHIB

alone or co-treated with 10 µM DTHIB and 20 µM liproxstatin-1 for 3 days. Erastin was

included as a positive control to induce canonical ferroptosis. (E) Immunoblotting detection of

autophagy markers in S462 cells with stable *TSC2* knockdown treated with 10 µM DTHIB alone

or co-treated with 10 µM DTHIB and 3 µM wortmannin for 3 days. Rapamycin was included as

a positive control to induce autophagy. (F) Quantitation of cytotoxicity by flow cytometry using

Live-or-Dye stains in S462 cells with and without stable *TSC2* knockdown. Cells were treated

with 10 µM DTHIB alone or co-treated with 10 µM DTHIB and 3 µM wortmannin for 3 days

(mean ± SD, n=3 independent experiments, Two-way ANOVA). (G) Schematic depiction of

instigation of cell death by severe proteomic imbalance, owing to simultaneous mTORC1

stimulation (**1**) and HSF1 inhibition (**2**). In cancer cells, constitutive HSF1 activation

provides extra chaperoning capacity to cope with elevated protein misfolding, partly due to

enhanced protein synthesis and widespread genetic mutations. Nevertheless, amyloids still

emerge, although at low levels. Importantly, HSF1 can neutralize highly toxic amyloid

oligomers, averting lethal consequences. By contrast, HSF1 inhibition diminishes chaperoning

capacity, insufficient to counterbalance the robust protein translation. mTORC1 stimulation

further aggravates this proteomic imbalance, which, in turn, promotes amyloidogenesis and

proteomic catastrophe. In consequence, the amounts of amyloid oligomers surpass the

neutralizing capacity of HSF1, leading to cell death; nonetheless, it remains unclear how this

proteomic catastrophe-induced death (PCID) occurs.

1146

1147

1148

1149

1150

1151

1152

1153

1154

1155

1156

1157

1158

1159

1160

1161

1162

Figure 7: Concurrent HSF1 inhibition and mTORC1 stimulation suppress *in vivo* tumor growth. (A) Combined KRIBB11 treatment and L-leucine stimulation impaired the *in vivo* growth of xenografted S462 human MPNST cells in NIH-III nude mice (mean ± SEM, n=8 mice per group, $\log_2$ transformation for Two-way ANOVA). (B) Combined KRIBB11 treatment and L-leucine stimulation impeded the *in vivo* growth of xenografted B16-F10 murine melanoma cells in C57BL/6J mice (mean ± SEM, n=8 mice per group, $\log_2$ transformation for Two-way ANOVA). (C) Quantitation of phospho-S6RP by immunoblotting in treated melanomas. Images show representative 3 tumors each group (mean ± SEM, n=5 tumors per group, One-way ANOVA). (D) Quantitation of *Hsp72* and *Hsp25* mRNAs in treated melanomas by RT-qPCR (mean ± SD, n=5 tumors per group, One-way ANOVA). (E) Quantitation of soluble AOs in treated B16F10 melanomas by ELISA ((mean ± SD, n=5 tumors per group, One-way ANOVA). (F) and (G) Quantitation of soluble AOs in brains and spleens from treated C57BL/6J mice by ELISA (mean ± SD, n=5 brains or spleens per group, One-way ANOVA). (H) Schematic depiction of the concept of provoking proteomic catastrophe to combat malignancy. On the one hand, in cancer cells, mTORC1 is inevitably activated to stimulate protein translation, markedly augmenting protein quantity. On the other hand, the extra chaperoning capacity governed by HSF1, albeit dispensable for normal life, becomes necessary to ensure sufficient protein quality in cancer cells, thereby counterbalancing augmented protein quantity and suppressing proteomic instability. Thus, proteomic balance promotes malignant growth. By contrast, disrupting proteomic balance, through HSF1 inhibition, is sufficient to elicit proteomic instability and tumor suppression. However, simultaneous mTORC1 stimulation can remarkably drive proteomic imbalance, causing proteomic catastrophe that leads to tumor suppression.