Photoreceptor loss does not recruit neutrophils despite strong microglial activation

Derek Power,[1,2] Justin Elstrott,[3] Jesse Schallek[1,2,4]

Corresponding author: Jesse Schallek, Flaum Eye Institute, University of Rochester, Rochester, NY 14642, USA

Email: jschall3@ur.rochester.edu

Phone number: +1-585-273-4848

[1]Center for Visual Science, University of Rochester, Rochester, NY 14627, USA

[2]Flaum Eye Institute, University of Rochester, Rochester, NY 14642, USA

[3]Department of Translational Imaging, Genentech Inc., South San Francisco, CA 94080, USA

[4]Department of Neuroscience, University of Rochester, Rochester, NY 14642, USA

Impact statement: Systemic neutrophils do not respond to laser-induced photoreceptor damage despite the robust response from resident microglia

Financial support: Research was supported by the National Eye Institute of the National Institutes of Health under Award No. R01 EY028293, and P30 EY001319. Work was also supported by a collaborative grant from Genentech Inc, a Career Development Award and Career Advancement Award (Schallek) and Unrestricted Grant to the University of Rochester Department of Ophthalmology from Research to Prevent Blindness, New York, New York and the Dana Foundation David Mahoney Neuroimaging Award (Schallek)

Conflict of Interest statement: Portions of this work were funded by a collaborative grant from Genentech, Inc. (Elstrott) to examine the extent to which immune cells could be studied in the deep retina using adaptive optics. Schallek also has six patents held through the University of Rochester on adaptive optics technology.

Data availability statement: All data collected (~3 TB) is archived at the University of Rochester and is available upon request.

Abstract:

In response to central nervous system (CNS) injury, tissue resident immune cells such as microglia and circulating systemic neutrophils are often first responders. The degree to which these cells interact in response to CNS damage is poorly understood, and even less so, in the neural retina which poses a challenge for high resolution imaging in vivo. In this study, we deploy fluorescence adaptive optics scanning light ophthalmoscopy (AOSLO) to study fluorescent microglia and neutrophils in mice. We simultaneously track immune cell dynamics using label-free phase-contrast AOSLO at micron-level resolution. Retinal lesions were induced with 488 nm light focused onto photoreceptor (PR) outer segments. These lesions focally ablated PRs, with minimal collateral damage to cells above and below the plane of focus. We used in vivo (AOSLO, SLO and OCT) imaging to reveal the natural history of the microglial and neutrophil response from minutes-to-months after injury. While microglia showed dynamic and progressive immune response with cells migrating into the injury locus within 1-day after injury, neutrophils were not recruited despite close proximity to vessels carrying neutrophils only microns away. Post-mortem confocal microscopy confirmed in vivo findings. This work illustrates that microglial activation does not recruit neutrophils in response to acute, focal loss of PRs, a condition encountered in many retinal diseases.

Introduction

In the mammalian retina, a rapid and coordinated immune response to infection

or injury is important for maintaining tissue homeostasis. This is especially critical

in the eye since mature retinal neurons do not typically regenerate, resulting in

long-term functional losses for the host.[1,2] The retina is considered immune-

privileged and is equipped with a resident population of innate immune cells

including microglia.[3,4] In healthy retina, microglia are distributed primarily in the

inner retina, residing within nerve fiber layer (NFL), inner plexiform layer (IPL)

and outer plexiform layers (OPL).[5,6], generally avoiding the nuclear layers.

Microglia tile the retina and like their counterparts in the brain, exhibit long, thin

processes that continually probe the neuro-glial microenvironment.[4,7,8]

In addition to phagocytosing debris[9,10], regulating synaptic

maintenance[11,12] and removing dead tissue[13,14], microglia can secrete

chemokines to recruit other leukocytes to help fight infection and repair damaged

tissue.[4,15–17] For many injuries, one of the first systemic responders recruited and

activated by microglia are neutrophils.[18,19] Neutrophils comprise a large fraction

of leukocytes in murine blood (7.4 – 33.9%, "young" C57BL/6J mice[20]). They

assist in maintaining tissue homeostasis by neutralizing foreign agents,

regulating the immune response, and phagocytosing dead tissue[21,22]. Under

inflammatory conditions, the spatiotemporal interplay between microglia and

neutrophils is poorly understood. A missed window of interaction is highly

problematic in histological study where a single time point reveals a snapshot of

the temporally complex immune response, which changes dynamically over time.

Here, we use in vivo imaging to overcome these constraints.

Documenting immune cell interactions in the retina over time has been

challenged by insufficient resolution and contrast to visualize single cells in the

living eye. The microscopic size of immune cells requires exceptional resolution

for detection. Recently, advances in AOSLO imaging have provided micron-level

resolution and enhanced contrast for imaging individual immune cells in the

retina and without requiring extrinsic dyes[7,23]. AOSLO provides multi-modal

information from confocal reflectance, phase-contrast and fluorescence

modalities, which can reveal a variety of cell types simultaneously in the living

eye. Here, we used confocal AOSLO to track changes in reflectance at cellular

scale. Phase-contrast AOSLO provides detail on highly translucent retinal

structures such as vascular wall, single blood cells[27–29], PR somata[30], and is well-

suited to image resident and systemic immune cells.[7,23] Fluorescence AOSLO

provides the ability to study fluorescently-labeled cells[25,31,32] and exogenous

dyes[27,33] throughout the living retina. These modalities used in combination have

recently provided detailed images of the retinal response to a model of human

uveitis.[23,34] Together, these innovations now provide a platform to visualize, for

the first time, the dynamic interplay between many immune cell types, each with

a unique role in tissue inflammation. We combine these innovative modalities

with conventional histology as well as commercial OCT to reveal the progressive

nature of the cellular response to acute retinal injury.

Here, we ask the question: "To what extent do microglia/neutrophils

respond to acute neural loss in the retina?" To begin unraveling the complexities

in this response, we deploy a deep retinal laser ablation model. Using AOSLO,

we track and characterize the changes in microglia, neutrophils and retinal

structure within hours, days and months after acute laser exposure.

Methods:

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, strain background (*Mus musculus*)	C57BL/6J	The Jackson Laboratory	Strain #: 000664	
strain, strain background (*Mus musculus*)	CX3CR1-GFP	The Jackson Laboratory	Strain #: 005582	
strain, strain background (*Mus musculus*)	Catchup	Laboratory of M. Gunzer	C57BL/6-*Ly6g*(tm2621(CretdTomato)Arte	(Hasenberg *et al.* 2015) [35]
antibody	Ly-6G-647	Biolegend	Cat #: 127610 RRID:AB_1134159	IF (1:200)
other	DAPI stain	Cell Signaling Technology	Cat #: 4083	1:500 of 10 mg/ml stock
chemical compound, drug	Lipopolysaccharide (LPS)	Sigma-Aldrich	Cat #: L4391	1 ng (1 μl) delivered intravitreally

software, algorithm	Cell counter plugin	Imagej FIJI	Cell counter plugin	Author: Kurt De Vos, Imagej version 1.53q

Mice

All experiments herein were approved by the University Committee on Animal

Resources (protocol #: UCAR-2010-052E) and according to the Association for

Research in Vision and Ophthalmology statement for the Use of Animals in

Ophthalmic and Vision Research as well as institutional approvals by the

University of Rochester. C57BL/6J (#000664, Jackson Labs, Bar Harbor, ME)

mice were used to track the retinal phenotype after laser exposure.

Heterozygous CX3CR1-GFP (#005582, Jackson Labs) mice were used to track

GFP-expressing microglia. Mice with heterozygous transgenic expression of

tdTomato in neutrophils ("Catchup" mice) were provided by the foundry lab of M.

Gunzer.[35] 18 mice (10 male, 8 female) in total were used for *in vivo* imaging. 10

additional mice (5 male, 5 female) were used for *ex vivo* histology. Age for all

mice used for this work were post-natal weeks 6 - 24.

Preparation for *in vivo* imaging

Mice were anesthetized with intra-perotineal injection of ketamine (100 mg/kg)

and xylazine (10 mg/kg). The pupil was dilated with 1% tropicamide (Sandoz,

Basel, Switzerland) and 2.5% phenylephrine (Akorn, Lake Forest, IL). A custom

contact lens (1.5 mm base curve, 3.2 mm diameter, +10 diopter power,

Advanced Vision Technologies, Lakewood, CO) was fitted to the eye. For a

subset of experiments, 50 mg/kg fluorescein (AK-FLUOR, Akorn, Decatur, IL)

was administered by intra-peritoneal injection to confirm vascular integrity and

perfusion status after injury. During AOSLO imaging, anesthesia was

supplemented with 1% (v/v) isoflurane in oxygen and mice were maintained at

37°C via electric heat pad. Eye hydration was maintained throughout imaging

with regular application of saline eye drops (Refresh tears, Allergan, Sydney,

Australia) and lubricating eye gel (Genteal, Alcon Laboratories Inc. Fort Worth,

TX). Mice were placed in a positioning frame with 6-degrees of freedom to allow

for stable animal positioning and aid in retinal navigation relative to the beam of

the AOSLO and OCT.

***In vivo* AOSLO imaging**

Four light sources were used for AOSLO imaging. A 904 nm diode source (12

μW, Qphotonics, Ann Harbor, MI) was used for wavefront sensing. A second

796 nm superluminescent diode (196 μW, Superlum, Cork, Ireland) was used for

reflectance imaging, including confocal and phase contrast modes.[27,31] A third

488 nm light source (56 μW, Toptica Photonics, Farmington, NY) was used to

visualize GFP-positive microglia in CX3CR1-GFP mice. A fourth 561 nm light

source (95 μW, Toptica Photonics) was used to visualize tdTomato-positive

neutrophils in Catchup mice. All light sources were fiber-coupled and axially

combined through the AOSLO system[31]. Fast (15.4 kHz) and slow (25 Hz)

scanners create a raster scan pattern, which is relayed through a series of afocal

telescopes to and from the eye. A Shack-Hartmann wavefront sensor (consisting

of a lenslet array and a Rolera XR camera, QImaging, Surrey, Canada)

measures the aberrations of the eye and a deformable mirror (ALPAO,

Montbonnot-Saint-Martin, France) provided the wavefront correction. Reflected

796 nm light was collected with a photomultiplier tube (H7422-50, Hamamatsu

Photonics, Hamamatsu, Japan). All confocal reflectance images were captured

with a 30 µm pinhole (1.3 Airy Disc Diameters, ADD). Phase-contrast was

achieved by displacing the pinhole relative to the principal axis of the detection

plane as previously described.[25] Fluorescence was captured with a

photomultiplier tube (H7422-40, Hamamatsu) either coupled with a 520Δ35

band-pass filter (FF01-520/35-25, Semrock, Rochester, NY) for GFP emission, or

a 630Δ92 band-pass filter (FF01-630/92-25, Semrock) for tdTomato emission. All

fluorescent images were captured with a confocal 50 µm pinhole (2.1 ADD).

Image field sizes were either 4.98° x 3.95° or 2.39° x 1.94°. NIR and visible

imaging channels were made coplanar, compensating for longitudinal chromatic

aberration by independently focusing each light source onto the same axial

structure in the retina. Through-focus stacks were acquired by sequentially

changing the focus from NFL to the PR outer segments by using the defocus

term on the custom adaptive optics control software.

As described previously, red blood cell (RBC) imaging was achieved by combining phase-contrast imaging with a strategy to arrest the slow galvanometer scanner and let the resonant scanner project a single "line" (0.71° scan angle) on the retina. This enabled RBC flux imaging by positioning this line orthogonal to the direction of flow for single capillaries. As blood cells moved through capillaries, they were "self-scanned" producing images of RBCs in space/time[27].

PR laser damage model and post-injury time points for imaging

488 nm light (Continuous wave Laser diode, ±4 nm bandwidth, Toptica Photonics) was used to create an acute laser injury. 785 µW of 488 nm light was projected through the AOSLO and focused onto the PR outer segments (Figure 1a) for 3 minutes in a single line on the retina subtending 24 x 1 µm to concentrate the power to a small region. Laser injuries were placed between 5-15° from the optic disc. To avoid absorption confounds, we refrained from placing lesions beneath large retinal vessels. For experiments examining neutrophil involvement, lesions were placed <100 µm away from retinal veins and within microns of capillaries to increase the chance of extravasation through these preferred pathways in the retina[36–38]. As many as four such lesions were placed per retina. This protocol produced a hyper-reflective phenotype in the >40 locations across 28 mice. In rare cases, the exposure yielded no hyper-reflective lesion and were often in mice with high retinal motion, where the light dosage

was spread over a larger retinal area. These locations were not included in the

in-vivo or histological analysis.

Throughout this work, we assessed the effects of the laser injury for the

following time points: baseline/control, 1 day (defined as 18 – 28 hours post-

damage), 3 days, 7 days and 2 months.

***In vivo* SLO, and OCT imaging**

To confirm global ocular health and changes imparted by the laser damage, a

commercial Heidelberg Spectralis system (Heidelberg, Germany) was used to

acquire scanning laser ophthalmoscopy (SLO) and optical coherence

tomography (OCT) images. 30° and 55° fields were used for SLO acquisitions.

The 30° field was used for OCT acquisitions. For some experiments, fluorescein

angiography was captured by imaging the retina within 10 minutes of fluorescein

administration (details above). The fluorescence mode of the SLO also enabled

wide-field images of GFP positive microglia.

OCT was acquired with the following settings: ART 16 (automatic real-time

tracking) was used to average B-scans and improve signal. To capture several

damage locations in a single field, we used a coarse scan area (61 B-scans, 1.02

x 0.85 mm). A dense 3D data cube was also captured (49 B-scans, 513 x 171

µm) to provide detailed information regarding the axial nature of the laser

damage. "Follow-up" mode, which allows the HRA software to return to the same

retinal location, was used whenever possible. To reveal the cross-sectional

profile for each lesion, several adjacent B-scans were spatially averaged (~30

μm).

Preparation for *ex vivo* imaging

Mice were euthanized by CO_2 asphyxiation followed by cervical dislocation.

Within 5 minutes of asphyxiation, eyes were enucleated and placed in 4%

paraformaldehyde (PFA, diluted from: #15714-S, Electron Microscopy Sciences,

Hatfield, Pennsylvania) in 1x phosphate buffered saline (PBS, #806552, Sigma-

Aldrich, St. Louis, MO) for one hour at room temperature. Eyes were dissected to

remove the cornea, lens and vitreous. Each eye cup was placed in one well of a

24-well plate containing 0.5 ml of 0.8% PFA and left overnight at 4°C. The retina

was separated from the retinal pigmented epithelium/choroid with attention to

preserve inner retinal layers "up" orientation. If not applying antibody, the tissue

was directly flat mounted (see below). For antibody staining, the retina was

placed in 1x BD perm/wash buffer (#554723, BD Biosciences, Franklin Lakes,

NJ) with 5% donkey serum (#D9663, Sigma-Aldrich) diluted in PBS for overnight

incubation at room temperature with gentle shaking. The following was

performed in the dark. Ly-6G-647 antibody (1:200, #127610, RRID:AB_1134159,

BioLegend, San Diego, CA) and DAPI (1:500 of 10 mg/ml stock, #4083, Cell

Signaling Technology, Danvers, MA) were diluted in 1x perm/wash buffer and

retinas were incubated for 3 days at room temperature with gentle shaking.

Retinas were washed with PBS three times over three hours. The retina was cut

into 4 radially symmetrical petals, flat-mounted on a glass slide in Vectashield

175 mounting buffer (H-1000-10, Vector labs, Newark, CA) with a #1.5 cover slip

176 (#260406, Ted Pella Inc., Redding, CA), sealed with nail polish and stored at 4°C

177 until imaged.

178 **Endotoxin-Induced Uveitis protocol**

179 To serve as a positive control and show evidence of known neutrophil invasion,

180 we adopted the endotoxin-induced uveitis (EIU) model to confirm that fluorescent

181 Catchup neutrophils could be observed in vivo with AOSLO. This was performed

182 in two mice. The EIU model has been described previously.[34,39] Briefly, we

183 performed intravitreal injections of lipopolysaccharide (LPS, #L4391, Sigma-

184 Aldrich). Mice were anesthetized with ketamine and xylazine. A 34 gauge

185 Hamilton needle was used to deliver 1 μL (1 ng) of LPS diluted in PBS into the

186 vitreous posterior to the limbus. 1 day post-LPS injection, mice were either

187 imaged with AOSLO (Catchup mice) or collected for *ex vivo* histology (Ly-6G-647

188 stained C57BL/6J mice) to confirm fluorescent neutrophil presence in the neural

189 parenchyma.

190 *Ex vivo* **confocal imaging**

191 Whole-mount retinas were imaged with a Nikon A1 confocal microscope

192 (Melville, NY). DAPI (405 nm ex, 441Δ66 nm em), CX3CR1-GFP (488 nm ex,

193 525Δ50 nm em), and Catchup/anti-Ly-6G-647 (635 nm ex, 665Δ50 nm em) were

194 simultaneously imaged. Z-stacks (0.1 or 0.5 μm step size) at control or laser-

195 damaged locations were acquired with a 60x oil objective, producing images that

196 were 295 x 295 µm. Larger (796 x 796 µm) z-stacks were acquired by blending

197 several 60x z-stack acquisitions (3x3, 15% overlap) using Nikon NIS Elements

198 software. z-stacks were re-sliced (Imagej FIJI[42], X-Z dimension) to visualize

199 fluorescence depth profiles.

AOSLO image processing

To correct residual motion from heart rate and respiration, AOSLO videos were

registered with a custom cross-correlation-based frame registration software.[40,41]

Motion correction was also applied to simultaneously collected fluorescence

videos. After registration, confocal, phase-contrast and fluorescence AOSLO

videos were temporally averaged (250 frames, 10 seconds). Blood perfusion

maps were computed by calculating the standard deviation of pixel intensity over

30 seconds[32].

INL + ONL nuclei quantification

Raw DAPI z-stacks were used for manual counting of nuclei (n = 10 mice, 3

regions per time point). Analysis regions were circular, with a 50 µm diameter

(corresponding to the size and shape of the damage region seen with confocal

AOSLO) and analyzed in depth producing volumetric cylinders through the INL or

outer nuclear layer (ONL, Figure 3d,e). Nuclei were counted manually using the

"Cell Counter" plugin in Imagej (Author: Kurt De Vos, Imagej version 1.53q).

PR + Microglia nuclei volume quantification

A single DAPI-stained z-stack (OPL + ONL) from a CX3CR1-GFP mouse was

used to quantify nuclear volume of PR's (n = 20) and microglia (n = 14). Data

was centered at a lesion location 3 days post-laser exposure. Imagej was used to

manually measure the en-face diameter of nuclei in their short and long axis. We

averaged the two measurements and assumed spherical shape for analysis.

These measurements aided in the differentiation of PR's, invading microglia and

PR phagosomes.

Quantification of RBC flux in capillaries

Using the capillary line-scan approach[27,43,44], we quantified RBC flux within an

epoch of 1 second. This spanned several cardiac cycles.[29,45] To improve SNR,

space-time images were convolved with a Gaussian spatio-temporal filter (σ =

7.5 pixels, 0.33 μm). This strategy did not interfere with spatial resolution as

pixels oversample the optical point-spread of the AOSLO by >20x.[27] RBC flux

was determined by manually marking blood cells using the "Cell counter" plugin

in Imagej.

Neutrophil density quantification

Ly-6G-647-stained retinal tissue was used to manually count neutrophils within

montaged 796 x 796 μm z-stacks. Cells were counted at control (n = 4 locations,

2 mice) and lesioned (n = 4 locations, 2 mice) locations using the "Cell Counter"

Imagej plugin. All values are reported as mean ± SD. There were no data points

omitted from any of the analysis reported in this work.

Results:

Characterization of deep focal laser damage

Four complementary imaging modalities were used to evaluate the nature and localization of the focal laser damage induced by 488 nm light: wide field SLO, OCT, AOSLO and post-mortem histology. Each is reported in turn below.

Laser damage induces focal hyperreflective lesions in outer retina imaged with wide-field SLO

To observe global and focal retinal health, we used commercial SLO. 1 day post-488 nm light exposure, both NIR and blue reflectance modalities showed hyperreflective lesions, most apparent at a deeper retinal focus position. When SLO focus was brought to the inner retina, lesions were not visible and retina appeared healthy (Figure 1b). Despite deep retinal damage, fluorescein angiography did not reveal dye leakage (Figure 1b, bottom), indicating the blood retinal barrier remained intact.

OCT B-scans reveal outer retinal hyperreflection without inner retinal damage

To assist in determination of which retinal layers are damaged by the 488 nm laser, we used OCT. Within 30 minutes post-laser exposure, OCT revealed a zone of hyperreflection within the ONL (Figure 1 – figure supplement 1). 1 day post-lesion, the focal hyperreflection remained localized to the ONL (~50 µm

wide, Figure 1c). Retina outside of damage foci appeared normal. The

hyperreflective phenotype persisted through 7 days and was cleared by 2 months

(Figure 2a). There did not appear to be any cellular excavation, "cratering" or

evidence of edema for any time point assessed. Despite the hyperreflective ONL

phenotype, Bruch's membrane appeared intact for all time points, evidenced by

lack of fluorescein leakage at the site of lesion. Additionally, retinal vessels

appeared normal in OCT B-scans.

AOSLO reveals outer retinal damage with confocal and phase-contrast

modalities

Confocal AOSLO provided micron-level detail of the lesioned area. Confirming

OCT findings, we observed hyperreflective changes localized within the ONL that

were, by 1 day, the brightest at the OPL/ONL interface (Figure 2b). At this plane,

the lesions manifest as an ellipse with the long axis in the direction of the line

scan used to create the lesion. The most prominent hyperreflective phenotype

was seen at 1 day post lesion, with diminishing size and brightness by day 3 and

7. The phenotype was largely diminished by 2 months (Figure 2b).

Using phase-contrast AOSLO also allowed visualization of translucent

cells within the retina which enabled us to image PR somas of the ONL[25,30].

Normally, the ONL is comprised of PR somata that are densely-packed[30,46]. We

found that the dense packing of individual somata was disrupted 1 day post-

exposure (~50 μm ovoid, Figure 2c), suggesting degradation or ablation of the

cell membrane.

Confirmation of outer retinal cell loss using post-mortem histology

To assess the extent of cell loss caused by focal laser exposure, we performed

DAPI staining on whole-mount retinal tissue using the same time points

assessed for *in vivo* imaging. 1 day after laser exposure, we observed mild

thickening of the ONL compared to unexposed locations only microns away. At 3

and 7 days, local ONL thinning was observed (Figure 3a). En-face planes within

the ONL revealed a loss of PR nuclei in the outer aspect of the ONL for 3 and 7

day time points, while inner ONL exhibited little evidence of cell loss (Figure 3b,

c), illustrating the precise axial confinement of the laser damage induced by this

method. By 2 months, the lesion's overall appearance and ONL thickness

returned to baseline (Figure 3a-c). This histological finding corroborates the OCT

findings observed *in vivo*.

ONL nuclear counts were reduced at locations of laser exposure. In

comparison to control locations (501317 nuclei/mm^2 ± 46198 nuclei/mm^2, mean

±SD, similar to previous work[47]), at 3 and 7 days post-exposure, we found a

corresponding reduction in the density of ONL nuclei of 18% (408965 nuclei/mm^2

± 25621 nuclei/mm^2) and 27% (367542 nuclei/mm^2 ± 9038 nuclei/mm^2)

respectively. Compared to control locations, the 3 and 7 day data resulted in p-

values of 0.17 and 0.07, respectively (Student's paired two-tailed t-test). 2

months after damage, PR nuclear densities were indistinguishable from that of

control (Figure 3f, gray bars). Despite losses of PR nuclei, total cell nuclei within

the INL remained unchanged for all time points (Figure 3f, black bars). These

data indicate that the laser exposure focally ablated PR's whilst leaving inner

retinal cells intact.

Retinal vasculature unaffected by deep retinal lesion

A concern with laser lesions of this type is that it may coagulate retinal vessels.

Motion contrast[26,27] images revealed that the vasculature remained perfused

from hours to months after laser damage suggesting that acute or long term

changes are not imparted by the laser injury (Figure 4). None of the primary

vascular stratifications within the NFL, IPL and OPL of the mouse retina[32]

showed stopped flow as a result of the laser lesion, reinforcing the findings above

regarding the axial confinement of the damage to the outer retina.

In addition to perfusion status, phase-contrast AOSLO also permitted

analysis of single blood cell flux within capillaries[27,48]. We tracked capillary flux at

different retinal depths within and above the lesion (IPL, OPL, 1 capillary each,

Figure 4 – figure supplement 1a). Blood cell flux for capillaries within lesion

locations were within the range of normal flux for the C57BL/6J mouse[48] (Figure

4 – figure supplement 1b, c). Flux tracked from hours to days in these capillaries

changed synchronously, displaying positive linear correlation (R^2 = 0.59, Figure 4

– figure supplement 1c, d). This suggested any such changes in flux were a

property of systemic perfusion, rather than locally imparted changes in flow due

to the lesion. As an additional control, we evaluated blood flux in two distant

capillaries (control, Figure 4 – figure supplement 1e). Both capillaries displayed

similar flux values from minutes to 2 months post-injury (Figure 4 – figure

supplement 1f, g) resulting in positive linear correlation (R^2 = 0.78, Figure 4 –

figure supplement 1h). Taken together, these findings suggest lesions do not

appreciably impact local RBC delivery in the capillary network.

PR laser injury promotes a robust response in nearby microglia

To observe the microglial response to PR laser injury, we imaged fluorescent

microglia in CX3CR1-GFP mice with both SLO and AOSLO. 1 day after injury,

SLO revealed bright, focal congregations of microglia at injury locations in

contrast to undamaged locations, which maintained a distribution of lateral tiling

(Figure 5a, b). The global visualization of microglia was augmented by high

resolution fluorescence AOSLO, providing enhanced detail of the microglial

response to laser injury[7,49]. Whereas AOSLO imaging of microglia in the healthy

retina displayed a distributed array of microglia with ramified processes (Figure

5c, left), laser damaged locations showed a congregation of cells 1 day post-

injury with less lateral ramification (Figure 5c, right). Within hours of the laser

exposure, we did not observe a photo-bleaching or death of regional microglia

suggesting that while the laser exposure was sufficient to damage PR's, it left

retinal microglia intact.

With phase-contrast imaging targeting the ONL, we documented a rare

event of putative pseudopod extension at a lesion site (Figure 5 – video 1). Given

the axial complexity of microglia in this layer, it is now possible that microglial

process dynamics may be revealed with this label-free approach.

We returned to the same laser-damaged locations to capture microglial

appearance at baseline, 1, 3, 7 day and 2 month time points with AOSLO to track

the natural history of the microglial response to PR damage. At 1 day post-injury,

damage locations displayed aggregations of microglia. The surrounding microglia

displayed process polarization with extensions projecting toward the injury

(Figure 6). By days 3 and 7, microglia exhibited fewer lateral projections and

somas have migrated into the ONL where they do not normally reside (Figure 6).

By 2 months post-injury, the hyperreflective phenotype was absent and microglia

once again occupied only the inner retina. The microglial distribution at 2 months

was similar to baseline, with cells exhibiting radially symmetrical branching

projections, similar to those prior to laser injury (Figure 6).

A standing question in OCT/confocal AOSLO lesion interpretation is

whether microglia contribute or directly produce the hyperreflective phenotype

seen in axial B-scans and en-face fundus images[50,51]. With confocal AOSLO, we

find that the hyperreflective phenotype is visible as early as 30 minutes,

becoming larger and brighter by 90 minutes. During these time points,

simultaneously imaged microglia remained ramified and maintained a tiled

arrangement, indicating that microglial are not the initial source of the lesion-

induced hyperreflective appearance (Figure 6 – figure supplement 1).

Despite robust microglial involvement, neutrophils do not extravasate

While we observed a robust microglial response to PR damage, there was no

evidence of neutrophil involvement at the times we examined (1, 3, 7 day, and 2

month follow-up).

In vivo fluorescence imaging allowed us to track neutrophils with AOSLO

in Catchup mice[35]. In healthy mice, we observed a sparse population of

circulating neutrophils flowing quickly within the largest retinal vessels (Figure 7 –

video 1). We also observed neutrophils moving through single capillary branches

(Figure 7 – video 2). We found that neutrophils show deformation within the small

confines of the capillary lumen, often resulting in tube or pill-shaped morphology

seen both in vivo and ex vivo (Figure 7b, top).

After laser lesion, we found no evidence of neutrophil aggregation or

extravasation for any time point assessed (Figure 8). There was also a notable

lack of rolling/crawling neutrophils (or any putative leukocyte) in large arterioles

or venules surrounding the injury (Figure 8 – video 1). Neutrophils closest to the

ONL lesion were occasionally detected within the deepest retinal capillaries.

However, these neutrophils stayed within the retinal vasculature, as evidenced

by their pill-shaped morphology and passage routes that follow the known

vascular paths seen in confocal/phase contrast images (Figure 8 – video 2).

Leukocytes, including neutrophils, often impede flow due to their large size (13.7

µm[52]) as they compress through capillaries <7 µm in diameter[48]. However, we

found no evidence of permanently stopped flow, as rare stalls would re-perfuse

similar to those previously characterized in healthy mice (Figure 7a).[48]

As a positive control, we used the EIU model in Catchup mice to show that

we could image extravasated neutrophils. This model is known to induce a strong

neutrophil response. With fluorescence AOSLO, we found Catchup-positive

neutrophils within the retinal parenchyma, providing confirmation that

extravasated neutrophils can be imaged with this modality. Moreover, we found

that neutrophils that have extravasated into the retinal parenchyma tended to

have a more spherical morphology rather than the compressed, pill-shaped

morphology of neutrophils within capillaries (Figure 7b).

Ex vivo **analysis confirms** *in vivo* **findings**

To confirm our *in vivo* findings, we examined fluorescent microglia and

neutrophils in laser-damaged retinal whole-mounts imaged with confocal

microscopy. The progressive nature of the microglial response to PR damage

and general lack of neutrophil participation corroborated *in vivo* findings.

*Ex vivo***: Microglia display dynamic morphological changes in lesion areas**

Without laser damage, microglia exhibited a tiled distribution and stellate

morphology with highly-ramified branching patterns (Figure 9a). 1 day after laser

exposure, microglial somas aggregated to the lesion location. They began to

migrate into the ONL and changed from the ramified, and pancake-flat

morphology seen in the healthy retina, to a dagger-like axial morphology (figure

9b). At 3 and 7 days after lesion, microglia migrated deeper into the outer retina

and remained aggregated with an axially elongated phenotype. 2 months after

lesion, microglial somas were no longer found in the ONL. Instead, microglia

redistributed similar to that of the healthy retina, once again co-stratifying

predominantly with the NFL and plexiform layers of the retina (Figure 9a, b,

Video 1).

Whereas we found a robust response of deep microglia at the OPL to PR

injury, we observed little migratory response of the microglia in the NFL or IPL

(Figure 9 + Figure 9 – figure supplement 1) suggesting axial and lateral

constraints on the extent of microglial recruitment.

Microglia form PR-containing phagosomes

A detailed examination of microglial involvement with PR somata in response to

PR injury revealed unique microglia-PR interactions within the ONL. In cross-

section, PR cells comprise a large portion of the retina reaching from the outer-

segment tip, inner segments, somata and spherule/pedicle synaptic contacts at

the OPL. Likewise, we saw microglial involvement with all of these layers. Within

1 day, we found microglial processes interspersed within the dense aggregation

of PR somata within the ONL. Amoeboid cells enveloped the somata of PR's and

phagocytosis was detected 1, 3 and 7 days post-laser exposure (Figure 9 –

figure supplement 2). Confocal microscopy revealed GFP-positive processes

surrounding PR nuclei, engulfing multiple somata (Figure 9 – figure supplement

2a, c). By 3 days, microglial processes and somas migrated deeper, with

processes extending into the distal portions of the PR cells including the

inner/outer segments. By two months microglial processes and somas had

retreated out of the ONL.

DAPI nuclear stain combined with confocal microscopy not only helped us

discern retinal neurons, but also allowed us to differentiate between microglia

and phagocytosed PR's. Two features were different from PR and microglial

nuclei. First, each PR displayed a uniform, homogeneous nuclear fluorescence,

while microglial nuclei appeared heterogenous and mottled (Figure 9 - figure

supplement 2a). Second, microglial nuclei were nearly 3x larger in volume

compared to PR nuclei. Microglia had a nuclear volume of 110 ± 42 μm^3 and PR

nuclei were 35 ± 5 μm^3 (p < 0.001, Figure 9 – figure supplement 2b). PR nuclei

within microglial phagosomes displayed similar nuclear volume compared to

adjacent PR's in undamaged locations (Figure 9 – figure supplement 2a).

Ex vivo: **At lesion sites, neutrophils remain within the retinal vasculature**

Corroborating in vivo AOSLO findings we did not find neutrophil aggregation or

extravasation in response to laser damage at any of the time points examined.

Ex vivo microscopy revealed that neutrophils were found within the vascular

network and did not extravasate into the neural retina. Very few neutrophils were

found in lesioned retinas, comparable healthy retinas. The few detected

neutrophils were remnants of those found within the vascular network at point of

death. Detailed 60x Z-stacks revealed "pill-shaped" morphology similar to that

seen *in vivo*, where neutrophils are compressed within capillaries. This

morphology contrasts with the spherical shape of neutrophils in response to the

EIU positive-control model (Video 2). In one exceptional example (1 day post-

damage), we observed two neutrophils in a single z-stack, one in a capillary (1)

and one at a capillary branch point (2), but none were observed to have left the

confines of the vasculature suggesting they were not recruited by activated

microglia (Figure 9c,d, Video 2).

Retinal neutrophil concentrations were quantified from larger z-stacks (796

x 796 μm, Figure 10a). Control locations (n = 2 mice, 4 z-stacks) had 15 ± 8

neutrophils per mm^2 of retina whereas lesioned locations (n = 2 mice, 4 z-stacks)

had 23 ± 5 neutrophils per mm^2 of retina (Figure 10b). The difference between

control and lesioned groups were not statistically significant (p = 0.19).

Finally, when we rendered confocal Z-stacks in cross-sectional view, we

discovered that the few fluorescent neutrophils present were found colocalized

within the tri-layered vascular stratifications of the mouse retina [32,53,54]. None

were found in non-vascular layers. Taken together, all *ex vivo* data indicates that

regardless of the robust phagocytic microglial response, neutrophils remain

localized within vessels, suggesting that neutrophils do not extravasate or

aggregate in response to this laser lesion model. These data corroborate the in

vivo findings.

Discussion:

Summary

In many retinal diseases, resident microglial populations are found to exhibit

cross-talk with systemic immune cells[15,18]. The complex temporal dynamics

between resident and systemic immune cells unfolds from seconds-to-months

and has been poorly characterized due to lack of resolution, sufficient contrast

and a non-destructive imaging approach that can track changes over time. Here,

we overcome these limitations by using phase-contrast and fluorescent imaging

with adaptive optics to visualize the interaction of multiple cell types. With this

advanced imaging technology, we reveal the absence of neutrophil involvement

in response to an acute injury, despite progressive changes in microglial activity

and morphology. Here we discuss the implications of such findings in the context

of retinal damage and more broadly, to retinal disease.

Focal retinal lesions for tracking the immune response

Seminal studies in the brain have used focal light exposure to induce targeted,

acute microglial responses.[59,60] Since then, it has been a popular damage model.

Focal lesions in the retina have been conducted in a number of popular animal

models including mouse[61,62], macaque[63–65] and cats.[66,67] Laser damage is also

clinically relevant as it is both experienced with accidental laser exposure[68,69] as

well as purposeful ablation used in photocoagulation and as a therapy for retinal

ischemic disease, retinopathy of prematurity[70] and diabetic retinopathy[71]. Thus

our results shed light toward the immune response that may be imparted due to

phototoxic damage.

Light damage offers a number of advantages for creating acute damage in

the retina. It offers placement precision that is better controlled than other acute

and invasive methods, such as incision/poke injuries[55], retinal detachment[56] and

chemical injury[57,58]. In the mouse, focal light dosage comes with greater axial

confinement within the retina by nature of the high numerical aperture of the

mouse eye[31] which imposes minimal collateral damage to layers above or below

the plane of focus (Figure 1a). The damage took on an elliptical form, likely due

to: 1) Eye motion from respiration and heart rate which spreads the light over a

larger integrative area (rather than line). 2) The impact of focal light scatter. 3) A

micron-thin line imparting damage on cells that are many microns across

manifesting as an ellipse. The majority of light exposures produced lesions of this

elliptical shape. In a few conditions, for the reasons described above, the

exposure failed to produce a strong, focal damage phenotype. To improve lesion

reproducibility, future experiments should control for subtle eye motion affecting

light damage, especially for long exposures.

Titration of the 488 nm laser leads to mild retinal damage

While it is beyond the scope of this report to catalog the vast parameter space in

which light may impose damage in the retina, it is noteworthy to discuss that the

exposure intensity, subtended angle and duration of the light source we used

delivered an intentionally mild damage to the retina. Using dosages too high

would run the risk of collateral damage to the blood retinal barrier or Bruch's

membrane[72] which could cause a choroidal neovascularization phenotype[73].

This would be undesirable for studying the native/systemic immune response as

it could confound the traditional extravasation pathways of the inflammatory

system[23,74]. Instead, we chose our laser exposure condition to impart a weak, but

reproducible loss of PRs while minimizing damage to the cells above and below

the plane of focus.

Progressive PR loss and morphology of the lesion tracked over time

The amount of light delivered did not immediately ablate PRs as peak cell loss

wasn't observed until 3-7 days after laser exposure, suggesting a progressive

damage phenotype of apoptosis or necroptotic death. Histology showed a ~27%

reduction in PR nuclei 7 days post lesion. The total number of PR nuclei within

the lesion location returned to values similar to baseline by 2 months, which was

surprising given that PRs do not regenerate. Our interpretation is that adjacent

PRs spill into the region of loss, similar to what several have described

previously[63,75]. Corroborating the histological finding of mild cell loss, OCT shows

no evidence of local edema, cavitation or excavation of the ONL after PR loss

(Figure 2a). The return of the appearance, thickness and redistribution of cells

within the ablated location indicates that retinal remodeling occurs within 2

months. The concurrence of microglia in these outer retinal bands (where they

are normally absent[76]) support the hypothesis that microglia localized to this

region to facilitate a phagocytic injury response and perhaps contribute toward

synaptic remodeling[11].

CX3CR1-GFP mice exhibit fluorescence not only in microglia

We recognize that the CX3CR1-GFP model can also label systemic cells such as

monocytes/macrophages[77]. While it is possible these cells could infiltrate the

retina in response to the lesion, we find it unlikely since there was no indication of

the leukocyte extravasation cascade (rolling/crawling/stalled cells) within the

nearest retinal vasculature. In addition to microglia, retinal perivascular

macrophages and hyalocytes also exhibit GFP fluorescence and thus that these

cells may also contribute toward damage resolution.

The hyperreflective phenotype does not arise from microglia or neutrophils

Light damage is known to create hyper-reflective bands in OCT imaging [64,78,79]. A

common speculation is that the increased backscatter may arise from local

inflammatory cells that activate or move into the damage location. In our data,

confocal AOSLO and OCT revealed a hyperreflective band at the OPL and ONL

after 488 nm light exposure (Figure 2a, b). We found that the hyperreflective

bands appeared within 30 minutes after the laser injury, preceding any

detectable microglial migration toward the damage location (Figure 2 – figure

supplement 1 and Figure 6 – figure supplement 1). We thus conclude that the

initial hyperreflective phenotype is not caused by microglial cell activity or

aggregation.

Direct activation of microglia from 488 nm light exposure was minimal

It is conceivable that 488 nm light used for either imaging (56 μW) or imparting

damage (785 μW) might activate the GFP-containing microglia used here.

However, several lines of evidence speak against this possibility. 1) We did not

observe photobleaching of CX3CR1 positive cells in response to the damage,

suggesting the fluorescence imaging light was insufficient to damage microglia.

2) For retinal injury, the focal plane was adjusted using the AOSLO, such that the

dose was axially concentrated onto the outer retina. Thus, the light dosage

received by microglia above was defocused and less than the targeted PR layer

(Figure 1a). 3) Histology showed no evident necrotic/apoptotic microglial

morphologies.

488 nm laser lesion does not photocoagulate or alter retinal circulation

Despite imparting damage to the PRs, the damage regime used here did not

alter the perfusion of the retinal circulation. We show three independent

measures that blood flow is uninterrupted, despite PR loss and activation of

microglia. 1) Fluorescein angiography (Figure 1b, bottom) revealed an absence

of vascular leakage. 2) AOSLO motion contrast vascular maps [24,27,32] displayed

persistent blood perfusion inside vessels near laser lesion sites (Figure 4). 3)

Capillary line scans indicate that RBC flux was not modified at lesion locations

and fell within the normal range[27,48] (Figure 4 – figure supplement 1). Altogether,

these three lines of evidence indicate that the lesion did not compromise the

blood-retinal barrier or impart perfusion changes within the retinal vasculature.

Resident microglia do not need systemic neutrophils for resolution of mild

laser-damage

The CNS and retina, unlike other peripheral tissues, cannot suffer from excess

inflammation as there may be dire functional consequences. Therefore it is

possible that microglia protect against exorbitant inflammation by modulating the

recruitment of systemic inflammatory cells.[4,15–17] Of these, neutrophils are often

one of the first systemic responders. Despite their helpful roles in other tissues,

neutrophils can secrete neurotoxic compounds that could present a danger to the

CNS.[80] Given their conflicted role in the body, we ask the question: to what

extent do neutrophils respond to acute neural loss in the retina? Retinal cells are

lost with age[81] and disease[82] and yet, for the organism, visual perception must

persist. There are limitations on how generalizable this mild damage to more

severe damage or disease phenotypes, but this acute damage model can begin

to provide clues about how immune cells interact in response to PR loss. In this

laser lesion model, we ablate 27% of the PRs in a 50 μm region.

We find that microglia undergo a rapid and progressive response to this

injury. We show evidence of PR phagocytosis (Figure 9 - figure supplement 2),

interaction with neighboring microglia (Figure 9 – figure supplement 1) and they

are also axially positioned to facilitate retinal remodeling (Figure 9). Furthermore,

throughout the temporal evolution of the microglial response, we find no evidence

of neutrophil recruitment despite the damage being within 10s of microns from

retinal vessels that carry them. At the onset of the neutrophil extravasation

cascade, endothelial cells in the vicinity of inflamed tissue typically elevate the

expression of adhesion molecules, facilitating the adherence and extravasation

of circulating neutrophils.[83] Furthermore, neutrophils are dependent on priming

events as prerequisite to further activation and engagement of their effector

functions.[84,85] Based on our data, we suggest that although microglia show a

strong and lasting activation, at no time point from seconds-to-months, are the

damage-associated molecular patterns or chemotactic gradients strong enough

to recruit neutrophils in response to this damage. This is evidenced in our data

from two key observations: 1) We saw no examples of systemic leukocytes

rolling in vessels adjacent to injury locations (Figure 8 – video 1). 2) Using

fluorescence, we did not observe adherent or extravasated neutrophils adjacent

to imparted PR loss (Figure 8). This may suggest that the region of insult is too

small, or that activated microglia are not sufficient for recruiting neutrophils with

damage of this magnitude. Perhaps a minimum threshold of neural damage must

be met before neutrophils will respond. It is possible that resident microglia

facilitate the necessary phagocytic and retinal remodeling response despite

release of cytokines from damaged retinal cells that would normally recruit

systemic immune cells in peripheral tissues. Such a strategy would benefit the

CNS.

Future work will explore whether there is a threshold magnitude of

neuronal cell loss required for recruitment of systemic cells that is unique to the

retina. Next studies will examine more severe or widespread injury regimes that

provide stronger activating molecular signals and interact with a larger population

of systemic cells.

Microglia may inhibit neutrophil activation

Microglia may be involved in a system that protects the CNS from propagating a

larger systemic response, potentially exacerbating disease pathologies that

would compromise overall CNS function. In another damage model[86], they report

that tissue-resident macrophages may exhibit the capacity to cloak tissue micro-

damage. This offers the possibility that resident immune cells, such as retinal

microglia, can handle small insults without inducing a chemokine cascade that

may invoke a larger systemic response that could further damage the precious

retinal tissue.[87] Regardless of the mechanism, we find that despite a robust

microglial activation that lasts for weeks, at no time point do they recruit

neutrophils. The nuance of this interaction likely represents the fine balance that

facilitates a helpful local response within the CNS that does not impart a

widespread cytokine storm that may otherwise exacerbate retinal damage.

Further work will explore whether microglia exhibit a cloaking response in the

retina, inhibiting neutrophil or other immune cell extravasation/chemotaxis toward

lesion sites. We expect such work to be pivotal in understanding the balance that

is broken or left unchecked in conditions of autoimmune disease and the

umbrella of diseases that comprise the uveitic response, a direct threat to lifelong

vision.[88]

Conclusion

Here, we have applied innovative *in vivo* imaging at the microscopic scale to

reveal the cellular immune response to a retina in jeopardy. The dynamic

environment of the retina includes a native population of resident microglia and

systemic immune cells delivered through the vasculature. These two lines of

defense work in concert in the mammalian body and are critical for maintaining

retinal homeostasis. In this work we directly study the interaction between

microglia and neutrophils, two major classes of immune cells that are implicated

in inflammatory initiation, escalation, propagation and debris removal in response

to acute geographical injury in the retina. Using cutting-edge retinal imaging

modalities, we find that resident microglia become locally activated and regionally

responsive to focal laser lesion. Cells migrate away from their stratified locations

near plexiform layers of the retina and migrate toward the site of damage within

hours to weeks after injury. However, systemic neutrophils, which are typically

regarded as first-line responders to tissue damage, are not recruited to this

damage despite neutrophils flowing within 10s of microns away from the location

of damage (Figure 8 – video 2). Beyond the context of this specific finding, we

share this work with the excitement that AOSLO cellular level imaging may reveal

the interaction of multiple immune cell types in the living retina. By using

fluorophores associated with specific immune cell populations, the complex

dynamics that orchestrate the immune response may be examined in this

specialized tissue. This work and future studies may reveal further insights to the

interactions of single immune cells in the living body in a non-invasive way.

Acknowledgements: Thanks to Colin Chu, Justin Elstrott and Tiffany Heaster

for useful intellectual conversations. Thanks to Minsoo Kim for generously

transferring the Catchup mouse strain.

References:

1. Stone, J. *et al.* Mechanisms of photoreceptor death and survival in mammalian retina. *Progress in Retinal and Eye Research* **18**, 689–735 (1999).

2. García-Ayuso, D., Di Pierdomenico, J., Vidal-Sanz, M. & Villegas-Pérez, M. P. Retinal Ganglion Cell Death as a Late Remodeling Effect of Photoreceptor Degeneration. *Int J Mol Sci* **20**, 4649 (2019).

3. Boycott, B. B. & Hopkins, J. M. Microglia in the retina of monkey and other mammals: its distinction from other types of glia and horizontal cells. *Neuroscience* **6**, 679–688 (1981).

4. Silverman, S. M. & Wong, W. T. Microglia in the Retina: Roles in Development, Maturity, and Disease. *Annual Review of Vision Science* **4**, 45–77 (2018).

5. Wang, X. *et al.* Requirement for Microglia for the Maintenance of Synaptic Function and Integrity in the Mature Retina. *J. Neurosci.* **36**, 2827–2842 (2016).

6. Zhang, Y. *et al.* Repopulating retinal microglia restore endogenous organization and function under CX3CL1-CX3CR1 regulation. *Science Advances* **4**, eaap8492 (2018).

7. Joseph, A. *et al.* Imaging the dynamics of individual processes of microglia in the living retina in vivo. *Biomed. Opt. Express, BOE* **12**, 6157–6183 (2021).

8. Nimmerjahn, A., Kirchhoff, F. & Helmchen, F. Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science* **308**, 1314–1318 (2005).

9. Neumann, H., Kotter, M. R. & Franklin, R. J. M. Debris clearance by microglia: an essential link between degeneration and regeneration. *Brain* **132**, 288 (2009).

10. Zabel, M. K. *et al.* Microglial phagocytosis and activation underlying photoreceptor degeneration is regulated by CX3CL1-CX3CR1 signaling in a mouse model of retinitis pigmentosa. *Glia* **64**, 1479–1491 (2016).

11. Tremblay, M.-È. & Majewska, A. K. A role for microglia in synaptic plasticity? *Commun Integr*
Biol **4**, 220–222 (2011).

12. Sipe, G. O. *et al.* Microglial P2Y12 is necessary for synaptic plasticity in mouse visual cortex.
Nat Commun **7**, 10905 (2016).

13. Márquez-Ropero, M., Benito, E., Plaza-Zabala, A. & Sierra, A. Microglial Corpse Clearance:
Lessons From Macrophages. *Front Immunol* **11**, 506 (2020).

14. Karlen, S. J., Miller, E. B. & Burns, M. E. Microglia Activation and Inflammation During the
Death of Mammalian Photoreceptors. *Annual Review of Vision Science* **6**, 149–169 (2020).

15. Okunuki, Y. *et al.* Retinal microglia initiate neuroinflammation in ocular autoimmunity. *Proc*
Natl Acad Sci U S A **116**, 9989–9998 (2019).

16. Kremlev, S. G., Roberts, R. L. & Palmer, C. Differential expression of chemokines and
chemokine receptors during microglial activation and inhibition. *Journal of*
Neuroimmunology **149**, 1–9 (2004).

17. Babcock, A. A., Kuziel, W. A., Rivest, S. & Owens, T. Chemokine Expression by Glial Cells
Directs Leukocytes to Sites of Axonal Injury in the CNS. *J. Neurosci.* **23**, 7922–7930 (2003).

18. Boyce, M. *et al.* Microglia–Neutrophil Interactions Drive Dry AMD-like Pathology in a Mouse
Model. *Cells* **11**, 3535 (2022).

19. Zhou, H., Lapointe, B. M., Clark, S. R., Zbytnuik, L. & Kubes, P. A Requirement for Microglial
TLR4 in Leukocyte Recruitment into Brain in Response to Lipopolysaccharide. *The Journal of*
Immunology **177**, 8103–8110 (2006).

20. physiological_data_000664.pdf. https://www.jax.org/-/media/jaxweb/files/jax-mice-and-
services/physiological_data_000664.pdf (2007).

21. Burn, G. L., Foti, A., Marsman, G., Patel, D. F. & Zychlinsky, A. The Neutrophil. *Immunity* **54**,

1377–1391 (2021).

22. Peiseler, M. & Kubes, P. More friend than foe: the emerging role of neutrophils in tissue

repair. *J Clin Invest* **129**, 2629–2639 (2019).

23. Joseph, A., Chu, C. J., Feng, G., Dholakia, K. & Schallek, J. Label-free imaging of immune cell

dynamics in the living retina using adaptive optics. *eLife* **9**, e60547 (2020).

24. Sulai, Y. N., Scoles, D., Harvey, Z. & Dubra, A. Visualization of retinal vascular structure and

perfusion with a nonconfocal adaptive optics scanning light ophthalmoscope. *J Opt Soc Am*

A Opt Image Sci Vis **31**, 569–579 (2014).

25. Guevara-Torres, A., Williams, D. R. & Schallek, J. B. Origin of cell contrast in offset aperture

adaptive optics ophthalmoscopy. *Opt. Lett., OL* **45**, 840–843 (2020).

26. Chui, T. Y. P., VanNasdale, D. A. & Burns, S. A. The use of forward scatter to improve retinal

vascular imaging with an adaptive optics scanning laser ophthalmoscope. *Biomedical Optics*

Express **3**, 2537 (2012).

27. Guevara-Torres, A., Joseph, A. & Schallek, J. B. Label free measurement of retinal blood cell

flux, velocity, hematocrit and capillary width in the living mouse eye. *Biomed. Opt. Express,*

BOE **7**, 4228–4249 (2016).

28. Joseph, A. *et al.* Label-free tracking of single blood cells in the retinal immune response.

Journal of Vision **19**, 20–20 (2019).

29. Joseph, A., Guevara-Torres, A. & Schallek, J. Imaging single-cell blood flow in the smallest to

largest vessels in the living retina. *eLife* **8**, e45077 (2019).

30. Guevara-Torres, A., Williams, D. R. & Schallek, J. B. Imaging translucent cell bodies in the

living mouse retina without contrast agents. *Biomedical Optics Express* **6**, 2106 (2015).

31. Geng, Y. *et al.* Adaptive optics retinal imaging in the living mouse eye. *Biomed. Opt. Express*

3, 715–734 (2012).

32. Schallek, J., Geng, Y., Nguyen, H. & Williams, D. R. Morphology and topography of retinal

pericytes in the living mouse retina using in vivo adaptive optics imaging and ex vivo

characterization. *Invest. Ophthalmol. Vis. Sci.* **54**, 8237–8250 (2013).

33. Pinhas, A. *et al.* In vivo imaging of human retinal microvasculature using adaptive optics

scanning light ophthalmoscope fluorescein angiography. *Biomed Opt Express* **4**, 1305–1317

(2013).

34. Chu, C. J. *et al.* Multimodal analysis of ocular inflammation using the endotoxin-induced

uveitis mouse model. *Disease Models & Mechanisms* **9**, 473–481 (2016).

35. Hasenberg, A. *et al.* Catchup: a mouse model for imaging-based tracking and modulation of

neutrophil granulocytes. *Nat Methods* **12**, 445–452 (2015).

36. Crane, I. J. & Liversidge, J. Mechanisms of leukocyte migration across the blood–retina

barrier. *Semin Immunopathol* **30**, 165–177 (2008).

37. Xu, H. *et al.* Involvement of CD44 in leukocyte trafficking at the blood-retinal barrier. *Journal*

of Leukocyte Biology **72**, 1133–1141 (2002).

38. Xu, H. *et al.* Reduction in shear stress, activation of the endothelium, and leukocyte priming

are all required for leukocyte passage across the blood—retina barrier. *Journal of Leukocyte*

Biology **75**, 224–232 (2004).

39. Rosenbaum, J. T., McDevitt, H. O., Guss, R. B. & Egbert, P. R. Endotoxin-induced uveitis in

rats as a model for human disease. *Nature* **286**, 611–613 (1980).

40. Dubra, A. & Harvey, Z. Registration of 2D Images from Fast Scanning Ophthalmic

Instruments. in *Biomedical Image Registration* (eds. Fischer, B., Dawant, B. M. & Lorenz, C.)

60–71 (Springer, Berlin, Heidelberg, 2010). doi:10.1007/978-3-642-14366-3_6.

41. Yang, Q. *et al.* Closed-loop optical stabilization and digital image registration in adaptive

optics scanning light ophthalmoscopy. *Biomedical Optics Express* **5**, 3174 (2014).

42. Fiji: an open-source platform for biological-image analysis | Nature Methods.

https://www.nature.com/articles/nmeth.2019.

43. Joseph, A., Guevara-Torres, A., Williams, D. R. & Schallek, J. B. Measurement of blood flow

in the largest vessels and smallest capillaries in the living mouse retina using an adaptive

optics scanning light ophthalmoscope. *Invest. Ophthalmol. Vis. Sci.* **56**, 3323–3323 (2015).

44. Dholakia, K., Guevara-Torres, A. & Schallek, J. B. Early changes in blood cell flux in retinal

capillaries of diabetic and healthy mice imaged in vivo. *Invest. Ophthalmol. Vis. Sci.* **61**, 806–

806 (2020).

45. Feng, G. *et al.* High-resolution structural and functional retinal imaging in the awake

behaving mouse. *Commun Biol* **6**, 1–15 (2023).

46. Carter-Dawson, L. D. & LaVail, M. M. Rods and cones in the mouse retina. I. Structural

analysis using light and electron microscopy. *J. Comp. Neurol* **188**, 245–262 (1979).

47. Jeon, C. J., Strettoi, E. & Masland, R. H. The major cell populations of the mouse retina. *J*

Neurosci **18**, 8936–8946 (1998).

48. Dholakia, K. Y., Guevara-Torres, A., Feng, G., Power, D. & Schallek, J. In Vivo Capillary

Structure and Blood Cell Flux in the Normal and Diabetic Mouse Eye. *Invest Ophthalmol Vis*

Sci **63**, 18 (2022).

49. Wahl, D. J., Ng, R., Ju, M. J., Jian, Y. & Sarunic, M. V. Sensorless adaptive optics multimodal

en-face small animal retinal imaging. *Biomed Opt Express* **10**, 252–267 (2019).

50. Puthenparampil, M. *et al.* Hyper-Reflecting Foci in Multiple Sclerosis Retina Associate With

Macrophage/Microglia-Derived Cytokines in Cerebrospinal Fluid. *Front Immunol* **13**, 852183

(2022).

51. Pilotto, E. *et al.* Hyper-reflective retinal foci as possible in vivo imaging biomarker of

microglia activation in von Hippel-Lindau disease. *PLoS One* **17**, e0272318 (2022).

52. Kornmann, L. M. *et al.* Echogenic perfluorohexane-loaded macrophages adhere in vivo to

activated vascular endothelium in mice, an explorative study. *Cardiovasc Ultrasound* **13**, 1

(2015).

53. Paques, M. *et al.* Structural and hemodynamic analysis of the mouse retinal

microcirculation. *Invest. Ophthalmol. Vis. Sci* **44**, 4960–4967 (2003).

54. Stahl, A. *et al.* The Mouse Retina as an Angiogenesis Model. *Investigative Ophthalmology &*

Visual Science **51**, 2813–2826 (2010).

55. Senut, M.-C., Gulati-Leekha, A. & Goldman, D. An Element in the α1-Tubulin Promoter Is

Necessary for Retinal Expression during Optic Nerve Regeneration But Not after Eye Injury

in the Adult Zebrafish. *J. Neurosci.* **24**, 7663–7673 (2004).

56. Maidana, D. E., Gonzalez-Buendia, L., Miller, J. W. & Vavvas, D. G. Local photoreceptor cell

death differences in the murine model of retinal detachment. *Sci Rep* **11**, 18798 (2021).

57. Wan, J. *et al.* Preferential regeneration of photoreceptor from Müller glia after retinal

degeneration in adult rat. *Vision Research* **48**, 223–234 (2008).

58. Rösch, S. *et al.* The effects of iodoacetic acid on the mouse retina. *Graefes Arch Clin Exp*

Ophthalmol **253**, 25–35 (2015).

59. Nimmerjahn, A., Kirchhoff, F. & Helmchen, F. Resting microglial cells are highly dynamic

surveillants of brain parenchyma in vivo. *Science* **308**, 1314–1318 (2005).

60. Davalos, D. *et al.* ATP mediates rapid microglial response to local brain injury in vivo. *Nat*

Neurosci **8**, 752–758 (2005).

61. Shah, R. S., Soetikno, B. T., Lajko, M. & Fawzi, A. A. A Mouse Model for Laser-induced

Choroidal Neovascularization. *J Vis Exp* 53502 (2015) doi:10.3791/53502.

62. Khan, A. H. *et al.* A laser-induced mouse model of progressive retinal degeneration with

central sparing displays features of parafoveal geographic atrophy. *Sci Rep* **13**, 4194 (2023).

63. Strazzeri, J. M. *et al.* Focal damage to macaque photoreceptors produces persistent visual

loss. *Exp Eye Res* **119**, 88–96 (2014).

64. Dhakal, K. R. *et al.* Localized Photoreceptor Ablation Using Femtosecond Pulses Focused

With Adaptive Optics. *Transl Vis Sci Technol* **9**, 16 (2020).

65. Schwarz, C. *et al.* Selective S Cone Damage and Retinal Remodeling Following Intense

Ultrashort Pulse Laser Exposures in the Near-Infrared. *Investigative Ophthalmology & Visual*

Science **59**, 5973–5984 (2018).

66. Calford, M. B., Schmid, L. M. & Rosa, M. G. P. Monocular focal retinal lesions induce short–

term topographic plasticity in adult cat visual cortex. *Proceedings of the Royal Society of*

London. Series B: Biological Sciences **266**, 499–507 (1999).

67. Dreher, B., Burke, W. & Calford, M. B. Chapter 15 Cortical plasticity revealed by

circumscribed retinal lesions or artificial scotomas. in *Progress in Brain Research* vol. 134

217–246 (Elsevier, 2001).

68. Vitellas, C. *et al.* CONE PHOTORECEPTOR INTEGRITY ASSESSED WITH ADAPTIVE OPTICS

IMAGING AFTER LASER POINTER-INDUCED RETINAL INJURY. *Retinal Cases and Brief Reports*

16, 586 (2022).

69. Linton, E. *et al.* Retinal burns from laser pointers: a risk in children with behavioural

problems. *Eye* **33**, 492–504 (2019).

70. Current ROP Therapies: How Laser and Anti-VEGF Compare. *American Academy of*

Ophthalmology https://www.aao.org/eyenet/article/current-rop-therapies-how-laser-

antivegf-compare (2014).

71. Reddy, S. V. & Husain, D. Panretinal Photocoagulation: A Review of Complications. *Semin*

Ophthalmol **33**, 83–88 (2018).

72. Haupert, C. L. *et al.* Optimal laser power to rupture Bruch's membrane and the retinal vein

in the pig. *Ophthalmic Surg Lasers Imaging* **34**, 122–127 (2003).

73. Shah, R. S., Soetikno, B. T., Lajko, M. & Fawzi, A. A. A Mouse Model for Laser-induced

Choroidal Neovascularization. *J Vis Exp* e53502 (2015) doi:10.3791/53502.

74. Ley, K., Laudanna, C., Cybulsky, M. I. & Nourshargh, S. Getting to the site of inflammation:

the leukocyte adhesion cascade updated. *Nat Rev Immunol* **7**, 678–689 (2007).

75. Busch, E. M., Gorgels, T. G. & Van Norren, D. Filling-in after focal loss of photoreceptors in

rat retina. *Exp Eye Res* **68**, 485–492 (1999).

76. Lee, J. E., Liang, K. J., Fariss, R. N. & Wong, W. T. Ex vivo Dynamic Imaging of Retinal

Microglia using Time-lapse Confocal Microscopy. *Invest Ophthalmol Vis Sci* **49**, 4169–4176

(2008).

77. Jung, S. *et al.* Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and

green fluorescent protein reporter gene insertion. *Mol Cell Biol* **20**, 4106–4114 (2000).

78. Miller, E. B., Zhang, P., Ching, K., Pugh, E. N. & Burns, M. E. In vivo imaging reveals transient

microglia recruitment and functional recovery of photoreceptor signaling after injury.

Proceedings of the National Academy of Sciences **116**, 16603–16612 (2019).

79. Wu, C. Y. *et al.* Acute Solar Retinopathy Imaged With Adaptive Optics, Optical Coherence

Tomography Angiography, and En Face Optical Coherence Tomography. *JAMA*

Ophthalmology **136**, 82–85 (2018).

80. Allen, C. *et al.* Neutrophil Cerebrovascular Transmigration Triggers Rapid Neurotoxicity

through Release of Proteases Associated with Decondensed DNA. *The Journal of*

Immunology **189**, 381–392 (2012).

81. Panda-Jonas, S., Jonas, J. B. & Jakobczyk-Zmija, M. Retinal Photoreceptor Density Decreases

with Age. *Ophthalmology* **102**, 1853–1859 (1995).

82. Margalit, E. & Sadda, S. R. Retinal and Optic Nerve Diseases. *Artificial Organs* **27**, 963–974

(2003).

83. Sadik, C. D., Kim, N. D. & Luster, A. D. Neutrophils cascading their way to inflammation.

Trends Immunol **32**, 452–460 (2011).

84. Worthen, G. S. *et al.* Neutrophil-mediated pulmonary vascular injury. Synergistic effect of

trace amounts of lipopolysaccharide and neutrophil stimuli on vascular permeability and

neutrophil sequestration in the lung. *Am Rev Respir Dis* **136**, 19–28 (1987).

85. Linas, S. L., Whittenburg, D., Parsons, P. E. & Repine, J. E. Mild renal ischemia activates

primed neutrophils to cause acute renal failure. *Kidney Int* **42**, 610–616 (1992).

86. Uderhardt, S., Martins, A. J., Tsang, J. S., Lämmermann, T. & Germain, R. N. Resident

Macrophages Cloak Tissue Microlesions to Prevent Neutrophil-Driven Inflammatory

Damage. *Cell* **177**, 541-555.e17 (2019).

87. Pfeifer, C. W. *et al.* Dysregulated CD200-CD200R signaling in early diabetes modulates

microglia-mediated retinopathy. *Proceedings of the National Academy of Sciences* **120**,

e2308214120 (2023).

88. Miserocchi, E., Fogliato, G., Modorati, G. & Bandello, F. Review on the Worldwide

Epidemiology of Uveitis. *European Journal of Ophthalmology* **23**, 705–717 (2013).

Figures & videos:

Figure 1. Laser injury assessed with commercial SLO and OCT. **(A)** 488 nm light is focused onto the photoreceptor outer segments using AOSLO. Created with Biorender.com. **(B)** 30° SLO images of NIR reflectance, blue reflectance and fluorescein angiography of a mouse retina 1 day after laser exposure. Three focal planes are shown. NIR and blue reflectance reveal small hyperreflective regions below the superficial plane. Fluorescein reveals intact vasculature with no sign of leakage. Arrows indicate regions with imparted laser damage (1-4). **(C)** OCT B-scans passing through laser-exposed regions indicated in B. Exposures produced a focal hyperreflective band within the ONL with adjacent retina appearing healthy. OCT images were spatially averaged (~30 µm, 3 B-scans). Scale bars = 200 µm horizontal, 200 µm vertical.

Figure 1 - figure supplement 1 Lesion location tracked from minutes to 1 day
with OCT. After baseline OCT acquisition, OCT was performed every 5-7
minutes for one hour after 488 nm light exposure. 6 hour and 1 day time points
were subsequently acquired. A band of hyperreflectivity forms near the OPL/ONL
interface within 30 minutes of 488 nm light exposure. Hyperreflective band,
spreads deeper into the ONL within ~1 hour. OCT images were spatially
averaged (~30 µm, 8 B-scans). Scale bar = 40 µm horizontal, 100 µm vertical.

Figure 2. Laser damage temporally tracked with AOSLO and OCT. Laser-exposed retina was tracked with OCT **(A)**, confocal **(B)** and phase-contrast **(C)** AOSLO for baseline, 1, 3, 7 day and 2 month time points. OCT and confocal AOSLO display a hyperreflective phenotype that was largest/brightest at 1 day and became nearly invisible by 2 months. Dashed oval indicates region targeted for laser injury. Phase-contrast AOSLO revealed disrupted photoreceptor soma's 1 day after laser injury. Phase contrast data was not acquired for time points 3 days-2 months due to development of cataract which obscured the phase contrast signal. OCT images were spatially averaged (~30 μm, 8 B-scans). Scale bars = 40 μm, OCT image vertical scale bar = 100 μm.

Figure 3. Retinal histology confirms photoreceptor ablation and preservation of inner retinal cells. Cross sectional view **(A)** and en-face **(B+C)** images of DAPI-stained whole-mount retinas (5 mice) at laser injury locations over time. By 1 day, ONL becomes thicker at lesion location, but thinner by 3 and 7 days. By 2 months, the ONL appeared similar to that of control. The inner (**B**, solid rectangle) and outer (**C**, dashed rectangle) stratum of ONL show axial differences in ONL loss**.** Most cell loss was seen in the outer aspect of the ONL **(C)**. Scale bars = 40 µm. **(D)** Cross section of DAPI-stained retina displaying INL and ONL regions for quantification. Each analysis region was 50 µm across and encompassed the entire INL or ONL. **(E)** En-face images show 50 µm diameter circles used for analysis. **(F)** Nuclei density for control and post-injury time points. ONL nuclei were reduced at 3 and 7 days (p = 0.17 and 0.07 respectively) while INL density remained stable (n = 10 mice, 3 unique regions per time point). Error bars display mean ± 1 SD.

Figure 4. Microvascular perfusion unchanged after laser damage. A single
location tracked over time and at three vascular plexuses using AOSLO. Motion-
contrast images were generated from confocal videos to reveal vascular
perfusion status. Retinal vasculature remained perfused for all time points
tracked and at all depths. White oval indicates damage location. Scale bar = 40
μm.

Figure 4 – figure supplement 1 Measurement of single-cell blood flux after
laser damage using phase contrast AOSLO. **Mouse 1: (A)** The vascular plexus
corresponding to IPL (cyan) and OPL (red) were targeted for flux determination. Blood
cell flux was measured for 2 capillaries within the same field, at different depths. Arrows
show the location for repeated line scan acquisitions. **(B)** RBC flux images were tracked
up to 7 days post-damage. Scale bars = 10 ms, horizontal and 5 µm vertical. **(C)**
Capillary flux measurements over 7 days. Despite the outer capillary displaying higher
flux, both inner and outer capillaries changed synchronously for each time point. **(D)**
Correlation of inner and outer capillary flux. Linear regression model displays a weak
positive correlation (black dotted line). **Mouse 2: (E)** Left: Representative 55° SLO
image showing regions targeted for capillary flux measurement. One region was subject
to 488 nm laser damage and the other was left unlasered (Control). Scale bar = 200 µm.
Right: Capillaries targeted for blood cell flux measurement. Arrows show the location for
repeated line scan acquisitions. Scale bar = 40 µm. **(F)** RBC flux images were tracked
up to 2 months post-damage. Scale bars = 10 ms horizontal and 5 µm vertical. **(G)**
Capillary flux remained similar at lesion and control locations over all time points
assessed. Gray shaded regions indicate range for normal capillary flux in the healthy
C57BL/6J mouse (Dholakia et al 2022). **(H)** Correlation of flux in lesion and control
locations. Linear regression model displays a positive correlation (black dotted line).

Figure 5. Microglial response 1 day after laser injury imaged in vivo with
fluorescence SLO and AOSLO. **(A)** Left: Deep-focus NIR SLO fundus image (55°
FOV) of laser-injured retina. White arrowheads point to damaged locations
showing hyperreflective regions. Inset scale bar = 40 µm. Right: Fluorescence
fundus image from same location. Fluorescent CX3CR1-GFP microglia are
distributed across the retina and show congregations at laser-damaged locations.
Scale bar = 200 µm. **(B)** Magnified SLO images of microglia at laser-damaged
and control locations (indicated in A, right, white boxes). Control location displays
distributed microglial whereas microglia at the lesion location are bright and
focally aggregated. **(C)** Fluorescence AOSLO images show greater detail of cell
morphology at the same scale. In control locations microglia showed ramified
morphology and distributed concentration whereas damage locations revealed
dense aggregation of many microglia that display less ramification. Scale bars =
40 µm.

971   *Refer to .avi file*

972   **Figure 5 – video 1** Dynamic pseudopodia imaged with phase-contrast AOSLO 1
973   day post-injury. At the OPL/ONL border, a putative microglial pseudopod
974   extension was captured among a field of static, disrupted PR somas. Video is 3
975   minutes elapsed. Scale bars = 20 μm.

Figure 6. Microglial response to laser injury tracked with AOSLO. Simultaneously acquired NIR confocal and fluorescence AOSLO images across different retinal depths. Data are from one CX3CR1-GFP mouse tracked for 2 months. Microglia swarm to hyperreflective locations within 1 day. Microglia maintain an aggregated density for days and resolve by 2 months after damage. Scale bar = 40 µm.

Figure 6 – figure supplement 1 Hyperreflective appearance emerges before
microglia swarm to damage location. AOSLO confocal and fluorescence images
were acquired for baseline, 30, 90 minute and 1 day post-laser exposure. The
hyperreflective phenotype appeared within 30 minutes post damage. Microglia
were not found to aggregate until ~1 day after. Scalebar = 40 μm.

Figure 7. Neutrophil morphology imaged in vivo using AOSLO. **(A)** Phase-
contrast, motion-contrast and fluorescence AOSLO reveal the impact of passing
neutrophils on single capillaries. A rare and exemplary event shows a neutrophil
transiently impeding capillary blood flow for minutes in healthy retina. Scale bar =
40 μm. **(B)** In-vivo AOSLO and ex-vivo fluorescence microscopy show
neutrophils in two states. Neutrophils within capillaries displayed elongated,
tubular morphology. Extravasated neutrophils were more spherical. Bottom
images show extravasated neutrophils in response to LPS model for comparison
(not laser damage model). Scale bar = 20 μm.

1002

Refer to .avi file

Figure 7 – video 1 Neutrophil dynamics within a primary retinal vessel in a
Catchup mouse. Confocal (left) and fluorescence (right) AOSLO were acquired
from a large primary retinal vessel simultaneously. Several quickly flowing
neutrophils are seen as streaks within the lumen of the large retinal vessel. Video
plays in real-time (25 FPS, 20 seconds). Scale bar = 20 μm.

Refer to .avi file

Figure 7 – video 2 Neutrophils imaged in capillaries of a Catchup mouse.
Motion-contrast (left) and fluorescence (right) AOSLO show a single neutrophil
moving through a branch of the OPL capillary network. Red circles/arrows
indicate neutrophil path within the vascular perfusion map. Rare neutrophils were
found to move slowly through capillaries. Video is 0.4 seconds elapsed. Scale
bar = 20 μm.

1016

1017

Figure 8. Retinal damage location tracked with AOSLO does not show
accumulation of neutrophils. A single retinal location was tracked in a Catchup
mouse from baseline to 2 months after lesion. Location of the lesion is apparent
at 1 and 3 days post injury with diminishing visibility after 1 week. Despite the
ability to detect neutrophils (Figure 7), we did not observe stalled, aggregated or
an accumulation of neutrophils at any time point evaluated. This evaluation was
confirmed at multiple depths ranging from the NFL to the ONL of the retina in
vivo. Scale bar = 40 μm.

1026

1027

Figure 8 – figure supplement 1 A single mouse was tracked for 0.5 - 3.5 hours after a deep retinal lesion was placed at the center of the imaged field, adjacent to a large venule. Neutrophils did not extravasate within this early window post-lesion (also see figure 8 – video 3).

1032

Refer to .avi file

Figure 8 – video 1 Neutrophil dynamics within a primary retinal vessel in a
Catchup mouse 1 day after a deep retinal laser lesion is placed nearby. Confocal
(left) and fluorescence (right) AOSLO images were acquired simultaneously.
Quickly flowing neutrophils are seen as streaks within the lumen of this large
retinal vessel and despite the adjacent deep retinal lesion (within 100 μm), there
is no indication of rolling or crawling neutrophils. Video plays in real-time (25
FPS, 20 seconds). Scale bar = 20 μm.

Refer to .avi file

Figure 8 – video 2 Neutrophil dynamics within an OPL capillary in a Catchup
mouse 1 day after deep laser injury. Motion-contrast (left) and fluorescence
(right) AOSLO are displayed in tandem. At 1 day post-laser injury, a single
neutrophil is seen moving through a capillary that runs directly through the injury
site (yellow oval). Despite injury, neutrophil does not slow or stall at the lesion
location. Video is 0.68 seconds elapsed. Scale bar = 20 μm.

1048

1049 **Figure 8 – video 3** Neutrophil dynamics 2.5 hours post-lesion. Neutrophil
1050 movement indicates they stayed within OPL capillaries and did not extravasate
1051 into the retinal parenchyma at this early timepoint.

1052

Figure 9. Neutrophil behavior under laser injury, as observed through ex vivo
confocal microscopy, remains unresponsive despite marked microglial activation
(A) En-face max intensity projection images of inner and outer (separated by
approximate INL center) retinal microglia/neutrophils in Ly-6G-647-stained
CX3CR1-GFP retinas. Microglia display focal aggregation in the outer retina for
1, 3 and 7 day time points that is resolved by 2 months. Neutrophils do not
aggregate or colocalize to the injury location at any time point. Z-stacks were
collected from 5 mice for the indicated time points. **(B)** Cross-sectional views of
en-face z-stacks presented in A, including DAPI nuclear label. White dotted line
indicates 100 µm region expanded below. Microglia migrate into the ONL by 1, 3
and 7 days post-laser injury and return to an axial distribution similar to that of
control by 2 months. The few neutrophils detected remained within the inner
retina. Scale bars = 40 µm. **(C)** Orthogonal view of DAPI-stained retina with Ly-
6G-647-labelled overlay 1 day post-laser-injury. In a rare example, 2 neutrophils
are found within the IPL/OPL layers despite a nearby outer retinal laser lesion.
Scale bar = 20 µm. **(D)** Magnified 3D cubes representing cell 1 and 2 in C. Cell 1
displays pill-shaped morphology and cell 2 is localized to a putative capillary
branch-point. Each are confined within vessels suggesting they do not
extravasate in response to laser injury.

1072

Figure 9 – figure supplement 1 Neutrophil/microglial response to laser injury
tracked with *ex vivo* confocal microscopy. Simultaneously acquired GFP-positive
microglia and Ly-6G-647-positive neutrophils were imaged with confocal
microscopy in 5 CX3CR1-GFP mice. En-face images for several retinal depths
are displayed. By 1, 3 and 7 days post-lesion, microglia have migrated into the
outer retina, many appearing amoeboid and displaying fewer laterally-branching
projections. Despite the deep microglial response, neutrophils stay within the
inner retina and are not found in the avascular outer retinal layers. Scale bar = 40
μm.

1082

1083

Figure 9 – figure supplement 2 Microglial PR phagosomes in the outer retina
assessed with *ex vivo* confocal imaging. **(A)** En-face images of outer ONL in a
DAPI-stained CX3CR1-GFP mouse 3 days post-laser-injury (top row). Microglia
have infiltrated deep into the ONL and several PR phagosomes were identified.
White arrows indicate locations for a single microglia (i), PR (ii) and PR
phagosome (iii). These locations were expanded and displayed below. Microglia
exhibited a heterogeneous nuclear staining pattern while PR nuclei exhibited
homogenous DAPI fluorescence pattern. PR's displayed this pattern regardless
of whether they were within a microglial phagosome or not. Top scale bar = 20
μm, bottom scale bar = 2 μm**. (B)** A finely-sliced (0.1 μm step size) outer retinal z-
stack of DAPI-stained CX3CR1-GFP retina was used to quantify the average
nuclear volume for infiltrated microglia (n = 14 nuclei) and PR's (n = 20 nuclei) for
the same lesion site presented in A. On average, microglia had a statistically
significant (p < 0.001) nuclear volume that was >3x that of PR's. These
measurements allowed us to discriminate microglial somas from PR
phagosomes. Error bars display mean + 1 SD. **(C)** Cross sections of DAPI-

stained outer retina in CX3CR1-GFP mice for 1, 3 and 7 days post-laser injury
(n=3 mice). 3 representative planes (X-Z) through the lesion are displayed for
each time point. Microglia form PR phagosomes within the ONL and microglial
processes were seen extended into the PR inner/outer segment layer. Arrows
label various morphological features seen at lesion sites: microglial somas
(yellow), diving microglial process (violet), PR phagosome (red), microglial
inner/outer segment process (cyan). Scale bar = 20 μm.

1107

1108

Figure 10. Quantification of neutrophils in laser damaged retinas assessed with
ex vivo confocal microscopy over a wide-field. **(A)** Representative image displays
neutrophils quantified using large-field (796 x 796 µm) z-stacks for control or 1
day after injury time points. In both control and laser injured retinas, neutrophils
were sparse and confined to locations within capillaries suggesting they were the
native fraction of circulating neutrophils at time of death. Inset displays expanded
image of a single neutrophil. Scale bar = 200 µm. **(B)** Neutrophils quantified and
displayed as the number of neutrophils per retinal area. The difference in number
of neutrophils in control vs lesioned retinas was not statistically significant (p =
0.19). Error bars display mean + 1 SD.

1119

1120

Refer to .avi file

Video 1 Rotating 3D cubes of outer retinal nuclei and microglia after focal laser
injury. Outer retinal Z-stacks of DAPI-stained whole-mount CX3CR1-GFP retinal
tissue were imaged for control, 1, 3, 7 day and 2 month time points (n = 5 mice).
DAPI + microglia composite cubes are displayed above and microglia cubes are
displayed below. By 1 day, microglia send projections into the ONL, by 3 and 7
days, microglial somas have migrated into the ONL. Microglia within the ONL
exhibit different morphology compared to the baseline condition. By 2 months,
microglia are found back within the OPL, exhibiting lateral projections, similar to
baseline. Scale bar = 40 μm.

1131

Refer to .avi file

Video 2 Rotating 3D cubes of single neutrophils after laser injury or endotoxin
induced uveitis. *Ex vivo* confocal z-stacks (0.1 μm steps) allowed detailed
visualization of single neutrophils at 1 day after laser injury or 1 day after
intravitreal LPS injection. After laser injury, neutrophils maintain a tubular, pill-
shaped morphology (left). Occasionally, they would come to rest at capillary
branch points (middle). In the EIU model, neutrophils extravasate into the retinal
parenchyma and exhibit more spheroid morphology (right). We did not observe
neutrophils to exhibit the extravasated morphology in response to laser injury.

1141